1 **Evaluation of graphene oxide-mediated NET formation using HL-60-derived**

2 **neutrophil-like cells**

4 Kiyoshi Fukuhara[1]*, Takashi Obama[2], Hitomi Ohinata[2], Takashi Takaki[3], Masaru Kato[4],

5 Yukari Totsuka[5], Hiroyuki Itabe[2], Akiko Ohno[6]

6
7 [1]Division of Organic and Medicinal Chemistry, Department of Pharmaceutical Sciences,
8 Showa Medical University School of Pharmacy, Tokyo, Japan
9
10 [2]Division of Biological Chemistry, Department of Pharmaceutical Sciences, Showa
11 Medical University School of Pharmacy, Tokyo, Japan
12
13 [3]Division of Bioanalytical Chemistry, Department of Pharmaceutical Sciences, Showa
14 Medical University School of Pharmacy, Tokyo, Japan
15
16 [4]Division of Electron Microscopy, Showa Medical University School of Medicine, Tokyo,
17 Japan
18
19 [5]Department of Environmental Health Sciences, Hoshi University, Tokyo, Japan
20
21 [6]Division of Genome Safety Science, National Institute of Health Sciences, Kawasaki,
22 Japan
23
24

25 *Corresponding author

26 E-mail: fukuhara@pharm.showa-u.ac.jp

Abstract

Neutrophil extracellular traps (NETs) are chromatin-based structures released by activated neutrophils in response to pathogens or chemical stimuli, contributing to host defense but also implicated in autoimmune disease and inflammation. As NET formation gains attention as an endpoint in in vitro immunotoxicity screening, the lack of reproducible and scalable systems hampers its broader application. Here, we developed an in vitro assay using HL-60-derived neutrophil-like cells (dHL-60) differentiated with all-trans retinoic acid to evaluate NET-inducing activity in a standardized, non-animal model. Graphene oxide (GO), a model nanomaterial known to trigger NETs in primary neutrophils, induced concentration-dependent NET formation in dHL-60 cells, with maximal induction at intermediate doses and attenuation at higher concentrations, likely due to particle aggregation. NET formation was validated by extracellular DNA staining and scanning electron microscopy. GO also induced superoxide-mediated ROS production, as confirmed by electron spin resonance, consistent with canonical NETosis pathways. Furthermore, GO suppressed PMA-induced NET formation, suggesting a dose- and context-dependent immunomodulatory effect. Collectively, our results demonstrate that dHL-60 cells recapitulate key features of NETosis observed in primary neutrophils and provide a practical, reproducible model for investigating immune responses to nanomaterials. This system supports the development of non-animal approaches for assessing immunological effects of chemical substances under controlled in vitro conditions.

Introduction

The evaluation of chemical and pharmaceutical safety is a critical step in product development, traditionally relying on animal models to assess toxicity before clinical or commercial application. However, ethical concerns over animal welfare, coupled with increasing regulatory and societal pressure, have accelerated the pursuit of alternative testing strategies. In response, in vitro toxicity testing platforms are being actively developed, with a particular emphasis on generating systems that are mechanistically informative, reproducible, and biologically relevant. Among the major challenges in this field is the establishment of robust assays capable of detecting immune-related toxicities, especially those associated with neutrophil activation.

A promising endpoint in this context is the formation of neutrophil extracellular traps (NETs), web-like structures composed of decondensed chromatin and neutrophil-derived proteins that are released in response to microbial pathogens or sterile stimuli. [1] [2] [3] As a defense mechanism of the innate immune system, NETs contribute to microbial clearance and modulation of inflammation. However, aberrant or excessive NET formation has been increasingly associated with adverse health outcomes, including autoimmune diseases, thrombosis, and chronic inflammatory disorders. [4] [5] [6] Various exogenous agents, ranging from environmental pollutants [5] [7] and pharmaceuticals [8] [9] to engineered nanomaterials, [10] [11] have been shown to induce NETosis, implicating it as a key event in the manifestation of immunotoxic effects.

The molecular process underlying NET formation involves a cascade of intracellular signaling events. Upon activation by pathogens or chemical stimuli, neutrophils generate reactive oxygen species (ROS) via NADPH oxidase and initiate chromatin

decondensation through peptidylarginine deiminase 4 (PAD4)-mediated histone

citrullination. [12] This leads to nuclear membrane disintegration, plasma membrane

rupture, and the eventual extrusion of chromatin into the extracellular space. [13] While

NETs serve a protective function, they also reflect an early immune activation state,

rendering NET formation a potentially sensitive and mechanistically informative

endpoint for in vitro toxicity testing, particularly in the context of immunotoxicity. [3]

Despite this promise, conventional in vitro assays have largely focused on endpoints such

as genotoxicity or cytotoxicity, which capture later-stage or downstream effects of

toxicants. These methods often overlook early immune interactions, even though immune

cells such as neutrophils are among the first to encounter xenobiotics after systemic

exposure. Thus, assessing how neutrophils respond, especially through NET formation,

provides a more physiologically relevant perspective on chemical bioactivity. However,

primary human neutrophils are limited in vitro by their short lifespan (6–8 hours), donor-

to-donor variability, and restricted scalability, posing significant barriers to assay

standardization.

To address these limitations, HL-60 cells, a human promyelocytic leukemia cell line, can

be differentiated into neutrophil-like cells (dHL-60) using all-trans retinoic acid (AtRA).

[14] These cells exhibit essential neutrophil functions, including ROS generation and

NET formation, and offer several practical advantages, such as consistent availability,

reproducibility, and suitability for high-throughput applications. dHL-60 cells therefore

represent a viable surrogate for primary neutrophils in NET-based toxicological

assessments.

Nanomaterials such as graphene oxide (GO) have attracted considerable attention in

biomedical and industrial contexts owing to their unique physicochemical properties.

However, these same properties also complicate the prediction of biological outcomes.

GO has been reported to induce NET formation in primary neutrophils, [15] [9] raising

concerns regarding its potential immunomodulatory activity. In this study, we utilized

GO as a model nanomaterial to validate an in vitro NET formation assay using dHL-60

cells. We demonstrate that GO elicits NET formation via canonical ROS-dependent

pathways and modulates NET responses to known stimulants such as PMA.

By establishing that dHL-60 cells respond to GO with reproducible NET formation and

capture complex modulatory effects, this study lays the foundation for a scalable, non-

animal assay system targeting NETs as a mechanistic endpoint. Such systems may not

only facilitate the mechanistic dissection of nanomaterial-immune interactions, but also

support the development of predictive and ethically responsible platforms for

immunotoxicity screening and chemical safety evaluation.

Materials and methods

Chemicals

AtRA and phorbol 12-myristate 13-acetate (PMA) were purchased from Fujifilm Wako

Pure Chemical Co. (Osaka, Japan). SYTOX Green was purchased from Thermo Fisher

Scientific (Rockford, IL, USA). Furthermore, 5,5-dimethyl-1-pyrroline N-oxide (DMPO)

was purchased from Labotec (Tokyo, Japan). Graphene oxide (777676) and poly-L-lysine

solution (P4707) were purchased from Sigma–Aldrich (Saint Louis, MO, USA).

Graphene oxide (GO) was suspended in a serum-free Roswell Park Memorial Institute

(RPMI)-1640 (phenol red free) medium at the appropriate concentration and underwent

sonication at room temperature for 1 min immediately before each experimental

procedure to ensure proper dispersion.

NET induction by HL-60 cell-derived neutrophils

HL-60 cells were treated with AtRA to induce their differentiation into neutrophil-like

cells, as previously described [16]. Briefly, 2 μM AtRA were added to HL-60 cells (2.0

$\times 10^6$ cells/dish) cultured in an RPMI-1640 medium supplemented with 5% deactivated

(for 30 min at 56 ℃) fetal bovine serum, 50 U/mL penicillin, and 50 μg/mL streptomycin

in 10 cm dishes for 4 days. HL-60-derived neutrophil-like cells were collected and

washed once with serum-free RPMI-1640 (phenol red free) medium, seeded in a 24-well

poly-l-lysine-coated plate (5.0×10^5 cells/well), and cultured for 30 min. The cells were

treated with aliquots of PMA or GO and incubated at 37 ℃ under 5% CO_2 for 2 h.

Fluorometric quantitation of NET-DNA

The culture medium was treated with 1 U/mL MNase to degrade the DNA exposed

extracellularly upon NET formation in the culture dishes. The culture medium from each

dish was recovered and centrifuged at $1,800 \times g$ for 10 min. After mixing the supernatant

with 1 μM SYTOX Green, the fluorescence intensity was measured as previously

described [16]. Fluorescence (ex: 485 nm, em: 525 nm) was measured using Varioskan

Flash (Thermo Fisher Scientific).

Dynamic light scattering (DLS) analysis

DLS measurements were conducted using a Nanotrac Wave II (Microtrac Inc.) at room

temperature. Before measurement, the nanoparticle suspension was prepared at the

appropriate concentration in an RPMI medium and underwent sonication for 1 min at

room temperature. The suspension was allowed to stand for 10 min before DLS analysis.

The particle size distribution was analyzed based on the intensity-weighted average

diameter.

Electron spin resonance (ESR)-spin trapping ROS determination

HL-60-derived neutrophil-like cells were seeded in a 24-well plate coated with poly-L-

lysine (5.0×10^5 cells/well) and incubated for 30 min at 37 ℃ under 5% CO_2. The cells

were treated with DMPO at a final concentration of 400 mM, and PMA or GO was added.

After incubation for 10 or 20 min at 37 ℃ under 5% CO_2, the culture medium was

transferred to a flat quartz ESR cell and placed in a JEOL JES-3X ESR spectrometer. The

ESR spectra were recorded with manganous oxide as an external standard at room

temperature under the following conditions: microwave power, 10.0 mW; modulation

frequency, 100 kHz; modulation amplitude, 0.63 G; scanning field, $3,360 \pm 50$ G; receiver

gain, 6.3×100; response time, 0.1 s; sweep time, 2 min.

Scanning electron microscopy (SEM) analysis

HL-60 cells were differentiated into neutrophil-like cells through treatment with AtRA.

The differentiated cells were seeded onto chamber slides (Cat. No. 154526PK; Thermo

Fisher) and incubated for 30 min at 37 ℃ in a 5% CO_2 atmosphere to ensure adherence.

Subsequently, the cells were stimulated with 50 nM PMA or 100 μg/mL GO and

incubated for 2 h under the same conditions. Following stimulation, the cells were fixed

with 4% paraformaldehyde in phosphate-buffered saline for 15 min at room temperature.

After fixation, samples were processed for SEM imaging.

Statistical analysis

Data were expressed as mean ± SD. The comparisons were executed through one-way

analysis of variance (ANOVA) followed by Dunnett's post hoc test. Statistically

significant difference was evaluated at $p < 0.05$.

Results

GO induces NET formation in dHL-60 cells

Phorbol 12-myristate 13-acetate (PMA) is a well-established chemical stimulant known

to activate protein kinase C (PKC), thereby triggering neutrophil extracellular trap (NET)

formation in various neutrophil models, including HL-60 cells differentiated with AtRA.

[17] To explore whether graphene oxide (GO), previously reported to induce NETs in

primary neutrophils, exerts a similar effect on dHL-60, we evaluated NET formation in

response to GO stimulation.

GO was ultrasonically dispersed in culture medium to achieve uniform particle

distribution and incubated with dHL-60 cells for 2 hours. The extent of NET formation

was quantitatively assessed by SYTOX Green fluorescence, which selectively stains

extracellular DNA released during NETosis (Figure 1A). GO stimulation led to

significant DNA release in dHL-60 cells, confirming NET formation. Notably, treatment

with 25 µg/mL GO elicited a stronger NETotic response than 50 nM PMA, a potent

positive control. GO-induced NET formation exhibited a clear concentration-dependent

increase, peaking at 50 µg/mL. However, at 100 µg/mL, a paradoxical reduction in NET

formation was observed.

This non-linear response suggests that higher GO concentrations may result in particle

aggregation, thereby impairing effective interaction with the cell surface. Indeed, the

bioactivity of nanomaterials is often dictated by their dispersion state, and loss of colloidal

stability at high concentrations can reduce cellular uptake and membrane association.

These findings underscore the importance of GO concentration and dispersion state in

modulating NET-inducing capacity in vitro.

194 Fig 1. Fluorometric quantification of DNA release from HL-60-derived

195 neutrophils (A).

Graphene oxide (GO; 12.5−100 µg/mL) and phorbol 12-myristate 13-acetate

(PMA; 50 nM) were added to HL-60-derived neutrophils, and the mixture was

incubated for 2 h. (B) HL-60-derived neutrophils were stimulated with 50 nM PMA

for 10 min and treated with 0−100 µg/mL GO for an additional 2 h. The collected

culture medium was treated with micrococcal nuclease and stained with 1 µM

SYTOX Green, after which DNA released from the cells was quantified

fluorometrically. Data are presented as the mean ± SD of four independent

experiments. Asterisks indicate statistical significance. **$p < 0.005$, ***$p < 0.001$.

205 GO modulates PMA-induced NET formation

PMA activates NETosis by engaging PKC at the plasma membrane, and its effect can be

modulated by co-exposure to pro-oxidant or antioxidant compounds. Previous studies

have shown that oxidized low-density lipoprotein (oxLDL) enhances, while resveratrol

inhibits, PMA-induced NET formation. [16] [18] To determine whether GO influences

NET formation indirectly through modulation of known stimuli, we examined the effect

of co-treatment with GO and PMA in dHL-60 cells.

Stimulation with 50 nM PMA alone robustly induced NET formation. However, the

addition of GO during PMA stimulation significantly suppressed NET formation in a

concentration-dependent fashion (Figure 1B). At 100 µg/mL, GO exhibited the most

pronounced inhibitory effect. These results suggest that GO may antagonize PKC-driven

NETosis, potentially by disrupting redox-sensitive signaling pathways or membrane-

associated processes. Such dual functionality, inducing NETs independently, while

suppressing chemically triggered NETosis, highlights the complex, context-dependent

bioactivity of GO.

GO aggregation at high concentrations affects its biological activity

To further probe the concentration-dependent effects of GO on NET formation, we

characterized its dispersion profile in cell culture medium RPMI via DLS (Figure 2). GO

at 20 µg/mL exhibited a unimodal size distribution with a median hydrodynamic diameter

(D50) of 838 nm, indicating well-dispersed nanosheets. As the concentration increased,

a second peak appeared, reflecting the formation of larger particle aggregates. At 200

µg/mL, D50 increased markedly to 2518 nm, consistent with extensive aggregation.

This aggregation likely reduces GO's surface area available for cellular interaction and

may hinder its internalization or membrane binding. The loss of NET-inducing capacity

at 100 µg/mL may therefore be attributed to this physical transformation. These results

reinforce the concept that nanomaterial dispersion is a critical determinant of biological

activity and must be carefully monitored in concentration-response evaluations.

Fig 2. Dynamic light scattering (DLS) analysis of GO (20−200 µM) in RPMI.

GO induces ROS production in dHL-60 cells

NET formation is tightly linked to intracellular ROS production, primarily driven by the NADPH oxidase complex during the respiratory burst. [19] To investigate whether GO induces ROS in dHL-60 cells, we performed ESR spectroscopy with the spin-trap 5,5-dimethyl-1-pyrroline N-oxide (DMPO) (Figure 3). dHL-60 cells stimulated with PMA displayed a characteristic ESR signal for the DMPO-OH adduct, a spin-trap product of hydroxyl radicals, with a 1:2:2:1 quartet hyperfine pattern (a_N = 1.49 mT, a_H = 1.49 mT). The signal was attenuated by superoxide dismutase (SOD), confirming that hydroxyl radicals originated via superoxide-dependent pathways.

Similarly, GO stimulation yielded a strong DMPO-OH signal, supporting the conclusion that GO activates ROS production through a comparable mechanism. Additionally, a weaker triplet-of-triplets signal (1:1:2:1:2:1:2:1:1) characteristic of DMPO-H, the adduct of hydrogen radicals (aN = 1.63 mT, aH = 2.53 mT), was observed in GO-treated cells. While the biological implications of hydrogen radical formation remain unclear, this finding points to the distinct redox properties of GO and warrants further mechanistic investigation.

Fig 3. Electron spin resonance (ESR) spectra of 5,5-dimethyl-1-pyrroline N-oxide (DMPO)-free radical spin adducts obtained from solutions containing dHL-60 treated with PMA or GO.

 (A) 50 nM PMA + 613 mM DMPO, 10 min incubation. (B) Similar to that in (A), with the addition of 100 U/mL superoxide dismutase (SOD). (C) 100 µM GO +

613 mM DMPO, 10 min incubation. (D) Similar to that in (C); however, with 20

min incubation.

Morphological validation of GO-induced NET structures

To confirm the morphological characteristics of GO-induced NET formation, we

conducted SEM analysis of dHL-60 cells exposed to GO or PMA (Figure 4). In PMA-

treated cells, extensive chromatin decondensation and extracellular web-like structures

were observed, consistent with classical NETosis. GO-treated dHL-60 cells exhibited

comparable phenotypes, including membrane rupture and the release of reticulated

chromatin fibers.

These ultrastructural findings provide visual evidence that GO induces NET formation

via a process phenotypically indistinguishable from canonical NETosis. When considered

alongside DNA quantification and ROS measurements, these results establish that GO

elicits a bona fide NET response in dHL-60 cells, validating the use of this cell model for

NET-oriented toxicity screening.

Fig 4. Scanning electron microscopy (SEM) images showing the

characteristic appearance of NETs in dHL-60 cells treated with PMA or GO.

277 Discussion

The biological impact of chemical substances and engineered nanomaterials extends

beyond classical toxicological endpoints, increasingly encompassing immunological

responses such as inflammation, immune suppression, and aberrant activation. Among

these, the formation of NETs has emerged as a highly sensitive indicator of immune cell

activation and dysregulation. Although originally identified as a host defense mechanism

against microbial invasion, excessive or misregulated NET formation has been implicated

in diverse pathologies, including autoimmune diseases, thrombosis, and tissue damage.

[20] [6] These insights underscore the need to incorporate NET formation as a

biologically relevant endpoint in next-generation in vitro toxicology platforms. However,

the practical implementation of NET-based assays has been hampered by the technical

limitations of primary neutrophils, such as their short lifespan, donor-to-donor variability,

and restricted scalability. To circumvent these issues, we employed HL-60-derived

neutrophil-like cells (dHL-60), which acquire key neutrophilic functions upon

differentiation with AtRA. Our study demonstrates that dHL-60 cells robustly reproduce

canonical features of NETosis, including extracellular DNA release, reactive oxygen

species (ROS) generation, and characteristic chromatin morphology, in response to both

phorbol ester stimulation and nanomaterial exposure.

GO was used as a model nanomaterial due to its reported ability to induce NET formation

in primary neutrophils. Consistent with previous findings, GO elicited pronounced NET

formation in dHL-60 cells in a concentration-dependent manner. The peak response

observed at 50 μg/mL, followed by attenuation at higher concentrations, suggests that

GO's bioactivity is modulated by its dispersion state. DLS analysis revealed significant

aggregation at elevated GO concentrations, which likely reduced its effective cellular

interactions. This phenomenon highlights the critical role of nanomaterial

physicochemical properties, particularly particle size and surface area, in determining

biological activity. Similar behaviors have been reported for other nanomaterials, such as

silica [21] and titanium dioxide, [22] indicating that this is a generalizable concern in

nanotoxicology.

Importantly, our assay system also captured the modulatory effects of GO on chemically

induced NETosis. Co-treatment with PMA and GO resulted in a significant, dose-

dependent suppression of NET formation. This dual functionality, namely NET induction

in naïve cells and inhibition of PMA-triggered NETosis, suggests that GO can interact

with both basal and activated signaling pathways. While the underlying mechanisms

remain to be fully elucidated, possible explanations include interference with PKC

signaling, alteration of redox balance, or modulation of membrane-associated receptors.

These results underscore the utility of the dHL-60 model not only for screening NET

inducers, but also for identifying potential modulators or inhibitors of NETosis, which

may have therapeutic relevance.

The involvement of ROS in NET formation is well established, with NADPH oxidase-

mediated superoxide production serving as a critical upstream trigger. ESR spectroscopy

revealed that GO induces hydroxyl radical formation via superoxide-dependent

pathways, mirroring the ROS signature observed with PMA. Notably, GO also generated

hydrogen radical signals, a phenomenon not typically reported in PMA-induced NETosis.

Although the functional implications of hydrogen radical generation remain unclear, this

observation suggests that GO may activate distinct redox chemistries, potentially

contributing to its unique biological effects.

Morphological analyses via scanning electron microscopy provided further validation of

NET-like structures induced by GO, confirming the rupture of nuclear and plasma

membranes and the release of extracellular chromatin webs. Taken together with

fluorescence-based DNA detection and ROS measurements, these findings establish that

dHL-60 cells can serve as a robust and reproducible surrogate model for NET-based

toxicity assessment.

Beyond the scope of GO, our previous unpublished data indicate that other particulate

materials, such as silica nanoparticles, also induce NET formation in dHL-60 cells. These

observations support the broader applicability of this model system across diverse

compound classes. Given its reproducibility, scalability, and non-reliance on animal-

derived primary cells, the dHL-60 NET assay is particularly suited for high-throughput

screening and mechanistic investigations in immunotoxicology. Nevertheless, several

limitations should be acknowledged. The mechanistic equivalence between NETosis in

dHL-60 cells and primary neutrophils requires further validation, particularly with respect

to downstream immunological consequences such as cytokine release, macrophage

activation, or tissue remodeling. Additionally, while ESR spectroscopy offers valuable

insights into ROS dynamics, complementary assays, including mitochondrial function,

calcium flux, and transcriptional profiling, may enrich mechanistic interpretation. Finally,

as nanomaterials display complex behaviors in biological fluids, standardization of

dispersion protocols will be essential for assay reproducibility and inter-laboratory

comparability. Collectively, these results support the use of HL-60-derived neutrophil-

like cells as a robust and reproducible platform for dissecting NET-related cellular

responses to chemical substances and nanomaterials. The ability of this system to detect

both NET-inducing and NET-modulating activities, together with its compatibility with

high-throughput formats, underscores its potential utility in mechanism-driven, non-

animal immunotoxicity testing strategies.

Conclusion

This study establishes a standardized and mechanistically informative in vitro platform for evaluating NET formation using HL-60-derived neutrophil-like cells (dHL-60). By employing GO as a representative nanomaterial, we demonstrated that dHL-60 cells recapitulate hallmark features of NETosis, including extracellular chromatin release, ROS production, and characteristic morphological changes, traditionally observed in primary neutrophils. Notably, GO induced NET formation in a concentration-dependent manner, and also suppressed PMA-induced NETosis, revealing its dual modulatory capacity and highlighting the complexity of nanomaterial-immune cell interactions. Our findings underscore the utility of dHL-60 cells as a viable alternative to primary neutrophils for immunotoxicological studies. The system provides consistent performance without the variability associated with donor-derived cells, and is amenable to high-throughput screening, mechanistic dissection, and standardization. Furthermore, the correlation between GO aggregation state and its bioactivity emphasizes the importance of physicochemical characterization in nanotoxicology assay design.

While the current work focused on GO, the assay platform is readily applicable to a broad spectrum of environmental, pharmaceutical, and industrial compounds known or suspected to modulate neutrophil function. Future studies incorporating structurally and mechanistically diverse substances will help to refine the system's predictive value and expand its regulatory relevance. As regulatory frameworks shift toward mechanism-based and animal-free safety assessment strategies, the integration of NET-focused assays using standardized cell systems like dHL-60 holds promise for transforming the evaluation of immune-related toxicities across a wide array of chemical exposures.

References

1. Brinkmann V, Reichard U, Goosmann C, Fauler B, Uhlemann Y, Weiss DS, et al.
Neutrophil extracellular traps kill bacteria. Science. 2004;303(5663):1532-5. doi:
10.1126/science.1092385.
2. Papayannopoulos V. Neutrophil extracellular traps in immunity and disease. Nat
Rev Immunol. 2018;18(2):134-47. Epub 20171009. doi: 10.1038/nri.2017.105.
3. Wang H, Kim SJ, Lei Y, Wang S, Wang H, Huang H, et al. Neutrophil extracellular
traps in homeostasis and disease. Signal Transduct Target Ther. 2024;9(1):235.
Epub 20240920. doi: 10.1038/s41392-024-01933-x.
4. Yang H, Marion TN, Liu Y, Zhang L, Cao X, Hu H, et al. Nanomaterial Exposure
Induced Neutrophil Extracellular Traps: A New Target in Inflammation and Innate
Immunity. J Immunol Res. 2019;2019:3560180. Epub 20190228. doi:
10.1155/2019/3560180.
5. Valderrama A, Ortiz-Hernandez P, Agraz-Cibrian JM, Tabares-Guevara JH,
Gomez DM, Zambrano-Zaragoza JF, et al. Particulate matter (PM(10)) induces in
vitro activation of human neutrophils, and lung histopathological alterations in a
mouse model. Sci Rep. 2022;12(1):7581. Epub 20220509. doi: 10.1038/s41598-
022-11553-6.
6. Wigerblad G, Kaplan MJ. Neutrophil extracellular traps in systemic autoimmune
and autoinflammatory diseases. Nat Rev Immunol. 2023;23(5):274-88. Epub
20221018. doi: 10.1038/s41577-022-00787-0.
7. Zhu Y, Xu H, Wang T, Xie Y, Liu L, He X, et al. Pro-inflammation and pro-
atherosclerotic responses to short-term air pollution exposure associated with
alterations in sphingolipid ceramides and neutrophil extracellular traps. Environ
Pollut. 2023;335:122301. Epub 20230802. doi: 10.1016/j.envpol.2023.122301.
8. Warren K, Vu K, Shergill K, Watson B, Faris M. Rare complication of a commonly
used antihypertensive agent: A case of hydralazine-induced ANCA-associated
vasculitis presenting as rapidly progressive glomerulonephritis. Clin Case Rep.
2022;10(2):e05411. Epub 20220207. doi: 10.1002/ccr3.5411.
9. Huang S, Li S, Liu Y, Ghalandari B, Hao L, Huang C, et al. Encountering and
Wrestling: Neutrophils Recognize and Defensively Degrade Graphene Oxide. Adv
Healthc Mater. 2022;11(8):e2102439. Epub 20211210. doi:
10.1002/adhm.202102439.
10. Manda-Handzlik A, Bystrzycka W, Sieczkowska S, Demkow U, Ciepiela O.
Antibiotics Modulate the Ability of Neutrophils to Release Neutrophil Extracellular
Traps. Adv Exp Med Biol. 2017;944:47-52. doi: 10.1007/5584_2016_59.
11. Vaseruk A, Bila G, Bilyy R. Nanoparticles for stimulation of neutrophil
extracellular trap-mediated immunity. Eur J Immunol. 2024;54(4):e2350582. Epub
20240126. doi: 10.1002/eji.202350582.
12. Zhu D, Zhang Y, Wang S. Histone citrullination: a new target for tumors. Mol
Cancer. 2021;20(1):90. Epub 20210611. doi: 10.1186/s12943-021-01373-z.

13. Vorobjeva NV, Chernyak BV. NETosis: Molecular Mechanisms, Role in
Physiology and Pathology. Biochemistry (Mosc). 2020;85(10):1178-90. doi:
10.1134/S0006297920100065.
14. Tasseff R, Jensen HA, Congleton J, Dai D, Rogers KV, Sagar A, et al. An Effective
Model of the Retinoic Acid Induced HL-60 Differentiation Program. Sci Rep.
2017;7(1):14327. Epub 20171030. doi: 10.1038/s41598-017-14523-5.
15. Lu YJ, Wang YH, Sahu RS, Chen JP, Dash BS, Chung PJ, et al. Mechanism of
Nanoformulated Graphene Oxide-Mediated Human Neutrophil Activation. ACS
Appl Mater Interfaces. 2020;12(36):40141-52. Epub 20200826. doi:
10.1021/acsami.0c12490.
16. Obama T, Ohinata H, Takaki T, Iwamoto S, Sawada N, Aiuchi T, et al. Cooperative
Action of Oxidized Low-Density Lipoproteins and Neutrophils on Endothelial
Inflammatory Responses Through Neutrophil Extracellular Trap Formation. Front
Immunol. 2019;10:1899. Epub 20190809. doi: 10.3389/fimmu.2019.01899.
17. Kawakami T, He J, Morita H, Yokoyama K, Kaji H, Tanaka C, et al. Rab27a is
essential for the formation of neutrophil extracellular traps (NETs) in neutrophil-
like differentiated HL60 cells. PLoS One. 2014;9(1):e84704. Epub 20140103. doi:
10.1371/journal.pone.0084704.
18. Ohinata H, Phimarn W, Mizuno M, Obama T, Fukuhara K, Makiyama T, et al.
Suppressive effect of resveratrol, catechin and their conformationally constrained
analogs on neutrophil extracellular trap formation by HL-60-derived neutrophils. J
Clin Biochem Nutr. 2024;75(1):17-23. Epub 20240305. doi: 10.3164/jcbn.23-80.
19. Kirchner T, Moller S, Klinger M, Solbach W, Laskay T, Behnen M. The impact of
various reactive oxygen species on the formation of neutrophil extracellular traps.
Mediators Inflamm. 2012;2012:849136. Epub 20120126. doi:
10.1155/2012/849136.
20. Wirestam L, Arve S, Linge P, Bengtsson AA. Neutrophils-Important
Communicators in Systemic Lupus Erythematosus and Antiphospholipid
Syndrome. Front Immunol. 2019;10:2734. Epub 20191122. doi:
10.3389/fimmu.2019.02734.
21. Inoue M, Sakamoto K, Suzuki A, Nakai S, Ando A, Shiraki Y, et al. Size and
surface modification of silica nanoparticles affect the severity of lung toxicity by
modulating endosomal ROS generation in macrophages. Part Fibre Toxicol.
2021;18(1):21. Epub 20210617. doi: 10.1186/s12989-021-00415-0.
22. Murugadoss S, Mulhopt S, Diabate S, Ghosh M, Paur HR, Stapf D, et al.
Agglomeration State of Titanium-Dioxide (TiO(2)) Nanomaterials Influences the
Dose Deposition and Cytotoxic Responses in Human Bronchial Epithelial Cells at
the Air-Liquid Interface. Nanomaterials (Basel). 2021;11(12). Epub 20211127. doi:
10.3390/nano11123226.

Figure 1

Figure 2

Figure 3

PMA 25nM   GO 12.5µg/ml   GO 50µg/ml

Figure 4