Remodelled cholesteryl ester enriched lipid droplets fuel flavivirus morphogenesis

Adrianna Banducci-Karp[1], Sophie Brixton[1], Pranav N. M. Shah[2], Ming-Yuan Li[3], James Bancroft[4], Joey Riepsaame[1], Raman Dhaliwal[1], Charlotte E. Melia[1], Gathsaurie Neelika Malavige[5] and Sumana Sanyal[1]*

[1]Sir William Dunn School of Pathology, University of Oxford, South Parks Road, Oxford, UK

[2]Division of Structural Biology, University of Oxford, Centre for Human Genetics, Roosevelt Drive, Oxford, UK

[3]Department of Chemical Pathology and Li Ka Shing Institute of Health Sciences, Chinese University of Hong Kong, Hong Kong SAR, China

[4]Centre for Human Genetics, Nuffield Department of Medicine (NDM), University of Oxford, Oxford, UK

[5]Allergy, Immunology and Cell Biology Unit, Department of Immunology and Molecular Medicine, University of Sri Jayewardenepura, Nugegoda, Sri Lanka

Correspondence: sumana.sanyal@path.ox.ac.uk

Abstract

Flaviviruses such as dengue and Zika viruses extensively remodel host cell membranes to create specialised replication organelles, but the molecular mechanisms governing lipid metabolism during infection remain poorly understood.

Through systematic screens of fatty acyl transferase enzymes (MBOAT and zDHHC families) and complementary approaches including CRISPR/Cas9 gene deletions, pharmacological inhibition, proteomics, and photo-crosslinkable cholesterol analogues, we identified Sterol O-acyltransferases 1 and 2 (SOAT1/SOAT2) as critical host dependency factors for flavivirus infection. SOAT1/2 activities were upregulated early during infection, coinciding with increased LD formation, which underwent transition to liquid crystalline phases. Genetic deletion or pharmacological inhibition of either enzyme resulted in a dramatic ~100-fold reduction in viral production. Mechanistically, SOAT1/2 generate cholesteryl ester-enriched lipid droplets with fundamentally altered proteomes, enriched in fatty acid remodelling enzymes, Rab-GTPases, lipid transport proteins and sphingomyelinases. Photo-crosslinking experiments demonstrated direct interactions between LD-derived cholesterol and viral prM, capsid and NS1. SOAT1/2 deficiency resulted in defective, viral RNA-free replication organelles and complete absence of immature virions. Supporting the clinical relevance of viral lipid exploitation, analysis of dengue patients from a Sri Lankan cohort revealed that central obesity significantly increased the risk of severe dengue haemorrhagic fever, compared to lean patients.

This study establishes SOAT1/2 as essential host factors that enable flavivirus morphogenesis through specialised cholesteryl ester-enriched LDs, revealing a previously unrecognised virus-host interaction mechanism and identifying host lipid metabolism as a promising therapeutic target for combating flavivirus infections.

Flaviviruses pose significant global health threats, with members such as Dengue, West Nile, and Zika affecting millions annually[1]. These mosquito-borne pathogens cause diseases ranging from mild febrile illness to severe manifestations including haemorrhagic fever, encephalitis and congenital abnormalities[1]. Despite their clinical importance, critical gaps persist in our understanding of the molecular mechanisms governing flavivirus biogenesis and host cell interactions[1]. This knowledge deficit has significantly impeded the development of effective antiviral strategies.

A hallmark of flavivirus infection is the extensive remodelling of host cell membranes to create specialised replication organelles (ROs)[1–3]. These virus-engineered compartments serve as protected factories for viral genome replication and assembly, effectively shielding viral components from host immune surveillance. RO biogenesis necessitates substantial modifications in lipid metabolism, particularly involving the biosynthesis, trafficking and redistribution of lipid species to support membrane proliferation and curvature[4–8]. However, the specific host factors mediating these critical lipid modifications during flavivirus infection remain largely uncharacterised.

Central to cellular lipid homeostasis are lipid droplets (LDs), dynamic organelles consisting of a hydrophobic core of neutral lipids (primarily triacylglycerols and cholesteryl esters) surrounded by a phospholipid monolayer embedded with a complement of regulatory proteins[9]. Emerging evidence indicates that flaviviruses strategically manipulate both the formation and turnover of LDs during infection, suggesting a potential role for these organelles in the viral lifecycle[4,10–13]. However, the molecular mechanisms governing flaviviruses exploitation of LDs and the viral and host factors involved in this process, remain poorly understood.

We previously reported that dengue virus (DENV) infection is accompanied by an early accumulation of LDs (between 10-18 hours) followed by their substantial turnover of lipid droplets at later stages (between 24-48 hours). This controlled degradation occurs primarily via AUP1-Ube2g2 mediated lipophagy, a selective autophagic pathway targeting LDs[4,10]. While the functional impact of these temporal events in the LD lifecycle is not known,

they can serve several purposes: (i) controlled release of fatty acids and cholesterol from LDs may supply building blocks for replication organelle biogenesis; (ii) released fatty acids may be used for protein acylation as a means of hijacking host factors; (iii) regulated lipophagy may prevent excessive LD accumulation, which could otherwise trigger inflammation; (iv) released fatty acids could also fuel β-oxidation, generating the ATP to meet the heightened energy demands of virus replication[14].

In this study, we determined that LD alterations were accompanied by an increase in both lipid and protein fatty acylation in Zika virus (ZIKV) infected cells. To identify relevant acyltransferases, we conducted a systematic screen of fatty acyl transferase enzymes (MBOAT[15] and zDHHC families[16]) and identified Sterol O-acyltransferase 1 and 2 (SOAT1 and SOAT2) as critical host factors in ZIKV and DENV infection. Also known as Acyl-CoA:cholesterol acyltransferases (ACAT1 and ACAT2), these enzymes serve as key regulators of cellular cholesterol metabolism[17]. These are ER-resident integral membrane proteins that catalyse the formation of cholesteryl esters from long-chain fatty acyl-CoA and cholesterol[18]. This esterification process generates transport-competent forms of cholesterol that are important for cholesterol homeostasis and LD formation.

Our findings revealed that SOAT1 and SOAT2 activities are upregulated early during ZIKV infection, coinciding with increased LD formation. Using complementary approaches including genetic deletion through CRISPR/Cas9 and dose-dependent pharmacological inhibition, we established that SOAT1 and SOAT2 are required for ZIKV production in cellular infection models, specifically affecting the formation of replication organelles, and virion assembly. The importance of these enzymes lies in their ability to generate cholesteryl-ester enriched LDs with altered proteomes, enriched in fatty acid remodelling enzymes and lipid transport proteins, thereby creating specialised metabolic hubs that support membrane remodelling necessary for virus replication and assembly. Using a photo-crosslinkable cholesterol analogue we identified direct interactions between cholesterol and the viral structural proteins prM and capsid, as well as the non-structural protein NS1. Importantly, deletion or pharmacological inhibition of either SOAT1 or SOAT2 impaired this

remodelling process, resulting in defective, vRNA-free replication organelles and a complete loss of immature virions, with subsequent degradation of viral proteins. This study establishes SOAT1 and SOAT2 as key host dependency factors for flaviviruses and provides mechanistic insights into the role of LDs as metabolic hubs during infection. These findings are clinically relevant, as reflected in dengue patients with obesity displaying increased disease severity, likely due to enhanced virus exploitation of lipid metabolism. Our findings open new avenues for understanding flavivirus-host interactions and may guide the development of novel therapeutic strategies targeting host lipid metabolism to combat flavivirus infections.

Results

Flavivirus infection alters lipid droplet dynamics and increased fatty acylation

We previously characterised Aup1/Ube2g2-dependent lipid droplet (LD) turnover during dengue infection[4,10]. Building on these observations, we investigated the temporal dynamics of LDs during ZIKV infection. We infected wild-type (WT) and Aup1 knockout (Aup1[-/-]) Huh7 cells and monitored LD turnover through the infection time-course (**Figure 1A**). In WT cells, we observed an initial increase in LD abundance during the first 10 hours post-infection, followed by a gradual decrease over 24 hours and substantial reduction at 48 hours. This pattern is consistent with previous studies on DENV infection, which showed early LD accumulation followed by lipophagy-mediated turnover[10]. In contrast, Aup1[-/-] cells displayed impaired LD turnover, maintaining elevated LD numbers throughout the infection time course (**Figure 1A**, **1B**).

Given that LD turnover releases free fatty acids, we hypothesised that this phenomenon might enhance lipid and/or protein acylation during ZIKV infection (**Figure 1C**). Comprehensive acylation profiling in infected cells revealed significant increases in both lipid and protein acylation in WT cells (**Figure 1D, 1E**), which was inhibited in Aup1[-/-] cells, establishing a link between LD turnover and enhanced acylation during infection (**Figure 1D, 1E**).

To identify the specific acyltransferases mediating these modifications, we conducted parallel targeted screens of two enzyme families: the ZDHHC S-palmitoyltransferases (**Figure 1F**) and the membrane-bound O-acyltransferases (MBOATs) (**Figure 1G**) in ZIKV infected cells. The zDHHC family members primarily mediate protein S-palmitoylation and have established roles in SARS-CoV-2 infection and immunomodulation[19–22]. MBOATs catalyse fatty acid transfer to diverse substrates including lipids and proteins, with several members previously implicated in viral replication[23].

To determine the functional significance of these acyltransferases during ZIKV infection, we performed siRNA-mediated gene silencing against individual MBOAT and ZDHHC family members. Knockdown efficiency was confirmed by mRNA levels (**Figure S1A**

and S1B). While depletion of several ZDHHC proteins resulted in increased virus production (**Figure 1H**), depletion of SOAT1 and SOAT2 (involved in cholesteryl ester synthesis) resulted in a substantial loss of virus production, suggesting a potential role for these enzymes in the virus lifecycle (**Figure 1I**). In contrast, depletion of DGAT2 (involved in triacylglycerol synthesis) resulted in no detectable impact on viral titres (**Figure 1I**). These results suggest that SOAT1 and SOAT2, which specialise in cholesterol esterification, play important roles in supporting efficient ZIKV replication.

SOAT1 and SOAT2 activities are required for productive viral infection

SOAT1 and SOAT2 (alternatively known as ACAT1 and ACAT2) are ER-resident cholesterol acyltransferases that convert free cholesterol to esterified cholesterol to be deposited into LDs[24] (**Figure 2A**). To investigate their potential involvement in flavivirus infection, we first examined their RNA expression profiles and activity during virus infection (**Figure 2B**). SOAT2 mRNA levels were significantly upregulated following ZIKV infection, while increase in SOAT1 mRNA levels was less pronounced than SOAT2, and occurred only at 24 hours post infection. In contrast, DGAT1 and DGAT2 mRNA levels remained relatively stable, with DGAT2 mRNA levels significantly decreasing at 48 hours post infection. In line with increased SOAT1 and SOAT2 activity, there were more cholesterol esters formed in virus-infected cells (**Figure 2C**).

To assess the functional impact of SOAT1 and SOAT2 on Zika virus production, we generated Huh7 cells deleted in these genes via CRISPR/Cas9 gene editing (**Figure 2D**). Confocal microscopy revealed a marked decrease in LD numbers in the SOAT-deficient cells compared to wild-type cells (**Figure S2A**), confirming the expected disruption of cholesteryl ester synthesis and storage. Virus production from SOAT1$^{-/-}$ and SOAT2$^{-/-}$ cells was dramatically reduced by ~100-fold at 24 and 48 hours post infection, in Zika infected cells (**Figure 2E**). To validate these findings through an independent approach, we treated wild-type cells with Avasimibe, a specific inhibitor of SOAT1/2 enzymatic activity[25] (**Figure 2F**). When treated with increasing concentrations, viral titres in supernatants were similarly

reduced at 24h and 48h post infection for both ZIKV and DENV without causing cytotoxicity at these concentrations (**Figure 2F, Figure S2B**). This is in line with a previous report showing reduced virus production in a range of cells and tissues upon treatment with a SOAT1 inhibitor[26]. On the other hand, inhibiting DGAT1/2 did not have any measurable impact on virus production (**Figure 2F**).

SOAT1 and SOAT2 activities are required for morphogenesis of virions

To determine the mechanism underlying impaired virus production in SOAT1[-/-] and SOAT2[-/-] cells, we visualised dsRNA (marking virus replication organelles) and viral E-protein (indicating viral assembly sites) in infected wild-type, SOAT1[-/-] and SOAT2[-/-] Huh7 cells (**Figure 3A**). Immunofluorescence microscopy revealed that cells deleted in either SOAT1 or SOAT2 displayed substantially reduced dsRNA abundance and an even more profound decrease in E-protein clusters (**Figure 3A**). We confirmed this reduction through RT qPCR quantitation of viral RNA (**Figure 3B**) and immunoblotting for viral proteins (**Figure S3A**). Interestingly, in cells treated with the SOAT1/2 inhibitor, while the structural proteins and NS1 levels decreased substantially (**Figure S3A, S3B**), viral RNA levels did not significantly change, suggesting that these enzymes most likely did not affect NS5-dependent replication directly (**Figure S3C**). To test whether assembly of virus particles is also affected, we employed our previously established virus-like-particle system expressing the prME alone (**Figure 3C**) to measure levels of structural proteins[11] independent of replication. Similar to live infection, prM and E levels were found to be profoundly reduced in SOAT1[-/-] and SOAT2[-/-] cells (**Figure 3D**).

To determine whether SOAT1/2-deficiency impacted viral RNA stability and virion assembly on account of defective replication organelle and assembly sites, we used transmission electron microscopy on chemically fixed and resin embedded sections to assess ultrastructural defects (**Figure 3E**). In wild-type cells, expanded ER containing vesicle packets and immature virus particles near the vesicle packets were frequently observed in virus-infected cells. Interestingly, these replication organelles contained electron-dense material, likely viral RNA, positioned adjacent to assembled virions in the

wild-type infected cells (**Figure 3E**). In contrast, in SOAT1[-/-] or SOAT2[-/-] cells, the number of replication organelles were substantially lower, and those present were predominantly devoid of electron-dense material (highlighted in inset) (**Figure 3F**). Most significantly, no assembled virions could be detected in the SOAT1[-/-] or SOAT2[-/-] cells (**Figure 3F**). Similarly, SOAT-inhibitor treated cells displayed impaired biogenesis of replication organelle and viral progenies (**Figure S3D**). Interestingly, inhibitor treated cells also displayed defective mitochondrial morphology, indicative of potentially altered β-oxidation. Collectively, these data indicate that deficiency of SOAT1[-/-] or SOAT2[-/-] likely results in defective replication organelles and assembly sites, causing degradation of the viral RNA produced and abolishing assembly of viral progenies.

Remodelled cholesteryl ester enriched LDs supply lipids for ROs and virus assembly sites

SOAT1 and SOAT2 are enzymes that convert long chain fatty acids and free cholesterol into cholesteryl esters that are deposited into lipid droplets (LDs). SOAT1 and SOAT2 are induced upon ZIKV infection, increasing the amount of cholesteryl esters (**Figure 2B, 2C**). In line with this, TEM analyses revealed striking morphological differences between LDs in uninfected versus ZIKV-infected cells indicative of a change in composition (**Figure 4A**). Interestingly, LDs in virus-infected cells were also frequently ER-enwrapped. When visualised in FIB-milled samples, LDs in virus-infected cells frequently displayed a liquid crystalline phase (with 3.90 nm spacing of the lattice), a characteristic feature of cholesteryl ester-enriched LDs[27,28] (**Figure 4B, 4C**).

To directly investigate this compositional remodelling of LDs, we applied a chemical biology approach using a cholesterol alkyne analogue. This probe is metabolically incorporated into cells, esterified and deposited into LDs. Through click chemistry, we attached a fluorescent tag onto alkyne cholesterol to visualise its subcellular localisation in virus-infected wild-type versus SOAT-deficient cells (**Figure 4D**). TLC analysis confirmed that infected cells displayed significantly higher conversion of free cholesterol to cholesteryl

esters compared to mock infected cells, in a SOAT-dependent manner (**Figure 4D**). Temporal tracking of cholesterol localisation revealed a dynamic redistribution pattern during infection (**Figure 4E**). At early timepoints, cholesteryl esters predominantly accumulated within LDs. However, as infection progressed, this cholesterol redistributed to viral replication and assembly sites, where it colocalised with dsRNA (**Figure 4E**). In contrast, cholesterol accumulated in the plasma membrane upon SOAT1/2 inhibition and did not colocalise with replication organelles during infection (**Figure 4E**).

To identify the viral proteins that directly interact with cholesterol, we employed a bifunctional cholesterol analogue featuring both photo-crosslinking capability (via a diazirine group) and click chemistry reactivity (via an alkyne group). Mock and ZIKV infected cells were metabolically labelled with this cholesterol analogue and exposed to UV-light to activate the photo-crosslinking group, followed by Cu-catalysed click reaction with azide-biotin (**Figure 5A, B**). Streptavidin pulldown followed by silver staining confirmed capture of cross-linked proteins in both mock and virus-infected cells, but with distinct profiles (**Figure 5C**). Immunoblotting for viral proteins indicated prM, capsid and NS1 interacted with cholesterol (**Figure 5D**), as reported previously[29,30]. Importantly, these interactions were abolished in cells treated with the SOAT1/2 inhibitor, indicating that LD-derived cholesteryl esters – rather than free cholesterol – represent the primary source of cholesterol for viral protein interactions (**Figure 5D**).

To test whether adding exogenous lipids could rescue viral titres in SOAT1/2-inhibited cells, we supplemented either free cholesterol or fatty acids in the form of oleic acid in time-of-addition experiments (**Figure 5E, 5F**). When oleic acid was added to SOAT1/2-inhibited cells, virus production was rescued compared to DMSO-treated cells. In contrast, addition of free cholesterol – either at the same time of inhibitor addition, or 24 hours after inhibition addition – exacerbated the difference in viral titres between DMSO and SOAT-inhibited conditions regardless of the time point (**Figure 5E, 5F**). This indicates that the specific remodelling of LDs and access from them during infection is crucial for the ER-

reorganisation required for productive virus morphogenesis, rather than free cholesterol, which can be detrimental to virus production.

CE-rich lipid droplets are enriched in fatty acid remodelling and lipid transport proteins

To determine the molecular mechanisms by which cholesteryl ester rich (CE)-LDs support membrane remodelling during flavivirus infection, we conducted proteomic analyses of isolated LDs from mock and virus-infected cells using established protocols (**Figure 6A**)[31,32]. The LD proteome from virus-infected cells was profoundly altered, showing enrichment of fatty acid remodelling enzymes, Rab-GTPases, and lipid transport proteins (LTPs) including the class of bridge-like LTPs, and sphingomyelinases (**Figure 6B**). These alterations collectively suggest a virus-induced remodelling of LDs at both the lipidome and proteome levels, transforming them into metabolically active hubs optimised for lipid mobilisation and delivery.

To further characterise this remodelling process, we examined key host proteins that participate in lipid mobilisation and membrane remodelling pathways. Remarkably, genetic deletion of either SOAT1 or SOAT2 resulted in complete loss of expression of specific proteins (data not shown) such as TMEM41B (a lipid scramblase previously implicated in flavivirus replication)[33,34]. This loss was recapitulated in SOAT1/2-inhibited cells (**Figure 6C**). These findings suggest a pathway in which SOAT1/2 enzymatic activity generates specialised CE-LDs with modified proteomes that include critical lipid transport machinery, such as LTPs and the scramblase TMEM41B. This remodelling most likely occurs at ER-LD contact sites, where regulated hydrolysis and lipid remodelling generate the lipid species necessary for virus morphogenesis.

Collectively, our study demonstrates that flaviviruses, particularly DENV and ZIKV induce extensive reprogramming of LDs that promotes virus production. These findings have important implications for understanding severe disease manifestations such as liver damage in individuals with obesity and related metabolic disorders[35]. To investigate the

clinical relevance of our LD remodelling, we analysed dengue infection outcomes in a Sri Lankan cohort of 561 patients stratified by central obesity status, defined as waist circumference ≥80 cm in women or ≥90 cm in men[36]. Analysis of disease severity distribution revealed a clear association between central obesity and severe dengue manifestations (**Figure 6D**). While both lean and obese patient groups showed predominantly dengue fever (DF) cases, obese patients demonstrated a markedly higher proportion of dengue haemorrhagic fever (DHF) compared to lean patients. This difference in disease severity distribution was statistically significant (p=0.0407, Fisher's exact test). Quantitative analysis of DHF risk showed that patients with central obesity had a substantially elevated risk of developing severe disease (**Figure 6E**). Central obesity was associated with a 33.3% risk of DHF compared to 25.4% in patients without central obesity, representing an absolute risk increase of 7.9 percentage points and a relative risk of 1.22 (95% CI: 1.01-1.46).

These clinical findings support our mechanistic discoveries regarding SOAT-mediated LD remodelling. The increased severe dengue susceptibility in obese patients is consistent with our demonstration that flaviviruses exploit cholesteryl ester-enriched lipid droplets for morphogenesis. Given that obesity is characterised by altered lipid metabolism and increased LD abundance, obese patients may harbour cellular environments that are more conducive to viral replication. Our data suggest BMI could serve as a valuable clinical indicator for identifying patients at elevated risk for severe dengue manifestations.

Discussion

This study establishes cholesteryl ester synthesis as a critical host dependency factor for flavivirus infection and provides mechanistic insights into how viruses exploit LD metabolism to support their replication and assembly. Our findings demonstrate that SOAT1/2-mediated cholesterol esterification generates specialised LDs with liquid crystalline phases that serve as essential metabolic hubs for viral morphogenesis, advancing our understanding of flavivirus-host interactions.

The dramatic ~100-fold reduction in viral production upon SOAT1/2 deficiency, coupled with the complete absence of assembled virions, underscores the requirement for cholesterol esterification in flavivirus morphogenesis. Importantly, our findings reveal specificity for cholesterol esterification over triacylglycerol synthesis, suggesting that flaviviruses have evolved to preferentially exploit cholesteryl ester-enriched LDs rather than conventional triacylglycerol-rich LDs.

Our proteomic analyses reveal that flavivirus infection transforms LDs into specialised organelles with fundamentally altered composition and function. The enrichment of fatty acid remodelling enzymes, Rab-GTPases, sphingomyelinases, and lipid transport proteins, establishes these remodelled LDs as metabolically active hubs rather than passive lipid storage depots. The recruitment of bridge-like lipid transfer proteins, and correlation with TMEM41B expression provides a potential molecular mechanism for directional lipid transfer from CE-LDs to viral replication and assembly sites. This finding links previous observations of TMEM41B's requirement for flavivirus replication with the broader context of virus-induced lipid mobilisation[33].

The direct interactions between cholesterol and viral proteins prM, capsid, and NS1, which are specifically dependent on LD-derived cholesteryl esters rather than free cholesterol, suggest that the cellular source of cholesterol is a critical determinant of its functional role in viral morphogenesis. The failure of exogenous free cholesterol supplementation to rescue viral production in SOAT-inhibited cells, and its apparent detrimental effects, reinforces the specificity of this dependency. The interaction of cholesterol with both structural proteins (prM, capsid) and the secreted glycoprotein NS1 suggests multiple roles for cholesterol throughout the viral lifecycle. Our work also highlights the importance of organelle contact sites, particularly ER-LD contacts, as potential platforms for virus-host interactions and therapeutic targeting. The temporal coordination between LD accumulation, turnover, and viral replication phases may represent a programmed sequence optimised to provide lipid building blocks precisely when needed for replication organelle biogenesis and virion assembly.

The clinical relevance of our mechanistic findings is supported by analysis of dengue patients from a Sri Lankan cohort, where adiposity showed significantly higher rates of severe dengue haemorrhagic fever compared to lean patients. This observation aligns with our demonstration that flaviviruses exploit cholesteryl ester-enriched LD, as obesity involves altered lipid metabolism and increased LD abundance that may provide enhanced cellular environments supporting viral replication.

Our findings contribute to the growing recognition that viruses extensively reprogramme host metabolism to support their replication[12–14]. Future studies will examine whether this dependency extends to other flaviviruses and related RNA viruses, and whether similar mechanisms operate in physiologically relevant primary cell types. The development of more potent and selective SOAT inhibitors will be essential for advancing therapeutic applications targeting host lipid metabolism to combat flavivirus infections.

Acknowledgements

The Sanyal lab is supported by the Wellcome Trust (grants 220776/Z/20/Z and 223107/Z/21/Z to SS), BBSRC (BB/Y000307/1 to SS), and MRC studentship (AB-K).

References

1. van Leur, S.W., Heunis, T., Munnur, D., and Sanyal, S. (2021). Pathogenesis and virulence of flavivirus infections. Virulence *12*, 2814–2838. https://doi.org/10.1080/21505594.2021.1996059.

2. Paul, D., and Bartenschlager, R. (2015). Flaviviridae Replication Organelles: Oh, What a Tangled Web We Weave. Annu. Rev. Virol. *2*, 289–310. https://doi.org/10.1146/annurev-virology-100114-055007.

3. Stancheva, V.G., and Sanyal, S. (2024). Positive-strand RNA virus replication organelles at a glance. Journal of Cell Science *137*, jcs262164. https://doi.org/10.1242/jcs.262164.

4. Lan, Y., Van Leur, S.W., Fernando, J.A., Wong, H.H., Kampmann, M., Siu, L., Zhang, J., Li, M., Nicholls, J.M., and Sanyal, S. (2023). Viral subversion of selective autophagy is critical for biogenesis of virus replication organelles. Nat Commun *14*, 2698. https://doi.org/10.1038/s41467-023-38377-w.

5. Pombo, J.P., and Sanyal, S. (2018). Perturbation of Intracellular Cholesterol and Fatty Acid Homeostasis During Flavivirus Infections. Front Immunol *9*, 1276. https://doi.org/10.3389/fimmu.2018.01276.

6. Zhang, J., Lan, Y., and Sanyal, S. (2017). Modulation of Lipid Droplet Metabolism—A Potential Target for Therapeutic Intervention in Flaviviridae Infections. Front. Microbiol. *8*, 2286. https://doi.org/10.3389/fmicb.2017.02286.

7. Mackenzie, J.M., Khromykh, A.A., and Parton, R.G. (2007). Cholesterol Manipulation by West Nile Virus Perturbs the Cellular Immune Response. Cell Host & Microbe *2*, 229–239. https://doi.org/10.1016/j.chom.2007.09.003.

8. Wong, H.H., Crudgington, D.R.K., Siu, L., and Sanyal, S. (2024). Flaviviruses induce ER-specific remodelling of protein synthesis. PLOS Pathogens *20*, e1012766. https://doi.org/10.1371/journal.ppat.1012766.

9. Bosch, M., and Pol, A. (2022). Eukaryotic lipid droplets: metabolic hubs, and immune first responders. Trends in Endocrinology & Metabolism *33*, 218–229. https://doi.org/10.1016/j.tem.2021.12.006.

10. Zhang, J., Lan, Y., Li, M.Y., Lamers, M.M., Fusade-Boyer, M., Klemm, E., Thiele, C., Ashour, J., and Sanyal, S. (2018). Flaviviruses Exploit the Lipid Droplet Protein AUP1 to Trigger Lipophagy and Drive Virus Production. Cell Host Microbe *23*, 819-831.e5. https://doi.org/10.1016/j.chom.2018.05.005.

11. Li, M.Y., Naik, T.S., Siu, L.Y.L., Acuto, O., Spooner, E., Wang, P., Yang, X., Lin, Y., Bruzzone, R., Ashour, J., et al. (2020). Lyn kinase regulates egress of flaviviruses in autophagosome-derived organelles. Nat Commun *11*, 5189. https://doi.org/10.1038/s41467-020-19028-w.

12. Chen, Q., Gouilly, J., Ferrat, Y.J., Espino, A., Glaziou, Q., Cartron, G., El Costa, H., Al-Daccak, R., and Jabrane-Ferrat, N. (2020). Metabolic reprogramming by Zika virus provokes inflammation in human placenta. Nat Commun *11*, 2967. https://doi.org/10.1038/s41467-020-16754-z.

13. Dias, S.S.G., Cunha-Fernandes, T., Souza-Moreira, L., Soares, V.C., Lima, G.B., Azevedo-Quintanilha, I.G., Santos, J., Pereira-Dutra, F., Freitas, C., Reis, P.A., et al.

(2023). Metabolic reprogramming and lipid droplets are involved in Zika virus replication in neural cells. J Neuroinflammation *20*, 61. https://doi.org/10.1186/s12974-023-02736-7.

14. Heaton, N.S., and Randall, G. (2010). Dengue Virus-Induced Autophagy Regulates Lipid Metabolism. Cell Host & Microbe *8*, 422–432. https://doi.org/10.1016/j.chom.2010.10.006.

15. Coupland, C.E., Ansell, T.B., Sansom, M.S.P., and Siebold, C. (2023). Rocking the MBOAT: Structural insights into the membrane bound *O*-acyltransferase family. Current Opinion in Structural Biology *80*, 102589. https://doi.org/10.1016/j.sbi.2023.102589.

16. Lemonidis, K., Werno, M.W., Greaves, J., Diez-Ardanuy, C., Sanchez-Perez, M.C., Salaun, C., Thomson, D.M., and Chamberlain, L.H. (2015). The zDHHC family of S-acyltransferases. Biochemical Society Transactions *43*, 217–221. https://doi.org/10.1042/BST20140270.

17. Qian, H., Zhao, X., Yan, R., Yao, X., Gao, S., Sun, X., Du, X., Yang, H., Wong, C.C.L., and Yan, N. (2020). Structural basis for catalysis and substrate specificity of human ACAT1. Nature *581*, 333–338. https://doi.org/10.1038/s41586-020-2290-0.

18. Wang, Y.-J., Bian, Y., Luo, J., Lu, M., Xiong, Y., Guo, S.-Y., Yin, H.-Y., Lin, X., Li, Q., Chang, C.C.Y., et al. (2017). Cholesterol and fatty acids regulate cysteine ubiquitylation of ACAT2 through competitive oxidation. Nat Cell Biol *19*, 808–819. https://doi.org/10.1038/ncb3551.

19. Wang, Z., Qiu, M., Ji, Y., Chai, K., Liu, C., Xu, F., Guo, F., Tan, J., Liu, R., and Qiao, W. (2024). Palmitoylation of SARS-CoV-2 Envelope protein is central to virus particle formation. Journal of Virology *98*, e01072-24. https://doi.org/10.1128/jvi.01072-24.

20. Mesquita, F.S., Abrami, L., Sergeeva, O., Turelli, P., Qing, E., Kunz, B., Raclot, C., Paz Montoya, J., Abriata, L.A., Gallagher, T., et al. (2021). S-acylation controls SARS-CoV-2 membrane lipid organization and enhances infectivity. Developmental Cell *56*, 2790-2807.e8. https://doi.org/10.1016/j.devcel.2021.09.016.

21. Sheng, J., You, X., Nie, D., Fu, Y., Ling, Q., Yang, X., Chen, Y., Ma, L., and Hu, S. (2025). ZDHHC2 promoted antimycobacterial responses by selective autophagic degradation of B-RAF and C-RAF in macrophages. Sci. Adv. *11*, eadq7706. https://doi.org/10.1126/sciadv.adq7706.

22. Yi, L., and Zheng, C. (2021). The emerging roles of ZDHHCs-mediated protein palmitoylation in the antiviral innate immune responses. Critical Reviews in Microbiology *47*, 34–43. https://doi.org/10.1080/1040841X.2020.1835821.

23. Thabet, K., Asimakopoulos, A., Shojaei, M., Romero-Gomez, M., Mangia, A., Irving, W.L., Berg, T., Dore, G.J., Grønbæk, H., Sheridan, D., et al. (2016). MBOAT7 rs641738 increases risk of liver inflammation and transition to fibrosis in chronic hepatitis C. Nat Commun *7*, 12757. https://doi.org/10.1038/ncomms12757.

24. Bairos, J.A., Njoku, U., Zafar, M., Akl, M.G., Li, L., Parlakgul, G., Arruda, A.P., and Widenmaier, S.B. (2024). Sterol O-acyltransferase (SOAT/ACAT) activity is required to form cholesterol crystals in hepatocyte lipid droplets. Biochimica et Biophysica Acta (BBA) - Molecular and Cell Biology of Lipids *1869*, 159512. https://doi.org/10.1016/j.bbalip.2024.159512.

25. Llaverías, G., Laguna, J.C., and Alegret, M. (2003). Pharmacology of the AC AT Inhibitor Avasimibe (CI-1011). Cardiovascular Drug Reviews *21*, 33–50. https://doi.org/10.1111/j.1527-3466.2003.tb00104.x.

26. Schöbel, A., Pinho Dos Reis, V., Burkhard, R., Hehner, J., Schneider, L., Schauflinger, M., Vieyres, G., and Herker, E. (2024). Inhibition of sterol O-acyltransferase 1 blocks Zika virus infection in cell lines and cerebral organoids. Commun Biol *7*, 1089. https://doi.org/10.1038/s42003-024-06776-4.

27. Rogers, S., Gui, L., Kovalenko, A., Zoni, V., Carpentier, M., Ramji, K., Ben Mbarek, K., Bacle, A., Fuchs, P., Campomanes, P., et al. (2022). Triglyceride lipolysis triggers liquid crystalline phases in lipid droplets and alters the LD proteome. Journal of Cell Biology *221*, e202205053. https://doi.org/10.1083/jcb.202205053.

28. Mahamid, J., Tegunov, D., Maiser, A., Arnold, J., Leonhardt, H., Plitzko, J.M., and Baumeister, W. (2019). Liquid-crystalline phase transitions in lipid droplets are related to cellular states and specific organelle association. Proceedings of the National Academy of Sciences *116*, 16866–16871. https://doi.org/10.1073/pnas.1903642116.

29. Goellner, S., Enkavi, G., Prasad, V., Denolly, S., Eu, S., Mizzon, G., Witte, L., Kulig, W., Uckeley, Z.M., Lavacca, T.M., et al. (2023). Zika virus prM protein contains cholesterol binding motifs required for virus entry and assembly. Nat Commun *14*, 7344. https://doi.org/10.1038/s41467-023-42985-x.

30. Gutsche, I., Coulibaly, F., Voss, J.E., Salmon, J., d'Alayer, J., Ermonval, M., Larquet, E., Charneau, P., Krey, T., Mégret, F., et al. (2011). Secreted dengue virus nonstructural protein NS1 is an atypical barrel-shaped high-density lipoprotein. Proc. Natl. Acad. Sci. U.S.A. *108*, 8003–8008. https://doi.org/10.1073/pnas.1017338108.

31. Jahan, A.S., Biquand, E., Muñoz-Moreno, R., Le Quang, A., Mok, C.K.-P., Wong, H.H., Teo, Q.W., Valkenburg, S.A., Chin, A.W.H., Man Poon, L.L., et al. (2020). OTUB1 Is a Key Regulator of RIG-I-Dependent Immune Signaling and Is Targeted for Proteasomal Degradation by Influenza A NS1. Cell Rep *30*, 1570-1584.e6. https://doi.org/10.1016/j.celrep.2020.01.015.

32. Teo, Q.W., Wong, H.H., Heunis, T., Stancheva, V., Hachim, A., Lv, H., Siu, L., Ho, J., Lan, Y., Mok, C.K.P., et al. (2023). Usp25-Erlin1/2 activity limits cholesterol flux to restrict virus infection. Developmental Cell *58*, 2495-2509.e6. https://doi.org/10.1016/j.devcel.2023.08.013.

33. Hoffmann, H.-H., Schneider, W.M., Rozen-Gagnon, K., Miles, L.A., Schuster, F., Razooky, B., Jacobson, E., Wu, X., Yi, S., Rudin, C.M., et al. (2021). TMEM41B Is a Pan-flavivirus Host Factor. Cell *184*, 133-148.e20. https://doi.org/10.1016/j.cell.2020.12.005.

34. Huang, D., Xu, B., Liu, L., Wu, L., Zhu, Y., Ghanbarpour, A., Wang, Y., Chen, F.-J., Lyu, J., Hu, Y., et al. (2021). TMEM41B acts as an ER scramblase required for lipoprotein biogenesis and lipid homeostasis. Cell Metabolism *33*, 1655-1670.e8. https://doi.org/10.1016/j.cmet.2021.05.006.

35. Kuruppu, H., Karunananda, M., Jeewandara, C., Gomes, L., Dissanayake, D.M.C.B., Ranatunga, C., Chathurangika, P.H., Senatilleke, N., Warnakulasuriya, N., Wickramanayake, R.H., et al. (2025). Oxidative stress induced liver damage in dengue is exacerbated in those with obesity. Preprint at medRxiv, https://doi.org/10.1101/2025.03.18.25324170.

36. Gunasinghe, D.N., Jayamali, J., Chathurangika, P.H., Nimasha, T., Perera, L., Amarasekara, U., Wickramanayake, R.H., Panditharatne, A., Malavige, G.N., and Jeewandara, C. (2025). Adiposity and lean muscle indices that associate with cardiometabolic risk factors and inflammatory mediators in a South Asian population. Preprint at medRxiv, https://doi.org/10.1101/2025.04.13.25325724.

37. Munnur, D., Teo, Q., Eggermont, D., Lee, H.H.Y., Thery, F., Ho, J., van Leur, S.W., Ng, W.W.S., Siu, L.Y.L., Beling, A., et al. (2021). Altered ISGylation drives aberrant macrophage-dependent immune responses during SARS-CoV-2 infection. Nat Immunol *22*, 1416–1427. https://doi.org/10.1038/s41590-021-01035-8.

38. Stemmer, M., Thumberger, T., Del Sol Keyer, M., Wittbrodt, J., and Mateo, J.L. (2015). CCTop: An Intuitive, Flexible and Reliable CRISPR/Cas9 Target Prediction Tool. PLoS ONE *10*, e0124633. https://doi.org/10.1371/journal.pone.0124633.

39. Eisenstein, F., Yanagisawa, H., Kashihara, H., Kikkawa, M., Tsukita, S., and Danev, R. (2023). Parallel cryo electron tomography on in situ lamellae. Nat Methods *20*, 131–138. https://doi.org/10.1038/s41592-022-01690-1.

40. Mastronarde, D.N. (2005). Automated electron microscope tomography using robust prediction of specimen movements. Journal of Structural Biology *152*, 36–51. https://doi.org/10.1016/j.jsb.2005.07.007.

41. Hagen, W.J.H., Wan, W., and Briggs, J.A.G. (2017). Implementation of a cryo-electron tomography tilt-scheme optimized for high resolution subtomogram averaging. Journal of Structural Biology *197*, 191–198. https://doi.org/10.1016/j.jsb.2016.06.007.

42. Guo, H., Franken, E., Deng, Y., Benlekbir, S., Singla Lezcano, G., Janssen, B., Yu, L., Ripstein, Z.A., Tan, Y.Z., and Rubinstein, J.L. (2020). Electron-event representation data enable efficient cryoEM file storage with full preservation of spatial and temporal resolution. IUCrJ *7*, 860–869. https://doi.org/10.1107/S205225252000929X.

43. Tegunov, D., and Cramer, P. (2019). Real-time cryo-electron microscopy data preprocessing with Warp. Nat Methods *16*, 1146–1152. https://doi.org/10.1038/s41592-019-0580-y.

44. Zheng, S., Wolff, G., Greenan, G., Chen, Z., Faas, F.G.A., Bárcena, M., Koster, A.J., Cheng, Y., and Agard, D.A. (2022). AreTomo: An integrated software package for automated marker-free, motion-corrected cryo-electron tomographic alignment and reconstruction. Journal of Structural Biology: X *6*, 100068. https://doi.org/10.1016/j.yjsbx.2022.100068.

STAR Methods

Cell Culture and Viral Infections

Huh7 cells were maintained in Dulbecco's Modified Eagle Medium (DMEM) high glucose (Sigma-Aldrich, D6546) supplemented with 10% foetal bovine serum (FBS) at 37°C with 5% CO_2. Zika virus (ZIKV, strains MR766, NC-14-5132) and Dengue virus (DENV, strain 16681) were propagated in Vero or C6/36 mosquito cells and titrated by plaque assay on Vero cells as previously described[4,37]. The Vero E6 cell line was maintained in DMEM high glucose with 1% FBS. All cell lines were maintained at 37°C with 5% CO_2.

For infection experiments, Huh7 cells were seeded at a density of $1x10^5$ cells per well in a 12 well plate. The following day, Huh7 cells were washed 1x with PBS and infected with ZIKV MR766 at a multiplicity of infection (MOI) of 2 in DMEM containing no FBS. 2 hours later, cells were washed 2x with PBS and replaced with DMEM containing 10% FBS. 24 and 48 hours post infection, the supernatant was collected and stored at -70°C for plaque assays. Cells were washed 1x with PBS before collecting cells for western blotting and qRT-PCR. For immunofluorescence, cells were washed 2x with PBS before fixation.

Generation of Knockout Cell Lines

Guide RNAs (gRNAs) were designed using CCtop[38]. CRISPR RNAs (crRNAs) and trans-activating CRISPR-RNAs (tracrRNAs) were annealed by heating to 95°C followed by slowly cooling to room temperature to form crRNA/tracrRNA duplexes. Exonic and intronic targeting duplexes were combined in a 1:1 ratio and complexed with HiFi Cas9 Nuclease V3 (Integrated DNA Technologies, 1081060) at a final concentration of 44 µM crRNA/tracrRNA hybrids and 36 µM Cas9 to generate ribonucleoprotein (RNP) complexes. 1 µl of ribonuclease proteins was mixed with Huh7 (200,000 cells total) and electroporated using the Neon Transfection System (MPK10096; HiTrans 1400V 20 ms width, one pulse). Electroporated cells were cultured to establish CRISPR-edited cell pools, and genomic DNA

was subsequently extracted using the DNeasy kit (Qiagen, 69504) for genotyping analysis. Pooled knockouts were single cell sorted using flow cytometry into 96 well plates containing DMEM with 20% FBS.

siRNA Transfection

Small interfering RNAs (siRNAs) targeting MBOAT and ZDHHC family members were obtained from Dharmacon (ON-TARGETplus SMARTpool). Huh7 cells were transfected with 25 nM siRNA using Lipofectamine RNAiMAX (Thermo Fisher Scientific) according to manufacturer's protocol. Scrambled siRNA was used as negative control. Knockdown efficiency was verified by qRT-PCR 48 hours post-transfection.

Chemical Inhibitors and Treatments

Inhibitors were added 2 hours after Zika virus infection in complete DMEM (+10% FBS). To inhibit SOAT1/2, avasimibe (Cayman Chemical, 18129) was used at a final concentration of 20 µM. DGAT1 and DGAT2 were inhibited with 80 µM PF-04620110 (Cayman Chemical, 16425) and 40 µM PF-06424439 (Cayman Chemical, 17680), respectively.

Oleic acid (Sigma-Aldrich, O1008) or cholesterol (Sigma-Aldrich, C4951) was added 2 hours or 24 hours after Zika virus infection in combination with DMSO or SOATi in DMEM + 10% FBS. A final concentration of 500 µM oleic acid in 0.6% bovine serum albumin (BSA) was used (stock 10mM oleic acid in 12% BSA). Cholesterol (Sigma-Aldrich, C4951) was used at a final concentration of 0.1mM.

Quantitative RT-PCR

Huh7 cells were washed 1x with PBS and then lysed with 350 µl RLT buffer (Qiagen, 74104) + 1% beta-mercaptoethanol for 30 min before storing at -70°C. RNA was isolated using the RNeasy kit (Qiagen, 74104) and subsequently used for qRT-PCR using the One-Step SYBR Green PrimeScript RT-PCR kit (Takara, RR066B).

Table 1 Primer sequences for qRT-PCR

Target	Primer sequences	Source
SOAT1	Fp: 5'-CAATACAATGGTGGGTGAAGAGAAG-3' Rp: 5'-AAAATCTTTCCTTGTTCTGGAGGTG-3'	https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4500072/
SOAT2	Fp: 5'-CAATGGACCCGACACATGGA-3' Rp: 5'-CAGGGACTTGCGGATGATGA-3'	This study
DGAT1	Fp: 5'-AACGCAGCCGTTGTCCTTGA-3' Rp: 5'-ACCTCCGGGCCCTAGACAA-3'	This study
DGAT2	Fp: 5'-ACGGCCTTACCTGGCTACAC-3' Rp: 5'-CTATGGTGTCCCGGCTGACA-3'	This study
ZIKV MR766 E	Fp: 5'-TTGGGTTGTGTACGGAACCTG-3' Rp: 5'-GTGCTTTGTGTATTCTCTTGA-3'	Reference[4]
Beta-actin	Fp: 5'-CACCATTGGCAATGAGCGGTTC-3' Rp: 5'-AGGTCTTTGCGGATGTCCACGT-3'	Reference[32]

Viral Titration Assays

Supernatant from ZIKV infections were thawed on ice and 10-fold diluted from 10^{-1} to 10^{-4} in DMEM. 100 µl of each dilution was added to a 24-well plate in triplicate, and 2.5 x 10^5 Vero cells in 500 µl DMEM + 1% FBS were added to each well. Cells were incubated at 37°C for 2 hours, before adding 500 µl 1.5% carboxymethylcellulose sodium salt (Sigma-Aldrich, C5678) in DMEM with 5% FBS. After incubation at 37°C for 3.5 days, cells were washed 1x with PBS before adding amido black stain (10 mM Naphthol Blue Black (Sigma-Aldrich, 195243), 1 M glacial acetic acid, 170 mM sodium acetate) for 1 hour at RT.

Immunofluorescence Microscopy

Huh7 cells were grown on 19 mm coverslips in 12 well plates for immunofluorescence staining. After Zika virus infection, Huh7 cells were washed 2x with PBS, fixed with 4% paraformaldehyde (Thermo Fisher, 43368) in PBS for 15 min at RT, washed 3x with PBS, and permeabilised with 0.1% Triton in PBS for 10 min at RT. Cells were then washed 1x with PBS before blocking with 10% normal goat serum (NGS) (Thermo Fisher, 31873) in PBS for 1 hour at RT. Cells were then incubated with primary antibodies in 5% NGS in PBS

overnight at 4°C. The following day, cells were washed 4x with PBS for 5 min each and incubated with the appropriate secondary antibodies and BODIPY 493/503 (Invitrogen, D3922, 1:500) in 5% NGS in PBS for 1 hour at RT. Cells were washed a further 4x with PBS for 5 min each and then mounted onto glass slides with mounting media containing DAPI (Abcam, ab104139). The following primary antibodies were used: rabbit ZIKV E (GeneTex, GTX133314, 1:200) and mouse dsRNA (Sigma-Aldrich, MABE1134, 1:100). The secondary antibodies used were Alexa Fluor 555 anti-mouse IgG (Invitrogen, A-21422, 1:500) and Alexa Fluor 647 anti-rabbit IgG (Invitrogen, A-21244, 1:500). Stained cells were imaged using a Zeiss 880 Airyscan confocal microscope at a magnification of 63x with oil immersion (NA: 1.40).

Transmission Electron Microscopy

Cells were seeded on 13 mm coverslips in 12 well plates. After 24 hours of Zika virus infection with an MOI 2, the supernatant was removed and pre-warmed fixative agent (4% formaldehyde, 2.5% glutaraldehyde, 0.1 M PIPES buffer) was immediately added to cells for 1 hour at RT. Samples were then stored at 4°C until processing.

Ultrathin sections (90 nm) were cut using a Leica UC7 ultramicrotome, post-stained with lead citrate and imaged on a Thermo Fisher Tecnai T12 TEM at 120 keV (with Gatan OneView CMOS camera).

Cryo-FIB milling and cryo-TEM data acquisition:

Lamellae were prepared from Huh7 cells grown on gold Quantifoil R2/1 and plunge frozen in liquid ethane, using the Arctis (ThermoFisher Scientific) plasma FIB. Prior to milling, the cells were coated with platinum followed by a layer of organoplatinum (trimethyl(methylcyclopentadienyl)platinum(IV)) using the gas injection system. Selected cells were successively thinned using a 30kV plasma beam and currents 4nA to 0.1nA before being polished using a current of 30pA.

Tilt series data was collected on Titan Krios (ThermoFisher) cryo-electron microscope equipped with a Falcon 4i detector and a SelectrisX energy filter. Data acquisition was performed using Pacetomo[39] scripts in SerialEM[40] covering an angular range of ±60˚ in 3˚

increments using a dose-symmetric scheme[41]. Each tilt series was collected with a nominal defocus of 5µm and movies were recorded in EER format[42] using a dose of 2.91 e/Å^2 per tilt resulting in a cumulative dose of 120 e/Å^2 over the full tilt series. Magnification was nominally 64000x resulting in a calibrated pixel size of 1.97 Å/px on the detector.

Tomogram reconstruction

WarpTools[43] was used to perform reference-free motion correction, CTF and defocus estimation on the individual frameseries data followed by the generation of dose-weighted tilt-series files. Tilt series were then aligned using AreTomo2[44] and the alignments were imported back into WarpTools for generating binned CTF-corrected tomograms as well as half-sets for denoising to boost contrast and aid the interpretability of the tomograms.

Lipid Droplet Isolation

Lipid droplets were isolated by sucrose density gradient centrifugation as previously described. Briefly, cells were homogenised in buffer A (20 mM Tricine-KOH pH 7.8, 250 mM sucrose, 1 mM EDTA) using a Potter-Elvehjem homogenizer. The homogenate was adjusted to 20% sucrose and overlaid with 5% sucrose in buffer A. After centrifugation at 152,000 × g for 1 hour at 4°C, the LD fraction was collected from the top interface.

Proteomics Analysis

Isolated lipid droplets were lysed in 8 M urea buffer, proteins were reduced with dithiothreitol, alkylated with iodoacetamide, and digested with trypsin. Peptides were analysed by liquid chromatography-tandem mass spectrometry (LC-MS/MS) using an Orbitrap Fusion Lumos mass spectrometer. Data were processed using MaxQuant software and analysed using Perseus for statistical analysis and volcano plot generation.

Virus-Like Particle (VLP) Assembly Assay

1 µg of a pcDNA plasmid containing ZIKV prM and E [11] was transfected into Huh7 WT, SOAT1 and SOAT2 knockout cells using Lipofectamine 3000 (Invitrogen, L3000015). Cell lysates were collected 48 hours in NP-40 cell lysis buffer (50 mM Tris-HCl pH 7.5, 150 mM NaCl, 2 mM MgCl2, 5 mM EDTA, 0.5% NP-40) with protease inhibitors after transfection.

Acylation Profiling

For lipid acylation analysis, cells were labelled with [^{14}C]-oleic acid (50 µCi/ml) for 4 hours. Lipids were extracted using chloroform:methanol (2:1) and separated by thin-layer chromatography (TLC) on silica gel plates. Radioactive signals were detected by autoradiography.

For protein acylation analysis, cells were labelled with alkyne-C16 (50 µM) for 4 hours, lysed, and subjected to click chemistry using azide-biotin. Biotinylated proteins were captured using streptavidin beads and analyzed by Western blot.

Cholesterol tracking

For cholesterol tracking experiments, cells were incubated with alkyne-cholesterol analogue (10 µM) for 16 hours before infection. At indicated time points, cells were fixed and subjected to copper-catalyzed azide-alkyne cycloaddition (CuAAC) click chemistry using azide-Alexa Fluor 594 to visualize cholesterol distribution.

For photo-crosslinking experiments, cells were labelled with diazirine-alkyne cholesterol analogue (10 µM) for 16 hours, infected with ZIKV, and exposed to UV light (365 nm, 15 minutes) at 24 hours post-infection. Crosslinked proteins were captured using CuAAC click chemistry with azide-biotin followed by streptavidin pulldown and Western blot analysis.

Clinical Data Analysis

Dengue patient data from a Sri Lankan cohort (n=561) were analysed with appropriate ethical approval. Central obesity was defined as waist circumference ≥80 cm in women or ≥90 cm in men. Disease severity was classified as dengue fever (DF) or dengue haemorrhagic fever (DHF) according to WHO criteria. Statistical analysis was performed using Fisher's exact test for categorical variables.

Statistical Analysis

Data are presented as mean ± standard error of the mean (SEM) from at least three independent experiments unless otherwise stated. Statistical significance was determined using unpaired Student's t-test for two-group comparisons or one-way ANOVA with Dunnet's

post-hoc test for multiple comparisons. P-values <0.05 were considered statistically significant. Statistical analyses were performed using GraphPad Prism 9 software.

Figure legends

Figure 1. Flavivirus infection alters lipid droplet dynamics and increases fatty acylation

(A) Confocal microscopy of WT and AUP1$^{-/-}$ Huh7 cells stained with BODIPY (LD, green), Sec61 (ER, red) and DAPI (nuclei, blue) at 10, 24, and 48h in control and ZIKV-infected cells. **(B)** Quantification of LD numbers per cell in WT and Aup1$^{-/-}$ cells over infection time course. n≥30 cells per condition across 3 independent experiments; data show mean ± SEM. **(C)** Schematic of AUP1-Ube2g2 mediated lipophagy releasing fatty acids for enhanced acylation and phospholid remodelling during infection. **(D-E)** Acylation profiling by TLC and silver staining in mock vs. ZIKV-infected WT and Aup1$^{-/-}$ cells at 48h p.i. **(D)** Lipid acylation in [^{14}C]oleate labelled cells and **(E)** protein acylation levels in alkyne C16 labelled cells subjected to biotinylation by click reaction. **(F-G)** Schematic of ZDHHC palmitoyltransferases and MBOAT acyltransferases (11 family members). **(H-I)** Viral titres (pfu/ml) at 24h p.i. following siRNA knockdown of ZDHHC family and MBOAT family members. n=3 replicates; **p<0.01, ***p<0.001 vs. scrambled control.

Figure 2. SOAT1 and SOAT2 activities are required for productive viral infection

(A) Schematic of LD biogenesis showing SOAT1/2-mediated conversion of free cholesterol and fatty acyl-CoA to cholesteryl esters (CE) and DGAT1/2-mediated synthesis of triacylglycerols (TAG) for LD formation. **(B)** qRT-PCR analysis of acyltransferase mRNA expression in Huh7 cells at 24h and 48h post-ZIKV infection (MOI 2). Data normalised to beta-actin; n=3 independent biological replicates; *p<0.05 vs. mock. **(C)** Cholesteryl esterification activity assay using [^{14}C]-cholesterol (left) and [^{14}C]-oleate incorporation in mock and ZIKV-infected cells. Lipids are extracted from cells at 10 hpi and resolved by TLC. Percentage of cholesteryl esters is measured by densitometry at 10h and 24h post infection. **(D)** Western blot validation of CRISPR/Cas9-generated SOAT1$^{-/-}$ and SOAT2$^{-/-}$ Huh7 cell lines. Vinculin is used as loading control. **(E)** Viral titres (PFU/ml) in culture supernatants from WT, SOAT1$^{-/-}$, and SOAT2$^{-/-}$ cells at 24h and 48h post-ZIKV infection. n=4 independent experiments; ***p<0.001 vs. WT. **(F)** Viral titres following pharmacological inhibition. Left panels: Avasimibe (20μM) treatment at 2h post infection in Huh7 cells infected with ZIKV compared to DMSO (n=3). Middle panel: Avasimibe (20μM) treatment in Huh7 cells after DENV infection, at 24h and 48h (n=3). Right panel: Individual and combined inhibition of DGAT inhibitor vs DMSO controls in ZIKV infected cells.

Figure 3. SOAT1 and SOAT2 activities are required for morphogenesis of virions

(A) Immunofluorescence microscopy of WT, SOAT1$^{-/-}$, and SOAT2$^{-/-}$ Huh7 cells at 24h post-ZIKV infection. dsRNA (green) marks viral replication organelles, ZIKV E protein (red) indicates assembly sites, DAPI (blue) shows nuclei. LDs are stained with BODIPY 493/503 in white. Insets show higher magnification of boxed regions. Scale bars 10μm. Right panel: Quantitative analysis from panel A. Total area of dsRNA per cell (μm²). n≥50 cells per condition across 4 independent experiments; ****p<0.0001. **(B)** Fold change in ZIKV E RNA expression normalised to WT controls. N=3 independent experiments; ****p<0.0001. **(C)** Schematic of virus-like particle (VLP) system expressing flavivirus structural proteins prM and E for assembly analysis independent of replication. **(D)** Western blot analysis and quantification of prM and E protein levels in VLP-transfected cells. Left: Representative blot showing prM (~17kDa), ZIKV E (~50kDa), and GAPDH loading control. Right: Normalised prM and E protein intensity in WT vs. SOAT knockout cells. n=3 biological replicates; **p<0.01, ***p<0.001. **(E)** Transmission electron microscopy of WT Huh7 cells at 24h post-ZIKV infection (MOI 2). Top row: Mock and infected cells. Bottom row: Magnified ZIKV-infected cells showing expanded ER with vesicle packets (replication organelles, red arrows) containing electron-dense viral RNA adjacent to assembled virions. Right: Quantification of indicated structures per field. n=20 fields per condition. **(F)** TEM analysis of SOAT1$^{-/-}$ and SOAT2$^{-/-}$ cells showing dramatically reduced replication organelle formation and complete absence of virus particles. Replication organelles present are predominantly empty (devoid

of electron-dense material). Right: Quantification showing reduced RO abundance and absent virus particles in knockout cells compared to WT.

Figure 4. Remodelled cholesteryl ester enriched lipid droplets supply lipids for replication organelles and virus assembly sites

(A) Transmission electron microscopy of lipid droplets (red arrows) in mock-infected and ZIKV-infected Huh7 cells (24h p.i.) across WT, SOAT1$^{-/-}$, and SOAT2$^{-/-}$ cell lines, showing ER-enwrapped LDs with altered LD morphology in infected WT cells. **(B)** cryo-EM of individual lipid droplets from FIB-milled samples showing internal structure. Mock-infected cells display homogeneous LD structure while ZIKV-infected cells exhibit liquid crystalline phases characteristic of cholesteryl ester-enriched LDs. **(C)** Quantitative analysis of LD internal organisation. Left: Radial intensity profile analysis measuring structural organisation within LDs (ring spacing ~3.9nm indicates liquid crystalline phase). Right: Histogram showing distribution of LD areas (nm²). **(D)** Chemical biology approach using cholesterol alkyne analogue for metabolic labelling. *Top*: Structure of alkyne cholesterol probe. *Bottom*: Time course of labelling followed by ZIKV infection -/+ SOAT inhibition. TLC analysis for resolving free cholesterol (Chol) and cholesteryl esters (CE). **(E)** Temporal tracking of cholesterol localisation during ZIKV infection using click chemistry-labelled alkyne cholesterol. Confocal microscopy at 10h and 24h p.i. in untreated and SOAT inhibitor-treated cells. DAPI (blue) shows nuclei, cholesterol signal (red) demonstrates redistribution from LDs (green) to viral replication/assembly sites (dsRNA, white) over time. Insets show higher magnification of boxed regions. Scale bars 10µm.

Figure 5. Cholesteryl ester-derived cholesterol directly interacts with viral proteins and cannot be substituted by free cholesterol

(A) Structure of bifunctional cholesterol analogue containing both photo-crosslinking capability (diazirine group) and click chemistry reactivity (alkyne group) for identifying cholesterol-protein interactions. **(B)** Experimental workflow for photo-crosslinking analysis. Cells are metabolically labelled with cholesterol analogue (16h), infected with ZIKV (24h), UV-irradiated for photo-crosslinking (3h), followed by Cu-catalysed azide-biotin click reaction and streptavidin capture. **(C)** Silver staining of streptavidin-captured proteins showing distinct profiles between mock and ZIKV-infected cells with different experimental conditions in a UV exposure and alkyne-cholesterol (alk-chol) dependent manner. Distinct protein profiles are captured between mock and ZIKV infection,10h and 24h post-infection. **(D)** Western blot analysis of cholesterol-crosslinked proteins from streptavidin captured fractions in control DMSO-treated cells (left) and SOAT inhibitor treated cells (right). Caveolin and NPC1 serve as positive controls and β-tubulin serves as the loading control. **(E-F)** Time-of-addition rescue experiments with oleic acid or free cholesterol in SOAT-inhibited cells. Experimental timeline showing SOAT inhibitor treatment with oleic acid (OA) or cholesterol (Chol) supplementation at the start of or 24h post infection. Viral titres (PFU/ml) measured in supernatants from control and lipid supplementation cells by plaque assays. (F) left panel: supernatant collected before lipid supplementation at 24h to confirm reduction in viral titres upon SOAT inhibition. n>3 biological replicates; *p<0.05, **p<0.01, ns=not significant vs. DMSO control.

Figure 6. Cholesteryl ester-rich LDs are enriched in fatty acid remodelling and lipid transport proteins

(A) Schematic of lipid droplet isolation protocol using sucrose density gradient centrifugation for proteomic analysis of LD fractions from mock and ZIKV-infected cells. **(B)** Volcano plots of LD proteome analysis comparing ZIKV-infected vs. mock cells. Top: Untreated conditions with significant enrichment of fatty acid remodelling enzymes, Rab-GTPases, lipid transport proteins (LTPs), and sphingomyelinases (red dots = upregulated, green dots = downregulated). Bottom: SOAT inhibitor-treated conditions with altered protein recruitment patterns. FDR-adjusted p-values plotted against log□(fold change); dotted line indicates significance threshold. **(C)** Western blot validation of key ER and membrane remodelling

proteins. Analysis of whole cell lysates at 24h and 48h post-infection comparing mock vs. ZIKV infection under DMSO control or SOAT inhibitor treatment. Proteins indicated: CIDEC (lipid transferase), TMEM41B (lipid scramblase), VMP1, and MLN64, Calnexin, Reticulon 4 and Plin2. Vinculin, and GAPDH serve as loading controls. **(D)** Disease severity distribution by central obesity status. Stacked bar chart showing the proportion of dengue fever (DF, gray) and dengue haemorrhagic fever (DHF, red) in lean (n=303) and centrally obese (n=258) patients. Central obesity was defined as waist circumference ≥80 cm in women or ≥90 cm in men. p=0.0407, Fisher's exact test. **(E)** DHF risk comparison by central obesity status. Bar chart displaying the percentage risk of developing dengue haemorrhagic fever in patients with central obesity (33.3%) versus those without central obesity (25.4%). The absolute risk difference of 7.9 percentage points represents a significant increase in severe disease risk associated with central obesity (relative risk 1.22, 95% CI: 1.01-1.46; odds ratio 1.47, 95% CI: 1.02-2.12).