Integrative Chemical Genetics Platform Identifies Condensate Modulators Linked to Neurological Disorders

Dylan Poch,[1] Chandrayee Mukherjee,[1] Sunanda Mallik,[1] Vanessa Todorow,[1] E.F. Elsiena Kuiper,[1,2] Nalini Dhingra,[3] Yulia V Surovtseva,[3] Christian Schlieker.[1,4,†]

[1]Department of Molecular Biophysics and Biochemistry, Yale University, New Haven, CT 06511, USA.

[2]Max Planck Institute of Biophysics, Mechanisms of Cellular Quality Control, Frankfurt a. M., Germany

[3]Yale Center for Molecular Discovery, Yale University, West Haven, CT 06516, USA.

[4]Department of Cell Biology, Yale School of Medicine, New Haven, CT 06520, USA.

[†]Correspondence: christian.schlieker@yale.edu

Abstract

Aberrant biomolecular condensates are implicated in neurological disorders including ALS, frontotemporal dementia, and DYT1 dystonia, yet approaches to systematically identify their modulators remain limited. Here we establish MLF2 as a versatile condensate biomarker and develop CondenScreen, an integrated high-content screening and bioinformatic pipeline enabling identification of condensate modulators across chemical and genetic space. Screening 1,760 FDA-approved compounds in a cellular DYT1 dystonia model, we identify drugs that alter aberrant condensate properties, validating the platform for condensate-targeted drug discovery. In parallel, a genome-wide CRISPR/Cas9 screen links condensate accumulation to microcephaly genes and more than 10 additional neurodevelopmental disorders. Machine learning and confocal imaging resolves distinct condensate phenotypes: loss of microcephaly-associated ZNF335 results in nucleoplasmic condensates, whereas RNF26 deletion produces nuclear envelope condensates that phenocopy hallmarks of torsin deficiency. Our study provides a scalable resource for identifying corrective modulators of condensates and establishes a link between nuclear condensate accumulation and neurodevelopmental disorders.

Keywords

Biomolecular condensates, high-content screening, microcephaly, dystonia, ubiquitin, MLF2, nuclear pore, pyrithione zinc, machine learning, neurodegeneration, stress granules, ubiquitin ligase.

Introduction

Dysregulation of physiological condensates is a driver for many neurological disorders, including Amyotrophic Lateral Sclerosis (ALS), Frontotemporal Dementia (FTD), and DYT1 dystonia.[1–5] Biomolecular condensates, often made from a milieu of hundreds of different proteins, consist of a thermodynamically favorable separation between a dense and dilute phase.[6,7] When properly regulated, biomolecular condensates allow efficient, fine-tuned microenvironments, ensuring maintenance of cellular homeostasis while buffering against external perturbations.[8–10] However, in pathological states referred to as condensatopathies, these condensates exhibit reduced dynamicity, leading to protein aggregation or amyloid formation that is not easily reversible.[11–13] It is estimated that up to one-third of common diseases can be linked to condensate-forming proteins.[14] Determining mechanisms and cellular activities of these processes is essential for devising therapeutic strategies that tune condensate properties to counteract pathological transitions.

One favorable phase-state is observed in the central channel of the nuclear pore complex (NPC). The NPC serves as the intermediary gate at which nuclear transcription is separated from cytoplasmic translation in eukaryotic organisms. Within the central channel of the NPC, phenylalanine-glycine nucleoporins (FG-nups) have a high proportion of intrinsically disordered regions that set up a selective phase, contributing to the selective barrier of the NPC.[15–18] Disruptions to NPC assembly or nucleocytoplasmic transport have been strongly implicated in neurological diseases, including DYT1 dystonia,[1,19,20] a debilitating movement disorder characterized by involuntary muscle contractions and painful sustained postures.[21,22]

DYT1 dystonia is caused by a mutation within the TOR1A gene that disrupts activation of the encoded AAA+ ATPase,[23–27] TorsinA, resulting in the formation of pathological nuclear envelope condensates.[1,19,20,28,29] As with many condensatopathies, there are only palliative interventions, with no curative or preventative treatments for dystonia.[21,22] These nuclear envelope condensates accumulate K48-linked polyubiquitinylated proteins that evade proteasomal degradation and sequester chaperone proteins, causing cellular proteotoxicity.[1,20,30] Thus, regulating aberrant

31   nuclear condensates by either direct dissolution or modulation of its proteotoxic

32   properties represents a practical therapeutic approach. However, current limitations to

33   high-throughput screening campaigns targeting condensatopathies include a lack of

34   well-defined condensate targets and broadly applicable condensate biomarkers.

35   In this study, we establish Myeloid leukemia factor 2 (MLF2) as a robust reporter

that is enriched in proteotoxic nuclear condensates and stress granules, as evidenced

by co-localization with TDP-43 and G3BP1. Using an MLF2 readout, we integrated

complementary chemical and genetic high-content screening strategies to define the

landscape underlying atypical nuclear condensates. First, a target-agonistic chemical

screen in a cell-based DYT1 dystonia model identified several FDA-approved small

molecules that modulate proteotoxic nuclear condensates, providing a tractable

methodological framework for identifying therapeutics for condensatopathies. Second, a

CRISPR/Cas9 knockout (KO) screen in wild-type cells, combined with deep learning

approaches, identified RNF26 and ZNF335 as protective factors against distinct

condensate phenotypes. Notably, 15 out of the top 50 genetic hits are associated with

neurodevelopmental disorders, including seven directly linked to primary microcephaly

or related congenital brain malformations. Together, these orthogonal approaches

converge on a shared connection between nuclear condensate accumulation, perturbed

nuclear proteostasis, and neurodevelopmental disorders, while establishing a scalable

platform for therapeutic discovery.

Results

MLF2 is a Biomarker for Condensates Associated with ALS/FTD and DYT1

Dystonia.

Having established the connection between impaired protein quality control and

condensates in a torsin-deficient DYT1 dystonia HeLa cell model (4TorKO cells) (Figure

1A),[1,20] we aimed to develop biomarkers to facilitate the identification of condensate

modulators across other condensatopathies. MLF2 represents a major protein

constituent of nuclear envelope condensates within 4TorKO cells[20] and readily

immerses into FG-nup condensates *in vitro*,[1] likely due to its highly enriched Methionine

and Arginine residues. Since many condensates share aromatic side chains as part of

their molecular grammar,[31] we previously suggested that MLF2 might localize to

additional aberrant condensates through π-cation interactions as MLF2 is highly

enriched in Arg residues that cannot be functionally replaced by charge-conserving Lys

residues.[1] Here, we have exploited these properties to visualize proteotoxic

condensates by designing doxycycline (Dox) inducible fluorophore-derivatized variants

of MLF2 as condensate biomarkers.

Dysregulation of stress granules (SGs) has been pathologically associated with

several neurological disorders, including ALS and FTD.[11] To assess MLF2 localization to

stress granules, we co-expressed the SG markers G3BP1 and TDP-43 in wild-type

(WT) HeLa cells exposed to 0.5 mM sodium arsenite (NaAsO$_2$) for 30 minutes, an

established condition for SG induction.[32–35] Upon NaAsO$_2$ exposure, MLF2 exhibited

strong co-localization with both G3BP1-GFP and TDP-43-tdTomato (Figure 1B). These

findings suggest that MLF2 serves as a robust reporter of condensates associated with

distinct neurological disorders.

Figure 1, *Poch et al.*

Fig. 1. MLF2 is a biomarker for aberrant condensates across neurological disorders. (**A**) MLF2 detects both nuclear condensates and stress granules. Nuclear envelope condensates that sequester MLF2, FG-nups (not shown for clarity), and K48-Ubiquitin-enriched proteins have been identified in DYT1 dystonia models. The ability of MLF2 to immerse into condensates can be leveraged for high-content imaging platforms to systematically profile additional disease-relevant condensates. (**B**) MLF2 co-localizes with G3BP1 (top panels) and TDP-43 (bottom panels) after stress granule formation in response to oxidative stress (0.5 mM NaAsO$_2$, 30 min).

Establishing MLF2-GFP as a Condensate Reporter for High-Content Screening

Leveraging the condensate biomarker properties of MLF2, we developed a miniaturized,

high-content assay using Dox-inducible MLF2-GFP cells in a 384-well format. This

facilitated the design of phenotypic, high-content arrayed screens to identify chemical

and genetic modulators of nuclear condensates. Specifically, each well contained either

a small molecule applied to 4TorKO cells or synthetic single guide RNAs (sgRNAs)

delivered to Cas9-integrated WT cells, designed to suppress or promote nuclear

condensates, respectively. To analyze the high-resolution images (over 10k

images/plate), we developed a computational condensate screening suite,

CondenScreen. This CellProfiler and R script was designed for the rapid quantification

and statistical analysis of condensates (Figure S1A).

88   While many robust computational resources are available for monitoring

condensates (see *PhaseMetrics* and *Punctatools*)[36,37], *CondenScreen* is specifically

designed for large-scale, high-content screening analysis using a customizable

CellProfiler 4 [38,39] pipeline and *R* analysis script. To this end, *CondenScreen* quantifies

condensates at both single-cell and population levels and provides a graphical user

interface (GUI) allowing users to select plate size and layout, as well as tailoring

downstream calculations for a signal "ON" vs signal "OFF" screening campaign. In

addition, built-in quality control features automatically assess screen robustness using

established statistical metrics, including the Z'-Score, BZ-Score, signal-to-background

(SB), normalized effect, and coefficient of variation (CV). In addition, the platform has

optional filters to flag potential false positives arising from either spatial plate effects

(Figure S1B-S1D) and filter candidate compounds with known false positive liabilities,

including pan-assay interference compounds (PAINS) and additional structural alert

categories (Figure S1E). Integration of the MLF2-GFP inducible system with our

CondenScreen analysis pipeline ensures rapid identification of condensate modulators.

High-Content Screen Identifies FDA-approved Drugs that Modulate Condensates

To identify chemical matter that can modulate aberrant nuclear envelope condensates,

we employed our MLF2-GFP inducible 4TorKO DYT1 cell model. We screened 1760

drugs from the Pharmakon library, comprising 1360 US and 400 international drugs, at a

10 µM working concentration. To minimize false positives due to differential reporter

expression, Dox (0.5 µg/ml) was added for four hours, followed by an additional 2 hours

with compound incubation.

CondenScreen reported an average Z'-score of 0.75 indicating pronounced

separation between the positive (0.1% DMSO-treated WT HeLa) and negative (0.1%

DMSO-treated 4TorKO HeLa) controls.[40] Additionally, the ratio between the negative

and positive controls (SB) was 950, while the SD and CV of DMSO-treated cells were

1.9% and 8.3%, respectively, further confirming assay robustness.

To control for plate-specific column, row, and edge effects, we assigned each

compound a BZ-score based on its ability to decrease the number of nuclear MLF2-

GFP foci (Figure 2A, see methods). The distribution of the average number of MLF2-

GFP foci per condition was near-normally distributed (Figures 2B, 2C). Using a cutoff

BZ-score of three, we identified 20 putative hit compounds that significantly depleted

MLF2-GFP condensate foci. Top compounds included plumbagin, sanguinarium

chloride, thiram, piroctone olamine, penfluridol, bronopol, thimerosal, triclosan,

disulfiram, celastrol, dihydrocelastryl diacetate, and pyrithione zinc (PZ). A full list of the

screened drugs can be found in the supplementary materials (Table S1).

Top-scoring compounds were resolubilized to construct a 12-point dose response

(n = 4 replicates), covering over 5 orders of magnitude to determine the calculated IC_{50}.

(Figure 2D, Figure S2A). The normalized percent effect and percent toxicity (calculated

as a function of viability, nuclear area, and nuclear morphology) were quantified for each

concentration to determine potential therapeutic windows. Four structurally diverse

drugs demonstrated a potent dose-dependent relationship, which included celastrol

(IC_{50} = 124.8 nM), PZ (IC_{50} = 601.6 nM), plumbagin (IC_{50} = 7.4 µM), and disulfiram (IC_{50}

= 10.8 µM) (Figure 2D).

Figure 2, *Poch et al.*

Fig. 2. Chemical screen identifies drugs that modulate nuclear envelope condensates. (A) Drugs from Pharmakon 1760 screen decrease nuclear MLF2-GFP accumulation in 4TorKO HeLa cells. Compounds (circles) passing significance threshold (dotted-line, BZ-score > 3) are labeled purple, while red-labeled dots denote drugs selected for downstream validation. (**B**) Distribution in the average number of MLF2-GFP foci/cell and (**C**) percent decrease of MLF2-GFP foci per cell, normalized to the screening population. (**D**) Hit compounds are dose dependent. Determined IC_{50} from 12-point dose response (four orders of magnitude, n = 4). Black line represents % effect (decrease in MLF2-GFP foci), red line represents % toxicity relative to DMSO (four-parameter-log-logistic model). (**E**) Drugs modulate accumulation of K48-Ub proteins. Dots depict the number of MLF2-GFP (left, 6h Dox) or α-K48-ubiquitin (right) foci/cell after treatment with drug or DMSO. Asterisks indicate Bonferroni-corrected p-values calculated by Mann-Whitney U testing (***: p <= 0.001 ****: p <= 0.0001). (**F**) Representative immunofluorescence images of MLF2-GFP (green) and α-K48-ubiquitin (red) in torsin-deficient cells treated with DMSO compared to either PZ (3 μM) or celastrol (6 μM). (**G**) Schematic of inducible MLF2-GFP-P2A construct without (top) or with (bottom) Dox. (**H**) Confocal images of torsin-deficient cells expressing MLF2-GFP-P2A-tdTomato treated with either DMSO or 1.5 μM PZ treatments (6h). Scale bars, 5 μm.

Pyrithione Zinc Disrupts K48-Ubiquitinylated Protein Accumulation within Nuclear Condensates

To determine the optimal concentration, we tested each of the four candidate

compounds—celastrol, PZ, plumbagin, and disulfiram—at four tighter concentrations,

including the IC_{50} itself, at 2x above/below the IC_{50}, and 8x above the IC_{50}

concentrations, using a 3 h compound treatment. The resulting boxplot reflects each

compound's effect at the single-cell level (Figure S2B). Consistent with the significant

cell-cycle dependency of NE condensate formation previously demonstrated,[20] we

observed significant variability in the number of MLF2-GFP foci/cells. However, all

compounds significantly reduced perinuclear MLF2-GFP accumulation at 2x and 8x

their respective IC_{50} (Figure S2B).

To ensure consistent reduction across different condensate markers, we used

K48-linked ubiquitin (K48) as an independent and endogenous biomarker based on our

previous observation that 4TorKO cells have reduced protein turnover, resulting in the

accumulation of polyubiquitinylated proteins within nuclear envelope condensates.[1,20]

Using immunofluorescence staining with α-K48 antibodies, all four compounds

demonstrated a statistically significant decrease in endogenous, perinuclear K48 foci,

replicating our results at reducing MLF2-GFP accumulation (Figure 2E). Between the

optimal concentration profile of PZ and celastrol (Figure 2D), and the undesirable

structural elements of plumbagin (quinone, Michael acceptor; Figure S1F), we

continued to validate PZ and celastrol by assessing MLF2-GFP and endogenous K48

foci using confocal microscopy (Figure 2F). Importantly, this data validates that the

mechanism of action either directly or indirectly mediates endogenous protein cargo

within condensates.

A significant limitation to phenotype screening is the prevalence of identification

of false positives that inhibit biomarker expression by non-specifically disrupting cellular

protein transcription or translation. To circumvent this, we constructed a HeLa cell line

with an inducible bicistronic MLF2-GFP-P2A-tdTomato reporter (Figure 2G). The P2A

site induces ribosomal skipping, allowing the translation of two separate polypeptides,

an N-terminal MLF2-GFP and a C-terminal tdTomato, both derived from the same

mRNA transcript.[41] We selected MLF2-GFP-P2A-tdTomato 4TorKO cells using

Fluorescence-Activated Cell Sorting (FACS) (Figure S2C), before inducing MLF2-GFP

expression and treating cells with either PZ (0.75 µM or 1.5 µM, 6 h) or celastrol (1.5 µM

or 3 µM, 6 h) . We quantified the number of MLF2-GFP foci and tdTomato expression

between either uninduced cells treated with DMSO vehicle control, induced cells treated

with DMSO, and induced cells treated with either 100 µg/mL cycloheximide, PZ, or

celastrol. Concentrations of both celastrol and PZ at 1.5 µM had a significant decrease

in the number of MLF2-GFP foci/cell compared with the DMSO control (Figure S2D).

While PZ did not reduce the expression of secondary-tdTomato reporter (Figures 2H,

S2E), celastrol slightly, albeit at a non-statistically significant level, decreased secondary

reporter expression compared with DMSO-treated cells.

To address and quantify the potentially confounding effects of a compound

affecting reporter abundance non-specifically, we developed an empirical metric, the

Condensate Normalization Score (CNS). The CNS is based on the differential

fluorescence between the condensate biomarker and secondary reporter. A high CNS-

value indicates a reduction in condensate biomarkers with minimal impact on overall

protein levels (see methods). Both celastrol- and PZ-treated cells showed a significantly

increased percentage of cells with a median CNS exceeding 5-fold that of the DMSO

control (Figure S2F), indicating a mechanism independent of reporter expression.

To assess the specificity of the top two compounds in modulating aberrant

condensates, we examined their effects on nuclear Promyelocytic Leukemia (PML)

bodies, physiological phase-separated compartments involved in protein quality control.

As expected, treatment with 5% 1,6-hexanediol (5 minutes), a known disrupter of liquid-

liquid phase-separated environments,[42] significantly reduced PML body number (Figure

S2H, S2G). In contrast, neither PZ nor celastrol reduced PML body number at a

concentration of 1.5 μM that profoundly reduced MLF2 foci in our disease model (Figure

S2H, S2G). Interestingly, treatment with PZ resulted in a statistically significant increase

to the number of PML body puncta. Overall, this indicates that the activity of PZ and

celastrol does not reflect nonspecific disruption of physiological nuclear condensates.

However, visualization of condensates in 4TorKO cells using transmission electron

microscopy (TEM) did not reveal full dissolution of nuclear envelope condensates after

a 3 h treatment of 2 μM PZ (Figure S2I).

To identify the structural features responsible for the condensate-modifying

activity of PZ, we examined whether this effect originates from its pyrithione moiety (2-

Mercaptopyridine N-oxide) or the zinc ion. To this end, we treated 4TorKO cells with

either DMSO vehicle control, PZ alone, or PZ with zinc chelator TPEN

(N,N,N',N'- *tetrakis*-(2-Pyridylmethyl)ethylenediamine) at a 3 molar equivalent.[43] After

zinc chelation with TPEN, the pyrithione compound failed to reduce MLF2-GFP

condensate accumulation in 4TorKO cells (Figures S3A, S3B). Furthermore, we tested

whether zinc alone is sufficient to deplete nuclear MLF2-GFP condensate foci. Due to

its lower membrane permeability, 4TorKO cells were treated with higher concentrations

of zinc sulfate monohydrate ($ZnSO_4$). Treatment with 2 mM $ZnSO_4$ alone decreased

MLF2-GFP foci (Figure S3C, S3D) while maintaining tdTomato expression (Figure S3E),

resulting in a higher relative CNS value (Figure S3F). Thus, PZ modulates nuclear

condensates through a zinc-dependent mechanism.

Genome-Wide CRISPR Screening Uncovers Condensate Modulators Implicated in

Microcephaly and Neurodevelopmental Disorders

Having established a robust screening pipeline, we next performed a genome-

wide CRISPR/Cas9 KO screen to identify genetic determinants of nuclear condensate

accumulation. Knowledge of the corresponding activities may facilitate the identification

of future pharmacological targets. We screened an arrayed library of 19,195 human

genes using the Horizon Discovery Edit-R synthetic sgRNA collection. To increase KO

efficiency, we designed and FACS-sorted a WT HeLa cell line that constitutively

expresses Cas9 with Dox-inducible MLF2-GFP. Each well was reverse-transfected in a

384-well plate format using a pool of three sgRNAs targeting different regions of the

same gene (Figure 3A). Cells were incubated for 72 h, and MLF2-GFP was induced by

Dox 16 h before fixation (all within the 72 h window). After running all high-content

images through our *CondenScreen* pipeline, the average Z' value was 0.52,

demonstrating a dynamic range between the positive (TOR1A/B sgRNAs) and negative

(non-targeting (NT) sgRNA) controls. The calculated SB was 4.8, and the SD and CV of

the NT control were 3.1 and 9.7, respectively. We determined the assay quality score to

be sufficiently robust as indicated by similar scores in the literature.[34,44]

We defined a candidate gene as any gene deletion that resulted in a Z-score > 2

(Figure 3B). The top 10-ranked candidate genes were: RNF26, WAC, ZNF335,

NCAPG2, NCAPH2, ECT2, ASRGL1, SETD1B, ASB6, and FOXS1 (see Table S2 for

comprehensive list). Overall, the screen resulted in a near-normal distribution of the

number of MLF2-GFP foci per cell (Figures 3C, 3D). Notably, we observed several

genes linked to neurodevelopmental disorders, with high enrichment of microcephaly-

associated genes (orange labeled dots, Figure 3B), according to human genetics

annotated from OMIM[45] and the NIH Genetic Testing Registry (GTR).[46] Both

homozygous and compound heterozygous mutations to one of our top hits, zinc finger

protein 335 (ZNF335, also identified as MCPH10 or NIF1) have been documented to

cause primary microcephaly.[47–54] ZNF335 has been tied to the regulation of

neurogenesis through modulation of the REST complex.[48]

Additional top hits of interest include, RNF26, an E3-ubiquitin ligase that has

been implicated in regulating the endosomal system, serving as a "gatekeeper" for

vesicles in the perinuclear ER cloud.[55,56] Furthermore, RNF26 ubiquitinylates

SQSTM1[55] and has recently been tied to ER-sheet morphology and maintenance.[56]

Other hits include APOL3 (Z-Score = 2.5), a putative regulator of membrane dynamics

with detergent-like bactericidal properties;[57,58] Ran (Z-Score = 2.8), a GTP-binding

nuclear protein; and EIF4A3/DDX48 (Z-Score 3.8), a DEAD-box ATPase. DEAD-box

ATPases have recently been linked to the regulation of phase-separated membraneless

organelles,[59] indicating a potential role of EIF4A3 in modulating condensates. These

condensate phenotypes observed across gene deletions highlight critical entry points

into unexplored biology and promising directions for therapeutic discovery.

Figure 3, *Poch et al.*

Fig. 3. CRISPR KO Screen reveals genes that modulate nuclear condensates. (**A**) Schematic workflow of the genome-wide CRISPR KO screen using WT HeLa cells. Plates represent direct outputs provided by *CondenScreen*, indicating the number of MLF2-foci/cell (top plate) and cell count per image (bottom plate). (**B**) Genes hits from CRISPR KO screen associated with microcephaly modulate nuclear condensates. Dotted line represents significance threshold (Z-Score > 2) while highlighted and labeled genes have a Z-Score > 3. Green dots represent the TorsinA and TorsinB KO conditions; orange dots represent genes linked to microcephaly; and larger, red dots indicate genes of additional interest. (**C**) Distribution of the average number of MLF2-GFP foci/cell and (**D**) percent increase in MLF2-GFP foci/cell, normalized to the screening population. (**E**) Heatmap of the 30 highest scoring GO biological processes (rows) and associated genes (columns) from the top 100 candidate CRISPR KO hits. Vertical scale legend indicates -log(10) Benjamini-Hochberg adjusted p-value, with lighter regions representing higher significance. Scaled boxes (top) indicate the Z-Score of each gene from the CRISPR Screen, with darker colors indicating a higher Z-score (horizontal scale bar). Euclidean distance hierarchical clustering for GO-enriched biological pathways and genes is shown with the left and top dendrogram, respectively.

To determine if candidate genes are clustered into enriched genetic pathways,

we hierarchically clustered the top 100 gene hits with their corresponding 30 enriched

biological processes (q-value < 0.05) using Gene Ontology (GO) analysis[60] (Figure 3E;

interactive webpage available, see methods) and illustrated pathway

interconnectedness (Figures S4A-S4C). We observed enrichment (10 out of 30 GO

terms) of chromosomal and chromatid condensation, separation, or segregation

pathways (Figures 3E, S4A). GO pathways were manually clustered into the following

five categories: Autophagy, Cell Cycle, Chromosomal Regulation, Protein Transport,

and Other (Figure S4B). Consistent with these enriched pathways, loss of protein

homeostasis,[1,61] a cell-cycle dependency of condensates,[20] and disruption to

nucleocytoplasmic transport[62,63] have been documented in condensatopathies.

Additionally, dysregulation of both cell cycle checkpoints and chromosomal regulation

has been linked to primary microcephaly.[64–66] Indeed, seven out of the top 50 gene hits

have been linked to microcephaly.[45,46] Beyond the enrichment of individual gene hits

linked to microcephaly, encephalopathies, and other neurodevelopmental disorders,

additional enriched pathways of interest included organelle fission and positive

regulation of neuroblast proliferation.

To validate the three highest candidate genes, we knocked out ZNF335, RNF26,

and the WW Domain Containing Adaptor with Coiled-Coil (WAC), and binned them

according to the number of nuclear MLF2-GFP foci/cell (5-15, 15-30, or >30 foci/cell)

(Figure S4D). While WAC depletion had only a modest effect, RNF26 KO and ZNF335

KO cells displayed pronounced nuclear condensate accumulation as indicated by

MLF2-GFP accumulation (Figure S4D), with ZNF335 deficient cells resulting in the

greatest number of condensates.

Supervised Machine Learning Links Genetic Perturbations to Distinct Condensate

Phenotypes

To resolve phenotypic distinctions and cluster genetic perturbations by their condensate

morphology, potentially highlighting divergent biological roles, we developed a deep

learning model to capture subtle morphological variations between condensates. Using

a residual convolutional neural network (Figure S5A),[67] we trained the model on single-

cell patches derived from our strongest validated genetic hits to determine whether

condensates differences could be computationally discriminated (Figure 4A). We

employed a ResNet18 neural network backbone (ImageNet)[68] to learn a regression

mapping across three reference states: RNF26 KO (-1), non-targeting (0), and ZNF335

KO (+1). The model assigned each unlabeled cell a continuous phenotypic similarity

score ranging from −1 (RNF26-like) to +1 (ZNF335-like)

After optimization, the model correctly classified 70% of individual cells to their

respective anchor conditions (Figure 4B), demonstrating strong performance metrics

(area under the curve [AUC] = 0.95, loss = 0.14, true vs. predicted correlation = 0.76;

Figures 4C, S5B–C). Notably, fewer than 2% of RNF26 KO cells were misclassified as

ZNF335 KO and vice versa (Figure 4E), demonstrating the model's ability to

discriminate and report biologically meaningful results. To confirm that predictions were

driven by biologically interpretable morphological features, we employed post-hoc

saliency mapping based on the integrated gradient (see methods, Figure S5F, S5G).[69]

Consistent with this, the RNF26 KO cells resulted in a higher number of smaller, more

heterogeneous nuclear condensates when compared to the ZNF335 KO condition

(Figure S5G).

Independent of the model's continuous regression output, we extracted raw

feature embeddings for each CRISPR hit with a Z-score > 2 and averaged single-cell

embeddings per condition to generate centroid vectors. These were projected into two

dimensions via Uniform Manifold Approximation and Projection (UMAP), [70] revealing

three distinct clusters by KMeans clustering (Figures S5D, S5E). The resulting UMAP

visualization (Figure 4F) separated genes into groups phenotypes resembling the

ZNF335 KO cells (cluster 1), RNF26 KO cells (cluster 2), or the NT control cells (cluster

3).

Figure 4, *Poch et al.*

Fig. 4. Deep learning model differentiates and clusters distinct condensate phenotypes. (**A**) RNF26 and ZNF335 have distinct MLF2-GFP condensate phenotypes. Representative confocal imaging of RNF26 KO (left) or ZNF335 KO (right) HeLa cells expressing MLF2-GFP (green). Scale bars, 10 µm. (**B**) Accuracy curve of model during training. Accuracy reflects the percentage of validation cells correctly classified as RNF26 KO, ZNF335 KO, or NT control. Peak validation accuracy (0.70) was achieved at epoch 21. (**C**) Receiver operating characteristic curve of optimized model. Distinguishes true positives (y-axis) vs false positives (x-axis) with a calculated AUC of 0.95. (**D**) Correlation histogram indicating the similarity for each trained cell (x-axis) to one of three conditions (-1 = RNF26 KO, 0 = NT Ctrl, +1 = ZNF335 KO). Cells are colored by their true knockout condition (light orange = RNF26 KO; red = NT Ctrl; purple = ZNF335 KO). (**E**) Confusion matrix depicting the predicted label (x-axis) of each cell in the validation set compared to its true label (y-axis), based on rounding its similarity score to the nearest integer. (**F**) Top scoring CRISPR KO genes cluster into three distinct groups. Gene-level centroid embeddings of the top 358 genes (Z-Score > 2) were extracted before UMAP dimensionality reduction. K-means clustering partitions genes into three phenotypic clusters: cluster 1 (ZNF335-like), cluster 2 (RNF26-like), and cluster 3 (NT-like). (**G**) Hierarchical clustering heatmap showing phenotypic similarity of the top 50 gene knockouts to RNF26 or ZNF335 KO anchors. Darker colors indicate greater similarity to the corresponding anchor (RNF26 KO, left; ZNF335 KO, middle; NT control, right). The RNF26 and ZNF335 KO conditions are labeled in green, while the non-SMC condensin II subunits are labeled in orange.

Finally, centroid feature vectors were used to compute the similarity metrics for

each gene against the RNF26 KO, ZNF335 KO, and NT control conditions. We

clustered gene-level similarity profiles using hierarchical clustering and constructed a

heatmap of the top 50 genes sharing high phenotypic similarity to RNF26 KO or

ZNF335 KO conditions (Figure 4G). Of interest, knocking out any of the non-SMC

subunits of the condensin II complex (NCAPG2, NCAPD3, and NCAPH2)[71], and to a

lesser extent, the condensin II SMC2 subunit, displayed condensates with high

phenotypic similarity to the ZNF335 KO condition (Figure 4G).

RNF26 and ZNF335 Knockouts Exhibit Distinct Phenotypes but Share

Accumulation of K48-Ubiquitin

To validate the deep learning results suggesting that condensates arising from

knocking out RNF26 and ZNF335 are phenotypically distinct, we used

immunofluorescence staining to determine if these condensates accumulate

polyubiquitinylated proteins. Similar to those observed in torsin-deficient cells (see

chemical screen), condensates in both RNF26 KO and ZNF335 KO cells displayed co-

localization with K48-Ub using immunofluorescence imaging (Figure 5A). Next, we

assessed whether both condensate conditions co-localize with nucleoporins by

performing immunofluorescence staining using α-mAb414, which detects NPC-localized

nucleoporins. RNF26 KO cells showed prominent MLF2-GFP co-localization with

mAb414 (Figure 5B), analogous to that observed in torsin-deficient cells with stalled

NPCs (Figure 5B). In contrast, analysis of confocal Z-stacks revealed that condensates

in ZNF335 KO cells rarely shared significant co-localization with mAb414 staining

(Figure 5B, 5C), supporting the prediction that RNF26 and ZNF335 KO conditions

promote condensates with distinct properties.

Consistent with immunofluorescence data, TEM imaging demonstrated that

RNF26 KO cells contain condensates between the nuclear envelope (Figure 5D,

shaded yellow). Additionally, RNF26 KO cells feature pronounced NE invaginations

reminiscent of nucleoplasmic reticulum formation (Figure 5D).[72,73] While we observed

several NE invaginations contiguous with the inner nuclear membrane (INM), further

electron tomographic studies are needed to determine if all invaginations are contiguous

with the NE. We then treated RNF26, ZNF335, or TOR1A/B depleted cells with 5% 1,6-

hexanediol for five minutes. Only 1,6-hexanediol, but not DMSO or 2,5-hexanediol,

dissolved all three types of condensates (Figure S6). Together, these results indicate

that while both RNF26 and ZNF335 have a role in safeguarding against nuclear

condensate accumulation, only lack of RNF26 results in a condensate phenotype

similar to the one observed upon torsin deletion.

Chemical hit Pyrithione Zinc Modulates Condensates Induced by Genetic

Knockouts.

Finally, we tested whether compounds identified in our chemical screen could

modulate the condensate phenotypes arising from the genetic perturbations, including

those linked to microcephaly. Due to the ability of PZ to potently modulate MLF2-GFP

condensate markers in torsin-deficient cells (Figures 2D, 2E), we tested whether it could

similarly modulate NE condensates in RNF26 KO and ZNF335 KO cells. In both KO

conditions, treatment with 2 µM PZ significantly reduced accumulation of nuclear MLF2-

GFP condensate foci (Figures 5E, S7). Thus, FDA-approved drugs, including PZ, can

be repurposed to modulate condensates across phenotypically distinct genetic

perturbations.

Figure 5, *Poch et al.*

Fig. 5. Knockout of RNF26 or ZNF335 induces nuclear condensate accumulation. (**A**) K48-ubiquitinylated proteins accumulate within nuclear condensates. Immunofluorescence confocal images comparing co-localization of MLF2-GFP (green) with α-K48-Ub staining (red) in Cas9-integrated HeLa cells transfected with either (from left to right) NT, TOR1A/B, RNF26 or ZNF335 sgRNAs. (**B**) Condensates of RNF26 KO cells co-localize with NPC markers. Both RNF26 and TOR1A/B KO conditions demonstrated strong co-localization (yellow) of MLF2-GFP (green) with α-Mab414 staining (red). (**C**) Z-stack confocal projections of cells treated with the same conditions as (B). (**D**) TEM images of RNF26 KO cells depicting pronounced nuclear invaginations. Condensates are shaded yellow, while the INM and ONM are outlined in magenta and cyan, respectively. Black arrows draw attention to the interconnectedness of finger-like nuclear protrusions with INM; black asterisks indicate putative NPCs; red asterisks indicate omega-shaped herniations N, nucleus; C, cytoplasm. (**E**) Quantification of MLF2-GFP foci in CRISPR KO cells after PZ treatment. HeLa expressing MLF2-GFP (green) were treated with NT, Tor1A/B, RNF26, or ZNF335 sgRNAs along with either 2 μM PZ or DMSO for 3 h. KO conditions treated with DMSO are shown on the left (black outline) while those treated with PZ are shown to the right (blue outline). Asterisks indicate Bonferroni-corrected p-values by Mann-Whitney U testing (****: p <= 0.0001). IF scale bars at 10 μm (main) and 2 μm (inset); EM scale bars at 250 nm (main) and 100 nm (inset).

Discussion

This study establishes a scalable high-content platform optimized for identifying

chemical and genetic modulators of aberrant condensates that are increasingly

implicated in neurological disorders.[12,14,74] We paired a chemical screen in a DYT1

dystonia disease model with a genome-wide CRISPR KO screen in WT cells enabling

complementary analysis of condensate modulation. This integrated approach identified

FDA-approved compounds that alter the properties of aberrant condensates in a

disease model and uncovered genes that are tied to nuclear condensate homeostasis in

WT cells. Notably, several genes identified through this study are mutated in patients

with microcephaly, dystonia, and related neurodevelopmental disorders, highlighting

previously unappreciated commonalities in the cellular pathology underlying these

conditions. In addition, our study identifies previously unknown condensate-modifying

genes which may inform both patient-based sequencing and mechanistic studies.

Using the known proclivity of MLF2 to immerse into aberrant condensates,[1] we

exploit MLF2 derivatives as phenotypic condensate reporters. Remarkably, our reporter

allows for detection of DYT1 dystonia-associated condensates under single torsin

deletion conditions (Fig 3B) that remain undetectable by conventional markers.[28] In

addition, we generalize the utility of MLF2-based reporters for stress granules linked to

ALS/FTD (Figure 1B). Notably, MLF2 was recently found as resident in Poly-GA and

phosphorylated-TDP-43 inclusions in C9orf72 ALS/FTD patient samples,[75] as well as

co-localization of its *Drosophila* homolog (dMLF) with polyglutamine aggregates[76,77],

suggesting that our readout can be applied to a variety of neuropathological conditions

in the future.

Importantly, our screening approach is target-agnostic, utilizing MLF2 solely as a

readout for condensate detection at low expression levels to avoid altering condensate

dynamics. This allows our platform to be readily adapted to support a range of

condensate screening campaigns, even when the affected disease allele is not itself a

condensate constituent. This is exemplified by TOR1A/B or ZNF335 KO conditions,

which induce condensates via indirect mechanisms, representing a key advantage

since condensates are often studied under conditions of disease-allele overexpression,

which may perturb physiological condensate equilibria.

Inspired by recent condensate-modifying (c-mod) campaigns,[34,35,78,79] we

screened drugs that could modulate condensates either directly or indirectly. Several of

our identified drug hits caused robust re-localization of both our MLF2-GFP biomarker

and endogenous (K48-Ub) nuclear condensate markers (Figures 2E, 2F), without

significantly altering reporter expression or non-specifically altering physiological phase-

environments (e.g., PML bodies, Figure S2G, S2H). To assess the ability to repurpose

our chemical hits for other condensatopathies, we compared our top 20 drug hits with

two recent screens that identified modulators of stress granules.[34,35] Half of our drug hits

(10 out of 20), including PZ, celastrol, and disulfiram were identified among the top 100

hits of these stress granule screens (Table S4).[34,35] The ability of identical chemical

matter to modulate disparate disease phenotypes underscores the potential of shared

etiological principles across aberrant condensates.

To address the incomplete understanding of the dynamics driving condensate

formation and the paucity of reliable targets, we repurposed our readout to conduct a

genome-wide CRISPR KO screen toward target-directed approaches. Using a deep

residual convolutional neural network, further analysis of the resulting condensate

phenotypes identified two distinct clusters around either the RNF26 KO or ZNF335 KO

condition (Figure 4F). To distinguish these clusters, we performed downstream confocal

and electron microscopy on the RNF26 KO and ZNF335 KO conditions. Previous work

has implicated RNF26 in ER positioning, endosome retention, and Sec62-mediated

recovERphagy.[55,56,80] While condensates in both knockout conditions accumulated K48-

linked ubiquitin (Figure 5A), indicating dysregulation of cellular protein quality control,

only RNF26-deficient cells showed strong condensate colocalization with nucleoporins

(Figure 5B, 5C) and prominent omega-shaped inner nuclear membrane invaginations

(Figure 5C, 5D).

The identification of RNF26—previously unrelated to Torsins or NE morphology—

strongly validates our unbiased genetic approach. Building on our prior discovery that

MLF2 is a resident of nuclear envelope blebs,[20] the hallmark phenotype of DYT1

dystonia,[1,19,20,28,29] we found here that RNF26 deletion provokes a nearly identical

cellular pathology characterized by accumulation of MLF2, nucleoporin-associated

nuclear condensates, and omega-shaped inner nuclear membrane invaginations, all of

which have previously been observed across dystonia models.[1,19,20,28,29,62] Thus, our

study strongly motivates future efforts to explore how RNF26, Torsins, and NE dynamics

are functionally connected.

Depletion of ZNF335, a C2H2-type zinc finger and component of the trithorax

H3K4-methylation complex,[48] resulted in the most pronounced condensate phenotype,

with the average number of condensate foci exceeding that found in either the RNF26

or TOR1A KO conditions (Figure S4D). Mutations or aberrant splice variants of ZNF335

result in severe microcephaly and frequently reduced life expectancy.[48–53] Moreover,

ZNF335 has been reported to regulate the REST/NRSF complex with its knockdown

inhibiting neuronal differentiation. [48]

In addition to ZNF335, pathogenic allelic variants of 15 out of the top 50 gene

hits are associated with neurodevelopmental disorders according to human test results

from OMIM[45] and the NIH GTR.[46] (Figure 6A). Strikingly, 7/15 of these genes (ZNF335

[MCPH10], CIT [MCPH17], NCAPD3 [MCPH22], CDK5, SIN3A, SHH, and SPDL1) are

linked to microcephaly, holoprosencephaly, or lissencephaly (OMIM, GTR), along with

additional findings implicating NCAPG2, NCAPH2, and EP300 with microcephaly.[48,64,81–

84] Given that whole-exome sequencing fails to identify causal disease-causing

mutations in more than 70% of microcephaly patients,[85] our curated list of condensate-

associated genes provides a valuable resource to prioritize candidates in future patient

genetic studies.

Several of the microcephaly-associated gene hits identified in our screen are also

involved in chromosomal condensation, including several independently identified

subunits of the condensin II complex. Microcephaly has previously been labeled as a

"condensinopathy",[64] not to be confused with condensatopathies, reflecting prior work

that established a relationship between chromosomal condensation defects and

reduced cerebral cortex size.[64] Our deep learning model uncovered an unusually similar

nuclear MLF2-containing condensate phenotype between cells deficient in ZNF335 and

four out of five of the condensin II subunits (NCAPG2, NCAPH2, NCAPD3, and SMC2;

Figures 3B, 4G), inspiring future functional studies at the intersection of MLF2,

condensin biology, and phase separation.

In conclusion, our study lays the foundation for exploring what cellular role

biomolecular condensates have in microcephaly or related neurodevelopmental

disorders (Figure 6B). We overcome the scarcity of generic phase-separation markers

by establishing MLF2 as a versatile condensate biomarker and report several chemical

modulators of aberrant condensates. This reinforces similar chemical-modulator

screening as a general strategy to identify therapeutic interventions across

condensatopathies. Our screen and analysis pipeline are readily adaptable for

screening campaigns targeting diverse condensate phenotypes, including stress

granules and other disease-relevant condensates. This study uncovers a connection

between altered condensate accumulation and neurodevelopmental disorders, inspiring

pharmacological strategies to correct aberrant condensates.

Limitations of the study

While our study highlights several chemical and genetic modulators of disease-linked

condensates, opportunities exist for further refinement. Pyrithione zinc markedly

resolved the proteotoxic condensate cargo in torsin-deficient cells, while complete

resolution of the underlying condensates was not observed at the tested concentration

(Figure S2I). Extended incubation times may result in condensate resolution, but

increased cytotoxicity limited this approach. Similarly, while our genetic screen

uncovered an enrichment of neurodevelopmentally-linked genes, redundant pathways

or those strictly requiring a neuronal context are likely to be underrepresented and we

filtered out conditions that resulted in severe cytotoxicity. Large-scale screening of

optimized compound libraries and testing genetic redundancy through combinatorial

perturbations will likely identify additional condensate modulators. Furthermore,

assessing the condensate burden associated with these neurodevelopmental conditions

would benefit from future studies in neuronal or in vivo systems.

A

Gene	Z-Score	Disorder	Gene	Z-Score	Disorder
RNF26	7.99	—	LGALS9C	3.29	—
WAC	6.31	Desanto-Shinawi Syndrome	NAPRT	3.27	—
ZNF335	5.46	Microcephaly	MAP3K1	3.25	46, XY sex reversal 6
NCAPG2	5.33	Khan-Khan-Katsanis Syndrome	C9orf114	3.23	Neurodevelopmental disorder with poor growth, seizures, and brain abnormalities
NCAPH2	4.88	Cardioencephalomyopathy, Myopia	ESPL1	3.21	—
ECT2	4.58	—	CAPN5	3.18	Vitreoretinopathy, neovascular inflammatory
ASRGL1	4.22	—	TOR1B	3.18	—
SETD1B	4.16	Intellectual Developmental Disorder with Seizures and Language Delay	POLD4	3.16	—
ASB6	4.15	—	EP300	3.16	Rubinstein-Taybi Syndrome 2, Menke-Hennekam Syndrome 2, Colorectal Cancer
FOXS1	4.08	—	PRPF40A	3.13	—
SIN3A	3.94	Witteveen-Kolk Syndrome (WITKOS)	CCDC157	3.13	—
BRSK2	3.93	Idiopathic Pulmonary Fibrosis Susceptibility*	ZBED8	3.11	—
EIF4A3	3.78	Robin Sequence with Cleft Mandible and Limb Anomalies	TMEM25	3.11	—
SPDL1	3.77	Severe Primary Microcephaly, Neonatal Death	GJA4	3.08	—
ZNF783	3.66	—	CNOT3	3.07	Intellectual developmental disorder with speech delay, autism, dysmorphic facies
MKI67	3.64	Long QT Syndrome	CD248	3.04	—
SLC29A2	3.6	Bipolar disorder susceptibility*	ACOD1	3.02	Lymphatic Malformation
VEGFC	3.57	Lymphatic malformation	BTBD18	3.01	—
ACY3	3.51	—	CIT	2.99	Primary Microcephaly
CDK5	3.42	Isolated Microcephaly, Nonsyndromic Microcephaly, Lissencephaly	PHF23	2.99	—
NCAPD3	3.4	Primary Microcephaly	AK2	2.98	Reticular Dysgenesis
PTGES	3.35	—	TCEA2	2.98	—
C9orf129	3.32	—	C5orf60	2.95	—
TUBB4B	3.29	Leber Congenital Amaurosis with Early-Onset Deafness	SHH	2.94	Holoprosencephaly, Schizencephaly, Solitary Median Maxillary Central Incisor Syndrome, Microphthalmia
ACTL7A	3.29	Spermatogenic failure; breast cancer susceptibility*	LDLRAD1	2.93	—
Microcephaly/Congenital Brain Malformations			**Other Neurodevelopmental Disorders**		

B

Figure 6, *Poch et al.*

Fig. 6. Targeting aberrant condensates as a therapeutic approach for neurodevelopmental disorders. (**A**) CRISPR KO hits enriched for genes implicated in neurodevelopmental disorders. Top 50 candidate genes (left columns), their significance score (Z-Score, middle columns), and associated conditions (right columns) are color-coded based on disorders. Rows labeled purple indicate established link to microcephaly/congenital brain malformations with dark gray indicating additional neurodevelopmental disorders. Genes denoted with an asterisk indicate evidence from genome-wide association studies. Gene relevance to disease was determined based on allelic variants obtained from OMIM and GTR. (**B**) Chemical and genetic modulation of aberrant condensates in neurodevelopmental disorders. Genes involved in the etiology of primary microcephaly and encephalopathies were highly enriched in our screen. Condensates can be modulated using small molecules or genetic perturbations. Targeting condensates could potentially lead to preserving neuronal homeostasis. Arrows indicate possible intervention points and potential outcomes.

491 **Supplementary Figures**

Figure S1, *Poch et al.*

Fig. S1. CondenScreen as a high-content condensate analysis pipeline (**A**) Schematic of the computational *CondenScreen*. Initial cell segmentation and foci detection uses custom *CellProfiler* script and while downstream statistical analysis is completed in *R*. (**B**) Distribution of cumulative hit counts from the Pharmakon 1760 library screen (n = 6 plates) summarized by plate (top left), row (top right), and column (bottom). (**C**) Spatial heatmap of total hits per well position across all plates. Color intensity reflects hit frequency, with darker purple indicating higher cumulative hit counts (a maximum of one hit per well was observed). (**D**) Comparison of hit detection using the Z-Score vs BZ-Score Normalization methods. A heatmap of the median (n = 6) of the Z-Score (top) vs the BZ-Score (bottom) per well. Color indicates score magnitude and direction (orange = negative, purple = positive, white ≈ 0; darker shading reflects larger absolute values). (**E**) Schematic of CondenScreen's built-in chemical structure hit triage workflow. (top) Using either the name or CAS numbers, the SMILES will be returned (PubChem/NCI CACTUS), followed by substructural pattern identification (SMARTS) flags using several industry alert flags, along with the tested cytotoxicity of each compound. (bottom) Example results using plumbagin. Three alerts were flagged for plumbagin, including a quinone, ketone, and Michael acceptor substructural elements. As quinones are listed as a PAINS substructure, plumbagin received a risk category ranking of "Elevated".

Figure S2, *Poch et al.*

Fig. S2. Validation of chemical hits from the Pharmakon 1760 screen. (**A**) Calculated IC_{50} values of additional compounds with dose-dependent relationship. The black and red curves represent % effect and % toxicity, respectively, relative to the DMSO control. (**B**) Drug treatment decreases nuclear MLF2-GFP accumulation. Torsin-deficient cells expressing MLF2-GFP (8 h) were treated at four different concentrations (2x below IC_{50}, at IC_{50}, 2x above IC_{50}, and 8x above IC_{50}) for 3 h with either PZ, celastrol, plumbagin, or disulfiram, compared against the DMSO control. Each dot represents the number of MLF2-GFP foci for a single cell, while the horizontal lines indicate the median foci/condition. (**C**) Distribution of FACS results for 4TorKO MLF2-GFP-P2A-tdTomato HeLa cell sorting. Cells within quadrant 2 were sorted. (**D**) Quantification of the number of MLF2-GFP foci/cell (**E**) and tdTomato integrated intensity/cell. Cells (4TorKO) were all (except the untreated condition) induced with Dox and treated with either DMSO, cycloheximide (CHX), celastrol, or PZ. (**F**). Binned CNS value per cell across treatment conditions. Orange indicates <2x the median DMSO CNS value, red indicates 2–5x and purple indicates >5x the median DMSO control. Quantification (**G**) and confocal imaging (**H**) show that PZ does not non-specifically dissolve physiological condensates. Celastrol and PZ reduced MLF2-GFP foci (green) without dissolving PML bodies (red), whereas 5% 1,6-hexanediol dissolved both. All three compounds decreased MLF2-GFP foci relative to DMSO; only 1,6-hexanediol reduced PML foci. PZ increased PML foci number, while celastrol had no significant effect. Scale bars, 5 µm. (**I**) TEM images of 4TorKO HeLa cells treated with either 0.1% DMSO or 2 µM PZ for 3 h. Insets show NE condensates between the inner and outer nuclear envelopes. Scale bars at 500 nm; inset scale bars at 100 nm. Asterisk, putative nuclear pores; black arrowhead, outer nuclear envelope; white arrow, inner nuclear envelope; C, cytoplasm; N, nucleus. Asterisks indicate Bonferroni-corrected p-values determined by Mann-Whitney U testing (**: $p <= 0.01$; ****: $p <= 0.0001$).

495
496

Figure S3, *Poch et al.*

Fig. S3. Zinc is sufficient for modulating nuclear condensates in 4TorKO cells. (**A**) TPEN negates the effect of PZ treatment. Confocal images of 4TorKO cells expressing MLF2-GFP (green) and stained with Hoechst (nucleus, blue). Cells were treated with either DMSO, 6 µM TPEN, 2 µM PZ, or 6 µM TPEN + 2 µM PZ (3 h). (**B**) Barplot quantification of (A) showing the average number of MLF2-GFP foci per cell for each condition. (**C**) Zinc sulfate is sufficient to modulate nuclear condensates. Dox-induced 4TorKO were treated with either DMSO, 2 µM PZ, 500 µM ZnSO4, or 2 mM ZnSO4. Representative confocal images demonstrating distribution of MLF2-GFP, tdTomato, and DAPI are shown by the green, red, and blue channels, respectively. The corresponding barplot quantifications of each condition are shown representing (**D**) the average number of MLF2-GFP foci per cell (**E**) the average tdTomato integrated intensity per cell, and (**F**) Binned CNS value per cell across treatment conditions. Orange indicates <2x the median DMSO CNS value, red indicates 2–5x and purple indicates >5x the median DMSO control. Scale bars, 5 µm. Asterisks indicate Bonferroni corrected p-values using a Mann-Whitney U test (*: p <= 0.05; **: p <= 0.01; ***: p <= 0.001; ****: p <= 0.0001).

Figure S4, *Poch et al.*

Fig. S4. Validation and enriched gene pathways of CRISPR KO hits that promote condensate accumulation. (A)
Enrichment of pathways related to chromosomal and chromatid regulation. Upset plot depicting relationships between the top 30
GO-enriched biological process pathways. Horizontal bars (left) indicate number of genes for each GO pathway, with terms
ranked by gene count. Vertical bars (top) indicate size of each intersecting gene set of genes sharing multiple GO terms. The dot
matrix below indicates shared genes among the corresponding GO terms (black dots connected by black line); single dots
represent GO terms with unique gene sets. Red dots highlight pathways related to chromosomal or chromatid-related activities,
(**B**) Manual clustering of GO biological pathways from the genome-wide KO screen in WT HeLa. The top 30 enriched pathways
for the 100 highest-scoring candidate genes were manually clustered within five categories. Percentage represents the fraction of
GO pathways within each cluster. (**C**) Chord diagram mapping each gene to its specific GO terms. Each gene (outer labels, top) is
linked to its corresponding GO terms (bottom labels), with colored ribbons denoting relationships. (**D**) Binned quantification of the
MLF2-GFP foci count after CRISPR KO of select candidate genes. RNF26 KO, ZNF335 KO, and WAC KO HeLa cells were
compared against NT control. Bars indicate percent of cells with 5-15 foci MLF2-GFP foci/cell (orange), 15-30 foci (violet), or >30
foci (dark purple).

Figure S5, *Poch et al.*

Fig. S5. Validation of deep learning model to detect subtle variations in condensate phenotypes. (**A**) Schematic workflow of the deep learning-based condensate phenotype analysis pipeline. The model was trained on raw images from the CRISPR screen, which were resized to single-cell patches and labeled. A residual convolutional neural network (ResNet18) was optimized to distinguish variations in condensate morphology. Centroid feature vectors from the top CRISPR hits (Z-score > 2) were extracted and mapped using UMAP and k-means clustering. (**B**) Training and validation loss curves based on Huber loss (nn.SmoothL1Loss). Loss of each epoch for both the training set (gray) and validation set (purple) are reported with a minimum loss (0.14) occurring at epoch 21. (**C**) Correlation scatter plot of validation set. Each dot represents the similarity score (-1 = RNF26 KO; 0 = NT Ctrl; +1 = ZNF335 KO) of a single cell from the validation set. The predicted value (y-axis) is compared against the true value (x-axis), with a Pearson correlation score of 0.76. (**D**) Elbow (scree) plot showing the within-cluster sum of squares across cluster numbers (k = 1-10). (**E**) Silhouette plot showing the silhouette coefficient (0.34, vertical dashed line) for three clusters. (**F**) Representative saliency maps prior to training (top, Epoch 0) vs post-training (Epoch 20). From left to right, non-targeting control cell (true label = 0, initial pred. = -0.45, final pred. = +0.02), RNF26 KO cell (true label = -1, initial pred. = +0.26, final pred. = -1.01), and ZNF335 KO cell (true label = +1, initial pred. = -0.46, final pred. = +0.84). (**G**) Boxplots of the hotspot count, mean size of hotspots, effective attribution area, and Shannon's Entropy across the single cells (dots). Asterisks indicate p-values using a Mann-Whitney U test (*: p <= 0.05; **: p <= 0.01).

Figure S6, *Poch et al.*

Fig. S6. Condensates resulting from CRISPR KO can be dissolved by 1,6-Hexanediol. (A) Fluorescent microscopy images of Cas9-integrated WT HeLa cells treated with sgRNAs against either NT control, TOR1A/B, RNF26, or ZNF335 (from left to right) for 72 h. Cells were then exposed to either DMSO (top), 5% 1,6-hexanediol (middle), or 5% 2,5-hexanediol (bottom) for 5 min. WT cells with ZNF335, RNF26, or TOR1A/TOR1B knocked out demonstrated dissolution of condensates marked by MLF2-GFP (green) only after exposure to 5% 1,6-hexanediol (middle), but not DMSO (top) or 5% 2,5-hexanediol (bottom) controls.

Figure 7, *Poch et al.*

Fig. S7. Pyrithione zinc modulates atypical nuclear condensates across genetic perturbations. Cas9-integrated WT HeLa cells treated with sgRNAs against either NT control, TOR1A/B, RNF26, or ZNF335 (from left to right) for 72 h. Cells were then induced with MLF2-GFP (6 h Dox) and treated with either 0.1% DMSO (top panel) or 2 μM PZ (bottom panel) for 3 h. Representative confocal images show MLF2-GFP (green) and DAPI (blue). Scale bars, 5 μm.

523
524

Materials and Methods

Tissue culture

Adherent cultured cell lines (HeLa and HEK293T) were maintained in Dulbecco's

modified Eagle's medium (DMEM) supplemented with 10% (vol/vol) heat-inactivated

fetal bovine serum (Thermo Fisher Scientific) and 100 units (U) /mL penicillin-

streptomycin (Thermo Fisher Scientific). Cells were maintained in either T75 flasks or 10

cm dishes at 37℃ (95% humidity; 5% CO_2) and passaged every third day. For splitting,

the culture medium was removed, washed once with 1x PBS (Gibco), trypsinized with

0.25% trypsin-EDTA (Gibco), and transferred to new dishes (diluted 1:5 or 1:10).

Plasmid Constructs

Unless otherwise stated, all plasmid constructs were subcloned into a pcDNA™3.1 (+)

Mammalian Expression Vector (Invitrogen, V79020) using standard Gibson Assembly

protocols. The cDNA for the MLF2-GFP-P2A-tdTomato construct (MLF2 GenBank

accession no. BC000898) was subcloned into the pRetroX-Tight-Pur-GOI (gene of

interest) vector (Takara Bio) and co-transfected with a pRetroX-Tet-On advanced

inducible expression construct (Takara Bio). Viral constructs were co-transfected with

both a VSV-G viral envelope plasmid (Addgene #12259) and an MMLV gag/pol viral

packaging plasmid. For generation of HeLa cells expressing Cas9-mCherry, an mCherry

tag was inserted into the lentiviral plasmid containing Cas9-Flag (Addgene #52962)

immediately after the Flag tag into the BamHI cloning site.

Viral Production

Lentivirus production was carried out as previously described.[86,87] Briefly, to generate

Cas9-flag lentivirus, low-passage HEK293T cells were plated in a 10 cm dish with a

density of $3.5x10^6$ cells per dish. Medium was replaced the next day with 6 mL fresh

(antibiotic-free) DMEM and transfected with three plasmids encoding: psPAX2

(Addgene #12260) (2 µg), viral envelope protein VSV-G (1 µg), and lenti-Cas9-Flag-

mCherry (6 µg) using Lipofectamine 2000 (Thermo Fisher Scientific) at a 3:1 ratio of

DNA to Lipofectamine. After 72 h, supernatants containing the virus encoding Cas9-

mCherry were harvested and filtered through a 0.45 µm Steriflip filter unit

(MilliporeSigma), aliquoted into screw-cap tubes, and stored at -80°C or used for

transduction immediately. For the Dox-inducible MLF2-GFP-P2A-tdTomato system, two

separate retroviruses were produced using three plasmids encoding: MMLV gag/pol (2

µg), viral envelope protein VSV-G (1 µg), and either pRetroX-Tight-MLF2-GFP-P2A-

tdTomato or pRetroX-Tet-On (6 µg) using Lipofectamine 3000 (Thermo Fisher Scientific)

at a 2:1.5 ratio supplemented with Lipofectamine P3000 (2.5:1). After 72 h,

supernatants were collected, stored, or used, as described.

Generation of stable HeLa cell lines

The detailed protocol regarding the Tor1A/1B/2A/3A KO and integration of a Dox-

inducible MLF2-GFP reporter in HeLa cells has been extensively described by our lab

(Laudermilch et al. 2016; Rampello et al. 2020).[20,28] The Cas9-flag-mCherry HeLa and

Dox-inducible MLF2-GFP-P2A-tdTomato 4TorKO HeLa cell lines were generated by

seeding Dox-inducible MLF2-GFP WT HeLa or 4TorKO cells in six-well plates. The next

day, media was replaced with DMEM (antibiotic-free) supplemented with 4 µg/mL

polybrene (Sigma-Aldrich), and 100 µl of either the Cas9-Flag-mCherry virus (WT HeLa)

or both 100 µL of the MLF2-GFP-P2A-tdTomato and pRetro-Tet-On viruses (4TorKO)

added dropwise to wells. Medium was replaced 48 hours after transduction (complete

DMEM) and grown for another 24 hours. Cells were collected in sorting buffer (ice-cold

1x PBS, Gibco) and filtered through a 40 µm nylon mesh filter. Using a BD FACS Aria

III, cells were sorted as single cells by selecting those high in either mCherry (WT HeLa)

or MLF2-GFP/tdTomato (4TorKO, 16 h after 0.5 µg/mL Dox induction) fluorescence

intensity. A stringent gating strategy was applied to enrich for cells positive for the

respective fluorophores while excluding those with extremely high signal intensity, to

obtain a pure yet physiologically representative population.

High-content phenotypic screening strategy

All high-content screening was performed at Yale's Center for Molecular Discovery

(YCMD). The Horizon Discovery Edit-R synthetic sgRNA library was used in the whole

genome CRISPR KO screen, targeting 19,195 human genes, and comprising of the

Drug Targets library (3,683 genes, GA-004650-E2, Lot 20126), the Druggable Genome

library (4,643 genes, GA-004670-E2, Lot 20120), and the Human Genome library

(10,869 genes, GA-006500-E2, Lot 20130). Each screening plate included 16 replicates

of the negative control (NT sgRNA) and 16 replicates of the positive control (combined

Tor1A and Tor1B sgRNAs). To ensure robust KO, a pool of three sgRNAs targeting the

same gene at different regions was pre-plated into each arrayed well of 384-well glass-

bottom plates (Cellvis, #384-1.5H-N). Addition of sgRNA to screening plates was done

using the Labcyte Echo 550 Acoustic Dispenser (Labcyte, now Beckman) at a final

concentration of 25 nM.

Reverse transfection was performed using DharmaFECT1 Transfection Reagent

(Horizon/Dharmacon), which was diluted at 1:500 in Opti-MEM (Thermo Fisher

Scientific) and mixed with sgRNA. The mixture of sgRNA and transfection reagent was

incubated for 30 minutes, after which, 2,000 cells/well (WT HeLa integrated with

Cas9/MLF2-GFP) in antibiotic-free DMEM were added using a Thermo MultiDrop Combi

(Thermo Fisher Scientific). The plates were then incubated for 72 hours at 37°C. To

induce MLF2-GFP reporter expression, 0.5 µg/mL Dox was added to each well 16 h

before fixation. Media was removed and cells washed once with PBS and fixed with 4%

paraformaldehyde (PFA) for 20 minutes. Cells were washed twice with PBS,

permeabilized with 0.1% Triton X-100 for 5 minutes, and washed again with PBS. Cells

were stained with Hoechst 33342 dye (1:1000) (Invitrogen, H1399) and HCS CellMask

Deep Red Stain (Invitrogen, H32721) (1:5000 of 10 mg/mL, Thermo Fisher Scientific)

diluted in PBS for 30 minutes, and washed a final time in PBS. The Molecular Devices

ImageXpress 4.0 high-throughput fluorescent microscope (40x magnification) was used

to obtain 9-image sets (each consisting of DAPI, MLF2-GFP, and Deep Red CellMask

images) per well, resulting in 10,368 TIFF images/plate. Images were uploaded to

Yale's McCleary high-performance computing cluster (Yale Center for Research

Computing) for analysis.

For chemical screening, the Pharmakon 1760 drug library (MicroSource

Discovery Systems, Inc.) was used, consisting of 1360 FDA-approved drugs and 400

International drugs on six 384-well plates. Each screening plate consisted of 16

replicates of the negative control (4TorKO HeLa cells treated with DMSO [0.1% v/v])

and 16 replicates of the positive control (WT HeLa cells treated with DMSO [0.1% v/v]).

Plates were seeded with 4,000 HeLa cells/well (Multidrop Combi) in 20 μL of DMEM

before incubating at 37°C for 24 h. MLF2-GFP expression was induced by adding 0.5

μg/mL of Dox into each well 8 h prior to fixation. Compounds were resuspended in

DMSO and added to each well 2 h prior to fixation using a Labcyte Echo 550 Acoustic

Dispenser (Labcyte, now Beckman), at a final concentration of 10 μM. Cells were

washed, fixed, stained, and imaged using the methods outlined above.

Image segmentation of phenotypic screen data

We created a custom analysis pipeline to segment condensate foci into single cells

using the open-source software *CellProfiler 4.2.6.*[38,39] and downstream statistical

analysis using *R Studio* (Version 2023.12.1+402).[88] Individual nuclei were distinguished

by shape for objects between 30-100 (+4) pixels using an adaptive (40 pixels) Minimum

Cross-Entropy thresholding (0.01-1.0 threshold) in *CellProfiler 4*. The cytoplasm of each

nucleus was delineated as a secondary object of each nucleus using a watershed

algorithm with global Otsu thresholding (three classes, foreground; 0-1.0 threshold,

threshold correction factor = 0.7).[39,89] Each GFP-channel image was enhanced using a

white top-hat transform (speckles; maximum 10 pixels), allowing each condensate foci

between 1-10 pixels to be detected using adaptive (50 pixels) Otsu thresholding (three

classes, foreground; 0.0175-1.0 threshold, threshold correction factor = 1.2). Groups of

condensate foci were then distinguished using local intensity and related to their

corresponding cytoplasm/nuclei. The same image analysis pipeline was applied across

all imaging experiments, with only minor adjustments (e.g., cell size or foci intensity

thresholds). Any parameter changes were applied uniformly across all conditions and

controls within a given experiment.

643   For parallel analysis, a Bash script was developed allowing the execution of

multiple headless CellProfiler instances on a Simple Linux Utility for Resource

Management (SLURM)-managed compute cluster. Specifically, after constructing the

analysis pipeline in the CellProfiler GUI, the CreateBatchFiles module is used to

generate a batch configuration file that defines image metadata and processing

instructions. This batch file is then imported to the cluster environment, and all images

are registered within a single CellProfiler project in array format, with images from each

plate corresponding to a separate SLURM job. This approach enables autonomous

analysis of over 1 million high-content images. TIFF images—amounting to multiple

terabytes of data—within a matter of hours. Both the CellProfiler pipeline and Bash

scripts are open source and available publicly on the SchliekerLab GitHub page as part

of the CondenScreen pipeline (https://github.com/SchliekerLab/Condenscreen.git).

Statistical analysis of screen data

We developed a condensate screening using a combined *CellProfiler* and *R* pipeline,

CondenScreen, to statistically analyze all screen data. The *R*-script features an

interactive GUI allowing the user to select plate size (6-well to 384-well format), plate

layout, and input whether the data was taken from a signal "ON" or a signal "OFF"

screen. With limited user input, it then automatically iterates through images from each

plate and assigns groups of image sets into their corresponding wells using string

parsing. To control for variations in transfection efficiencies that could inadvertently

impact downstream scoring, the program has an optional filter to select only the top

25% of cells showing the highest (or lowest) signal using the modest assumption that at

least 1 out of 4 cells, on average, will have received the sgRNAs of interest. Additionally,

for both screens, the image with the highest score was filtered out, as there were

several conditions where one image-set had screening aberrations skewing the

condition toward a false positive result. For each plate, aggregate statistics of the

respective positive and negative controls are compared, including signal-to-background

(SB), standard deviation (SD) of the controls, and the coefficient of variation (CV) of the

controls, among others. To determine separation between positive and negative

controls, a Z' assay quality score was calculated: $Z' = 1 - (3(\sigma_{PC} + \sigma_{NC})/|\mu_{PC} -$

$\mu_{NC}|)$,[40] where σ_{PC} and σ_{NC} is the SD and μ_{PC} and μ_{NC} is the condensate mean of the

positive and negative controls, respectively.

The percent effect of each condition was quantified by either $E\% = \left(\frac{\mu - \overline{x}}{\mu}\right) \cdot 100$

(for signal "OFF" selection) or $E\% = \left(\frac{\overline{x} - \mu}{\mu}\right) \cdot 100$ (for signal "ON" selection) where $\overline{x}$ is

the average number of condensate foci/cell per tested condition, and μ is the average

number of condensate foci/cell across the screening population of the plate. The

significance scoring function is automatically changed depending on selection of a

signal "ON" vs a signal "OFF" screen, although this can be adjusted.

For signal "ON" screens, the significance of each condition was calculated using

the Z-Score to normalize for intraplate variations: $Z = (E\% - \mu_{E\%})/\sigma_{E\%}$, where $\mu_{E\%}$ and

$\sigma_{E\%}$ represent the population average percent effect and standard deviation,

respectively. Since signal "OFF" screens can often have increased variability, the

significance of each well was quantified by calculating the BZ-Score. First, B-Scores

were computed by applying Tukey's median polish to each plate to remove row, column,

and edge effects. The residuals from the median polish were then scaled by the median

absolute deviation (MAD) to produce a robust per-well B-score. Then, the BZ-Score was

calculated using the following formula: $BZ = (BScore - \mu_{BScore})/\sigma_{BScore}$, where μ_{BScore}

and σ_{BScore} are the mean and standard deviation of the B-Scores over the entire

plate.[90,91]

To determine how chemical or genetic perturbations alter cell viability, we derived

a function to extrapolate differential cellular health by morphological characterization of

the nucleus, which was used during validation experiments. Specifically, we added

several modules to *CellProfiler* to measure the size, shape, intensity, and texture of

each nucleus.[38] This is crucial as several compound concentrations had a non-

significant effect in changing the total cell count, but greatly disrupted nuclear

morphology based on visual inspection. After testing the effects of multiple parameters,

we derived the following empirical formula which best represented the data to estimate

nuclear health: $NucHealth = \Delta_{CC} + ((100 + \Delta_{CC})/100) \cdot \sum(\Delta_{Area} + \Delta_{FF})/2$, where Δ_{CC},

Δ_{Area}, and Δ_{FF} are the normalized difference in cell count, nuclear area, and nuclear

FormFactor (measurement of nuclear irregularity) compared to the screening population

(or DMSO for dose-response), respectively. Normalization was calculated according to:

$\Delta_v = ((V - \mu_V)/\mu_V) \cdot 100$, in which *V* is the variable of interest. The equation was derived

with the following assumptions and limitations. (1) Changes to the area and FormFactor

should be weighted equally, and (2) cell death should be given a higher impact weight

over morphological changes.

Two optional *CondenScreen* quality control features allow for flagging potential

false-positives or problematic hits arising from screening large libraries. In addition to

detecting spatial intra-plate bias, the program flags screen actives arising from global

systematic inter-plate spatial biases. These include shifts in continuous significance

score metrics (Z-Scores and BZ-Scores) and enrichment of binary hits across rows,

columns, individual well positions, and plates with a statistically enriched hit rate.

Continuous metric shifts are assessed using a Wilcoxon rank-sum (Mann–Whitney U)

test comparing each group to all other wells, while hit enrichment is assessed using

Fisher's exact test. P-values are Benjamini–Hochberg adjusted across groups and

flagged at an adjusted p-value < 0.05.

CondenScreen enables flagging potentially problematic chemical structures

using compound SMILES (Simplified Molecular Input Line Entry System), either

provided manually (recommended) or can attempt to be retrieved automatically from

PubChem[92] using the CAS number, with CACTUS[93] (NCI Chemical Identifier Resolver,

https://cactus.nci.nih.gov/chemical/structure) used as a fallback. Smiles are processed

using RDKit (via Reticulate in R), and compounds are evaluated against SMARTS

(SMILES Arbitrary Target Specification) substructures appearing in alert libraries

compiled from RDKit and ChEMBL resources. We utilized a pre-compiled list of these

structural alert filters from a csv document allowing SMARTS identification within the

RDKit environment, compiled by Pat Walters (see GitHub for more details:

https://github.com/PatWalters/rd_filters.git). We defined compounds as having an

"Elevated" risk if they include SMARTS overlapping with the Pan-Assay Interference

Compounds (PAINS) alerts,[94,95] NIH Molecular Libraries Small Molecule Repository

(MLSMR) alerts, or have a significant cytotoxicity (≥25% viability loss); "Moderate" risk if

overlapping with the Bristol-Myers Squibb (BMS), GlaxoSmithKline (GSK/Glaxo), Pfizer

LINT alerts, or the University of Dundee filter sets, or tdTomato reporter suppression

(≥25% decrease); and "Minor" risk if violating the Lipinski rule-of-five[96] or appearing in

the Inpharmatica or SureChEMBL medicinal chemistry alert libraries. Compounds

lacking these features were classified as "Clean." These categories are intended to

support chemical triage rather than exclude compounds.

CondenScreen provides additional downstream inter-plate and intra-plate

visualizations using open-source libraries, including *ggplot2*[97] and *ggplate*

(10.32614/CRAN.package.ggplate). The output includes a ranked list of hits, plots of the

screening distributions (raw condensate foci/cell and %Effect), a Z/BZ-score plot, and

individual plate heatmaps for each 384-well plate. Finally, top hits in both screens were

inspected manually to ensure quality control and rule out any additional potential

screening aberrations. Notably, manual inspection was rarely a reliable measure of hit

robustness. In several cases, hits that were initially overlooked during first-pass

screening (false negatives) were later validated and correctly identified by the program.

The *CondenScreen* platform is publicly available on the SchliekerLab GitHub page:

https://github.com/SchliekerLab/Condenscreen.git

Development of phenotypic machine learning model to distinguish condensate

phenotypes

We developed a supervised machine learning model to differentiate between subtle

phenotypic differences in nuclear condensates. The model was built around the

ResNet18 architecture[67] with pretrained initial starting weights borrowed from

ImageNet1k IMAGENET1K_V1[68] using torchvision[98] on Python 3.11.8 (conda-forge).

Using images directly from the CRISPR KO screen, 4,523 single cells were extracted,

cropped (100x100) and resized (224x224) before centering (DAPI-channel) each cell.

Single cells were labeled in accordance with their file names, with NT Ctrl cells = 0,

RNF26 KO cells = -1, and ZNF335 KO cells = +1. To ensure even sampling numbers

across conditions, cells from the overrepresented NT Ctrl conditioned were

downsampled to match KO conditions, with only the MLF2-GFP channel selected for

analysis. Data were split 80:20 into train and validation sets (sklearn.model_selection:

train_test_split)[99]. Tensors were packaged (torch.utils.data) before employing the

following parameters during model training (PyTorch[100]): AdamW (lr = 1e-4, weight

decay = 5e-2), dropout = 0.5, hidden dimension = 64, SmoothL1Loss,

CosineAnnealingLR, 100 max epochs, and early stopping (patience = 25). Model

performance was evaluated using several modules from sklearn.metrics and

scipy.stats,[99,101] including: mean_absolute_error(); r2_score(); pearsonr();

accuracy_score(); confusion_matrix(); roc_curve(); roc_auc_score(); and

ConfusionMatrixDisplay(). The integrated gradient[69] for the validation data was used for

post-hoc saliency mapping in order to ensure the model was learning biological features

compared to technical noise. Briefly, for each cell, an attribution map was calculated

using integrated gradients, which averages model gradients across 24 interpolation

steps between the input image and a zero baseline. This attribution map was min-

maxed normalized and converted to a single-cell 2D heatmap. For each cell, the spatial

distribution of saliency attribution (Shannon entropy;[102] heat_entropy), effective number

of pixels contributing to the saliency distribution (heat_effective_npix), and hotspot

number and mean size (heat_hotspot_components), where hotspots were defined as

connected components within the top 2% of saliency intensities. The per-cell

distributions for ZNF335 vs RNF26 cells were statistically assessed using a two-sided

Wilcoxon rank-sum (Mann–Whitney U) test.

After optimization, all CRISPR KO gene hits with a Z-Score > 2 were cropped,

resized, and centered before randomly downsampled to 500 cells per condition

(accounting for differences in cell viability). Cells were then analyzed by the optimized

model and their latent representation embeddings (512-dimensional latent embeddings

per cell, e.g., condensate morphology, size, intensity, etc.) were extracted. Relative

centroid vector scores were computed (mean embedding across cells), and KMeans

clustering with k = 3 was applied to UMAP (n_neighbors = 15, min_dist = 0.1)

coordinates to assign genes to discrete clusters. The optimal cluster number was

selected using the elbow method (KMeans, k = 1–10) and silhouette analysis

(sklearn.metrics: silhouette_samples | silhouette_score), yielding three distinct clusters.

The difference in the cosine similarity for each gene's centroid vector embedding was

compared with the three anchor centroids (RNF26/NT Ctrl/ZNF335) and exported to an

.xlsx file (see Table S3). A hierarchical heatmap was constructed using the top 50 genes

with highest similarity to the RNF26 and ZNF335 KO conditions, via pandas,

seaborn.clustermap, and matplotlib.pyplot. For a full list of modules imported and model

training parameters, please refer to our source script:

https://github.com/SchliekerLab/Condenscreen.git

Solubilization of hit compounds

All putative hit compounds were resuspended up to 100 mM in DMSO. The following

compounds were solubilized: pyrithione zinc (PZ) (CAS 13463-41-7), celastrol (CAS

34157-83-0), nerol (CAS 106-25-2), spiramycin (CAS 8025-81-8), choline chloride (CAS

67-48-1), disulfiram (CAS 97-77-8), bronopol (CAS 52-51-7), penfluridol (CAS 26864-

56-2), plumbagin (CAS 481-42-5), dyphylline (CAS 479-18-5), econazole nitrate (CAS

24169-02-6), ethacrynic acid (CAS 58-54-8), piroctone olamine (CAS 68890-66-4),

dihydrocelastryl (CAS 80-97-7), fenipentol (CAS 583-03-9), triclosan (CAS 3380-34-5),

eprodisate disodium (CAS 36589-58-9), L-phenylalanine (CAS 63-91-2), nabumetone

(CAS 42924-53-8), edrophonium chloride (CAS 116-38-1), josamycin (CAS 16846-24-

5), mevastatin (CAS 73573-88-3), clofibric acid (CAS 882-09-7), docosahexaenoic acid

(CAS 6217-54-5).

Dose-dependent validation of small molecules

Based on the change in MLF2-GFP foci, viability, and structural diversity, the 22 top-

scoring putative hit compounds were resuspended to 100 mM in DMSO. Using

analogous methods described in the "High-content screening strategy" section, a 12-

point dose response was constructed with concentrations ranging from < 100 nM to 100

µM of each compound, with a final working DMSO concentration of 0.1%. All

compounds were tested using four replicates, and 12-positive (WT HeLa) and 12-

negative (DMSO) controls were included on each screening plate to ensure assay

robustness. MLF2-GFP was induced with 0.5 µg/mL of Dox 8 h before fixation, and

compounds were added 3 h prior to fixation (5 h after initial Dox addition). Cells were

washed, fixed, stained, and imaged as described above. All dose-response images

were analyzed using an analogous *CellProfiler* + *R*-script pipeline.

825   The effect of each concentration was interpolated (*R*, *drc* & *ggplot*) using a four-

parameter log-logistic function for each compound: $f(x) = \min + (\max - \min)/(1 +$

$e^{h(\log(x) - \log(IC_{50}))})$, where x is the concentration, min represents the lower asymptote,

max represents the higher asymptote, IC_{50} is the concentration at which response is

50% of the way between min and max, and h is the hill coefficient.[97,103] The percent

toxicity was calculated by fitting the additive inverse of the nuclear health function

values to the same model.

Pathway enrichment analysis

Gene Ontology (GO) term enrichment analysis was used for the top 100 CRISPR gene

hits.[60] The gene/chemical hits were separately analyzed in *R* to determine enriched

biological process pathways using the enrichGO function (*clusterProfiler*).[104] Enriched

pathways were compared against an annotation database (org.Hs.eg.db). The top 30

pathways were then plotted using the pheatmap function and additionally plotted to an

interactive .html graphic using heatmaply.[105] Pathways were grouped by hierarchical

clustering using Euclidean distance with complete linkage. To explore the interactive

heatmap, please visit the [Schlieker Lab GitHub page](#) and download the corresponding

.html files under the "Interactive Figures" folder.

Immunofluorescence and confocal microscopy:

HeLa cells were grown on 13 mm round coverslips and fixed with 4% (v/v)

paraformaldehyde (Thermo Fisher Scientific), washed with 1x PBS (Gibco),

permeabilized with 0.1% (v/v) Triton X-100 (Sigma-Aldrich) for 5 minutes, and blocked

with a PBS mixture containing 2% (w/v) BSA and 2% (w/v) glycine. Primary antibodies

were diluted in blocking buffer and added dropwise to coverslips for a one-hour

incubation. After washing 3x with blocking buffer, fluorescent secondary antibodies

diluted in blocking buffer were added for an additional hour, followed by a 1x PBS wash.

Coverslips were then stained using a 1:2000 dilution of Hoechst 33342 (Life

Technologies). Cells were then washed 3x with PBS before mounting onto slides

(Fluoromount-G, Southern Biotech). Primary antibodies were diluted 1:500: anti-HA

produced in rat (11867423001, Roche), anti-K48 produced in rabbit (ZRB2150, Millipore

Sigma), and anti-Mab414 produced in mouse. Secondary antibodies were conjugated to

Alexa Fluor 568 (Life Technologies).

Epifluorescence images were captured using a Zeiss AXIO Observer Z1 inverted

fluorescence microscope using a ×63/1.40 oil objective. All confocal images were

acquired at Yale's Imaging Facility on Science Hill using a Zeiss LSM 880 laser

scanning confocal microscope with a C Plan-Apochromat ×63/1.40 oil objective. Images

were analyzed with *CondenScreen* and compiled for publication using Zeiss v3.4 and

Image J (Fiji) software.

Validation of hit compounds

Based on the dose-response results, we focused validation efforts on four compounds:

PZ (IC_{50} = 601.6 nM), celastrol (IC_{50} = 124.8 nM), disulfiram (IC_{50} = 10.8 µM), and

plumbagin (IC_{50} = 7.41 µM). To determine the optimal concentrations using a tighter set

of concentrations, MLF2-GFP expression was induced in 4TorKO HeLa cells (0.5 µg/mL

Dox, 8 h) and treated with either DMSO or one of four different

concentrations/compounds: 2x below the IC_{50}, IC_{50}, 2x above the IC_{50}, or 8x above the

IC_{50}. Compounds were added 3 h before fixation. For each condition, images (n > 50

cells/condition) were captured on a Zeiss AXIO Observer Z1 inverted fluorescence

microscope and analyzed using our *CellProfiler* pipeline. *R Studio* was used to cluster

MLF2-GFP foci into single cells grouped by treatment condition and construct jittered

boxplots. To determine the effect of compounds on K48-ubiquitinylated proteins

accumulation within condensates, we incubated cells with compounds for 3h (PZ, 3 µM;

celastrol, 3 µM; plumbagin, 15 µM; and disulfiram, 50 µM), followed by

immunofluorescent labeling with α-K48 primary antibody (1:500). Images were captured

and analyzed as described above. High-resolution representative images were acquired

using confocal microscopy (Zeiss LSM 880).

To determine differential protein expression, we constructed a 4TorKO HeLa cell

line with a Dox inducible MLF2-GFP-P2A-tdTomato marker, where tdTomato serves as

a diffuse secondary reporter for global protein synthesis. Cells were induced with 0.5

µg/mL Dox and treated with their respective compound for 6 h before fixation. Imaging

was carried out on the Zeiss AXIO and the Zeiss LSM 880 for quantification and high-

resolution, respectively. The integrated intensity of both GFP and tdTomato signal was

quantified (n > 50) with *CellProfiler* and *R*. To ascertain the relative influence of global

protein expression inhibition on the reduced number of MLF2-GFP foci, we derived the

following empirical equation to determine the condensate normalization score ($CNS =$

$(1 - (\overline{F}_x/\overline{F}_{DMSO})) \cdot (\overline{II}_x/\overline{II}_{DMSO})^2$, where $\overline{F}$ and $\overline{II}$ are the average number of NE

condensate foci and the average integrated intensity of the secondary marker,

respectively, for either the tested condition (x) or the DMSO control. The scoring system

is designed to prioritize compounds that robustly reduce condensate foci without

affecting tdTomato reporter expression and penalizes those that have little effect on foci

or substantially suppress reporter expression. It penalizes modulators that have either

little effect on foci or those that substantially suppress reporter expression. The

weighting is particularly sensitive to changes in tdTomato intensity due to the squared

exponent, thereby strongly penalizing compounds which impair protein expression. A

low CNS-score suggests non-specific or off-target activities, such as decreased

transcription, translation, or increased biomarker turnover.

Guide RNA-mediated KO experiments:

Each gene KO consisted of a pool of 3 distinct, synthetic sgRNAs, each targeting a

different portion of the same gene (Horizon Discovery). Gene knock-out was conducted

using either DharmaFECT or DharmaFECT Duo Transfection Reagent (Horizon

discovery) using Tor1A KO + Tor1B KO as a positive control and NT sgRNAs as a

negative control (Horizon discovery). The sgRNAs were transfected into stably-

integrated Cas9 expressing HeLa cells and incubated for 72 h with a final sgRNA

concentration of 25 nM. The cells were treated with 0.5 µg/mL Dox to induce the MLF2-

GFP expression 6 h before fixation.

The pooled sgRNA sequences used were: ZNF335 sgRNAs (SG-016173,

Horizon Discovery), GATGATGGTGGCGCCGCCCA, GTGCCAGGGTGATCTGCGGT, ACCAAATGGACCAGGTGCCA;

RNF26 sgRNAs (SG-007060, Horizon Discovery), TATCATGGAGGCAGTGTACC,

GCATGACATCCTCTCGAAGA, CAGTGTACCTGGTAGTGAAT; TOR1A sgRNAs (SG-011023, Horizon

Discovery) TATGAACATGGCTTTCTGGT, TATGAAGATGGACCTCGCAC, AGTGCAATGTGGCCACAAAC; and

TOR1B sgRNAs (SG-014203-01, Horizon Discovery), AGTGCAGAGTCGATACAAAC,

CAAGAGCCGTCTTAGTTATA, TTGGCGGGCCGGAAGAAAGA.

Sodium arsenite and 1,6-hexanediol treatment

To determine the co-localization of MLF2 into stress granules, Lipofectamine 2000 was

used to co-transfect MLF2 with either G3BP1-GFP (Addgene #135997) + MLF2-HA,

TDP-43-tdTomato-HA (Addgene #28205) + MLF2-GFP-Flag (Addgene #230986), or

pcDNA3.1 empty vector control. Next day (16 h post-transfection), HeLa cells were

treated with $NaAsO_2$ (0.5 mM) and incubated for 30 min. Cells were fixed for 20 min with

PFA, stained, and imaged as outlined. For hexanediol treatment, cells were incubated in

complete DMEM medium containing 5% (v/v) 1,6-hexanediol (CAS 629-11-8) or 5%

(v/v) 2,5-hexanediol (CAS 2935-44-6) for 5 min before fixation and immunofluorescence

staining as described.

Electron Microscopy:

Electron microscopy was performed at the Center for Cellular and Molecular Imaging,

Yale School of Medicine, using our previously described workflow (Laudermilch et al.,

2016).[28] Briefly, cells were fixed for 1 h in 2.5% glutaraldehyde in 0.1 M sodium

cacodylate buffer, pH 7.4. Cells were rinsed and scraped in 1% gelatin and centrifuged

in a 2% agar solution. Chilled cell blocks were processed with osmium and

thiocarbohydrazide-osmium ligand as described (West et al., 2010), and samples were

embedded in Durcupan ACM resin (Electron Microscopy Science). Polymerization was

performed by incubating samples at 60°C overnight. These blocks were cut into 60 nm

sections with a Leica UltraCut UC7 and stained with 2% uranyl acetate and lead citrate

on Formvar/carbon-coated grids. Samples were visualized with an FEI Tecnai Biotwin

TEM at 80 kV, and pictures were taken with Morada CCD and iTEM (Olympus)

software.

Statistical Analysis:

Genetic and chemical hits were ranked based on their Z-Score or BZ-Score,

respectively, as described in detail within the *Statistical analysis of screen data* section.

GO pathways were ranked using a p-value cutoff of 0.05, after controlling for false

discovery rate using the Benjamini-Hochberg (BH) procedure. The significance of all

chemical and genetic validation experiments was assessed using a Bonferroni multiple

comparisons corrected Mann–Whitney U test with > 50 cells per condition individually

analyzed. Unless otherwise stated, asterisks indicate Bonferroni corrected p-values

using a Mann-Whitney U test where *: $p <= 0.05$; **: $p <= 0.01$; ***: $p <= 0.001$; and ****:

$p <= 0.0001$.

Acknowledgements

Funding:

This work was funded by the DOD PR200788 (C.S.) awarded by the U.S. Department

of Defense, Department of the Army, along with support from the Dystonia Medical

Research Foundation. Additionally, this work was supported by the NIH T32 GM145469

(D.P.), the European Molecular Biology Organization (EMBO) ALTF 910-2022 (E.F.F.K),

and the Dutch Research Council (NWO) 019.222EN.007 (E.F.E.K.) This work utilized

ImageXpress Micro 4 high content imager that was purchased with funding from a

National Institutes of Health SIG grant 1S10OD032384. We thank Morven Graham and

the Center for Cellular and Molecular Imaging (CCMI) Electron Microscopy Facility for

the collection of TEM images and Shawn Ferguson for sharing stress granule marker

reagents. The funders had no role in study design, data collection and analysis,

decision to publish or preparation of the manuscript.

Author contributions:

Conceptualization: DP, CM, EFEK, YVS, CS
Methodology: DP, CM, SM, EFEK, VT, ND
Investigation: DP, CM, SM, EFEK, VT, ND, CS
Computational Pipeline: DP, VT
Machine Learning: DP
Data Analysis: DP
Supervision: CS
Writing—original draft: DP
Writing—review & editing: DP, CS, EFEK, CM, SM, ND, YVS

Competing interests:

Authors declare that they have no competing interests.

Data and materials availability:

All data are available in the main text or the supplementary materials. The

computational *CondenScreen* pipeline and machine learning source code are available

on the Schlieker Lab GitHub page:https://github.com/SchliekerLab/Condenscreen.git .

988   At the time of publication, all raw screening images will be uploaded to the EMBL-EBI

989   BioImage Archive: https://www.ebi.ac.uk/bioimage-archive/.

1000

1001

1002

1003

1004

1005

1006

1007

1008

1009

1010

1011

1012

1013

1014

1015

1016

1017

1018

1019

References

1. Prophet, S.M., Rampello, A.J., Niescier, R.F., Gentile, J.E., Mallik, S., Koleske, A.J.,
and Schlieker, C. (2022). Atypical nuclear envelope condensates linked to
neurological disorders reveal nucleoporin-directed chaperone activities. Nat. Cell
Biol. *24*, 1630–1641. https://doi.org/10.1038/s41556-022-01001-y.

2. Mackenzie, I.R., Nicholson, A.M., Sarkar, M., Messing, J., Purice, M.D., Pottier, C.,
Annu, K., Baker, M., Perkerson, R.B., Kurti, A., et al. (2017). TIA1 Mutations in
Amyotrophic Lateral Sclerosis and Frontotemporal Dementia Promote Phase
Separation and Alter Stress Granule Dynamics. Neuron *95*, 808-816.e9.
https://doi.org/10.1016/j.neuron.2017.07.025.

3. Alberti, S., and Dormann, D. (2019). Liquid-Liquid Phase Separation in Disease.
Annu. Rev. Genet. *53*, 171–194. https://doi.org/10.1146/annurev-genet-112618-
043527.

4. Kedersha, N., Stoecklin, G., Ayodele, M., Yacono, P., Lykke-Andersen, J., Fritzler,
M.J., Scheuner, D., Kaufman, R.J., Golan, D.E., and Anderson, P. (2005). Stress
granules and processing bodies are dynamically linked sites of mRNP remodeling.
J. Cell Biol. *169*, 871–884. https://doi.org/10.1083/jcb.200502088.

5. Portz, B., Lee, B.L., and Shorter, J. (2021). FUS and TDP-43 Phases in Health and
Disease. Trends Biochem. Sci. *46*, 550–563.
https://doi.org/10.1016/j.tibs.2020.12.005.

6. Alberti, S., Gladfelter, A., and Mittag, T. (2019). Considerations and Challenges in
Studying Liquid-Liquid Phase Separation and Biomolecular Condensates. Cell *176*,
419–434. https://doi.org/10.1016/j.cell.2018.12.035.

7. Mittag, T., and Pappu, R.V. (2022). A conceptual framework for understanding
phase separation and addressing open questions and challenges. Mol. Cell *82*,
2201–2214. https://doi.org/10.1016/j.molcel.2022.05.018.

8. Brangwynne, C.P., Eckmann, C.R., Courson, D.S., Rybarska, A., Hoege, C.,
Gharakhani, J., Jülicher, F., and Hyman, A.A. (2009). Germline P granules are
liquid droplets that localize by controlled dissolution/condensation. Science *324*,
1729–1732. https://doi.org/10.1126/science.1172046.

9. Schmidt, H.B., and Görlich, D. (2016). Transport Selectivity of Nuclear Pores, Phase
Separation, and Membraneless Organelles. Trends Biochem. Sci. *41*, 46–61.
https://doi.org/10.1016/j.tibs.2015.11.001.

10. Li, P., Banjade, S., Cheng, H.-C., Kim, S., Chen, B., Guo, L., Llaguno, M.,
Hollingsworth, J.V., King, D.S., Banani, S.F., et al. (2012). Phase transitions in the
assembly of multivalent signalling proteins. Nature *483*, 336–340.
https://doi.org/10.1038/nature10879.

11. Wolozin, B., and Ivanov, P. (2019). Stress granules and neurodegeneration. Nat.
Rev. Neurosci. *20*, 649–666. https://doi.org/10.1038/s41583-019-0222-5.

12. Alberti, S., and Hyman, A.A. (2021). Biomolecular condensates at the nexus of
cellular stress, protein aggregation disease and ageing. Nat. Rev. Mol. Cell Biol.
22, 196–213. https://doi.org/10.1038/s41580-020-00326-6.

13. Yan, X., Kuster, D., Mohanty, P., Nijssen, J., Pombo-García, K., Morato, J.G.,
Rizuan, A., Franzmann, T.M., Sergeeva, A., Ly, A.M., et al. (2025). Intra-
condensate demixing of TDP-43 inside stress granules generates pathological
aggregates. Cell *188*. https://doi.org/10.1016/j.cell.2025.04.039.

14. Vendruscolo, M., and Fuxreiter, M. (2022). Protein condensation diseases:
therapeutic opportunities. Nat. Commun. *13*, 5550. https://doi.org/10.1038/s41467-
022-32940-7.

15. Ng, S.C., and Görlich, D. (2022). A simple thermodynamic description of phase
separation of Nup98 FG domains. Nat. Commun. *13*, 6172.
https://doi.org/10.1038/s41467-022-33697-9.

16. Kozai, T., Fernandez-Martinez, J., van Eeuwen, T., Gallardo, P., Kapinos, L.E.,
Mazur, A., Zhang, W., Tempkin, J., Panatala, R., Delgado-Izquierdo, M., et al.
(2023). Dynamic molecular mechanism of the nuclear pore complex permeability
barrier. bioRxiv, 2023.03.31.535055. https://doi.org/10.1101/2023.03.31.535055.

17. Cowburn, D., and Rout, M. (2023). Improving the hole picture: towards a consensus
on the mechanism of nuclear transport. Biochem. Soc. Trans. *51*, 871–886.
https://doi.org/10.1042/BST20220494.

18. Frey, S., Richter, R.P., and Görlich, D. (2006). FG-rich repeats of nuclear pore
proteins form a three-dimensional meshwork with hydrogel-like properties. Science
314, 815–817. https://doi.org/10.1126/science.1132516.

19. Kim, S., Phan, S., Tran, H.T., Shaw, T.R., Shahmoradian, S.H., Ellisman, M.H.,
Veatch, S.L., Barmada, S.J., Pappas, S.S., and Dauer, W.T. (2024). TorsinA is
essential for neuronal nuclear pore complex localization and maturation. Nat. Cell
Biol. *26*, 1482–1495. https://doi.org/10.1038/s41556-024-01480-1.

20. Rampello, A.J., Laudermilch, E., Vishnoi, N., Prophet, S.M., Shao, L., Zhao, C.,
Lusk, C.P., and Schlieker, C. (2020). Torsin ATPase deficiency leads to defects in
nuclear pore biogenesis and sequestration of MLF2. J. Cell Biol. *219*.
https://doi.org/10.1083/jcb.201910185.

21. Fan, Y., Si, Z., Wang, L., and Zhang, L. (2023). DYT-TOR1A dystonia: an update on
pathogenesis and treatment. Front. Neurosci. *17*, 1216929.
https://doi.org/10.3389/fnins.2023.1216929.

22. Balint, B., Mencacci, N.E., Valente, E.M., Pisani, A., Rothwell, J., Jankovic, J.,
Vidailhet, M., and Bhatia, K.P. (2018). Dystonia. Nat. Rev. Dis. Primer *4*, 1–23.
https://doi.org/10.1038/s41572-018-0023-6.

23. Ozelius, L.J., Hewett, J.W., Page, C.E., Bressman, S.B., Kramer, P.L., Shalish, C.,
de Leon, D., Brin, M.F., Raymond, D., Corey, D.P., et al. (1997). The early-onset
torsion dystonia gene (DYT1) encodes an ATP-binding protein. Nat. Genet. *17*, 40–
48. https://doi.org/10.1038/ng0997-40.

24. Liang, C.-C., Tanabe, L.M., Jou, S., Chi, F., and Dauer, W.T. (2014). TorsinA
hypofunction causes abnormal twisting movements and sensorimotor circuit
neurodegeneration. J. Clin. Invest. *124*, 3080–3092.
https://doi.org/10.1172/JCI72830.

25. Zhao, C., Brown, R.S.H., Chase, A.R., Eisele, M.R., and Schlieker, C. (2013).
Regulation of Torsin ATPases by LAP1 and LULL1. Proc. Natl. Acad. Sci. U. S. A.
110, E1545–E1554. https://doi.org/10.1073/pnas.1300676110.

26. Brown, R.S.H., Zhao, C., Chase, A.R., Wang, J., and Schlieker, C. (2014). The
mechanism of Torsin ATPase activation. Proc. Natl. Acad. Sci. U. S. A. *111*,
E4822–E4831. https://doi.org/10.1073/pnas.1415271111.

27. Demircioglu, F.E., Sosa, B.A., Ingram, J., Ploegh, H.L., and Schwartz, T.U. (2016).
Structures of TorsinA and its disease-mutant complexed with an activator reveal
the molecular basis for primary dystonia. eLife *5*, e17983.
https://doi.org/10.7554/eLife.17983.

28. Laudermilch, E., Tsai, P.-L., Graham, M., Turner, E., Zhao, C., and Schlieker, C.
(2016). Dissecting Torsin/cofactor function at the nuclear envelope: a genetic
study. Mol. Biol. Cell *27*, 3964–3971. https://doi.org/10.1091/mbc.E16-07-0511.

29. Goodchild, R.E., Kim, C.E., and Dauer, W.T. (2005). Loss of the Dystonia-
Associated Protein TorsinA Selectively Disrupts the Neuronal Nuclear Envelope.
Neuron *48*, 923–932. https://doi.org/10.1016/j.neuron.2005.11.010.

30. Kuiper, E.F.E., Gallardo, P., Bergsma, T., Mari, M., Kolbe Musskopf, M., Kuipers, J.,
Giepmans, B.N.G., Steen, A., Kampinga, H.H., Veenhoff, L.M., et al. (2022). The
chaperone DNAJB6 surveils FG-nucleoporins and is required for interphase
nuclear pore complex biogenesis. Nat. Cell Biol. *24*, 1584–1594.
https://doi.org/10.1038/s41556-022-01010-x.

31. Martin, E.W., Holehouse, A.S., Peran, I., Farag, M., Incicco, J.J., Bremer, A., Grace,
C.R., Soranno, A., Pappu, R.V., and Mittag, T. (2020). Valence and patterning of
aromatic residues determine the phase behavior of prion-like domains. Science
367, 694–699. https://doi.org/10.1126/science.aaw8653.

32. Colombrita, C., Zennaro, E., Fallini, C., Weber, M., Sommacal, A., Buratti, E., Silani,
V., and Ratti, A. (2009). TDP-43 is recruited to stress granules in conditions of

oxidative insult. J. Neurochem. *111*, 1051–1061. https://doi.org/10.1111/j.1471-
4159.2009.06383.x.

33. Palangi, F., Samuel, S.M., Thompson, I.R., Triggle, C.R., and Emara, M.M. (2017).
Effects of oxidative and thermal stresses on stress granule formation in human
induced pluripotent stem cells. PLOS ONE *12*, e0182059.
https://doi.org/10.1371/journal.pone.0182059.

34. Fang, M.Y., Markmiller, S., Vu, A.Q., Javaherian, A., Dowdle, W.E., Jolivet, P.,
Bushway, P.J., Castello, N.A., Baral, A., Chan, M.Y., et al. (2019). Small-Molecule
Modulation of TDP-43 Recruitment to Stress Granules Prevents Persistent TDP-43
Accumulation in ALS/FTD. Neuron *103*, 802-819.e11.
https://doi.org/10.1016/j.neuron.2019.05.048.

35. Uechi, H., Sridharan, S., Nijssen, J., Bilstein, J., Iglesias-Artola, J.M., Kishigami, S.,
Casablancas-Antras, V., Poser, I., Martinez, E.J., Boczek, E., et al. (2025). Small-
molecule dissolution of stress granules by redox modulation benefits ALS models.
Nat. Chem. Biol. *21*, 1–12. https://doi.org/10.1038/s41589-025-01893-5.

36. Bergsma, T., Steen, A., Kamenz, J.L., Otto, T.A., Gallardo, P., and Veenhoff, L.M.
(2024). Imaging-based quantitative assessment of biomolecular condensates
in vitro and in cells. J. Biol. Chem. *301*, 108130.
https://doi.org/10.1016/j.jbc.2024.108130.

37. Baggett, D.W., Medyukhina, A., Tripathi, S., Shirnekhi, H.K., Wu, H., Pounds, S.B.,
Khairy, K., and Kriwacki, R. (2022). An Image Analysis Pipeline for Quantifying the
Features of Fluorescently-Labeled Biomolecular Condensates in Cells. Front.
Bioinforma. *2*. https://doi.org/10.3389/fbinf.2022.897238.

38. Stirling, D.R., Swain-Bowden, M.J., Lucas, A.M., Carpenter, A.E., Cimini, B.A., and
Goodman, A. (2021). CellProfiler 4: improvements in speed, utility and usability.
BMC Bioinformatics *22*, 433. https://doi.org/10.1186/s12859-021-04344-9.

39. Carpenter, A.E., Jones, T.R., Lamprecht, M.R., Clarke, C., Kang, I.H., Friman, O.,
Guertin, D.A., Chang, J.H., Lindquist, R.A., Moffat, J., et al. (2006). CellProfiler:
image analysis software for identifying and quantifying cell phenotypes. Genome
Biol. *7*, R100. https://doi.org/10.1186/gb-2006-7-10-r100.

40. Zhang, J.-H., Chung, T.D.Y., and Oldenburg, K.R. (1999). A Simple Statistical
Parameter for Use in Evaluation and Validation of High Throughput Screening
Assays. J. Biomol. Screen. *4*, 67–73.
https://doi.org/10.1177/108705719900400206.

41. Liu, Z., Chen, O., Wall, J.B.J., Zheng, M., Zhou, Y., Wang, L., Ruth Vaseghi, H.,
Qian, L., and Liu, J. (2017). Systematic comparison of 2A peptides for cloning
multi-genes in a polycistronic vector. Sci. Rep. *7*, 2193.
https://doi.org/10.1038/s41598-017-02460-2.

42. Kroschwald, S., Maharana, S., and Simon, A. (2017). Hexanediol: a chemical probe
to investigate the material properties of membrane-less compartments. Matters.
https://doi.org/10.19185/matters.201702000010.

43. Wang, W.-M., Liu, Z., Liu, A.-J., Wang, Y.-X., Wang, H.-G., An, D., Heng, B., Xie, L.-
H., Duan, J.-L., and Liu, Y.-Q. (2015). The Zinc Ion Chelating Agent TPEN
Attenuates Neuronal Death/apoptosis Caused by Hypoxia/ischemia Via Mediating
the Pathophysiological Cascade Including Excitotoxicity, Oxidative Stress, and
Inflammation. CNS Neurosci. Ther. *21*, 708–717.
https://doi.org/10.1111/cns.12428.

44. Martin, H.L., Adams, M., Higgins, J., Bond, J., Morrison, E.E., Bell, S.M., Warriner,
S., Nelson, A., and Tomlinson, D.C. (2014). High-Content, High-Throughput
Screening for the Identification of Cytotoxic Compounds Based on Cell Morphology
and Cell Proliferation Markers. PLOS ONE *9*, e88338.
https://doi.org/10.1371/journal.pone.0088338.

45. Amberger, J.S., Bocchini, C.A., Schiettecatte, F., Scott, A.F., and Hamosh, A.
(2015). OMIM.org: Online Mendelian Inheritance in Man (OMIM®), an online
catalog of human genes and genetic disorders. Nucleic Acids Res. *43*, D789–
D798. https://doi.org/10.1093/nar/gku1205.

46. Rubinstein, W.S., Maglott, D.R., Lee, J.M., Kattman, B.L., Malheiro, A.J., Ovetsky,
M., Hem, V., Gorelenkov, V., Song, G., Wallin, C., et al. (2013). The NIH genetic
testing registry: a new, centralized database of genetic tests to enable access to
comprehensive information and improve transparency. Nucleic Acids Res. *41*,
D925–D935. https://doi.org/10.1093/nar/gks1173.

47. Patel, D.D., Gripp, K.W., Wadman, E., Mishra, I., and Kandula, V. (2024). Novel
biallelic ZNF335 variant causing primary microcephaly: A case report and
radiological review. Am. J. Med. Genet. A. *194*, e63593.
https://doi.org/10.1002/ajmg.a.63593.

48. Yang, Y.J., Baltus, A.E., Mathew, R.S., Murphy, E.A., Evrony, G.D., Gonzalez, D.M.,
Wang, E.P., Marshall-Walker, C.A., Barry, B.J., Murn, J., et al. (2012).
Microcephaly Gene Links Trithorax and REST/NRSF to Control Neural Stem Cell
Proliferation and Differentiation. Cell *151*, 1097–1112.
https://doi.org/10.1016/j.cell.2012.10.043.

49. Li, D., Quan, Z., Ni, J., Li, H., and Qing, H. (2023). The many faces of the zinc finger
protein 335 in brain development and immune system. Biomed. Pharmacother.
165, 115257. https://doi.org/10.1016/j.biopha.2023.115257.

50. Stouffs, K., Stergachis, A.B., Vanderhasselt, T., Dica, A., Janssens, S., Vandervore,
L., Gheldof, A., Bodamer, O., Keymolen, K., Seneca, S., et al. (2018). Expanding
the clinical spectrum of biallelic ZNF335 variants. Clin. Genet. *94*, 246–251.
https://doi.org/10.1111/cge.13260.

51. Sato, R., Takanashi, J., Tsuyusaki, Y., Kato, M., Saitsu, H., Matsumoto, N., and
Takahashi, T. (2016). Association Between Invisible Basal Ganglia and ZNF335
Mutations: A Case Report. Pediatrics *138*, e20160897.
https://doi.org/10.1542/peds.2016-0897.

52. Caglayan, A.O., Yaghouti, K., Kockaya, T., Kemer, D., Cankaya, T., Ameziane, N.,
Cogulu, O., Coker, M., and Yalcinkaya, C. (2021). Biallelic ZNF335 mutations
cause basal ganglia abnormality with progressive cerebral/cerebellar atrophy. J.
Neurogenet. *35*, 23–28. https://doi.org/10.1080/01677063.2020.1833006.

53. Nishida, T., Mihara, T., Horiki, T., and Ka, K. (2016). Anaesthesia and orphan
disease: primary autosomal recessive microcephaly-10 caused by a mutation in the
ZNF335 gene. Eur. J. Anaesthesiol. *33*, 543–545.
https://doi.org/10.1097/EJA.0000000000000358.

54. Theusch, E., Ting, F.Y., Qin, Y., Stevens, K., Naidoo, D., King, S.M., Yang, N.V.,
Orr, J., Han, B.Y., Cyster, J.G., et al. (2024). Participant-derived cell line
transcriptomic analyses and mouse studies reveal a role for ZNF335 in plasma
cholesterol statin response. Genome Med. *16*, 93. https://doi.org/10.1186/s13073-
024-01366-9.

55. Cremer, T., Jongsma, M.L.M., Trulsson, F., Vertegaal, A.C.O., Neefjes, J., and
Berlin, I. (2021). The ER-embedded UBE2J1/RNF26 ubiquitylation complex exerts
spatiotemporal control over the endolysosomal pathway. Cell Rep. *34*.
https://doi.org/10.1016/j.celrep.2020.108659.

56. Cremer, T., Voortman, L.M., Bos, E., Jongsma, M.L., ter Haar, L.R., Akkermans,
J.J., Talavera Ormeño, C.M., Wijdeven, R.H., de Vries, J., Kim, R.Q., et al. (2023).
RNF26 binds perinuclear vimentin filaments to integrate ER and endolysosomal
responses to proteotoxic stress. EMBO J. *42*, e111252.
https://doi.org/10.15252/embj.2022111252.

57. Lecordier, L., Heo, P., Graversen, J.H., Hennig, D., Skytthe, M.K., Cornet d'Elzius,
A., Pincet, F., Pérez-Morga, D., and Pays, E. (2023). Apolipoproteins L1 and L3
control mitochondrial membrane dynamics. Cell Rep. *42*, 113528.
https://doi.org/10.1016/j.celrep.2023.113528.

58. Gaudet, R.G., Zhu, S., Halder, A., Kim, B.-H., Bradfield, C.J., Huang, S., Xu, D.,
Mamiñska, A., Nguyen, T.N., Lazarou, M., et al. (2021). A human apolipoprotein L
with detergent-like activity kills intracellular pathogens. Science *373*.
https://doi.org/10.1126/science.abf8113.

59. Hondele, M., Sachdev, R., Heinrich, S., Wang, J., Vallotton, P., Fontoura, B.M.A.,
and Weis, K. (2019). DEAD-box ATPases are global regulators of phase-separated
organelles. Nature *573*, 144–148. https://doi.org/10.1038/s41586-019-1502-y.

60. Ashburner, M., Ball, C.A., Blake, J.A., Botstein, D., Butler, H., Cherry, J.M., Davis,
A.P., Dolinski, K., Dwight, S.S., Eppig, J.T., et al. (2000). Gene Ontology: tool for
the unification of biology. Nat. Genet. *25*, 25–29. https://doi.org/10.1038/75556.

61. Shroff, K., Caffall, Z.F., and Calakos, N. (2021). DYT-TOR1A subcellular proteomics
reveals selective vulnerability of the nuclear proteome to cell stress. Neurobiol. Dis.
158, 105464. https://doi.org/10.1016/j.nbd.2021.105464.

62. Ding, B., Tang, Y., Ma, S., Akter, M., Liu, M.-L., Zang, T., and Zhang, C.-L. (2021).
Disease Modeling with Human Neurons Reveals LMNB1 Dysregulation Underlying
DYT1 Dystonia. J. Neurosci. *41*, 2024–2038.
https://doi.org/10.1523/JNEUROSCI.2507-20.2020.

63. Pappas, S.S., Liang, C.-C., Kim, S., Rivera, C.O., and Dauer, W.T. (2018). TorsinA
dysfunction causes persistent neuronal nuclear pore defects. Hum. Mol. Genet. *27*,
407–420. https://doi.org/10.1093/hmg/ddx405.

64. Martin, C.-A., Murray, J.E., Carroll, P., Leitch, A., Mackenzie, K.J., Halachev, M.,
Fetit, A.E., Keith, C., Bicknell, L.S., Fluteau, A., et al. (2016). Mutations in genes
encoding condensin complex proteins cause microcephaly through decatenation
failure at mitosis. Genes Dev. *30*, 2158–2172.
https://doi.org/10.1101/gad.286351.116.

65. Trimborn, M., Bell, S.M., Felix, C., Rashid, Y., Jafri, H., Griffiths, P.D., Neumann,
L.M., Krebs, A., Reis, A., Sperling, K., et al. (2004). Mutations in Microcephalin
Cause Aberrant Regulation of Chromosome Condensation. Am. J. Hum. Genet.
75, 261–266. https://doi.org/10.1086/422855.

66. Jean, F., Stuart, A., and Tarailo-Graovac, M. (2020). Dissecting the Genetic and
Etiological Causes of Primary Microcephaly. Front. Neurol. *11*.
https://doi.org/10.3389/fneur.2020.570830.

67. He, K., Zhang, X., Ren, S., and Sun, J. (2015). Deep Residual Learning for Image
Recognition. arXiv. https://doi.org/1512.03385.

68. Deng, J., Dong, W., Socher, R., Li, L.-J., Li, K., and Fei-Fei, L. ImageNet: A large-
scale hierarchical image database. In 2009 IEEE Conference on Computer Vision
and Pattern Recognition, pp. 248–255.
https://doi.org/10.1109/CVPR.2009.5206848.

69. Sundararajan, M., Taly, A., and Yan, Q. (2017). Axiomatic Attribution for Deep
Networks. arXiv.

70. McInnes, L., Healy, J., and Melville, J. (2020). UMAP: Uniform Manifold
Approximation and Projection for Dimension Reduction. Preprint at arXiv,
https://doi.org/10.48550/arXiv.1802.03426
https://doi.org/10.48550/arXiv.1802.03426.

71. Hirano, T. (2005). Condensins: Organizing and Segregating the Genome. Curr. Biol.
15, R265–R275. https://doi.org/10.1016/j.cub.2005.03.037.

72. Drozdz, M.M., and Vaux, D.J. (2016). Shared mechanisms in physiological and
pathological nucleoplasmic reticulum formation. Nucleus *8*, 34–45.
https://doi.org/10.1080/19491034.2016.1252893.

73. Drozdz, M.M., Jiang, H., Pytowski, L., Grovenor, C., and Vaux, D.J. (2017).
Formation of a nucleoplasmic reticulum requires de novo assembly of nascent
phospholipids and shows preferential incorporation of nascent lamins. Sci. Rep. *7*,
7454. https://doi.org/10.1038/s41598-017-07614-w.

74. Mathieu, C., Pappu, R.V., and Taylor, J.P. (2020). Beyond aggregation: Pathological
phase transitions in neurodegenerative disease. Science *370*, 56–60.
https://doi.org/10.1126/science.abb8032.

75. Schludi, M.H., Becker, L., Garrett, L., Gendron, T.F., Zhou, Q., Schreiber, F.,
Popper, B., Dimou, L., Strom, T.M., Winkelmann, J., et al. (2017). Spinal poly-GA
inclusions in a C9orf72 mouse model trigger motor deficits and inflammation
without neuron loss. Acta Neuropathol. (Berl.) *134*, 241–254.
https://doi.org/10.1007/s00401-017-1711-0.

76. Kazemi-Esfarjani, P., and Benzer, S. (2002). Suppression of polyglutamine toxicity
by a Drosophila homolog of myeloid leukemia factor 1. Hum. Mol. Genet. *11*,
2657–2672. https://doi.org/10.1093/hmg/11.21.2657.

77. Kim, W.-Y., Fayazi, Z., Bao, X., Higgins, D., and Kazemi-Esfarjani, P. (2005).
Evidence for sequestration of polyglutamine inclusions by Drosophila myeloid
leukemia factor. Mol. Cell. Neurosci. *29*, 536–544.
https://doi.org/10.1016/j.mcn.2005.04.005.

78. Mitrea, D.M., Mittasch, M., Gomes, B.F., Klein, I.A., and Murcko, M.A. (2022).
Modulating biomolecular condensates: a novel approach to drug discovery. Nat.
Rev. Drug Discov. *21*, 841–862. https://doi.org/10.1038/s41573-022-00505-4.

79. Patel, A., Mitrea, D., Namasivayam, V., Murcko, M.A., Wagner, M., and Klein, I.A.
(2022). Principles and functions of condensate modifying drugs. Front. Mol. Biosci.
9. https://doi.org/10.3389/fmolb.2022.1007744.

80. Jongsma, M.L.M., Berlin, I., Wijdeven, R.H.M., Janssen, L., Janssen, G.M.C.,
Garstka, M.A., Janssen, H., Mensink, M., van Veelen, P.A., Spaapen, R.M., et al.
(2016). An ER-Associated Pathway Defines Endosomal Architecture for Controlled
Cargo Transport. Cell *166*, 152–166. https://doi.org/10.1016/j.cell.2016.05.078.

81. Shaheen, R., Maddirevula, S., Ewida, N., Alsahli, S., Abdel-Salam, G.M.H., Zaki,
M.S., Tala, S.A., Alhashem, A., Softah, A., Al-Owain, M., et al. (2019). Genomic
and phenotypic delineation of congenital microcephaly. Genet. Med. *21*, 545–552.
https://doi.org/10.1038/s41436-018-0140-3.

82. Pavinato, L., Carestiato, S., Trajkova, S., Sorasio, L., Mantovani, G., De Sanctis, L.,
Kerkhof, J., McConkey, H., Rzasa, J., Todd, E., et al. (2025). Skipping of Exon 20
in EP300: A Novel Variant Linked to Rubinstein–Taybi Syndrome With Atypical and
Severe Clinical Manifestations. Clin. Genet. *107*, 354–358.
https://doi.org/10.1111/cge.14654.

83. Roessler, E., Belloni, E., Gaudenz, K., Jay, P., Berta, P., Scherer, S.W., Tsui, L.C.,
and Muenke, M. (1996). Mutations in the human Sonic Hedgehog gene cause
holoprosencephaly. Nat. Genet. *14*, 357–360. https://doi.org/10.1038/ng1196-357.

84. Loo, C.K.C., Pearen, M.A., and Ramm, G.A. (2021). The Role of Sonic Hedgehog in
Human Holoprosencephaly and Short-Rib Polydactyly Syndromes. Int. J. Mol. Sci.
22, 9854. https://doi.org/10.3390/ijms22189854.

85. Rump, P., Jazayeri, O., van Dijk-Bos, K.K., Johansson, L.F., van Essen, A.J.,
Verheij, J.B.G.M., Veenstra-Knol, H.E., Redeker, E.J.W., Mannens, M.M.A.M.,
Swertz, M.A., et al. (2016). Whole-exome sequencing is a powerful approach for
establishing the etiological diagnosis in patients with intellectual disability and
microcephaly. BMC Med. Genomics *9*, 7. https://doi.org/10.1186/s12920-016-0167-
8.

86. Tsai, P.-L., Cameron, C.J.F., Forni, M.F., Wasko, R.R., Naughton, B.S., Horsley, V.,
Gerstein, M.B., and Schlieker, C. (2022). Dynamic quality control machinery that
operates across compartmental borders mediates the degradation of mammalian
nuclear membrane proteins. Cell Rep. *41*, 111675.
https://doi.org/10.1016/j.celrep.2022.111675.

87. Tsai, P.-L., Zhao, C., and Schlieker, C. (2019). Methodologies to monitor protein
turnover at the inner nuclear membrane. Methods Enzymol. *619*, 47–69.
https://doi.org/10.1016/bs.mie.2018.12.033.

88. Ihaka, R., and Gentleman, R. (1996). R: A Language for Data Analysis and
Graphics. J. Comput. Graph. Stat. *5*, 299–314.
https://doi.org/10.1080/10618600.1996.10474713.

89. Vincent, L., and Soille, P. (1991). Watersheds in digital spaces: an efficient algorithm
based on immersion simulations. IEEE Trans. Pattern Anal. Mach. Intell. *13*, 583–
598. https://doi.org/10.1109/34.87344.

90. Brideau, C., Gunter, B., Pikounis, B., and Liaw, A. (2003). Improved Statistical
Methods for Hit Selection in High-Throughput Screening. SLAS Discov. *8*, 634–
647. https://doi.org/10.1177/1087057103258285.

91. Shun, T.Y., Lazo, J.S., Sharlow, E.R., and Johnston, P.A. (2011). Identifying Actives
from HTS Data Sets: Practical Approaches for the Selection of an Appropriate HTS
Data-Processing Method and Quality Control Review. J. Biomol. Screen. *16*, 1–14.
https://doi.org/10.1177/1087057110389039.

92. Wang, Y., Xiao, J., Suzek, T.O., Zhang, J., Wang, J., and Bryant, S.H. PubChem: a
public information system for analyzing bioactivities of small molecules.

93. NCI/CADD Chemical Identifier Resolver
https://cactus.nci.nih.gov/chemical/structure.

94. Baell, J.B., and Holloway, G.A. (2010). New Substructure Filters for Removal of Pan
Assay Interference Compounds (PAINS) from Screening Libraries and for Their
Exclusion in Bioassays. J. Med. Chem. *53*, 2719–2740.
https://doi.org/10.1021/jm901137j.

95. Baell, J.B., and Nissink, J.W.M. (2018). Seven Year Itch: Pan-Assay Interference
Compounds (PAINS) in 2017-Utility and Limitations. ACS Chem. Biol. *13*, 36–44.
https://doi.org/10.1021/acschembio.7b00903.

96. Lipinski, C.A. (2004). Lead- and drug-like compounds: the rule-of-five revolution.
Drug Discov. Today Technol. *1*, 337–341.
https://doi.org/10.1016/j.ddtec.2004.11.007.

97. Wickham, H. (2011). ggplot2. WIREs Comput. Stat. *3*, 180–185.
https://doi.org/10.1002/wics.147.

98. Marcel, S., and Rodriguez, Y. (2010). Torchvision the machine-vision package of
torch. In Proceedings of the 18th ACM international conference on Multimedia
(ACM), pp. 1485–1488. https://doi.org/10.1145/1873951.1874254.

99. Pedregosa, F., Varoquaux, G., Gramfort, A., Michel, V., Thirion, B., Grisel, O.,
Blondel, M., Müller, A., Nothman, J., Louppe, G., et al. (2018). Scikit-learn:
Machine Learning in Python. Preprint at arXiv,
https://doi.org/10.48550/arXiv.1201.0490 https://doi.org/10.48550/arXiv.1201.0490.

100. Paszke, A., Gross, S., Massa, F., Lerer, A., Bradbury, J., Chanan, G., Killeen, T.,
Lin, Z., Gimelshein, N., Antiga, L., et al. (2019). PyTorch: An Imperative Style,
High-Performance Deep Learning Library. Preprint at arXiv,
https://doi.org/10.48550/arXiv.1912.01703
https://doi.org/10.48550/arXiv.1912.01703.

101. Virtanen, P., Gommers, R., Oliphant, T.E., Haberland, M., Reddy, T., Cournapeau,
D., Burovski, E., Peterson, P., Weckesser, W., Bright, J., et al. (2020). SciPy 1.0--
Fundamental Algorithms for Scientific Computing in Python. Nat. Methods *17*, 261–
272. https://doi.org/10.1038/s41592-019-0686-2.

102. Shannon, C.E. (1948). A Mathematical Theory of Communication. Bell Syst. Tech.
J. *27*, 379–423. https://doi.org/10.1002/j.1538-7305.1948.tb01338.x.

103. Ritz, C., Baty, F., Streibig, J.C., and Gerhard, D. (2015). Dose-Response Analysis
Using R. PLOS ONE *10*, e0146021. https://doi.org/10.1371/journal.pone.0146021.

104. Wu, T., Hu, E., Xu, S., Chen, M., Guo, P., Dai, Z., Feng, T., Zhou, L., Tang, W.,
Zhan, L., et al. (2021). clusterProfiler 4.0: A universal enrichment tool for
interpreting omics data. The Innovation *2*.
https://doi.org/10.1016/j.xinn.2021.100141.

105. Galili, T., O'Callaghan, A., Sidi, J., and Sievert, C. (2018). heatmaply: an R
package for creating interactive cluster heatmaps for online publishing.
Bioinformatics *34*, 1600–1602. https://doi.org/10.1093/bioinformatics/btx657.

1401