Spectroscopic and morphological study of sinusoidal mechanical vibration exposed human RBC in vitro

Vijay Ghodake[1,2], Sarika Hinge*[1], Arun Banpurkar[1], and Gauri Kulkarni[1]

[1]Department of Physics, Savitribai Phule Pune University, Pune, Maharashtra, India

[2]Central Water and Power Research Station, Pune - 411024, Maharashtra, India.

*Corresponding Author: Sarika Hinge

alwaysonsara@yahoo.co.in; sarika21hinge@gmail.com

Key points:

- Effect of mechanical vibration on RBC highlights the importance of blood transportation and handling related issues.
- Spectroscopic and Morphological analysis revealed the modification in RBC due to sinusoidal mechanical vibration.

Abstract

Humans undergoing mechanical vibration of frequency ranging from mHz to a few kHz in daily life. Human body contains about 60% of water. Blood is one of the most important body fluids and RBC are functional constituents of blood. RBC, due to their deformable nature, can undergo physical deformation or membrane damage under the influence of external mechanical vibrations over extended periods. In the present work, we studied in vitro the effect of sinusoidal mechanical vibration of 2 mm amplitude and frequency ranging from 5 to 100 Hz for 10 min duration. Physicochemical properties of controlled and mechanically vibrated RBC were examined using UV-visible, Fourier Transform Infrared (FTIR), Raman spectroscopy, and morphology was examined by Scanning Electron Microscopy (SEM). Results showed that mechanical vibrations affect oxygenation level at the Q band (543 nm and 578 nm), and variation in transmission intensity is observed in the Amide region. RBC showed a change in morphology, where the elongation index increases along with vibration frequency. Current findings are useful to various stages of blood management and supply chain like collection, storage, transportation, handling, and donation.

Keywords: vibrations, RBC, spectroscopy, SEM, elongation index

Introduction

Daily physical activities like running, walking, playing, driving, travelling cause excessive mechanical stress viz. compression, tension, and shear stress on human body.[1-3] During road transportion, human body experience mechanical vibrations in the range of 1 to 500 Hz [4] which indirectly affects the organs, tissues and soft matter in human body. As a result, damage to the cytoskeleton, proteins, cells, and tissues occur which alters cell cycle, its metabolism and blood supply to various vital organs. [5,6] Mechanical vibrations can be destructive or constructive in nature and generate abiotic stresses on the body cells[7-8] which depends on its waveform, frequency, amplitude, and exposure time. In some cases, vibrational frequencies affect the bone density, nerve, muscle, vision and skin microcirculation. [9-13] High frequency vibrations can also create multifracture damage to the bone. [14] However, whole human body treated with external vibrations during physical workout showed better performance in bone strength and bone mass in adult persons, astronauts, etc.[15] Vibrational frequency is used in physiotherapy for treatments of joints and tissues, hemodialysis, wound healing in diabetic foot ulcers, [16] morphogenesis of neuronal cells and fibroblast. [17,18] In recent years, it is shown that the mechanical vibration affects the blood circulation, blood constituents and it activates. [19] Changes in the physicochemical characteristics of RBC and its biological pathways might result in circulation disorders, and cardiovascular problem due to mechanical vibrations. [20,21] About 45% blood volume is occupied by RBC and it is a major constituent of whole blood mainly used as a gas transporter and drug carrier. Blood components like RBC, lymphocytes, platelets and blood plasma get affected due to mechanical vibration [22,23] which influences the viscosity of blood to avoid aggregation [24] and increases the erythrocyte volume fraction (EVF) of arm hand

and Hb in index fingers.[25]   Other studies have also shown that vibrations and

mechanical shock can increase the hemolysis of RBC. External local vibration therapy

delivered to impaired part of the human body increases blood flow without any RBC

aggregation. However, most of the vibrations show adverse effect on peroxidase activity

of hemoglobin at low frequency vibrations within the range of 8 to 32 Hz. [26] Earlier, it

was reported that road transportation stress on RBC causes osmotic fragility of RBC,

hemoglobin level and blood rheology. [27]

Although the effects of mechanical vibrations on biological systems have been

documented, it is not known how mechanical vibrations affect morphological and

biophysical properties of RBC. In the present study, RBC were exposed to 5 Hz,10 Hz,

20 Hz, 30 Hz, 40 Hz, 50 Hz, 60 Hz, 70 Hz, and 100 Hz sinusoidal vibrational

frequency. Comparison between vibrated and non-vibrated RBC was done on the basis

of UV-visible spectroscopy, FTIR spectroscopy, Raman spectroscopy and Scanning

Electron Microscopy.

Materials and Methods

Sample preparation

Blood sample (10 ml) was collected from venous of healthy human by in

ethylenediamine tetraacetic acid ~EDTA, (3g/l) and all methods were performed with

the approval and following the relevant guidelines and regulations set by the Institute

Ethical Committee (Ref. No. SPPU/IEC/2021/129). Collected blood was stored at 4 °C

and used for the experiment within 48 hr. RBC were extracted from whole blood at

3000 revolutions per minute (RPM) for 10 minutes at 25 °C, using centrifugation

process (Centrifuge machine: Superspin R-V/FM Plasto Crafts). Blood components like

RBC, WBC, plasma were separated relative to density gradient. Isolated RBC were

washed thrice by using 0.9% saline and used for experiments. All experiments were

conducted three times using pure RBC samples obtained from adult, healthy volunteers

(Age = 25-30 yr, N = 5).

Experimental setup for mechanical vibration

A vibration test system (Sine force – peak 100 kgf, frequency range 5 Hz to 4500 Hz,

maximum acceleration, velocity, and displacement are 65 g, 1.6 m/s, and pk-pk 25 mm,

respectively) using an electrodynamic shaker from Spectral Dynamics, USA was used to

induce mechanical vibrations of various frequencies to the sample holder the sample.

The schematic of the mechanical vibration system is shown in Figure 1.

Figure 1. Schematic of the experimental setup for inducing mechanical vibration

system.

The driver signal from the function generator is input to the amplifier. The Amplifier

converts low voltage and mA currents from the controller to higher voltages and

currents require for the driving shaker unit. The moving element (armature) of the

shaker is suspended in a strong magnetic field. The current output from the amplifier

flows through the wire in a magnetic field. The suspended moving element is restrained

and allowed to move up and down. Movement of sample table is sensed by the

piezoelectric accelerometer. The conditioned signal is fed into the closed control loop

from the charge amplifier to a pre-defined reference level. The control loop adjusts the

drive signal to the amplifier in order to maintain the reference level. The motion of the

table follows the characteristics of the current flowing through a coil produces sinusoidal

vibration motion of the sample holder.

To examine the vibration effect on blood samples (0.05 ml) in vial was fixed vertically

on vibration platform of electrodynamic shaker. RBC samples were vibrated at different

frequencies viz. 5, 10, 20, 30, 40, 50, 60, 70, and 100 Hz with constant amplitude of 2

mm and 10 minute exposure.

Experimental procedure for UV-visible, FTIR and Raman spectroscopy

Effect of mechanical vibrations on RBC was characterized by using UV-visible

(JASCO-670, Japan), FTIR (JASCO-6100, Japan) and Raman spectroscopy (inVia^{TM}

Renishaw). UV-visible spectroscopy of RBC was studied by diluting the fluids in the

ratio of 1:100 within the spectral range 200 - 800 nm. For FTIR spectra, 15 µl of RBC

added into KBr powder and dried at room temperature and measurement were recorded

in 400 – 4000 cm^{-1}. To obtained the Raman spectra , 10 µl of RBC solution was dried

on glass slide and spectra were monitored in the spectral range of 400 -1800 cm^{-1}.

Experimental procedure for Scanning Electron Microscopy of RBC

Isolated RBC (2 ml) were added in a vial containing buffer solution (0.4 ml of 2%

glutaraldehyde and 19.6 ml saline). These samples were slowly shaken for 2 to 3

minutes and kept in refridgerator 10-12 hr at temperature 4 °C. Again 0.5 ml saline was

added to each vial, then these vials were centrifuged and supernatant plasma was

removed with micropipette. The ethanol treatment was given to the leftover red blood

cells. For ethanol treatment, the solution was prepared by mixture of ethanol and saline

with concentration 10, 20, 30, 40, 50, 60, 70 , 80, 90, and 100 % by using the standard

protocol (Add 0.2 ml ethanol to 1.8 ml saline, the total solution is 10%. Add 0.4 ml

ethanol to 1.6 ml saline the total solution is 20%, similar way 30%, 40%, 50%, 60%,

70%, 80%,90% and 100% solution were prepared). Three glass slides for each sample

were prepared and RBC smeared uniformly on the glass substrate. Total thirty slides

were prepared and kept in vacuum for whole night. SEM for control and vibrated

samples were performed for x3000 magnification.

Results

UV-visible spectra, FTIR spectra, Raman spectra and Scanning Electron Microscopy

measured for control RBC and mechanically vibrated RBC.

UV-visible spectroscopy

The absorption spectra measured from the vibrated RBC at different frequencies and

non-vibrated (control)   isolated RBC from whole blood is shown in Figure 2.

Absorption from vibrated and non-vibrated RBC were seen at 281 nm, 346 nm, and

418 nm, 543 nm, and 578 nm.

Figure 2. Changes in the absorption spectra of **(a)** RBC vibrated at 5, 10, 20, 30, 40, 50, 60, 70 and 100 Hz with amplitude of 2 mm for 10 minute (b) cumulative absorption spectra of RBC, inset shows intensity of 418 nm sorat band. (c) magnified view of the Q-band (d) plot shows ratio of Q1/Q2 versus vibrated frequency.

The absorbance at 281 nm and 346 nm is a determinant indicator of protein content in RBC. The peak at obtained at 281 nm is associated with π-π* transition and peak at 346 nm showed signature of aromatic amino residues (Tryptophan, Tyrosine, and Phenelanine). Absorption at haem group of vibartaed and non-vibrated RBC is seen at 418 nm. The alteration in intensity and wavelength is related to haem group, and its interaction with polypeptide bonds surrounded by Fe group in porphyrin ring. However, Q band (alpha and beta) showed changes in ligand-porphyrin which is

associated with oxygen binding and transport. Absorption peak related to oxygenation

Q1(alpha) ~543 nm and Q2 (beta) ~578 nm showed increasing trend with increase in

the vibrational frequencies from 5, 10, 20, 30, 40, 50, 60, 70, and 100 Hz as compared

to control sample. Variation in intensity at Q1 and Q2 bands was observed for vibrated

RBC which led to change in the oxygenation state.

FTIR Spectroscopy

The FTIR spectra of control and vibrated RBC were measured at 10, 30, 50, 70 and

100 Hz with amplitude of 2 mm at 10-minute duration.

Figure 3. FTIR spectra of RBC vibrated at 5, 10, 20, 30, 40, 50, 60, 70, and 100 Hz

with amplitude of 2 mm for 10 minute duration.

FTIR spectra of control and vibrated RBC show peaks at 1551-1558 cm^{-1} which

corresponds to Amide II (C=O stretching of proteins), 1632-1659 cm^{-1} corresponds to

Amide I (mainly C=O stretching of proteins) and 3357 - 3443 cm^{-1} corresponds to

Amide A (N-H stretching of secondary amide proteins). [28] Figure 3 shows the peaks

obtained from the control and vibrated RBC. FTIR spectra of vibrated RBC shows

variation in the transmission intensity and changes in the functional group particularly

amide band. Alterations in Amide groups of RBC was observed due to vibrations of

different frequency.

Raman Spectroscopy

Raman spectra of RBC were recorded with laser of 785 nm wavelength (power : 3 mW)

within the range of 400 cm^{-1} to 1800 cm^{-1} as shown in **Figure 4**.

Figure 4. Raman spectra of RBC vibrated at 5, 10, 20, 30, 40, 50, 60, 70, and 100 Hz

with amplitude of 2 mm for 10 minute duration.

Raman spectra of RBC and mechanically vibrated RBC showed the Raman peaks at

569 cm^{-1} (Fe-O$_2$) stretch), 676 cm^{-1} (δ(pyrrole deformation) symmetric in plane

deformation (ν_7)), 755 cm^{-1} (ν (pyrrole breathing)stretch (ν_{15})), 793 cm^{-1} (ν(pyrrole

deformation)ν_6), 825 cm^{-1} (γ (C$_m$H)out of plane deformation (γ_{10}, Tyrosine), 964 cm^{-1}

(C-C stretch in unordered protein),1003 cm^{-1} (ν (C$_\beta$C$_1$) asymmetric stretch (ν_{45}) and

Phenylalanine), 1130 cm^{-1} (C-C stretch (trans) lipids), 1175 cm^{-1} (ν(pyrrole half ring)

asymmetric stretch (ν_{30})), 1224 cm^{-1} (δ(CmH) in plane deformation (ν_{13})), 1341 cm^{-1} (N

(pyrrole half ring) symmetric stretch (ν_{41})), 1396 cm^{-1} (N (pyrrole quarter-ring) stretch

(ν_{20})), 1450 cm^{-1} (δ(CH$_2$/CH$_3$)), 1564 cm^{-1} (ν(C$_\beta$C$_\beta$) stretch (ν_2)),1580 cm^{-1} (ν (C$_\alpha$C$_m$)

asymmetric stretch (ν_{37})),1620 cm^{-1} (ν (C$_a$=C$_b$) of vinyl groups). [29] Normalized Raman

spectra of vibrated RBC showed decrease in intensity with increase in mechanical

vibration. However, oxidative state marker region (1300-1400 cm^{-1}) showed slightly

varition in peak position and intensity.

Morphological study

Scanning electron microscopic images of RBC isolated from vibrated whole blood were

obtained at different frequencies (Figure 5).

Figure 5. (a) Morphology of isolated RBC (Control) and vibrated RBC at different
frequencies of vibrations i.e. 10, 30, 50, 70, and 100 Hz with constant amplitude (2
mm); red line indicates the length of RBC (b)   Elongation index Vs Vibrational
frequency; inset   image presents the   measurement of RBC length and width for
elongation index

Scanning electron microscope (JEOl JSM-6360A) was used to obtain RBC images
before and after mechanical vibration. Public domain (Image J, 1.54f) software is used
for data analysis. From the control, the average diameter of RBC was measured as 4.76
μm and thickness around 1.84 μm. In the case of vibrated samples, the diameter of RBC
increases from 4.76 μm ± 0.08 to 5.64 ± 0.06 μm with increasing vibrating frequencies
and the width of RBC decreases from 1.84 ± 0.03 μm to 1.32 ± 0.05 μm with increasing
vibrating frequencies. Change in the elongation index of RBC at different vibration
frequencies is seen in Figure 5(b).   Scanning electron microscopic images show
deformation in RBC after exposure to sinusoidal mechanical vibrations, diameter of
RBC increased along with a decrease in thickness, and width. The same trend was
observed in isolated RBC from vibrated whole blood at different frequencies. More
details of the physicochemical properties of vibrated whole blood and its
characterizations are available in the electronic supplementary note (**SP)**.

Discussion

The advancement in the research of vibration on RBC has potential in health and risk
management in blood transfusion. During blood transportation via road, many external
factors affect the physiology of RBC. To maintain the efficacy and integrity of RBC, it
is necessary to understand the effect of sinusoidal vibration particularly for blood
transfusion.

Hemoglobin is a major part of RBC carrying and binding oxygen to an encapsulated porphyrin ring with two alpha and beta chains. Absorption at oxygen binding sites is examined at 543 nm (Q1) and 578 nm (Q2) using UV-visible spectroscopy. Mechanical vibration affects the oxygenation state of RBC which is determined by the ratio of Q1/Q2. Change in the microenvironment of the RBC was observed with variation in vibrational frequency which led to a decrease in absorbance intensity and FWHM of the peak. For quantitative measurement and studying the effect of sinusoidal vibration on RBC, FTIR spectroscopy is used to investigate functional groups mainly associated with Amide I and Amide II.

Sinusoidal vibrations induce stress on the secondary structure present at Amide I and II region . RBC is composed of hemoglobin, spectrin network, cytoskeleton membrane proteins, and associate secondary structure viz. α-helix, β-sheet, random coil, and aggregates. The second derivative and difference in amide bands I and II provide information related to secondary structure. Molecular level investigation of RBC was performed by using Raman spectrsocsopy. Spectra shows changes in intensities in CH_3 amino acid deformation mode, hemeagreegation, $Fe - O_2$ stretch mode. Intensity of spectra increases, but no significant change in Raman shift was observed due to mechanical vibrations.

During microcirculilation, RBC deform from biconcave shape to dumbbell shape and passess through 3-5 micron size blood capillaries. Due to elastic and flexible nature, RBC regain its original shape and size. Elongation index of RBC was determined by measuring the diameter and width of RBC using public domain software (ImageJ 1.54f). Vibartion affects the morphology of RBC which causes an increase in diameter with a decrease in thickness. Overall elongation index is increased as a function of

vibrational frequency. Elongation may affect the deformation property of RBC. Hence,

there is more research is needed to understand the deformability of vibration exposed

individual live RBC.

Conclusions

In this work, it is seen that sinusoidal mechanical vibration affects the shape and

physicochemical properties of RBC. Such change is vital in microcirculation and

decides oxygenation state of RBC affecting oxygen carrying capacity. Mechanical

vibrational stress beyond the natural frequency of RBC affects the morphology of RBC

which led to elongation of RBC, and variation in peak intensity of Raman peak . Hence,

it is advisable to avoid exposure of human body to continuos mechanical vibrations

within certain range of frequency amd amplitude and type of vibrations. Vibrations may

cause change in mechanical properties of RBC and affects their dynamics. This work

has important significance in safety handling and risk management during blood

transportation. .

Authors Information

Authors and Affiliations

Department of Physics, Savitribai Phule Pune University

Central Water and Power Research Station, Pune

Contributions

V.G. and S.H. contributes equally. V.G: contributes to Conceptualization,

Methodology, Validation, Writing; S.H: contributes to Methodology, Investigation,

Data curation, writing – draft; A.B.: Data curation, writing - review & editing,

Supervision; G.K: Data curation, writing - review & editing, Supervision

Corresponding author

Correspondance to **Sarika Hinge**

Declarations

Ethics approval

The entire protocol for collection of blood sample is approved by Institute Ethical

Committee (Ref. No. SPPU/IEC/2021/129).

Consent to participate

Informed consent was obtained from all individual participants included in the study.

Competing Interests

"The authors declare that they have no competing interests".

Acknowledgement

Authors are grateful to Director, Central Water and Power Research Station, Pune, India

and Director of Basic Medical Science, Department of Physics, Savitribai Phule, Pune

University, Pune - 411007, MH, INDIA.

Funding : None

Data Availability

All data generated or analyzed during this study are included in this published article.

References

1. Gattner, H., Adamiak, J., Piotrowska, A., Czerwińska-Ledwig, O., Mętel, S., Kępińska-
Szyszkowska, M. and Pilch, W.
Effect of Whole-Body Vibration Training on Hemorheological Blood Indices in Young,
Healthy Women.
International Journal of Environmental Research and Public Health, 20(4), (2023),
p.3232.
2. Bernardo-Filho, M., Bemben, D., Stark, C. and Taiar, R.

Biological consequences of exposure to mechanical vibration.

Dose-Response, 16(3), (2018),p.1559325818799618.

3. 294   Monteiro, M.O., Pinto, N., Marin, P.J., Santos-Filho, S.D. and Bernardo-Filho, M.

295   Effect on osmotic fragility of red blood cells of whole blood submitted to vibrations in an

296   oscillating platform.

297   African Journal of Biotechnology, 10(64), (2011),pp.14197-14202.

4. 298   Lalanne, C.

299   Mechanical vibration and shock analysis, fatigue damage (Vol. 4), (2014), John Wiley &

300   Sons.

5. 301   Kumar, A., Ali, M., Pandey, B.N., Hassan, P.A. and Mishra, K.P.

302   Role of membrane sialic acid and glycophorin protein in thorium induced aggregation and

303   hemolysis of human erythrocytes.

304   Biochimie, 92(7), (2010), pp.869-879.

6. 305   De Zepetnek, J.T., Giangregorio, L.M. and Craven, B.C.

306   Whole-body vibration as potential intervention for people with low bone mineral density

307   and osteoporosis: a review.

308   Journal of Rehabilitation Research & Development, *46*(4), 2009.

7. 309   Hassanpour, H. and Niknam, V.

310   Effect of different sinusoidal-vibration frequencies on some physiological parameters,

311   antioxidant activity and total phenol of Anthemis gilanica Bornm & Gauba.

312   Journal of Plant Process and Function, 8(30), (2019), pp.185-194.

8. 313   Darwish, S.M., Darwish, A.S. and Darwish, D.S.

314   An extremely low-frequency magnetic field can affect CREB protein conformation which

315   may have a role in neuronal activities including memory.

316   Journal of Physics Communications, 4(1), 2020, p.015009.

9. 317   Liao, F., Zhang, K., Zhou, L., Chen, Y., Elliott, J. and Jan, Y.K.

Effect of different local vibration frequencies on the multiscale regularity of plantar skin

blood flow.

Entropy, 22(11),(2020), p.1288.

10. Johansson, N., Ragnebro, O., Stjernbrandt, A., Graff, P., Bryngelsson, I.L. and Vihlborg,

P. Effects on blood parameters from hand-arm vibrations exposure.

Toxicology and industrial health, 39(6), (2023), pp.291-297.

11. Pinto, N.S., Monteiro, M.B., Arthur, A.P., Paiva, D.N., Meyer, P.F., Santos-Filho, S.D.,

Marín, P.J. and Bernardo-Filho, M.

Effectiveness of a protocol involving acute whole-body vibration exercises in an adult and

health individual with delayed-onset muscle soreness observed after running: a case

report.

J. Med. Med. Sci, 2(1),(2011), pp.612-7.

12. Pelmear, P.L., Kusiak, R. and Leong, D.

Hand-arm vibration syndrome associated with impact vibration.

Journal of Low Frequency Noise, Vibration and Active Control, 14(2), (1995), pp.73-79.

13. Ando, H., Nieminen, K., Toppila, E., Starck, J. and Ishitake, T.

Effect of impulse vibration on red blood cells in vitro.

Scandinavian journal of work, environment & health, (2005), pp.286-290.

14. Prisby, R.D., Lafage-Proust, M.H., Malaval, L., Belli, A. and Vico, L.

Effects of whole body vibration on the skeleton and other organ systems in man and

animal models: what we know and what we need to know.

Ageing research reviews, 7(4), (2008), pp.319-329.

15. De Zepetnek, J.T., Giangregorio, L.M. and Craven, B.C.

Whole-body vibration as potential intervention for people with low bone mineral density

and osteoporosis: a review.

Journal of Rehabilitation Research & Development, 46(4), (2009).

Ren, W., Pu, F., Luan, H., Duan, Y., Su, H., Fan, Y. and Jan, Y.K.

Effects of local vibration with different intermittent durations on skin blood flow responses

in diabetic people.

Frontiers in Bioengineering and Biotechnology, 7, (2019), p.310.

16. Santiago Júnior, O., Huebner, R. and Duarte, M.L.

Effects of mechanical vibration on bone–a critical review.

Jaw Functional Orthopedics and Craniofacial Growth, 2(3), (2022), pp.141-150.

17. Syabariyah, S., Nurachmah, E., Widjojo, B.D., Prasetyo, S., Sanada, H., Irianto,

Nakagami, G., Suriadi, Kardiatun, T. and Hisan.

The Effect of Vibration on the Acceleration of Wound Healing of Diabetic Neuropathic Foot

Ulcer: A Prospective Experimental Study on Human Patients.

In Healthcare , 11(2), (2023), p. 191.

18. Nakagami, G., Sanada, H., Matsui, N., Kitagawa, A., Yokogawa, H., Sekiya, N., Ichioka,

S., Sugama, J. and Shibata, M.

Effect of vibration on skin blood flow in an in vivo microcirculatory model.

Bioscience trends, 1(3),(2007), pp.161-166.

19. Shin, S., Jang, J.H., Park, M.S., Ku, Y.H. and Suh, J.S.

A noble RBC aggregometer with vibration-induced disaggregation mechanism.

Korea-Australia Rheology Journal, 17(1), (2005), pp.9-13.

20. Shin, S., Ku, Y., Suh, J.S., Moon, S.Y. and Jang, J.Y.

Characteristics of blood flow resistance under transverse vibration: red blood cell

suspension in Dextran-40.

Annals of Biomedical Engineering, 31,(2003), pp.1077-1083.

21. Dybas, J., Alcicek, F.C., Wajda, A., Kaczmarska, M., Zimna, A., Bulat, K., Blat, A.,

Stepanenko, T., Mohaissen, T., Szczesny-Malysiak, E. and Perez-Guaita, D.

Trends in biomedical analysis of red blood cells–Raman spectroscopy against other

spectroscopic, microscopic and classical techniques.

TrAC Trends in Analytical Chemistry, *146*, (2022), p.116481.

22. Liao, F., Zhang, K., Zhou, L., Chen, Y., Elliott, J. and Jan, Y.K.

Effect of different local vibration frequencies on the multiscale regularity of plantar skin

blood flow.

Entropy, 22(11), (2020), p.1288.

23. Andrzejak, R. and Smolik, R.

Effect of vibration on red cell metabolism.

International archives of occupational and environmental health, 54, (1984), pp.303-308.

24. Shin, S., Ku, Y.H., Moon, S.Y. and Suh, J.S.

The effect of vibration on the hemorheological characteristics of non-aggregated blood.

KSME international journal, 17, (2003), pp.1104-1110.

25. Johansson, N., Ragnebro, O., Stjernbrandt, A., Graff, P., Bryngelsson, I.L. and Vihlborg,

P.Effects on blood parameters from hand-arm vibrations exposure.

Toxicology and industrial health, 39(6), (2023), pp.291-297.

26. Dotsenko, O.I., Taradina, G.V. and Mischenko, A.M.

Peroxidase Activity of Erythrocytes Hemoglobin Under Action of Low-Frequency

Vibration.

Studia Biologica, 15(4),(2021), pp.3-16.

27. Pramudita, J. J., Widjanarko, B., Munadi, M., & Suwondo, A.

Effect of vibration on the quality of blood transfusion in whole blood.

Journal of Medicinal and Chemical Sciences, 6(12), (2023), 2913–2921.

https://doi.org/10.26655/JMCHEMSCI.2023.12.6

28. Miller, L.M., Bourassa, M.W. and Smith, R.J.

FTIR spectroscopic imaging of protein aggregation in living cells.

Biochimica et biophysica acta (BBA)-biomembranes, 1828(10),(2013), pp.2339-2346.

29. Krishna, R. and Colak, I.

Advances in Biomedical Applications of Raman Microscopy and Data Processing: A Mini

Review.

Analytical Letters, 56(4), (2023), pp.576-617.