Placental Iron Utilisation in Fetal Growth Restriction: Alterations in Mitochondrial Heme Synthesis and Iron-Sulfur Cluster Assembly Pathways

Veronica B Botha[1,2], Heather C Murray[1,3], Siddharth Acharya[1,2], Kirsty G Pringle[1,4], Roger Smith*[2,5], Joshua J Fisher*[2,5].

[1]School of Biomedical Sciences and Pharmacy, University of Newcastle, Callaghan, New South Wales, Australia

[2]School of Medicine and Public Health, University of Newcastle, Callaghan, New South Wales, Australia

[3] Precision Medicine Program, Hunter Medical Research Institute, New Lambton Heights, New South Wales, Australia

[4]Women's Health Research Program, Hunter Medical Research Institute, New Lambton Heights, New South Wales, Australia

[5]Mothers and Babies Research Program, Hunter Medical Research Institute, New Lambton Heights, New South Wales, Australia

*denotes equal contribution as senior author

Corresponding Author:

Dr Joshua Fisher

Postdoctoral Research Fellow

Mothers and Babies Research Centre

Hunter Medical Research Institute

Faculty of Health and Medicine | School of Medicine and Public Health

Joshua.fisher@newcastle.edu.au, https://orcid.org/0000-0002-0505-0598

Key Points

- Iron plays a critical role in placental function, while iron-dependent pathway components are well-characterised, their integrated response and adaptive reprogramming in fetal growth restriction (FGR) remain poorly understood.

- In FGR, maternal iron status was unchanged, however, placental iron uptake proteins were increased and ferroportin reduced, suggesting that the placenta retains iron.

- FGR placentae showed altered de novo mitochondrial iron-sulfur cluster (Fe-S) formation and a bottleneck in late-stage Fe-S cluster assembly.

- This shift in Fe-S synthesis prioritises mitochondrial and cytosolic heme synthesis pathways, consistent with increased heme utilisation and breakdown.

- Globin subunits were lower in protein abundance and impaired placental erythrocyte
structure in FGR.
- Dysregulation of erythrocyte membrane proteins in FGR placentae suggests altered
erythrocyte structure, potentially representing an adaptive response to inadequate
vascularisation, attempting to optimise oxygen delivery to the fetus.

Abstract

Fetal growth restriction (FGR) affects ~10% of pregnancies worldwide and is often associated
with placental insufficiency. Iron is essential for maternal haematopoietic adaptations and
placental processes such as mitochondrial iron–sulfur (Fe-S) cluster assembly, heme synthesis,
and erythropoiesis. This study aimed to characterise iron transport and downstream utilisation
in FGR. Placental tissues from term uncomplicated (n=19) and FGR (n=18) pregnancies were
analysed. Maternal iron status was retrospectively assessed from clinical records. Placental
mRNA and protein expression of iron-dependent pathways were analysed via RT-qPCR, LC-
MS, and western blotting. Placental iron content was assessed histologically, and heme levels
were measured by activity assay. FGR pregnancies showed significantly elevated maternal
serum ferritin and lower red cell distribution width, although remained within normal clinical
values. Placental iron uptake transporters TFRC and DMT1 were significantly upregulated,
while the iron exporter to the fetus, ferroportin, was reduced, indicating increased iron retention
in the FGR placenta. Despite altered transporter expression, Fe^{3+} iron levels were unchanged,
suggesting iron utilisation over storage. Subsequent investigations identified reduced
mitochondrial Fe-S synthesis components (FDXR, FDX2, NDUFAB1, HSPA9), and a
prioritisation of mitochondrial and cytosolic heme synthesis enzymes in FGR. Protein levels
of haemoglobin subunits (HBG1, HBG2, HBB, HBA1) and erythrocyte membrane markers
(EPB41, EPB42, SPTA1, SPTB, ANK1) were decreased. These findings reveal a compensatory
response in FGR placentae, with increased iron uptake and utilisation favouring heme synthesis
over Fe-S cluster formation, possibly to support oxygen handling under poor placental
vascularisation and reduced fetal oxygenation, with potential consequences for mitochondrial
energy metabolism.

Introduction

Fetal growth restriction (FGR) occurs in 10% of pregnancies globally and is associated with increased perinatal and neonatal morbidity and mortality (Miller *et al.*, 2008). FGR is defined by the inability of the fetus to achieve its genetically predetermined growth potential relative to its gestational age (Gordijn *et al.*, 2016). Although maternal and fetal complications are recognised causes of FGR, 60-70% of FGR cases are associated with placental insufficiency, where the placenta fails to adequately deliver nutrients and oxygen, arising from inadequate uterine spiral artery remodelling (Brooker *et al.*, 2025; Higashijima *et al.*, 2013; Tang *et al.*, 2017). In FGR placentae, the cellular landscape is altered, with impaired cytotrophoblast maturation and decreases in trophoblast volume. This results in a reduction in the number of placental villi and a decreased placental diameter (Sun *et al.*, 2020). Consequently, the surface area available for oxygen and nutrient exchange is diminished, limiting substrate transfer to the fetus (Zygmunt *et al.*, 2003). In response, the fetus adapts by redistributing blood flow, prioritising essential fetal organs, such as the brain and heart, resulting in asymmetrical growth and slowing the fetal growth rate (Miller *et al.*, 2008; Roberts *et al.*, 2020; Wu *et al.*, 2023).

The placenta is a highly vascularised, potent haematopoietic organ. The placenta performs primitive haematopoiesis during embryogenesis, establishing a large reservoir of haematopoietic stem cells (HSCs) that not only support erythrocyte activity during early gestation, but also provide progenitor cells for the fetal liver and bone marrow when haematopoiesis transitions away from the placenta (Gekas *et al.*, 2010; Palis & Segel, 1998). Hematopoietic function is critically reliant on iron availability, as iron is essential for erythropoiesis and mitochondrial functions that facilitate the biosynthesis of iron-sulfur clusters (Fe-S) and subsequent heme synthesis in erythroid cells (Chiabrando *et al.*, 2014; Hentze *et al.*, 2004; Mastrogiannaki *et al.*, 2009).

Iron is an essential micronutrient for both fetoplacental development and mitochondrial bioenergetics (Dev & Babitt, 2017; Guo *et al.*, 2019). During pregnancy, the requirement for iron alters as gestation advances. In early pregnancy, placental development occurs in a physiologically hypoxic environment, with relatively low iron demands (Guo *et al.*, 2016). This low-oxygen environment stabilises hypoxia inducible factors (HIFs), which transcriptionally activate genes involved in oxygen and iron regulation, such as erythropoietin (EPO), hepcidin and erythroferrone (Semenza, 2014). In the first trimester, rising EPO levels support both maternal erythropoiesis and transient placental hematopoietic activity (Cindrova-Davies & Sferruzzi-Perri, 2022; Delaney *et al.*, 2021; Fairchild Benyo & Conrad, 1999). Consequently,

hepcidin is suppressed, enhancing maternal iron mobilisation and supporting the increase in
maternal iron demands to meet increased cardiovascular, renal, and haematological
adaptations, as well as fetoplacental development (Zaugg *et al.*, 2022). These adaptations
require sufficient iron for enhanced haematopoiesis and haemoglobin synthesis. Subsequently,
this supports optimal oxygen transfer, thereby sustaining fetal development, while also
establishing fetal iron reserves in the later stages of gestation (Cao & Fleming, 2016; Roberts
et al., 2020). Maternal iron deficiency, anaemia and excess iron alike have been implicated in
adverse pregnancy outcomes, including FGR (Andrews, 2000), highlighting the important role
of iron in healthy pregnancies.
111
Iron transport during pregnancy occurs unidirectionally from the mother to the fetus, primarily
via transferrin-bound iron in the maternal circulation (McArdle *et al.*, 2003). At the maternal
placental interface, transferrin receptor 1 (TfR1/TFRC) binds the maternal transferrin-iron
complex, inducing endocytosis (McArdle *et al.*, 2011; Roberts *et al.*, 2020). Within the
endosome, the acidic pH induces a conformational change in TfR1, releasing ferric (Fe^{3+}) iron.
Following the conversion of Fe^{3+} to ferrous (Fe^{2+}) iron, iron is transported into the cytosol via
divalent metal transporter 1 (DMT1) (McArdle *et al.*, 2011; Roberts *et al.*, 2020; Waldvogel-
Abramowski *et al.*, 2014). Once in the cytosol, iron has three potential fates, it is either: (i)
stored in ferritin, (ii) shuttled to the mitochondria for iron-sulfur (Fe-S) cluster biogenesis and
heme synthesis, or (iii) transported across the basolateral (fetal) membrane via ferroportin for
entry into the fetal circulation following reoxidation to Fe^{3+} (McArdle *et al.*, 2011; Roberts *et*
al., 2020). Approximately 20-50% of cellular iron is directed towards the mitochondria to
support two critical pathways, the biosynthesis of Fe-S clusters and heme (Qi *et al.*, 2023;
Roberts *et al.*, 2020)
126
Fe-S clusters are diverse molecular structures that perform various biological functions within
the cell. Specifically, Fe-S clusters play a role within the mitochondrial electron transport chain
(ETC), via redox reactions that transfer electrons and generate a proton gradient essential for
ATP synthesis (Mostajabi Sarhangi & Matyushov, 2023; Paul *et al.*, 2017). De novo Fe-S
cluster synthesis begins with the formation of [2Fe-2S] clusters, which serve as essential
precursors in the Fe-S cluster pathway. These de novo [2Fe-2S] clusters undergo further
processing to generate more complex [3Fe-4S] and [4Fe-4S] clusters downstream. These late-
stage [3Fe-4S] and [4Fe-4S] clusters are critical for the function and stability of the
mitochondrial ETC, particularly in complexes I, II and III, where they facilitate efficient
electron transfer during oxidative phosphorylation (Maio & Rouault, 2020; Mostajabi Sarhangi

& Matyushov, 2023; Read *et al.*, 2021). Additionally, the terminal enzyme in heme
biosynthesis, ferrochelatase (FECH), requires [2Fe-2S] clusters for its activity, linking Fe-S
cluster synthesis with heme production. Heme is essential for oxygen transport through its
incorporation into haemoglobin and cytochromes, which facilitate electron transfer in the ETC,
particularly complexes II, III, and IV (Anderson & Frazer, 2017; Kim *et al.*, 2012).
142
The primary utilisation of iron in fetal development is in erythropoiesis (Cerami, 2017), and
heme stimulates the production of α- and β-like globin genes with distinct haemoglobin profiles
expressed at specific stages of gestation (Kim *et al.*, 2012; Moksnes *et al.*, 2022) . Embryonic
($\zeta 2\varepsilon 2$) and fetal ($\alpha 2\gamma 2$) haemoglobin exhibit higher oxygen affinity compared to adult ($\alpha 2\beta 2$)
haemoglobin, demonstrated by a leftward shift in their oxygen dissociation curves. These
haemoglobin variants exhibit higher oxygen affinities during early development, enabling
adaptation to lower oxygen availability and efficient maternal–fetal oxygen transfer (Manning
et al., 2020).
151
Despite the well-described crucial role of iron in placental and fetal development, the specific
alterations in iron-dependent pathways in FGR placentae remain uncharacterised. The complex
interplay between iron transport, Fe-S cluster assembly, and heme synthesis is particularly
relevant in the context of FGR, where placental insufficiency and impaired oxygen delivery
are defining clinical characteristics. We have therefore examined genes and proteins involved
in placental iron transport, as well as iron-related mitochondrial proteins involved in Fe-S
assembly, heme synthesis, and placental erythropoiesis in FGR compared to healthy placentae.
Collectively, these results provide molecular insights into how altered iron transport and iron-
dependent pathways may contribute to mitochondrial dysfunction and placental insufficiency
observed in FGR.
162
<u>Methodology</u>
Ethical Approval
Ethical approval for this research was obtained from the University of Newcastle Human
Research Ethics Committee (H-382-0602), Hunter New England Health Human Research
Ethics Committee (02/06/12/3.13), and Site-Specific Assessment (SSA/15/HNE/291), in
compliance with the Declaration of Helsinki. Informed consent was obtained from all
participants.
170
171

Maternal and Placental Sample Collection

Placental villous cores were collected from 19 healthy, uncomplicated and 18 FGR (percentiles: $<3^{rd}$ n=7, $<5^{th}$ n=2 and $<10^{th}$ n=9) pregnancies within 45 minutes of birth as previously described by Fisher *et al.* (2024) and washed in ice-cold PBS to limit maternal blood contamination. Tissues were either preserved in formalin at 4°C overnight and stored in 0.1 M PBS with 0.05% sodium azide for histological analysis, or snap-frozen and stored at -80°C for later gene and protein analysis. FGR was diagnosed with consensus criteria as published by Gordijn *et al.* (2016). All placentae collected for this study met the eligibility criteria. Healthy term pregnancies were from singleton deliveries between 37- 40 weeks of gestation. FGR cases were from singleton pregnancies delivered by caesarean section at 34-40 weeks of gestation with an estimated fetal weight <10th centile and birthweight confirmed $<10^{th}$ centile. Exclusion criteria applied to all samples with premature membrane rupture, hypertension, diabetes, placental abnormalities, chorioamnionitis, immune deficiencies, infectious diseases, smoking, non-steroidal anti-inflammatory drug use, illicit drug use, and metformin use were not eligible within our study. Maternal blood samples collected as part of routine care were analysed retrospectively to assess clinical iron parameters. Reference ranges were obtained using New South Wales Health Pathology (NSWHP) Hospital Laboratory Clinical Projects Laboratory Manual (Brighton, 2019) in conjunction with Pregnancy and Laboratory Studies: A Reference Table for Clinicians (Abbassi-Ghanavati *et al.*, 2009).

Histological analysis of iron deposition by Prussian Blue staining

To detect $Fe^{3+,}$ samples were embedded in paraffin and sectioned at a thickness of 4 μm using a Leica RM2135 Microtome (Leica Biosystems, Germany). Fixed slides were dewaxed in a series of xylene and ethanol submersions and then washed in MilliQ water. A subset of collected placental villous tissue slides (Control n=7 and FGR n=7; : $<3^{rd}$ n=3, $<5^{th}$ n=2, $<10^{th}$ n=2) were stained with 10% (weight/weight) Prussian Blue (potassium hexacyanoferrate (II) trihydrate, Sigma-Aldrich, AUS) for 5 minutes, then 5% (weight/weight) Prussian Blue stain in 10% HCl for 30 minutes, and counterstained with nuclear fast red for 5 minutes (Sigma Aldrich, AUS) as described by Zaugg *et al.* (2024). Slides were then washed and dehydrated following gradient ethanol exposure of 70% and 100% for 3 minutes, finishing with a 10-minute xylene exposure. Slides were mounted with a coverslip and left to dry overnight. Slides were then imaged on Aperio GT 450 DX (Leica Biosystems, Germany) at 40X magnification. The intensity of the Prussian blue staining was quantified using H-scores obtained from Halo

205 Quantitative Pathology analysis software (Indica Labs; USA). All images were processed using
206 identical parameters to ensure valid comparisons between healthy and FGR placental samples.
207

Placental RNA Preparation and Gene Expression Studies

Placental tissue was crushed on dry ice with liquid nitrogen and subsequently, lysed and
homogenised with 1 mL Trizol® Reagent (Life Technologies; Thermo Fisher Scientific, AUS)
using the Precellys24 instrument (Bertin Technologies, France) for two cycles of 30 seconds at
$5000 \times g$, incubated on ice for 30 minutes, and then repeated. Samples were centrifuged for 10
minutes at $10,000 \times g$ using the Microfuge 20R (Beckman Coulter, AUS). The remainder of
the protocol was performed in line with the manufacturer's instructions outlined in the Direct-
zol^{TM} RNA mini-prep kit (Zymo Research, California, USA). Following quantification on the
Nanodrop One (Thermo Scientific, AUS), 1000 ng of cDNA was synthesised using
SuperScript^{TM} IV First-Strand Synthesis System (Thermo Fisher Scientific, AUS). RT-qPCR
was conducted using the PowerUpTM SYBR Green Master Mix (Applied Biosystems, Thermo
Fisher, AUS) with KiCqstart primers (Sigma Aldrich, AUS; Table 1). RT-qPCR was performed
using the QuantStudio6 Flex Real-Time PCR system (Thermo Fisher Scientific, MA, USA).
Relative gene expression was determined by normalising to the geometric mean of three
housekeeping genes (β-Actin, TBP, and YWHAZ), which were quantified using the $2^{-\Delta\Delta CT}$
method to assess relative gene expression within placental tissue samples.
224

225 **Table 1: Kicqstart Primers for Gene Expression Studies of Targeted Iron Transporters**
226 **and Intermediaries.**

Target Gene Name	Gene Abbreviation	Primer sequence
Cellular Iron Transporters		
Transferrin Receptor 1	TFRC	F 5'-AAGATTCAGGTCAAAGACAG-3'
		R 5'-CTTACTATACGCCACATAACC-3'
Solute Carrier Family 11 Member 2	SLC11A2	F5'-GAGTATGTTACAGTGAAACCC-3'
		R 5'-GACTTGACTAAGGCAGAATG-3'
Solute Carrier Family 40 Member 1	SLC40A1	F 5'-AAAGATACTGAGCCAAAACC-3'
		R 5'-GTTGTAGTAGGAGACCCATC-3'
Mitochondrial Iron transporters		
Solute Carrier Family 25 Member 37/ Mitoferrin 1	SLC25A37	F 5'-GGTAATGAATCCAGCAGAAG-3'
		R 5'-AGGAACTCATAGGTGATGAAG-3'

Solute Carrier Family 25 Member 28/ Mitoferrin 2	SLC25A28	F 5'-ACCTTTGATAACCTCTCTCC-3' R 5'-TATTAGGAACCAGGGGAATG-3'

Mitochondrial Heme Synthesis

Coproporphyrinogen Oxidase	CPOX	F 5'-CATGGAAATCTTTCAGAGGAAG-3' R 5'-ATTCTTGGGGTGGATAACAG-3'
Ferrochelatase	FECH	F 5'-GGAAGAATATCCTCTTGGTTC-3' R 5'-CACTCCTTGGCTAAAACTTG-3'

Erythropoiesis

Hemogen	HEMGN	F 5'-CAAGATGGATTTGGGAAAGG-3' R 5'-TCTGGAGAATGGTTCTCTTC-3'
Haemoglobin Subunit Alpha	HBA	F 5'-GACCTCCAAATACCGTTAAG-3' R 5'-ACTTTATTCAAAGACCACGG-3'
Haemoglobin Subunit Beta	HBB	F 5'-ATTTGCTTCTGACACAACTG-3' R 5'-CAAAGGACTCAAAGAACCTC-3'
Haemoglobin Subunit Gamma 1	HBG1	F 5'-ACTTCCTTGGGAGATGCCAC-3' R 5'-GCCTATCCTTGAAAGCTCTGA-3'
Haemoglobin Subunit Gamma 2	HBG2	F 5'-ACTTCCTTGGGAGATGCCAT-3' R 5'-GCCTATCCTTGAAAGCTCTGC-3'
Erythrocyte Membrane Protein Band 4.1	EPB41	F 5'-AAAGGACTTGGAAGGAGTAG-3' R 5'-GCGGTTAATTCTCAGCTTATC-3'
Solute Carrier Family 4 Member 1	SLC4A1	F 5'-CAAGAACAGCTCCTATTTCC-3' R 5'-CTGAATGAAGAAATCCACCAG-3'

Iron-Sulfur Cluster biosynthesis

LYR Motif Containing 4	LYRM4/ISD11	F 5'-AGGATAAGAGATGCCTTCAG-3' R 5'-ATCTGTCGACGAATTACTCC-3'
NADH: Ubiquinone Oxidoreductase Subunit AB1	NDUFAB1	F 5'-CAGGTTCCTTGGTAGAGTTAC-3' R 5'-TGGGTCAATCTTGTCATAGAG-3'
Ferredoxin 2	FDX1L/FDX2	F 5'-CATGTGTATGTGAGTGAAGAC-3' R 5'-TAGAAGGTTCCTGGTGATCTTG-3'
Frataxin	FXN	F 5'- GAAAGACTAGCAGAGGAAAC-3' R 5'-TCCCAAAGGAGACATCATAG-3'

Iron-Sulfur Cluster Assembly Enzyme	ISCU	F 5'- GTGTAGACCTTTCTGCTCAGG-3' R 5'-TAGATGTCTTGTCAAGGGAC-3'
Glutaredoxin 5	GLRX5	F 5'-AAGAAGGACAAGGTGGTG-3' R 5'-TAGTCTTTAATGCCTTGTCG-3'
The NFU1 Iron-Sulfur Cluster Scaffold	NFU1	F 5'-CCCTAGTTCAATCATTACTCTG -3' R 5'-CATCATCCATAACCTGTTCTAC- 3'
NUBP Iron-Sulfur Cluster Assembly Factor	NUBPL	F 5'-TGAAGTTCTAGGAGACATTCC-3' R 5'-TCTTCTTACCACTTCCACAG-3'
BolA Family Member 3	BolA3	F 5'-CGAGCTACAGCTATAAAAGTC-3' R 5'-TTGGGGACAGAGGTAAATATC-3'
Housekeeper Genes		
Beta Actin	β-Actin	F 5'-GACGACATGGAGAAAATCTG-3' R 5'-ATGATCTGGGTCATCTTCTC-3'
Tryptophan 5-Monooxygenase Activation Protein Zeta	YWHAZ	F 5'-CCTGCATGAAGTCTGTAACTGAG-3' R 5'-GACCTACGGGCTCCTACAACA-3'
TATA-Box Binding Protein	TBP	F 5'-GCCAAGAGTGAAGAACAG-3' R 5'-GAAGTCCAAGAACTTAGCTG-3'

227 KiCqStart Primers targeting nuclear DNA involved in placental transport and iron-dependent
228 pathways. Forward (F) and Reverse (R) primers were reconstituted to a 100 μM final
229 concentration following the manufacturer's instructions. Gene expression was analysed using
230 the $2^{-\Delta\Delta CT}$. Genes were examined in response to the physiological function of Iron in healthy
231 and FGR placentae.
232
Proteomics Studies:
Placental tissue was crushed on dry ice and stored at -80°C until Liquid Chromatography-Mass
Spectrometry (LC-MS) based proteomic analysis could be performed. LC-MS sample
preparation followed previously published methodology by Mulhall *et al.* (2024), with
placental tissue samples lysed in chilled 4% (weight/volume) sodium deoxycholate (SDC), 100
mM Tris HCl (pH.8.5) and passed through an 18G blunt-end needle before sonication.
Samples were quantified and concentrations equalised to 200 μg/μl before digestion, alkylation
and overnight trypsin incubation. Following trypsin exposure, peptides were diluted 1:1 with
100 mM Tris-HCl (ensuring SDC concentration < 1%) and vortexed briefly before adding 99%
ethylacetate /1% Trifluoroacetic acid (TFA; 1:1) and thoroughly vortexed at 2000 rpm using

ThermoMixer. Peptides were then desalted using styrene-divinylbenzene reverse-phase sulfonated StageTips.

Peptide separation was performed on an Aurora Ultimate C18 column (25 cm × 75μm, IonOpticks) using a 90-minute linear gradient from 3% to 41% of solvent B (acetonitrile in 0.1% formic acid) at a flow rate of 400 nL/min. MS analysis was conducted on an Eclipse mass spectrometer (Thermo Fisher Scientific, AUS), and MS scans were acquired over a mass range of 375-1400 m/z at a resolution of 60,000, with an automatic gain control (AGC) target of 100% and maximum injection time of 50 ms. MS/MS scans were acquired at a resolution of 15,000, with an AGC target of 100%, normalised collision energy of 30%, and dynamic maximum injection time. Raw data was analysed using Proteome Discoverer 2.5 (Thermo Fisher Scientific, AUS), outlined by Murray *et al.* (2023).

Western Blotting

To validate our proteomics data, western blotting was performed. Protein was extracted from healthy (n=7) and FGR (n=7) placental tissue using 1 mL RIPA buffer (50 nM Tris-HCl (pH 7.4), 150 nM NaCl, 1% NP-P40, 1% sodium deoxycholate, 1% SDS, Complete Mini Protease Inhibitor Cocktail tablets (Roche Diagnostics, North Ryde AUS) and 1 nM PMFS). Samples were homogenised using Precellys24 and centrifuged as outlined previously. The supernatant was collected and stored at -80°C for subsequent western blotting analysis. Following Fisher *et al.* (2019) published protocol, samples (40 μg/μL) were loaded onto 12% Tris gels (Invitrogen, AUS), electrophoresed, and transferred to Polyvinylidene fluoride (PVDF) membranes. Membranes were blocked with Odyssey Intercept buffer (LICORBio, USA) for 1 hour, then incubated overnight with primary antibody, LYMR4 at 1:250 dilution (ab253001, Abcam, AUS) and Beta Actin (ab8226, Abcam, AUS) at a 1:1000 dilution. After washing with PBST, membranes were incubated for 1 hour with secondary antibodies anti-mouse IRDye®800CW and anti-rabbit IRDye®680LT (LICORBio, USA) at 1:20,000 dilution and washed with PBST. Membranes were imaged using the LI-COR Odyssey M system (LICORBio, USA) and analysed with Studio Image v5.2.

Assessment of Heme

Placental heme concentrations were measured using a Heme Assay kit (MAK316, Sigma Aldrich, AUS). Following the manufacturing instructions, a total of 250 μL of water was added to a 96-well plate, creating a blank well. A standard well was created by mixing 50 μL of heme calibrator and 200 μL of water. Subsequently, 50 μL of homogenised term and FGR samples

suspended in glycerol buffer (50 mM Tris HCl, 1 mM EDTA, 0.5% Glycerol) was mixed with
200 μL of reagent respectively. The plate was left to incubate for 5 mins at room temperature,
and the absorbance of well contents was determined at 400 nm. All samples and standards were
performed in duplicate, and total heme concentration was determined using the following
equation:

$$[\text{Total Heme}]\left(\frac{\mu\text{mol}}{\text{L}}\right) = \frac{(\text{ODsample} - \text{ODblank})}{(\text{ODcalibrator} - \text{ODblank})} \text{ x } 62.5 \left(\frac{\mu\text{mol}}{\text{L}}\right)$$

284
Statistical Analysis
All data were analysed using GraphPad Prism 10.0. Outliers were identified and removed using
Grubbs and ROUT tests, and data normality was assessed using the Shapiro-Wilk test. For
normally distributed data, student t-tests were performed. For data that did not follow a normal
distribution, Welch t-tests were conducted. Statistical significance was set at $p < 0.05$. John
Hunter Hospital clinical data are presented with 95% confidence intervals (CI) and interquartile
ranges (IQR). Gene expression data were presented as mean ± standard error of the mean
(SEM), and histological H-scores, accounting for blue-stained iron deposits within placental
tissue, were quantified using Halo Quantitative Pathology Analysis Software (Indica Labs,
USA) and calculated using the following formula (Wen *et al.*, 2024):

$$\text{Hscore} = (3 \times \% \text{ heavily stained}) + (2 \times \% \text{ moderately stained})$$
$$+ (1 \times \% \text{ lightly stained})$$

This formula weights the staining intensity, generating a composite score ranging from 0-300,
reflecting both the extent and intensity of iron deposits. Proteomic analysis was expressed as
$\log_2$-fold change (FC) to determine differential protein expression between healthy controls
and FGR, visualised using heatmaps. The $\log_2$-fold was derived from the ratio between healthy
term placentae to assess changes in FGR placentae, so a positive $\log_2$-FC indicates higher
protein abundance in healthy control placentae, whereas a negative $\log_2$-FC indicates higher
protein abundance in FGR placentae. Peptides with fewer than 2 Peptide Spectrum Matches
(PSMs) were excluded from the analysis.
305
Results
Clinical Insights into Maternal Iron Bioavailability, Birth and Gestational Characteristics
To better understand the potential role of iron throughout gestation, we evaluated clinical
characteristics encompassing maternal physical attributes, birth-related factors, and gestational
characteristics (Table 2). Analysis included maternal parameters (age and BMI), birth outcomes
(gestational age at delivery, birth weight, birth centile, placental weight, fetal sex, mode of

delivery) and fetal biometry measurements following sonography assessment (estimated fetal weight (EFW%), abdominal circumference, femur length, head circumference, and amniotic fluid volume). FGR babies were significantly smaller (p<0.0001), with a median birth weight of 2415g (95% CI: 1760g-2356g), compared to healthy term babies, who had a median birth weight of 3555g (95% CI: 3346g- 3676g). Additionally, FGR babies had smaller placentae (p<0.0001), with a median placental weight of 556g (95% CI: 395g-519g), compared to healthy term babies, who had a median placental weight of 689g (95% CI: 628g-757g). Furthermore, fetal biometric measurements, including abdominal circumference (p<0.0001), femur length (p=0.002), and head circumference (p=0.0007), were all significantly lower in FGR pregnancies. Although the difference in amniotic fluid volume was not statistically significant, FGR pregnancies had lower amniotic fluid volume (p=0.054) compared to healthy term controls.

Maternal haematological indices and maternal haematocrit and anaemia screening (Table 2) were performed on healthy and FGR pregnancies. No clinical or statistical significance was observed between FGR and healthy control groups in haemoglobin (Hb), mean corpuscular haemoglobin (MCH), mean corpuscular haemoglobin concentration (MCHC), mean corpuscular volume (MCV), red cell count (RCC), hematocrit (HCt), systemic iron levels and transferrin (Tf). Maternal ferritin levels were higher (p=0.029) and red cell distribution width (RDW) lower (p=0.006) in FGR cases compared to healthy control, however, both parameters fell within clinical reference ranges.

Table 2: Maternal, Fetal and Birth Clinical Characteristics from Pregnancies Diagnosed as FGR And Healthy Term Controls.

	Term Control	FGR	P value
n	19	18	
a. Maternal and Birth Characteristics			
Maternal Age (years)	31 (27.8, 33.0)	28.5 (26.9, 31.0)	0.3631
Maternal Body mass index; BMI (kg/m^2)	26.0 (24.2, 28.2)	25.0 (22.4, 30.6)	0.8836
Birth weight (g)	3555 (3346, 3676)	2415 (1760, 2356)	**< 0.0001**

Placental weight (g)	689 (628,757)	556 (395, 519)	**< 0.0001**
Sex ratio	9 Males, 10 Females	8 Males, 10 Females	-

b. Fetal Biometry Measurements following Sonography Assessment

Fetal abdominal circumference (mm)	330 (321, 339)	275 (257, 293)	**< 0.0001**
Fetal Femur Length (mm)	68.4 (66.3, 73.6)	61.6 (58.0, 65.2)	**0.002**
Fetal Head Circumference (mm)	328.0 (315.2, 347.6)	295.5 (282.4, 307.8)	**0.0007**
Amniotic Fluid Volume (cm)	12.6 (10.8, 15.4)	10.9 (7.99, 12.1)	0.054
UAD Pulsatility Index	0.950 (0.820,1.50)	1.10 (0.860, 1.41)	0.823
UAD Resistance Index	0.585 (0.530, 0.790)	0.620 (0.560, 0.730)	0.5476

c. Maternal Haematological Parameters

Haemoglobin (Hb) *RI: 10.5-15.0g/dL*	12.7 (12.1, 13.0)	12.2 (11.8,12.7)	0.325
Mean Corpuscular Haemoglobin (mcH) *RI: 29.0-33.0pg/cell*	30.0 (28.4, 30.4)	30.0 (28.9, 30.8)	0.519
Mean Corpuscular Haemoglobin concentration (mcHc) *RI:310-360g/L*	345 (338, 352)	344 (338, 346)	0.437
Haematocrit (HCt) *RI: 38.0-41.0%*	37.6 (36.1,39.5)	37.1 (35.9, 38.7)	0.620
Mean Corpuscular volume (McV) *RI: 81-99 (μm^3)*	88.0 (86.1, 90.0)	88.0 (86.4, 90.4)	0.8
Red Cell Count (RCC) *RI: 2.81-4.55 $10^6/mm^3$*	4.09 (4.02, 4.34)	4.00 (3.86, 4.13)	0.081
Red Cell Distribution Width (RDW)	14.2 (13.8, 15.0)	13.4 (13.1, 13.7)	**0.006**

RI: 12.5-15.3%			

d. Maternal Haematocrit and Anaemia Screening			
Iron (Fe) RI: 5.37-34.55µmol/L	22.0 (14.5, 24.4)	15.50 (12.1, 20.4)	0.29
Transferrin (Tf) RI: 2.2-5.3g/dL	3.50 (2.97, 4.12)	3.20 (2.82, 3.76)	0.513
Ferritin (F) RI: 30-200µg/L	20.0 (14.8, 33.3)	46.0 (25.4, 67.7)	**0.029**

This table presents data from consenting participants at John Hunter Hospital. Data are presented as median and interquartile range (IQR). Shapiro-Wilk normality tests were conducted, with student t-tests or Welch's tests used where appropriate. Statistically significant values ($p<0.05$). Missing data are indicated by –.

Placental Iron Transport

Analysis of iron transporters between FGR and control placentae revealed significant alterations in gene expression and protein levels of iron import and export components. Transferrin Receptor 1 (TFRC) showed significantly increased mRNA expression ($p<0.005$, Figure 1A) and upregulated protein levels (-0.117 log_2-FC; Figure 1D) in FGR placentae compared to healthy term placentae. Divalent Metal Transporter 1 (DMT1/SLC11A2) exhibited significantly increased mRNA expression in FGR ($p<0.05$, Figure 1B) compared with controls. In contrast, Ferroportin (FPN/SLC40A1; Figure 1C), the iron receptor responsible for transporting iron out of the placenta into fetal circulation, showed no significant difference in mRNA expression between FGR and healthy controls. FPN protein analysis (Figure 1D) revealed lower protein levels (0.646 log_2-FC) in FGR placental villous tissue. These findings suggest FGR placentae are retaining iron for placental function. Following the clinical characterisation of maternal iron parameters and mRNA expression of iron transport into the placenta, the placental iron content between healthy and FGR placentae was assessed. Histological analysis by Prussian blue staining revealed intense blue regions with no statistical difference between healthy control (Figure 1F-H) and FGR (Figure 1I-K) placentae.

Figure 1: mRNA expression and protein levels of cellular iron transporters and histological Prussian blue staining in FGR and healthy control placental villous tissue. (A) TFRC, Transferrin Receptor 1; (B) SLC11A2, Solute Carrier Family 11 Member 2/ Divalent Metal Transporter 1 (DMT1); (C) SLC40A1, Solute Carrier Family 40 Member 1/Ferroportin (FPN). Data presented as mean ± SEM, control n=11-19 (white and circles), FGR n=10-18 (grey and triangles). Student t-tests were performed, *p<0.05, **p<0.01, ***p<0.001, ****p<0.0001. (D) Proteomes of cellular iron transport depicted as a heatmap based on the log2fold, with transporters indicated on top. The log_2-FC was derived from the ratio between healthy term placentae to assess changes in FGR placentae, so a positive log_2-FC indicates higher protein abundance in healthy control placentae (orange), and a negative log_2-FC indicates higher protein abundance in FGR placentae (blue). (E) Halo Pathology Analysis Software was used to quantify the percentage of Prussian blue staining relative to the tissue area. Heavy, moderate, and light intensity values were used to calculate the H-score in healthy control (n=7) and FGR (n=7) placental tissue. An unpaired t-test revealed no significant difference between the groups. (F) A representative merged image of healthy control placental tissue showing Fe^{3+} deposits (blue staining indicated by arrows). (G) Control placental tissue stained with nuclear fast red counterstain only and (H) Prussian blue staining isolated from

counterstained tissue in (F), upon which intensity calculations were performed. (I) A representative merged image of FGR placental tissue, with (J-K) shown similarly to the row above (G-H). The scale bar represents 100μm.

Mitochondrial Iron Transport

Given the pivotal role of iron in mitochondrial function, we examined mitochondrial iron transporters located within the inner mitochondrial membrane, Mitoferrin 1 (Solute Carrier Family 25 Member 37, SLC25A37) and Mitoferrin 2 (Solute Carrier Family 25 Member 28, SLC25A28). No significant difference in mitoferrin 1 (Figure 2A) mRNA expression was observed in FGR placentae compared to healthy controls. In contrast, mitoferrin 2 (Figure 2B; $p < 0.005$) mRNA expression was significantly lower in FGR placental villous tissue than in healthy controls.

Mitochondrial Iron Utilisation: De Novo Fe-S Cluster Synthesis

The utilisation of iron within the mitochondria converges on the synthesis of Fe-S clusters. This process comprises two main stages: de novo synthesis of [2Fe-2S] clusters and late-stage genesis of [4Fe-4S] clusters vital to mitochondrial function and cellular viability. In de novo Fe-S cluster synthesis, the mRNA expression of LYR Motif 4 (LYRM4/ISD11; Figure 2C), NADH: Ubiquinone Oxidoreductase Subunit AB1 (NDUFAB1/ACP; Figure 2D) and Frataxin (FXN; Figure 2E) were similar in FGR and healthy control placental villous tissue. Conversely, protein levels of NDUFAB1 (0.123 $\log_2$-FC; Figure 2L) were lower in FGR placentae compared with healthy controls. Ferredoxin 2 (FDX2/FDX1L) mRNA expression (Figure 2G) was significantly lower ($p < 0.05$), while ferredoxin reductase (FDXR; Figure 2L) was 1.08 $\log_2$-fold lower in protein levels in FGR compared to controls. The mRNA expression of Iron-Sulfur Cluster Assembly Scaffold Enzyme (ISCU; Figure 2F), which is essential for [2Fe-2S] assembly and transfer, was significantly lower ($p < 0.05$) within FGR placental tissue compared with healthy controls. However, ISCU protein expression (-0.372 $\log_2$-FC; Figure 2L) was higher in FGR placental villous tissue when compared with healthy controls. An important transporter of the [2Fe-2S] cluster, Glutaredoxin 5 (GLRX5; Figure 2H), had significantly lower mRNA expression ($p < 0.05$) in FGR placentae; however, no difference in protein abundance between FGR and healthy term placentae was observed.

Late-Stage Fe-S Cluster Synthesis

In late-stage Fe-S cluster assembly, Heat Shock Protein Family A (HSP70) Member 9 (HSPA9; Figure 2L) observed a 0.219 $\log_2$-fold increase at the protein level in healthy controls compared

to FGR placentae. IBA57 (Figure 2L) was -0.073 $\log_2$-fold higher in protein level with statistical significance ($p < 0.005$) confirmed by western blotting (Figure 2M). [4Fe-4S] cluster transporters showed increased mRNA expression of NFU1 Iron-Sulfur Cluster Scaffold (NFU1; $p < 0.005$, Figure 2I) and BOLA Family member 3 (BOLA3; $p < 0.05$, Figure 2J) in FGR placental villous tissue when compared to healthy controls. No significant changes in mRNA expression were observed in NUBP Iron-Sulfur Cluster Assembly factor (NUBPL; Figure 2K).

Figure 2: mRNA expression and proteomes of targeted genes in the iron-sulfur (Fe-S) cluster biosynthesis pathway. (A) SLC25A37, Solute Carrier Family 25 Member 37/ Mitoferrin 1; (B) SLC25A28, Solute Carrier Family 25 Member 28/Mitoferrin 2 (C) LYRM4, LYR Motif Containing 4 (ISD11); (D) NDUFAB1, NADH: Ubiquinone Oxidoreductase Subunit AB1; (E) ISCU, Iron-Sulfur Cluster Assembly Enzyme; (F) FXN, Frataxin; (G) FDX1l, Ferredoxin 2 (FDX2); (H) GLRX5, Glutaredoxin 5; (I) NUBPL, NUBP Iron-Sulfur Cluster Assembly Factor; (J) NFU1, NFU1 Iron-Sulfur Cluster Scaffold; (K) BolA3, BolA Family Member 3; Data are presented as mean ± SEM, control n=11-19 (white and circles), FGR n=10-18 (grey and triangles). Student t-tests were performed *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$, ****$p < 0.0001$. (L) Fe-S cluster proteomes, depicted as a heatmap based on $\log_2$-

FC with proteome identification on top. Proteomes of Fe-S cluster assembly depicted as a

heatmap based on the $\log_2$-FC, with transporters indicated on top. The $\log_2$-FC was derived

from the ratio between healthy term placentae to assess changes in FGR placentae, so a positive

$\log_2$-FC indicates higher protein abundance in healthy control placentae (orange), and a

negative $\log_2$-FC indicates higher protein abundance in FGR placentae (blue). (M) Protein

abundance of putative transferase CAF17 (IBA57; 38.155kDa) normalised to β-Actin

(41.737kDa) in healthy and FGR placentae. Representative immunoblot shown below.

Heme Synthesis

Analysis of the heme synthesis pathway showed significantly higher mRNA expression of

coproporphyrinogen oxidase (CPOX; p=0.002; Figure 3A) and ferrochelatase (FECH;

p=0.005; Figure 3B) at the gene and protein level, -0.163 $\log_2$-fold and -0.356 $\log_2$-fold,

respectively (Figure 3D) within FGR placental tissue compared to healthy tissue. Furthermore,

protein levels of additional enzymes important to heme synthesis, Uroporphyrinogen

Decarboxylase (UROD; Figure 3D) Hydroxymethylbilane Synthase (HMBS; Figure 3D), were

-0.175 $\log_2$-fold and -0.196 $\log_2$-fold greater in FGR, respectively. In contrast, Aminolevulinate

dehydratase (ALAD/PBGS; Figure 3D) was 0.541 $\log_2$-fold lower protein expression in FGR

compared with term control placental villous tissue. Although heme synthesis enzymes were

upregulated, this was not conserved when assessing heme activity (p<0.005) in FGR placentae.

Heme oxygenase (HMOX) -1 protein abundance was -1.13 $\log_2$-fold higher in FGR placentae

compared to control placentae (Figure 3D). Additionally, NADPH-cytochrome P450

oxidoreductase (POR) protein abundance was -0.157 $\log_2$-fold higher in FGR placentae than

healthy controls.

Figure 3: mRNA expression of targeted genes involved in the heme synthesis pathway and

heme content in FGR and healthy term placentae. (A) CPOX, Coproporphyrinogen

Oxidase; (B) FECH, Ferrochelatase. Data presented as mean ± SEM, control n=11-19 (white

and circles), FGR n=10-18 (grey and triangles), with sample loss due to outliers' tests

conducted. Student t-tests were performed *p<0.05, **p<0.01, ***p<0.001, ****p<0.0001.

(C) Heme Assay measuring total Heme Concentration (μM) present. **(D)** Heme synthesis

proteomes depicted as a heatmap based on $\log_2$-FC, with proteome identification on top.

Proteomes of heme synthesis depicted as a heatmap based on $\log_2$-FC, with transporters

indicated on top. The $\log_2$-FC was derived from the ratio between healthy term placentae to

assess changes in FGR placentae, so a positive $\log_2$-FC indicates higher protein abundance in

healthy control placentae (orange), and a negative $\log_2$-FC indicates higher protein abundance

in FGR placentae (blue).

The placenta is a potent haemopoietic site where heme synthesis stimulates globin production,

forming a haemoglobin tetramer of alpha and beta-like chains critical for binding and

transporting oxygen. Our analysis revealed no changes in the mRNA expression of

Haemoglobin Subunit Alpha 1 (HBA1) between controls and FGR placental villous tissue,

although protein levels were 0.210 $\log_2$-fold lower in FGR compared to healthy controls

(Figure 4A and G). Within the beta-like globin chains, we observed significantly greater mRNA

expression of Haemoglobin Subunit Gamma 1 (HBG1; $p<0.05$, Figure 4B) in FGR placentae,

although Haemoglobin Subunit Gamma 2 showed no differences between control and FGR

placentae. Protein abundance of HBG1 (0.325 $\log_2$-FC) and HBG2 (0.092 $\log_2$-FC) was lower

in FGR compared with term control placental tissue samples (Figure 4G). Haemoglobin

Subunit Beta (HBB) mRNA expression was significantly higher ($p<0.05$) in FGR compared to

healthy controls (Figure 4D). However, HBB protein abundance was lower (0.585 $\log_2$-FC)

within FGR placental tissue when compared to healthy placental tissue samples (Figure 4G).

Levels of Erythrocyte structural proteins, specifically Erythrocyte Protein Band 4.1 (EPB41),

EPB42, Spectrin Alpha 1 (SPTA1), Spectrin Beta (SPTB), Ankyrin1 (ANK1) and Solute

Carrier Family 4 Member 1 (SLC4A1), were subsequently assessed. The mRNA expression

($p<0.05$) and protein abundance (0.254 $\log_2$-FC) of EPB41 were lower in FGR placental tissue

compared to healthy tissue controls (Figure 4E and G). EPB42 protein abundance was 0.562

$\log_2$-fold lower within FGR placentae than healthy placentae (Figure 4G). Analysis of spectrin

protein levels, SPTA1 and SPTB (Figure 4G), was 0.156 $\log_2$-fold and 0.491 $\log_2$-fold lower

in FGR placenta compared with healthy term placenta. Similarly, ANK1 and SLC4A1 protein

abundance were 0.661 $\log_2$-fold and 0.493 $\log_2$-fold lower in FGR placentae than healthy

placentae, suggesting compromised erythrocyte structure. However, analysis of Hemogen

(HEMGN; Figure 4F), a regulator of haematopoietic cell differentiation, mRNA expression

was similar in healthy and FGR placental villous tissue, indicating that these changes were not

occurring at the onset of hematopoiesis but rather at erythrocyte maturation.

Figure 4: mRNA expression and proteome of targeted genes involved in Haematopoiesis. (A) HBA1. Haemoglobin Subunit Alpha 1; (B) HBG1, Haemoglobin Subunit Gamma 1; (C) HBG2, Haemoglobin Subunit Gamma 2; (D) HBB, Haemoglobin Subunit Beta; (E) EPB41, Erythrocyte Membrane Protein Band 4.1; (F) HEMGN, Hemogen; Data presented as mean ± SEM, control n=11-19 (white and circles), FGR n=10-18 (grey and triangles). Student t-tests test performed *p<0.05, **p<0.01, ***p<0.001, ****p<0.0001.(G) Haematopoiesis proteomes, depicted as a heatmap and based on log$_2$-FC with proteome identification on top. Proteomes of haematopoiesis depicted as a heatmap based on log$_2$-FC, with transporters indicated on top. The log$_2$-FC was derived from the ratio between healthy term placentae to assess changes in FGR placentae, so a positive log$_2$-FC indicates higher protein abundance in healthy control placentae (orange), and a negative log$_2$-FC indicates higher protein abundance in FGR placentae (blue).

Discussion

This project sought to elucidate the complex role of iron in placental and mitochondrial function in fetal growth restriction (FGR). As expected, we observed reduced placental and birth weights in FGR compared with healthy controls (Table 2). Similarly, fetal biometric measurements showed significantly reduced abdominal circumference, head circumference, and femur length in FGR pregnancies compared with healthy-term pregnancies. Mothers with FGR babies also showed reduced red cell distribution width (RDW) in conjunction with elevated ferritin levels, although these fell within clinical reference ranges. No changes were

observed in circulating maternal iron or haemoglobin levels, and all maternal measurements stayed within the clinical reference ranges (Table 2). Subsequent analysis of FGR placental tissue identified altered iron transporters (Figure 1). Furthermore, FGR placentae showed changes in mitochondrial iron-sulfur (Fe-S) cluster biosynthesis, specifically a reduced capacity for converting [2Fe-2S] to [4Fe-4S] clusters due to lower levels of key [2Fe-2S] proteins ferredoxin reductase (FDXR) and ferredoxin 2 (FDX2) required for synthesis of both [2Fe-2S] and late-stage [4Fe-4S] clusters (Figure 2). The observed reduction in proteins involved in [4Fe-4S] cluster assembly suggests a metabolic redirection from incorporation into the mitochondrial electron transport chain (ETC) toward increased heme production, as evidenced by increased cytosolic and mitochondrial proteins of the heme synthesis proteins (Figure 3). Despite the apparent shift toward heme synthesis, total heme concentrations in FGR placentae were decreased. We propose that this arises from increased regulation of the heme degradation pathway in FGR placentae, evidenced by elevated protein levels of heme oxygenases and NADPH-cytochrome P450 oxidoreductase. Consequently, we observed reduced haemoglobin proteins within the FGR placentae (Figure 4). Additionally, our findings identified alterations in levels of erythrocyte structural proteins in FGR placentae (Figure 4). These alterations may further compromise oxygen delivery to the fetus or simultaneously represent an adaptive response to the poor vasculature of FGR placentae. Together, these findings enhance our understanding of the fundamental biological processes underlying FGR and highlight how adaptations in iron utilisation contribute to mitochondrial dysfunction observed in FGR, perpetuating inadequate fetal growth.

Maternal Haematological Assessment in FGR

Pregnancy is a dynamic process that is associated with significant alterations in the maternal cardiovascular system to support fetoplacental development (Sanghavi & Rutherford, 2014). Maternal cardiovascular adaptations, including increased plasma volume (20-50%) and red blood cell mass (20-30%), begin rising from week 5 of pregnancy, well before the maternal and fetal circulatory systems are functionally established via the placenta (Allerkamp *et al.*, 2022; Mégier *et al.*, 2022; Söhnchen *et al.*, 2011). Additionally, studies have shown that erythropoiesis is the largest consumer of iron and as gestation advances, the demand for iron increases substantially to support expanded maternal red cell mass, placental development, and fetal growth (Gambling *et al.*, 2001; Kautz & Nemeth, 2014). Maternal cardiovascular function and haemodynamic failures have significant implications for the health of both the mother and the fetus.

In FGR pregnancies, normal maternal haemodynamic responses are known to be compromised,

including inadequate blood volume expansion and impaired reduction in vascular resistance

(Mecacci *et al.*, 2021; Perry *et al.*, 2020). Therefore, our study assessed whether maternal

haemodynamic responses in FGR pregnancies might be reflected in maternal blood parameters

(Table 2). We observed reduced maternal red blood cell distribution width (RDW) in FGR

mothers compared with controls. RDW is the measurement of variation in red blood cell size

and volume relative to the mean red cell volume (MCV) (Fava *et al.*, 2019; Shehata *et al.*,

1998). Despite being statistically significant, the reduction in maternal RDW remained within

clinical reference ranges, indicating decreased anisocytosis, less variability in erythrocyte size

and a narrower distribution around the mean cell volume (de Freitas *et al.*, 2019; Kurt *et al.*,

2015; Lippi & Plebani, 2014). We speculate that the statistical significance in maternal RDW

between FGR and healthy control mothers, while within clinical reference ranges, may be a

maternal adaptation to overcome poor spiral artery remodelling. Improved erythrocyte

uniformity may enhance blood rheology (Nader *et al.*, 2019; Reddy *et al.*, 2016; Zhang *et al.*,

2009), which may provide advantages in the context of increased vascular resistance and

narrowed maternal spiral arteries, contributing to the reduced perfusion of the intervillous space

often observed in FGR (Burton & Jauniaux, 2018).

Maternal serum ferritin levels were significantly higher in FGR pregnancies than in healthy

controls (Table 2), although ferritin levels fell within clinical reference ranges (30-200 µg/L).

Our observations are substantiated by previous studies showing that elevated serum ferritin

levels increased the risk of developing FGR (Ozgu-Erdinc *et al.*, 2014). Furthermore, Hou *et*

al. (2000) demonstrated that FGR pregnancies with elevated serum ferritin levels presented

with asymmetric fetal growth during the second and third trimesters. Asymmetrical growth is

a key hallmark of late-onset FGR (diagnosis >32 weeks), accounting for 70-80% of FGR cases

(Sun *et al.*, 2020). This may explain why serum ferritin levels were higher in FGR cases within

our study, as 83% of our samples were delivered >32 weeks. Moreover, serum ferritin's

specificity (true negative rate in identifying iron sufficiency) and sensitivity (true iron

deficiency) are essential for assessing biological iron status when chronic inflammation is

absent (Mégier *et al.*, 2022). The increase in maternal serum ferritin may, therefore, be

attributed to an inflammatory response as ferritin acts as an acute-phase reactant during

inflammation (Camaschella, 2015), rather than truly indicating maternal iron status, as

pregnancy is inherently pro-inflammatory. This is supported by increases in oxidative stress

and inflammation observed in the maternal circulation and placenta of FGR pregnancies (Hou
et al., 2000; Ozgu-Erdinc *et al.*, 2014).
582
583 Despite statistically elevated maternal serum ferritin levels and reduced RDW in FGR, the lack
584 of maternal changes in iron haematological parameters, outside of normal reference ranges,
585 suggests that subsequent findings were not due to impaired iron delivery but rather demonstrate
586 that maternal adaptations are likely secondary responses to compromised placental function.
587
Iron transport in the FGR placenta.
Our study revealed that iron transporter proteins were altered in FGR placentae, characterised
by an enhanced capacity for iron uptake. Specifically, we observed a 1.08-fold increase in
Transferrin receptor 1 (TfR1/TFRC) levels (-0.117 log2-fold), the major importer of non-heme
iron into cells, in FGR placenta (Figure 1A and D). While maternal iron deficiency has been
shown to increase TfR1 expression in placental tissue, ensuring appropriate iron uptake from
maternal circulation to support fetal and placental iron requirements, we found no evidence of
maternal iron deficiency within our FGR group (Table 2). However, Burton and Fowden (2012)
and Sibley *et al.* (2010) report that when placental size and capacity are reduced, the placenta
increases nutrient transporter activity as a compensatory mechanism to suboptimal conditions,
rather than failing. This may explain our observed increases in iron transporters in significantly
smaller FGR placentae (Table 2). Collectively, this suggests that the increase in TfR1 in our
FGR placentae might represent a compensatory response to maximise iron uptake in a
compromised placental vascular environment.
602
Since iron transport is transcellular and unidirectional from mother to fetus, the placenta serves
as the "gatekeeper" for iron transfer to the fetus (O'Brien, 2022; Roberts *et al.*, 2020). The fetus
depends entirely on maternal iron transfer through the placenta, mediated by ferroportin (FPN),
which exports iron into the fetal circulation (Figure 1C-D) (Sangkhae *et al.*, 2020). We
observed 1.56-fold lower FPN (0.646 log2-fold) levels in FGR placental tissue, suggesting that
the placenta is retaining iron for its own use (O'Brien, 2022; Sangkhae *et al.*, 2020). This is
supported by an increase in divalent metal transporter 1 (DMT1/SLC11A2) mRNA expression,
the transporter of iron into the cell cytoplasm following endocytosis of the TfR1-iron complex
(Figure 1B). Subsequent Prussian blue staining showed no difference in iron deposit content
for ferric (Fe^{3+}) iron between FGR and healthy controls. Collectively, these findings suggest
that iron is being prioritised for intracellular function in FGR placentae, rather than for storage
or being exported into fetal circulation.
615

Mitochondrial Iron Transport

The mitochondria utilise between ~20-50% of cellular iron (Qi *et al.*, 2023), facilitating integral mitochondrial function and bioenergetic pathways to support fetoplacental growth (Aye *et al.*, 2022; Paul *et al.*, 2017). Mitoferrin-1 (SLC25A39) and mitoferrin-2 (SLC25A28) facilitate iron transport across the inner mitochondrial membrane but exhibit different cell expression profiles and cell-specific functions (Figure 2A-B) (Ali *et al.*, 2022). We observed no changes in mitoferrin-1 mRNA expression between our groups. In line with published literature in mice and zebrafish haematopoietic tissue (liver, bone marrow, spleen, and yolk sac islands) and embryonic tissue that determined mitoferrin-1 is essential for erythroid development to meet the higher demand for mitochondrial iron to support heme production (Paradkar *et al.*, 2009; Shaw *et al.*, 2006). As the placenta is a potent haematopoietic tissue containing erythroid progenitors from as early as 6 weeks of gestation through to term (Gekas *et al.*, 2005; Robin *et al.*, 2009). We propose that the expression of mitoferrin-1 is likely conserved, attributed to the placenta's fundamental role as a haematopoietic organ, rather than solely as a facilitator of maternal-fetal exchange. This is further demonstrated by Shaw *et al.* (2006) who observed that knockout (KO) of mitoferrin-1 in murine models results in embryonic lethality at the initiation of definitive erythrocyte development.

In contrast to mitoferrin-1, which is predominantly expressed in erythroid tissue, and minimal expressed in other tissues (Ali *et al.*, 2022), Shaw *et al.* (2006) and Paradkar *et al.* (2009) found that mitoferrin-2 is ubiquitously expressed in non-erythroid tissue and is responsible for the iron import required for haemoproteins and Fe-S cluster synthesis (Ali *et al.*, 2022). Our study observed a decrease in mitoferrin-2 mRNA expression in placentae from FGR pregnancies. These findings highlight the complex regulation of iron transport in FGR placentae, suggesting a selective downregulation of mitoferrin-2 while preserving mitoferrin-1 iron uptake for haematopoietic use.

Altered Iron-Sulfur Cluster Biosynthesis Machinery in FGR Placental Mitochondria

Given that our data suggested a conservation of intracellular iron and altered mitochondrial iron uptake within FGR placentae, we went on to investigate iron-sulfur (Fe-S) cluster biosynthesis. Fe-S clusters are apoproteins essential for cellular activities, including heme and are directly incorporated into the mitochondrial electron transport chain (ETC) to support mitochondrial structure and function (Guan *et al.*, 2022; Voltarelli *et al.*, 2023). Fe-S assembly

650   is a complex process that can be categorised into two main components: (i) de novo synthesis
651   of [2Fe-2S] cluster and (ii) late Fe-S ([4Fe-4S]) cluster synthesis (Figure 3). Disruptions in Fe-
652   S biogenesis impair cellular and mitochondrial function, and have been shown in literature to
653   lead to metabolic myopathies such as Friedreich's ataxia, sideroblastic anaemia, and ISCU
654   myopathy (Lee et $al.$, 2011).
655

Mitochondrial De Novo Fe-S Synthesis in FGR

De novo Fe-S cluster synthesis is a complex series of reactions, initiated by the catalysis of a

pyridoxal-phosphate, cysteine desulfurase NFS1 (Maio & Rouault, 2020). Stabilising the

activity of NFS1, NADH: Ubiquinone Oxidoreductase Subunit AB1 (NDUFAB1) forms a

heterodimer with LYR Motif Containing 4 (LYRM4), initiating Fe-S cluster synthesis via the

complex NFS1:LYRM4:NDUFBA1 (Figure 2C-D) (Herrera et $al.$, 2019). Our data showed a

1.09-fold lower abundance of NDUFAB1 (0.123 $\log_2$-fold) in FGR placentae, despite the

heterodimer conjugate LYRM4 mRNA expression remaining unchanged. The subsequent

synthesis of the NFS1:LYRM4:NDUFBA1 complex results in the binding of the iron-sulfur

cluster assembly enzyme (ISCU) and frataxin (FXN), forming a hetero-octamer that consists

of two copies of the NFS1:LYRM4:NDUFBA1:FXN:ISCU complex (Figure 2C-F and L)

(Maio & Rouault, 2020).
668

In our study, FXN mRNA expression remained unchanged between the two groups (Figure

2E), suggesting any deficits in Fe-S assembly likely occur downstream of initial FXN-mediated

iron and sulfide delivery onto ISCU, as loss of FXN results in widespread depletion of Fe-

containing proteins and attenuation of mitochondrial protein synthesis (Figure 2E) (Zhong et

$al.$, 2023). We observed a decrease in ISCU mRNA, which provides the structural scaffold for

[2Fe-2S] formation, in FGR placental mitochondria (Figure 2F). However, this observation

was not conserved in our proteomics data, with a 1.29-fold higher protein abundance in FGR

(-0.372 $\log_2$-FC; Figure 2L). In addition, key [2Fe-2S] proteins ferredoxin reductase (FDXR)

and ferredoxin 2 (FDX2), which are required to supply electrons to complete [2Fe-2S] and

[4Fe-4S] assembly, had 2.11-fold lower protein (1.08 $\log_2$-FC) and decreased gene expression

($p<0.05$) in FGR placentae, respectively. As we did not observe global suppression, we suggest

the FGR placenta prioritises [2Fe-2S] biogenesis at the detriment of [4Fe-4S] cluster synthesis

(Figure 2G and L). This is consistent with previous knockdown studies showing that loss of

FDX2 globally disrupts mitochondrial Fe-S assembly, leading to dysregulated oxidative

phosphorylation (OXPHOS) and reduced antioxidant production (Shi et $al.$, 2012).
684

Following [2Fe-2S] cluster formation, GLRX5 receives the [2Fe-2S] cluster from ISCU for

transport to apoproteins required for various cellular and mitochondrial functions (Paul *et al.*,

2017). We observed a significant decrease in mRNA expression of GLRX5, yet GLRX5 protein

abundance remained unchanged between the two groups (Figure 2H and L), again preserving

the capacity for [2Fe-2S] cluster transfer to downstream cellular and mitochondrial functions

such as heme synthesis or late-stage Fe-S cluster formation.

Mitochondrial Late-Stage Fe-S Synthesis in FGR

Given the changes observed in de novo synthesis, specifically reductions in protein abundance

of FDXR and mRNA expression of FDX2 and their sequential roles in electron transfer in the

formation of [2Fe-2S] and [4Fe-4S] clusters, we investigated late-stage Fe-S synthesis. In

contrast to de novo Fe-S cluster synthesis, late-stage [4Fe-4S] clusters are critical cofactors that

are incorporated into mitochondrial ETC complexes, and apoproteins such as iron-regulatory

proteins, and antioxidants (Maio & Rouault, 2020).

The formation of [4Fe-4S] clusters requires the transfer of [2Fe-2S] intermediaries to the A-

type scaffold proteins, Iron-Sulfur Cluster Assembly 1 and 2 (ISCA1 and ISCA2) in

conjugation with Iron-Sulfur Cluster Assembly Factor IBA57 (Melber *et al.*, 2016; Paul *et al.*,

2017). In FGR placentae, we observed a small 1.05-fold increase in IBA57 protein (-0.073

$\log_2$-FC; Figure 2L), which was confirmed by western blotting to be significant (p<0.002;

Figure 2M). However, we did not detect ISCA1 and ISCA2 at sufficient levels within our

proteomic analysis. Given that ISCA1 and ISCA2 are known to contain [2Fe-2S] clusters

(Beilschmidt *et al.*, 2017; Nasta *et al.*, 2019) but IBA57 does not contain [2Fe-2S] clusters

(Amick *et al.*, 2014; Dores-Silva *et al.*, 2015), the inability to detect sufficient levels of ISCA

proteins in FGR may suggest that the newly synthesised [2Fe-2S] clusters cannot be effectively

matured into [4Fe-4S] clusters.

Newly synthesised [4Fe-4S] clusters are released from the ISCA1:ISCA2:IBA57 complex by

a chaperone/co-chaperone system comprising heat shock protein family A (HSP 70) member 9

(HSPA9) and mitochondrial Fe-S Cluster co-chaperone (HSC20). HSPA9 protein abundance

was 1.16-fold lower in FGR placentae (0.219 $\log_2$-FC; Figure 2L). Following [4Fe-4S] cluster

release, HSPA9 mediates the transfer [4Fe-4S] clusters directly to apoproteins or through

targeting factors such as NFU1 Iron-Sulfur Cluster Scaffold (NFU1), BolA Family Member 3

(BOLA3) and NUBP Iron-Sulfur Cluster Assembly Factor (NUBPL) (Maio & Rouault, 2020;

Melber *et al.*, 2016; Zhong *et al.*, 2023). Incorporation of [4Fe-4S] into the mitochondrial ETC

complexes provides electron tunnelling through complexes I-IV, generating energy and a proton gradient crucial for ATP synthesis (Mostajabi Sarhangi & Matyushov, 2023; Vanlander & Van Coster, 2018). Despite increased mRNA expression of BOLA3 and NFU1 in FGR placentae (Figure 2I-J), we observed no changes in NUBPL mRNA expression (Figure 2K), which may suggest an attempt to maximise the delivery of a limited pool of [4Fe-4S] clusters.

Collectively, the reduced FDX2 mRNA expression and lower abundance of FDXR and HSPA9 suggest a bottleneck in mitochondrial Fe-S cluster biogenesis in the FGR placenta. While these proteins participate in both [2Fe-2S] and [4Fe-4S] cluster formation (Maio & Rouault, 2020), their decreased abundance in FGR likely prioritises the direction of limited resources toward essential [2Fe-2S] cluster formation, for heme synthesis (Shi *et al.*, 2012; Weiler *et al.*, 2020). This disruption in mitochondrial Fe-S assembly provides a molecular basis for the mitochondrial dysfunction observed in FGR placentae, characterised by impaired respiratory chain complex activity, altered ATP production, and increased oxidative stress (Holland *et al.*, 2017; Xu *et al.*, 2021).

Altered Heme Synthesis and Erythrocyte Structure in FGR Placentae: Implications for Fetal Oxygenation and Development

Based on our data showing prioritisation of [2Fe-2S] clusters in FGR placentae, we further investigated the heme synthesis pathway. Heme synthesis critically depends on the insertion of [2Fe-2S] clusters into ferrochelatase (FECH), the terminal enzyme of heme biosynthesis that resides within the mitochondria. The insertion of [2Fe-2S] clusters into FECH ensures its stability and activity for heme production (Al-Karadaghi *et al.*, 1997; Obi *et al.*, 2022). Our data revealed significantly higher levels of enzymes across the heme synthesis pathway in FGR placentae (Figure 3D). Specifically, elevated mRNA and protein levels of enzymes coproporphyrinogen oxidase (CPOX; 1.12-fold; -0.163 $\log_2$ FC) and FECH (1.28-fold; -0.356 $\log_2$-fold), located within the mitochondria, were accompanied by greater protein abundance of heme enzymes Hydroxymethylbilane Synthase (HMBS; 1.15-fold), and Uroporphyrinogen Decarboxylase (UROD: 1.13-fold). Supporting our suggestion that the FGR placentae prioritises [2Fe-2S] clusters and shifts utilisation towards heme synthesis. Additionally, FECH has been reported to be complexed to ATP Binding Cassette Subfamily B Member 10 (ABCB10), conferring stability to mitoferrin-1. This FECH-ABCB10 conjugation may explain why mitoferrin-1 expression remains conserved in FGR placentae despite reduced mitoferrin-2.

Beyond its role in oxygen transport and haemoglobin production, heme also contributes to mitochondrial respiration by facilitating electron transfer and proton translocation within ETC complexes II, III, and IV through its incorporation as heme a, b, and c moieties (Kim et al., 2012). Despite the increased levels of heme synthesis enzymes, the assessment of heme concentration in FGR placentae was lower (Figure 3C). This may be due to increased incorporation of heme a, b and c into ETC complexes II, III and IV to increase mitochondrial respiration resulting from a lack of [4Fe-4S] assembly, although this hypothesis requires confirmation.

Alternatively, heme is a known pro-oxidant, inducing oxidative stress, and heme accumulation can induce its catabolism via heme oxygenase 1 (HMOX1), degrading heme into carbon monoxide, iron and biliverdin (Figure 3D). We observed 2.19-fold higher HMOX1 abundance in FGR placentae, suggesting that despite upregulation of the heme biosynthesis pathway, increased heme breakdown may reflect heightened oxidative stress or a compensatory mechanism to regulate "free" heme levels. The catabolism of heme by the placenta is not well understood (Liu et al., 2016), although Peoc'h et al. (2020) have suggested that HMOX expression in the placenta could be a protective response against oxidative stress, which may explain why HMOX1 was higher in FGR placentae.

Our findings suggest a delicate balance in placental heme metabolism in FGR, whereby the upregulation of heme synthesis pathways likely represents an adaptive response to fetoplacental hypoxia and maintenance of placental bioenergetics. Simultaneously, the increase in HMOX1-mediated heme catabolism serves as a protective mechanism against heme-induced oxidative stress.

Haematopoietic function in FGR placentae

In addition to its role in mitochondrial function, heme is integral for the production of various globin chains, including gamma (γ), alpha (α) and beta (β) (Figure 4). As we observed an increase in heme-producing proteins, but no corresponding change in available heme in our FGR placental tissue, we examined the intricate relationship between heme and globin synthesis. This fundamental process is central to the oxygen-carrying capacity of blood and, by extension, aerobic respiration essential for cellular function and viability (Hamza & Dailey, 2012; Voltarelli et al., 2023). During embryonic development, primitive embryonic erythrocytes are located in blood islands of the secondary yolk sac (Manning *et al.*, 2020). Subsequently, a crucial transition occurs, coinciding with placental formation and the

emergence of definitive erythropoiesis (Palis & Segel, 1998). This developmental shift results
in the production of fetal haemoglobin ($\alpha2\gamma2$), which becomes the predominant form
throughout the remainder of fetal development (Hardison, 2012). This haemoglobin switching
process represents a critical milestone in fetal haematological development and is tightly
regulated by complex genetic and epigenetic factors (Hardison, 2012; Manning *et al.*, 2020).
Fetal haemoglobin production begins at around 10 weeks of gestation, peaking at 20 weeks and
remaining constant until the final month of fetal development when gamma-globin synthesis
rapidly decreases (Cantú & Philipsen, 2014; Mavilio *et al.*, 1983). Our findings revealed lower
protein abundance of gamma-1 (HBG1; 1.25-fold; 0.325 $\log_2$-FC) and gamma-2 (HBG2: 1.09-
fold; 0.09 $\log_2$-FC), key beta-like globin components of fetal haemoglobin, along with
haemoglobin subunit alpha 1 (HBA1; 1.16-fold; 0.21 $\log_2$-FC), in FGR placental tissue
compared to healthy term placental tissue (Figure 4). Additionally, the mRNA expression of
haemoglobin subunit beta (HBB; $p<0.05$), a critical component of adult haemoglobin ($\alpha_2\beta_2$),
was significantly increased in FGR placentae, however, protein abundance for HBB was 1.50-
fold lower (Figure 4D and G). The presence of HBB in the placenta aligns with current
literature stating that β-globin synthesis begins during the last month of fetal development
(Cantú & Philipsen, 2014). The switch from fetal to adult haemoglobin occurs when
erythropoiesis shifts from the fetal liver to the bone marrow at approximately (Khandros &
Blobel, 2024).
809
While fetal hypoxia in FGR is typically associated with elevated erythropoietin (EPO)
production and increased fetal haemoglobin synthesis (Chang *et al.*, 2018), our findings
indicate a disruption in globin chain production. The lower protein abundance of key globin
subunits, HBG1, HBG2, HBB and HBA1, in FGR placental tissue, suggests that globin chain
synthesis or stability may be impaired, potentially limiting the assembly of functional
haemoglobin. Although recent studies describe the placenta as a transient hematopoietic niche,
particularly during early to mid-gestation (Ivanovs *et al.*, 2017; Rhodes *et al.*, 2008), this
function may be compromised in FGR. Chronic hypoxia, inflammation, and oxidative stress,
all hallmarks of FGR pathophysiology, can disrupt haematopoietic stem and progenitor cell
(HSPC) function and erythroid differentiation (Dzierzak & Bigas, 2018). While hypoxia is
known to drive the upregulation of fetal haemoglobin via HIF1α activation (Khandros &
Blobel, 2024), we instead observed lower levels of subunits in both fetal and adult
haemoglobin. Together, suggesting a dysfunctional placental erythropoietic environment in
FGR.
824

Given the substantial alterations in both heme and globin subunits, we further investigated the erythropoietic structure in the FGR placenta. Our study revealed significant alterations in key structural proteins within FGR placentae. Notably, we observed reductions in erythrocyte membrane protein band 4.1 (EPB41) expression at both mRNA ($p < 0.05$) and protein levels (1.19-fold; 0.253 $\log_2$-FC) in FGR placentae (Figure 4E and G). The crucial role of EPB41 is in stabilising the actin-spectrin cytoskeleton of erythrocytes, thereby maintaining the characteristic biconcave disc shape of erythrocytes. The biconcave shape is fundamental to erythrocytes' function, optimising surface area to volume ratio and facilitating efficient oxygen diffusion and laminar flow in circulation (Uzoigwe, 2006; Vorn *et al.*, 2023). Huang et al (2017) showed that EPB41 null mice failed to develop a biconcave shape, resulting in compromised oxygen diffusion capacity. Furthermore, we observed reduced protein abundance in EPB42 (1.48-fold; 0.562 $\log_2$-FC), SPTA1 (1.12-fold; 0.153 $\log_2$-FC), SPTB (1.40-fold; 0.0491 $\log_2$-FC), ANK1 (1.57-fold; 0.661 $\log_2$-FC) and SLC4A1 (1.40-fold; 0.493 $\log_2$-FC) in FGR placentae when compared to healthy controls (Figure 4G). Similarly, these proteins are integral components of the erythrocyte membrane, conferring flexibility and stability (Alper, 2009; Guo *et al.*, 2022; Vorn *et al.*, 2023) Mutations in the aforementioned genes have been implicated in hereditary spherocytosis (Xu *et al.*, 2023), a condition characterised by erythrocyte membrane instability and deformation. Our findings suggest altered erythropoietic structure during erythrocyte maturation, as HEMGN mRNA expression remained unchanged, indicating that the HSPC population and early erythroid commitment remain preserved.

We propose that changes in the erythropoietic structure reflects a hypoxia-driven alteration in erythroid maturation, skewing erythropoiesis toward a rapid, stress-induced phenotype regulated in part by HIFs (Haase, 2013). Hypoxia-induced oxidative stress, as characteristic of FGR (Xu *et al.*, 2021), has been shown to affect the post-translational folding or stability of erythrocyte membrane components, which may further affect the abundance of EPB41 and EPB42 (Mohandas & Gallagher, 2008) and other erythrocyte membrane-bound proteins. Stress-induced erythropoiesis favours accelerated erythropoiesis, which may compromise erythrocyte cytoskeletal assembly and membrane protein integration (Paulson *et al.*, 2020). Collectively, these alterations may represent an attempt to increase erythrocyte deformability to facilitate circulation within the structurally compromised vasculature of the FGR placenta, characterised by narrower blood vessels and increased vascular resistance (Boss *et al.*, 2023; Kovo *et al.*, 2013; Sun *et al.*, 2020). Thus, enabling efficient uptake and transport of the limited oxygen available to the fetus.

Despite the structural and functional changes observed in erythrocytes of FGR placental tissue, the mRNA expression of hemogen (HEMGN) remained unchanged. Hemogen regulates the proliferation and differentiation of haematopoietic cells (Jiang *et al.*, 2023) indicating that the architectural insult observed in FGR placentae is likely not occurring during haematopoiesis but rather during erythrocyte maturation (Vorn *et al.*, 2023) or as a result of stress-induced erythrocyte maturation (Paulson *et al.*, 2020).

<u>Limitations</u>

While previous research indicates that maternal iron status is a critical determinant of fetal iron acquisition through placental transport (Cao & Fleming, 2016; Fisher & Nemeth, 2017; Roberts *et al.*, 2020), our study has provided new insights into intrinsic placental changes that are independent of maternal iron status, revealing potential intracellular alterations that have remained elusive in the development of FGR. Since, the placenta is the critical interface between maternal and fetal circulations, building upon the biomolecular foundation established in this study incorporating fetal cord blood analysis in future studies would provide further insight into how modifications in placental iron transport and utilisation impact fetal iron status. In this study, maternal patient records were retrieved from mothers who delivered either healthy or FGR infants, offering valuable insights into maternal iron status during pregnancy. The clinical records revealed comparable iron parameters between groups, strengthening our conclusion that the placental iron alterations we observed represent intrinsic placental adaptations rather than responses to maternal iron availability. Although the retrospective nature of clinical records means that iron parameters were measured based on clinical need rather than a standardised gestational timepoint. While this approach provides real-time data collected during routine clinical care, data uniformity across participants may be potentially limiting (Talari & Goyal, 2020).

This study employed a bottom-up liquid chromatography mass spectrometry (LC-MS) proteomic approach to successfully identify significant alterations in placental iron transport and mitochondrial iron utilisation. While the LC-MS methodology chosen is well-suited to global profiling performed within this study to assess total cellular protein content, some proteins were difficult to detect due to protein inference ambiguity, underrepresentation of low-abundance, membrane-bound proteins, and incomplete coverage of post-translational modifications (Bruderer *et al.*, 2015; Mulhall *et al.*). Notably, key Fe-S cluster proteins essential for de novo and late-stage synthesis were difficult to detect, likely due to their hydrophobic nature and the inherent limitations of the protocol (Mulhall *et al.*). Future studies

incorporating targeted approaches may improve the detection and quantification of such

proteins.

Our study primarily focused on non-heme iron pathways within the placenta, which represents

85% of iron obtained from the maternal diet through plant-based sources of iron (Piskin *et al.*,

2022; Yiannikourides & Latunde-Dada, 2019). Heme-iron, accounting for 15% of dietary iron

from animal-based sources, requires further investigation to understand heme-iron absorption

during pregnancy. Current literature suggests that heme-iron absorption follows the same

pattern as non-heme iron absorption in the placenta and is mainly sourced via maternal

haemolysis rather than absorbed dietary iron (Milman, 2006; Roberts *et al.*, 2020).

<u>Conclusion</u>

Our study reveals specific alterations in placental iron transport and utilisation pathways in

FGR, that are independent of maternal iron status. We identified increased placental iron

transporters alongside reduced ferroportin, suggesting an adaptive iron retention within FGR

placentae. Assessment of mitochondrial iron utilisation pathways in FGR placentae suggested

a preference for [2Fe-2S] cluster formation and heme synthesis over [4Fe-4S] cluster assembly

in FGR. Collectively, these findings of altered placental iron transporters and modifications in

mitochondrial Fe-S cluster and heme synthesis pathways likely reflect the placenta's response

to the compromised vascular environment characteristic of FGR. We also observed reduced

haemoglobin subunits and changes in erythrocyte structural proteins in FGR placentae,

indicative of altered placental hematopoietic function. We propose these data are a programmed

response to hypoxia and stress-induced haematopoiesis to overcome the poor placental

vasculature in FGR. Together, these findings provide new insights into how alterations in iron

transport and iron-dependent pathways contribute to the development of FGR.

Data Availability Statement

The authors will make the raw data supporting the conclusions of this article available, without

undue reservation, to any qualified researcher.

Funding

This research was supported by the National Health and Medical Research Council Investigator

Grant GNT2026065.

Author Contributions

VBB, KGP, RS and JJF contributed to the conception and design of the study. VBB, HM and SA contributed to the acquisition, analysis and interpretation of data for the study. VBB, KGP, RS, and JJF contributed to drafting the work and critically revising the intellectual content. All authors have approved the final version of the manuscript. All authors agree to be accountable for all aspects of the work. All persons designated as authors qualify for authorship and are listed.

Competing Interests

The authors declare that they have no conflict of interest in this study.

Acknowledgements

We gratefully acknowledge the research midwives at John Hunter Hospital, Skye Doel and Bridget McCleery, who were involved in participant recruitment and clinical data collection, for their invaluable support and contribution to this study.

References

Abbassi-Ghanavati M, Greer LG & Cunningham FG. (2009). Pregnancy and laboratory studies: a reference table for clinicians. *Obstet Gynecol* **114,** 1326-1331.

Al-Karadaghi S, Hansson M, Nikonov S, Jönsson B & Hederstedt L. (1997). Crystal structure of ferrochelatase: the terminal enzyme in heme biosynthesis. *Structure* **5,** 1501-1510.

Ali MY, Oliva CR, Flor S & Griguer CE. (2022). Mitoferrin, Cellular and Mitochondrial Iron Homeostasis. *Cells* **11**.

Allerkamp HH, Pole T, Boukham A, James JL & Clark AR. (2022). Pregnancy-specific uterine vascular reactivity: a data-driven computational model of shear-dependent, myogenic, and mechanical radial artery features. *American Journal of Physiology-Heart and Circulatory Physiology* **323,** H72-H88.

Alper SL. (2009). Molecular physiology and genetics of Na+-independent SLC4 anion exchangers. *J Exp Biol* **212,** 1672-1683.

Amick J, Schlanger SE, Wachnowsky C, Moseng MA, Emerson CC, Dare M, Luo WI,
Ithychanda SS, Nix JC, Cowan JA, Page RC & Misra S. (2014). Crystal structure of
the nucleotide-binding domain of mortalin, the mitochondrial Hsp70 chaperone.
Protein Sci **23,** 833-842.

Anderson GJ & Frazer DM. (2017). Current understanding of iron homeostasis. *Am J Clin*
Nutr **106,** 1559s-1566s.

Andrews NC. (2000). Iron metabolism: iron deficiency and iron overload. *Annu Rev*
Genomics Hum Genet **1,** 75-98.

Aye I, Aiken CE, Charnock-Jones DS & Smith GCS. (2022). Placental energy metabolism
in health and disease-significance of development and implications for
preeclampsia. *Am J Obstet Gynecol* **226,** S928-s944.

Beilschmidt LK, Ollagnier de Choudens S, Fournier M, Sanakis I, Hograindleur MA,
Clémancey M, Blondin G, Schmucker S, Eisenmann A, Weiss A, Koebel P,
Messaddeq N, Puccio H & Martelli A. (2017). ISCA1 is essential for mitochondrial
Fe(4)S(4) biogenesis in vivo. *Nat Commun* **8,** 15124.

Boss AL, Chamley LW, Brooks AES & James JL. (2023). Human placental vascular and
perivascular cell heterogeneity differs between first trimester and term, and in
pregnancies affected by foetal growth restriction. *Mol Hum Reprod* **29**.

Brighton E. (2019). New South Wales Hopsital Laboratory clinical projects laboratory
manaual ed. 2019 NRRA. NSW Health Pathology.

Brooker I, Fisher J, Delforce S, Endacott S, Lumbers E, Sutherland J & Pringle K. (2025).
Characterising the role of placental angiotensin-converting enzyme 2 (ACE2)
during the onset of oxidative insult by hypoxia/reoxygenation: Implications for
fetal growth restriction. *Placenta* **165**.

Bruderer R, Bernhardt OM, Gandhi T, Miladinović SM, Cheng L-Y, Messner S,
Ehrenberger T, Zanotelli V, Butscheid Y, Escher C, Vitek O, Rinner O & Reiter L.
(2015). Extending the Limits of Quantitative Proteome Profiling with Data-
Independent Acquisition and Application to Acetaminophen-Treated Three-
Dimensional Liver Microtissues*[S]. *Molecular & Cellular Proteomics* **14,** 1400-
1410.

1005

1006 Burton G & Fowden A. (2012). The placenta and developmental programming: balancing
1007   fetal nutrient demands with maternal resource allocation. *Placenta* **33,** S23-S27.

1008

1009   Burton GJ & Jauniaux E. (2018). Pathophysiology of placental-derived fetal
1010   growth restriction. *Am J Obstet Gynecol* **218,** S745-s761.

1011
1012   Camaschella C. (2015). Iron-deficiency anemia. *N Engl J Med* **372,** 1832-1843.

1013
1014   Cantú I & Philipsen S. (2014). Flicking the switch: Adult hemoglobin expression in
1015   erythroid cells derived from cord blood and human induced pluripotent stem
1016   cells. *Haematologica* **99,** 1647-1649.

1017
1018   Cao C & Fleming MD. (2016). The placenta: the forgotten essential organ of iron
1019   transport. *Nutr Rev* **74,** 421-431.

1020
1021   Cerami C. (2017). Iron Nutriture of the Fetus, Neonate, Infant, and Child. *Ann Nutr*
1022   *Metab* **71 Suppl 3,** 8-14.

1023
1024   Chang Y-L, Chao A-S, Peng H-H, Chang S-D, Chen K-J, Cheng P-J & Wang T-H. (2018).
1025   Placental erythropoietin expression is upregulated in growth-restricted fetuses
1026   with abnormal umbilical artery Doppler findings: a case–control study of
1027   monochorionic twins. *BMC Pregnancy and Childbirth* **18,** 1-7.

1028
1029   Chiabrando D, Mercurio S & Tolosano E. (2014). Heme and erythropoieis: more than a
1030   structural role. *Haematologica* **99,** 973-983.

1031
1032   Cindrova-Davies T & Sferruzzi-Perri AN. (2022). Human placental development and
1033   function. *Semin Cell Dev Biol* **131,** 66-77.

1034
1035   de Freitas MAR, da Costa AV, Medeiros LA, Cunha LM, Coutinho Filho U, Garrote Filho
1036   MdS, Diniz ALD & Penha-Silva N. (2019). The role of the erythrocyte in the
1037   outcome of pregnancy with preeclampsia. *PLOS ONE* **14,** e0212763.

1038
1039   Delaney KM, Guillet R, Pressman EK, Ganz T, Nemeth E & O'Brien KO. (2021). Serum
1040   Erythroferrone During Pregnancy Is Related to Erythropoietin but Does Not
1041   Predict the Risk of Anemia. *The Journal of Nutrition* **151,** 1824-1833.

1042
1043   Dev S & Babitt JL. (2017). Overview of iron metabolism in health and disease. *Hemodial*
1044   *Int* **21 Suppl 1,** S6-s20.

1045
1046   Dores-Silva PR, Barbosa LR, Ramos CH & Borges JC. (2015). Human mitochondrial
1047   Hsp70 (mortalin): shedding light on ATPase activity, interaction with adenosine
1048   nucleotides, solution structure and domain organization. *PLoS One* **10,**
1049   e0117170.

1050

1051 Dzierzak E & Bigas A. (2018). Blood development: hematopoietic stem cell dependence
1052 and independence. *Cell stem cell* **22,** 639-651.

1053

1054 Fairchild Benyo D & Conrad KP. (1999). Expression of the erythropoietin receptor by
1055 trophoblast cells in the human placenta. *Biology of reproduction* **60,** 861-870.

1056

1057 Fava C, Cattazzo F, Hu ZD, Lippi G & Montagnana M. (2019). The role of red blood cell
1058 distribution width (RDW) in cardiovascular risk assessment: useful or hype? *Ann*
1059 *Transl Med* **7,** 581.

1060

1061 Fisher AL & Nemeth E. (2017). Iron homeostasis during pregnancy. *Am J Clin Nutr* **106,**
1062 1567s-1574s.

1063

1064 Fisher JJ, Grace T, Castles NA, Jones EA, Delforce SJ, Peters AE, Crombie GK, Hoedt EC,
1065 Warren KE, Kahl RG, Hirst JJ, Pringle KG & Pennell CE. (2024). Methodology for
1066 Biological Sample Collection, Processing, and Storage in the Newcastle 1000
1067 Pregnancy Cohort: Protocol for a Longitudinal, Prospective Population-Based
1068 Study in Australia. *JMIR Res Protoc* **13,** e63562.

1069

1070 Fisher JJ, McKeating DR, Cuffe JS, Bianco-Miotto T, Holland OJ & Perkins AV. (2019).
1071 Proteomic analysis of placental mitochondria following trophoblast
1072 differentiation. *Frontiers in physiology* **10,** 1536.

1073

1074 Gambling L, Danzeisen R, Gair S, Lea RG, Charania Z, Solanky N, Joory KD, Srai SK &
1075 McArdle HJ. (2001). Effect of iron deficiency on placental transfer of iron and
1076 expression of iron transport proteins in vivo and in vitro. *Biochem J* **356,** 883-889.

1077

1078 Gekas C, Dieterlen-Lièvre F, Orkin SH & Mikkola HK. (2005). The placenta is a niche for
1079 hematopoietic stem cells. *Dev Cell* **8,** 365-375.

1080

1081 Gekas C, Rhodes KE, Van Handel B, Chhabra A, Ueno M & Mikkola HK. (2010).
1082 Hematopoietic stem cell development in the placenta. *Int J Dev Biol* **54,** 1089-
1083 1098.

1084

1085 Gordijn SJ, Beune IM, Thilaganathan B, Papageorghiou A, Baschat AA, Baker PN, Silver
1086 RM, Wynia K & Ganzevoort W. (2016). Consensus definition of fetal growth
1087 restriction: a Delphi procedure. *Ultrasound in Obstetrics & Gynecology* **48,** 333-
1088 339.

1089

1090 Guan S, Zhao L & Peng R. (2022). Mitochondrial Respiratory Chain Supercomplexes:
1091 From Structure to Function. *Int J Mol Sci* **23**.

Guo S, Frazer DM & Anderson GJ. (2016). Iron homeostasis: transport, metabolism, and regulation. *Curr Opin Clin Nutr Metab Care* **19,** 276-281.

Guo X, Zhao Y, Kim J & Dean A. (2022). Hemogen/BRG1 cooperativity modulates promoter and enhancer activation during erythropoiesis. *Blood* **139,** 3532-3545.

Guo Y, Zhang N, Zhang D, Ren Q, Ganz T, Liu S & Nemeth E. (2019). Iron homeostasis in pregnancy and spontaneous abortion. *American journal of hematology* **94,** 184-188.

Haase VH. (2013). Regulation of erythropoiesis by hypoxia-inducible factors. *Blood Reviews* **27,** 41-53.

Hardison RC. (2012). Evolution of hemoglobin and its genes. *Cold Spring Harbor perspectives in medicine* **2,** a011627.

Hentze MW, Muckenthaler MU & Andrews NC. (2004). Balancing Acts: Molecular Control of Mammalian Iron Metabolism. *Cell* **117,** 285-297.

Herrera MG, Noguera ME, Sewell KE, Agudelo Suárez WA, Capece L, Klinke S & Santos J. (2019). Structure of the Human ACP-ISD11 Heterodimer. *Biochemistry* **58,** 4596-4609.

Higashijima A, Miura K, Mishima H, Kinoshita A, Jo O, Abe S, Hasegawa Y, Miura S, Yamasaki K, Yoshida A, Yoshiura K-i & Masuzaki H. (2013). Characterization of placenta-specific microRNAs in fetal growth restriction pregnancy. *Prenatal Diagnosis* **33,** 214-222.

Holland O, Dekker Nitert M, Gallo LA, Vejzovic M, Fisher JJ & Perkins AV. (2017). Review: Placental mitochondrial function and structure in gestational disorders. *Placenta* **54,** 2-9.

Hou J, Cliver SP, Tamura T, Johnston KE & Goldenberg R. (2000). Maternal serum ferritin and fetal growth. *Obstet Gynecol* **95,** 447-452.

Ivanovs A, Rybtsov S, Ng ES, Stanley EG, Elefanty AG & Medvinsky A. (2017). Human haematopoietic stem cell development: from the embryo to the dish. *Development* **144,** 2323-2337.

1132   Jiang Y, Yu L, Hu Q, Kang Y, You J, Huang C, Xu X & Chen L. (2023). Cancer/testis antigen
1133   HEMGN correlated with immune infiltration serves as a prognostic biomarker in
1134   lung adenocarcinoma. *Mol Immunol* **153,** 226-237.

1135

1136   Kautz L & Nemeth E. (2014). Molecular liaisons between erythropoiesis and iron
1137   metabolism. *Blood* **124,** 479-482.

1138

1139   Khandros E & Blobel GA. (2024). Elevating fetal hemoglobin–recently discovered
1140   regulators and mechanisms. *Blood*.

1141

1142   Kim HJ, Khalimonchuk O, Smith PM & Winge DR. (2012). Structure, function, and
1143   assembly of heme centers in mitochondrial respiratory complexes. *Biochimica*
1144   *et Biophysica Acta (BBA)-Molecular Cell Research* **1823,** 1604-1616.

1145

1146   Kovo M, Schreiber L, Ben-Haroush A, Cohen G, Weiner E, Golan A & Bar J. (2013). The
1147   placental factor in early- and late-onset normotensive fetal growth restriction.
1148   *Placenta* **34,** 320-324.

1149

1150   Kurt RK, Aras Z, Silfeler DB, Kunt C, Islimye M & Kosar O. (2015). Relationship of Red Cell
1151   Distribution Width With the Presence and Severity of Preeclampsia. *Clinical and*
1152   *Applied Thrombosis/Hemostasis* **21,** 128-131.

1153

Lee D-C, Romero R, Kim J-S, Tarca AL, Montenegro D, Pineles BL, Kim E, Lee J, Kim SY,
Draghici S, Mittal P, Kusanovic JP, Chaiworapongsa T, Hassan SS & Kim CJ. (2011).
miR-210 Targets Iron-Sulfur Cluster Scaffold Homologue in Human Trophoblast
Cell Lines: Siderosis of Interstitial Trophoblasts as a Novel Pathology of Preterm
Preeclampsia and Small-for-Gestational-Age Pregnancies. *The American Journal*
of Pathology **179,** 590-602.

1160

1161   Lippi G & Plebani M. (2014). Red blood cell distribution width (RDW) and human
1162   pathology. One size fits all. *Clinical Chemistry and Laboratory Medicine (CCLM)*
1163   **52,** 1247-1249.

1164

1165   Maio N & Rouault TA. (2020). Outlining the Complex Pathway of Mammalian Fe-S
1166   Cluster Biogenesis. *Trends Biochem Sci* **45,** 411-426.

1167

1168   Maio N & Rouault TA. (2020). Outlining the Complex Pathway of Mammalian Fe-S
1169   Cluster Biogenesis. *Trends in Biochemical Sciences* **45,** 411-426.

1170

1171   Manning JM, Manning LR, Dumoulin A, Padovan JC & Chait B. (2020). Embryonic and
1172   Fetal Human Hemoglobins: Structures, Oxygen Binding, and Physiological Roles.
1173   In *Vertebrate and Invertebrate Respiratory Proteins, Lipoproteins and other Body*

1174 *Fluid Proteins*, ed. Hoeger U & Harris JR, pp. 275-296. Springer International
1175   Publishing, Cham.

1176
1177 Mastrogiannaki M, Matak P, Keith B, Simon MC, Vaulont S & Peyssonnaux C. (2009). HIF-
1178   2alpha, but not HIF-1alpha, promotes iron absorption in mice. *J Clin Invest* **119,**
1179   1159-1166.

1180
1181 Mavilio F, Giampaolo A, Carè A, Migliaccio G, Calandrini M, Russo G, Pagliardi GL,
1182   Mastroberardino G, Marinucci M & Peschle C. (1983). Molecular mechanisms of
1183   human hemoglobin switching: selective undermethylation and expression of
1184   globin genes in embryonic, fetal, and adult erythroblasts. *Proc Natl Acad Sci U S*
1185   *A* **80,** 6907-6911.

1186
1187 McArdle HJ, Danzeisen R, Fosset C & Gambling L. (2003). The role of the placenta in iron
1188   transfer from mother to fetus and the relationship between iron status and fetal
1189   outcome. *Biometals* **16,** 161-167.

1190
1191 McArdle HJ, Lang C, Hayes H & Gambling L. (2011). Role of the placenta in regulation of
1192   fetal iron status. *Nutr Rev* **69 Suppl 1,** S17-22.

1193
1194 Mecacci F, Avagliano L, Lisi F, Clemenza S, Serena C, Vannuccini S, Rambaldi MP,
1195   Simeone S, Ottanelli S & Petraglia F. (2021). Fetal Growth Restriction: Does an
1196   Integrated Maternal Hemodynamic-Placental Model Fit Better? *Reprod Sci* **28,**
1197   2422-2435.

1198
1199 Mégier C, Peoc'h K, Puy V & Cordier AG. (2022). Iron Metabolism in Normal and
1200   Pathological Pregnancies and Fetal Consequences. *Metabolites* **12**.

1201
1202 Melber A, Na U, Vashisht A, Weiler BD, Lill R, Wohlschlegel JA & Winge DR. (2016). Role
1203   of Nfu1 and Bol3 in iron-sulfur cluster transfer to mitochondrial clients. *Elife* **5,**
1204   e15991.

1205
1206 Miller J, Turan S & Baschat AA. (2008). Fetal Growth Restriction. *Seminars in*
1207   *Perinatology* **32,** 274-280.

1208
1209 Milman N. (2006). Iron and pregnancy--a delicate balance. *Ann Hematol* **85,** 559-565.

1210
1211 Mohandas N & Gallagher PG. (2008). Red cell membrane: past, present, and future.
1212   *Blood* **112,** 3939-3948.

1213

Moksnes MR, Graham SE, Wu K-H, Hansen AF, Gagliano Taliun SA, Zhou W, Thorstensen K, Fritsche LG, Gill D, Mason A, Cucca F, Schlessinger D, Abecasis GR, Burgess S, Åsvold BO, Nielsen JB, Hveem K, Willer CJ & Brumpton BM. (2022). Genome-wide meta-analysis of iron status biomarkers and the effect of iron on all-cause mortality in HUNT. *Communications Biology* **5,** 591.

Mostajabi Sarhangi S & Matyushov DV. (2023). Electron Tunneling in Biology: When Does it Matter? *ACS Omega* **8,** 27355-27365.

Mulhall JE, Trigg NA, Bernstein IR, Anderson AL, Murray HC, Sipilä P, Lord T, Schjenken JE, Nixon B & Skerrett-Byrne DA. (2024). Immortalized mouse caput epididymal epithelial (mECap18) cell line recapitulates the in-vivo environment. *PROTEOMICS* **24,** 2300253.

Murray HC, Miller K, Brzozowski JS, Kahl RGS, Smith ND, Humphrey SJ, Dun MD & Verrills NM. (2023). Synergistic Targeting of DNA-PK and KIT Signaling Pathways in KIT Mutant Acute Myeloid Leukemia. *Molecular & Cellular Proteomics* **22,** 100503.

Nader E, Skinner S, Romana M, Fort R, Lemonne N, Guillot N, Gauthier A, Antoine-Jonville S, Renoux C, Hardy-Dessources M-D, Stauffer E, Joly P, Bertrand Y & Connes P. (2019). Blood Rheology: Key Parameters, Impact on Blood Flow, Role in Sickle Cell Disease and Effects of Exercise. *Frontiers in Physiology* **Volume 10 - 2019**.

Nasta V, Da Vela S, Gourdoupis S, Ciofi-Baffoni S, Svergun DI & Banci L. (2019). Structural properties of [2Fe-2S] ISCA2-IBA57: a complex of the mitochondrial iron-sulfur cluster assembly machinery. *Scientific Reports* **9,** 18986.

O'Brien KO. (2022). Maternal, fetal and placental regulation of placental iron trafficking. *Placenta* **125,** 47-53.

Obi CD, Bhuiyan T, Dailey HA & Medlock AE. (2022). Ferrochelatase: Mapping the Intersection of Iron and Porphyrin Metabolism in the Mitochondria. *Frontiers in Cell and Developmental Biology* **Volume 10 - 2022**.

Ozgu-Erdinc AS, Cavkaytar S, Aktulay A, Buyukkagnici U, Erkaya S & Danisman N. (2014). Mid-trimester maternal serum and amniotic fluid biomarkers for the prediction of preterm delivery and intrauterine growth retardation. *J Obstet Gynaecol Res* **40,** 1540-1546.

Palis J & Segel GB. (1998). Developmental biology of erythropoiesis. *Blood Reviews* **12,** 106-114.

Paradkar PN, Zumbrennen KB, Paw BH, Ward DM & Kaplan J. (2009). Regulation of mitochondrial iron import through differential turnover of mitoferrin 1 and mitoferrin 2. *Molecular and cellular biology* **29,** 1007-1016.

Paul BT, Manz DH, Torti FM & Torti SV. (2017). Mitochondria and Iron: current questions. *Expert Rev Hematol* **10,** 65-79.

Paulson RF, Hariharan S & Little JA. (2020). Stress erythropoiesis: definitions and models for its study. *Experimental hematology* **89,** 43-54. e42.

Perry H, Lehmann H, Mantovani E, Thilaganathan B & Khalil A. (2020). Are maternal hemodynamic indices markers of fetal growth restriction in pregnancies with a small-for-gestational-age fetus? *Ultrasound in Obstetrics & Gynecology* **55,** 210-216.

Piskin E, Cianciosi D, Gulec S, Tomas M & Capanoglu E. (2022). Iron Absorption: Factors, Limitations, and Improvement Methods. *ACS Omega* **7,** 20441-20456.

Qi Y, Hu M, Wang Z & Shang W. (2023). Mitochondrial iron regulation as an emerging target in ischemia/reperfusion injury during kidney transplantation. *Biochemical Pharmacology* **215,** 115725.

Read AD, Bentley RE, Archer SL & Dunham-Snary KJ. (2021). Mitochondrial iron-sulfur clusters: Structure, function, and an emerging role in vascular biology. *Redox Biol* **47,** 102164.

Reddy SG, Kumar H, Nagaraj G & Kottur PP. (2016). Red cell distribution width as a marker of preeclampsia severity. *Journal of Evolution of Medical and Dental Sciences* **5,** 3104-3108.

Rhodes KE, Gekas C, Wang Y, Lux CT, Francis CS, Chan DN, Conway S, Orkin SH, Yoder MC & Mikkola HK. (2008). The emergence of hematopoietic stem cells is initiated in the placental vasculature in the absence of circulation. *Cell stem cell* **2,** 252-263.

Roberts H, Bourque SL & Renaud SJ. (2020). Maternal iron homeostasis: effect on placental development and function. *Reproduction* **160,** R65-r78.

1296 Robin C, Bollerot K, Mendes S, Haak E, Crisan M, Cerisoli F, Lauw I, Kaimakis P, Jorna R,
1297   Vermeulen M, Kayser M, van der Linden R, Imanirad P, Verstegen M, Nawaz-
1298   Yousaf H, Papazian N, Steegers E, Cupedo T & Dzierzak E. (2009). Human
1299   placenta is a potent hematopoietic niche containing hematopoietic stem and
1300   progenitor cells throughout development. *Cell Stem Cell* **5,** 385-395.

1301

1302 Sanghavi M & Rutherford JD. (2014). Cardiovascular Physiology of Pregnancy.
1303   *Circulation* **130,** 1003-1008.

1304

1305 Sangkhae V, Fisher AL, Wong S, Koenig MD, Tussing-Humphreys L, Chu A, Lelić M, Ganz
1306   T & Nemeth E. (2020). Effects of maternal iron status on placental and fetal iron
1307   homeostasis. *J Clin Invest* **130,** 625-640.

1308

1309 Semenza GL. (2014). Oxygen sensing, hypoxia-inducible factors, and disease
1310   pathophysiology. *Annu Rev Pathol* **9,** 47-71.

1311

1312 Shaw GC, Cope JJ, Li L, Corson K, Hersey C, Ackermann GE, Gwynn B, Lambert AJ,
1313   Wingert RA, Traver D, Trede NS, Barut BA, Zhou Y, Minet E, Donovan A, Brownlie
1314   A, Balzan R, Weiss MJ, Peters LL, Kaplan J, Zon LI & Paw BH. (2006). Mitoferrin is
1315   essential for erythroid iron assimilation. *Nature* **440,** 96-100.

1316

1317 Shehata HA, Ali MM, Evans-Jones JC, Upton GJG & Manyonda IT. (1998). Red cell
1318   distribution width (RDW) changes in pregnancy. *International Journal of
1319   Gynecology & Obstetrics* **62,** 43-46.

1320

1321 Shi Y, Ghosh M, Kovtunovych G, Crooks DR & Rouault TA. (2012). Both human
1322   ferredoxins 1 and 2 and ferredoxin reductase are important for iron-sulfur cluster
1323   biogenesis. *Biochimica et Biophysica Acta (BBA) - Molecular Cell Research* **1823,**
1324   484-492.

1325

1326 Sibley CP, Brownbill P, Dilworth M & Glazier JD. (2010). Adaptation in placental nutrient
1327   supply to meet fetal growth demand: implications for programming. *Placenta* **31,**
1328   S70-S74.

1329

1330 Söhnchen N, Melzer K, Tejada BMd, Jastrow-Meyer N, Othenin-Girard V, Irion O,
1331   Boulvain M & Kayser B. (2011). Maternal heart rate changes during labour.
1332   *European Journal of Obstetrics & Gynecology and Reproductive Biology* **158,**
1333   173-178.

1334

1335 Sun C, Groom KM, Oyston C, Chamley LW, Clark AR & James JL. (2020). The placenta in
1336   fetal growth restriction: What is going wrong? *Placenta* **96,** 10-18.

1337

1338 Talari K & Goyal M. (2020). Retrospective studies - utility and caveats. *J R Coll Physicians*
1339   *Edinb* **50,** 398-402.

1340

1341 Tang L, He G, Liu X & Xu W. (2017). Progress in the understanding of the etiology and
1342   predictability of fetal growth restriction. *Reproduction* **153,** R227-R240.

1343

1344 Uzoigwe C. (2006). The human erythrocyte has developed the biconcave disc shape to
1345   optimise the flow properties of the blood in the large vessels. *Med Hypotheses*
1346   **67,** 1159-1163.

1347

1348 Vanlander AV & Van Coster R. (2018). Clinical and genetic aspects of defects in the
1349   mitochondrial iron–sulfur cluster synthesis pathway. *JBIC Journal of Biological*
1350   *Inorganic Chemistry* **23,** 495-506.

1351

1352 Voltarelli VA, Alves de Souza RW, Miyauchi K, Hauser CJ & Otterbein LE. (2023). Heme:
1353   The Lord of the Iron Ring. *Antioxidants* **12,** 1074.

1354

Vorn R, Devoto C, Meier TB, Lai C, Yun S, Broglio SP, Mithani S, McAllister TW, Giza CC,
Kim H-S, Huber D, Harezlak J, Cameron KL, McGinty G, Jackson J, Guskiewicz
KM, Mihalik JP, Brooks A, Duma S, Rowson S, Nelson LD, Pasquina P, McCrea MA
& Gill JM. (2023). Are EPB41 and alpha-synuclein diagnostic biomarkers of sport-
related concussion? Findings from the NCAA and Department of Defense CARE
Consortium. *Journal of Sport and Health Science* **12,** 379-387.

1361

1362 Waldvogel-Abramowski S, Waeber G, Gassner C, Buser A, Frey BM, Favrat B & Tissot JD.
1363   (2014). Physiology of iron metabolism. *Transfus Med Hemother* **41,** 213-221.

1364

1365 Weiler BD, Brück M-C, Kothe I, Bill E, Lill R & Mühlenhoff U. (2020). Mitochondrial [4Fe-
1366   4S] protein assembly involves reductive [2Fe-2S] cluster fusion on ISCA1–ISCA2
1367   by electron flow from ferredoxin FDX2. *Proceedings of the National Academy of*
1368   *Sciences* **117,** 20555-20565.

1369

1370 Wen Z, Luo D, Wang S, Rong R, Evers BM, Jia L, Fang Y, Daoud EV, Yang S, Gu Z, Arner EN,
1371   Lewis CM, Solis Soto LM, Fujimoto J, Behrens C, Wistuba, II, Yang DM, Brekken
1372   RA, O'Donnell KA, Xie Y & Xiao G. (2024). Deep Learning-Based H-Score
1373   Quantification of Immunohistochemistry-Stained Images. *Mod Pathol* **37,**
1374   100398.

1375

1376 Wu BA, Chand KK, Bell A, Miller SL, Colditz PB, Malhotra A & Wixey JA. (2023). Effects of
1377   fetal growth restriction on the perinatal neurovascular unit and possible
1378   treatment targets. *Pediatric Research*.

1379

1380 Xu L, Wei X, Liang G, Zhu D, Zhang Y, Zhang Y & Shang X. (2023). A novel splicing
1381   mutation of ANK1 is associated with phenotypic heterogeneity of hereditary
1382   spherocytosis in a Chinese family. *Biochimica et Biophysica Acta (BBA) -*
1383   *Molecular Basis of Disease* **1869,** 166595.
1384
1385 Xu Y-Y, Liu Y, Cui L, Wu W-B, Quinn MJ, Menon R & Zhang H-J. (2021). Hypoxic effects on
1386   the mitochondrial content and functions of the placenta in fetal growth
1387   restriction. *Placenta* **114,** 100-107.
1388
1389 Yiannikourides A & Latunde-Dada GO. (2019). A Short Review of Iron Metabolism and
1390   Pathophysiology of Iron Disorders. *Medicines (Basel)* **6**.
1391
1392 Zaugg J, Lopez-Tello J, Musial B, Vaughan OR, Fowden AL, Albrecht C & Sferruzzi-Perri
1393   AN. (2024). Obesogenic diet in pregnancy disrupts placental iron handling and
1394   ferroptosis and stress signalling in association with fetal growth alterations.
1395   *Cellular and Molecular Life Sciences* **81,** 151.
1396
1397 Zaugg J, Solenthaler F & Albrecht C. (2022). Materno-fetal iron transfer and the emerging
1398   role of ferroptosis pathways. *Biochem Pharmacol* **202,** 115141.
1399
1400 Zhang J, Johnson PC & Popel AS. (2009). Effects of erythrocyte deformability and
1401   aggregation on the cell free layer and apparent viscosity of microscopic blood
1402   flows. *Microvasc Res* **77,** 265-272.
1403
1404 Zhong H, Janer A, Khalimonchuk O, Antonicka H, Shoubridge Eric A & Barrientos A.
1405   (2023). BOLA3 and NFU1 link mitoribosome iron–sulfur cluster assembly to
1406   multiple mitochondrial dysfunctions syndrome. *Nucleic Acids Research* **51,**
1407   11797-11812.
1408
1409 Zygmunt M, Herr F, Münstedt K, Lang U & Liang OD. (2003). Angiogenesis and
1410   vasculogenesis in pregnancy. *European Journal of Obstetrics & Gynecology and*
1411   *Reproductive Biology* **110,** S10-S18.
1412
1413