Optimization of Genome-wide CRISPR Screens using dual-guide RNA Infection with Cas9

Electroporation (DICE)

Caitlin Purman[1,2]*, Charles Lu[1]*, Apexa Modi[1], Vinny Vijaykumar[1], Michael J. Flister[1,3], Anneke I. den

Hollander[1], Sabah Kadri[1], and Joshua D. Stender[1]

*=Equal contributions

[1]Genomics Research Center, AbbVie, Inc., North Chicago, IL 60064, USA; [2]Oncology Discovery, AbbVie,

Inc., North Chicago, IL 60064, USA; [3]Oncology Research Unit, Pfizer Inc, Bothell, WA

Author Emails:

Caitlin Purman: caitlin.purman@abbvie.com

Charles Lu: charles.lu@abbvie.com

Apexa Modi: apexa.modi@abbvie.com

Vinny Vijaykumar: vinny.vijaykumar@abbvie.com

Michael Flister: michael.flister@pfizer.com

Anneke den Hollander: anneke.denhollander@abbvie.com

Sabah Kadri: sabah.kadri@abbvie.com

Joshua Stender: Joshua.stender@abbvie.com

Abstract
Single-guide RNA lentiviral infection with Cas9 protein electroporation (SLICE) enables CRISPR screening
in primary cell types that require transient Cas9 expression yet is limited by scalability and robustness.
Here, we introduce dual-guide RNA infection with Cas9 electroporation (DICE), which expresses two
guides from the same lentiviral construct that target the same gene. In genome-wide screens, DICE
outperformed SLICE in defining essential genes and modulators of PD-L1 expression in IFN gamma
activated THP1 cells. Collectively, these data demonstrate that DICE can be utilized for reduced-scale
CRISPR screens in cell types with transient Cas9 protein expression without sacrificing screening quality.

Keywords: Dual-Cas9, DICE, SLICE, CRISPR, PDL1

Background
CRISPR-Cas systems have emerged as a potent technology to provide readily programmable and highly
efficient gene perturbations on single gene, multi-gene, and genome-wide scales. CRISPR-Cas dependent
gene editing requires a guide RNA (gRNA) to recruit a Cas DNA nuclease (typically Cas9) to areas of
homology in the genome where Cas9 creates a double-stranded break [1-3]. The imperfect repair of this
lesion through non-homologous end joining results in an insertion or deletion (indel) of bases in the
genome subsequently leading to a knockout of gene function [4]. While genome-wide CRISPR screens in
immortalized cell lines have become routine, CRISPR screens in some biologically relevant models, like
primary immune cells, present challenges that require modification of the standard approach [5-9].
Compared to immortalized cell lines, primary cells cannot be easily engineered to express Cas9
endogenously, and cell numbers are often limited, precluding the use of standard genome-wide CRISPR
libraries which require hundreds of millions of cells for adequate gRNA coverage. To overcome the
challenge of expressing Cas9 in primary immune cells, the single guide RNA (sgRNA) lentiviral infection
with Cas9 protein electroporation (SLICE) method for CRISPR genome editing was recently developed
[10]. SLICE enables CRISPR-mediated gene editing in cells that don't express Cas9 protein and has been
applied for genome-wide CRISPR screening in primary human T cells [10, 11]. While the development of
the SLICE approach represents a major innovation for CRISPR screening in primary cells, to our
knowledge, no head-to-head comparison of CRISPR screening with transient Cas9 (SLICE) versus
constitutive Cas9 expression has been performed.
62

Here we compare CRISPR screening using the Brunello library [5] in THP1 cells with constitutively Cas9
expression or in THP1 cells requiring Cas9 protein electroporation and demonstrate that screening
performance was more robust in THP1 cells constitutively expressing Cas9. To improve transient Cas9
screening approaches, we developed DICE, a multiplex gRNA lentiviral system expressing two gRNAs
targeting the same gene from the same construct followed by Cas9 electroporation. We designed a
miniaturized, genome-wide dual-Cas9 CRISPR library (21,034 constructs) and benchmarked it to the
gold-standard Brunello library (76,441 constructs). We observe that, despite the reduction in library size,
the dual-Cas9 library performs comparably to the Brunello genome-wide library in Cas9-expressing cells.
Moreover, we show that the dual-Cas9 library (DICE) outperforms the Brunello library (SLICE) when Cas9
protein is introduced via electroporation. We thus present a novel approach and library that improves

the outcome of genome-wide screens when Cas9 is exogenously introduced and drastically reduces the
number of cells and resources needed to perform a genome-wide screen.

Results

Genome-wide SLICE has limited performance relative to traditional screening
Genome-wide CRISPR screens are routinely executed in immortalized cell lines, as these cells can be
engineered to express Cas9 protein and expanded to essentially unlimited numbers; however, screening
in cell lines does not always yield disease relevant targets. Therefore, there is an emphasis on developing
approaches to enable CRISPR screening in primary cell models that are more likely to reflect disease
biology and donor diversity. A hybrid CRISPR approach termed Single guide RNA Lentiviral Infection with
Cas9 protein Electroporation (SLICE) was recently developed for pooled screening in primary human T
cells wherein cells are transduced with a lentiviral CRISPR sgRNA library and Cas9 is subsequently
introduced by electroporation [10]. Given that Cas9 activity within a cell persists for only 24-48 hours
[12, 13] when Cas9 is introduced via electroporation and that most published screens allow 1+ weeks for
genetic perturbation in Cas9-expressing cells, we sought to evaluate the global effect of Cas9 expression
modes on CRISPR screen performance.
To evaluate the contribution of Cas9 expression on screening performance, we executed genome-wide
screens for regulators of PD-L1 in THP-1 macrophages expressing Cas9 and in parental cells
electroporated with Cas9 protein (**Figure 1A**) in response to IFNγ stimulation. Briefly, we transduced
Cas9-expressing or parental THP-1 cells with the Brunello genome-wide library at 500X coverage with a
multiplicity of infection of 0.3 and selected for cells transduced with gRNAs. THP-1 parental cells
expressing gRNAs were electroporated with Cas9 protein to induce genome editing following the SLICE
protocol. Edited cells were then differentiated into macrophages with phorbol 12-myristate 13-acetate
(PMA) and treated with IFN gamma to induce PD-L1 surface expression. Cells were FACS sorted for the
top and bottom 10% of PD-L1 expressors. Thus, our screening workflow allowed us to assess library
performance using both a viability and FACS-based readout.

Receiver operating characteristic (ROC)-area under the curve (AUC) analysis of a logistic regression
classifier that was trained with reference sets containing essential and nonessential genes was
performed to estimate true positive and false positive rates for the Cas9 expressing cells and SLICE. This
analysis revealed that the screen executed in Cas9 expressing cells (AUC=0.94) outperformed the screen
with transient Cas9 expression (AUC=0.74) (**Figure 1B**). To further understand the difference in library
performance between the stable and transient Cas9 screens, we compared the performance of the four
gRNAs for each common essential gene in both screens. While the Z-score for guides targeting the same
gene were well-correlated in the integrated Cas9 screen, this correlation was markedly reduced in the
SLICE screen, indicating that the consistency of some individual guides is dependent on the mode and
duration of Cas9 expression (**Figure 1C**).

FACS-based screening identified key regulators of IFNy-induced PD-L1 expression (**Figure 1D-E**) [14-16].
As expected, in Cas9-expressing cells, the core IFNy signaling pathway genes (JAK1/2, STAT1, IFNGR1/2)
and CD274 itself were identified as top enriched genes in the PD-L1^{low} population (**Figure 1D**). In the
SLICE screen, some of the positive controls were identified as significant hits including STAT1 (LFC=1.4)

and JAK2 (LFC=1.2); however, strikingly, several core IFNy signaling pathway genes including IFNGR1
(LFC=0.479) IFNGR2 (LFC=0.504) and CD274 (LFC=0.429) were not significantly enriched in the PD-L1[low]
population (**Figure 1E**). The Z-scores of each of the four gRNA constructs for the known PD-L1 regulator
genes were enriched to similar levels in the integrated Cas9 screen (**Figure 1F**). In contrast, in the SLICE
screen, the variability in Z-scores for guides targeting the same gene increased dramatically, with some
guides showing no enrichment (**Figure 1G**). Taken together, these results indicate gRNAs enrichment
within genome-wide screens is highly dependent on the mode of Cas9 expression and delivery.

Dual-guide RNA lentiviral infection followed by Cas9 electroporation (DICE) improves editing efficiency
compared to SLICE
Multiplex Cas9 systems have emerged which express multiple sgRNAs which target the same gene or
multiple genes from the same lentiviral construct. These approaches improve gene editing efficiencies
compared to single guide systems [17]. Therefore, we reasoned that we could increase gene editing
efficiencies in experimental contexts requiring nucleofection of Cas9 protein by leveraging a recently
developed spCas9/dual gRNA approach wherein each lentiviral construct harbors two gRNAs targeting
separate genes. This dual gRNA Cas9 (dual-Cas9) system features lentiviral constructs designed with
separate U6-driven and H1-driven gRNA expression cassettes in opposing orientations. U6-VCR1 and H1-
WCR3 promoter-tracr pairings were chosen based on superior efficacy and reduced recombination rate
as previously reported [18]. Using this system, we designed lentiviral constructs with guides targeting
both CD47 and CD63 expressed from both proximal and distal positions (**Figure 2A**). We transduced cells
with both lentiviral constructs and then introduced Cas9 by electroporation. We observed greater than
75% knockout of both CD47 and CD63 with both construct orientations indicating that the U6 and H1
promoters express sgRNAs for similar gene editing efficiencies with transient Cas9 (**Figure 2A**).
134
Previous reports have demonstrated that utilizing multiple guides targeting the same gene increases the
probability of complete knockout by inducing deletions or indels [19, 20]. To increase the editing
efficiency of the single-gene perturbation Cas9 system, we designed a dual-Cas9 construct containing
two guide RNAs targeting the cell surface protein CD45 (**Figure 2B**). We assessed CD45 editing in cells
transduced with lentiviral single- or dual sgCD45 constructs followed by electroporation of Cas9. Single-
guide expression resulted in ~75% CD45 editing efficiency. However, editing efficiency was augmented
(~87%) using the dual guide expression construct with transient Cas9. We thus present a dual-guide RNA
lentiviral infection followed by Cas9 electroporation (DICE) approach which shows improved editing
efficiency compared to SLICE.

Design and benchmark of dual-Cas9 genome-wide library against the gold-standard Brunello library
To apply the DICE approach for genome-wide CRISPR screening, we designed a dual-Cas9 genome-wide
CRISPR library (**Figure 3A**). Given the observed increase in editing efficiency with two guides per
perturbation as opposed to one, we reasoned that we could efficiently target each gene with one dual-
Cas9 construct as opposed to the four separate single-guide constructs per gene in the standard Brunello
genome-wide library, reducing the total library size by 75%. Guide RNA sequences for the dual-Cas9
library were chosen based on the top two gRNAs for each target gene as predicted by Rule Set 3 [21],
which incorporates variations in tracrRNA sequences and improves accuracy in predicting on-target

efficiency over previous models. As the Brunello library was generated prior to the development of Rule

Set 3, ~70% of guides in the dual-Cas9 library are unique, while ~30% are also found in the Brunello

library **(Figure 3A).**

Dual-Cas9 and Brunello libraries perform comparably to identify common essential genes and PD-L1

regulators in Cas9-expressing THP-1 macrophages

We first compared the dual-Cas9 and Brunello library screening performance in THP-1 macrophages with

stable Cas9 expression. The Pearson correlation for essential genes between the dual-Cas9 and Brunello

was 0.806 and AUC score from ROC curve analyses indicated high and comparable performance between

the two libraries in the detection of essential genes (Brunello, AUC 0.94; dual-Cas9 AUC 0.93) (**Figure 3B**

and 3C). In addition to common essential genes, we assessed whether our dual-Cas9 library was able to

identify known regulators of PD-L1 expression in response to IFNγ treatment (**Figure 1A and 3D).** We

plotted the median LFC of genes in our sorted PD-L1^{low} versus our PD-L1^{high} populations and found that

all five of the known regulators as well as PD-L1 (CD274) itself were enriched in the PD-L1^{low} population

and depleted in PD-L1^{high} population in dual-Cas9 screen (**Figure 3D**). When we compared the Z-score of

each sgRNA construct in the Brunello (red) and dual-Cas9 (blue) libraries for five known regulator hits in

our PD-L1^{low} population, we observed high enrichment and tight distribution of all constructs targeting

each gene (**Figure 3E**). Collectively these results demonstrate that the novel dual-Cas9 library identifies

regulators of PDL1 expression at similar robustness as the gold-standard Brunello library.

DICE outperforms SLICE for genome-wide CRISPR screening

Given that our dual-Cas9 library performed comparably to the Brunello library in cells that stably express

Cas9 and that gene editing efficiency is higher with the DICE approach compared to SLICE, we

hypothesized that the dual-Cas9 library would outperform the Brunello library in screens where Cas9

expression is transient. The DICE platform would thus provide more robust gene editing to improve

screening quality as well as enable genome-wide scale screens in cell models where scale is limited. We

compared the performance of SLICE (Brunello) and DICE (dual-Cas9) genome-wide screens in our

viability and FACS-based screens (**Figure 1A**). Depletion of essential genes was more robust in the DICE

screen compared to SLICE, indicating improved editing efficiency with the dual-Cas9 library compared to

Brunello (**Figure 4A**). Similarly, the DICE approach significantly outperformed the SLICE approach as

shown by receiver operating characteristic (ROC)–area under the curve (AUC) analysis (**Figure 4B**). We

next asked if our DICE screening platform was better able to identify known regulators of PD-L1

compared to SLICE. In the FACS-based screens, DICE identified all five known regulators of PD-L1 as

significantly enriched in the PD-L1^{low} population whereas LFCs for known PD-L1 regulators were

significantly lower in the SLICE screen, and PD-L1 itself was not identified as a statistically significant hit

(**Figures 1E and 4C**). Moreover, the Z-score of each sgRNA construct in the Brunello (red) were generally

lower than the guides in the dual-Cas9 (blue) library for known regulators of our PD-L1 (**Figure 4D**). As

the sgRNA composition of the two libraries is not identical, performance of each library could depend on

the repertoire of sgRNAs in each library and/or the dual compared to the single sgRNA library format. Of

the five known regulators, both guides present in the dual-Cas9 library for JAK1 are also represented in

the Brunello library. Interestingly, while all three constructs performed comparably in the integrated

Cas9 screens, neither individual guide constructs in the Brunello library performed as well as the dual

guide construct in the transient Cas9 screens, pointing to a likely increase in editing efficiency with two

sgRNAs verses one when Cas9 is not stably expressed (**Figure 3E** and **Figure 4D**). These data suggest that

the DICE approach significantly outperforms the SLICE in viability- and FACS-based CRISPR screens.

Discussion

In this study, we present a new approach for improved CRISPR editing using single-target, dual sgRNA

expressing lentiviral constructs and Cas9 protein electroporation. We designed a genome-wide dual

sgRNA (dual-Cas9) library wherein each gene is targeted by two separate sgRNAs expressed from a single

construct, reducing the library size by 75% compared to the gold standard Brunello genome-wide library.

We compared the performance of the dual-Cas9 genome-wide CRISPR library to the Brunello library

using both viability- and FACS-based screening readouts. We confirm that the dual-Cas9 library performs

comparably to the Brunello library in cells expressing Cas9 but significantly outperforms the Brunello

library in screens where Cas9 is delivered exogenously. The observation that both libraries perform

similarly in the context of constitutive Cas9 suggests that genome-wide screens can be scaled down by

75% without missing valuable biological information. Miniature libraries can enable genome-wide

screening in cell types where cell numbers are limited such as primary immune and neuronal cells. At the

same time, the dual-Cas9 library, which expresses two sgRNAs targeting the same gene, provides a

performance advantage over Brunello library in screens where Cas9 must be introduced by

electroporation. A potential explanation for improved screening quality with the dual-Cas9 despite

reduced library size is that under transient expression, Cas9 protein has twice as many opportunities to

edit the genome in each cell with the dual-Cas9 system compared to a single guide library. This

explanation is supported by the tight distribution of LFCs for Brunello and Dual-Cas9 library constructs

targeting known PD-L1 regulators in screens in THP-1 Cas9+ cells, but relative loss of enrichment of many

Brunello constructs compared to Dual-Cas9 constructs in the SLICE screens. In SLICE experiments key

members of the JAK/STAT pathway were not identified as significant hits when using the Brunello library

suggesting that single guide libraries have the potential to miss important regulators of the targeted

biology. Our dual-Cas9 platform fills critical gaps in CRISPR screening capabilities to enable improved

screen performance when Cas9 is transient and reduces genome-wide screening scale, both of which are

critical for screening in primary human immune and neuronal cells.

Conclusions

This study presents a novel dual-Cas9 genome-wide CRISPR library consisting of one gRNA construct per

gene and compares its performance to the four-construct-per-gene Brunello library. In both viability- and

FACS-based screens, the dual-Cas9 library performs similarly to Brunello when Cas9 is constitutively

expressed and outperforms Brunello when Cas9 is delivered exogenously. The dual-Cas9 platform

enables screening with 75% fewer resources without compromising quality, making it ideal for primary

cells with limited numbers. Moreover, its superior performance in contexts with exogenous Cas9

highlights the potential of the dual-Cas9 library to capture biologically valuable information missed by

single-guide libraries.

Methods

Cell culture
THP-1 cells were maintained in RPMI 1640 (Gibco) supplemented with 10% heat inactivated fetal bovine
serum (FBS, Gibco), 1% Pen/Strep (Gibco), 1% L-glutamine (ThermoFisher Scientific, Cat# 25030081), 1%
Sodium pyruvate (ThermoFisher Scientific, Cat# 11360070), 1% HEPES (ThermoFisher Scientific, Cat#
15630-080), and 0.05 mM 2-mercaptoethanol (ThermoFisher Scientific, Cat# 21985023) in a 37°C
humidified 5% CO_2 incubator. Cells were differentiated by the addition of 10 ng/ml phorbol 12-myristate
13-acetate (Sigma, P1585) for 24 hours and treated with 10 ng/ml IFN gamma (PeproTech, Cat# 300-02)
for 48 hours.
243

Generation and characterization of Cas9-stable cells
Cas9-stable cells were generated by infecting parental cell lines with a lentiviral construct expressing
Cas9 and blasticidin-resistance gene in 12-well plates at 1000 x g for 1 h, in the presence of 8 µg/µL
polybrene (Sigma-Aldrich). Plates were then returned to 37 °C with 5% CO_2. Cells were incubated
overnight and then selected by 10 µg/ml blasticidin (Thermo Fisher Scientific). The activity of Cas9 was
confirmed to be greater than 75% through introducing guides targeting CD274, CD45, and confirming
loss of cell surface expression compared to a non-targeting control by flow cytometry.
251

Design of dual-Cas9 library
Guide RNAs for each human genes were designed using Broad's Rule Set 3 [21] and are listed in
Supplementary Table 1. The oligo library was synthesized at Twist Biosciences and cloned using a two-
step process where the guides were cloned into pvAA0017 backbone followed by introduction of the
TRACR RNA cassettes. The TRACR RNA sequences downstream of the U6 and H1 promoters are VCR1
and WCR3 respectively [18].
258

Genome-wide CRISPR screening
For the Brunello library transductions, 120 million THP1 cells or THP1 stably express Cas9 were
transfected with the sgRNA lentiviral library at multiplicity of infection (MOI) = 0.3 in three biological
replicates. The Brunello CRISPR-Cas9 knockout library contains 76,441 sgRNAs targeting 19,114 genes
with 1,000 non-targeting controls. For Dual-Cas9 library transductions, 30 million THP1 cells or THP1
stably express Cas9 were transfected with a lentiviral library at multiplicity of infection (MOI) = 0.3 in
three biological replicates. Following transduction, transduced cells were selected with puromycin (2
µg/ml, ThermoFisher Scientific, Cat# A1113802) and dead cells were removed (Miltenyi Biotec, Cat# 130-
090-101). For SLICE experiments, 800 pmol Cas9 was nucleofected per 20E6 cells using the Expanded T
cell program 2 on MaxCyte STx nucleofector. Cells were then differentiated with PMA and treated with
10 ng/ml IFN gamma for 24 hours. 10 (dual Cas9) and 40 million (Brunello) cells were collected as
presort for genomic DNA extraction for each replicate. Greater than 40 or 10 million cells per replicate
for Brunello or Dual-Cas9 experiments respectively were then processed and stained for anti-Human
PDL1 cell surface expression (Biolegend, Cat# A1113802) at 10 million cells/ml according to

manufacturer's recommendation. Cell populations expressing the top and bottom 10% PDL1 levels were
enriched through FACS sorting using Aria Fusion (BD Biosciences). At least one million PDL1^{low} and
PDL1^{high} cells were collected for genomic DNA extraction using Quick-DNA FFPE Kit (Zymo Research). PCR
reactions containing up to 10 μg genomic DNA in each reaction were performed using Titanium Taq DNA
Polymerase with primers to amplify the sgRNAs. Samples were then purified with SPRIselect beads,
mixed and sequenced by 75-bp single-end sequencing for Brunello libraries and 2x150 bp paired-end
sequencing for dual-Cas9 libraries.
280

CRISPR screen analysis
Pair-end sequencing data were demultiplexed and paired reads each with a sgRNA were quantified using
custom Perl scripts. Given that the position of the sgRNA could vary per read, the primer sequences
flanking the sgRNA were considered. Sequences in between the flanking primers were extracted and
then compared to sequences in the sgRNA library. Only sequences with no mismatches were used in the
calculation of guide-level read counts. Due to possibilities of recombination events, only pair of sgRNAs
that maps to the same gene or only 1 sgRNA were identified in each read-pair were used for
downstream analysis. Overall, samples with a minimum of 75% reads mapping to a sgRNAs were
considered for further analysis. Through the *edgeR* Bioconductor package, reads were normalized
between samples using the trimmed mean of M values (TMM) method [22]. Differential analysis of guide
level counts was performed using the *limma* R Bioconductor package [23]. Gene level effects were
considered as the median log fold change of the four guides associated with each gene. Statistically
significant genes were identified by aggregating differential sgRNAs ranked consistently higher at gene
level using robust rank aggregation (RRA) method. Benchmarking the performance of the various CRISPR
platform was described in [24].
296

297
298
299

307   **Figures:**

Figure 1: Performance of Brunello genome-wide CRISPR library with constitutive and transient Cas9 expression

A) Schematic overview of CRISPR screens to identify regulators of CD274 expression in THP-1 macrophages. Four screens were performed to enable a comparison of the dual-Cas9 and Brunello (single guide) platforms in the context of constitutive Cas9 expression and SLICE (electroporated Cas9). B) Single-gene receiver operating characteristic-area under the curve (ROC-AUC) plots derived from pan-essential and nonessential genes for Brunello screens in THP-1 or THP1-Cas9+ cells. C) Heatmap for the performance of each of four guide RNAs for common essential genes in THP1 cells with constitutive (integrated Cas9) or transient (SLICE) Cas9 expression. Data is displayed as Z-score normalized comparison between the guide counts in presort populations compared to pDNA levels. Comparison of gene enrichment in CD274^Low vs CD274^High sorted population in THP1 cells constitutively expressing Cas9 (D) or transient Cas9 expression (E). Density strip plots showing performance of each of the four guides targeting the core IFNγ signaling pathway in CD274^Low sorted population in THP1 cells constitutively expressing Cas9 (F) or transient Cas9 (G).

Figure 2: Development of DICE as strategy for multiplex gene editing

A) Schematic of dual-guide RNA expression cassettes targeting CD47 and CD63 (above). Evaluation of gene editing for CD47 and CD63 from guides driven by either the U6 and H1 promoters by flow cytometry (below). B) Comparison of targeting CD45 using single guides (SLICE, left) and dual-Cas9 constructs (DICE, right) using flow cytometry. CD45 levels were assessed by FACS at the indicated times post-transduction. Editing efficiency is indicated in the gray box in each FACS plot.

Figure 3: Miniature dual-Cas9 library provides comparable performance to Brunello library

A) Comparison and overlap of Brunello (single-guide) and dual-Cas9 genome-wide libraries. B) Scatter plot showing the correlation between lethality scores for essential genes in the Brunello and dual-Cas9 screens in THP-1 Cas9+ cells. C) Single-gene receiver operating characteristic-area under the curve (ROC-AUC) plots derived from pan-essential and nonessential genes for Brunello and dual-Cas9 screens in THP-1 Cas9+ cells. D) Scatter plots showing screening results for the dual-Cas9 screens. Each dot represents the median LFC of all sgRNAs targeting a particular gene. Dots for known CD274 regulator genes are labeled and shown in red. E) Density strip plots showing performance of guides targeting the core IFNγ signaling pathway in CD274^{Low} sorted population in THP1 cells constitutively expressing Cas9 and transduced with either the Brunello or Dual-Cas9 libraries.

Figure 4. dual-Cas9 outperforms Brunello to identify essential genes and PD-L1 regulators in THP-1 macrophages when Cas9 is introduced transiently

A) Scatter plot showing the correlation between lethality scores for common essential genes in SLICE vs DICE screening formats where the two Brunello library guides overlap with Dual-Cas9 library. B) Single-gene receiver operating characteristic-area under the curve plots derived from pan-essential and nonessential genes for Brunello and dual-Cas9 in Cas9+ THP1 cells or in either the SLICE or DICE screening format. C) Comparison of SLICE vs DICE for CD274^{low} populations. The core IFN pathway genes are highlighted in red. D) Density strip plots showing performance of guides targeting the core IFNγ signaling pathway in CD274^{Low} sorted population in SLICE (Red) and DICE (Blue). Guides that are shared between Brunello and Dual-Cas9 are dotted red lines.

Acknowledgements

We thank David Root, Olivia Bare, John Doench, and other members of the Genetic Perturbation Platform of The Broad Institute (Cambridge, MA) for the Brunello library as well as the design and cloning of the dual-Cas9 sgRNA library. We thank Jing Wang, Si Wu and Fedik Rahimov for critical review of the manuscript.

Author contributions

C.P. C.L., M.J.F, and J.D.S conceived, designed and led the study and drafted the manuscript. C.P., V.V., performed experiments, C.L., A.M., C.P., J.D.S., analyzed the data, S.K., M.J.F., A.I.D, provided resources and critical feedback. All authors reviewed the manuscript, provided feedback, and approved the final version of the manuscript.

References:

1. Jinek M, Chylinski K, Fonfara I, Hauer M, Doudna JA, Charpentier E: **A programmable dual-RNA-guided DNA endonuclease in adaptive bacterial immunity.** *Science* 2012, **337:**816-821.
2. O'Connell MR, Oakes BL, Sternberg SH, East-Seletsky A, Kaplan M, Doudna JA: **Programmable RNA recognition and cleavage by CRISPR/Cas9.** *Nature* 2014, **516:**263-266.
3. Sapranauskas R, Gasiunas G, Fremaux C, Barrangou R, Horvath P, Siksnys V: **The Streptococcus thermophilus CRISPR/Cas system provides immunity in Escherichia coli.** *Nucleic Acids Res* 2011, **39:**9275-9282.
4. Ran FA, Hsu PD, Wright J, Agarwala V, Scott DA, Zhang F: **Genome engineering using the CRISPR-Cas9 system.** *Nat Protoc* 2013, **8:**2281-2308.
5. Doench JG, Fusi N, Sullender M, Hegde M, Vaimberg EW, Donovan KF, Smith I, Tothova Z, Wilen C, Orchard R, et al: **Optimized sgRNA design to maximize activity and minimize off-target effects of CRISPR-Cas9.** *Nat Biotechnol* 2016, **34:**184-191.
6. Kampmann M: **CRISPR-based functional genomics for neurological disease.** *Nat Rev Neurol* 2020, **16:**465-480.
7. Katti A, Diaz BJ, Caragine CM, Sanjana NE, Dow LE: **CRISPR in cancer biology and therapy.** *Nat Rev Cancer* 2022, **22:**259-279.
8. Li YR, Lyu Z, Tian Y, Fang Y, Zhu Y, Chen Y, Yang L: **Advancements in CRISPR screens for the development of cancer immunotherapy strategies.** *Mol Ther Oncolytics* 2023, **31:**100733.
9. Shi H, Doench JG, Chi H: **CRISPR screens for functional interrogation of immunity.** *Nat Rev Immunol* 2023, **23:**363-380.
10. Shifrut E, Carnevale J, Tobin V, Roth TL, Woo JM, Bui CT, Li PJ, Diolaiti ME, Ashworth A, Marson A: **Genome-wide CRISPR Screens in Primary Human T Cells Reveal Key Regulators of Immune Function.** *Cell* 2018, **175:**1958-1971 e1915.

11.   Chen KY, Kibayashi T, Giguelay A, Hata M, Nakajima S, Mikami N, Takeshima Y, Ichiyama K, Omiya
R, Ludwig LS, et al: **Genome-wide CRISPR screen in human T cells reveals regulators of FOXP3.**
Nature 2025.

12.   Kim S, Kim D, Cho SW, Kim J, Kim JS: **Highly efficient RNA-guided genome editing in human cells**
via delivery of purified Cas9 ribonucleoproteins. *Genome Res* 2014, **24:**1012-1019.

13.   Liang X, Potter J, Kumar S, Zou Y, Quintanilla R, Sridharan M, Carte J, Chen W, Roark N,
Ranganathan S, et al: **Rapid and highly efficient mammalian cell engineering via Cas9 protein**
transfection. *J Biotechnol* 2015, **208:**44-53.

14.   Burr ML, Sparbier CE, Chan YC, Williamson JC, Woods K, Beavis PA, Lam EYN, Henderson MA, Bell
CC, Stolzenburg S, et al: **CMTM6 maintains the expression of PD-L1 and regulates anti-tumour**
immunity. *Nature* 2017, **549:**101-105.

15.   Mezzadra R, Sun C, Jae LT, Gomez-Eerland R, de Vries E, Wu W, Logtenberg MEW, Slagter M,
Rozeman EA, Hofland I, et al: **Identification of CMTM6 and CMTM4 as PD-L1 protein regulators.**
Nature 2017, **549:**106-110.

16.   Platanias LC: **Mechanisms of type-I- and type-II-interferon-mediated signalling.** *Nat Rev*
Immunol 2005, **5:**375-386.

17.   McCarty NS, Graham AE, Studena L, Ledesma-Amaro R: **Multiplexed CRISPR technologies for**
gene editing and transcriptional regulation. *Nat Commun* 2020, **11:**1281.

18.   Li R, Klingbeil O, Monducci D, Young MJ, Rodriguez DJ, Bayyat Z, Dempster JM, Kesar D, Yang X,
Zamanighomi M, et al: **Comparative optimization of combinatorial CRISPR screens.** *Nat*
Commun 2022, **13:**2469.

19.   Lukasiak S, Kalinka A, Gupta N, Papadopoulos A, Saeed K, McDermott U, Hannon GJ, Ross-
Thriepland D, Walter D: **A benchmark comparison of CRISPRn guide-RNA design algorithms and**
generation of small single and dual-targeting libraries to boost screening efficiency. *BMC*
Genomics 2025, **26:**198.

20.   Replogle JM, Bonnar JL, Pogson AN, Liem CR, Maier NK, Ding Y, Russell BJ, Wang X, Leng K, Guna
A, et al: **Maximizing CRISPRi efficacy and accessibility with dual-sgRNA libraries and optimal**
effectors. *Elife* 2022, **11**.

21.   DeWeirdt PC, McGee AV, Zheng F, Nwolah I, Hegde M, Doench JG: **Accounting for small**
variations in the tracrRNA sequence improves sgRNA activity predictions for CRISPR screening.
Nat Commun 2022, **13:**5255.

22.   Robinson MD, Oshlack A: **A scaling normalization method for differential expression analysis of**
RNA-seq data. *Genome Biol* 2010, **11:**R25.

23.   Ritchie ME, Phipson B, Wu D, Hu Y, Law CW, Shi W, Smyth GK: **limma powers differential**
expression analyses for RNA-sequencing and microarray studies. *Nucleic Acids Res* 2015,
43:e47.

24.   DeWeirdt PC, Sanson KR, Sangree AK, Hegde M, Hanna RE, Feeley MN, Griffith AL, Teng T, Borys
SM, Strand C, et al: **Optimization of AsCas12a for combinatorial genetic screens in human cells.**
Nat Biotechnol 2021, **39:**94-104.

446