Heterologous expression of carbonic anhydrase in *Acinetobacter* sp. Tol 5 for

whole-cell biocatalysis

Shogo Yoshimoto[1], Hiroya Oka[1,#], Yuki Ohara[2], Yan-Yu Chen[1], Masahito Ishikawa[1,3],

Katsutoshi Hori[1,*]

[1]Department of Biomolecular Engineering, Graduate School of Engineering, Nagoya

University, Nagoya, Aichi, Japan

[2]Friend Microbe Inc., Nagoya, Aichi, Japan

[3]Department of Bioscience, Nagahama Institute of Bio-Science and Technology,

Nagahama, Shiga, Japan

[#]Present address: US Department of Energy Joint Genome Institute, Lawrence Berkeley

National Laboratory, Berkeley, CA 94720, USA

*Corresponding author: Katsutoshi Hori (khori@chembio.nagoya-u.ac.jp)

Phone: +81-52-789-3339

Email: khori@chembio.nagoya-u.ac.jp

Abstract

Carbonic anhydrase accelerates the hydration of carbon dioxide (CO_2) and is an

attractive biocatalyst for carbon capture and utilization processes. *Acinetobacter* sp. Tol

5 shows high adhesiveness to materials due to its cell surface protein AtaA. Previously,

we developed a method for immobilizing bacteria using AtaA and demonstrated its

application to gas-phase bioproduction. In this study, we aimed to develop Tol 5 cells

expressing carbonic anhydrase and to evaluate CO_2 conversion ability as a whole-cell

biocatalyst. A codon-optimized carbonic anhydrase from *Sulfurihydrogenibium*

yellowstonense (SyCA) was first produced in the cytoplasm. Western blotting confirmed

expression, but the Wilbur–Anderson unit assay showed little carbonic anhydrase

activity as a whole-cell biocatalyst. To enhance activity, we fused six signal peptides

(SPs) to SyCA for periplasmic expression: PelB and TorA from *Escherichia coli*, the

Sec-type SP of Tol 5 outer-membrane protein Omp38, and three putative Tat-type SPs

from Tol 5. All constructs except the one fused to the SP of PelB produced SyCA

protein. The Omp38-SP fused SyCA was correctly processed to the expected molecular

size and exhibited increased whole-cell carbonic anhydrase activity, whereas other SPs

were not processed and showed smaller activities. These results show that periplasmic

expression of SyCA with a suitable SP is important for efficient CO_2 hydration in Tol 5.

The integration of the adhesive properties of Tol 5 with engineered periplasmic

expression of SyCA will lead to a simple platform for whole-cell biocatalysts in gas-

phase carbon capture and utilization applications.

Keywords

carbonic anhydrase, whole-cell biocatalyst, periplasmic expression, *Acinetobacter*

1. Introduction

The amount of carbon dioxide (CO_2) emissions as a result of human activity increased from 24.9 to 41 billion tons per year in the period from 1990 to 2022 emphasizing the urgent need for efficient, scalable carbon capture and utilization technologies [1]. Biocatalytic approaches that operate under ambient temperature and pressure without organic solvents have attracted particular attention for their sustainability. Carbonic anhydrase (CA; EC 4.2.1.1) is a zinc-dependent metalloenzyme that accelerates the reversible hydration of CO_2 to bicarbonate (HCO_3^-) with catalytic efficiencies approaching 10^6 s^{-1}, thereby enhancing CO_2 solubility by several orders of magnitude [2,3]. Solubilized bicarbonate (HCO_3^-) can be captured as calcium carbonate or utilized by microorganisms as a metabolic substrate [2,4].

The gram-negative bacterium *Acinetobacter* sp. Tol 5 exhibits high adhesiveness to various material surfaces, including hydrophobic plastics, hydrophilic glass, and metals, through its fibrous cell surface protein AtaA [5]. AtaA is a member of the trimeric autotransporter adhesin (TAA) family [6], and polypeptide chains of AtaA consisting of 3630 amino acids form a large homotrimeric fibrous structure that is over 1 MDa in molecular weight and 260 nm in length. TAAs, including AtaA, are secreted to the outer membrane (OM) by the type Vc secretion system [7]. We previously developed a method for immobilizing bacterial cells using AtaA; bacterial cells transformed with the *ataA* gene exhibit nonspecific high adhesiveness, and large amounts of growing and resting cells can be quickly and firmly immobilized onto various material surfaces [8,9]. The immobilized cells directly on surfaces through AtaA show better tolerance and increased chemical reaction rates and can be repeatedly used in reactions without inactivation [8]. Furthermore, cells immobilized via AtaA can be

placed directly in the gas phase, enabling efficient reactions with gaseous substrates

[10,11]. Therefore, Tol 5 is expected to be useful for CO_2 utilization. However, to the best

of our knowledge, there have been no reports of expressing a heterologous carbonic

anhydrase in *Acinetobacter*.

In this study, we aimed to develop Tol 5 expressing carbonic anhydrase and to

evaluate CO_2 conversion ability as a whole-cell biocatalyst.

2. Materials and methods

2.1. Bacterial strains

The restriction-modification system and *ataA*-deficient strain of *Acinetobacter* sp. Tol 5

REK123Δ*ataA* (Tol 5REK) was used as the host strain [12,13]. *Escherichia coli* DH5α was

used for plasmid construction. Tol 5REK was cultured in LB medium at 28°C, and DH5α

was cultured in LB medium at 37°C.

2.2. Plasmid construction

The oligonucleotides and synthetic DNA used in this study are listed in Table S1. An

Acinetobacter codon-optimized gene encoding SyCA (carbonic anhydrase from *S.*

yellowstonense) amino acid residues 21–246 was synthesized with a ribosome-binding

site immediately upstream and a C-terminal hexahistidine tag followed by a stop codon

and the T7 terminator. The synthesized gene was inserted into the EcoRI and XbaI sites

of pARP3, generating pARP3-SyCA. To construct a plasmid expressing SyCA fused to

a signal peptide (SP), the SyCA coding region was amplified by PCR from pARP3-

SyCA using primers SyCA-F and SyCA-R, producing Fragment-1. For PelB, TorA, and

Omp38, complementary oligonucleotides (SP-F and SP-R) were annealed to yield the

double-stranded fragment (Fragment-2). For Tat-1, Tat-2, or Tat-3, complementary

oligonucleotides (SP-F1 and SP-R1) were annealed, and a DNA fragment was amplified

by PCR using primers SP-F2 and SP-R2 with the double-stranded oligonucleotides as a

template, yielding Fragment-2′. The backbone of pARP3-SyCA was linearized with

EcoRI and XbaI to obtain Fragment-3. Fragments 1, 2 (or 2′), and 3 were assembled

using HiFi DNA Assembly Master Mix (New England Biolabs, Ipswich, United

Kingdom), generating the expression plasmid for SP-fused SyCA. The constructed

plasmid was introduced into Tol 5_{REK} cells by electroporation as previously described[13].

2.3. Protein expression

One milliliter of the overnight culture of Tol 5_{REK} harboring the expression plasmid was

inoculated into 100 ml of LB medium supplemented with gentamicin (10 µg/mL), 0.5%

arabinose, and 0.5 mM zinc sulfate and cultured at 28°C for 18 h at 115 rpm. Cells were

harvested by centrifugation at 5,000 × g for 5 min at 4°C. The cell pellet was

resuspended in SDS-sample buffer (5% (v/v) 2-mercaptoethanol, 2% (w/v) SDS, 0.02%

(w/v) bromophenol blue, 62.5 mM Tris-HCl, pH 6.8) and subjected to SDS-PAGE

followed by western blotting using an anti-His-tag antibody to verify expression.

2.4. Wilbur-Anderson unit (WAU) assay

CO_2 hydration activity was evaluated using the Wilbur-Anderson unit (WAU) assay [14]

with slight modifications. Cultured cells were harvested by centrifugation at 5,000 × g

for 5 min at 4°C, and the cell pellet was resuspended in pure water at an optical density

of 1.0 at 660 nm. Eight milliliters of 20 mM Tris-HCl (pH 8.3) and 1 ml of the cell

suspension were added to a 30-ml beaker on ice. Immediately after 6 ml of chilled CO_2-

saturated water was added to the beaker, the pH drop was recorded from 8.3 to 6.3 with stirring on ice using a pH sensor (PH-208; SATOTECH, Kanagawa, Japan) equipped with a logger (MJ-LOG2; SATOTEC). The obtained times for the cell suspension sample and blank (water) were designated T and T_0, respectively. The WAU was defined as $(T_0 - T)/T$. Differences among the samples were evaluated using the Brunner–Munzel test with Holm's step-down adjustment.

3. Results

3.1. Cytoplasmic expression of SyCA

The α-type carbonic anhydrase from *Sulfurihydrogenibium yellowstonense* (SyCA) [15] was chosen as the model enzyme because of its high catalytic efficiency and remarkable thermostability. A synthetic SyCA gene, codon-optimized for *Acinetobacter*, was cloned into the *Acinetobacter* expression plasmid pARP3 under the control of the arabinose-inducible promoter, with a C-terminal His-tag for detection. The constructed plasmid was introduced into Tol 5_{REK}, a restriction-modification system and *ataA*-deficient strain of Tol 5, for accurate enzymatic activity measurement due to its non-agglutination nature. SDS-PAGE of whole cell samples followed by western blotting with an anti-His antibody revealed a clear band corresponding to SyCA, confirming successful cytoplasmic expression (Fig. 1A).

Whole-cell carbonic anhydrase activity was assessed with the Wilbur–Anderson unit (WAU) assay [14], which measures the rate of CO_2 hydration to bicarbonate via the associated pH drop in CO_2-saturated water. A representative pH–time profile is shown in Fig. 1B. Cells expressing cytoplasmic SyCA produced a similar pH–time curve as the blank, which was water instead of cell suspension, and the time required to

reach pH 6.3 was almost the same. These results indicate that the cytoplasmic SyCA

exhibited little activity as a whole-cell biocatalyst under the conditions.

3.2. Selection of signal peptides for the periplasmic expression of SyCA

A previous study has shown that cytoplasmic expression of carbonic anhydrase leads to

poor enzymatic activity because the cell membrane limits CO_2 access and can even

cause intracellular toxicity, presumably via pH imbalance [16]. Therefore, periplasmic

expression is regarded as an alternative strategy for efficient heterologous expression of

carbonic anhydrase. We thus attempted to express SyCA in the periplasm of Tol 5_{REK}.

Proteins that need to be secreted to the periplasm are mediated by either the

general secretion (Sec) pathway or the twin-arginine translocation (Tat) pathway [17,18]. In

the Sec pathway, the unfolded polypeptide is secreted from the cytoplasm to the

periplasm through the SecYEG channel. A signal peptidase then cuts off its N-terminal

signal peptide (SP), and the protein folds in the periplasm. In the Tat system,

polypeptides that have already folded are secreted from the cytoplasm to the periplasm

through the TatABC complex, and then their SPs are cleaved by a signal peptidase. To

identify suitable signal peptides (SPs) for exporting SyCA, we first selected two well-

established *E. coli* SPs[16,19]: PelB for the Sec pathway and TorA for the twin-arginine

translocation (Tat) pathway, as candidates. The Sec-type SP of Omp38 [20], which is an

abundant OM protein known as OmpA [21], was also selected as a native Sec-type SP

from Tol 5. Because no Tat-type SPs have been characterized in Tol 5, open reading

frames of the Tol 5 genome were screened with SignalP [22], yielding three non-

lipoproteins (BCX74450, BCX75992, and BCX75163), two lipoproteins (BCX75681

and BCX73201), and one protein of uncertain classification (BCX72454). We selected

one representative from each of the three groups possessing putative Tat-type SPs,

resulting in three candidates: BCX72454, BCX75163, and BCX75681, which we named

Tat-1, Tat-2, and Tat-3, respectively. In total, six SPs were selected (Table 1). Each SP

sequence, spanning from the start methionine to its predicted cleavage site, was fused to

the N-terminus of SyCA through a short triglycine (GGG) linker (Fig. 2A). The six

fusion genes were cloned into the pARP3 vector, with a C-terminal His-tag, and were

introduced into Tol 5$_{REK}$.

3.3. Expression of SyCA fused to each signal peptide

Tol 5$_{REK}$ cells harboring the expression plasmid were cultured at 28 °C for 18 h with

0.5 % arabinose to induce the expression of SyCA. After cell lysis, proteins were

analyzed by western blotting with an anti-His antibody. SyCA was detected in every

construct except when fused to the SP of PelB (Fig. 2B). The constructs fused to SPs

from TorA, Tat-1, Tat-2, and Tat-3 exhibited slightly higher molecular weights than the

cytoplasmically expressed SyCA, suggesting that their SPs were not removed. In

contrast, the SyCA fused to the SP of Omp38 showed a band at the same size as the

cytoplasmic SyCA, indicating proper cleavage of its SP and secretion into the

periplasm. These results suggest that the Omp38-SP is the most suitable for periplasmic

expression of SyCA in Tol 5$_{REK}$.

3.4. Catalytic activity of cells expressing SyCA fused to each signal peptide

We evaluated whether fusion with the SPs improved whole-cell carbonic anhydrase

activity. All SyCA-producing constructs, except the one fused to PelB-SP, were tested

using the WAU assay. Figure 3A shows the representative pH-time profile: cells

expressing SyCA fused to each SP lowered the pH faster than the blank, which was water instead of a cell suspension. This suggests that these cells exhibit carbonic anhydrase activity as a whole-cell biocatalyst. The WAU values calculated from three independent measurements were summarized in Figure 3B. The Omp38-SP, which follows the Sec pathway, yielded the highest activity. Tat-type SPs from either *E. coli* (TorA) or Tol 5 gave lower activity. The Brunner–Munzel test was applied with Omp38 as the reference. After Holm adjustment for four comparisons, Omp38 showed significantly higher WAU than TorA (adjusted $p < 0.01$), Tat-2 (adjusted $p < 0.01$), Tat-3 (adjusted $p < 0.01$), and Tat-1 (adjusted $p = 0.013$). These results indicate that periplasmic expression with a suitable SP is important for achieving effective whole-cell biocatalysis of carbonic anhydrase in Tol 5.

4. Discussion

In this study, we investigated the expression of the carbonic anhydrase SyCA in Tol 5_{REK} and its CO_2 conversion activity as a whole-cell biocatalyst. Fusing SyCA with the SP of Tol 5 Omp38 led to efficient processing of its SP, resulting in the highest WAU value (Figs. 2, 3). These results highlight that periplasmic expression and the selection of an appropriate SP are important for the catalytic performance of carbonic anhydrase as a whole-cell biocatalyst. To the best of our knowledge, this is the first report on the heterologous expression of carbonic anhydrase in *Acinetobacter* for carbon utilization.

Several studies have explored the periplasmic expression of carbonic anhydrase in *E. coli*. Jo *et al.* expressed NgCA using the PelB (Sec) and TorA (Tat) signal peptides and found that the TorA construct yielded higher catalytic activity [16]. Patel *et al.* expressed the Cab and Cam with PelB (Sec) and gIII (Sec) signals, reporting that PelB

conferred slightly higher activity for both enzymes [23]. More recently, Jo *et al.* showed

that the periplasmic expression of TdCA with the PelB (Sec) signal peptide enhanced

enzyme activity [24]. Collectively, these findings indicate that the optimal signal peptide

for periplasmic expression appears to be specific to each carbonic anhydrase even in *E.*

coli. In Tol 5, the *E. coli* PelB-SP did not yield detectable SyCA production (Fig. 2), and

the *E. coli* TorA-SP produced only weak activity (Fig. 3). These results suggest that SPs

from *E. coli* do not match the secretion machinery of Tol 5 and are consistent with

previous reports that variations in signal peptide sequences influence secretion

efficiency and that some signal peptides exhibit species specificity [25,26]. The three Tat-

type SPs from Tol 5 were not fully cleaved and also yielded lower activity (Figs. 2, 3).

In contrast, the native Sec-type Omp38-SP from Tol 5 was cleaved efficiently and gave

the best activity, demonstrating its suitability for expressing SyCA in this host.

Tol 5 cells can be immobilized on many materials through AtaA and can be

applied to gas-phase reactions [10,11]. Combining the bacterial cell immobilization with

the periplasmic expression of SyCA may lead to stable, reusable catalysts for gas-phase

CO_2 hydration or for supplying bicarbonate to downstream biosynthetic steps. Such

systems could contribute to efficient biological carbon capture and conversion

technologies.

5. Conclusion

In this study, we demonstrated that periplasmic expression and selection of an

appropriate SP are important for the catalytic performance of carbonic anhydrase in Tol

5 as a whole-cell biocatalyst. Integrating the adhesive properties of Tol 5 with

engineered periplasmic expression of SyCA could lead to efficient CO_2 capture and

conversion processes.

CRediT authorship contribution statement

Shogo Yoshimoto: Conceptualization, Validation, Writing - Original Draft. **Hiroya**

Oka: Investigation. **Yuki Ohara**: Investigation. **Yan-Yu Chen**: Investigation. **Masahito**

Ishikawa: Investigation. **Katsutoshi Hori**: Conceptualization, Writing - Review &

Editing, Funding Acquisition

Acknowledgements

The authors thank Kensho Abe, Chiharu Kitano, and Eriko Kawamoto for their

technical assistance. This research was supported by the Japan Society for the

Promotion of Science (JSPS) KAKENHI (Grant Number JP24H00043) to KH and SY.

Declaration of competing interest

The authors declare no competing interests.

Data availability

All data generated or analyzed during this study are included in this published article

and its supplementary materials.

References

1   Yurak, V. V. & Fedorov, S. A. Review of natural and anthropogenic emissions of
256   carbon dioxide into the earth's atmosphere. *Int J Environ Sci Tech* **22**, 2719-2736
257   (2025).
258   2   Talekar, S., Jo, B. H., Dordick, J. S. & Kim, J. Carbonic anhydrase for CO_2

259  capture, conversion and utilization. *Curr Opin Biotech* **74**, 230-240 (2022).

260 3 Shao, P. J., Ye, J. X., Shen, Y., Zhang, S. H. & Zhao, J. K. Recent advancements

261  in carbonic anhydrase for CO_2 capture: A mini review. *Gas Sci Eng* **123**, 205237

262  (2024).

263 4 Liu, Z. H., Wang, K., Chen, Y., Tan, T. W. & Nielsen, J. Third-generation

264  biorefineries as the means to produce fuels and chemicals from CO_2. *Nat Catal*

265  **3**, 274-288 (2020).

266 5 Ishikawa, M., Nakatani, H. & Hori, K. AtaA, a new member of the trimeric

267  autotransporter adhesins from *Acinetobacter* sp. Tol 5 mediating high

268  adhesiveness to various abiotic surfaces. *PLoS One* **7**, e48830 (2012).

269 6 Linke, D., Riess, T., Autenrieth, I. B., Lupas, A. & Kempf, V. A. J. Trimeric

270  autotransporter adhesins: variable structure, common function. *Trends Microbiol*

271  **14**, 264-270 (2006).

272 7 Leo, J. C., Grin, I. & Linke, D. Type V secretion: mechanism(s) of autotransport

273  through the bacterial outer membrane. *Philos T R Soc B* **367**, 1088-1101 (2012).

274 8 Ishikawa, M., Shigemori, K. & Hori, K. Application of the adhesive

275  bacterionanofiber AtaA to a novel microbial immobilization method for the

276  production of indigo as a model chemical. *Biotechnol Bioeng* **111**, 16-24 (2014).

277 9 Hori, K., Ohara, Y., Ishikawa, M. & Nakatani, H. Effectiveness of direct

278  immobilization of bacterial cells onto material surfaces using the

279  bacterionanofiber protein AtaA. *Appl Microbiol Biotechnol* **99**, 5025-5032

280  (2015).

281 10 Yoshimoto, S., Ohara, Y., Nakatani, H. & Hori, K. Reversible bacterial

282  immobilization based on the salt-dependent adhesion of the bacterionanofiber

283  protein AtaA. *Microb Cell Fact* **16**, 123 (2017).

284 11 Usami, A., Ishikawa, M. & Hori, K. Gas-phase bioproduction of a high-value-

285  added monoterpenoid (*E*)-geranic acid by metabolically engineered

286  *Acinetobacter* sp. Tol 5. *Green Chem* **22**, 1258-1268 (2020).

287 12 Inoue, S., Yoshimoto, S. & Hori, K. A new target of multiple lysine methylation

288  in bacteria. *J Bacteriol* **207**, e00325-00324 (2025).

289 13 Ishikawa, M. & Hori, K. The elimination of two restriction enzyme genes allows

290  for electroporation-based transformation and CRISPR-Cas9-based base editing

291  in the non-competent Gram-negative bacterium *Acinetobacter* sp. Tol 5. *Appl*

292  *Environ Microb* **90**, e0040024 (2024).

293 14 Wilbur, K. M. & Anderson, N. G. Electrometric and colorimetric determination

294  of carbonic anhydrase. *Journal of Biological Chemistry* **176**, 147-154 (1948).

15 Capasso, C., De Luca, V., Carginale, V., Cannio, R. & Rossi, M. Biochemical
properties of a novel and highly thermostable bacterial α-carbonic anhydrase
from *Sulfurihydrogenibium yellowstonense* YO3AOP1. *J Enzym Inhib Med Ch*
27, 892-897 (2012).
16 Jo, B. H., Kim, I. G., Seo, J. H., Kang, D. G. & Cha, H. J. Engineered
Escherichia coli with periplasmic carbonic anhydrase as a biocatalyst for CO_2
sequestration. *Appl Environ Microb* **79**, 6697-6705 (2013).
17 Natale, P., Brüser, T. & Driessen, A. J. M. Sec- and Tat-mediated protein
secretion across the bacterial cytoplasmic membrane - Distinct translocases and
mechanisms. *Bba-Biomembranes* **1778**, 1735-1756 (2008).
18 Mergulhao, F. J. M., Summers, D. K. & Monteiro, G. A. Recombinant protein
secretion in *Escherichia coli*. *Biotechnol Adv* **23**, 177-202 (2005).
19 Matos, C. F. R. O. *et al.* High-yield export of a native heterologous protein to the
periplasm by the tat translocation pathway in *Escherichia coli*. *Biotechnology*
and Bioengineering **109**, 2533-2542 (2012).
20 Ishikawa, M., Yoshimoto, S., Hayashi, A., Kanie, J. & Hori, K. Discovery of a
novel periplasmic protein that forms a complex with a trimeric autotransporter
adhesin and peptidoglycan. *Mol Microbiol* **101**, 394-410 (2016).
21 Confer, A. W. & Ayalew, S. The OmpA family of proteins: Roles in bacterial
pathogenesis and immunity. *Vet Microbiol* **163**, 207-222 (2013).
22 Teufel, F. *et al.* SignalP 6.0 predicts all five types of signal peptides using
protein language models. *Nat Biotechnol* **40**, 1023-1025 (2022).
23 Patel, T. N., Park, A. H. A. & Banta, S. Periplasmic expression of carbonic
anhydrase in *Escherichia coli*: A new biocatalyst for CO_2 hydration.
Biotechnology and Bioengineering **110**, 1865-1873 (2013).
24 Jo, B. H. & Hwang, I. S. Characterization and high-level periplasmic expression
of thermostable α-carbonic anhydrase from *Thermosulfurimonas dismutans* in
Escherichia coli for CO_2 capture and utilization. *International Journal of*
Molecular Sciences **21**, 103 (2020).
25 Blaudeck, N., Sprenger, G. A., Freudl, R. & Wiegert, T. Specificity of signal
peptide recognition in Tat-dependent bacterial protein translocation. *J Bacteriol*
183, 604-610 (2001).
26 Freudl, R. Signal peptides for recombinant protein secretion in bacterial
expression systems. *Microbial Cell Factories* **17**, 52 (2018).
329
330

 Figure captions

 Figure 1. Cytoplasmic expression of SyCA. (A) Western blotting of whole-cell lysates

 from Tol 5$_{REK}$ cells carrying the SyCA expression plasmid (+) or carrying no plasmid

 (−). Proteins were separated by SDS-PAGE, transferred to a PVDF membrane, and

 probed with an anti-His antibody; the arrowhead marks the band corresponding to

 SyCA. (B) Time course of the pH changes during the WAU assay. Carbonic anhydrase

 activity of cells expressing cytoplasmic SyCA or blank (water only, no cells) was

 measured.

344 **Figure 2. Expression of SyCA variants fused to each signal peptide.** (A) Schematic

345 representation of SyCA fused to a signal peptide (SP). (B) Protein expression was

346 detected by western blotting using an anti-His antibody. The arrowhead indicates the

347 band corresponding to the cytoplasmically expressed SyCA (Non-SP).

Figure 3. Carbonic anhydrase activity of cells expressing SyCA variants fused to each signal peptide. (A) Time course of the pH changes during the WAU assay of cells expressing SyCA variants fused to each signal peptide. (B) The WAU values calculated from three independent measurements. Data are expressed as mean ± SEM (n = 3). Statistical significance, *adjusted p = 0.013, **adjusted p < 0.01.

356 **Table 1. Signal peptides used in this study.**

Source	Gene	Sec/Tat	Sequence
E. coli	*pelB*	Sec	MKYLLPTAAAGLLLLAAQPAMA
E. coli	*torA*	Tat	MNNNDLFQASRRRFLAQLGGLTVAGMLGPSLLTPRRATA
Tol 5	*omp38*	Sec	MKLSRIALAMLVAAPLAAANAG
Tol 5	BCX72454 (Tat-1)	Tat	MTDLTPYHEDQELDNNTSNNIHFRDILEQRISRRSLITKTASGAAALALASSLTG
Tol 5	BCX75163 (Tat-2)	Tat	MNKKVDFDPAKRDFLKTMGSLAAFGAVTTWMPVQSAKA
Tol 5	BCX75681 (Tat-3)	Tat	MTNSANQHTRREVLTWLASIPFLPLGAMATSATLAG