**Heterologous expression of carbonic anhydrase in *Acinetobacter* sp. Tol 5 for**

**whole-cell biocatalysis**

Running head: Expression of carbonic anhydrase in *Acinetobacter*

Shogo Yoshimoto[1], Hiroya Oka[1,#], Yuki Ohara[2], Yan-Yu Chen[1], Masahito Ishikawa[1,3],

Katsutoshi Hori[1,*]

[1]Department of Biomolecular Engineering, Graduate School of Engineering, Nagoya

University, Nagoya, Aichi, Japan

[2]Friend Microbe Inc., Nagoya, Aichi, Japan

[3]Department of Bioscience, Nagahama Institute of Bio-Science and Technology,

Nagahama, Shiga, Japan

[#]Present address: US Department of Energy Joint Genome Institute, Lawrence Berkeley

National Laboratory, Berkeley, CA 94720, USA

*Corresponding author: Katsutoshi Hori (khori@chembio.nagoya-u.ac.jp)

Phone: +81-52-789-3339

Email: khori@chembio.nagoya-u.ac.jp

**Abstract**

Carbonic anhydrase accelerates the hydration of carbon dioxide ($CO_2$) and is an attractive biocatalyst for carbon capture and utilization. *Acinetobacter* sp. Tol 5 shows high adhesiveness via its cell-surface protein AtaA. We previously demonstrated its application to bacterial immobilization and gas-phase bioproduction. Here, we developed Tol 5 cells expressing carbonic anhydrase and evaluated $CO_2$ conversion ability as whole-cell biocatalysts. A codon-optimized carbonic anhydrase from *Sulfurihydrogenibium yellowstonense* (SyCA) was produced in the cytoplasm, but the cells showed little activity as a whole-cell biocatalyst. To enhance activity, we fused six signal peptides (SPs) to SyCA for periplasmic expression. The Omp38-SP fusion of SyCA was properly processed to the mature size, yielding higher whole-cell activity. By contrast, the other constructs were either undetectable or remained unprocessed, resulting in lower activities. These results show that periplasmic expression of SyCA is important for efficient $CO_2$ hydration in Tol 5 cells as whole-cell biocatalysts.

**Keywords**

carbonic anhydrase, whole-cell biocatalyst, periplasmic expression, *Acinetobacter*

**Introduction**

The amount of carbon dioxide ($CO_2$) emissions as a result of human activity increased from 24.9 to 41 billion tons per year in the period from 1990 to 2022 emphasizing the urgent need for efficient, scalable carbon capture and utilization technologies (Yurak and Fedorov, 2025). A variety of strategies, including physical absorption, chemical solvent scrubbing, membrane separation, and physical and chemical adsorption, have been explored to mitigate $CO_2$ emissions (Hanifa et al., 2023; Kammerer et al., 2023; Nagireddi et al., 2023). Among them, biocatalytic and bio-integrated approaches that operate under ambient temperature and pressure without organic solvents are methods currently attracting significant attention because of their sustainability (Carceller et al., 2023; Marra et al., 2024; Sieborg et al., 2024).

Carbonic anhydrase (EC 4.2.1.1) is a zinc-dependent metalloenzyme that accelerates the reversible hydration of $CO_2$ to bicarbonate ($HCO_3^-$). Solubilized bicarbonate can be captured as calcium carbonate or utilized by microorganisms as a metabolic substrate (Liu et al., 2020; Talekar et al., 2022). The catalytic efficiency of carbonic anhydrase is very high, with a $k_{cat}$ value of $10^6$ $s^{-1}$, thereby enhancing $CO_2$ solubility by several orders of magnitude (Khalifah, 1971; Talekar et al., 2022; Shao et al., 2024). Many studies have shown that purified carbonic anhydrase markedly accelerates $CO_2$ capture (Talekar et al., 2022); however, the scale required for industrial carbon capture makes the use of purified enzymes economically impractical. Consequently, whole-cell catalysts that do not require a costly purification process are gaining attention as a cost-effective alternative (Moon et al., 2020; Ting et al., 2022; Zhu et al., 2022; Baidya et al., 2024).

The gram-negative bacterium *Acinetobacter* sp. Tol 5 exhibits high

adhesiveness to various material surfaces, including hydrophobic plastics, hydrophilic

glass, and metals, through its fibrous cell surface protein AtaA (Ishikawa et al., 2012).

We previously developed a method for immobilizing bacterial cells using AtaA;

bacterial cells transformed with the *ataA* gene exhibit nonspecific high adhesiveness,

and large amounts of growing and resting cells can be quickly and firmly immobilized

onto various material surfaces (Ishikawa et al., 2014; Hori et al., 2015). The

immobilized cells directly on surfaces through AtaA show better tolerance and increased

chemical reaction rates and can be repeatedly used in reactions without inactivation

(Ishikawa et al., 2014). Furthermore, cells immobilized via AtaA can be placed directly

in the gas phase, enabling efficient reactions with gaseous substrates (Yoshimoto et al.,

2017; Usami et al., 2020). Therefore, Tol 5 is expected to be useful for $CO_2$ utilization.

However, to the best of our knowledge, there have been no reports of expressing a

heterologous carbonic anhydrase in *Acinetobacter*.

In this study, we aimed to develop Tol 5 expressing carbonic anhydrase and to

evaluate $CO_2$ conversion ability as a whole-cell biocatalyst.

**Materials and methods**

**Bacterial strains**

The restriction-modification system and *ataA*-deficient strain of *Acinetobacter* sp. Tol 5

REK123Δ*ataA* (Tol 5$_{REK}$) was used as the host strain (Ishikawa and Hori, 2024; Inoue et

al., 2025). *Escherichia coli* DH5α was used for plasmid construction. Tol 5$_{REK}$ was

cultured in LB medium at 28°C, and DH5α was cultured in LB medium at 37°C.

**Plasmid construction**

The oligonucleotides and synthetic DNA used in this study are listed in Table S1. An

*Acinetobacter* codon-optimized gene encoding SyCA (carbonic anhydrase from *S.*

*yellowstonense*) amino acid residues 21–246 was synthesized with a ribosome-binding

site immediately upstream and a C-terminal hexahistidine tag followed by a stop codon

and the T7 terminator. The synthesized gene was inserted into the EcoRI and XbaI sites

of pARP3, generating pARP3-SyCA. To construct a plasmid expressing SyCA fused to

a signal peptide (SP), the SyCA coding region was amplified by PCR from pARP3-

SyCA using primers SyCA-F and SyCA-R, producing Fragment-1. For PelB, TorA, and

Omp38, complementary oligonucleotides (SP-F and SP-R) were annealed to yield the

double-stranded fragment (Fragment-2). For Tat-1, Tat-2, or Tat-3, complementary

oligonucleotides (SP-F1 and SP-R1) were annealed, and a DNA fragment was amplified

by PCR using primers SP-F2 and SP-R2 with the double-stranded oligonucleotides as a

template, yielding Fragment-2′. The backbone of pARP3-SyCA was linearized with

EcoRI and XbaI to obtain Fragment-3. Fragments 1, 2 (or 2′), and 3 were assembled

using HiFi DNA Assembly Master Mix (New England Biolabs, Ipswich, United

Kingdom), generating the expression plasmid for SP-fused SyCA. The constructed

plasmid was introduced into Tol 5$_{REK}$ cells by electroporation as previously

described(Ishikawa and Hori, 2024).

**Protein expression**

One milliliter of the overnight culture of Tol 5$_{REK}$ harboring the expression plasmid was

inoculated into 100 ml of LB medium supplemented with gentamicin (10 µg/mL), 0.5%

arabinose, and 0.5 mM zinc sulfate and cultured at 28°C for 18 h at 115 rpm. Cells were

harvested by centrifugation at 5,000 × g for 5 min at 4°C. The cell pellet was

resuspended in SDS-sample buffer (5% (v/v) 2-mercaptoethanol, 2% (w/v) SDS, 0.02%

(w/v) bromophenol blue, 62.5 mM Tris-HCl, pH 6.8) and subjected to SDS-PAGE

followed by western blotting using an anti-His-tag antibody to verify expression.

**Wilbur-Anderson unit (WAU) assay**

$CO_2$ hydration activity was evaluated using the Wilbur and Anderson unit (WAU) assay

(Wilbur and Anderson, 1948) with slight modifications. Cultured cells were harvested

by centrifugation at $5,000 \times g$ for 5 min at 4°C, and the cell pellet was resuspended in

pure water at an optical density of 1.0 at 660 nm. Eight milliliters of 20 mM Tris-HCl

(pH 8.3) and 1 ml of the cell suspension were added to a 30-ml beaker on ice.

Immediately after 6 ml of chilled $CO_2$-saturated water was added to the beaker, the pH

drop was recorded from 8.3 to 6.3 with stirring on ice using a pH sensor (PH-208;

SATOTECH, Kanagawa, Japan) equipped with a logger (MJ-LOG2; SATOTEC). The

obtained times for the cell suspension sample and blank (water) were designated T and

$T_0$, respectively. The WAU was defined as $(T_0 - T)/T$. Differences among the samples

were evaluated using the Brunner–Munzel test with Holm's step-down adjustment.

**Results**

**Cytoplasmic expression of SyCA**

The α-type carbonic anhydrase from *Sulfurihydrogenibium yellowstonense* (SyCA)

(Capasso et al., 2012) was chosen as the model enzyme because of its high catalytic

efficiency and remarkable thermostability. A synthetic SyCA gene, codon-optimized for

*Acinetobacter*, was cloned into the *Acinetobacter* expression plasmid pARP3 under the

control of the arabinose-inducible promoter, with a C-terminal His-tag for detection.

The constructed plasmid was introduced into Tol $5_{REK}$, a restriction-modification system

and *ataA*-deficient strain of Tol 5, for accurate enzymatic activity measurement due to

its non-agglutination nature. SDS-PAGE of whole cell samples followed by western

blotting with an anti-His antibody revealed a clear band corresponding to SyCA,

confirming successful cytoplasmic expression (Fig. 1A).

137            Whole-cell carbonic anhydrase activity was assessed with the Wilbur–

Anderson unit (WAU) assay (Wilbur and Anderson, 1948), which measures the rate of

$CO_2$ hydration to bicarbonate via the associated pH drop in $CO_2$-saturated water. A

representative pH–time profile is shown in Fig. 1B. Cells expressing cytoplasmic SyCA

produced a similar pH–time curve as the blank, which was water instead of cell

suspension, and the time required to reach pH 6.3 was almost the same. These results

indicate that the cytoplasmic SyCA exhibited little activity as a whole-cell biocatalyst

under the conditions.

**146    Selection of signal peptides for the periplasmic expression of SyCA**

A previous study has shown that cytoplasmic expression of carbonic anhydrase leads to

poor enzymatic activity because the cell membrane limits $CO_2$ access and can even

cause intracellular toxicity, presumably via pH imbalance (Jo et al., 2013). Therefore,

periplasmic expression is regarded as an alternative strategy for efficient heterologous

expression of carbonic anhydrase. We thus attempted to express SyCA in the periplasm

of Tol $5_{REK}$.

153            Proteins that need to be secreted to the periplasm are mediated by either the

general secretion (Sec) pathway or the twin-arginine translocation (Tat) pathway

(Mergulhao et al., 2005; Natale et al., 2008). In the Sec pathway, the unfolded

polypeptide is secreted from the cytoplasm to the periplasm through the SecYEG

channel. A signal peptidase then cuts off its N-terminal signal peptide (SP), and the

protein folds in the periplasm. In the Tat system, polypeptides that have already folded

are secreted from the cytoplasm to the periplasm through the TatABC complex, and

then their SPs are cleaved by a signal peptidase. To identify suitable signal peptides

(SPs) for exporting SyCA, we first selected two well-established *E. coli* SPs(Matos et

al., 2012; Jo et al., 2013): PelB for the Sec pathway and TorA for the twin-arginine

translocation (Tat) pathway, as candidates. The Sec-type SP of Omp38 (Ishikawa et al.,

2016), which is an abundant OM protein known as OmpA (Confer and Ayalew, 2013),

was also selected as a native Sec-type SP from Tol 5. Because no Tat-type SPs have

been characterized in Tol 5, open reading frames of the Tol 5 genome were screened

with SignalP (Teufel et al., 2022), yielding three non-lipoproteins (BCX74450,

BCX75992, and BCX75163), two lipoproteins (BCX75681 and BCX73201), and one

protein of uncertain classification (BCX72454). We selected one representative from

each of the three groups possessing putative Tat-type SPs, resulting in three candidates:

BCX72454, BCX75163, and BCX75681, which we named Tat-1, Tat-2, and Tat-3,

respectively. In total, six SPs were selected (Table 1). Each SP sequence, spanning from

the start methionine to its predicted cleavage site, was fused to the N-terminus of SyCA

through a short triglycine (GGG) linker (Fig. 2A). The six fusion genes were cloned into

the pARP3 vector, with a C-terminal His-tag, and were introduced into Tol 5$_{REK}$.

**Expression of SyCA fused to each signal peptide**

Tol 5$_{REK}$ cells harboring the expression plasmid were cultured at 28 °C for 18 h with

0.5 % arabinose to induce the expression of SyCA. After cell lysis, proteins were

analyzed by western blotting with an anti-His antibody. SyCA was detected in every

construct except when fused to the SP of PelB (Fig. 2B). The constructs fused to SPs

from TorA, Tat-1, Tat-2, and Tat-3 exhibited slightly higher molecular weights than the

cytoplasmically expressed SyCA, suggesting that their SPs were not removed. In

contrast, the SyCA fused to the SP of Omp38 showed a band at the same size as the

cytoplasmic SyCA, indicating proper cleavage of its SP and secretion into the

periplasm. These results suggest that the Omp38-SP is the most suitable for periplasmic

expression of SyCA in Tol $5_{REK}$.

**Catalytic activity of cells expressing SyCA fused to each signal peptide**

We evaluated whether fusion with the SPs improved whole-cell carbonic anhydrase

activity. All SyCA-producing constructs, except the one fused to PelB-SP, were tested

using the WAU assay. Figure 3A shows the representative pH-time profile: cells

expressing SyCA fused to each SP lowered the pH faster than the blank, which was

water instead of a cell suspension. This suggests that these cells exhibit carbonic

anhydrase activity as a whole-cell biocatalyst. The WAU values calculated from three

independent measurements were summarized in Figure 3B. The Omp38-SP, which

follows the Sec pathway, yielded the highest activity. Tat-type SPs from either *E. coli*

(TorA) or Tol 5 gave lower activity. The Brunner–Munzel test was applied with Omp38

as the reference. After Holm adjustment for four comparisons, Omp38 showed

significantly higher WAU than TorA (adjusted $p < 0.01$), Tat-2 (adjusted $p < 0.01$), Tat-3

(adjusted $p < 0.01$), and Tat-1 (adjusted $p = 0.013$). These results indicate that

periplasmic expression with a suitable SP is important for achieving effective whole-cell

biocatalysis of carbonic anhydrase in Tol 5.

**Discussion**

In this study, we investigated the expression of the carbonic anhydrase SyCA in Tol

$5_{REK}$ and its $CO_2$ conversion activity as a whole-cell biocatalyst. Fusing SyCA with the

SP of Tol 5 Omp38 led to efficient processing of its SP, resulting in the highest WAU

value (Figs. 2, 3). These results highlight that periplasmic expression and the selection

of an appropriate SP are important for the catalytic performance of carbonic anhydrase

as a whole-cell biocatalyst. To the best of our knowledge, this is the first report on the

heterologous expression of carbonic anhydrase in *Acinetobacter* for carbon utilization.

Several studies have explored the periplasmic expression of carbonic anhydrase

in *E. coli*. Jo *et al.* expressed NgCA using the PelB (Sec) and TorA (Tat) signal peptides

and found that the TorA construct yielded higher catalytic activity (Jo et al., 2013). Patel

*et al.* expressed the Cab and Cam with PelB (Sec) and gIII (Sec) signals, reporting that

PelB conferred slightly higher activity for both enzymes (Patel et al., 2013). More

recently, Jo *et al.* showed that the periplasmic expression of TdCA with the PelB (Sec)

signal peptide enhanced enzyme activity (Jo and Hwang, 2020). Collectively, these

findings indicate that the optimal signal peptide for periplasmic expression appears to

be specific to each carbonic anhydrase even in *E. coli*. In Tol 5, the *E. coli* PelB-SP did

not yield detectable SyCA production (Fig. 2), and the *E. coli* TorA-SP produced only

weak activity (Fig. 3). These results suggest that SPs from *E. coli* do not match the

secretion machinery of Tol 5 and are consistent with previous reports that variations in

signal peptide sequences influence secretion efficiency and that some signal peptides

exhibit species specificity (Blaudeck et al., 2001; Freudl, 2018). The three Tat-type SPs

from Tol 5 were not fully cleaved and also yielded lower activity (Figs. 2, 3). In

contrast, the native Sec-type Omp38-SP from Tol 5 was cleaved efficiently and gave the

best activity, demonstrating its suitability for expressing SyCA in this host.

Tol 5 cells can be immobilized on many materials through AtaA and can be

applied to gas-phase reactions (Yoshimoto et al., 2017; Usami et al., 2020). Combining

the bacterial cell immobilization with the periplasmic expression of SyCA may lead to

stable, reusable catalysts for gas-phase $CO_2$ hydration or for supplying bicarbonate to

downstream biosynthetic steps. Such systems could contribute to efficient biological

carbon capture and conversion technologies.

**Conclusion**

In this study, we demonstrated that periplasmic expression and selection of an

appropriate SP are important for the catalytic performance of carbonic anhydrase in Tol

5 as a whole-cell biocatalyst. Integrating the adhesive properties of Tol 5 with

engineered periplasmic expression of SyCA could lead to efficient $CO_2$ capture and

conversion processes.

**Author contribution**

**Shogo Yoshimoto**: Conceptualization, Validation, Writing - Original Draft. **Hiroya**

**Oka**: Investigation. **Yuki Ohara**: Investigation. **Yan-Yu Chen**: Investigation. **Masahito**

**Ishikawa**: Investigation. **Katsutoshi Hori**: Conceptualization, Writing - Review &

Editing, Funding Acquisition

**Acknowledgements and Funding information**

The authors thank Kensho Abe, Chiharu Kitano, and Eriko Kawamoto for their

technical assistance. An expression plasmid for SyCA used in preliminary experiments

was kindly provided by I-Son Ng (National Cheng Kung University). This research was

supported by the Japan Society for the Promotion of Science (JSPS) KAKENHI (Grant

Number JP24H00043) to KH and SY.

**Disclosure statement**

The authors declare no competing interests.

**Data availability**

All data generated or analyzed during this study are included in this published article

and its supplementary materials.

**References**

Baidya, P., Zhang, M., Xiao, Y.T., Zhang, H., Yu, L.J., and Li, W. (2024). Genetically
engineered whole-cell biocatalyst for efficient $CO_2$ capture by cell surface
display of carbonic anhydrase from *Bacillus cereus* GLRT202 on *Escherichia*
*coli*. *Biochem Eng J* 211, 109446.
Blaudeck, N., Sprenger, G.A., Freudl, R., and Wiegert, T. (2001). Specificity of signal
peptide recognition in Tat-dependent bacterial protein translocation. *J Bacteriol*
183(2), 604-610.
Capasso, C., De Luca, V., Carginale, V., Cannio, R., and Rossi, M. (2012). Biochemical
properties of a novel and highly thermostable bacterial α-carbonic anhydrase
from *Sulfurihydrogenibium yellowstonense* YO3AOP1. *J Enzym Inhib Med Ch*
27(6), 892-897.
Carceller, A., Guillén, M., and Alvaro, G. (2023). Lactic acid from $CO_2$: A carbon
capture and utilization strategy based on a biocatalytic approach. *Environ Sci*
*Technol* 57(51), 21727-21735.
Confer, A.W., and Ayalew, S. (2013). The OmpA family of proteins: Roles in bacterial
pathogenesis and immunity. *Vet Microbiol* 163(3-4), 207-222.

Freudl, R. (2018). Signal peptides for recombinant protein secretion in bacterial

expression systems. *Microb Cell Fact* 17, 52.

Hanifa, M., Agarwal, R., Sharma, U., Thapliyal, P.C., and Singh, L.P. (2023). A review

on $CO_2$ capture and sequestration in the construction industry: Emerging

approaches and commercialised technologies. *J CO_2 Util* 67, 102292.

Hori, K., Ohara, Y., Ishikawa, M., and Nakatani, H. (2015). Effectiveness of direct

immobilization of bacterial cells onto material surfaces using the

bacterionanofiber protein AtaA. *Appl Microbiol Biotechnol* 99(12), 5025-5032.

Inoue, S., Yoshimoto, S., and Hori, K. (2025). A new target of multiple lysine

methylation in bacteria. *J Bacteriol* 207(1), e00325-24.

Ishikawa, M., and Hori, K. (2024). The elimination of two restriction enzyme genes

allows for electroporation-based transformation and CRISPR-Cas9-based base

editing in the non-competent Gram-negative bacterium *Acinetobacter* sp. Tol 5.

*Appl Environ Microb* 90(6), e0040024.

Ishikawa, M., Nakatani, H., and Hori, K. (2012). AtaA, a new member of the trimeric

autotransporter adhesins from *Acinetobacter* sp. Tol 5 mediating high

adhesiveness to various abiotic surfaces. *PLoS One* 7(11), e48830.

Ishikawa, M., Shigemori, K., and Hori, K. (2014). Application of the adhesive

bacterionanofiber AtaA to a novel microbial immobilization method for the

production of indigo as a model chemical. *Biotechnol Bioeng* 111(1), 16-24.

Ishikawa, M., Yoshimoto, S., Hayashi, A., Kanie, J., and Hori, K. (2016). Discovery of a

novel periplasmic protein that forms a complex with a trimeric autotransporter

adhesin and peptidoglycan. *Mol Microbiol* 101(3), 394-410.

Jo, B.H., and Hwang, I.S. (2020). Characterization and high-level periplasmic

expression of thermostable α-carbonic anhydrase from *Thermosulfurimonas*

*dismutans* in *Escherichia coli* for $CO_2$ capture and utilization. *Int J Mol Sci*

21(1), 103.

Jo, B.H., Kim, I.G., Seo, J.H., Kang, D.G., and Cha, H.J. (2013). Engineered

*Escherichia coli* with periplasmic carbonic anhydrase as a biocatalyst for $CO_2$

sequestration. *Appl Environ Microb* 79(21), 6697-6705.

Kammerer, S., Borho, I., Jung, J., and Schmidt, M.S. (2023). Review: $CO_2$ capturing

methods of the last two decades. *Int J Environ Sci Technol* 20(7), 8087-8104.

Khalifah, R.G. (1971). The carbon dioxide hydration activity of carbonic anhydrase. I.

Stop-flow kinetic studies on the native human isoenzymes B and C. *J Biol Chem*

246(8), 2561-2573.

Liu, Z.H., Wang, K., Chen, Y., Tan, T.W., and Nielsen, J. (2020). Third-generation

biorefineries as the means to produce fuels and chemicals from $CO_2$. *Nat Catal* 3, 274-288.

Marra, L., Russo, M.E., Knuutila, H.K., Marzocchella, A., and Salatino, P. (2024). $CO_2$ capture and utilization by biocatalytic synthesis of carboxylic acids: a process design study. *Chem Eng Sci* 285, 119618.

Matos, C.F.R.O., Branston, S.D., Albiniak, A., Dhanoya, A., Freedman, R.B., Keshavarz-Moore, E., et al. (2012). High-yield export of a native heterologous protein to the periplasm by the tat translocation pathway in *Escherichia coli*. *Biotechnol Bioeng* 109(10), 2533-2542.

Mergulhao, F.J.M., Summers, D.K., and Monteiro, G.A. (2005). Recombinant protein secretion in *Escherichia coli*. *Biotechnol Adv* 23(3), 177-202.

Moon, H., Kim, S., Jo, B.H., and Cha, H.J. (2020). Immobilization of genetically engineered whole-cell biocatalysts with periplasmic carbonic anhydrase in polyurethane foam for enzymatic $CO_2$ capture and utilization. *J CO2 Util* 39, 101172.

Nagireddi, S., Agarwal, J.R., and Vedapuri, D. (2023). Carbon dioxide capture, utilization, and sequestration: Current status, challenges, and future prospects for global decarbonization. *ACS Eng Au* 4(1), 22-48.

Natale, P., Brüser, T., and Driessen, A.J.M. (2008). Sec- and Tat-mediated protein secretion across the bacterial cytoplasmic membrane - Distinct translocases and mechanisms. *BBA Biomemb* 1778(9), 1735-1756.

Patel, T.N., Park, A.H.A., and Banta, S. (2013). Periplasmic expression of carbonic anhydrase in *Escherichia coli*: A new biocatalyst for $CO_2$ hydration. *Biotechnol Bioeng* 110(7), 1865-1873.

Shao, P.J., Ye, J.X., Shen, Y., Zhang, S.H., and Zhao, J.K. (2024). Recent advancements in carbonic anhydrase for $CO_2$ capture: A mini review. *Gas Sci Eng* 123, 205237.

Sieborg, M.U., Nielsen, A.K.H., Ottosen, L.D.M., Daasbjerg, K., and Kofoed, M.V.W. (2024). Bio-integrated carbon capture and utilization: at the interface between capture chemistry and archaeal $CO_2$ reduction. *Nat Commun* 15, 7492.

Talekar, S., Jo, B.H., Dordick, J.S., and Kim, J. (2022). Carbonic anhydrase for $CO_2$ capture, conversion and utilization. *Curr Opin Biotech* 74, 230-240.

Teufel, F., Armenteros, J.J.A., Johansen, A.R., Gíslason, M.H., Pihl, S.I., Tsirigos, K.D., et al. (2022). SignalP 6.0 predicts all five types of signal peptides using protein language models. *Nat Biotechnol* 40, 1023-1025.

Ting, W.W., Yu, J.Y., Lin, Y.C., and Ng, I.S. (2022). Enhanced recombinant carbonic anhydrase in T7RNAP-equipped *Escherichia coli* W3110 for carbon capture

storage and utilization (CCSU). *Bioresour Technol* 363, 128010.

Usami, A., Ishikawa, M., and Hori, K. (2020). Gas-phase bioproduction of a high-value-

added monoterpenoid (*E*)-geranic acid by metabolically engineered

*Acinetobacter* sp. Tol 5. *Green Chem* 22(4), 1258-1268.

Wilbur, K.M., and Anderson, N.G. (1948). Electrometric and colorimetric determination

of carbonic anhydrase. *J Biol Chem* 176(1), 147-154.

Yoshimoto, S., Ohara, Y., Nakatani, H., and Hori, K. (2017). Reversible bacterial

immobilization based on the salt-dependent adhesion of the bacterionanofiber

protein AtaA. *Microb Cell Fact* 16(1), 123.

Yurak, V.V., and Fedorov, S.A. (2025). Review of natural and anthropogenic emissions

of carbon dioxide into the earth's atmosphere. *Int J Environ Sci Technol* 22(4),

2719-2736.

Zhu, Y.Z., Liu, Y.R., Ai, M.M., and Jia, X.Q. (2022). Surface display of carbonic

anhydrase on *Escherichia coli* for $CO_2$ capture and mineralization. *Synth Syst*

*Biotechnol* 7(1), 460-473.

**Figure captions**

**Figure 1. Cytoplasmic expression of SyCA.** (A) Western blotting of whole-cell lysates

from Tol 5$_{REK}$ cells carrying the SyCA expression plasmid (+) or carrying no plasmid

(–). Proteins were separated by SDS-PAGE, transferred to a PVDF membrane, and

probed with an anti-His antibody; the arrowhead marks the band corresponding to

SyCA. (B) Time course of the pH changes during the WAU assay. Carbonic anhydrase

activity of cells expressing cytoplasmic SyCA or blank (water only, no cells) was

measured.

**Figure 2. Expression of SyCA variants fused to each signal sequence.** (A) Schematic

representation of SyCA fused to a signal peptide (SP). (B) Protein expression was

detected by western blotting using an anti-His antibody. The arrowhead indicates the

band corresponding to the cytoplasmically expressed SyCA (Non-SP).

**Figure 3. Carbonic anhydrase activity of cells expressing SyCA variants fused to each signal sequence.** (A) Time course of the pH changes during the WAU assay of cells expressing SyCA variants fused to each signal sequence. (B) The WAU values calculated from three independent measurements. Data are expressed as mean ± SEM (n = 3). Statistical significance, *adjusted $p$ = 0.013, **adjusted $p$ < 0.01.

395 **Table 1. Signal peptides used in this study.**

| Source | Gene | Sec/Tat | Sequence |
|--------|------|---------|----------|
| *E. coli* | *pelB* | Sec | MKYLLPTAAAGLLLLAAQPAMA |
| *E. coli* | *torA* | Tat | MNNNDLFQASRRRFLAQLGGLTVAGMLGPSLLTPRRATA |
| Tol 5 | *omp38* | Sec | MKLSRIALAMLVAAPLAAANAG |
| Tol 5 | BCX72454 (Tat-1) | Tat | MTDLTPYHEDQELDNNTSNNIHFRDILEQRISRRSLITKTASGAAALALASSLTG |
| Tol 5 | BCX75163 (Tat-2) | Tat | MNKKVDFDPAKRDFLKTMGSLAAFGAVTTWMPVQSAKA |
| Tol 5 | BCX75681 (Tat-3) | Tat | MTNSANQHTRREVLTWLASIPFLPLGAMATSATLAG |