Effects of Extruder Dynamics and Noise on Simulated Chromatin Contact Probability Curves

Konstantinov V.[1,2*], Shadskiy A.[3,4], Lagunov T.[1,2]

[1] Institute of Cytology and Genetics, Novosibirsk, Russia
[2] Novosibirsk State University, Novosibirsk, Russia
[3] AXXX, Moscow, Russia
[4] Sirius University of Science and Technology, Sirius, Russia

* Corresponding author

E-mail: v.konstantinov.work@gmail.com

Abstract

Loop extrusion by SMC complexes is a key mechanism underlying chromatin folding during both interphase and mitosis. Despite this shared mechanism, computational models of loop extrusion often rely on fundamentally different assumptions: interphase models typically use dynamic extruders with finite lifetimes, whereas mitotic models employ static extruders placed according to loop size distributions. In this work, we investigate whether these modeling paradigms are interchangeable or yield intrinsically incompatible results. Using publicly available Hi-C data from mitotic chicken cells, we systematically compare dynamic and static loop extrusion models implemented in the Polychrom framework. We evaluate how key parameters such as the extruder lifetime, extrusion velocity, and spatial noise affect the simulated contact probability curves P(s) and loop size distributions. Our results reveal that while both model types can be tuned to approximate the general shape of P(s), they produce distinct internal structures and divergent relationships between loop size and contact decay. We also show that increased extruder lifetimes lead to excessive nested loop formation, which alters both loop statistics and P(s) derivatives. Introducing spatial exclusion constraints between extruders partially restores consistency with static models. These findings highlight that differences in extruder behavior and polymer noise levels can significantly impact chromatin model outcomes and must be carefully accounted for when interpreting or comparing simulation results across biological conditions.

Keywords: loop extrusion, chromatin contact maps, computational modeling, SMC complexes, Hi-C simulation, polymer dynamics

Author summary

Chromatin organization plays a crucial role in gene regulation and cellular function, yet our understanding of its three-dimensional structure relies heavily on computational modeling and the interpretation of complex experimental data. In this study, we use coarse-grained modeling approaches to simulate chromatin folding and systematically investigate how different analysis metrics and data processing methods influence the conclusions drawn from such models. By comparing widely used metrics and exploring the effects of normalization and noise, we highlight potential pitfalls and biases that can arise in chromatin modeling studies. Our findings provide practical recommendations for researchers in the field, aiming to improve the robustness and reproducibility of computational analyses of chromatin architecture. This work will help guide future studies toward more reliable interpretations of chromatin structure and its biological implications.

Background

In recent years, a range of chromosome conformation capture techniques have been developed to study the spatial organization of the genome (1–5). Among them, Hi-C has emerged as one of the most informative and widely used methods, providing genome-wide maps of chromatin contacts and enabling detailed insights into three-dimensional chromosome architecture (3). However, Hi-C data are subject to inherent limitations, including restriction-enzyme-dependent biases arising from the choice of restriction enzyme (e.g., DpnII used in this study), and modest genomic resolution (~1–10 kbp). These factors may influence the observed contact patterns

and thus potentially affect model calibration and interpretation. While other methods

such as Micro-C or approaches combined with ChIP-seq offer higher resolution and

capture higher-order interactions, the broad availability and established analytic

frameworks of Hi-C make it a reliable foundation for computational modeling (6–10).

One of the most prominent of these is the loop extrusion (LE) model, which posits

that ring-like proteins of the structural maintenance of chromosomes (SMC) family

assemble and load onto chromatin. Once loaded, these complexes actively

translocate in opposite directions along the DNA fibre, extruding loops of chromatin

in a directional manner (7,11). This leads to the emergence of loop organization in

chromatin. Current evidence suggests that this mechanism plays a critical role in

both interphase chromatin organization (7) and mitotic chromosome condensation

(8,12).

While the core biochemical mechanism of loop extrusion is thought to be shared

between interphase and mitosis (13,14), computational implementations of SMC

activity in these two contexts often differ substantially. For interphase chromatin, a

widely used model assumes dynamically loaded cohesin complexes that bind

randomly, extrude loops by sliding their legs in opposite directions, and dissociate

with a constant probability (7,15). These extruders are halted upon collision with

neighboring complexes or oriented CTCF proteins, forming an ensemble of loops

with stochastic sizes and limited lifetimes.

In contrast, mitotic chromosome models commonly adopt one of two approaches. A

popular static framework posits that condensins are sequentially loaded onto

chromatin, where they extrude loops until they are blocked by neighboring extruders,

after which they become fixed. In this scenario, condensin II first establishes a dense

array of long-range loops, followed by condensin I, which forms nested loops within

this scaffold (8,12). This hierarchical assembly captures the experimentally observed

loop sizes and density, and the model's validity is supported by a good match

between predicted and measured condensin occupancy per megabase (12).

Furthermore, the rapid stabilization of loop sizes observed in the experimental data

lends additional support to the static model. However, the static approach typically

requires imposing artificial constraints, such as cylindrical confinement and helical

axis generation, to recapitulate the global mitotic chromosome morphology (8,12).

Other modeling strategies attempt to circumvent the need for cylindrical confinement

by introducing additional physical features, such as increased interextruder affinity

(16) or hypothetical "superextruders" (17) capable of generating compact mitotic

structures through enhanced activity.

In contrast to the static approach, dynamic models of mitotic loop extrusion assume

that extruders bind and unbind chromatin with a fixed probability, analogous to their

behavior in interphase (11,18,19). Importantly, the residence time of these extruders

is considerably shorter than the duration of mitosis. This framework reaches a

dynamic equilibrium in which statistical properties—such as loop size distributions,

the number of nested loops, and other structural metrics—stabilize over time. To

reproduce mitotic chromosome patterns under this scheme, it is often assumed that

extruders are capable of bypassing one another, forming so-called "z-loops" (18,19) .

This assumption is motivated by in vitro observations (20) and experiments involving

bacterial condensins (21).

Despite this shared core mechanism, computational implementations of SMC

complexes differ substantially between interphase and mitosis, reflecting distinct

biological contexts and objectives. In interphase, dynamic loading and stochastic

dissociation of cohesin are motivated by the need to simulate steady-state chromatin

architecture that continuously remodels over hours. In contrast, mitotic

chromosomes must rapidly condense and remain stable throughout mitosis (~45 min

in mammalian cells), suggesting a quasistable loop architecture established by early-

loaded condensins that persist with minimal dynamics. Furthermore, in mitosis, the

high density of condensin loading and tight coordination with topological constraints

and chromosome geometry place different computational demands on the model.

Static extruder frameworks, which capture the final-state loop architecture, often

provide enough phenomenological descriptions of mitotic organization, even though

the underlying SMC mechanism remains fundamentally dynamic at the molecular

level.

In the present study, we sought to evaluate whether a dynamic extrusion model,

operating without z-loop formation, is sufficient to recapitulate mitotic chromosome

architecture under conditions of SMC3 and CAP-H2 subunit knockout. These

specific experimental conditions were selected because of the reported absence of

cylindrical chromosome morphology (12), thereby eliminating the need to impose

external constraints in the simulation.

One commonly analysed feature is the contact probability as a function of genomic

distance, P(s). This dependence is sensitive to both the polymer's fractal

dimensions—which characterize the self-similar, scale-invariant folding patterns

inherent to polymer physics and the specific compaction regimes and loop

architectures that emerge at different genomic scales depending on loop-extrusion

parameters. These two factors interact: whereas an unconstrained polymer exhibits

a universal fractal dimension (~1.5 for a linear random coil), the introduction of loop

extrusion modifies the effective fractal dimension at different scales, resulting in

scale-dependent compaction patterns. Thus, analysing P(s) provides a sensitive

readout of both fundamental polymer properties and extrusion-driven organization,

making it a powerful but nontrivial method for model validation (8,9,18).

In this study, we used publicly available data and baseline parameters from (12) to

assess how P(s) is influenced by the following:

- 136 ● the choice of modeling framework (we reimplemented the model via the
- 137   Polychrom toolkit as opposed to the original platform);
- 138 ● the type of SMC model (dynamic vs static extrusion);
- 139 ● the level of Gaussian noise added to the simulated contact maps.

Additionally, we compare several metrics for evaluating model-experiment

agreement and demonstrate the strong dependence on the normalization point used

in P(s) comparisons for most of them.

The focus of this study is a systematic evaluation of the reproducibility and

comparability of chromatin modeling results under varying assumptions and

implementations. Rather than limiting our investigation to a single mechanistic

hypothesis, we conducted a multilevel comparison across commonly used modeling

strategies.

First, we assessed the influence of different repulsive interaction potentials—

specifically, the Step (22) and DPD (23) potentials—on the simulation outcomes.

This comparison was performed using parameters closely aligned with those

reported in prior studies to isolate the effect of the potential itself.

Second, we contrasted two fundamentally distinct extrusion models under matched

conditions: a static model, in which loops are defined by a preassigned distribution,

and a dynamic model, in which extruders stochastically bind and dissociate from

chromatin.

Our results indicate that even when nearly equivalent input parameters are used, the

choice of modeling framework and software implementation (in this case, the

Polychrom toolkit versus the original platform) can substantially influence simulation

outcomes.

This highlights the critical importance of careful validation and reporting of all the

details of the computational model to ensure reproducibility in the field of

computational modeling of chromosome architecture.

Results

Used simulations and experimental data

In this study, we utilized experimental Hi-C data generated using the DpnII restriction

enzyme from mitotic chicken DT40 cell cultures, in which a degron system targets

subunits of both the condensin II and cohesin complexes (12; see Methods). Under

these conditions, the resulting contact probability curves P(s) lack statistically stable

genomic contacts beyond 2 Mbp, as more distant interactions are presumably

influenced by mechanisms unrelated to direct loop extrusion. As noted in the original

study, such experimental perturbation is interpreted to limit the effective range of

loop extrusion activity to approximately 2 Mbp.

Simulations were conducted via a modified version of the open2c/polychrom

software package (22), which supports modeling chromosome architecture via

dynamic loading and unloading of extruders. The extruder parameters were selected

based on those reported in (12). As in that study, we did not use cylindrical

confinement or include compartmentalization effects in our simulations. However, in

contrast to that study, we did not impose any length constraints on the

chromosomes, allowing the simulated chromosomes to reside in a diluted regime

without external geometric restrictions.

Baseline Outcomes Depend on Potential and Implementation

181
182

We began by evaluating the reproducibility of the experimental data via a model of a

40 Mbp chromosome with statically placed condensin I motors. The extruders were

distributed according to an exponential loop size distribution with a mean of 100 kbp

and an interextruder distance of 400 bp, parameters previously suggested as optimal

for recapitulating contact density profiles in experiments (12). To assess the

consistency of modeling results across different implementations, we employed two

parameter sets and interaction potentials within the Polychrom framework: the Step

potential, used as the default in dynamic extrusion models (24), and the DPD

potential, utilized in (12,25). The Step potential characterizes beads as nearly 'hard

spheres' with a steep potential gradient at the bead radius, minimizing overlaps

outside the interaction zone. Conversely, the DPD potential models 'softer' beads

with a smoothly distributed potential gradient over the interaction range. These two

potentials are representative of typical 'hard' and 'soft' coarse-grained interactions,

and are broadly comparable to other potentials used in polymer modeling. Further

details are provided in the Methods section.

The resulting contact probability curves (Fig 1A) revealed discrepancies between the

simulated contact distributions (as a function of genomic distance) and the

experimental data for both potentials. These differences can be summarized as

follows:

● Poor agreement was observed in the short-range contact region (up to 10

kbp), suggesting a mismatch between the chosen interaction potentials and

the underlying physical parameters (see Methods).

● A generally reduced contact density was noted across the 3 kbp to 2 Mbp

range.

● The fluctuations in the derivative of the contact probability are consistent with

the stochastic noise typical of complex molecular systems under limited

sampling.

● The Step potential produced results more consistent with the experimental

data than the DPD potential, primarily on the basis of closer visual agreement

of the simulated P(s) curves with the experimental profiles and superior

performance according to quantitative metrics assessing differences between

simulated and experimental contact probability distributions (see Methods).

Fig 1. Comparison of experimental and simulated contact probability P(s)

profiles.

A) P(s) curves in the range of 3 kbp - 2 Mbp for experimental data for the chicken

cell line DT40 [(12)] (black line), raw model single cells (transparent lines),

aggregated raw model data (solid lines). Static extruders with exponential distribution

of loop sizes: with Step potential (red line) and DPD potential (blue line); B) Results

of comparing segments experiment P(s) vs model data with shifting normalisation

points on different metrics. The normalisation point shifts within the range of 3 kbp to

2 Mbp. The comparable segment of the plot is 3 kbp-2 Mbp.

Although both models using Step and DPD potentials generally reproduce the shape

of the experimental P(s) curve, significant differences in detail, especially in the

short-range range and the overall level of contact density, point to the limitations of

the parameters and potentials used. These differences highlight the need for careful

selection of physical parameters and normalization approaches to achieve better

agreement between simulation and experiment.

Quantitative Agreement Metrics are Sensitive to Normalization Point in Model Fits

In addition to direct visual comparison, we employed several quantitative metrics to

assess the similarity between the simulated and experimental contact probability

curves. However, a key methodological issue arises: P(s) curves are relative by

nature and require normalization at a chosen genomic distance. For example, in

(12), normalization was performed at 3 kbp, and comparisons were made over the

range of 3 kbp to 2 Mbp. In contrast, (18) applies normalization at 10 kbp or, in some

cases, 100 kbp to emphasize longer-range contacts.

These differences in normalization introduce significant variability in commonly used

metrics, including the coefficient of determination (R^2) for P(s), root mean square

error (RMSE), mean absolute error (MAE), maximum error (ME), and average ratio.

In contrast, R^2 calculated for the derivative of P(s) remains unaffected by

normalization since the derivatives are independent of the absolute scaling of P(s)

(Fig 1B).

We observed that the R², RMSE, and MAE values generally increase with

normalization points approaching ~100 kbp, reaching a maximum near 1 Mbp. The

ME metric exhibits a similar trend, although its maximum occurs at approximately

100 kbp. Beyond these points, all the metrics based on P(s) begin to decline in

quality, especially as the analysis approaches the 2 Mbp upper boundary. In

contrast, the average ratio metric peaks at 20–30 kbp and shows a pronounced

decrease beyond 1 Mbp.

This analysis was repeated across all simulation conditions presented in this study

(see Appendix). For consistency, we selected 3 kbp as the normalization point for

determining the best-fit parameters throughout the paper, aligning with the

normalization strategy used in (12).

It is important to define and consistently apply a normalization point when analysing

P(s) curves, as variations in this choice can substantially affect the values of

similarity metrics and hinder direct comparisons across different studies. Moreover,

in certain cases, the choice of normalization point can even influence which model

appears to provide a better fit to the experimental data (S1-S2 Figs). Fixing the

normalization point throughout the analysis enhances the robustness and

reproducibility of model benchmarking. Unlike metrics based on the absolute values

of P(s), this derivative-based R² is independent of overall scaling and normalization,

providing a more rigorous and unbiased assessment of the model's ability to

reproduce the shape of contact probability profiles. Furthermore, this metric

quantifies how well the model approximates the experimental derivative relative to

the average slope observed in the experimental data over the comparison range.

Since the average slope of the P(s) derivative is directly linked to the underlying

fractal dimension of the polymer backbone in the absence of extrusion-driven loops,

the derivative-based R^2 can be regarded as a physically grounded and informative

measure of model fidelity.

Dynamic Behavior of Condensin I is Inconsistent with Experimental Signatures

The use of static extruders with an exponential loop size distribution raises important

questions regarding their validity for modeling the dynamic behavior of SMC

complexes. In (12), static extruders are employed in conjunction with externally

imposed large-scale helical constraints, which promote chromosome compaction. In

contrast, (18) demonstrated that a similar helical structure can emerge

spontaneously in simulations with dynamic extruders, even in the absence of

external spatial constraints. In our simulations, however, we did not observe the

formation of helical structures under either static or dynamic extrusion regimes (see

Appendix), in contrast to the results reported by (18) for dynamic models.

A related consideration is the choice of parameters for modeling dynamic extruders

(mitotic condensins). Loop size can be limited either by the intrinsic processivity of

condensin or by natural roadblocks (for example, other extruders, cohesin-specific

barriers such as CTCF, transcription, and related obstacles). Notably, condensin has

been reported to bypass even large DNA-bound obstacles (21). To date, there are

no well-established strong barriers to condensin motion along DNA, and

transcriptional activity during mitosis is minimal (26). Within this context, the key

parameter is the extruder lifetime (the mean residence time on DNA), which,

together with the extruder loading density, directly determines the characteristic loop

size . We estimated the baseline lifetime via the following relation:

lt = {*average loop size*} / ({*velocity on naked DNA*} x 6)   (3)

where the average loop size for condensin I is taken to be 100 kbp, the extrusion

velocity is set to 1 kbp/sec (27), and the 1/6 factor accounts for the effective scaling

of the polymer modeled as a chain of 10 nm nucleosome beads. Since the beads are

10 nm, which corresponds to 30 bp of naked DNA, and each nucleosome accounts

for approximately 200 bp of naked DNA, and since the extruder makes spatial steps,

the effective step distance is approximately 200/30 = ~6 times greater than that of

naked DNA, the velocity on chromatin is 6 times greater than that on naked DNA.

This yields a baseline lifetime of approximately 17 seconds, which is sufficient for

~63% of extruders to reach the next adjacent extruder along the polymer. In the

simulations, each condensin complex has a probability of 1/lt to detach from the

polymer at each time step and is immediately reloaded at a random location. SMC

complexes perform this translocation through discrete spatial steps: recent single-

molecule studies have demonstrated that condensin and cohesin translocate DNA in

steps with a median size of ~40 nm on naked DNA (28,29), with each step

accompanied by a DNA twist of approximately −0.6 rotations (30). This mechanistic

foundation (discrete spatial stepping coupled with DNA wrapping) supports our

coarse-grained representation of extrusion as a sequence of bead-displacement

events in our simulations.

To explore the effect of extruder dynamics, we simulated three different lifetimes:

● **17 seconds**, corresponding to the time required for condensin I to extrude its

average loop size (~100 kbp);

● **68 seconds**, corresponding to the time required for condensin II to extrude its

average loop size (~400 kbp);

● **136 seconds**, which was used as a control to investigate the effects of

prolonged binding.

As shown in Fig 2, we observe marked differences in the contact density profiles

between the dynamic and static extrusion regimes. For extruders with a 17-second

lifetime, the contact density is elevated relative to the experimental data at distances

in the 10–100 kbp range, followed by a steep decline. This reflects the short binding

duration, during which many extruders fail to encounter obstacles before detaching;

approximately one-third dissociate within 17 seconds.

Fig 2. Effect of extruder lifetime on contact probability P(s) profiles.

A) P(s) curves in the range of 3 kbp - 2 Mbp for experimental data for the chicken

cell line DT40 (black line), aggregated model data. Static (gray line) and dynamic

extruders with three lifetimes and base physical parameters (red, blue, purple lines).

The large dots in the derivative graphs correspond to the mean loop size; B) Loop

size distributions for different lifetimes. All the graphs are normalised on the area

under the all-loops graph. The purple line represents all extruders, the red line

represents unnested extruders, and the blue line represents nested extruders. 1)

Static extruders, 2) Dynamic extruders lt = 17 sec, 3) Dynamic extruders lt = 68 sec,

4) Dynamic extruders lt = 136 sec. Black line - corresponding derivative peak size.

For the 68-second condition, the number of short-range contacts (<40 kbp) is

reduced relative to the 17-second case because fewer short-lived extruders exist.

However, the contact density increases significantly in the 50–200 kbp range,

aligning more closely with the experimental data in this interval. Beyond this range,

the model again overestimates contact decay.

The 136-second extruders exhibit similar behavior up to ~20 kbp, but the contact

density is slightly lower than that in the 68-second case up to ~70 kbp. However,

between 70 and 600 kbp, the contact probability profile closely matches the

experimental curve. Beyond 1 Mbp, discrepancies reemerged across all conditions,

which is consistent with the finite influence range of loop extrusion observed in the

experimental system.

To further investigate the impact of extruder lifetimes on the three-dimensional

organization of chromatin, we analysed the resulting distributions of loop sizes under

different dynamic conditions. This analysis explicitly considers the formation of

nested loops, which occur when new loops are extruded within preexisting loops.

Importantly, that z-loop formation (21) does not occur in our model, as extruders are

not permitted to pass through one another.

For extruders with a 17-second lifetime, the loop size distribution follows an

approximately exponential profile (Fig 2B), closely resembling that observed for

static extruders. In this regime, nonnested loops significantly outnumber nested

loops. As the extruder lifetime increases, the average loop size increases

accordingly. Because the total number of extruders on chromatin remains constant,

the probability of new extruders binding within existing loops increases, leading to an

increase in the number of nested loops. With further increases in lifetime (e.g., 136

seconds), the ratio of nested to nonnested loops continues to rise, indicating

progressive structural complexity.

Previous studies have established a correlation between the average loop size and

the position of the peak in the derivative of the P(s) curve (7,31,32). While our

findings support the overall trend — namely, that larger average loop sizes are

associated with a rightward shift of the derivative peak — we also observe

quantitative discrepancies between these two measures. Specifically, under the 136-

second lifetime condition (Fig 2), the position of the derivative peak substantially

exceeds the corresponding average loop size.

A similar discrepancy is observed in the case of static extruders when Gaussian

noise exceeds 50 nm (S1 Appendix). Although the loop size is fixed at 100 kbp, the

derivative peak shifts from ~90 kbp to nearly 300 kbp as the Gaussian noise

increases (S1 Appendix). These findings suggest that while the derivative of P(s) is a

useful qualitative indicator of loop size, it may be quantitatively unreliable under

certain conditions, such as high noise levels or extensive nested loop formation.

Accordingly, the position of the derivative peak should be treated only as a first-order

approximation of the extruder-imposed loop size in the experimental data. When

greater precision is needed, a targeted parameter sweep in the vicinity of the

derivative peak is recommended to refine the estimate.

As shown in Fig 2, increasing the extruder lifetime leads to a marked rise in the

number of nested loops. This effect is associated with a reduction in the effective

number of active extruders, resulting from spatial overlap and the hierarchical nature

of nested loop organization. To address this limitation, we implemented an

alternative strategy that introduced spatial exclusion rules, whereby condensin

complexes are restricted to loading only within interloop regions, and not within

existing loops (Fig 3).

Fig 3. Impact of constrained extruder dynamics on contact probability P(s)

profiles across lifetimes.

P(s) curves in the range of 3 kbp - 2 Mbp for experimental data for the chicken cells

line DT40 (black line) and aggregated model data. The large dots in the derivative

graphs correspond to the mean loop size.

Under this modified framework, the loop size distribution for extruders with a 17-

second lifetime remained unchanged, in agreement with theoretical expectations.

However, at longer lifetimes, the distributions shifted substantially, becoming more

similar to those observed in the static extruder configuration, as reflected in both the

P(s) curves (Fig 3) and loop size distributions (S3 Fig).

The rationale for this exclusion model stems from experimental observations that

condensins can induce chromatin twisting or torsional stress (30), which may affect

the physical properties of the substrate and reduce the likelihood of additional

extruders binding to already extruded loops. It is therefore theoretically plausible that

torsional tension contributes to a reduced probability of condensin loading in regions

already occupied by other extruders.

None of the tested models outperformed the static extruder model in terms of

quantitative agreement with the experimental data. Specifically, the average ratio

metric for the static model with optimal Gaussian noise (σ = 200 nm, see Methods)

was 1.16. For the DPD potential model with its best-performing noise level (σ = 100

nm), the value was 1.23 (S1 Appendix). The dynamic extruder model with its optimal

lifetime (136 sec) yielded a value of 1.37 (S1 Fig), and the dynamic model with

restricted loading (i.e., extruders prohibited from binding within existing loops)

achieved a value of 1.35 (S2 Fig) under the optimal lifetime (136 sec).

A similar trend was observed for the derivative R^2 metric, which reflects the

agreement in slope between the modeled and experimental P(s) curves. The static

model with σ = 200 nm achieved a derivative R^2 value of 0.98; the DPD model with σ

= 100 nm produced 0.97; the unrestricted dynamic extruder model with the optimal

lifetime (136 sec) yielded 0.72; and the restricted dynamic model yielded 0.77 under

the optimal lifetime (136 sec).

Discussion and conclusions

In the present study, we systematically compared static and dynamic loop extrusion

models for modeling mitotic chromosome folding via chicken Hi-C data. Our work

aimed to clarify whether these two approaches are interchangeable or whether they

lead to fundamentally different results.

We showed that although both models can be tuned to approximate the general

shape of the contact probability curve P(s), the static extruder model demonstrated

the best quantitative agreement with the experimental data compared with all the

tested dynamic models. In particular, the static model with an optimally chosen noise

level (σ = 200 nm) and interaction potential Step achieved the best performance on

metrics such as the average ratio and coefficient of determination (R^2) for the

derivative of P(s). However, given the breadth of the hyperparameter space

(including extruder abundance, translocation velocity, and lifetime; bead–bead

interaction energies; the presence or absence of PBS boxes; and the level of

Gaussian smoothing (σ)), it remains plausible that an alternative interaction potential

could yield a superior approximation to the experimental data under a different, yet

biologically reasonable, parameterization.

Our results highlight key limitations of dynamic models in this context. The increase

in extruder lifetime required to reproduce contacts at large genomic distances

resulted in the formation of an excessive number of nested loops. This, in turn,

distorted both the loop size distribution and the P(s) curve. In addition, we found that

under high noise conditions or with many of nested loops, the position of the peak of

the P(s) derivative ceases to be a reliable indicator of the average loop size.

We qualitatively and quantitatively compared these two modeling paradigms within a

single platform (Polychrom), which allowed us to identify their intrinsic differences.

We also demonstrated a strong dependence of the results on the choice of

parameters such as the normalization point of the P(s) curves and the level of spatial

noise, and proposed using the R^2 of the P(s) derivative as a more reliable metric for

comparing the models.

The novelty of our study is that we perform, for the first time, a direct comparison of

static and dynamic models of extrusion within a single software framework using

parameters from a recent work.

Thus, our study confirms that models with static extruders, despite their apparent

simplicity, remain powerful and adequate tools for describing the final structure of

mitotic chromosomes. Across the conditions examined, the static model of the

mitotic chromosome provided the closest agreement with the experimental data. In

contrast, dynamic extrusion models in which extruders stall upon collision

consistently yield poorer fits. Attaining comparable performance with a dynamic

framework requires extruder lifetimes on the order of the mitotic phase, which

effectively produces loop size distributions indistinguishable from those of the static

model. Even with an added spatial-exclusion rule that prohibits loading within

preexisting loops, satisfactory agreement still necessitates long lifetimes, again

rendering the behavior functionally equivalent to the static configuration.

An alternative model assumes that condensins bind DNA once at mitotic entry,

extrude loops until they encounter other complexes, and remain stably associated

until the end of mitosis. However, this model would require the introduction of

additional mechanisms. For example, the experimentally observed nonzero gaps

between loops (12) could be explained by stalling forces arising from chromatin

tension exceeding ~2 pN. Similarly, the progressive compaction of chromosomes

and the upwards shift in the second diagonal of contact maps, despite relatively

stable loop sizes for condensin II (12), may reflect the influence of loop extrusion–

independent mechanisms, such as the activity of topoisomerase II (10).

Methods

Polymer Models

We constructed polymer models of chromosomes via polymer dynamics simulations.

The simulations were performed via a package based on open2c/polychrom

(Imakaev et al., 2019), which integrates parameters from both static and dynamic

simulations as described in Samejima et al. (2025).

The simulation workflow consists of two main stages:

1.   One-dimensional (1D) simulation

The polymer chain is represented as a 1D lattice composed of beads, each 10

nm in size and corresponding to 200 base pairs (bp).

1.1.   Dynamic extruders, representing condensin I complexes, are modeled

481   as walker objects with two legs that move probabilistically in opposite

482   directions. At each time step, only one leg is active and attempts to

483   move up to approximately 50 nm (5 beads) in its respective direction.

484   The active leg switches with a 50% probability, effectively simulating

485   two-sided loop extrusion. The positions of the legs define the loop

486   boundaries extruded by condensins.

487   1.2.   Condensins bind randomly to pairs of adjacent free lattice sites and

488   extrude loops until unbinding occurs, after which they immediately

489   rebind at random locations. When two legs of different extruders

490   collide, extrusion stalls until one leg unbinds.

491   1.3.   Static extruders are also modeled as walker objects, but with legs

492   loaded onto lattice sites following an exponential distribution with a

493   mean loop size of 100 kbp. These static extruders do not perform

494   extrusion or unbinding.

495   1.4.   The condensins identified in the 1D simulation are subsequently

496   implemented as additional harmonic constraints in the 3D polymer

497   model.

498   1.5.   The simulation time step is 0.1 seconds. The condensin I velocity is set

499   to 1 kbp/s on naked DNA (27), corresponding to a step size of 50 nm

500   (28); this translates to approximately 6 kbp/s on condensed chromatin.

501   The total simulation duration is 1800 seconds.

502   1.6.   All 1D modeling was performed via our scripts

503   (https://github.com/Three-point/1D-polymer-models)

504 2.   Three-dimensional (3D) simulation

505   This stage employs molecular dynamics simulations using normalized units,

506   where particles have a mass of 1 unit and a bead radius of 0.5 units, with one

507   length unit corresponding to 10 nm.

508   2.1.   The monomers are connected by harmonic springs

509   (polychrom.forces.harmonic_bonds) with a bond length of 1 unit and a

510   wiggle distance of 0.1 units.

511   2.2.   The polymer stiffness (polychrom.forces.angle_force) is modeled with a

512   persistence length of 150 bp, equivalent to three nucleosome linkers of

513   50 bp each, implemented via an angular force with k=3.

514   2.3.   Repulsive forces prevent spatial overlap via polynomial repulsion

515   (polychrom.forces.polynomial_repulsive) with Erep = 3.0, mimicking

516   real polymer behavior (Formula 1). The collision rate is set to 0.01

517   inverse picoseconds, and the Langevin integrator is used with an error

518   tolerance of 0.01. Each time step comprises 1000 molecular dynamics

519   steps.

520   2.4.   No external spatial constraints are applied to the polymer in this work.

521   This differs from (12), where cylindrical and pulling constraints were

522   used to achieve the required densities and patterns.

Each dataset comprises five independent 40 Mbp chains, each initialized with

distinct condensin positions and starting conformations generated by random walks.

Two different potential energies were used for nonbonded bead repulsion.

● "Step potential" (22):

$U\_12 = R^{12}*(R^2-1)*E/m + E$ (2)

● "DPD potential" (23):

$Udpd = (R-1)^2*E/2$ (3)

where R is the distance between beads in dimensionless units, E is the repulsion

energy coefficient, and m is the normalisation constant for this degree of equation.

Experiment types

There 11 types of model parameters were used:

● Static extruders. The positions of the extruders placed on the lattice with an

exponential distribution of loops once do not change with time (no move,

binding/unbinding):

○ Step potential

○ DPD potential (Groot and Warren, 1997, Formula 2), where Erep = 3.0

kT, k = 0 (polymer stiffness), and the other parameters are equal to the

Step potential.

● Dynamic extruders. All the physical parameters are taken from the Step

potential. They unbind from the lattice with a probability inversely proportional

to the lifetime and then instantly bind back at random site. Then, they make

one step by one of the legs chosen randomly (50% chance to switch direction)

on 5 beads each time step with a probability sufficient for the target extrusion

speed:

○ Lifetime = 17 sec for each extruder. This is enough for ~60% of the

extruders to extrude 100 kbp (if no obstacles are encountered), forming

a distribution closest to exponential. There are low numbers of nested

loops (loops on other loops) and a dominating number of unnested

loops.

○   Lifetime = 68 sec for each extruder. Nested loops started dominating in

numbers at short distances.

○   Lifetime = 136 sec for each extruder. Nested loops are dominant in

terms of number for all loop sizes.

● Dynamic extruders that are unable to load at already existing loops, they load

only at gaps between loops.

○   Lifetime = 17 sec for each extruder.

○   Lifetime = 68 sec for each extruder.

○   Lifetime = 136 sec for each extruder.

561 **Generation of Hi-C Maps**

For each polymer chain in the simulations, we extracted conformations represented

by the spatial coordinates of beads at the following time points: 1080 s, 1440 s, and

1800 s. This selection is based on the assumption that a 360 s interval (exceeding

three condensin lifetimes) suffices to establish a new loop distribution under dynamic

conditions.

Each bead position within a conformation was subsequently perturbed by adding a

random Gaussian displacement with standard deviation σ along each spatial axis. By

repeating this perturbation process, we generated 100 new conformations from each

original conformation, resulting in a total of 5×3×100=1500 conformations per

dataset.

For each conformation, pairs of 200 bp beads located within 11 nm (contact radius,

1.1 of the bead radius) were identified as contacts, from which a contact map in

.mcool format at 200 bp resolution was constructed (https://github.com/Three-

point/1D-polymer-models). Unless otherwise specified, the default value of σ was set

to 200 nm, as this value yielded the best agreement with the experimental P(s) curve

for static extruder conditions (see S1 Appendix).

Processed Hi-C data (.mcool) from the DT40 cell line of Gallus gallus

(https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE262525), sample

GSM8171168, were used as experimental data.

The P(s) curves were computed via the cooltools (version 0.5.1) and cooler (version

0.9.1) packages with curve Gaussian smoothing (σ=0.12) and excluding the first two

diagonals of the Hi-C contact matrix.

Quantitative Comparison of P(s) Curves

To quantitatively compare the contact probability curves P(s), we focused on the

genomic distance range from 3 kbp to 2 Mbp, which encompasses the effective

range of condensin I activity (typical loop size ~100 kbp). Prior to metric evaluation,

all curves were normalized at a selected genomic coordinate within this range. Given

the logarithmic scaling of the genomic distance axis, discrepancies at larger

distances exert a disproportionately strong influence on agreement metrics.

Therefore, the analysed range was subdivided into 100 uniformly spaced intervals on

a log scale.

The following metrics were used to assess the agreement between experimental and

simulated P(s) curves in log10–log10 space:

- **Coefficient of determination (R^2):** Values approaching 1 indicate strong agreement; negative values suggest that the model performs worse than a horizontal mean line.
- **Root Mean Square Error (RMSE):** Lower values denote better fit and reduced variance from the experimental curve.
- **Mean Absolute Error (MAE):** Measures the average magnitude of differences; lower values indicate better agreement.
- **Maximum Error (ME):** Captures the largest single deviation between curves; lower values are preferred.
- **Average Ratio:** At each point, the modeled P(s) value is divided by the corresponding experimental value. If the ratio is <1, its reciprocal is taken to ensure symmetry. The final value is the arithmetic mean of these normalized quotients across all distances. A value close to 1 reflects excellent fit, whereas larger values indicate greater overall deviation.

To compare the derivatives of the P(s) curves, only the **R^2 metric** was employed. This choice is justified by its invariance to normalization, as it captures the similarity in slope and shape of the curves, independent of absolute scaling.

Loop Distribution Analysis

Loop positions were extracted at identical time points (1080 s, 1440 s, and 1800 s) for each simulated cell and categorized into the following groups:

- **Nested loops**: loops fully enclosed within the genomic span of another loop.
- **Unnested loops**: loops that are not nested within any other loop.
- **All loops**: the complete set of loops detected, regardless of nesting status.

618   For each extruder, the genomic span—defined as the distance between its two

619   extrusion legs in base pairs was calculated. The resulting distributions of loop sizes

620   for each category were visualized via Kernel Density Estimation (KDE) plots,

621   generated with the Seaborn Python library.

References

1. Chromosome Conformation Capture Carbon Copy (5C): A massively parallel
solution for mapping interactions between genomic elements [Internet]. [cited
2025 May 29]. Available from: https://genome.cshlp.org/content/16/10/1299

2. Dekker J, Rippe K, Dekker M, Kleckner N. Capturing Chromosome
Conformation. Science [Internet]. 2002 Feb 15 [cited 2025 May 29]; Available
from: https://www.science.org/doi/10.1126/science.1067799

3. Lieberman-Aiden E, van Berkum NL, Williams L, Imakaev M, Ragoczy T, Telling
A, et al. Comprehensive mapping of long range interactions reveals folding
principles of the human genome. Science. 2009 Oct 9;326(5950):289–93.

4. Simonis M, Klous P, Splinter E, Moshkin Y, Willemsen R, de Wit E, et al. Nuclear
organization of active and inactive chromatin domains uncovered by
chromosome conformation capture–on-chip (4C). Nat Genet. 2006
Nov;38(11):1348–54.

5. Zhao Z, Tavoosidana G, Sjölinder M, Göndör A, Mariano P, Wang S, et al.
Circular chromosome conformation capture (4C) uncovers extensive networks of
epigenetically regulated intra- and interchromosomal interactions. Nat Genet.
2006 Nov;38(11):1341–7.

6. Banigan EJ, Tang W, van den Berg AA, Stocsits RR, Wutz G, Brandão HB, et al.
Transcription shapes 3D chromatin organization by interacting with loop
extrusion. Proceedings of the National Academy of Sciences. 2023 Mar
14;120(11):e2210480120.

7. Fudenberg G, Imakaev M, Lu C, Goloborodko A, Abdennur N, Mirny LA.
Formation of Chromosomal Domains by Loop Extrusion. Cell Reports. 2016 May
31;15(9):2038–49.

8. Gibcus JH, Samejima K, Goloborodko A, Samejima I, Naumova N, Nuebler J, et
al. A pathway for mitotic chromosome formation. Science. 2018 Feb
9;359(6376):eaao6135.

9. Nuebler J, Fudenberg G, Imakaev M, Abdennur N, Mirny LA. Chromatin
organization by an interplay of loop extrusion and compartmental segregation.
Proceedings of the National Academy of Sciences. 2018 July 17;115(29):E6697–
706.

10. Hildebrand EM, Polovnikov K, Dekker B, Liu Y, Lafontaine DL, Fox AN, et al.
Mitotic chromosomes are self-entangled and disentangle through a
topoisomerase-II-dependent two-stage exit from mitosis. Molecular Cell. 2024
Apr 18;84(8):1422-1441.e14.

11. Banigan EJ, Van Den Berg AA, Brandão HB, Marko JF, Mirny LA. Chromosome

organization by one-sided and two-sided loop extrusion. eLife. 2020 Apr
6;9:e53558.

12. Samejima K, Gibcus JH, Abraham S, Cisneros-Soberanis F, Samejima I, Beckett
AJ, et al. Rules of engagement for condensins and cohesins guide mitotic
chromosome formation. Science [Internet]. 2025 Apr 11 [cited 2025 May 12];
Available from: https://www.science.org/doi/10.1126/science.adq1709

13. Kong M, Cutts EE, Pan D, Beuron F, Kaliyappan T, Xue C, et al. Human
Condensin I and II Drive Extensive ATP-Dependent Compaction of Nucleosome-
Bound DNA. Molecular Cell. 2020 July 2;79(1):99-114.e9.

14. Pradhan B, Barth R, Kim E, Davidson IF, Bauer B, Van Laar T, et al. SMC
complexes can traverse physical roadblocks bigger than their ring size. Cell
Reports. 2022 Oct;41(3):111491.

15. Mach P, Kos PI, Zhan Y, Cramard J, Gaudin S, Tünnermann J, et al. Cohesin
and CTCF control the dynamics of chromosome folding. Nat Genet. 2022
Dec;54(12):1907–18.

16. Forte G, Boteva L, Gilbert N, Cook PR, Marenduzzo D. Bridging-mediated
compaction of mitotic chromosomes. Nucleus. 2025 Dec 31;16(1):2497765.

17. Norouzi D. A Super-Loop Extrusion Mechanism Shapes the 3D Mitotic
Chromosome Folding [Internet]. bioRxiv; 2025 [cited 2025 Oct 13]. p.
2025.05.05.652288. Available from:
https://www.biorxiv.org/content/10.1101/2025.05.05.652288v1

18. Dey A, Shi G, Takaki R, Thirumalai D. Structural changes in chromosomes
driven by multiple condensin motors during mitosis. Cell Reports. 2023 Apr
25;42(4):112348.

19. Beckwith KS, Brunner A, Morero NR, Jungmann R, Ellenberg J. Nanoscale DNA
tracing reveals the self-organization mechanism of mitotic chromosomes. Cell.
2025 May 15;188(10):2656-2669.e17.

20. Kim E, Kerssemakers J, Shaltiel IA, Haering CH, Dekker C. DNA-loop extruding
condensin complexes can traverse one another. Nature. 2020
Mar;579(7799):438–42.

21. Brandão HB, Ren Z, Karaboja X, Mirny LA, Wang X. DNA-loop extruding SMC
complexes can traverse one another in vivo. Nat Struct Mol Biol. 2021
Aug;28(8):642–51.

22. Imakaev M, Goloborodko A, hbbrandao. mirnylab/polychrom: v0.1.0 [Internet].
Zenodo; 2019 [cited 2024 Nov 29]. Available from:
https://zenodo.org/records/3579473

23. Groot RD, Warren PB. Dissipative particle dynamics: Bridging the gap between
atomistic and mesoscopic simulation. The Journal of Chemical Physics. 1997
Sept 15;107(11):4423–35.

24. Imakaev MV, Fudenberg G, Mirny LA. Modeling chromosomes: beyond pretty
pictures. FEBS Lett. 2015 Oct 7;589(20 0 0):3031–6.

25. Anderson JA, Glaser J, Glotzer SC. HOOMD-blue: A Python package for high-
performance molecular dynamics and hard particle Monte Carlo simulations.
Computational Materials Science. 2020 Feb 15;173:109363.

26. Contreras A, Perea-Resa C. Transcriptional repression across mitosis:
mechanisms and functions. Biochem Soc Trans. 2024 Feb 19;52(1):455–64.

27. Tortora MMC, Fudenberg G. The physical chemistry of interphase loop extrusion
[Internet]. bioRxiv; 2024 [cited 2025 May 29]. p. 2024.08.23.609419. Available
from: https://www.biorxiv.org/content/10.1101/2024.08.23.609419v1

28. Ryu JK, Rah SH, Janissen R, Kerssemakers JWJ, Bonato A, Michieletto D, et al.
Condensin extrudes DNA loops in steps up to hundreds of base pairs that are
generated by ATP binding events. Nucleic Acids Res. 2022 Jan 25;50(2):820–32.

29. Kaur P, Lu X, Xu Q, Irvin EM, Pappas C, Zhang H, et al. High-speed AFM
imaging reveals DNA capture and loop extrusion dynamics by cohesin-NIPBL.
Journal of Biological Chemistry [Internet]. 2023 Nov 1 [cited 2025 Nov
26];299(11). Available from: https://www.jbc.org/article/S0021-9258(23)02324-
4/abstract

30. Janissen R, Barth R, Davidson IF, Peters JM, Dekker C. All eukaryotic SMC
proteins induce a twist of –0.6 at each DNA loop extrusion step. Science
Advances [Internet]. 2024 Dec 13 [cited 2025 June 2]; Available from:
https://www.science.org/doi/10.1126/sciadv.adt1832

31. Kim J, Jimenez DS, Ragipani B, Zhang B, Street LA, Kramer M, et al. Condensin
DC loads and spreads from recruitment sites to create loop-anchored TADs in C.
elegans. Tyler JK, editor. eLife. 2022 Nov 4;11:e68745.

32. Polovnikov KE, Brandão HB, Belan S, Slavov B, Imakaev M, Mirny LA. Crumpled
Polymer with Loops Recapitulates Key Features of Chromosome Organization.
Phys Rev X. 2023 Nov 13;13(4):041029.

726 Funding

The degron analysis in this work was supported by the grant of the state program of

the «Sirius» Federal Territory «Scientific and technological development of the

«Sirius» Federal Territory» (Agreement №26-03, 27/09/2024) (A.S.). Modelling part

of the study was supported by the Ministry of Science and Higher Education of the

Russian Federation, grant no. FSUS-2024-0018 (T.L.). The funders had no role in

the study design, data collection and analysis, decision to publish, or preparation of

the manuscript.

Acknowledgements

735 Computational resources were provided by HPC facilities at the collaborative center

736 «Bioinformatics» of ICG SB RAS (funded by the Ministry of Education and Science

737 of the Russian Federation, state project FWNR-2022-0019). The original manuscript

738 text was composed by authors; proofreading was conducted with the assistance of

739 ChatGPT version 5, after which the text was corrected and edited by the authors.

Author contributions

741 T.L. conceptualized the study. Bioinformatic analyses and polymer simulations were

742 carried out by V.K., with contributions from T.L. and A.S. V.K. drafted the manuscript

743 with contributions by T.L. and A.S. All authors reviewed, revised, and approved the

744 final manuscript.

Ethics approval and consent to participate

746 Not applicable

Consent for publication

748 Not applicable

Data Availability Statement

750 All the data and code underlying the findings are fully available. The simulation code

751 is deposited at https://github.com/Three-point/Chrom_polymer_models. The

752 simulation outputs (contact maps, .csv files and configuration files) are deposited at

753 Zenodo (DOI: 10.5281/zenodo.17828395). Figure source data are provided in the

754 Supporting Information.

Competing interests

756 The authors declare that they have no competing interests

Supporting information

S1 Fig. *Effect of extruder lifetime on contact probability P(s) profiles with*

different   *metrics*

A) P(s) curves in the range of 3 kbp - 2 Mbp for experimental data for the chicken

cell line DT40 (black line) and aggregated model data. Static (gray line) and dynamic

extruders with three lifetimes and base physical parameters (red, blue, and purple

lines). The large dots in the derivative graphs correspond to the mean loop size; B)

Results of comparing the segment experiment DT40 P(s) with the model data for the

dynamic extruders over three lifetimes with shifting normalisation points on different

metrics. The normalization point shifts within the range of 3 kbp to 2 Mbp. The

comparable segment of the plot is 3 kbp-2 Mbp. The sigma-noise is 200 nm.

S2 Fig. *Effects of extruder lifetime free loops on contact probability P(s)*

profiles with different metrics

A) P(s) curves in the range of 3 kbp - 2 Mbp for experimental data for the chicken

cell line DT40 (black line), aggregated model data. Static (gray line) and dynamic

extruders, which are unable to load at existing loops, on three lifetimes with base

physical parameters (red, blue, purple lines). The large dots on the derivative graphs

correspond to the mean loop size; B) Results of comparing the segments experiment

DT40 P(s) with the model data on different metrics. The normalization point shifts

within the range of 3 kbp to 2 Mbp. The comparable segment of the plot is 3 kbp-2

Mbp. The sigma-noise is 200 nm.

S3 Fig. *Effect of extruder lifetime free loops on contact probability P(s) profiles*

with loop size distributions.

A) P(s) curves in the range of 3 kbp - 2 Mbp for experimental data for the chicken

cell line DT40 (black line), aggregated model data. Static (gray line) and dynamic

extruders, which are unable to load at existing loops, on three lifetimes for Step

potential parameters (red, blue, purple lines). The large dots in the derivative graphs

correspond to the mean loop size; B) Loop size distributions for different lifetimes. All

the graphs are normalised on the area under the all-loops graph. Purple line - all

extruders. 1) Static extruders, 2) Dynamic extruders lt = 17 sec, 3) Dynamic

extruders lt - 68 sec, 4) Dynamic extruders 136 sec.

S1 Appendix. Artificial Noise Effects and Their Role in Model-Data

Reconciliation; Analysis of Spiralization in Chromosome Models Using C(s)

Correlation.