Investigating the repeatability and behavioral relationships of acuity, contrast sensitivity, form, and motion perception measurements using a novel tablet-based vision test tool

Jonathan Green [1], Jan Skerswetat *[2,3], Peter J. Bex [3], Gunnar Schmidtmann [1]

[1] School of Health Professions, University of Plymouth, Plymouth, United Kingdom
[2] Department of Psychology, Northeastern University, 360 Huntington Ave, Boston, Massachusetts, 02115, United States of America
[3] Department of Ophthalmology, University of California, 850 Health Sciences Road, Irvine, California, 9269, United States of America

iD

Jonathan Green: https://orcid.org/0000-0001-5738-7515
Jan Skerswetat: https://orcid.org/0000-0002-2470-8404
Peter J. Bex: https://orcid.org/0000-0001-7561-7695
Gunnar Schmidtmann: https://orcid.org/0000-0001-8180-7751

*Corresponding author email: jskerswe@hs.uci.edu
Emails: jon.green@plymouth.ac.uk, jskerswe@hs.uci.edu,
p.bex@northeastern.edu, gunnar.schmidtmann@plymouth.ac.uk

Abstract (280 words)

Visual function tests are important in basic and clinical vision research but are typically limited to very few aspects of human vision, coarse diagnostic resolution, and require an administrator. Recently, the generalizable, response-adaptive, self-administered Angular Indication Measurement (AIM) and Foraging Interactive D-prime (FInD) methods were developed to assess vision across different visual functions. The AIM and FInD paradigms show a range of visual stimuli per display (4x4 stimuli) spanning ±2σ around an adaptively estimated perceptual threshold across multiple displays. Here, we investigated the repeatability and behavioral relationships of the AIM and FinD paradigms for near visual acuity, contrast sensitivity function (CSF), form, and motion coherence threshold measurements using a novel tablet-based vision test tool. 31 healthy participants were recruited and repeated two repetitions of each experiment in random order. Bland-Altman analyses were performed to calculate the Coefficient of Repeatability (precision) and Mean Bias (accuracy). Linear regressions and hierarchical cluster analysis were used to investigate the relationship between outcome parameters. Results show that AIM Form coherence and FInD Form horizontal coherence showed significant retest bias; all other tests were bias-free. Cluster analysis revealed overall clustering of CSF, form and motion outcomes. We further show significant correlations within CSF and between motion coherence outcomes but few significant correlations between form coherence outcomes. AIM and FInD near vision tests are generalizable across multiple visual functions and are precise and reliable. Most functions tested were

bias-free. CSF, form, and motion outcomes clustered together, and CSF and motion
outcomes correlated with one another. The combination of a generalizable,
response-adaptive, and self-administered approach may be a suitable set of tests for
basic science and clinical use cases.

Keywords: visual acuity, contrast sensitivity, form perception, motion perception,
psychophysical methods, response-adaptive, self-administered, tablet-based, vision
test, angular indication measurement, foraging interactive D-prime,repeatability

Introduction

Identification and measurement of visual dysfunction is a global challenge. In excess of 1 billion people have visual impairments that could have been or are yet to be addressed (World Health Organisation (WHO), 2019; Bourne *et al.*, 2019) with the estimated associated loss of productivity valued at US$ 410.7 billion (WHO, 2023). This challenge is exacerbated by the disproportionate prevalence of visual deficiency in lower and middle-income countries (90%) (WHO, 2023; Rizzo *et al.*, 2020; WHO, 2022) and in higher income countries, amongst those from ethnic minorities, with lower socio-economic status, lower educational attainment (Cumberland *et al.* 2016; Elam *et al.* 2022), and poorer general and mental health (Cumberland *et a*l. 2016). Ageing is also strongly associated with increased risk for many visual conditions (Killeen *et al.*, 2023; Martinez-Roda *et al.*, 2016) - including subtle and under-recognised presentations, such as those often associated with mild traumatic brain injury (Rasdall *et al.*, 2024) (over 1-million annually in the US alone). Globally, the number of people aged over 65 is predicted to double over the next quarter of a century (United Nations, 2022), representing a substantial generator of pressure on often already overstretched health services.

Innovative digital technologies offer a range of opportunities to mitigate such challenges (Chen, Ding & Wang, 2023). Many digital vision assessment tools are available, some of which can now be operated using conventional portable devices such as laptop and tablet computers, or even smartphones (Bano, Wolffsohn & Sheppard, 2024). Advances in functionality support rapid, non-specialist, point-of care and domiciliary assessment, as well as self-administration and remote operation (telemedicine (Samanta *et al.*, 2023)). These technologies promise to drive more efficient assessment and monitoring, at earlier disease stages, reducing inconvenience and expense, and increasing access to underserved patient groups (Khurana *et al.*, 2021). As yet, these novel systems generally target a limited range of visual functions – and many are yet to be subjected to robust, peer-reviewed validation of their accuracy and reliability (Bano, Wolffsohn & Sheppard, 2024).

This study focuses on new suites of computer-based visual assessment tools which apply novel psychophysical paradigms and algorithms; namely the Angular Indication Measurement (AIM) (Skerswetat *et al.*, 2024a; Neupane, Skerswetat & Bex, 2024) and the Foraging Interactive D-prime (FInD) (Bex & Skerswetat, 2021; He, Bex & Skerswetat, 2023; Skerswetat *et al.*, 2024b). Tests employing these generalisable, self-administered, multi-dimensional (spatial/temporal frequency/contrast), and response-adaptive (personalised) psychophysical methods have been developed to measure more than 15 visual functions (Skerswetat *et al.*, 2024b) including visual acuity with and without glare source (Skerswetat, Boruta & Bex, 2022), colour (He, Bex & Skerswetat, 2023) and depth perception (Neupane, Skerswetat & Bex, 2024), spatio- and spatio-temporal contrast sensitivity (Skerswetat *et al.*, 2024b), threshold versus contrast suprathreshold perception (Bex

& Skerswetat, 2021; Bex & Skerswetat, 2023), form and motion perception (Bex &
Skerswetat, 2021; Skerswetat *et al.*, 2024b; Merabet *et al.*, 2023), and face
discrimination (Walter & Bex, 2024). Advantages of these methods compared to
traditional clinical tests and current laboratory computer-based methods include
flexible testing ranges, fine outcome resolution and improved detection of subtle
changes, a bias-free test protocol due to randomised stimulus orientations and
variables that eliminate memorisation risks, rapid testing through adaptive algorithms
and instant result generation, reduction in administrative burden and time due to self-
administration, and a generalizable method to measure multiple visual functions
using the same principal i.e., orientation judgement for AIM and signal presence
detection for FInD.

The AIM method can be applied with different stimulus and response formats,
generating orientation judgements. Psychometric functions are fit to continuous
orientation report errors (as a function of stimulus intensity) (Skerswetat *et al.*,
2024a). The FInD method measures the identification of cells containing signal
stimuli of varying intensities, amongst null stimuli cells. Thresholds are estimated
from *d'* analysis of hits, misses, false alarms, and correct rejection responses
(Skerswetat *et al.*, 2024b).

Previous studies have supported the refinement and administration of many of these
assessment tools. One study demonstrated the systematic application of a battery of
eleven tests, across both paradigms, for the broad visual assessment of 21 research
participants with visual dysfunction (people with albinism), compared with 21 typically
sighted controls (Skerswetat *et al.*, 2024b). Anticipated variance between groups
was observed – and the incorporation of testing pertinent to different levels of visual
processing provided insight into a lack of differentiation for some mid-level
functioning (pattern recognition, and radial and circular motion coherence). The FInD
global motion coherence test has also been demonstrated to identify an anticipated
significant reduction in performance amongst a cohort of 20 young people with
cerebral visual impairment, compared with 30 controls (mean difference 32.34%) -
concurrently providing evidence of usability by neurodiverse participants and for the
targeted assessment of cerebral visual processing (Merabet *et al.*, 2023).
Participants were able to use both form and motion tests, taking an average of
around 6 minutes to complete both.

In a study of 22 young adults with induced defocus and astigmatic blur, results from
the AIM visual acuity (VA) test demonstrated comparable sensitivity to conventional
Early Treatment Diabetic Retinopathy Study (ETDRS) letter chart assessment and
comparable repeatability when using three or more adaptive steps - whilst providing
additional personalised information regarding minimal error level (noise), bias and
slope from a psychometric function fit (Skerswetat *et al.*, 2024a). Mean AIM VA test
duration was only around 30 seconds. Elsewhere, a study of 24 children (including
11 amblyopes) demonstrated the usability of AIM VA for children as young as 5

years, taking an average of around 1.5 minutes (Ahmed *et al.*, 2023). The results
demonstrated good agreement with Snellen and Rosenbaum chart use and identified
interocular VA differences.

The AIM Contrast Sensitivity Function (CSF) test was appraised in a study of 10
participants (8 typically sighted, 1 amblyopic and 1 Ortho-K contact lens wearer with
defocus blur) (Skerswetat & Bex, 2022). Between eye results were similar, and less
than for both eyes together, consistent with binocular contrast summation.
Anticipated interocular difference and overall performance reduction was observed
for the respective atypically sighted participants. CSF testing was generally
completed within 2 minutes. Elsewhere, results of FInD CSF and Threshold versus
Contrast functions (TvC) tests were found to be comparable to the results of
equivalent standard temporal two-alternative forced choice (AFC) paradigms for up
to 14 participants (Bex & Skerswetat, 2021). The FInD tests were significantly
quicker to administer and no significant test retest difference or *bias* were detected
(Bex & Skerswetat, 2021).

The aim of the current study is to investigate the repeatability and behavioral
relationships of the AIM and FInD paradigms for acuity, contrast sensitivity, form,
and motion perception measurements implemented as a novel tablet-based vision
test tool.

Methods

Participants

31 observers (mean age: 39.7, σ: ±11.7; 13 females) participated in the initial test-
retest study. Seven additional observers (mean age: 39, σ: ±15; 4 females) were
recruited for a control experiment (AIM CSF with five displays, see details below). 36
observers were naive as to the purpose of the experiment. All participants had
normal or corrected-to-normal visual acuity. Further participant details are
summarized in Table 1. Informed, written consent was obtained from each observer,
and the study was approved by the University of Plymouth ethics committee. All
experiments were conducted in accordance with the Declaration of Helsinki.
160
161
162 **Table 1** Self-reported participant demographic and medical details. Ethnicity: White (UK, Irish)
163 includes English, Welsh, Scottish, Irish, British. Sex: f = female; m = male
164 ADHD & Autism Thresholds: Adler's Adult ADHD Self-Report Scale (ASRS-v1.1) Part A (screener) ≥4
165 grey boxes = highly consistent with ADHD; Baron-Cohen's Autism Spectrum Quotient (AQ): ≥32 =
166 Clinically significant (80% of those diagnosed with autism, but <2% of controls), ≥26 = Potential traits

ID	Age	Ethnicity	Sex	Medical history	Ocular conditions	ASRS	AQ
1	48	White (UK, Irish)	m		Myopia	0	20
2	44	White (UK, Irish)	m			0	14
3	48	White (UK, Irish)	m	Hemiplegic migraine	Astigmatism	1	23
4	22	Black	m			0	11
5	44	White (UK, Irish)	f			0	7
6	35	White (UK, Irish)	f			2	15
7	42	White (UK, Irish)	m	Concussion		1	18
8	49	White (UK, Irish)	m			0	23
9	39	White (UK, Irish)	m			0	11
10	29	White (UK, Irish)	f	Anxiety, Emotional dysregulation, Depression		3	10
11	28	White (UK, Irish)	m			2	22
12	23	Asian	m		Color-Vision-Defect	0	12
13	20	Black	f			0	17
14	28	White (UK, Irish)	f			2	10
15	28	White (UK, Irish)	m			5	39
16	48	White (UK, Irish)	m			2	21

17	26	Asian	m			2	25
18	32	White (UK, Irish)	m	Concussion		3	28
19	29	White (UK, Irish)	m	Concussion		2	21
20	19	White (UK, Irish)	f			2	12
21	57	White (UK, Irish)	f			0	11
22	54	White (any other)	f			0	10
23	48	White (UK, Irish)	f	Dyslexia		2	14
24	43	White (UK, Irish)	m			2	20
25	58	White (UK, Irish)	f			1	15
26	35	White (UK, Irish)	m			0	8
27	50	White (UK, Irish)	f		Astigmatism	0	15
28	55	White (UK, Irish)	f			2	24
29	46	White other)	m			0	13
30	51	White (UK, Irish)	f			0	10
31	40	mixed	m			0	30

167
168

Stimuli and psychophysical procedure

Each display chart for each test contained a 4x4 set of cells. All tests used three charts per experimental run, including the first initial chart and two adaptive steps for all tests. The range of stimulus test levels on each display spanned ±2σ around an adaptively estimated perceptual threshold across displays (e.g., Skerswetat *et al.*, 2024b; Skersetat, Boruta & Bex, 2022). For form and motion perception testing, the default display mode, which presents all stimuli simultaneously and also the hidden cell approach, which shows only the stimulus over which the user positions the cursor (Merabet *et al.*, 2023) was used. Orientations or directions of stimuli within cells were randomized for all AIM tests. Stimulus examples are shown in Figure 1.

AIM Acuity: C optotypes were presented within each indication ring (diameter, Ø: 5°) with an inter-cell distance of 1°. Stimulus contrast was 0.97. The minimum range of stimuli sizes was set to 0.20 logMAR. The minimum visual acuity was kept at the minimum renderable size for -0.05 logMAR and maximum logMAR starting value was set to 1.00 logMAR. More details are published elsewhere (Skerswetat *et al.*, 2024a).

AIM CSF 2D+: Gabors (σ: 0.5°) were presented within each indication ring (Ø: 5°) with an inter-cell distance of 1˚. More details are published elsewhere (Skerswetat *et al.*, 2024b)

AIM Form Coherence (with and without Hidden Cells): Glass Pattern (Glass, 1969; Glass & Perez, 1973) coherence for translational form coherence was measured with 100 dipoles (50 black, 50 white; cell Ø: 5°; dot Ø=0.1°; intra-dipole spacing=0.2°).

AIM Motion Coherence (with and without Hidden Cells): Motion coherence for translational global dot motion was measured. Each cell (Ø: 5°) contained 100 black or white dots (Ø=0.1°, lifetime=100ms with random starting age, speed=2°/sec, dot contrast: 1.0). Stimulus examples are presented in the supplementary AIM_Motion.mp4 video file.

FInD Form (with and without Hidden Cells): Glass Pattern coherence (Merabet *et al.*, 2023; Glass, 1969; Glass & Perez, 1973) for translational, circular, and expanding (radial) form coherence was measured with 100 dipoles (50 black, 50 white; cell Ø: 3x3° squares; dot Ø=0.05°; inter-dot spacing=0.2°). The inter-cell distance was 3°.

FInD Motion (with and without Hidden Cells): Global motion coherence (Merabet *et al.*, 2023) for translational, circular, and expanding form coherence was measured with 100 dipoles (50 black, 50 white; dot contrast: 1.0; cell Ø: 3*3° squares; dot Ø=0.05°; speed=1°/sec, lifetime=100ms with random starting age). The inter-cell distance is 3°.

AIM specific feedback: Auditory feedback was provided after each chart based on a statistical probability analysis of per-chart report errors, if the mean error report was indistinguishable from random responses (i.e., 90° error for C optotypes and motion directions, 45° for gratings and Glass patterns) the application altered the user to report the orientation carefully.

AIM Visual Acuity

AIM CSF+

AIM Form and Motion

FInD Form and Motion

220
221 **Figure 1** The images show charts containing stimulus examples for the AIM Visual Acuity (VA), AIM
222 CSF+, AIM Form and Motion Coherence and FInD Form and Motion Coherence tests as seen by the
223 observers on the tablet.

Apparatus

Stimuli were presented on a Microsoft Surface Pro 8 (Core™ i5 CPU 8GB, 256GB
Go RAM) with display dimensions of 27.4 x 18.3 cm. The display's spatial resolution
was 2880 x 1920 pixels with a 60Hz frame rate. All experiments were carried out at a
viewing distance of 40 cm. At this distance one pixel subtends 0.0136° visual angle.
The screen was gamma-corrected using data color Spyder (Datacolor, Lucerne,
Switzerland) photometer. The input for the AIM tests was via the tablet's
touchscreen. The AIM hidden cell and FinD paradigm reports were made using a
common computer mouse.

Procedure

235 Repeatability of near visual acuity, contrast sensitivity function (CSF), and form &
236 motion coherence tasks were deployed on a tablet (see above). Observations were
237 made under binocular viewing conditions. The test order was randomised, each
238 experiment included a test and a retest session and took in total ~90 min.

Results

Psychophysical outcomes

The PerZeption application generated automatically outcomes as described below.

AIM: A cumulative Gaussian function was fit to the data of the AIM Visual Acuity, AIM Form Coherence, and AIM Motion Coherence tests (Neupane *et al.*, 2024; Skerswetat *et al.*, 2024b). Threshold, slope, and minimum report error parameter were calculated. For the AIM CSF 2D+, the area under the log CSF (AULCSF) and the high spatial frequency cut off, or CSF Acuity were analysed.

FInD: Response of each cell was classified as either Hit, Miss, False Alarm, or Correct Rejection, to calculate *d'* as a function of motion or form coherence. The probability of a Yes response as a function of signal intensity was subsequently calculated (Merabet *et al.*, 2023). Threshold (*d'*=1), slope, and false alarm rate parameters were calculated.

Bland-Altman Analysis

Test-retest reliability for all tests was assessed using Bland-Altman (Altman & Bland, 1983) plots using MATLAB (R2024b, Matworks). Figure A1 in the appendix shows an example (see Supplementary Material for all Bland-Altman plots).

We subsequently calculated the Coefficient of Repeatability (precision, *CoR*) according to

$$CoR = 1.96 \sqrt{2} \, \sigma, \tag{1}$$

where σ refers to the within-subject standard deviation. The mean bias ($\bar{x}$, accuracy) in Table 2 was calculated as

$$\bar{x} = \frac{\sum_{i=1}^{n}(m_{1i} - m_{2i})}{n}. \tag{2}$$

where m_1 and m_2 denote the test and retest measurements, respectively. One-sample t-tests were conducted to assess whether the bias is statistically significant. Test-retest data (m_1, m_2), *CoR*, mean *Bias* and *p*-values are summarized in Table 2, where $p \leq 0.05$ is considered statistically significant and indicated by asterisks.

Results show that AIM Form coherence results and FInD Form horizontal coherence results showed significant retest biases, but all other tests were bias-free.

270   **Table 2** Test Retest results for threshold values (Coh. = coherence threshold, HC = hidden cell mode,
271   ☐ = standard deviation, *CoR* = coefficient of repeatability).

Visual function test	Basic Statistics		Bland-Altman Statistics		
	Test (m_1) Mean ±σ	Retest (m_2) Mean ±σ	*CoR*	Mean *Bias*	Mean *Bias p*
AIM-VA [logMAR] (*fixed at limit (-0.104))	-0.06 ±0.05	-0.06 ±0.06	±0.19	-0.001	*p*=.94
AIM CSF+: AULCSF (3 charts) 5 charts *fixed at limit	2.55 ±0.54 3.00 ±0.014	2.41 ±0.52 2.98 ±0.066	±1.68 ±0.14	0.14 0.02	*p*=.21 *p*=.36
AIM CSF+: Acuity (3 charts) [c/°]	26.0 ±5.8	23.7 ±5.9	±20.1	2.3	*p*=.09
5 charts *fixed at limit	30.9 ±0.00	30.6 ±0.79	±2.26	0.3	*p*=.36
AIM-Form Coh. [%]	44.6 ±24.3	36.9 ±20.3	±44.8	7.6	*p*=.013*
AIM-Form Coh. [%] Hidden cell mode	49.9 ±27.7	44.5 ±24.7	±66.5	5.4	*p*=.21
AIM-Motion Coh. [%]	41.8 ±27.3	41.5 ±27.2	±85.3	0.3	*p*=.96
AIM-Motion Coh. [%] HC	41.2 ±28.3	40.4 ±28.3	±78.9	0.8	*p*=.87
FInD-Form Coh. - Horizontal [%]	41.1 ±25.8	32.4 ±22.7	±75.4	8.7	*p*=.08
FInD-Form Coh. - Expanding [%]	38.4 ±16.9	35.5 ±18.9	±75.0	2.8	*p*=.56
FInD-Form Coh. - Circular [%]	39.7 ±19.5	36.2 ±17.9	±61.8	3.5	*p*=.39
FInD-Form Coh. - Horizontal[%] HC	43.1 ±26.8	26.8 ±21.0	±68.3	16.0	*p*<.001***
FInD-Form Coh. - Expanding [%] HC	32.9 ±19.0	30.7 ±22.0	±72.3	2.2	*p*=.64
FInD-Form Coh. - Circular [%] HC	38.5 ±14.0	38.2 ±19.0	±58.4	0.3	*p*=.93
FInD-Motion Coh. - Horizontal [%]	25.6 ±19.5	35.3 ±25.5	±66.0	-9.7	*p*=.03*
FInD-Motion Coh. - Expanding [%]	31.3 ±15.5	33.7 ±15.7	±37.9	-2.4	*p*=.33
FInD-Motion Coh. - Circular [%]	19.1 ±12.9	22.4 ±15.8	±35.5	-3.3	*p*=.16
FInD-Motion Coh. - Horizontal [%] HC	21.3 ±19.8	27.2 ±25.5	±57.1	-6.0	*p*=.12
FInD-Motion Coh. - Expanding [%] HC	32.7 ±18.8	33.2 ±18.7	±59.4	-0.5	*p*=.89
FInD-Motion Coh. - Circular [%] HC	18.3 ±11.7	20.1 ±13.0	±29.1	-0.02	*p*=.35

282 **Table 3** Test Retest results for slope values.

Slopes	Test (m_1) Mean ±σ	Retest (m_2) Mean ±σ	CoR	Mean Bias	Mean Bias p
AIM-VA	0.12 ±0.12	0.12 ±0.13	0.49	-0.01	p=.8694
AIM CSF+: AULCSF (3 charts)	0.34 ±0.11	0.26 ±0.14	0.46	0.08	p=.0169*
AIM CSF+ Acuity (3 charts)	0.34 ±0.11	0.26 ±0.14	0.46	0.08	p=.0169*
AIM-Form Coh.	0.20 ±0.15	0.17 ±0.13	0.50	0.03	p=.3597
AIM-Form Coh. HC	0.23 ±0.18	0.21 ±0.17	0.56	0.01	p=.6926
AIM-Motion Coh.	0.16 ±0.16	0.12 ±0.11	0.47	0.04	p=.2310
AIM-Motion Coh. HC	0.16 ±0.13	0.14 ±0.12	0.50	0.02	p=.4814
FInD-Form Coh. - Horizontal	1.94 ±1.32	1.82 ±1.33	5.34	0.12	p=.7282
FInD-Form Coh. - Expanding	2.65 ±1.32	2.28 ±1.28	4.30	0.37	p=.1975
FInD-Form Coh. - Circular	2.63 ±1.22	2.16 ±1.22	3.59	0.47	p=.0520
FInD-Form Coh. - Horizontal HC	1.99 ±1.14	1.71 ±1.29	5.50	0.28	p=.4380
FInD-Form Coh. - Expanding HC	2.31 ±1.41	1.88 ±1.34	5.13	0.43	p=.2064
FInD-Form Coh. - Circular HC	2.55 ±1.24	2.79 ±1.35	5.10	-0.24	p=.4704
FInD-Motion Coh. - Horizontal	2.34 ±1..49	2.55 ±1.47	5.42	-0.20	p=.5656
FInD-Motion Coh. - Expanding	2.55 ±1.24	3.03 ±1.13	4.47	-0.49	p=.1033
FInD-Motion Coh. - Circular	2.27 ±1.40	2.48 ±1.21	5.38	-0.22	p=.5349
FInD-Motion Coh. - Horizontal HC	1.95 ±1.33	2.14 ±1.39	5.81	-0.19	p=.6260
FInD-Motion Coh. - Expanding HC	2.84 ±1.15	2.78 ±1.43	4.45	0.06	p=.8306
FInD-Motion Coh. - Circular HC	2.61 ±1.29	2.53 ±1.34	4.27	0.08	p=.7628

287 **Table 4** Test-retest results for minimum report errors values.

Errors [°]	Test (m_1) Mean ±σ	Retest (m_2) Mean ±σ	CoR	Mean Bias	Mean Bias p
AIM-VA	12.05 ±5.77	11.43 ±5.65	22.27	0.62	p=.6695
AIM CSF+: AULCSF	8.32 ±3.14	8.41 ±3.01	9.96	-0.09	p=.8900
AIM CSF+ Acuity	8.32 ±3.14	8.41 ±3.01	9.96	-0.09	p=.8900
AIM-Form Coh.	10.10 ±6.74	12.36 ±7.58	23.20	-2.26	p=.1430
AIM-Form Coh. HC	9.41 ±7.87	11.14 ±8.89	27.64	-1.73	p=.3410
AIM-Motion Coh.	15.94 ±15.71	15.40 ±15.80	47.84	0.54	p=.8631
AIM-Motion Coh. HC	16.29 ±13.39	13.25 ±15.78	53.69	3.04	p=.3894
FInD-Form Coh. - Horizontal	0.09 ±0.11	0.07 ±0.09	0.33	0.01	p=.5649
FInD-Form Coh. - Expanding	0.08 ±0.09	0.09 ±0.13	0.33	0.00	p=.8917
FInD-Form Coh. - Circular	0.04 ±0.07	0.07 ±0.08	0.15	-0.03	p=.0097*
FInD-Form Coh. - Horizontal HC	0.12 ±0.14	0.08 ±0.11	0.38	0.04	p=.0896
FInD-Form Coh. - Expanding HC	0.14 ±0.18	0.07 ±0.08	0.49	0.07	p=.0336*
FInD-Form Coh. - Circular HC	0.08 ±0.10	0.07 ±0.10	0.29	0.01	p=.4588
FInD-Motion Coh. - Horizontal	0.13 ±0.17	0.12 ±0.17	0.48	0.01	p=.8697
FInD-Motion Coh. - Expanding	0.05 ±0.07	0.03 ±0.06	0.23	0.02	p=.2093
FInD-Motion Coh. - Circular	0.04 ±0.07	0.07 ±0.14	0.24	-0.03	p=.0982
FInD-Motion Coh. - Horizontal HC	0.13 ±0.20	0.13 ±0.16	0.50	0.00	p=.8969
FInD-Motion Coh. - Expanding HC	0.08 ±0.09	0.08 ±0.13	0.29	0.00	p=.8124
FInD-Motion Coh. - Circular HC	0.03 ±0.04	0.05 ±0.13	0.33	-0.02	p=.3096

Table 5 Summary of overall testing time and per-chart time.

Visual function test	Total Duration [s]		Chart duration [s]	
	Test (m_1) Mean ±σ	Retest (m_2) Mean ±σ.	Test (m_1) Mean ±σ	Retest (m_2) Mean ±σ
AIM-Visual Acuity	109 ±40	92 ±34	36 ±13	31 ±11
AIM CSF+ (3 charts)	145 ±64	111 ±46	48 ±21	37 ±15
AIM- Form Coherence	196 ±111	177 ±21	65 ±40	59 ±82
AIM- Form Coherence HC	244 ±191	164 ±107	81 ±64	55 ±36
AIM- Motion Coherence	222 ±90	176 ±34	74 ±30	59 ±34
AIM- Motion Coherence HC	283 ±126	216 ±83	94 ±42	72 ±28
FInD- Form Coherence	263 ±288	182 ±124	29 ±32	20 ±14
FInD- Form Coherence HC	365 ±332	225 ±140	40 ±37	25 ±16
FInD- Form Coherence	306 ±165	248 ±99	34 ±18	28 ±11
FInD- Form Coherence HC)	360 ±139	301 ±149	40 ±15	33 ±17

Correlation and Cluster Analysis

Python, including the Seaborn toolbox (Waskom, 2021) was used to perform and visualize the linear correlation and hierarchical cluster analysis. Data were normalized using z-score normalization. The marginal cluster trees show a hierarchical cluster analysis using Ward's Minimum Variance Clustering Method based on the Euclidean distance between objects (i.e., test results). Linear correlations using Kendall coefficients were used. We conducted a linear regressions and hierarchical cluster analysis to investigate the relationship between outcome parameters. Kendall's correlation coefficients between all tests were calculated. The matrix in Figure 2 shows the correlation coefficients between all tests as color-code and value in each cell of the matrix. The warmer colors indicate higher and the colder colors lower correlations (see color bar). The asterisks indicate whether the correlation was significantly different from zero. This analysis reveals significant

correlations within CSF and between motion coherence outcomes but few significant
correlations between form coherence outcomes.

The marginal cluster trees in Figure 2 show results from a hierarchical cluster
analysis using Ward's Minimum Variance Clustering Method based on the Euclidean
distance between objects (i.e., tests). This analysis revealed overall clustering of
CSF, Form and Motion tests.

315
316

317
318 **Figure 2** (i) The figure shows Kendall's correlation coefficients between all tests as color-code and
319 value in each cell of the matrix, where warmer colors indicate higher correlations (see color bar). The
320 asterisks indicate whether the correlation was significantly different from zero. (ii) The marginal cluster
321 trees show results from a hierarchical cluster analysis).
322

Correlation between performance and ASRS and AQ scores

In order to investigate whether performance is affected by Asperger syndrome (i.e.,
high-functioning autism) or attention deficit hyperactivity disorder (ADHD), all
subjects filled out the Adult ADHD Self-Report Scale (ASRS, Kessler *et al.*, 2005)
and autism-spectrum quotient (AQ, Baron-Cohen *et al.*, 2001) questionnaires. The
results from these questionnaires are summarized in Table 1. We subsequently
calculated Pearson's linear correlation coefficients between all tests and the ASRS
and AQ scores. The correlation plots, including the correlation coefficients and p-
values, are shown in the supplementary ASRS_AQ_correlations.pdf. None of these
correlations was significant.
333
334

Discussion

The first goal of this study was to investigate the repeatability of the AIM and FInD paradigms for a variety of visual functions (acuity, contrast sensitivity, form- and motion coherence) performed on a tablet computer.

Comparison with previous FInD and AIM studies

AIM Visual Acuity: Previous studies investigated the repeatability of AIM VA for a viewing distance of 4m instead of near testing as performed in the current study (Skerswetat *et al.*, 2024a). That study showed that increasing the number of adaptive charts reduces the repeatability variance as expressed using limits of agreement which for 3 charts i.e., one initial and two adaptive steps ±0.24 logMAR was comparable to the ±0.14 logMAR (CoR: ±0.19 logMAR) of the current study. Also, the current study and the previous study found no systematic retest bias.

Form and Motion: A study in patients with and without cerebral visual impairment (Merabet *et al.*, 2023) reported 27% for circular motion and 30% for form thresholds using the hidden cell FInD tasks in the control cohort, which is slightly different to 18% and 39%, respectively, as reported in the current study's first test session (Table 2). This may be due to the different stimulus properties of the tablet used here and the 32" computer display used in Merabet *et al.* (2023). We are not aware of any published estimates of test-retest repeatability of coherence measured with other paradigms, such as 2AFC. In a study concerning vision in people with and without albinism (Skerswetat *et al.*, 2024b), thresholds for the control group, using FInD circular Motion and Form Coherence with hidden cells mode, were 22% and 52%, respectively. Again, higher sensitivity in motion compared to form coherence was shown, but lower sensitivity for form coherence, which can be explained due to the higher age of the cohort since form coherence thresholds increase with age (McKendrick, Weymouth & Battista, 2013). FInD complex motion and form coherence tasks showed no significant mean biases, whereas bias was found here for translation, simple form and motion signals (Table 2). Future research may investigate whether the bias reflects learning and might disappear between the second and third sessions. In the meantime, it is suggested to run an initial training session prior to an actual task for these stimulus types. The study also revealed a wide inter-individual difference in the ability to perceive global motion and form percepts (Figure 3).

AIM CSF 2D + showed no retest bias but had significantly larger variance when using three charts compared to five charts. Hence, it is suggested to use five charts as the default to ensure minimum retest variance.

Psychometric parameters

Some previous studies have investigated slope parameters (e.g., Wallis *et al*., 2013), however to our best knowledge no study specifically looked at the repeatability of slope parameters. Table 3 shows the slope and table 4 the minimum report error or *noise* (i.e., result of all noise factors) with the corresponding repeatability results. With the exception of CSF results using three charts, all slopes were bias free. The noise parameters were mostly bias-free except for FInD Form circular coherence and hidden cell expanding coherence. These results show that the methods employed in the AIM and FInd paradigms produce consistent and repeatable measurements.

Testing durations

The average time for visual acuity (Table 5) was less than two minutes in total and an approximate chart duration of 30 seconds, which is comparable to previously reported times when testing for distance visual acuity (Skerswetat *et al*., 2024a). The mean chart duration was less than one minute for the CSF task, hence the total duration for 5 charts is approximately 4 minutes, which is faster than conventional tests that may take 25 minutes (Kelly & Savoie, 1973) and comparable to other adaptive rapid procedures (Liu *et al*., 2021).

The test durations for the Form and Motion task varied more wieldy specifically between 3 and 6 minutes per test, one because the FInD task tested three subtypes of form and motion percepts, namely horizontal, circular and expanding percepts, compared to translational, simple form or motion percepts. The hidden cell paradigms caused longer testing durations than the open cell paradigms. This may be due to either the use of a mouse instead of finger touches, different gaze strategies used to detect signals, or a combination of these reasons. Since we did not collect gaze position data, future studies may investigate whether different gaze strategies were used for open and hidden cell modes.

Functional relationship between psychometric outcomes

The second goal of the study was to investigate relationships among thresholds. We used a combination of linear models and hierarchical cluster analysis, unlike in a previous study where these analyses were performed separately (Skerswetat *et al*., 2024b). The current study shows overall within-function outcomes clustering, specifically for contrast - contrast, form - form, and motion - motion outcomes (Figure 2). Similarly, contrast outcomes correlated with each other, as well as motion outcomes. Skerswetat *et al.* (2024b) and Bosten *et al.* (2017) also found significant correlations among contrast outcomes, but not among motion values. We speculate

that this difference may be due to the more heterogeneous age range and smaller
sample within the control cohort of that study compared to the current study. The
same study (Skerswetat *et al.*, 2024b) also showed clustering within contrast and
within form and motion outcomes for the control cohort, which is in agreement with
the current study. Overall, these results suggest distinct processing pathways of
contrast and mid-level form and motion perception.

Comparison between horizontal, expanding and circular form and motion

The results from the AIM and FInD Form and Motion tests have implications
regarding the visual processing of global texture and motion. There is ample
psychophysical and physiological evidence for the hypothesis that the representation
of objects and shapes is achieved by a hierarchical feedforward process along the
ventral pathway (e.g., Cadieu *et al.*, 2007; Serre *et al.*, 2005; Serre, Oliva, & Poggio,
2007; Van Essen, Anderson, & Felleman, 1992). A key question in vision science is
how local information detected in the early visual areas (primary visual cortex) is
integrated to represent more complex, behaviorally relevant stimuli (e.g., faces,
objects) at subsequent stages. Psychophysically, this has been tested in different
ways. One approach, also employed in the Form and Motion tests, is to determine
the minimum number of signal elements required to detect the presence of a global
pattern embedded in an array of noise elements (coherence thresholds). This
measurement of coherence thresholds has been used in many studies of motion
(Braddick *et al.*, 2000; Newsome, 1988), texture (Dakin, 1997; Schmidtmann *et al.*,
2015; Wilson & Wilkinson, 1998; Wilson, Wilkinson, & Asaad, 1997), and contour
perception (Achtman, Hess, & Wang, 2003; Braddick *et al.*, 2000; Schmidtmann *et
al.*, 2013; 2015). Coherence thresholds in Glass patterns (Glass, 1969; Glass &
Perez, 1973) have been employed to investigate signal integration in texture
perception (Dakin, 1997; Dakin & Bex, 2002; Dickinson, Broderick, & Badcock, 2009;
Wilson, Wilkinson & Asaad, 1997; Wilson & Wilkinson, 1998). Models of Glass
pattern detection propose a two-stage process, in which the first stage extracts local
orientation information while the second stage integrates the orientation information
in order to extract the global structure (Wilson, Wilkinson & Assad, 1997; Wilson &
Wilkinson, 1998, see Schmidtmann *et al.* (2015) for a revised model). This model is
supported by physiological data showing that neither the classical receptive fields
nor surround mechanisms in early visual areas (V1, V2) are sufficient to process the
Glass patterns (Smith, Bair, & Movshon, 2002; Smith *et al.*, 2007). Some previous
psychophysical studies found no difference in sensitivities for detecting circular and
radial structure (e.g., see Kelly *et al.*, 2001; Kurki & Saarinen, 2004; Pei *et al.*, 2005;
Seu & Ferrera, 2001; Wilson, Wilkinson & Asaad, 1997; Wilson & Wilkinson, 1998),
while others reported a significantly increased sensitivity of human observers for
circular, followed by radial, spiral, and parallel structure (Kelly *et al.*, 2001; Kurki &
Saarinen, 2004; Seu & Ferrera, 2001; Wilson, Wilkinson & Asaad, 1997; Wilson &
Wilkinson, 1998), suggesting that the global structure in Glass patterns is processed
by specialized detectors for circular and radial textures. However, there is also

contradictory evidence. For instance, Dakin and Bex (2002) found that the
advantage for circular Glass patterns was only evident if the stimulus was presented
within a circular window as opposed to a square window. Similarly, Schmidtmann *et*
al. (2015) were not able to replicate the above mentioned hierarchy. Their results
showed that detection sensitivity was independent of texture type, hence, no
evidence for specialized detectors for circular structure.
470   The results presented here also do not show the clear sensitivity hierarchy
(i.e., circular > expanding > horizontal) described above. Figure 3 shows Box and
Whisker plots comparing coherence thresholds achieved for horizontal, expanding
and circular structure for the FInD Form, FInD Form HC, FInD Motion, and FInD
Motion HC tests. Following Shapiro Wilk tests for normality which showed that some
of the conditions were not normally distributed, the data were analyzed using a non-
parametric Friedman ANOVA (Durbin-Conover) across all conditions. This analysis
revealed a statistically significant global effect (χ^2(119)=11, p<.001). Subsequent
pairwise comparisons (Bonferroni-corrected for multiple comparisons) showed that
for the FInD Form paradigms, sensitivity was largely independent of texture type,
except a statistically significant increase in thresholds (i.e. decrease in sensitivity)
between FInD Form HC horizontal and FInD Form HC circular. In general,
performance for the FInD Motion paradigms was slightly (sometimes significantly
better; see Table A1) compared to the static FInD Form stimuli. Results show that
sensitivity for FInD Motion horizontal and FInD Motion expanding stimuli was
significantly lower compared to FInD Motion circular. Finally, sensitivity for FInD
Motion HC expanding was significantly lower compared to FInD Motion HC circular
and FInD Motion HC horizontal. No superior performance for circular motion is
evident.

Figure 3 Box and Whisker plots with individual data (grey data points) and the mean (black solid data point) for the FInD Form (top left), FInD Form HC (top right), FInD Motion (bottom left) and FInD Motion HC (bottom right) tests (coherence thresholds are expressed in %, 0-1). The outliers (red diamonds) with coherence thresholds of ~1 (~100%) were from two subjects who could not do the task, independent of global structure.

Future directions

The current study investigated the repeatability and functional relationship of the tablet based psychophysical tests in young healthy participants. Based on the high repeatability of these results and the normative data, future research can be adequately powered to test the effect of age and the effect in clinically relevant cohorts on the test performance. The number of cells within each chart and the number of charts determine the testing duration, and the present data can guide future research to investigate their relationship to optimise the time-repeatability tradeoff.

Conclusion

AIM and FInD near vision tests are generalizable across multiple visual functions and are precise and reliable. Most functions tested were bias-free, except translational form and motion coherence, which may be improved with additional

instructions or training charts. CSF, form, and motion outcomes clustered together,
and CSF and motion outcomes correlated with one another, indicating separate
processing mechanisms for achromatic spatial and global coherence perception.
Together, the results demonstrate the reliability of the tests, which may be useful for
basic and clinical research.

Acknowledgements
Conflict of interest: JS and PJB are co-founders and shareholders of PerZeption Inc.
JS and PJB are the inventors of AIM and FInD technology (P). Northeastern
University is the owner of the intellectual property and granted exclusive licensing
rights to PerZeption Inc.
GS and JG were supported by the College of Optometrists (UK).
PJB and JS were supported by NIH R01EY032162.
524

References

Achtman, R., Hess, R. F., & Wang, Y. Z. (2003). Sensitivity for global shape detection. Journal of Vision, 3(10):4, 616–624 https://doi.org/10.1167/3.10.4

Ahmed Z., Skerswetat J., Bex P.J. & Wiecek E.K. (2023) Assessment of Visual Acuity in Children With and Without Amblyopia Using the Novel Method AIM (Angular Indication Measurement) Acuity. Investigative Ophthalmology & Vision Science 2023;64(8):1505 Available at: https://iovs.arvojournals.org/article.aspx?articleid=2790964

Altman, D. G., & Bland, J. M. (1983). Measurement in medicine: the analysis of method comparison studies. Journal of the Royal Statistical Society Series D: The Statistician, *32*(3), 307-317.http://dx.doi.org/10.2307/2987937

Baron-Cohen, S., Wheelwright, S., Skinner, R., Martin, J., & Clubley, E. (2001). The autism-spectrum quotient (AQ): Evidence from asperger syndrome/high-functioning autism, males and females, scientists and mathematicians. Journal of autism and developmental disorders, 31, 5-17.

Bano, T., Wolffsohn, J.S. & Sheppard, A.L. (2024) Assessment of visual function using mobile Apps. Eye 38:2406–2414 https://doi.org/10.1038/s41433-024-03031-2

Bex P. & Skerswetat J. (2021) FInD - Foraging Interactive D-prime, a rapid and easy general method for visual function measurement. Journal of Vision 21(9):2817 https://doi.org/10.1167/jov.21.9.2817

Bex P.J. & Skerswetat J. (2023) AIM+ (Angular Indication Measurement) Efficient Assessment of Multidimensional Functions;45th European Conference on Visual Perception (ECVP) 2023 Paphos, Cyprus Perception (Collections: ECVP Abstracts) pp202-3. Available at: https://journals.sagepub.com/page/pec/collections/ecvp-abstracts/index/ecvp-2023

Bosten, J. M., Goodbourn, P. T., Bargary, G., Verhallen, R. J., Lawrance-Owen, A. J., Hogg, R. E., & Mollon, J. D. (2017). An exploratory factor analysis of visual performance in a large population. Vision Research, 141, 303-316. https://doi.org/10.1016/j.visres.2017.02.005

Bourne, R. and GBD 2019 Blindness and Vision Impairment Collaborators (2021) Trends in prevalence of blindness and distance and near vision impairment over 30 years: an analysis for the Global Burden of Disease Study. The Lancet Global Health 9(2):e130 - e143 https://doi.org/10.1016/S2214-109X(20)30425-3

Braddick, O. J., O'Brien, J. M., Wattam-Bell, J., Atkinson, J., & Turner, R. (2000). Form and motion coherence activate independent, but not dorsal/ventral segregated, networks in the human brain. Current Biology, 10(12), 731–734 https://doi.org/10.1016/s0960-9822(00)00540-6

Britten, K.H. & Newsome, W.T. (1998) Tuning Bandwidths for Near-Threshold Stimuli in Area MT. Journal of Neurophysiology 80:2, 762-770 https://doi.org/10.1152/jn.1998.80.2.762

Cadieu, C., Kouh, M., Pasupathy, A., Connor, C. E., Riesenhuber, M., Poggio, T., & Pasupathy, A. (2007). A model of V4 shape selectivity and invariance. Journal of Neurophysiology, 98(3), 1733–1750 https://doi.org/10.1152/jn.01265.2006

Chen, C., Ding, S. & Wang, J. (2023) Digital health for aging populations. Nature Medicine 29:1623–1630 https://doi.org/10.1038/s41591-023-02391-8

Cumberland PM, Rahi JS, for the UK Biobank Eye and Vision Consortium (2016) Visual Function, Social Position, and Health and Life Chances: The UK Biobank Study. JAMA Ophthalmology 134(9):959–966 https://doi:10.1001/jamaophthalmol.2016.1778

Dakin, S. C., & Bex, P. J. (2002). Summation of concentric orientation structure: Seeing the Glass or the window? Vision Research, 42(16), 2013–2020 https://doi.org/10.1016/S0042-6989(02)00057-3

Dakin, S. C. (1997). The detection of structure in glass patterns: Psychophysics and computational models. Vision Research, 37(16), 2227–2246 https://doi.org/10.1016/S0042-6989(97)00038-2

Dickinson, J.E., Broderick C., Badcock D.R. (2009) Selective attention contributes to global processing in vision. Journal of Vision 9(2):6 https://doi.org/10.1167/9.2.6

Elam, A. R., Tseng V. L., Rodriguez T. M., Mike E. V., Warren A. K., Coleman A. L. and American Academy of Ophthalmology Taskforce on Disparities in Eye Care (2022) Disparities in Vision Health and Eye Care. Ophthalmology 129 (10):e89-e113 https://doi.org/10.1016/j.ophtha.2022.07.010

Glass, L., & Perez, R. (1973). Perception of random dot interference patterns. Nature, 246, 360–362 https://doi.org/10.1038/246360a0

Glass, L. (1969). Moiré effect from random dots. Nature, 223, 578–580 https://doi.org/10.1038/223578a0

He J., Bex P.J. & Skerswetat J. (2023) Rapid measurement and machine learning classification of colour vision deficiency. Ophthalmic and Physiological Optics 43:1379–1390 https://doi.org/10.1111/opo.13210

Kelly, D. M., Bischof, W. F., Wong-Wylie, D. R., & Spetch, M. L. (2001). Detection of glass patterns by pigeons and humans: Implications for differences in higher-level

processing. Psychological Science, 12(4), 338–342 https://doi.org/10.1111/1467-
9280.00362

Kelly D. H., Savoie R. E. (1973). A study of sine-wave contrast sensitivity by two
psychophysical methods. Perception & Psychophysics, 14, 313–318.
Kessler, R. C., Adler, L., Ames, M., Demler, O., Faraone, S., Hiripi, E. V. A., ... &
Walters, E. E. (2005). The World Health Organization Adult ADHD Self-Report Scale
(ASRS): a short screening scale for use in the general population. Psychological
medicine, 35(2), 245-256.

Khurana R. N., Hoang C., Khanani A. M., Steklov N. & Singerman L. J. (2021) A
Smart Mobile Application to Monitor Visual Function in Diabetic Retinopathy and
Age-Related Macular Degeneration: The CLEAR Study. American Journal of
Ophthalmology 227:222-230 https://doi.org/10.1016/j.ajo.2021.03.033

Killeen OJ, De Lott LB, Zhou Y, Hu M., Rein D., Reed N., Swenor B.K. & Ehrlich J.R.
(2023) Population Prevalence of Vision Impairment in US Adults 71 Years and Older:
The National Health and Aging Trends Study. JAMA Ophthalmology 141(2):197–204
https://doi:10.1001/jamaophthalmol.2022.5840

Kurki, I., & Saarinen, J. (2004). Shape perception in human vision: Specialized
detectors for concentric spatial structures? Neuroscience Letters, 360(1-2), 100–102
https://doi.or/10.1016/j.neulet.2004.01.053

Liu J., Bex P.J., Jnawali A. & Vera-Diaz F.A. (2021) Evaluation of the novel
Foraging Interactive D-prime (FInD) CSF Test. Investigative Ophthalmology &
Vision Science 62(8):525. Available at:
https://iovs.arvojournals.org/article.aspx?articleid=2774587

Martínez-roda, J. A., Vilaseca, M., Ondategui, J. C., Aguirre, M., & Pujol, J. (2016)
Effects of aging on optical quality and visual function. Clinical and Experimental
Optometry 99(6):518–525. https://doi.org/10.1111/cxo.12369

McKendrick, A.M., Weymouth, A.E. & Battista, J. (2013) Visual Perception from Age
20 through 80 Years. Investigative Ophthalmology & Visual Science 54(3):1730-
1739 https://10.1167/iovs.12-10974

Merabet L.B., Manley C.E., Pamir Z., Bauer C.M., Skerswetat J. & Bex P.J. (2023)
Motion and form coherence processing in individuals with cerebral visual impairment.
Developmental Medicine & Child Neurology 65:1379–1386
https://doi.org/10.1111/dmcn.15591

Neupane S., Skerswetat J. & Bex P.J. (2024) Comparison of Foraging Interactive D-
prime and Angular Indication Measurement Stereo with different methods to assess
stereopsis. PLOS ONE 19(6):e0305036
https://doi.org/10.1371/journal.pone.0305036

Newsome, W. T., & Pare, E. B. (1988). A selective impairment of motion perception
following lesions of the middle temporal visual area (MT). Journal of Neuroscience,
8(6), 2201-2211.
643
644 Pei, F., Pettet, M. W., Vildavski, V. Y., & Norcia, A. M. (2005). Event-related
645 potentials show configural specificity of global form processing. NeuroReport, 16(13),
646 1427–1430 https://doi.org/10.1097/01.wnr.0000177003.12322.9b
647
Rasdall M.A., Cho C., Stahl A.N., Tovar D.A., Lavin P., Kerley C.I., Chen Q., Ji X.,
Colyer M.H., Groves L., Longmuir R., Chomsky A., Gallagher M.J., Anderson A.,
Landman B.A. & Rex T.S. (2024) Primary Visual Pathway Changes in Individuals
With Chronic Mild Traumatic Brain Injury. JAMA Ophthalmology
https://doi:10.1001/jamaophthalmol.2024.5076
Rizzo, J. R., Beheshti, M., Hudson, T. E., Mongkolwat, P., Riewpaiboon, W., Seiple,
W., Ogedegbe O.G. & Vedanthan, R. (2020) The global crisis of visual impairment:
an emerging global health priority requiring urgent action. Disability and
Rehabilitation: Assistive Technology, 18(3):240–245.
https://doi.org/10.1080/17483107.2020.1854876
Samanta A, Mauntana S, Barsi Z, Yarlagadda B, Nelson PC. (2023) Is your vision
blurry? A systematic review of home-based visual acuity for telemedicine. Journal of
Telemedicine and Telecare 29(2):81-90 https://doi:10.1177/1357633X20970398
Schmidtmann, G., Gordon, G. E., Bennett, D. M., & Loffler, G. (2013). Detecting
shapes in noise: tuning characteristics of global shape mechanisms. Frontiers in
Computational Neuroscience, 7(37), 1–14 https://doi.org/10.3389/fncom.2013.00037
664
665 Schmidtmann, G., Jennings, B. J., Bell, J., Kingdom, F.A.A. (2015). Probability, not
666 linear summation, mediates the detection of concentric orientation-defined textures.
667 *Journal of Vision*, *15*(16), 1-19 https://doi:10.1167/15.16.6
668
669 Serre, T., Kouh, M., Cadieu, C., Knoblich, U., Kreiman, G. & Poggio, T. (2005). A
670 Theory of Object Recognition: Computations and Circuits in the Feedforward Path of
671 the Ventral Stream in Primate Visual Cortex. AI Memo 2005–036/CBCL Memo 259.
672 Cambridge, MA: MIT. Available at: http://hdl.handle.net/1721.1/36407
673
674 Serre, T., Oliva, A., & Poggio, T. (2007). A feedforward architecture accounts for
675 rapid categorization. Proceedings of the National Academy of Sciences, USA,
676 104(15), 6424–6429 https://doi:10.1073/pnas.0700622104
677
678 Seu, L., & Ferrera, V. P. (2001). Detection thresholds for spiral Glass patterns.
679 Vision Research, 41(28), 3785–3790 https://doi.or/10.1016/s0042-6989(01)00235-8
680

Skerswetat J. & Bex P.J. (2022) AIM+(Angular Indication Measurement Plus)
enables rapid and self-administered assessment of visual perception dependency
across multiple stimulus dimensions; 44th European Conference on Visual
Perception (ECVP) 2022 Nijmegen, The Netherlands, Perception 51(1_Suppl):34
https://doi.org/10.1177/03010066221141167

Skerswetat J., Boruta A. & Bex P.J. (2022) Disability Glare Quantified Rapidly with
AIM (Angular Indication Measurement) Glare Acuity. Investigative Ophthalmology &
Vision Science 63(7):2558 – F0512 Available at:
https://iovs.arvojournals.org/article.aspx?articleid=2783354

Skerswetat J., He J., Shah J.B., Aycardi N., Freeman M. & Bex P.J. (2024a) A new,
adaptive, self-administered, and generalizable method used to measure visual acuity
Optometry and Vision Science 101:451–463
https://doi.org/10.1097/OPX.0000000000002160

Skerswetat J., Ross N.C., Idman-Rait C., Sun K., Wynn O. & Bex P.J. (2024b) Visual
Performance of People With Albinism Assessed With Generalizable and Adaptive
AIM and FInD Methods. Investigative Ophthalmology & Visual Science 65(11):34
https://doi.org/10.1167/iovs.65.11.34

Smith, M. A., Bair, W., & Movshon, J. A. (2002). Signals in macaque striate cortical
neurons that support the perception of glass patterns. Journal of Neuroscience,
22(18), 8334–8345 https://doi.org/10.1523/JNEUROSCI.22-18-08334.2002
701
702   Smith, M. A., Kohn, A., Movshon, J. A., & Movshon, J. A. (2007). Glass pattern
703   responses in macaque V2 neurons. Journal of Vision, 7(3):5, 1–15 Available at:
704   https://doi.org/10.1167/7.3.5
705
706   United Nations (2022) World Population Prospects 2022: Summary of Results.
707   United Nations. Available at:
708   https://www.un.org/development/desa/pd/sites/www.un.org.development.desa.pd/file
709   s/wpp2022_summary_of_results.pdf
710
711   Van Essen, D., Anderson, C., & Felleman, D. (1992). Information processing in the
712   primate visual system: An integrated systems perspective. Science, 255(5043), 419–
713   423 https://doi.or/10.1126/science.1734518
714
715   Wallis, S. A., Baker, D. H., Meese, T. S., & Georgeson, M. A. (2013). The slope of
716   the psychometric function and non-stationarity of thresholds in spatiotemporal
717   contrast vision. *Vision research*, *76*, 1-10.
718   https://doi.org/10.1016/j.visres.2012.09.019
719
Walter K, Bex P (2024) A novel, rapid, quantitative method for face discrimination.
PLOS ONE 19(12): e0315998 https://doi.org/10.1371/journal.pone.0315998

Waskom, M. L., (2021). seaborn: statistical data visualization. Journal of Open
Source Software, 6(60), 3021, https://doi.org/10.21105/joss.03021

Wilson, H., Wilkinson, F., (1998) Detection of global structure in Glass patterns:
Implications for form vision. Vision Research, 38, 2933–2947
https://doi.or/10.1016/S0042-6989(98)00109-6

Wilson, H., Wilkinson, F., & Asaad, W. (1997). Concentric orientation summation in
human form vision. Vision Research, 37(17), 2325–2330
https://doi.org/10.1016/s0042-6989(97)00104-1

World Health Organisation (2019) World report on vision. World Health Organisation.
Available at: https://www.who.int/publications/i/item/world-report-on-vision

World Health Organisation (2022) Report of the 2030 targets on effective coverage
of eye care. Available at: https://www.who.int/publications/i/item/9789240058002

World Health Organization (2023) Increasing eye care interventions to address
vision impairment: Technical Brief. World Health Organization. Available at:
https://www.who.int/publications/m/item/increasing-eye-care-interventions-to-
address-vision-impairment

Appendix

Figure A1 Left: Example data showing Box and Whisker plots with individual data (grey data points)
and the mean (black solid data point) for the AIM Form Coherence test (coherence thresholds are
expressed in %, 0-1). Right: Bland-Altman plots for AIM Form Coherence. Figures for all tests can be
found in the supplementary material (Bland_Altman_plots.pdf).

755   '

757 **Table A1** non-parametric Friedman ANOVA pairwise comparison (Durbin-Conover).
758 Statistically significant conditions are indicated by *.

Pairwise Comparisons (Durbin-Conover)		Statistic	p
FInD Form circ	FInD Motion HC circ	7.0448	<.001*
FInD Form HC hori	FInD Form HC exp	0.218	0.827
FInD Form HC hori	FInD Form HC circ	2.8615	0.004*
FInD Form HC hori	FInD Motion hori	0.1635	0.87
FInD Form HC hori	FInD Motion exp	1.022	0.307
FInD Form HC hori	FInD Motion circ	4.2923	<.001*
FInD Form HC hori	FInD Motion HC hori	3.2431	0.001*
FInD Form HC hori	FInD Motion HC exp	1.4308	0.153
FInD Form HC hori	FInD Motion HC circ	4.5784	<.001*
FInD Form HC exp	FInD Form HC circ	2.6435	0.008*
FInD Form HC exp	FInD Motion hori	0.3815	0.703
FInD Form HC exp	FInD Motion exp	0.804	0.422
FInD Form HC exp	FInD Motion circ	4.5103	<.001*
FInD Form HC exp	FInD Motion HC hori	3.4611	<.001*
FInD Form HC exp	FInD Motion HC exp	1.2127	0.226
FInD Form HC exp	FInD Motion HC circ	4.7965	<.001*
FInD Form HC circ	FInD Motion hori	3.025	0.003*
FInD Form HC circ	FInD Motion exp	1.8396	0.066
FInD Form HC circ	FInD Motion circ	7.1538	<.001*
FInD Form HC circ	FInD Motion HC hori	6.1046	<.001*
FInD Form HC circ	FInD Motion HC exp	1.4308	0.153
FInD Form HC circ	FInD Motion HC circ	7.44	<.001*
FInD Motion hori	FInD Motion exp	1.1855	0.236
FInD Motion hori	FInD Motion circ	4.1288	<.001*
FInD Motion hori	FInD Motion HC hori	3.0795	0.002*
FInD Motion hori	FInD Motion HC exp	1.5943	0.111
FInD Motion hori	FInD Motion HC circ	4.4149	<.001*
FInD Motion exp	FInD Motion circ	5.3143	<.001*
FInD Motion exp	FInD Motion HC hori	4.265	<.001*
FInD Motion exp	FInD Motion HC exp	0.4088	0.683
FInD Motion exp	FInD Motion HC circ	5.6004	<.001*
FInD Motion circ	FInD Motion HC hori	1.0492	0.294
FInD Motion circ	FInD Motion HC exp	5.7231	<.001*
FInD Motion circ	FInD Motion HC circ	0.2862	0.775
FInD Motion HC hori	FInD Motion HC exp	4.6738	<.001*
FInD Motion HC hori	FInD Motion HC circ	1.3354	0.182
FInD Motion HC exp	FInD Motion HC circ	6.0092	<.001*

You are searching for
faint versions of this:
Horizontal

NEXT