Dynamics of cortico-subthalamic neuronal patterns during dyskinesia in Parkinson's disease

J. Habets [1,2], T. Merk [1], V. Mathiopoulou [1], R. Lofredi [1,2], T. Binns [1,2,3,4], J. L. Busch [1,2], R. Köhler [1], A.

Memarpouri [1], K. Faust [5], P. Krause [1], GH. Schneider [5], WJ. Neumann [1,2,3], P. Tovote [6], A. Kühn [1,2,3,7,8]

1 Movement Disorders and Neuromodulation Unit, Department of Neurology, Charité –
Universitätsmedizin Berlin, Germany
2 Berlin Institute of Health (BIH), Berlin, Germany
3 Einstein Center for Neurosciences Berlin, Charité – Universitätsmedizin Berlin, Germany
4 Bernstein Center for Computational Neuroscience, Berlin, Germany
5 Department of Neurosurgery, Charité – Universitätsmedizin Berlin, Germany
6 Institute of Clinical Neurobiology, Universitätsklinikum Würzburg, Germany
7 NeuroCure Clinical Research Centre, Charité – Universitätsmedizin Berlin, Germany
8 Berlin School of Mind and Brain, Humboldt-Universität zu Berlin, Berlin, Germany

Abstract

People with Parkinson's disease suffer from levodopa-induced dyskinesia, adversely to chronic
dopaminergic medication. These involuntary movements affect quality of life and are linked to
lowered motor inhibition in the cortico-basal-ganglia motor network during hyperdopaminergic
states. Several cortical and subcortical spectral biomarkers are associated with dyskinesia, but the
oscillatory patterns associated with the behavioral complexity of involuntary movement and
voluntary movement suppression that defines dyskinesia, is not studied so far. To elucidate these
spatio-temporal dynamics and its behavioral relevance, we studied cortico-subthalamic oscillations in
21 Parkinson's patients during a dyskinesia-provoking experiment. We differentiated between non-
dyskinetic and dyskinetic periods, and split data based on a kinematic movement registration, leading
to respectively resting states and voluntary movement, and active movement suppression and
dyskinetic movement. Elevated theta-activity and attenuated beta-activity was found in the
subthalamic nucleus during voluntary movement suppression within dyskinetic periods, while
cortico-subthalamic gamma-activity was only elevated during dyskinetic movement presence.
Subthalamic spectral changes significantly predicted dyskinesia presence, and the amount of
registered movement significantly affected the predictive value of dyskinesia predictions. We
propose a dyskinetic macrostate to consist of distinct behavioral microstates of dyskinetic
movements versus movement suppression, both associated with distinct oscillations in the cortico-
subthalamic motor network.

Main

Dopaminergic replacement therapy is the cornerstone to treat hypokinetic symptoms in Parkinson's disease (PD), but also adversely causes disabling abnormal involuntary movements, defined as levodopa induced dyskinesia (LID) [1–4]. As disease progression continues and requires larger and more frequent dopaminergic intakes, LID occurs in up to 80% of PD patients, impacting quality of life [1,3,5]. LID primarily emerges during moments of dopaminergic peaks and has a fluctuating heterogeneous clinical phenotype of involuntary movements, alternating with episodes of voluntary motor suppression and subjective internal motor restlessness [2]. Typically, LID is treated with either pharmacological interventions that improve levodopa's efficacy and allow dopamine reduction [4] or target specific striatal projecting neurotransmitter-receptors [6]; or with device-aided therapies such as deep brain stimulation (DBS) [5,7–10]. Although overstimulation with DBS can induce dyskinesia [11], DBS generally alleviates LID by allowing a reduction of dopaminergic medication [12] and possibly by a long-term desensitization of dopamine receptors [13].

Current understandings of the pathophysiology of LID hypothesize that a reduced cortical long-term depotentiation alters cortico-striatal synaptic plasticity [14,15]. The impaired synaptic plasticity fails to balance dopaminergic cortico-basal-ganglia excitation and increases the likelihood of movement [16]. On a striatal level, chronic dopamine depletion is hypothesized to cause compensatory structural changes in striatal projection neuron architecture and fast spiking interneurons [17–19]. These compensatory neural changes shift the balance between the canonical direct and indirect motor pathways to restore homeostasis during hypo-dopaminergic periods [17,18,20,21]. Moreover, long term dopaminergic depletion leads to abnormal spiking patterns of SPN neurons with loss of spatial and temporal selectivity [22]. Neural dynamics can partially be restored by dopaminergic treatment, but during hyper-dopaminergic periods, inverse abnormalities in striatal activity pattern with overcompensated reduction in motor inhibition may occur, facilitating LID [22–26]. Since the subthalamic nucleus (STN) is specifically involved in movement planning, execution, and cancellation [27–29], STN DBS can directly modulate physiological motor stopping [30], but also dyskinetic movement [11]. Invasive neurophysiological studies explored biomarkers that represent spectral network oscillations during different motor, cognitive, and emotional states in PD [31]. Subthalamic beta activity has been established as a symptom-specific biomarker for bradykinesia [28], independent of disease [32,33]. Furthermore, cortico-subthalamic gamma activity has been used to decode movement initiation, velocity and stopping [34–36], whereas increased theta band activity has been described in conflicting cognitive tasks during decision making and in impulse control disorders [37,38]. Regarding LID, elevated subthalamic and cortical gamma-band (60 – 90 Hz) activity has been associated with LID presence and severity [39–41], similar to elevated subthalamic theta- and alpha-band (resp. 4 – 8 and 8 – 12 Hz) activity [37], and decreased subthalamic beta-band (13 – 35 Hz) activity [40,42]. The spatiotemporal patterns of oscillations associated with LID so far, have not been differentiated with respect to the highly variable behavioral dynamics of LID [2], i.e. movement execution and movement suppression, and therefore their behavioral relevance within LID remains unclear [31]. This highly variable behavioral pattern within LID, containing both presence and absence of movement execution [2], may modulate neural oscillations by the degree of successful active motor suppression [43]. In this vein, we hypothesize that LID in PD patients consists of distinct behavioral states, with distinct neural oscillations [43], in line with a first rodent study that differentiated between distinct dyskinetic movements in rodents with distinct neural patterns [21].

A recent concept developed to explain integrated systemic, cardio-behavioral states in the context of emotional challenge [44] allows to characterize temporal dynamics of behavioral and physiological changes over variable time-periods [44]. Herein, slow transitions are termed as macrostate changes and contrast fast-changing, momentary microstates, which reflect short-lasting distinct behavioral or

neural patterns.
To gain insight into the behavioral relevance of LID-related biomarkers, we hypothesize that the
heterogeneous behavior during dyskinetic periods can be explained by distinguishable neural and
behavioral states that differ in spatio-temporal spectral characteristics. First, a prerequisite for LID to
occur is the slow transition from the hypodopaminergic OFF state to the hyperdopaminergic ON
state in PD, which we consider a macrostate change [44]. On the background of the hyperdopaminergic
ON state, fast-changing microstates may occur that lead to the execution of dyskinetic movements.
To investigate the hypothesized states-concept, we characterized spectral patterns in invasive
subthalamic and cortical neurophysiological recordings from the hypodopaminergic OFF state to the
hyperdopaminergic ON state with LID, during the execution of dyskinetic movements and during
voluntary movement suppression. The latter behavioral states we defined based on the presence and
absence of movement execution during a dyskinetic state. We identified distinct subthalamic theta
and beta oscillations associated with the slow-changing dopaminergic periods considered as a
macrostate. Fast-changing patterns of cortico-subthalamic gamma oscillations that depend on the
presence of dyskinetic movement could be defined as microstate changes. The identification of the
behavioral relevance of these spatio-temporal spectral changes during dyskinetic periods will help us
to better understand LID's pathophysiology and cortico-subthalamic biomarkers relevant for motor
control during LID. This may lead to optimized use of biomarkers for adaptive DBS tailored to clinical
demands.

Dyskinetic periods contain moments with and without objectified movement execution

To investigate LID-related spectral patterns, we included twenty-one PD patients implanted with
bilateral STN DBS electrodes, of which thirteen received a temporary unilateral electrocorticography
(ECoG) electrode (mean age: 61 years, standard deviation (sd): 9.2 years, mean disease duration: 10
years (sd: 4 years)) (see Table S1). All participants performed a dyskinesia-provoking experiment
during the interval (two to seven days) between their surgical implantation of DBS-electrodes and its
connection to the internal pulse generator [45]. Participants started the experimental protocol in the
medication-OFF condition (at least 12 hours of dopamine withdrawal) followed by a supra-threshold
dosage of fast-acting levodopa (circa 1.5 morning dose of Madopar LT ®, Roche Pharma). STN and
cortical activity were recorded during repetitive rest and unilateral self-paced hand tapping tasks,
intermingled with unscripted pauses to avoid fatigue and experiment abortion (see Figure 1 and Fig.
S1).
A clinical movement disorders specialist assessed the presence and severity of LID based on video
recordings, using the Clinical Dyskinesia Rating Scales (CDRS). [46] The presence of LID was observed in
eighteen out of twenty-one participants (86%), of which fifteen had a preoperative history of LID
(83%) (see Methods). One participant, that did not develop LID, had a preoperative LID history. The
eighteen patients that did develop LID had mean peak dyskinesia severities of 5 CDRS points (sd: 3,
range: 2 to 13 CDRS points), out of a maximum of 28 points (the latter corresponds to severe
dyskinetic movement in all seven rated body parts).
The distributed repetitions of resting and tapping tasks blocks led to the recording of movement
presence and movement absence in both dyskinetic states and non-dyskinetic states (Figure 1). We
defined a dyskinetic period as a time interval that starts at the first observed dyskinetic movement
and prolongs as long as the patient regularly shows abnormal involuntary movements. Dyskinetic
periods will also contain moments of suppression of involuntary movements, usually achieved by
active movement suppression with voluntary muscular co-contraction in between dyskinetic
movements (Figure 2). Fluctuations in the clinically assessed severity of a dyskinetic period were
assessed on a minute level, while the observation of actual movement was objectively assessed on a

sub second level with accelerometer data. Consequently, we differentiated between 1) movement
presence during the non-dyskinetic period (i.e. voluntary movements), 2) movement absence during
the non-dyskinetic period (i.e. the resting state), 3) movement presence during the dyskinetic period
(i.e., execution of both voluntary and involuntary movements, during episodes clinically confirmed as
dyskinetic), and 4) movement absence during the dyskinetic period (i.e., epochs of motor
suppression, lacking objectified movement execution, within periods clinically confirmed as
dyskinetic). This behavioral concept, that dyskinetic periods contain epochs of both movement
presence and absence, is essential to assess different neural microstates, i.e. dyskinetic movements,
during the hyperdopaminergic macrostate characterized by a dyskinetic period.

Figure 1: Single subject example of spectral changes and behavioral heterogeneity during dyskinetic periods. Neural, behavioral, and clinical data from one subject is shown. The first upper row (ACC) represents the participant's activity,
recorded via bilateral
accelerometers placed on the index-fingers. The line shows the average of the z-scored root mean squares of
both accelerometers. The second upper row (LID) represents the clinically assessed dyskinesia presence and
severity (defined as sum CDRS) over time. The first white part represents the non-dyskinetic period. A
dyskinetic period is represented by different shades of orange, where darker orange represents more severe
LID. LID occurred bilateral in this subject, with more severe dyskinetic movements in the right body side
compared to the left. The third upper row represents the task performed (TASK) during the experiment, in
which *rest* contained seated rest, *tap* contained a seated self-paced, unilateral hand-tapping task, and *pause*
contained an unscripted pause for the participant to prevent fatigue.
The three neural time-frequency plots represent spectral unilateral cortical (ECoG) and bilateral subthalamic
(STN) activity. Note the decrease in beta activity bilaterally in the STN at about 10 minutes after levodopa
intake and during LID, and irregular gamma synchronization during LID, mostly in the presence of movement.
*: The recording was interrupted around minute 45 for a toilet break and the x-axis is adjusted accordingly.
ECoG: electrocorticography, Hz: Hertz, L-Dopa: levodopa, LID: levodopa induced dyskinesia, STN: subthalamic
nucleus.
165
166

Multi-frequency cortico-subthalamic oscillatory patterns reflect dopamine-induced dyskinetic macrostate changes

Time-frequency analyses of the subthalamic and cortical recordings demonstrated changes in spectral patterns with levodopa-intake in different frequency ranges of interest (see the exemplary case in Figure 1). Spectral activity was calculated using an unbiased data-dimensionality reduction method (spatio-spectral decomposition) on the multi-channel recordings and resulted in single time-frequency data per recording site [47]. These data are optimized independently regarding signal-to-noise ratios per frequency band of interest (theta, beta, and gamma band; see Methods). We hypothesized that distinct neural oscillatory patterns occur during hyperdopaminergic ON periods and correlate with the severity of dyskinetic periods. To this end, we first descriptively compared local STN and cortical changes occurring with the transition from OFF to ON dopaminergic periods and during different LID severity grades. All neural data from the rest and tap task were included, without specifying movement presence and absence. CDRS raters were blinded for neurophysiological results and scores were categorized into mild (less than 4 points), moderate (4 to 8 points), and severe LID (8 or more points) [46]. Power spectral densities calculated over 10-second epochs showed an increase in subthalamic theta activity that occurred during the dopaminergic ON period without LID and gradually increased with LID-severity, whereas subthalamic low-beta activity showed a gradual decrease. Similarly, this low-beta decrease extended the low-beta decrease observed in both all non-dyskinetic periods as well as in ON-periods periods without LID after 30 minutes post-levodopa-intake (Figure 2, upper row). A large increase in narrow band gamma activity was observed in both subthalamic and cortical data predominantly during the dyskinetic periods but did not show a strong scaling effect with LID severity. Additionally, we calculated squared coherences in the same epochs to investigate how LID affects spectral connectivity between the two STNs and between the cortex and the ipsilateral STN. Inter-subthalamic coherence showed small and variable increases in theta-coherence and decreases in beta-coherence during the dyskinetic period but no obvious scaling effect (Figure 2, lower left). Gamma-activity was coherent between the cortex and the ipsilateral STN only in the two patients with severe LID.

Together, these spectral changes represent distinct neural patterns that represent medication induced macrostate changes: importantly, we describe the differences not only for the dopaminergic transition to the ON medication state without LID, but the dyskinetic state demonstrates a further spectral modulation.

Figure 2: Spectral subthalamic and cortical patterns change during dyskinetic periods depending on dyskinesia severity.

All four panels show the changes in spectral patterns as percentage changes relative to the dopaminergic OFF condition at rest up to five minutes after levodopa-intake. Three grades of dyskinesia severity are visualized: mild LID in yellow (1 to 3 CDRS points), moderate in red (4 to 7 CDRS points), and severe in purple (8 or more CDRS points). The OFF and ON conditions are visualized in light and dark green. The numbers in the legend represent the number of subjects that contributed neural data to a category.

The upper row shows power spectral densities of subthalamic nucleus (STN) (upper left) and cortical (upper right) recordings. Note the increase in STN theta and decrease in beta activity that scale with severity of dyskinesia, the scaling effect is less prominent for gamma synchronization with dyskinesia in STN and cortex. Spectral powers are calculated using Welch's method per ten-second windows and the panels show means and standard error of the means for individual averages per category. Spectral powers of bilateral STNs are averaged per window. The lower row shows squared coherences between the two STNs (lower left) and between the cortex and the ipsilateral STN (lower right) and demonstrates a decrease in beta coherence and increase in cortico-STN gamma coherence during LID, with less prominent changes. Squared coherence spectra are calculated in ten-second epochs using Welch's method. All lines were smoothed with a 3 Hz bin for visualization.

Hz: Hertz, LID: levodopa-induced dyskinesia, OFF: medication Off, STN: subthalamic nucleus, sq.: squared.

Subthalamic theta- and beta-oscillations are associated with movement absence in between dyskinetic movements

The observed changes in oscillatory cortical and subcortical neuronal pattern have been associated with LID presence in prior studies [37,40,42]. To explore the behavioral relevance of these long-lasting, dopamine-dependent macrostate changes (Figure 2) and their interaction with short-lasting movements, we split the neural data based on the presence and absence of movement. We first

defined dyskinetic versus non-dyskinetic periods based on the clinical expert assessments of video
recordings of the full protocol. Dyskinetic periods were identified as behavioral states that lasted at
least several minutes, and did not change on a (sub)-second resolution. Within these periods (i.e.,
macrostates), movement presence and absence were defined using bilateral index-finger worn
accelerometer data (TMSi Saga®, TMSi International, Oldenzaal, The Netherlands). Movement
absence was verified by video protocol to exclude movements on other parts of the body. Movement
start- and end-points were defined on a sub-second resolution based on a thresholding procedure
and led to epochs of movement presence and absence (i.e., microstates). Combining the sub second
movement detection with the clinical assessments of LID presence, resulted in the categorization of
all neurophysiological data into the four mentioned behavioral categories, i.e. movement presence
versus absence, and dyskinetic versus non-dyskinetic periods (Figure 3). We did not differentiate
between voluntary and involuntary movements during dyskinetic period, since these two concepts
are not reliably distinguishable. During dyskinetic periods, patients often try to mask involuntary
movements with a voluntary movement, and voluntary movements vice versa often evolve into or
provoke involuntary movements. All data were merged within the respective clinically relevant
behavioral categories to identify the distinct neural pattern and its modulation during microstates of
movement presence. Additionally, we defined an ON condition without LID, i.e., time points after 30
minutes post-levodopa-intake without LID (n = 15), to allow comparison of dyskinetic effects versus
non-dyskinetic dopaminergic medication effects. Neural data were analyzed per recording source,
i.e. per STN or ECoG electrode, using the first component of an un-biased spatial spectral
decomposition method to reduce the data-dimensionality while optimizing the signal-to-noise ratio
[47]. Neural data were merged within their respective microstate clusters for LID- and movement-
presence and underwent time-frequency decompositions allowing for spectral comparisons per
canonical frequency band of interest (Figure 4) (see Methods).

Figure 3: Schematic overview of dyskinesia-related spectral patterns during the presence and absence of executed dyskinetic movement.
This schematic visualization illustrates that the clinical manifestation of both dyskinetic and non-dyskinetic periods contain moments of movement absence and presence. The upper line plot visualizes the rising dopamine blood-levels after medication intake. In the hyperdopaminergic state, dyskinetic periods occur when levodopa-levels are elevated sufficiently. Green PSD panels (left) represent non-dyskinetic periods, classified as a total Clinical Dyskinesia Rating Scale of zero at all time points. Purple PSD panels (right) represent dyskinetic periods. Dyskinetic periods fluctuate slowly over time with typically durations in a minute to half hour range, depending on individual disease progression, pharmacodynamics, and levodopa dosage.

Movement presence and absence fluctuate
more frequently and faster, on a seconds-resolution.
Green PSDs represent cortical recordings and yellow PSDs represent subthalamic recordings. Spectral bands are
shown on the x-axes: LF: low-frequency activity (theta and alpha, 4 – 12 Hz), beta activity (13 – 35 Hz), gamma
activity (60 – 90 Hz). The y-axes represent the magnitude of spectral powers. LF: low-frequency, min: minutes.
Spectral elevations and attenuations in this graphic are based on the spectral results, but the heights of the
peaks are relative and not absolute.
286

Spectral power densities and squared coherences were compared between the behavioral
categories, within every frequency band of interest (Figure 4 and 5). First, we describe changes in
neuronal network activity that occur **within dyskinetic periods in the absence of movement**
execution. *Subthalamic theta and alpha band increase and low-beta band reduction* were
significantly larger in the dyskinetic period compared to the non-dyskinetic periods (resp. coefficients
of linear mixed effect models (lmm-coeff.) 0.17, p=0.0001, 0.11, p<0.0001 and -0.14, p<0.0001)
(Figure 4B). Importantly, these LID-related changes extended changes occurring during the ON
without LID condition. When statistically testing the dyskinetic period versus the ON condition,
subthalamic low-beta decrease and theta and alpha increase remained significant (resp. lmm-coeff.: -
0.07, p<0.0001; 0.09, p<0.0001; and 0.09, p<0.0001). A significant low-beta reduction also occurred
in the ON versus the OFF condition, this was however a smaller effect size than the dyskinetic-related
changes (lmm-coeff. -0.11, p>0.0001). Theta and alpha increases did not differ between ON without
LID and OFF conditions (lmm-coeff 0.10, p=0.20, 0.08, p=0.03).
Both individual peak gamma-activity, as well as broad-gamma activity, showed a small but significant
increase during dyskinetic periods compared to all non-dyskinetic periods (lmm-coeff. resp: 0.03,
p<0.0001; 0.03, p<0.0001). However, these increases did not remain significant when comparing the
dyskinetic periods with the non-dyskinetic ON condition (lmm. coeffs.: 0.00, p=.98 and -0.01, p=0.04)

but were significant in comparison with the OFF conditions (resp. lmm-coeff.: 0.03, p=0.001; 0.03,
p<0.0001).

Regarding cortical power modulation during dyskinetic periods in movement absence, we observed
small but significant low-beta increase (lmm-coeff. 0.04, p=0.0015) and gamma-activity decreases
(peak- and broad-gamma resp. lmm-coeff. -0.02 and -0.08, p<0.0001 and p<0.0001) (Figure 4, lower
panels). These cortical changes did not remain significant when comparing the dyskinetic periods
with the non-dyskinetic ON condition (resp. lmm.-coeffs.: 0.04, p=0.03; 0.00, p=0.87; and -0.01,
p=0.28). We cannot exclude that these small changes are overemphasized following the
normalization influenced by the changes during movement presence that are in higher magnitudes of
order.

In line with the observed STN power patterns, also *inter-subthalamic coherences significantly*
increased in the theta band and decreased in the low-beta band in the absence of movement during
dyskinetic periods compared to non-dyskinetic periods (Figure 5, upper row, resp. lmm-coeff. 0.35,
p<0.0001 and -0.08, p<0.0001). This LID-associated theta and low-beta modulation remained
significant compared to ON periods without LID (lmm.-coeffs.: 0.36, p<0.001 and -.08, p<0.001) but
no difference occurred between ON and OFF conditions suggesting that the theta- and beta-
coherence modulation is specific to the dyskinetic periods (lmm-coeff. resp. 0.00, p=0.935 and 0.00,
p=0.855). Inter-subthalamic peak-gamma showed a minor but significant difference between
dyskinetic and non-dyskinetic periods (lmm-coeff. 0.02, p=0.0004), whereas alpha and broad-gamma
bands did not differ significantly (lmm-coeff. resp. -0.01, p=0.74; 0.01, p=0.08).
Cortico-subthalamic theta- and beta-coherences did not change significantly between dyskinetic and
non-dyskinetic periods in the absence of movement (lmm-coeff. resp. 0.06, p=0.02; and 0.01,
p=0.79), whereas alpha activity showed a trend for an increase (lmm-coeff. 0.08, p=0.001). C*ortico-*
subthalamic gamma coherences in peak- and broad-frequencies showed a small but significant
increases in the dyskinetic periods compared to non-dyskinetic periods (Figure 5 lower row, lmm-
coeff. resp. 0.04, p<0.0001, and 0.06, p<0.0001) that remained increased compared to ON conditions
(lmm-coeffs. resp. 0.04, p<0.001, and 0.06, p<0.001).

Taking together, these results demonstrate that movement absence during dyskinetic periods is
characterized by increased subthalamic theta and alpha activity and decreased beta activity that
significantly differ from the non-dyskinetic ON condition and thus may represent a distinct
macrostate (Figures 4 and 5). In contrast, the small STN gamma power increase was more related to
a general dopamine ON condition and only the increase in cortico-STN coherence was associated
with the dyskinetic period.

Dyskinetic movements are characterized by increased gamma activity in STN and cortex

To investigate whether **movement presence during dyskinetic periods** can be identified and
objectified as a **microstate** in neural recordings, we isolated movements during dyskinetic periods
based on accelerometer data. To ensure that network pattern changes during movement presence
are specific for dyskinetic movements, we compared them with movement presence during non-
dyskinetic periods, extracted from the voluntary tapping task. Subthalamic theta and alpha activity
showed a trend for an increase during dyskinetic movement compared to all recorded non-dyskinetic
movement (Figure 4 right upper panel, lmm-coeff. resp. 0.16, p=0.002 and 0.06, p=0.035). Low-beta
activity decreased significantly during dyskinetic movement presence compared to non-dyskinetic
movement presence (lmm-coeff. -0.15, p<0.001). Subthalamic broad-gamma activity, but not peak-

gamma activity, increased significantly during dyskinetic movement compared to non-dyskinetic voluntary movements (resp. lmm-coeff. 0.03, p=0.0001; 0.01, p=0.044). Similarly, cortical broad-gamma activity significantly increased during dyskinetic movement compared to non-dyskinetic movement (lmm-coeff. 0.13, 0.0012, Figure 4, upper right panel). Cortical theta, beta, and peak-gamma activity did not show significant changes during dyskinetic versus non-dyskinetic movements. To examine that these changes are specific for movement during dyskinetic periods, instead of merely associated with movement during the dopaminergic ON state, we isolated all voluntary movements that occurred in non-dyskinetic states more than 30 minutes after levodopa-intake (n = 9 subjects). In line with the assumption that gamma activity codes motor parameter, peak-gamma activity showed a significant increase during dyskinetic movement compared to ON medication movement (lmm-coeff. 0.08, p=0.011), whereas theta, low-beta, and broad-gamma did not (lmm-coeffs. resp. 0.03, -0.05, and 0.0, p all > 0.12). Due to limited data availability of ON medication movement data, we only compared these main 4 frequency bands and could not assess cortical spectral changes.

Inter-subthalamic coherence during dyskinetic periods showed similar changes during movement presence as mentioned during movement absence. Theta coherence increased and low-beta coherence decreased during movement in dyskinetic periods, although only the latter reached statistical significance (lmm-coeff. resp. 0.20, p=0.006 and -0.13, p<0.001). Inter-subthalamic gamma coherence did not change significantly. Although larger cortico-subthalamic gamma coherence was observed during dyskinetic movements, none of the changes reached statistical significance, possibly due to high inter-individual variance (Figure 5, lower right).

In order to characterize the movement-related microstate changes during dyskinesia, we lastly compared the neural patterns during movement absence and presence within the dyskinetic period. Here, we found both subthalamic and cortical peak-gamma to be significantly higher during dyskinetic movement presence compared to dyskinetic movement absence (resp. lmm-coeff. 0.08, p<0.0001; and 0.24, p<0.0001). Also, cortical broad-gamma activity showed a significant increase during dyskinetic movement presence compared to dyskinetic movement absence (lmm-coeff. 0.18, p<0.001). Subthalamic broad-gamma activity however, showed a small but significant decrease in dyskinetic movement presence compared to dyskinetic movement absence (lmm-coeff. -0.02, p=0.0001). We argue that this small coefficient in the opposite direction as hypothesized, can be explained by the parallel increases in broad gamma during both dyskinetic movement absence and presence compared to non-dyskinetic periods. Modulations in cortical theta were significantly higher during dyskinetic movement presence versus absence (lmm-coeff. 0.3, p<0.001), whereas subthalamic theta and both subthalamic and cortical beta activity did not differ significantly (lmm-coeff. resp. 0.02, p=0.04; -0.01, p=0.20; and 0.06, p=0.03).

Together, when comparing movement presence with movement absence, both during dyskinetic periods, dyskinetic movement is characterized by peak-gamma increases in the STN and the cortex, and broad-gamma increases in the cortex. Subthalamic broad-gamma already shows an increase during dyskinetic periods in movement absence, which even slightly extends the increase during movement presence. These findings demonstrate that the hypothesized cortical gamma activity related to LID is specific to movement presence during LID, and that increased activity in cortical gamma as well as subthalamic peak-gamma extend gamma modulation during non-dyskinetic, dopaminergic-ON movement presence.

Figure 4: Movement modulates local subthalamic and cortical patterns during dyskinetic periods

Overall, theta increases and beta decreases in the STN were significant larger during dyskinetic states in both movement absence and presence compared to non-dyskinetic states. Increased gamma activity in the STN during dyskinetic states was mainly observed during the dyskinetic microstates with movement presence.

The left four panels show power spectral densities (PSD) of subthalamic recordings (upper row) and cortical recordings (lower row). The upper and lower left PSD-panels show microstates of movement absence, upper and lower right PSD-panels show microstates of movement presence. Movement presence was assessed based on accelerometer-data according to a thresholding procedure. All green lines represent the mean over all individual average-PSDs during non-dyskinetic periods, whereas purple lines represent the dyskinetic periods. Only during movement absence, we differentiated the OFF and ON conditions within non-dyskinetic periods with sufficient sample sizes (light and dark green line respectively). Shadings show the standard error of the mean of the individual averages and legends show the number of included subjects per category.

The two boxplots in the right panel visualize the mean spectral power values focused on the main spectral bands of interest (theta, low-beta, and gamma), originating from the same data as shown in the four PSD panels. For visualization purposes we decided to show these 4 main frequency bands of interest. Green and purple boxes represent respectively non-dyskinetic and dyskinetic periods. The diagonally striped backgrounds indicate movement presence data, whereas movement absence data do not have a background shading. Statistically significant differences between non-dyskinetic and dyskinetic periods, either within movement absence data or within movement presence data, are visualized as full-colored purple boxes. Transparent purple boxes indicate no significant difference compared to a neighboring green box. Alpha-values for significance testing are 0.01 and corrected for six multiple comparison (alpha is 0.0016).

Hz: Hertz, LID: levodopa induced dyskinesia, min: minutes after levodopa-intake, n: number of unique subjects included in category, STN: subthalamic nucleus. Spectral band symbols: θ: theta (4 – 8 Hz), α: alpha (8 – 12 Hz, not shown), β-low: low-beta (12 – 20 Hz), β-high: high-beta (20 – 30 Hz, not shown), γ-peak: individual gamma peak (5 Hz bin around peak between 60 and 90 Hz), and γ-broad: broad-gamma (60 – 90 Hz).

Figure 5: Cortico-subthalamic connectivity changes during dyskinetic periods

The left four panels show squared coherences between bilateral subthalamic recordings (upper row) and ipsilateral cortico-subthalamic recordings (lower row). The upper and lower left coherence-panels show movement absence, the upper and lower right coherence-panels show microstates of movement presence. All green lines represent the mean over all individual average- coherences during non-dyskinetic periods, whereas purple lines represent the dyskinetic periods. Only during movement absence, we could differentiate the OFF and ON conditions within non-dyskinetic periods with sufficient sample sizes (light and dark green lines, respectively). Shadings show the standard error of the mean of the individual averages and legends show the number of included subjects per category. Squared coherences are calculated based on one-second epochs using power spectral densities (psd) and cross spectral densities (csd) by Welch's method (|csd(source-1 and source-2)|^2/ (psd(source-1) * psd(source-2)), see Methods).

The two boxplots in the right panel visualizes the mean spectral power values focused on the main spectral bands of interest (theta, low-beta, and gamma), originating from the same data as shown in the four coherence panels. Green and purple boxes represent respectively non-dyskinetic and dyskinetic periods. The diagonally striped backgrounds indicate movement presence data, whereas movement absence data do not have a background shading. Statistically significant differences between non-dyskinetic and dyskinetic periods, either within movement absence data or within movement presence data, are visualized as full-colored purple boxes. Transparent purple boxes indicate no significant difference between a neighboring green box. Alpha-values for significance testing are 0.01 and corrected for six multiple comparison (alpha is 0.0016).

Hz: Hertz, LID: levodopa induced dyskinesia, min: minutes after levodopa-intake, n: number of unique subjects included in category, STN: subthalamic nucleus. Spectral band symbols: θ: theta (4 – 8 Hz), α: alpha (8 – 12 Hz), β-low: low-beta (12 – 20 Hz), β-high: high-beta (20 – 30 Hz), γ-peak: individual gamma peak (5 Hz bin around peak between 60 and 90 Hz), and γ-broad: broad-gamma (60 – 90 Hz). Coherence plots on the left were smoothed with a 3 Hz bin for visualization.

Dyskinetic patterns lateralize contralateral in the cortex but spread bilateral in the STN

So far, we have observed relevant changes in spectral oscillatory activities during dyskinetic periods in both the absence and presence of movement (Table 1 and Figures 4-5). To better understand the pathophysiological role of these LID-related changes within the cortico-basal-ganglia-motor network, we investigated their laterality with respect to clinically observed LID. For this sub analysis, we selected all dyskinetic periods of unilateral LID. Unilateral LID was defined as a sum CDRS of 1 or higher for one body side, while the sum of the other body side remained 0 CDRS points. Typically, unilateral LID marks the beginning of a dyskinetic period and often occurs at the more affected side of the body. Sixteen subjects experienced unilateral LID, of which eleven received a unilateral ECoG-electrode. This led to cortical recordings from five ipsilateral and six contralateral hemispheres with respect to the dyskinetic body side and subthalamic recordings from sixteen bilateral hemispheres. Ten-second epochs during unilateral dyskinetic periods containing both movement presence and

absence were selected to ensure a sufficient sample size.

STN recordings showed bilateral theta and gamma power elevation and beta power reduction during
unilateral LID (Figure 6, left panel). In contrast, cortical recordings only showed narrow gamma
activity in hemispheres contralateral to the dyskinetic body side (Figure 6, right panel). Cortical
recordings ipsilateral to the dyskinetic body side displayed an elevated mean gamma activity without
a distinct peak. Modulation of cortical theta and beta activity did not show a consistent pattern. We
suggest this to be due to the limited number of included subjects per group and the lack of
movement selection. The observed results suggest that the specific subthalamic oscillatory activities
during dyskinetic periods occur bilateral, whereas cortical LID-related oscillatory activity occurs only
contralateral to the dyskinetic body side. Based on these findings, we excluded all cortical recordings
during unilateral dyskinetic periods, originating from the hemisphere ipsilateral to the unilateral
dyskinetic body side for further analyses.

Figure 6: Lateralization of spectral patterns during unilateral dyskinetic periods

Percentage-changes in power spectral densities during clinical observations of unilateral dyskinetic periods are
shown in subthalamic nucleus (STN) data (left panel) and cortical data (right panel). Subthalamic theta and
gamma elevations and beta-reductions occur in both contra- and ipsilateral data, while cortical gamma
elevations only occur in the contralateral hemisphere opposed to the dyskinetic body side. All data result from
Welch-transformations per ten-second epochs, without objectified movement selection. All percentage-
changes are relative to the medication "OFF" state, defined as all non-dyskinetic-moments within the first five
minutes after levodopa-intake, in the absence of movement, objectified with bilateral index-finger
accelerometers. The blue lines represent spectral power changes in the hemispheres ipsilateral to the
dyskinetic body side; sand-color lines represent spectral power changes in the hemispheres contralateral to the
dyskinetic body side. Due to the presence of bilateral STN-electrodes in all sixteen subjects with unilateral
dyskinetic periods, and only unilateral ECoG-electrodes in the eleven subjects with unilateral dyskinetic periods
and an ECoG-electrode, sample sizes differ per visualized group. We smoothed all lines with a 3 Hz bin for
visualization.
Hz: Hertz, LID: levodopa-induced dyskinesia, OFF: medication Off, STN: subthalamic nucleus.

Theta- and gamma-increase and beta-decrease scale with dyskinesia severity

We demonstrated how neuronal patterns during dyskinetic periods differ regarding the absence or
presence of movement. In this sub-analysis, we investigated the dependence of LID-related
oscillatory activity on the severity of the dyskinetic periods. For this, we included the spectral powers
and coherences originating from all three experimental tasks (i.e., resting states, tapping, and
unscripted pauses, Figure 1) and quantified movement presence per ten-second epoch. Movement
was quantified as mean root mean squares (ACC-RMS) of bilateral accelerometer data and ACC-RMS
values were individually z-scored. LID severities were expressed as total sum CDRS scores.

Subthalamic theta-, beta-, and gamma-activity changed gradually with increasing LID severity
(horizontal gradients in Figure 7, left column). The gradual theta increases and beta decreases
occurred over the whole range of movement presence, showing no clear gradient related to the
amount of movement presence (Figure 7, left, upper two panels). In contrast, gamma increases
during more severe LID did show a gradient depending on movement presence. During low
movement presence (i.e., ACC-RMS below 0) gamma activity only increased at high LID severities
(i.e., 8 CDRS points), whereas gamma activity during higher movement presence (i.e., ACC-RMS
above 0) showed increases already at lower LID severities (i.e., 4 CDRS points). These findings
confirm that the LID-severity-related subthalamic gamma elevations are movement dependent
whereas subthalamic theta and beta attenuations are movement independent (Figure 7, left lowest
panel).
Cortical spectral patterns demonstrated similar but smaller associations with the severity of the
dyskinetic period, as expected based on the cortical patterns during the dyskinetic period regardless
of LID severity (Figure 2 and 4). Cortical theta activity increased and beta activity decreased gradually
with increasing LID severity, without difference between the amounts of movement presence (Figure
7, right upper panel). Similar to the subthalamic findings, elevations in cortical gamma activity were
only seen in mild dyskinetic periods (i.e., up to 4 CDRS points) with higher movement presence (i.e.,
ACC-RMS above 0). Cortical gamma in more severe dyskinetic periods (i.e., more than 4 CDS points)
occurred also with less movement presence (Figure 7, right lower panel).
The observed inter-subthalamic coherences demonstrated similar patterns as the local subthalamic
spectral powers (Figure S3, left column). Inter-subthalamic theta- and gamma-coherences showed a
gradual increase with increasing severity of the dyskinetic period. In addition, inter-subthalamic
gamma-coherence showed the strongest dependence on the amount of objectified movement
presence. The coherences between the cortex and the ipsilateral STN were most clearly influenced
by both the severity of dyskinetic periods, as well as the amount of movement presence in the
gamma-range (Figure S3, right panels).

Figure 7: Cortical and subthalamic spectral powers and their linear dependencies on dyskinesia-severity and movement presence

The scatterplots demonstrate the modulation of theta (first row), low-beta (second row), and peak-gamma (third row) activity by dyskinesia severity (x-axis) and movement presence (y-axis). Subthalamic spectral activity is demonstrated in the left column, cortical spectral activity in the right column. Gamma activity seems to be more dependent on movement presence, whereas theta and beta modulations enhance with more severe dyskinesia. Local spectral power features were extracted from ten-second epochs of subthalamic data and averaged over bilateral STNs per epoch (left column) and cortical data (right column). Mean spectral powers are calculated per ten-second epoch (applying a 50%-overlap with consecutive ten-second epochs) and shown for the three main frequency bands of interest, i.e., theta band (4 – 8 Hz, first row), the low-beta range (12 – 20 Hz, second row), and the peak gamma range (5 Hz bin around an individual gamma-peak frequency, third row). Power values are individually z-scored and color-coded with yellow representing high values and blue representing low values. X-axes represent the absolute dyskinesia severity as total sum of the clinical dyskinesia rating scale (CDRS), including bilateral and axial dyskinetic manifestations. Y-axes represent the amount of movement present, as individually z-scored root mean squares from the bilateral accelerometer data.

Multivariate subthalamic features decode dyskinesia and movement influences predictions

Following the consistent modulation of subthalamic theta, beta, and gamma activity during dyskinetic periods (Figures 2, 4, and 7), we utilized each spectral band's direction of modulation to compose one single biomarker to decode dyskinesia periods:

Oscillatory Dynamics Ratio $= \dfrac{\theta_{STN} * \gamma_{STN}}{\beta_{STN}}$.

Formula 1: Composed multi-frequency biomarker to decode dyskinesia presence based on
subthalamic theta-, beta-, and gamma-power.
The product of subthalamic theta power (4 – 8 Hz) and subthalamic peak-gamma power (individual gamma
peak +/- 2.5 Hz) is divided by subthalamic low-beta power (12 – 20 Hz). All powers are calculated over ten-
second epochs using Welch's method and are individually standardized over all recorded data. Bilateral
subthalamic spectral power values were averaged per ten-second epoch and individually z-scored per
frequency band over the full recording.

The Oscillatory Dynamics Ratio (ODR) biomarker (see Formula 1) leverages LID-associated
subthalamic theta- and gamma-elevations and low-beta-attenuations and is expected to increase
gradually during the rise of dopaminergic levels. To test the predictive potential of the ODR, all
subthalamic recordings of subjects who experienced dyskinetic periods were included. All individual
timelines were aligned to individual LID onset moments (Figure 8A). ODR values increased gradually
after the intake of dopaminergic medication and showed a further increase after LID-onset compared
to before LID onset. Then, ODR were calculated for all ten-second windows of all 21 subjects and a
leave-one-subject-out cross validation was applied. For every iteration, an optimal linear ODR
threshold for LID detection was determined in each training set (n=20) and tested on the remaining
test subject. Using a linear regression, the ODR could predict LID presence with a mean balanced
accuracy of 0.61 (sd: 0.14), with statistically significant predictions in eight out of twenty-one
subjects.

The ODR (Formula 1 and Figure 8A) successfully differentiated between dyskinetic and non-dyskinetic
periods in a sub population without requiring complex computational methods, which is
advantageous for future clinical applications. However, it does not leverage the full spectrum of
spectral features associated with LID observed before (Table 1, Figures 4 and 5). Therefore, we tested
multivariate classifiers for LID detection with different combinations of spectral features. We
compared classifiers based on subthalamic features, cortical features, and cortico-subthalamic
features to identify the additional value of cortical data. For each feature set, both a linear logistic
regression model (LM) and a non-linear convolutional neural network using a novel latency
embedding method (CEBRA [48]) were applied (Fig S4). Since we recorded cortical data in only 13 out of
21 subjects, we first performed a head-to-head comparison of LID decoding based on only
subthalamic (STN), only cortical (ECoG) and cortico-subthalamic (STN-ECoG) features in the 13
subjects with cortical recordings (Figure S4). All features were individually z-scored using the average
and standard deviation from non-dyskinetic periods, in movement absence during the first 5 minutes
after levodopa-intake. Both the subthalamic and cortico-subthalamic models, for both LM and
CEBRA, outperformed the cortical models (Figures 8B and S4) (mean balanced accuracies: STN-LM:
0.76, sd: 0.12, STN-CEBRA: 0.77, sd: 0.14, STN-ECoG-LM: 0.75, sd: 0.12, and STN-ECoG-CEBRA: 0.72,
sd: 0.15, ECOG-LM: 0.56, sd: 0.10, and ECOG-CEBRA: 0.58, sd: 0.11).
In a comparison between the ODR and the multivariate LM and CEBRA models, while maximizing our
available data, the STN-CEBRA classifier detected LID presence with a significantly higher balanced
accuracy than the STN-ODR (paired t-test stat: -3.87, p = 0.001) and the STN-LM (paired t-test stat: -
4.18, p = 0.001) (Figures 7C and S4).

To better understand the predictive potential of the best performing classifier (STN-CEBRA), we
investigated the influence of LID severity and the presence of movement on the LID detection. We
categorized all subjects (n=21) based on their maximum LID severity observed, and compared the
balanced accuracy distributions for different LID severities (Figure 8D). Although the relatively small
number of included subjects per category limited statistical comparisons (two to six per category),

we did not see a difference in LID detection with respect to LID severity.
To assess the influence of movement presence on LID prediction, we sorted all ten-second windows
based on the amount of activity captured by the accelerometers (i.e., individual z-scored
accelerometer root mean squares, split in eight categories ranging from -2 to 4 sd). Within each
movement activity range, we calculated individual positive and negative predictive values
(respectively PPV and NPV). PPVs indicate what part of positive predictions (i.e. "LID is present") is
correct. NPVs indicate what part of negative predictions (i.e., "LID is not present") is correct.
Accordingly, the PPVs showed significantly more correct "LID present"-predictions with increasing
movement presence (Spearman R = 0.27, p < 0.001), and NPVs showed a decreasing trend of the
correctness with increasing movement presence (Spearman R = -0.07, p = 0.06) (Figure 8E).

Figure 8: Performances of dyskinesia detection using different recording sites and models

The presence of LID was predicted using different classifier models combined with different combinations of data. Panel A shows the temporal course of a single neural biomarker, composed of the product of subthalamic theta and gamma activity divided by subthalamic beta activity (Formula 1). Ratio values were calculated per second window. The time axis is individually corrected for LID onset. Subjects without LID are not included here. The shaded area represents the standard deviation.

Panel B shows the distributions of individual balanced accuracies for multivariate neural network-classifiers (CEBRA, see Methods) using only subthalamic data, only cortical data, or both. Only subjects with an ECoG electrode are included (n=13). Horizontal black lines between boxes indicate statistical difference in a paired t-test (alpha = 0.01, divided by 3 for multiple comparison correction.

Panel C shows the distributions of individual balanced accuracies for classifiers using the ratio biomarker (as shown in Panel A), a multivariate linear model (STN-LM), or a

multivariate neural network (STN-CEBRA). Based on superiority of the STN data, only STN data for all subjects
(n=21) are included.
Panel D shows the relation between the individual balanced accuracy of the LID prediction and the maximum
LID severity observed within each subject. Only results of the STN CEBRA model are shown due to its superior
performance. LID severity categories contain respectively 3, 3, 3, 6, 4, and 2 subjects. The horizontal black lines
in the boxes represent the median and the colored triangles represent the mean of each box. The position on
the x-axis categorizes the subjects on maximum LID severity as the maximum CDRS sum score assessed.
Panel E shows the distribution of positive and negative predictive values (respectively PPV and NPV) over
different amounts of movement objectified in the accelerometer data. Only the STN CEBRA model is included
based on its highest average performance. The x-axis is categorized based on z-scored individual accelerometer
root mean squares and ranges from "No movement" (z-score -2 and less) to "Many movements" (z-score +4
and higher). The lines represent the distribution of individual PPV (green) and NPV (purple) values within the
respective movement quantity. The NPV values show that the reliability, or the correctness, of "no LID present"
predictions, shows a decreasing trend with increasing movement presence (Spearman R = -0.07, p = 0.06).
Conversely, the PPV values show that the correctness of "LID present" predictions significantly increases with
increasing movement presence (Spearman R = 0.27, p < 0.001).
a.u.: arbitrary units, bal. accuracy: balanced accuracy, CDRS: Clinical Dyskinesia Rating Scale, CEBRA: neural
network using latent embedding (see Methods), ECoG: electrocorticography, LID: Levodopa-induced
dyskinesia, LM: linear model, PPV: positive predictive value, STN: subthalamic nucleus, NPV: negative predicted
value.

Discussion

We found a specific modulation of oscillatory theta, beta, and gamma band activity in STN and over motor cortical areas that was consistently associated with dyskinesia-provoking dopamine levels, and that showed specific modulation patterns in movement presence and absence during LID in PD patients. The respective oscillatory patterns could predict LID presence or absence, with STN activity being more accurate for LID prediction as compared to cortical neuronal patterns. Here, we propose an interpretation of our findings within a framework of systemic state changes [44]. A systemic state, of which there is only one at any given time for the entire system, integrates many other, more specific states, such as neural, behavioral, cardiac and others. These sub-states can vary in their spatial and temporal dynamics, which we can acknowledge by differentiating them into microstates (short-lasting) and macrostates (long-lasting). As such, all different sub-states form an integrated system responsible for specific behavior. Using neural deep brain recordings over an entire levodopa cycle that extends the dopaminergic dosage required for clinical benefit in our patients, we were able to measure neural and behavioral sub-states within periods of levodopa-induced dyskinesia (LID) for a defined time period by repeated measurements and as entire time-series. Our findings propose that dyskinetic periods can be interpreted as hyperdopaminergic macrostates that contain distinct microstates of movement presence and movement absence which can be objectified in cortico-subthalamic oscillatory activities. These microstates fit well to the clinically observed phenotype of LID.

Neural biomarkers have been identified in PD patients during dopaminergic modulation and LID [40,49–52], and are in line with the general pattern of oscillatory modulation of theta, beta and gamma band activity observed in our patients. Herein, increased theta and decreased beta band activity have been associated with periods of reduced motor inhibition, whereas gamma synchronization has been related to motor execution, in line with our concept of long-lasting macrostate changes and intermingled movement-specific microstate changes. Nevertheless, these temporal and behavioral aspects of neural biomarkers during LID with respect to involuntary movement periods have not been elucidated before. In order to interpret the specific behavioral relevance of the neuronal activity modulation observed during dyskinetic periods, we distinguished LID periods containing both movement presence and absence during the hyperdopaminergic state. We observed movement-independent bilateral theta increase and beta reduction of subthalamic neuronal activity that occur with the dopaminergic ON period but further increase during the dyskinetic period (Figure 4). Importantly, this slowly fluctuating macrostate change occurred bilaterally and was more prominent in periods without movements compared to movement presence suggesting that these changes are movement-independent and may rather balance motor inhibition and execution on a broader level.

Indeed, enhanced STN beta activity in PD has been characterized as a biomarker of bradykinesia and rigidity, i.e. serving as a 'anti-kinetic' rhythm within the cortical-STN network [53], that is reduced with levodopa treatment in parallel with motor improvement in PD [54,55]. During active behavioral motor inhibition [28], including reactive motor stopping in the non-dyskinetic state [43,56], beta activity has been described to increase [57–59], while beta desynchronization occurs during movement preparation, -execution and -imagery [28,55,60] suggesting that beta desynchronization is a prerequisite change in network activity for movements to occur [61]. In line with this concept, we observed a significant decrease in subthalamic beta activity during dyskinetic periods both during movement absence and movement presence (Figure 4), which fits previous reports of beta attenuation during LID in animals und patients [28,40,42,49–51]. Importantly, the beta activity attenuation in dyskinetic periods significantly extended the well-known beta attenuation during the medication ON state without LID (Figure 4) [31,55,62] and thus, is hypothesized to represent a hyperdopaminergic macrostate change of lowered motor inhibition that facilitates involuntary dyskinetic movements. This is further supported by the

significantly higher beta suppression with movement during LID as compared to movement during
non-dyskinetic periods and the significant decoupling of inter-STN beta activity. The robustness of
the LID-related beta attenuation regardless of movement presence, suggests that physiological and
dopamine-dependent movement-related low beta synchronization and desynchronization [43,63] may
adapt their magnitude during the hyperdopaminergic state. To further investigate the latter
however, the applied data-merging method (see Methods) limits the evaluation of single movement
effects. A described relation between LID severity and attenuations in cortical beta activity previously
observed in rodents [42], was not observed in our data. This may be explained by different recording
techniques, i.e. subdural ECoG recordings versus intracerebral recordings of cortical layers 5 and 6 in
the rodent study.
702

Theta activity has been found to increase with levodopa [64] and during dyskinesia in PD [37,40,50,51]. The
latter has been described as a constant activity change during biphasic as well as peak-dose
dyskinesia that was considered not movement-evoked [50,51]. In line with this, we found movement-
independent subthalamic theta activity increase during dyskinetic periods- underlining that the theta
increase is not associated with the abnormal movement itself. Strikingly, the subthalamic theta
increase in dyskinetic periods was more pronounced during movement absence. Additionally, our
data show that STN theta increase correlated with LID severity (Figure 4 and 7). This is similar to
observations in other hyperkinetic disorders such as dystonia, where theta activity is an established
biomarker [65] with pallidal activity driving EMG in dystonic muscles [66] and increased theta power and
coherence within the striato-pallidal loop correlates with severity of dystonic symptoms [67]. On the
cortical level, this relationship has so far only been described using EEG data in a few patients [50], but
a recent ECoG study did not show improved dyskinesia prediction incorporating cortical theta activity
in the model [40]. Similar to the latter, we did not find theta modulation with dyskinesia on the cortical
level, but cortico-STN coherence increased in the theta band during dyskinetic periods in our patients
supporting an involvement of the cortico-STN network in this frequency band. This would also be in
line with functional MRI data reporting reduction in dyskinesia during inhibitory pre-SMA repetitive
transcranial stimulation, an area that was hyperactive after levodopa intake in PD prior to occurrence
of dyskinesia [45,68].

Nevertheless, subthalamic theta elevation and increased cortico-STN theta coherence has also been
described during task-related successful active movement suppression in PD [43] and successful
suppression of involuntary tics were associated with increased cortical theta band coherence [69].
Furthermore, elevations of subthalamic theta and inferior frontal cortex EEG power are reported
during cognitive states of decision making during conflict [37,38,70], where STN activity has been
associated with active suppression of premature responses [71,72]. Accordingly, stimulation of the STN
can increase the number of incorrect responses under conflict [73] or premature responses in motor
inhibition tasks [74,75]. These findings support an association of subthalamic theta activity with motor
and non-motor inhibition including motor intention or involuntary movements. The functional role of
theta activity seems more complex and its topographical distribution in STN recordings is more
variable including non-motor ventral contacts during cognitive processing but also dyskinesia [37,50].
Whether task-related theta modulation more prominent in the motor-cognitive domain has a
different functional role as compared to theta activity recorded within the motor cortical-basal
ganglia loop during hyperkinesia cannot be answered with the current data available. Our analysis
suggests that an increase in theta oscillatory activity can serve as a biomarker for the slow
macrostate shift towards the dyskinetic state independent of movement presence per se.

Within this macrostate of lowered motor inhibition, single dyskinetic movements occur (Figure 1)
that showed an association with the increase in local subthalamic and cortical gamma activity and

cortico-subthalamic gamma coherence in our patients (Figure 4 and 5). Narrowband gamma
synchronization is the most established LID-related biomarker in both subthalamic [37,40,42,49,50] and
cortical [39–41,76] recordings and is considered a 'pro-kinetic' activity [53]. Importantly, we show for the
first time a distinct movement-dependence of gamma activity associated with LID as we
differentiated between movement presence and movement absence in dyskinetic periods. Previous
findings on gamma synchronization during LID lack this information as current dyskinetic animal
models typically lack movement absence recordings during dyskinetic periods [77], similar to recent
naturalistic patient recordings without continuous movement monitoring on a second-scale
resolution [40]. Within our framework of macro- and microstate integrated state dynamics, we
interpret these movements during dyskinetic periods as fast-changing microstates integrated in the
complex phenotype of LID, which are represented by gamma elevations in bilateral STNs and the
cortex contralateral to dyskinetic body sides. Movement execution has been closely associated with
gamma synchronization on basal ganglia and cortical level suggesting that gamma increase may scale
with the speed or vigor of voluntary movements [35,36,60,78–81]. During a hyperdopaminergic state of
reduced motor inhibition an imbalance of reduced beta activity and excessive prokinetic gamma
synchronization may lead to dyskinesia. Nevertheless, it is not yet clear if narrowband gamma
oscillations cause dyskinesia or are simply a physiological short-lived correlate of (excessive)
movements. In our patients, the strictly contralateral cortical gamma synchronization scaled with the
dyskinesia severity and further supports that it can be used as a fast-tracking biomarker for
movement activity [40]. However, its contralateral expression opposed to dyskinetic movement makes
it less robust as a biomarker in daily life and would suggest bilateral cortical recording hardware.

Subthalamic gamma activity has been argued to not merely effect movement execution, but also
motor stopping [82]. Since the clinical phenotype of LID includes active motor stopping as patients'
typically intent to terminate involuntary movements, the observed decrease in cortical gamma
activity during movement absence in the dyskinetic state may also be explained in parallel to
reported decreases in subthalamic and thalamic gamma activity during successful intentional motor
inhibition [43,83].

Together, fast changing movement-dependent gamma-magnitudes could be modulated by
underlying slower bilateral theta and beta rhythm fluctuations,[84] which would comply with the here
suggested framework of interpreting the complex motor network activity associated with LID into
macro- and microstate changes. The movement-independence of the bilateral subthalamic theta and
beta patterns supports our hypothesis that they represent a slow fluctuating macrostate of lowered
motor inhibition that leads to an increased likelihood of movement to occur. The movement
dependence of the STN and cortical gamma patterns suggest a more volatile and fast-changing
pathophysiological role that is more lateralized within the cortico-STN network and on the cortical
level. Thus, we suggest dyskinetic movements to occur as fast-changing microstates represented by
gamma activity changes, parallel to a slow-changing hyperdopaminergic macrostate of lowered
motor inhibition represented by a reduction in beta and increase in theta activity within the cortico-
STN network.

The hypothesized macro- and microstates fit into current fundamental principles and frameworks of
motor inhibition within the cortico-basal-ganglia motor network. The striatal motor network lowers
its motor inhibition under chronic dopaminergic depletion by adapting the SPN-architecture [17] and
the GABAergic [18,20] and glutamatergic [19,85] striatal projections. The hypothesized relevance of these
striatal changes for the pathophysiology of LID [86,87], was recently confirmed by studies showing that
imbalanced clusters of direct- and indirect-SPN and structural hypertrophy of SPNs allow the
occurrence of LID [22,25]. More specifically, parvalbumin-expressing (PV+) FSIs have been attributed
importance in motor control [88,89], striatal neuronal ensembles reconfiguration [90], and gamma

oscillation plasticity [91]. A recently discovered PV+ selective subthalamo-striatal projection [92] and the
presence of PV+ glutamatergic interneurons specifically in the dorsolateral STN [93,94], may underline
the importance of PV+ FSIs in STN DBS' sweet spot for pro-kinetic effects [9].
The abovementioned studies provide fundamental evidence on compensatory motor disinhibition in
PD and its association with LID occurrence. Moreover, the FSI-changes may explain the characteristic
gamma activity during dyskinetic movement.

Since LID is a common target for adaptive DBS [95], the demonstrated clinical relevance of
neurophysiological biomarkers can inform future adaptive DBS (aDBS) strategies, especially to
improve spatial and temporal characteristics of different closed-loop feedback mechanisms such as
fast versus slow aDBS [96]. The demonstrated differences between cortical and basal ganglia spectral
patterns during LID suggest that cortical biomarkers are sensible for fast aDBS algorithms that react
to microstate changes, but may lose the specificity and effect with slow adapting algorithms. For the
latter, subthalamic biomarkers may suit better using beta and theta band as biomarkers of
macrostate changes. For example, in case a fast aDBS algorithm is targeted at movement-dependent
cortical gamma activity, the volatile, non-continuous nature of movements has to be incorporated in
the aDBS decision making process. The superiority of our subthalamic multivariate prediction model
over the single ratio-biomarker and the cortical multivariate models (Figure 8BC), suggests that slow
aDBS models (i.e., ten-second windows) do not profit from additional cortical features and perform
best using computational heavier multivariate classifiers. The superiority of subthalamic features
over cortical features to detect LID presence we hypothesize to be a result of the ten-second window
length that favors slow network activity changes. We could therefore hypothesize that the
prerequisite for dyskinesia to occur depends on the slower neural activity change at the level of the
basal ganglia, which is in line with described discharge pattern changes within the striato-pallidal
loop [22]. Microstate changes will not be preferentially detected in our prediction model due to the
selected relatively long window length that inevitably leads to the merging of microstates of
movement presence with periods of movement absence within the same window for prediction.
Cortical features are therefore expected to have more predictive value in fast-aDBS strategies,
applying shorter window lengths, and thus isolating separate microstates of movement presence.
The latter is supported by earlier work showing that cortical features had more predictive value than
subthalamic features in movement prediction on a sub-second temporal scale [35,97].
Additionally we could demonstrate that the amount of movement presence within a ten-second
window significantly affected the predictive performance of LID detection, since positive LID-present
predictions were significantly more reliable in windows containing more movement (Figure 8E).
These findings underline the distinct character of the behavioral states of movement presence and
absence within a dyskinetic state and suggest that future daily life LID detection models will benefit
from a separate movement classifier.
The demonstrated level of neurophysiological and behavioral detail was the result of invasive cortico-
subthalamic recordings in PD patients in an in-clinic controlled environment. We addressed several
analytical challenges resulting from heterogeneous individual pharmacodynamics during a LID-
provoking protocol in PD patients. The before mentioned methodological challenges in rodent and
naturalistic, resp. a lack of movement absence during LID [21,77] or a lack of dyskinetic movement
monitoring [98], underlines the importance of this translational human research to bridge the gap to
valid real-world applications. Multimodal naturalistic monitoring methods such as wearables sensors
and ecological valid patient reported outcomes can be applied as naturalistic proxies for kinematic
and clinical observations [99–101].

Our findings propose a conceptual adaptation of the interpretation of neurophysiological biomarkers
for LID. Distinct spatial, temporal, and spectral neural patterns are associated with distinct clinically

relevant behavioral states. A slow movement-independent macrostate change is hypothesized to
occur with levodopa that is represented by bilateral modulation in theta and beta subthalamic
patterns, whereas a volatile, movement-dependent microstate is associated with fast-adapting
modulation in cortico-subthalamic gamma activity occurring with the dyskinetic movement itself. The
observed movement-independent spatial and spectral findings characterizing the dyskinetic state fit
well into current pathophysiological concepts of an overcompensated motor disinhibition in PD that
can allow for LID to occur. Finally, this improved behavioral understanding of these biomarkers aims
to tailor future aDBS to their specific clinical purposes to optimally serve PD patients during real-life.

Acknowledgements

We would like to thank the patients that participated in the study. We would like to thank Ulrike
Uhlig for her help with organizing and assisting during the recordings. A.A.K. received funding from
the Lundbeck Foundation as part of the collaborative project grant "Adaptive and precise targeting of
cortex-basal ganglia circuits in Parkinson´s Disease" (Grant Nr. R336-2020-1035), from the
Forschungsgemeinschaft (D.F.G., German Research Foundation) – Project-ID 424778381 – TRR 295,
and from the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under
Germany´s Excellence Strategy – EXC-2049 – 390688087. P.T. received funding from the D.F.G. in
Project-IDs: 433490190, 426503586, 424778381, and 426503586. WJN received funding from the
European Union (ERC, ReinforceBG, project 101077060), DFG – Project-ID 424778381 – TRR 295 and
the Bundesministerium für Bildung und Forschung (BMBF, project FKZ01GQ1802). J.H., R.L., and J.L.B.
are fellows of BIH Charité Clinician Scientist Programs. J.H. is a postdoctoral fellow of the Parkinson
Foundation.

Author contributions

J.H., R.L., and A.A.K. conceptualized and designed the study. J.H., R.L., V.M., R.K., J.B. performed the
electrophysiological recordings. K.F. and G.S. performed the neurosurgical procedures. P.K. assessed
all clinical dyskinesia scores. J.H. conducted the electrophysiological, movement, and statistical
analysis and drafted the study. T.B. assisted in the coherence analyses. J.H. and T.M. performed the
machine learning analyses, A.A.K. and J.N. supervised these analyses. J.H., P.T., and A.A.K.,
contributed to the interpretation of the data, revised, and edited the work critically. J.H., V.M., and
A.M. created the visualizations and performed clinical data acquisition. A.A.K. contributed critically to
the supervision, project organization and funding acquisition. All coauthors have reviewed and
approved the content of the manuscript.

References

1. Voon, V. *et al.* Impulse control disorders and levodopa-induced dyskinesias in Parkinson's disease: an update. *Lancet Neurol.* **16**, 238–250 (2017).

2. Jankovic, J. Motor fluctuations and dyskinesias in Parkinson's disease: clinical manifestations. *Mov Disord* **20 Suppl 11**, S11-6 (2005).

3. Hechtner, M. C. *et al.* Quality of life in Parkinson's disease patients with motor fluctuations and dyskinesias in five European countries. *Parkinsonism Relat. Disord.* **20**, 969–974 (2014).

4. de Bie, R. M. A., Clarke, C. E., Espay, A. J., Fox, S. H. & Lang, A. E. Initiation of pharmacological therapy in Parkinson's disease: when, why, and how. *Lancet Neurol.* **19**, 452–461 (2020).

5. di Biase, L., Pecoraro, P. M., Carbone, S. P., Caminiti, M. L. & Di Lazzaro, V. Levodopa-Induced Dyskinesias in Parkinson's Disease: An Overview on Pathophysiology, Clinical Manifestations, Therapy Management Strategies and Future Directions. *J. Clin. Med.* **12**, 4427 (2023).

6. Espa, E., Song, L., Skovgård, K., Fanni, S. & Cenci, M. A. Dopamine Agonist Cotreatment Alters Neuroplasticity and Pharmacology of Levodopa-Induced Dyskinesia. *Mov. Disord.* **38**, 410–422 (2023).

7. Lachenmayer, M. L. *et al.* Subthalamic and pallidal deep brain stimulation for Parkinson's disease-meta-analysis of outcomes. *NPJ Park. Dis.* **7**, 77 (2021).

8. Kübler, D. *et al.* Gender-specific outcomes of deep brain stimulation for Parkinson's disease — results from a single movement disorder center. *Neurol. Sci.* **44**, 1625–1631 (2023).

9. Tödt, I. *et al.* The Contribution of Subthalamic Nucleus Deep Brain Stimulation to the Improvement in Motor Functions and Quality of Life. *Mov. Disord. Off. J. Mov. Disord. Soc.* **37**, 291–301 (2022).

10. Pürner, D. *et al.* Nationwide Retrospective Analysis of Combinations of Advanced Therapies in Patients With Parkinson Disease. *Neurology* **101**, e2078–e2093 (2023).

11. Sriram, A. *et al.* Brittle Dyskinesia Following STN but not GPi Deep Brain Stimulation. **4**, 242 (2014).

12. Mainardi, M. *et al.* Deep brain stimulation of globus pallidus internus and subthalamic nucleus in Parkinson's disease: a multicenter, retrospective study of efficacy and safety. *Neurol. Sci. Off. J. Ital. Neurol. Soc. Ital. Soc. Clin. Neurophysiol.* **45**, 177–185 (2024).

13. Weiss, D. *et al.* Changing Gears - DBS For Dopaminergic Desensitization in Parkinson's Disease? *Ann. Neurol.* **90**, 699–710 (2021).

14. Picconi, B. *et al.* Loss of bidirectional striatal synaptic plasticity in L-DOPA–induced dyskinesia. *Nat. Neurosci.* **6**, 501–506 (2003).

15. Bove, F. *et al.* Neuroplasticity in levodopa-induced dyskinesias: An overview on pathophysiology and therapeutic targets. *Prog. Neurobiol.* **232**, 102548 (2024).

16. Cenci, M. A. & Kumar, A. Cells, pathways, and models in dyskinesia research. *Curr. Opin. Neurobiol.* **84**, 102833 (2024).

17. Gomez, G. *et al.* Changes in Dendritic Spine Density and Inhibitory Perisomatic Connectivity onto Medium Spiny Neurons in L-Dopa-Induced Dyskinesia. *Mol. Neurobiol.* **56**, 6261–6275 (2019).

18. Fino, E., Vandecasteele, M., Perez, S., Saudou, F. & Venance, L. Region-specific and state-dependent action of striatal GABAergic interneurons. *Nat. Commun.* **9**, 3339 (2018).

19. Campanelli, F., Natale, G., Marino, G., Ghiglieri, V. & Calabresi, P. Striatal glutamatergic hyperactivity in Parkinson's disease. *Neurobiol. Dis.* **168**, 105697 (2022).

20. Mallet, N., Ballion, B., Moine, C. L. & Gonon, F. Cortical Inputs and GABA Interneurons Imbalance Projection Neurons in the Striatum of Parkinsonian Rats. *J. Neurosci.* **26**, 3875–3884 (2006).

21. Alcacer, C. *et al.* Abnormal hyperactivity of specific striatal ensembles encodes distinct dyskinetic behaviors revealed by high-resolution clustering. 2024.09.06.611664 Preprint at https://doi.org/10.1101/2024.09.06.611664 (2024).

22. Parker, J. G. *et al.* Diametric neural ensemble dynamics in parkinsonian and dyskinetic states. *Nature* **557**, 177–182 (2018).

23. Thiele, S. L. *et al.* Selective loss of bi-directional synaptic plasticity in the direct and indirect striatal output pathways accompanies generation of parkinsonism and l-DOPA induced dyskinesia in mouse models. *Neurobiol. Dis.* **71**, 334–344 (2014).

24. Alcacer, C. *et al.* Chemogenetic stimulation of striatal projection neurons modulates responses to Parkinson's disease therapy. *J. Clin. Invest.* **127**, 720–734 (2017).

25. Abe, Y. *et al.* Shared GABA transmission pathology in dopamine agonist- and antagonist-induced dyskinesia. *Cell Rep. Med.* **4**, 101208 (2023).

26. Ryan, M. B. *et al.* Excessive firing of dyskinesia-associated striatal direct pathway neurons is gated by dopamine and excitatory synaptic input. *Cell Rep.* **43**, (2024).

27. Polyakova, Z., Hatanaka, N., Chiken, S. & Nambu, A. Subthalamic Activity for Motor Execution and
Cancelation in Monkeys. *J. Neurosci. Off. J. Soc. Neurosci.* **44**, e1911222024 (2024).
28. Kühn, A. A. *et al.* Event-related beta desynchronization in human subthalamic nucleus correlates with
motor performance. *Brain J. Neurol.* **127**, 735–746 (2004).
29. Callahan, J. W. *et al.* Movement-related increases in subthalamic activity optimize locomotion. *Cell Rep.*
43, 114495 (2024).
30. Lofredi, R. *et al.* Subthalamic stimulation impairs stopping of ongoing movements. *Brain J. Neurol.* **144**, 44–
52 (2021).
31. Neumann, W.-J., Gilron, R., Little, S. & Tinkhauser, G. Adaptive Deep Brain Stimulation: From Experimental
Evidence Toward Practical Implementation. *Mov. Disord.* **38**, 937–948 (2023).
32. Kühn, A. A., Kupsch, A., Schneider, G.-H. & Brown, P. Reduction in subthalamic 8-35 Hz oscillatory activity
correlates with clinical improvement in Parkinson's disease. *Eur. J. Neurosci.* **23**, 1956–1960 (2006).
33. Lofredi, R. *et al.* Pallidal Beta Activity Is Linked to Stimulation-Induced Slowness in Dystonia. *Mov. Disord.*
Off. J. Mov. Disord. Soc. **38**, 894–899 (2023).
34. Herz, D. M. *et al.* Dynamic control of decision and movement speed in the human basal ganglia. *Nat.*
Commun. **13**, 7530 (2022).
35. Merk, T. *et al.* Electrocorticography is superior to subthalamic local field potentials for movement
decoding in Parkinson's disease. *eLife* **11**, e75126 (2022).
36. Lofredi, R. *et al.* Dopamine-dependent scaling of subthalamic gamma bursts with movement velocity in
patients with Parkinson's disease. *eLife* **7**, e31895 (2018).
37. Rodriguez-Oroz, M. C. *et al.* Involvement of the subthalamic nucleus in impulse control disorders
associated with Parkinson's disease. *Brain* **134**, 36–49 (2011).
38. Zavala, B. A. *et al.* Midline frontal cortex low-frequency activity drives subthalamic nucleus oscillations
during conflict. *J. Neurosci. Off. J. Soc. Neurosci.* **34**, 7322–7333 (2014).
39. Halje, P. *et al.* Levodopa-Induced Dyskinesia Is Strongly Associated with Resonant Cortical Oscillations. *J.*
Neurosci. **32**, 16541–16551 (2012).
40. Olaru, M. *et al.* Motor network gamma oscillations in chronic home recordings predict dyskinesia in
Parkinson's disease. *Brain* awae004 (2024) doi:10.1093/brain/awae004.
41. Swann, N. C. *et al.* Gamma Oscillations in the Hyperkinetic State Detected with Chronic Human Brain
Recordings in Parkinson's Disease. *J Neurosci* **36**, 6445–58 (2016).
42. Dupre, K. B. *et al.* Effects of L-dopa priming on cortical high beta and high gamma oscillatory activity in a
rodent model of Parkinson's disease. *Neurobiol. Dis.* **86**, 1–15 (2016).
43. Alegre, M. *et al.* The subthalamic nucleus is involved in successful inhibition in the stop-signal task: a local
field potential study in Parkinson's disease. *Exp. Neurol.* **239**, 1–12 (2013).
44. Signoret-Genest, J. *et al.* Integrated cardio-behavioral responses to threat define defensive states. *Nat.*
Neurosci. **26**, 447–457 (2023).
45. Herz, D. M. *et al.* The acute brain response to levodopa heralds dyskinesias in Parkinson disease. *Ann.*
Neurol. **75**, 829–836 (2014).
46. Hagell, P. & Widner, H. Clinical rating of dyskinesias in Parkinson's disease: use and reliability of a new
rating scale. *Mov. Disord. Off. J. Mov. Disord. Soc.* **14**, 448–455 (1999).
47. Nikulin, V. V., Nolte, G. & Curio, G. A novel method for reliable and fast extraction of neuronal EEG/MEG
oscillations on the basis of spatio-spectral decomposition. *NeuroImage* **55**, 1528–1535 (2011).
48. Schneider, S., Lee, J. H. & Mathis, M. W. Learnable latent embeddings for joint behavioural and neural
analysis. *Nature* **617**, 360–368 (2023).
49. Fogelson, N. *et al.* Reciprocal interactions between oscillatory activities of different frequencies in the
subthalamic region of patients with Parkinson's disease. *Eur. J. Neurosci.* **22**, 257–266 (2005).
50. Alonso-Frech, F. *et al.* Slow oscillatory activity and levodopa-induced dyskinesias in Parkinson's disease.
Brain **129**, 1748–1757 (2006).
51. Alegre, M. *et al.* Subthalamic activity during diphasic dyskinesias in Parkinson's disease. *Mov. Disord.* **27**,
1178–1181 (2012).
52. Gilron, R. *et al.* Long-term wireless streaming of neural recordings for circuit discovery and adaptive
stimulation in individuals with Parkinson's disease. *Nat. Biotechnol.* (2021) doi:10.1038/s41587-021-
00897-5.
53. Brown, P. *et al.* Dopamine Dependency of Oscillations between Subthalamic Nucleus and Pallidum in
Parkinson's Disease. *J. Neurosci.* **21**, 1033–1038 (2001).
54. Lofredi, R. *et al.* Subthalamic beta bursts correlate with dopamine-dependent motor symptoms in 106
Parkinson's patients. *Npj Park. Dis.* **9**, 1–9 (2023).

55. Kuhn, A. A., Kupsch, A., Schneider, G. H. & Brown, P. Reduction in subthalamic 8-35 Hz oscillatory activity correlates with clinical improvement in Parkinson's disease. *Eur J Neurosci* **23**, 1956–60 (2006).

56. Horst, J. ter, Boillot, M., Cohen, M. X. & Englitz, B. Decreased Beta Power and OFC–STN Phase Synchronization during Reactive Stopping in Freely Behaving Rats. *J. Neurosci.* **44**, (2024).

57. Cao, C. *et al.* Low-beta versus high-beta band cortico-subcortical coherence in movement inhibition and expectation. *Neurobiol. Dis.* **201**, 106689 (2024).

58. Wessel, J. R. *et al.* Stop-related subthalamic beta activity indexes global motor suppression in Parkinson's disease. *Mov. Disord.* **31**, 1846–1853 (2016).

59. Aron, A. R., Herz, D. M., Brown, P., Forstmann, B. U. & Zaghloul, K. Frontosubthalamic Circuits for Control of Action and Cognition. *J. Neurosci.* **36**, 11489–11495 (2016).

60. Androulidakis, A. G. *et al.* Dopaminergic therapy promotes lateralized motor activity in the subthalamic area in Parkinson's disease. *Brain J. Neurol.* **130**, 457–468 (2007).

61. Williams, D. *et al.* Dopamine-dependent changes in the functional connectivity between basal ganglia and cerebral cortex in humans. *Brain* **125**, 1558–1569 (2002).

62. Kuhn, A. A. *et al.* High-frequency stimulation of the subthalamic nucleus suppresses oscillatory beta activity in patients with Parkinson's disease in parallel with improvement in motor performance. *J Neurosci* **28**, 6165–73 (2008).

63. Mathiopoulou, V. *et al.* Modulation of subthalamic beta oscillations by movement, dopamine, and deep brain stimulation in Parkinson's disease. *Npj Park. Dis.* **10**, 1–7 (2024).

64. Priori, A. *et al.* Rhythm-specific pharmacological modulation of subthalamic activity in Parkinson's disease. *Exp Neurol* **189**, 369–79 (2004).

65. Neumann, W.-J. *et al.* A localized pallidal physiomarker in cervical dystonia. *Ann. Neurol.* **82**, 912–924 (2017).

66. Sharott, A. *et al.* Is the synchronization between pallidal and muscle activity in primary dystonia due to peripheral afferance or a motor drive? *Brain J. Neurol.* **131**, 473–484 (2008).

67. Lofredi, R. *et al.* Striato-pallidal oscillatory connectivity correlates with symptom severity in dystonia patients. *Nat. Commun.* **15**, 8475 (2024).

68. Lohse, A. *et al.* Low-frequency transcranial stimulation of pre-supplementary motor area alleviates levodopa-induced dyskinesia in Parkinson's disease: a randomized cross-over trial. *Brain Commun.* **2**, fcaa147 (2020).

69. Serrien, D. J., Orth, M., Evans, A. H., Lees, A. J. & Brown, P. Motor inhibition in patients with Gilles de la Tourette syndrome: functional activation patterns as revealed by EEG coherence. *Brain* **128**, 116–125 (2005).

70. Cavanagh, J. F. *et al.* Subthalamic nucleus stimulation reverses mediofrontal influence over decision threshold. *Nat. Neurosci.* **14**, 1462–1467 (2011).

71. Frank, M. J., Samanta, J., Moustafa, A. A. & Sherman, S. J. Hold Your Horses: Impulsivity, Deep Brain Stimulation, and Medication in Parkinsonism. *Science* **318**, 1309–1312 (2007).

72. Herz, D. M., Frank, M. J., Tan, H. & Groppa, S. Subthalamic control of impulsive actions: insights from deep brain stimulation in Parkinson's disease. *Brain* **147**, 3651–3664 (2024).

73. Green, N. *et al.* Reduction of Influence of Task Difficulty on Perceptual Decision Making by STN Deep Brain Stimulation. *Curr. Biol.* **23**, 1681–1684 (2013).

74. Ballanger, B. *et al.* Release your horses: Deep Brain Stimulation of the subthalamic nucleus improves motor functions at the expense of response inhibition. A H215O PET study. *NeuroImage* **47**, S115 (2009).

75. Wylie, S. A. *et al.* Subthalamic nucleus stimulation influences expression and suppression of impulsive behaviour in Parkinson's disease. *Brain* **133**, 3611–3624 (2010).

76. Güttler, C. *et al.* Levodopa-Induced Dyskinesia Are Mediated by Cortical Gamma Oscillations in Experimental Parkinsonism. *Mov. Disord. Off. J. Mov. Disord. Soc.* **36**, 927–937 (2021).

77. Ye, T., Bartlett, M. J., Sherman, S. J., Falk, T. & Cowen, S. L. Spectral signatures of L-DOPA-induced dyskinesia depend on L-DOPA dose and are suppressed by ketamine. *Exp. Neurol.* **340**, 113670 (2021).

78. Alegre, M. *et al.* Movement-related changes in oscillatory activity in the human subthalamic nucleus: ipsilateral vs. contralateral movements. *Eur. J. Neurosci.* **22**, 2315–2324 (2005).

79. Brücke, C. *et al.* Scaling of movement is related to pallidal γ oscillations in patients with dystonia. *J. Neurosci. Off. J. Soc. Neurosci.* **32**, 1008–1019 (2012).

80. Crone, N. E. *et al.* Functional mapping of human sensorimotor cortex with electrocorticographic spectral analysis. I. Alpha and beta event-related desynchronization. *Brain J. Neurol.* **121 (Pt 12)**, 2271–2299 (1998).

81. Cassidy, M. *et al.* Movement-related changes in synchronization in the human basal ganglia. *Brain* **125**, 1235–1246 (2002).

82. Fischer, P. *et al.* Subthalamic nucleus gamma activity increases not only during movement but also during
movement inhibition. *eLife* **6**, e23947 (2017).
83. Brücke, C. *et al.* Thalamic gamma oscillations correlate with reaction time in a Go/noGo task in patients
with essential tremor. *NeuroImage* **75**, 36–45 (2013).
84. Buzsáki, G. & Wang, X.-J. Mechanisms of gamma oscillations. *Annu. Rev. Neurosci.* **35**, 203–225 (2012).
85. Hutchison, M. A. *et al.* Genetic inhibition of neurotransmission reveals role of glutamatergic input to
dopamine neurons in high-effort behavior. *Mol. Psychiatry* **23**, 1213–1225 (2018).
86. Gittis, A. H., Nelson, A. B., Thwin, M. T., Palop, J. J. & Kreitzer, A. C. Distinct Roles of GABAergic
Interneurons in the Regulation of Striatal Output Pathways. *J. Neurosci.* **30**, 2223–2234 (2010).
87. Gittis, A. H. *et al.* Selective Inhibition of Striatal Fast-Spiking Interneurons Causes Dyskinesias. *J. Neurosci.*
31, 15727–15731 (2011).
88. Koós, T. & Tepper, J. M. Inhibitory control of neostriatal projection neurons by GABAergic interneurons.
Nat. Neurosci. **2**, 467–472 (1999).
89. Jung, K. *et al.* An adaptive behavioral control motif mediated by cortical axo-axonic inhibition. *Nat.*
Neurosci. **26**, 1379–1393 (2023).
90. Duhne, M. *et al.* Activation of parvalbumin-expressing neurons reconfigures neuronal ensembles in murine
striatal microcircuits. *Eur. J. Neurosci.* **53**, 2149–2164 (2021).
91. Hadler, M. D., Tzilivaki, A., Schmitz, D., Alle, H. & Geiger, J. R. P. Gamma oscillation plasticity is mediated
via parvalbumin interneurons. *Sci. Adv.* **10**, eadj7427 (2024).
92. Kondabolu, K. *et al.* A Selective Projection from the Subthalamic Nucleus to Parvalbumin-Expressing
Interneurons of the Striatum. *eNeuro* **10**, ENEURO.0417-21.2023 (2023).
93. Jeon, H. *et al.* Topographic connectivity and cellular profiling reveal detailed input pathways and
functionally distinct cell types in the subthalamic nucleus. *Cell Rep.* **38**, 110439 (2022).
94. Lévesque, J.-C. & Parent, A. GABAergic interneurons in human subthalamic nucleus. *Mov. Disord. Off. J.*
Mov. Disord. Soc. **20**, 574–584 (2005).
95. Oehrn, C. R. *et al.* Chronic adaptive deep brain stimulation versus conventional stimulation in Parkinson's
disease: a blinded randomized feasibility trial. *Nat. Med.* 1–12 (2024) doi:10.1038/s41591-024-03196-z.
96. Little, S. & Brown, P. Debugging Adaptive Deep Brain Stimulation for Parkinson's Disease. *Mov. Disord. Off.*
J. Mov. Disord. Soc. **35**, 555–561 (2020).
97. Köhler, R. M. *et al.* Dopamine and deep brain stimulation accelerate the neural dynamics of volitional
action in Parkinson's disease. *Brain J. Neurol.* awae219 (2024) doi:10.1093/brain/awae219.
98. Ancona, S. *et al.* Wearables in the home-based assessment of abnormal movements in Parkinson's
disease: a systematic review of the literature. *J. Neurol.* **269**, 100–110 (2022).
99. Heijmans, M. *et al.* Monitoring Parkinson's disease symptoms during daily life: a feasibility study. *NPJ Park.*
Dis **5**, 21 (2019).
100. Feldmann, L. K., Roudini, J., Kühn, A. A. & Habets, J. G. V. Improving naturalistic neuroscience with patient
engagement strategies. *Front. Hum. Neurosci.* **17**, (2024).
101. Kamo, H. *et al.* A proof of concept: digital diary using 24-hour monitoring using wearable device for
patients with Parkinson's disease in nursing homes. *Front. Neurol.* **15**, 1356042 (2024).
1085
1086

Methods

Patient inclusions and characteristics

We recruited 21 patients that underwent implantation of bilateral STN DBS electrodes as a treatment for idiopathic Parkinson's disease at the Movement Disorders and Neuromodulation Unit of the Charité University Medicine Hospital. We excluded patients with increased perioperative bleeding risks, intracranial structural abnormalities, a poor understanding of German or English language, and patients suffering from postoperative complications. This study was in agreement with the Declaration of Helsinki and was approved by the Medical Ethical Committee of Charité University Medicine (record number EA2 127 19). We described patient characteristics of our included population in Supplementary Table 1.

Surgical protocol and invasive neural recordings

Standard clinical care subthalamic electrode implantations took place according to stereotactic gold standard procedures. All patients received Medtronic SenSight electrodes (Medtronic ©, Minneapolis, MI, US), except one patient who received Boston Scientific Cartesia X (Boston Scientific ©, Marlborough, MA, US) electrodes. 13 out of 21 patients agreed to be implanted with an additional unilateral electrocorticography (ECoG) electrode containing 6 linearly arranged contacts (Ad-tech Medical Instrument ©, Oak Creek, WI, US). To minimize additional perioperative risks, the ECoG-electrode implantation took place at the intraoperative second hemisphere. To minimize cardiac artefacts during future chronic sensing with implanted internal pulse generators [102], the majority of ECoG-electrodes was implanted over the right hemisphere. All recordings took place in between the first surgery for DBS electrodes implantation and the second surgery for internal pulse generator (IPG) implantation. Details on implanted neural recording hardware is summarized in Supplementary Table 2.

Cortical, subthalamic, and kinematic recordings

Neural and kinematic recordings were sampled via one amplifier (TMSi SAGA, TMSi International, Oldenzaal, The Netherlands) with a sampling frequency of 4000 Hz. All subthalamic and cortical local field potential (LFP) recordings had a bipolar configuration, using the lowest subthalamic ring-contact contralateral to the ECoG-electrode as a common reference. In case of no ECoG-electrode implanted, the lowest subthalamic ring-contact ipsilateral to the most affected body side was used as a common reference. A patch-electrode on the right shoulder was used as a ground-electrode. We recorded kinematic data using two tri-axial accelerometers, one attached to the distal end of each index finger. We videotaped all recordings continuously to for blinded post hoc clinical assessments.

Behavioral protocol and clinical assessments

Patients started the experimental protocol in the "OFF"-condition, deprived from dopamine for at least 12 hours. At virtual time point zero, they took 1,5 times their dopaminergic morning dosage (100 to 300 milligrams) as fast-acting L-Dopa (Madopar LT, Roche ©, Basel, CH) (Table S1). The participants showed the first clinical signs of LID after 21.0 minutes on average (sd: 15.5 minutes). Five out of eighteen subjects that developed LID showed some first, i.e., transient, dyskinetic signs within 5 minutes after levodopa-intake. The remaining thirteen subjects had an average LID onset of 29.0 minutes (sd: 10.0 minutes). The three participants that did not develop LID received higher preoperative levodopa equivalent daily dosages (LEDD) (1873 milligrams (mg), sd: 534 mg, versus 1084 mg, sd: 455 mg, p = 0.039, Mann-Whitney-U (mwu)), and received higher experimental dopaminergic doses (250 mg, sd: 40 mg, versus 179 mg, sd: 39 mg, p = 0.037). This comparison is limited by unbalanced group sizes. The eighteen patients that did develop LID had

mean peak dyskinesia severities of 5 CDRS points (sd: 3, range: 2 to 13 CDRS points), out of a
maximum of 28 points (the latter corresponds to severe dyskinetic movement in all seven rated body
parts).
The patients started with a 3 to 5 minutes rest recording at the beginning of the experiment around
levodopa intake, which was repeated 2 to 4 times throughout the protocol. In between the seated
rest recordings, the patients performed a self-paced hand tap task. Data collection continued for 60 –
75 minutes after levodopa intake and unscripted pauses were incorporated to avoid experiment
abortion due to exhaustion of the patient. Data collection was continued during the unscripted
pauses. During the rest recordings, we instructed the patients to sit still with opened-eyes and not
talk. During the self-paced hand-tap tasks, patients remained seated and got the instruction to raise
one hand and tap it one their upper leg. We instructed the patients to self-pace their taps roughly
every ten seconds, without internally counting seconds. When patients paused for shorter than 7
seconds, or longer than 12 seconds in-between taps, we corrected their tapping pace.

We performed a UPDRS III assessment before levodopa intake and at the end of the recording. We
collected preoperative UPDRS III scores in the dopaminergic OFF- and ON-conditions from
preoperative DBS evaluations. The presence and severity of dyskinesia was assessed using the Clinical
Dyskinesia Rating Scale (CDRS) throughout the whole experiment based on the video recordings [46].
CDRS scores were assessed every five minutes, to account for fluctuations in LID severity. We chose
the CDRS as clinical dyskinesia assessment based on its differentiation between LID in the upper and
low extremities, and between the left and right body side. An experienced movement disorders
specialist (PK) independently rated all CDRS scores, while being blinded for clinical information and
neural data. To categorize LID severities in mild, moderate, and severe, we applied thresholds at 4
and 8 CDRS points. These thresholds were decided upon based on the clinical expert's opinion after
being familiar with the CDRS distribution and its correlation with clinical severity of LID.

Kinematic signal processing and movement selection

Accelerometer time series were band passed filtered between 1 and 48 Hz (Finite Impulse Response
(FIR) filter with a hamming-window) and notch filtered (FIR, notch width 4 Hz and transition width 10
Hz) to remove line noise. We down sampled accelerometer time series to 512 Hz. The filtered
accelerometer signals were transformed into signal vectors magnitudes (svm) according to standard
accelerometer practices [103]. A custom movement detection was applied which combined the
presence of activity and identified acceleration-peaks to categorize the accelerometer data into
movement presence and movement absence. The incorporation of acceleration peaks within this
categorization, helped to identify ongoing movements. Based on the variation within the svms,
thresholds for movement detection were defined at 0.1 mG (milli-gravity, G: 9.8 m/s^2) and 0.5 mG for
respectively ongoing movement and acceleration peaks. The performance of this method was
controlled with individual visualizations of detected movements over the course of the experiments
(Figure S1 panel C and D). After confirming that the movement selection was successful for all
individual subjects, the resulting time stamps of single movement start- and end-points were
extracted. These time points were applied to 1) calculate the amount of movement presence within
each ten-second window of neural data (as applied in the analysis showed in Figure 7), and 2) to
isolate moments of movement presence and movement absence from the ten-second windows.
Neural data belonging to movement presence and movement absence moments were merged
separately, allowing for the neurophysiological comparison between movement presence and
movement absence, as used for the analyses showed in Figures 4 and 5.

1175

1176

Neural signal processing

We band pass filtered all raw multi-channel subthalamic and cortical neural time series, recorded respectively from the STN DBS electrodes and the ECoG electrodes, between 2 and 198 Hz using a FIR bandpass-filter with a hamming-window. A notch filter (FIR, notch width 4 Hz and transition width 10 Hz) removed line noise of 50 Hz including its harmonics. A threshold at eight standard deviations automatically detected segments as artefacts and visual inspections of all recordings confirmed the absence of artefacts after signal cleaning. Down sampling ensured a common sampling frequency of 2048 Hz in all neural time series.

All data were split in ten-second windows with a 50% overlap. We discarded windows with more than 33% of missing data. To reduce the data-dimensionality of the multi-channel neural recordings in an unbiased, data-driven manner, and to increase spectral signal to noise ratios, we applied the spatial-spectral-decomposition (SSD) method. One SSD transformation was calculated per frequency band of interest, on every ten-second window, including all neural channels from one neural recording source (either one STN-, or ECoG-electrode) using the Modular EEg Toolkit (https://github.com/neurophysics/meet) [47]. SSD transforms multi-channel time series into a number of independent, singular component time series optimized for variation within a specified frequency band and uses a singular value decomposition. We selected the first SSD-component per frequency band, which holds the maximum variance within the frequency band of interest. The defined frequency bands of interest are theta (4 – 8 Hz), alpha (8 – 12 Hz), low-beta (12 – 20 Hz), high-beta (20 – 35 Hz), and gamma (60 –90 Hz). We limited the gamma range to 60 and 90 Hz based on previous literature on dyskinesia-related neurophysiological findings [39,41,76]. We divided the selected gamma-band in three SSDs (resp. 60 – 70, 70 – 80, and 80 – 90 Hz), to prevent too large loss of broad-band activity based on the optimization based on an individual narrow gamma peak. The SSD-formula internally normalizes the resulting time series to the power in the neighboring frequencies (defined as 'flanks'). To generalize this normalization step and to improve comparability of SSD results over different frequency ranges of interest, we defined the flank-frequencies for every SSD formula up to 2 Hz and 198 Hz. The only exception are the three gamma-sub-bands that were calculated with identical flank ranges between 2 and 60 Hz, and 90 and 98 Hz.
The windowing and SSD transformations resulted in ten-second windows with one optimized neural time series per frequency range of interest (delta, alpha, low-beta, high-beta, and gamma-I, -II, and -III).

Descriptive spectral analysis

Power spectral densities (PSD) were calculated with a one-second window and a 1 Hz frequency resolution using a welch time-frequency decomposition (applying a Hahn-window and a 50% segment-overlap). Only the powers within the frequency range of interest were extracted for every time series resulting from the SSD transformations (e.g., low-beta (12 – 20 Hz) time series generated the spectral power per 1-Hz bin between 12 and 20 Hz). Finally, we combined the results of all SSD time series to create a full spectrum PSD from 4 to 35 Hz and from 60 to 90 Hz, as visualized in Figures 2 and 3. To allow for inter-individual comparison and group-level analysis, we baseline corrected all spectral powers as the percentage change with respect to the OFF-condition. Using a baseline correction compared to the OFF-condition, the spectral powers contained the same temporal relevance throughout the protocol, which would be lost when a relative spectral power for each time point was calculated. Baseline corrections were performed per 1-Hz bin.
To classify spectral power changes based on the presence and severity of clinically observed LID, we labeled every ten-second window to the nearest CDRS-assessment available. Since CDRS assessments were performed every 3 to 5 minutes, to maximal distance between a spectral power value and a

CDRS assessments was 1.5 to 2.5 minutes. Sum CDRS scores including all bilateral and axial items
were included, except for the lateralization analysis. The latter made use of unilateral sums of both
body side, excluding the axial CDRS parts in the lateralized LID score.
Squared coherence was applied as a functional connectivity metric [52]. Similar to the spectral power
calculation, we applied Welch's methods for PSD and cross-spectral density (CSD) calculation in the
formula 2. Squared coherence values were calculated between the two bilateral STNs, and the
ipsilateral STN and cortex, composed from the SSD-time series and baseline corrected similarly to the
spectral power values.

1231

1232
$$\text{COHsquared} = \left| \text{csd}_{source1 \, x \, source2} \right|^2 / (\text{psd}_{source-1} * \text{psd}_{source-2})$$

1233   **Formula 2: Squared coherence calculation.**

1234

Feature extraction and predictive modeling

To extract neural features for the predictive analyses, we used the SSD'd ten-second windows and
calculated mean spectral power and spectral variation per frequency range of interest. Spectral
powers were calculated using a Welch's decomposition with a Hanning window and one second
segments with 50% overlap. Spectral variation was calculated as the coefficient of variation of the
envelope of the filtered time series (Hilbert transformation). Since SSD calculations were performed
for three sub-bands of broadband gamma within 60 to 90 Hz, we included the maximum value of the
three ranges as gamma value. In feature sets including subthalamic data, we included mean values of
coherences between the bilateral STNs. In the cortico-subthalamic feature sets we included
coherences between the ipsilateral STN and ECoG-recordings. Mean coherences values were
calculated for each band width of interest using the respectively SSD'd time series, per ten-second
windows.
This feature extraction resulted in individual data sets containing one value per feature for every ten-
second window. Every ten-second window was labeled with the nearest dyskinesia assessment.

For the predictive analyses, we applied two different classification models, i.e., a linear model and a
neural network leveraging latency embedding strategies [48]. Both models decoded binary LID
presence (0: non-dyskinetic states, 1: dyskinetic states) based on the above described features. STN
models included local subthalamic features and subthalamic-subthalamic coherences. ECoG models
included cortical features. STN-ECoG model included both local subthalamic and cortical features, as
well as subthalamic-subthalamic and cortico-subthalamic coherences. A leave one subject out cross-
validation was applied for all available subjects. For head-to-head model comparisons, only subjects
with an ECoG-electrode (n=13) were included (Figure 8B). For the remaining classifiers (Panel C-E) all
available subjects were included (n=21). First, we applied a linear model that decoded LID presence.
For this, a ridge-regularized class-weight balanced logistic regression model, implemented in scikit-
learn [104] (L-BFGS solver, 100 iterations) was applied. In addition to linear decoding, we utilized non-
linear deep learning for classifying binary dyskinesia presence. We used the recently presented
CEBRA package (Consistent Embeddings of high-dimensional recordings using auxiliary variables) [48].
The CEBRA contrastive learning method was implemented using the "offset10-model", a five-layer
convolutional neural network implemented in PyTorch [105]. The first layer consists of 32 hidden units,
followed by three convolutional skip layers, each with 32 hidden units, and a final four-dimensional
convolutional output layer. The skip layers created a bottleneck in the temporal filter dimension,
reducing it from 10 to 3 samples. A Gaussian Error Linear Unit (GELU) activation function was applied
to each layer, with normalization on the output layer. The model used the "constant" temperature

mode, a learning rate of 0.001, and was trained for 1000 iterations with a batch size of 300.
Contrastive learning sampling was performed with the InfoNCE loss function [106], using a cosine
distance between samples, by selecting time samples within a "time_offset" of 10, along with
samples corresponding to the auxiliary dyskinesia binary training target variable. The four-
dimensional embedding was following by training a ridge-regularized class-weight balanced logistic
regression model, implemented in scikit-learn [104] (L-BFGS solver, 100 iterations), which was also
applied for the non-cebra linear classifiers.

Analysis software

We performed all analysis with custom-written code in Python in which we made use of standard
libraries as numpy [107], pandas [108], scipy [109], sklearn. The custom libraries tmsi-python-interface [110],
meet [111] were used for specific parts of signal processing as referred to in the methods.

Code availability

The full code used for the processing of neural and kinematic data, performing descriptive and
predictive analyses, and plotting is available on online:
https://github.com/jgvhabets/dyskinesia_neurophys.

Data availability

Data can be made available in anonymized form on reasonable request.

1285

References

102. Neumann, W.-J. *et al.* The sensitivity of ECG contamination to surgical implantation site in brain computer
interfaces. *Brain Stimulat.* **14**, 1301–1306 (2021).
103. Habets, J. G. V. *et al.* A First Methodological Development and Validation of ReTap: An Open-Source
UPDRS Finger Tapping Assessment Tool Based on Accelerometer-Data. *Sensors* **23**, 5238 (2023).
104. Pedregosa, F. *et al.* Scikit-learn: Machine learning in Python. *J. Mach. Learn. Res.* **12**, 2825–2830 (2011).
105. Paszke, A. *et al.* PyTorch: An Imperative Style, High-Performance Deep Learning Library. Preprint at
https://doi.org/10.48550/arXiv.1912.01703 (2019).
106. Oord, A. van den, Li, Y. & Vinyals, O. Representation Learning with Contrastive Predictive Coding. Preprint
at https://doi.org/10.48550/arXiv.1807.03748 (2019).
107. Walt, S. van der, Colbert, S. C. & Varoquaux, G. The NumPy Array: A Structure for Efficient Numerical
Computation. *Comput. Sci. Eng.* **13**, 22–30 (2011).
108. Reback, W. M., jbrockmendel, Joris Van den Bossche, Tom Augspurger, Phillip Cloud, …. Martin Winkel Jeff.
Pandas 1.1.1. https://doi.org/10.5281/zenodo.3993412 (2020).
109. The SciPy Project. SciPy.
110. TMSi International. tmsi-python-interface.
111. Waterstraat, G. Modular EEg Toolkit.

1303