Context-dependent reshaping of defensive responses to predators in head-fixed and freely moving mice

Marti Ritter[1], Lara Mariel Chirich Barreira[1], Lara Sach[1], Aileen Hakus[2,3,4], Sanra Kumsal Öktem[1], Ronny Bergmann[5], Anne Voigt[5], Dietmar Schmitz[5,6,7,8], Panayiota Poirazi[9], Matthew E. Larkum[1], Robert N. S. Sachdev[1]

[1]Inst. of Biol., Humboldt University of Berlin, Berlin, Germany;
[2]Grad. Ctr. of Life Sci., Humboldt University of Berlin, Berlin, Germany;
[3]Dept. of Psychiatry and Psychotherapy, Charité Universitätsmedizin Berlin, Germany;
[4]Intl. Grad. Program Med. Neurosciences, Charité Universitätsmedizin Berlin, Germany;
[5]Neurosci. Res. Ctr., Charité Universitätsmedizin Berlin, Germany;
[6]Bernstein Center for Computational Neuroscience Berlin;
[7]Einstein Center for Neurosciences Berlin, Charité-Universitätsmedizin Berlin, Germany;
[8]Max-Delbrück Center for Molecular Medicine in the Helmholtz Association Berlin, Germany;
[9]Institute of Molec., Biol., and Biotech., Foundation for Research and Technology -- Hellas, Heraklion, Greece.

Correspondence to: bs387ster@gmail.com; Matthew.Larkum@gmail.com; Poirazi@imbb.forth.gr

Running head/Short title: Innate defensive responses in mice

Keywords: Innate Fear, Predator, Defensive Behaviors, Rats

Number of figures: 8

Multimedia videos: 6

Total number of words: 8506 (not including figure legends or references)

Additional Information: The following funding sources have supported this project: (1) Deutsche Forschungsgemeinschaft (DFG), Grant Nos. 246731133, 250048060 and 267823436 to ML; (2) DFG Project number 327654276 – SFB 1315 to ML; (3) European Commission Horizon 2020 Research And Innovation Program and Euratom Research and Training Program 2014–2018 (under grant agreement No. 670118 to ML); (4) Human Brain Project, EU Commission Grant 720270 (SGA1), 785907 (SGA2) and 945539 (SGA3) to ML; (5) Einstein Foundation Berlin EVF-2017-363 to ML; Einstein Foundation Visiting Fellowship EVF-2019-508 to PP; Humboldt Foundation Friedrich Wilhelm Bessel Research Award to PP.

Acknowledgements: We thank the Charité Workshop for technical assistance, especially Alexander Schill, Jan-Erik Ode and Daniel Deblitz. We also want to thank Melissa Long and the team of the ABPF for their help with animal handling and room planning. Finally, we thank Laura Schwarz of the Sainsbury Wellcome Centre and members of the Larkum lab for useful discussions about earlier versions of this manuscript.

Abstract

Behavioral responses to threat — such as fleeing, freezing, or fighting—can be innate, learned, and strongly shaped by context or competing goals. Here, we asked whether exposure to an ecologically relevant predator obligatorily elicits canonical defensive behaviors across behavioral contexts. We examined predator responses in mice across four experimental conditions: one novel head-fixed reward-driven foraging task and three established paradigms in freely moving animals. In the head-fixed condition, water-deprived mice were trained to walk on a treadmill controlling a virtual environment and water reward delivery and were subsequently exposed to a live rat positioned above the lick spout. Despite the presence of the predator, most mice (5 of 7) maintained foraging performance at baseline levels. However, individual mice exhibited significant, coordinated changes in running speed, pupil diameter, eye movements, and posture, indicating engagement with the threat. To assess how context influences predator responses, we exposed 36 naïve, freely moving mice to fear-inducing stimuli, including looming visual cues, rat odor, and a live rat. Even under these conditions, defensive behaviors were variable: only a subset of mice displayed avoidance or escape, and when presented with a freely moving rat, approximately half of the mice avoided the predator. Together, these findings show that predator threat does not elicit a uniform or obligatory defensive repertoire in mice. Instead, defensive responses are expressed flexibly, and are shaped by environmental constraints, task demands, and individual variability. These results challenge the assumption that innate fear behaviors are automatically triggered by predator encounters.

61   **Graphical Abstract**

62   Using a head-fixed, reward-based foraging task and complementary freely moving

63   paradigms, we show that mouse responses to predator-related stimuli are variable and context

64   dependent, with limited expression of canonical defensive behaviors such as freezing or flight.

Introduction

Threat perception and appropriate reaction to a threat are necessary for survival. Overestimating dangers, for example misinterpreting a benign sound as a predator, or underestimating a risk, for example misjudging the strength of a barrier separating prey from a predator, can lead to unwanted outcomes ranging from unnecessary anxiety to injury or death. Accordingly, threat processing has been studied extensively across species, with work traditionally focusing on three interrelated components: the detection of a threat, the behavioral and physiological responses it evokes, and the subjective experience of fear (Beckers et al. 2013; Fanselow and Pennington 2017; LeDoux and Brown 2017; Mobbs et al. 2019; Silva et al. 2016; Tovote et al. 2015). Much of what is known about threat processing derives from Pavlovian fear conditioning paradigms (McDannald 2023), in which a neutral stimulus—often an auditory cue— is paired with an aversive outcome such as a mild foot shock (Mowrer and Lamoreaux 1946). Through repeated pairing, the previously neutral cue comes to elicit defensive behaviors, including freezing, fleeing, postural changes, and accompanying physiological responses such as pupil dilation and increased heart rate (Blanchard and Blanchard 1989). These classes of paradigms offer precise experimental control and have been instrumental in delineating the neural circuits underlying threat detection and defensive behavior (Phillips and LeDoux 1992; reviewed in Tovote et al. 2015). In parallel, a range of ethologically inspired paradigms have been developed to probe so-called "innate" fear responses using evolutionarily relevant stimuli. Looming visual stimuli that mimic an approaching aerial predator can evoke rapid escape or freezing responses in rodents (Schiff 1965; Yilmaz and Meister 2013). Similarly, exposure to predator-associated odors (e.g., fox or cat urine), heights, loud sounds, or larger conspecifics has been used to elicit defensive behaviors by introducing naturalistic threats (Gibson and Walk 1960; Mongeau et al. 2003; Farmer-Dougan et al. 2005; Silva et al. 2016). These approaches are often assumed to engage hardwired defensive circuits that produce stereotyped behavioral outputs.

However, accumulating evidence suggests that defensive behaviors are not purely reflexive but are modulated by context, internal state, and available action (Maren et al. 2013). When threats occur in environments that impose physical constraints or introduce competing motivations, such as the need to obtain food or water, defensive responses may be reshaped rather than being simply expressed or suppressed (Burnett et al. 2016). Under such conditions, animals may engage in graded threat assessment or alternative coping strategies instead of canonical freezing or flight. In

this study, we sought to test whether exposure to an ecologically relevant predator obligatorily

elicits stereotyped defensive behaviors, or whether such responses are flexibly expressed

depending on behavioral context and physical constraints. To address this, we developed a head-

fixed, reward-based foraging paradigm in which mice ran on a treadmill to obtain water while

exposed to a live rat positioned directly above the lick spout. The head fixed preparation is the

state of the art and standard approach for circuit mapping, calcium imaging, and high-density

neural recordings to understand the neural basis of sensory motor behaviors and is often used for

studying fear conditioning (Lovett-Barron et al. 2014; Abs et al. 2018; Ahmad et al. 2020; Dalmay

et al. 2019). Here our goal was to develop a novel paradigm that could be easily combined with

state-of-the-art imaging and recording methods. To achieve this goal, we examined predator

responses under conditions of constrained movement and competing motivational demands. We

compared behavior in this novel context to responses observed in freely moving mice exposed to

established fear-inducing stimuli, including looming visual cues, predator odor, and a live rat.

Previous work has suggested that rats can act as natural predators of mice, eliciting innate

defensive responses (Karli 1956; Panksepp 1971; Blanchard et al. 1998). Here, we asked whether

mice would alter their behavior in the presence of a rat when foraging for reward was required, or

whether they would continue goal-directed behavior while expressing alternative, non-canonical

indicators of threat engagement. Surprisingly, although head-fixed mice showed clear changes in

physiological and postural measures during predator exposure, most continued to forage.

Moreover, even in freely moving paradigms designed to elicit robust fear responses, mice did not

consistently display avoidance, freezing, or escape. Together, these observations suggest a

reassessment of how "innate" defensive behaviors are expressed across different contexts.

Methods and Materials

All experiments were conducted in accordance with the guidelines of animal welfare of the Charité Universitätsmedizin Berlin and the local authorities, the 'Landesamt für Gesundheit und Soziales'. Adult mice (n=43) on a C57BL / J6 background and 5 rats were used in this study. For experiments involving head fixation, seven mice (4 male, 3 female, 8-12 months old) were sourced from the internal breeding facilities. We selected naive, untreated mice available in our facility. This resulted in the given mix of ages and sexes. For experiments with freely moving mice (n=36), male mice (~8 weeks old at start of experiments) were sourced from our in-house breeding colony, Janvier, and Charles River (n=12 each). Their origin was blinded prior to experiments, by having a separate researcher uninvolved with the experiments and analysis anonymize the mouse IDs and groups during the separation of the mice. Mice were housed socially with at least one other littermate. Five female Wistar rats served as the predator stimulus and were kept in the same facility as the mice, but in another room. For experiments with head-fixed mice, we used a different rat each day. In the freely-moving paradigms, a random rat was selected by the experimenter for each session. Both the mice and rats were kept in a 12h reversed light cycle (mice 6pm to 6am, rats 5pm to 5am). Except for mice that were actively used in the head-fixed simulated foraging paradigms and were on water restriction, all animals had ad lib access to food and water.

Surgical procedures for head-fixation

On the day of the surgery, adult C57Bl6 mice (n=7) weighing 20-40 grams were injected with Carprofen (5 mg/kg) intraperitoneally pre-surgery, then deeply anesthetized with a mixture of Ketamine and Xylazine (Ketamine 12.5mg/ml, Xylazine 1 mg/ml, 10 µl/g dose) and placed on a heating pad maintained at 37°C. A local analgesic, Lidocaine (100 µl), was injected under the skin at the site of the incision. Lightweight aluminum headposts (3 cm long, 2mm thick, weighing 0.4 gm) were affixed to the skull using Rely X and Jet Acrylic (Ortho-Jet) black cement (Dominiak et al., 2019; Ebner et al., 2019). Post-surgery analgesia was provided over 3 days, with Carprofen injections (5 mg/kg) intraperitoneally and, if there were signs of post-operative pain, Buprenorphine (0.05-0.1 mg/kg) subcutaneously in addition.

Simulated foraging experiment

The simulated foraging environment consisted of a rigid-foam-based circular treadmill and a 30 cm long plexiglass tube with a 7 cm diameter (**Figure 1, Video 1**). The treadmill was

lightweight enough that mice could move it effortlessly. It had a rubberized surface to increase the
grip of the mouse on the wheel. The movement of the treadmill was read by an encoder. The output
of the encoder was used to control the position of the plexiglass tube, which was mounted on a rail
and controlled by a small stepper motor (**Figure 1**). To ensure safe and stable movement, and to
minimize the stress on a rat held in the tube, the top speed of the tube was limited to 4 cm per
second. The output of the encoder was also linked to the PC that controlled two large monitors
positioned on each side of the treadmill that were used to stream visual stimuli tethered to the
movement of the treadmill.

The tube holding the rat was coated with a red film limiting the ability of mice to see the
position of the rat, or to see whether there was a rat in the tube (**Figure 1A**). Additionally, the front
of the tube was covered with a disk which could be rotated by an electronic input (**Figure 1B**).
When a rat was in the tube, the disk could be used to selectively hide or show the rat during the
simulated predator encounter. The disk could be rotated to allow odors through, or to block all
sensory stimuli from inside the tube. When the disk was in the open position, the rat could stick
its nose out of the tube and almost touch the mouse when it was at the lick spout.

Electronics and control of behavior

For an overview of the connections in the setup, see **Supplementary Figure 1**.

Trial control

A finite state machine (Bpod r2, Sanworks LLC) monitored trial states (see
Supplementary Figure 2), and the sequence of reward delivery and data acquisition. This state
machine also controlled the high-resolution acquisition of movement traces from the encoder in
the treadmill and triggered the cameras (recording at 100Hz).

Motor control

In addition to the state machine, a single-board PC (Raspberry Pi 4 by Raspberry Pi
Foundation) was used for real-time control. It received a 1kHz positional signal from the treadmill
(also received by the Bpod), downsampling it to 25 Hz and then transferring it to a microcontroller
(Arduino Nano by Arduino.cc) which controlled stepper motors to translate this into the
synchronized motion of the plexiglass tube. If the mouse moved the treadmill faster than 4 cm / s
then the downsampled signal was truncated to ensure that the tube followed smoothly. Note that
the treadmill moved as fast as the mice needed; but if the treadmill moved at a high speed, the

output was translated into a manageable speed for moving the lick spout, and the associated large
tube that could contain a living rat.

Rat stimulus & Visual stimuli

The position of the disk at the front of the tube was decided by the Bpod, and then updated
by the single-board PC and the microcontroller (Arduino Nano by Arduino.cc), which controlled
the stepper motors that rotated the disk. Finally, the single-board PC was used to generate the
virtual visual environment tethered to the movement signal of the treadmill. The monitors used for
displaying the streaming stimuli were also used as a go-cue at the beginning of the trial and an
end-cue at the end. The streaming stimuli consisted of symbols that moved in synchrony with the
treadmill. The screen flashed green at the start of a new trial and red at the end of a failed trial.

Data acquisition

The treadmill sent out a 1 kHz signal encoding the current position. This signal was
recorded on the state machine and stored to disk on a central workstation, along with the specific
settings used in each trial. Two cameras were located below one of the screens used for the visual
stimulus and set up such that they recorded a side view of the entire body of the mouse and its
face, respectively, at 100Hz and sent this data to the central workstation. The workstation also
provided centralized control over the single-board PC and state machine.

Habituation and training

After surgical operation to implant a head post, mice were habituated to being handled and
head-fixed (**Figure 1C, D**). Training on the treadmill began, once mice tolerated head fixation for
~ 20 minutes. At this point, water intake for mice was monitored and restricted, but ensured to be
at a level resulting in no more than 20% of weight loss per mouse compared to the weight before
onset of water control. To ensure that all mice were available on the same experimental day when
a rat was introduced, training continued for 3-4 weeks.

The initial training consisted of rewarding mice manually with Saccharose-sweetened
water when they were head-fixed and they moved the treadmill in the correct / forward direction.
Once mice moved the treadmill in the correct direction, training was automated. In the first days,
there were no time constraints. Mice simply had to move at least 28.5 cm forward on the treadmill
to receive the sweetened water reward, with no other limitation to speed or movement patterns.
The total distance to the treadmill was 30 cm, but for safety reasons we automated the closing in
of the spout on the last 1.5 cm. Five seconds after reward delivery, the lickspout / tube contraption

automatically moved to its starting position, away from the mouse and the next trial began. The

beginning of a new trial was indicated by a sound cue (played by a buzzer at 50Hz) and flashing

of the virtual reality screens.

In the next phase of training, a time limit was introduced. Mice had 30 seconds to move

the treadmill at least 28.5 cm and then another 2 seconds to lick the spout in front of them. Once

mice learned to keep moving the treadmill to obtain a reward, the duration of the trial was

shortened to 12 seconds. When the cohort of mice achieved >65% successful trials (with at least

30 trials per day) under these conditions for 3 days, the control and experimental data were

collected.

Live predator encounter

Once baseline control data had been collected on the next day (**Figure 1C-F**), a rat that had

been habituated to handling and to the apparatus, was placed in the tube. The mouse was then

head-fixed to the treadmill and experimental data were collected. The day after the encounter with

the rat, a second day of baseline data was collected without the rat.

Experimental paradigm with freely moving mice

One cohort of twelve male C57Bl6/J mice and a second cohort of twenty-four male

C57Bl6/J mice were used in these experiments. Mice were ordered in equal numbers from the in-

house breeding colony, Janvier, and Charles River. For the first cohort the 3 female Wistar-rats as

in the simulated foraging task were used, and for the second cohort 2 additional female Wistar-

rats were used. Female rats were used to minimize the possibility of a rat biting through the holder

or injuring a mouse (Karli 1956). We chose to use animals from multiple sources to verify whether

mice obtained from our in-house colony showed unusually low threat responses. For each session

involving rats with freely-moving mice the experimenter randomly selected a rat to be used. To

ensure that mice were naive to the stimulus in each of the paradigms, we ordered them by

ascending complexity. We applied first the looming stimulus (artificial visual), then presented rat

odor (natural olfactory), and finally a live rat (natural olfactory/visual/potentially tactile).

Looming stimuli

To evaluate innate threat responses independent of the presence of a live predator, we

applied the well-established looming stimulus paradigm. This paradigm is a standard for inducing

fear in mice. The mice were solitary housed and habituated to changes in day-night-cycle for 5

days, then over the next four days, one mouse per source and day (resulting in 3 mice per day) was

placed in a darkened room for an hour before experiments began. The purpose of the solitary
housing of these mice was to ensure a minimum of unintended interaction between mice before
each recording session. On the experimental day, mice were moved into a plexiglass arena (0.5 m
x 0.29 m) and left to explore it for 10 minutes. The arena was cleaned with 15% Ethanol before
each mouse was introduced. It contained a shelter, made of red plexiglass. When mice entered a
zone encompassing the quarter of the arena floor most distant from the shelter (~25 cm from the
shelter edge), a looming stimulus (a shadow that expanded above the mouse) was presented. The
looming stimulus consisted of 5 presentations of a small black dot (3°visual angle) expanding
rapidly over 200ms to its full size (50°) and then remaining at this size for 250 ms. The stimulus
was repeated at least 90 s after the last presentation. The sequence of looming stimuli was repeated
10-14 times for each mouse over 45 minutes. Video data was collected at 60 Hz. Trials ceased
when a mouse stopped leaving the shelter for at least 15 minutes. Each stimulus presentation was
triggered manually by a researcher supervising the experiment through a monitor located out of
sight of the mouse, but within the same room as the setup. Consistency was ensured by defining a
common criterion for activation of the stimulus. The trigger moment was defined as the moment
when a mouse crossed a visible demarcation line (marking the distant quarter of the arena relative
to the shelter) with all 4 paws.

Test with Rat odor

The mice used in the looming experiments were then used to examine place preference in
the presence or absence of rat odors. Over two days two mice from each source, i.e. in-house,
Charles River, and Janvier, per day (6 mice per day) were moved into the same plexiglass arena
as used in the looming stimulus test one at a time. The arena was split in half with a small connector
between the two sides. Mice were left to explore both sides of the arena for 10 minutes. Then the
mouse was removed, and a rat was brought into the arena, but could only explore one side of the
arena. During this time the rat could move freely on that side of the arena. The rat was removed
after 5 minutes, and the mouse was returned to the arena. If the rat urinated in the arena, the floor
was wiped dry with a paper towel. Movement of the mouse in the two conditions was tracked
offline with video acquired at 60Hz for the first cohort and 25Hz for the second cohort. The change
in sampling rate was decided based on the data collected with the first cohort, as higher frame rates
than 25Hz were not necessary for tracking the location of the mice. The arena was cleaned before
each mouse was introduced into the arena.

Assaying response to live rat

Finally, the same mice were placed in a modified version of the plexiglass arena (again 2 mice per group per day). The arena was divided in half by a 1mm thick metal mesh, with gaps that were just large enough for mice or rats to stick the tips of their noses through the mesh. Control data was acquired for five minutes, then mice were removed, and a rat was brought into the arena. Two minutes after introducing the rat, mice were introduced into the other side of the arena. Movement of the mouse was tracked offline with video acquired at 25-60Hz. The arena was cleaned before each mouse was introduced into the arena.

Behavioral analysis

Simulated foraging with a live rat encounter predator encounter

Behavioral measures including overall performance, speed of movement and a variety of movement parameters were tracked using high speed video at 100 Hz, over the course of the 3 days. Analog traces from the treadmill encoder, and the movement sequences of the mice were captured on two Basler cameras, one aimed at the body, and one aimed at the face. Movement and posture of the mice was tracked using DeepLabCut (Mathis et al. 2018). Behavioral phenotypes that were available for analysis in the control condition and in the presence of the rat included: 1) Flight, apparent as a backward movement on the treadmill, and here detected specifically when a mouse moved at least 3cm backwards when directly in front of the tube. This movement pushed the tube holding the rat away from the mouse and showed up as negative deflection in the encoder output and led to a failure in licking. 2) Freezing, i.e. the mouse stops moving, which would be apparent in the speed traces and lead to a failure of the trial. This was detected here by either failing to complete a trial in time, or by failing to lick during the reward phase, when the mouse was directly in front of the tube. 3) Changes in mouse posture, hunching or stretching. 4) Changes in pupil size, detected through constriction or dilation of a circle fitted to the DeepLabCut-tracked edge of the pupil. 5) Changes in nose movement, which were also tracked with keypoints detected with DeepLabCut. Light conditions were verified to be generally stable and unrelated to pupil size in our recordings.

We performed the experiments in such a way that we were able to pair the behavioral phenotype of each mouse on the first recording day (before a rat was introduced) as a baseline to all further recordings of the same mouse.

Assaying threat response in freely moving mice

One of the three assays, the looming stimuli, was dependent on vision, two of them had
olfactory components.   In these experiments —looming, rat odor avoidance, and live rat
exposure— the position of the mice was tracked using SLEAP (version 1.3.3). Three key body
points were annotated along the midline of each mouse. For positional tracking, the pixel
coordinates of the neck point were extracted and transformed into millimeter-based coordinates
relative to the layout of each experimental arena. To reduce noise, positional data were clipped to
the arena boundaries and smoothed using a sliding half-second window. Due to the side-view
recording setup in the looming experiment (necessitated by an overhead screen), additional
geometric corrections were applied. The X-coordinate was taken directly from the neck point,
while the Y-coordinate was defined as the lowest Y-value among the three tracked points. An
empirical downward shift of 20 pixels was then applied to the Y-coordinate. The adjusted
coordinates were subsequently clipped to the nearest point within the arena polygon visible in each
frame. After these corrections the pixel-based position is converted to millimeter-scale
coordinates.

Because the perspective of the looming videos enforced a strong correlation between the
X- and Y-position, only the X-position of the mouse relative to the shelter was used for further
analysis. The absolute X-axis speed (relative to the shelter) was smoothed with an 11-frame
window before being used to classify looms as either "fleeing" or "non-fleeing" events.

Statistical analysis

Simulated foraging with live a rat

All statistical comparisons for eye position, pupil diameter, locomotor speed, posture, and
facial movement were performed within the animal, comparing each mouse's baseline session to
its corresponding rat-exposure session. Time-resolved analyses were summarized into half-second
bins, but statistical testing was conducted at the level of per-animal distributions rather than
treating individual trials or bins as independent samples. Thus, each mouse served as its own
control, avoiding pseudoreplication due to within-subject correlations. For time-resolved analyses,
significance thresholds were adjusted for multiple comparisons across bins using a Bonferroni
correction corresponding to the number of consecutive bins tested.

Here we first assayed mouse behavior for evidence of innate fear (i.e. freezing, fleeing)
which would be evident in performance. This study was also designed to capture subtle changes
in the behavior of mice (speed of movement as mice approach rats, success rate, pupil size, body

elongation, nose movement) in the presence of rats. To assay these changes in behavior, the data

related to each behavioral feature (pupil size, speed of movement, nose movement, etc.) were first

filtered with a rolling z-score across a one second window. Values with an absolute score above

three were removed. The time series data was then smoothed with a half second window by way

of a rolling mean.

In the case of the video data (DeepLabCut output) it was also necessary to perform initial

filtering steps, according to the detected likelihood that a feature was discovered in a frame, and

with an initial z-score filtering across the x- and y-position of the features. In statistics where we

considered the pupil-diameter and pupil position, we also filtered the eight detected markers

around the pupil with a modified z-score (Iglewicz and Hoaglin 1993).

We compared the behavioral items pairwise per mouse, between the day of the baseline

recording and the following days. To achieve this, we summarized half-second wide bins relative

to the beginning of the trial or the delivery of reward of the time series and compared these between

the two conditions. These data were plotted to show the mean and standard deviation per feature

and condition. Binned data for sessions with and without the rat were compared, and the difference

between the mean and standard deviation and the significance of this shift shown as a bar plot.

Significance was assessed for each half second bin. Additionally, for each behavioral parameter

(i.e. position, speed pupil size, eye position) differences between sessions with and without the rat

had to be significantly different for 6 consecutive bins. In the reward phase, which lasted for a

short duration, there had to be a significant difference over at least 3 consecutive half-second bins

for the pupil x-position, pupil diameter, body length, and nose speed.

The continuous mean and standard deviation of the time series was calculated with the

formulas $\mu = \frac{\sum_{i=1}^{n} x_i}{n}$ and $\sigma = \sqrt{\frac{1}{n}\sum_{i=1}^{n} (x_i - \mu)^2}$. These values were then used to calculate the

difference between both statistics. We used the following formulas for this: $\mu_{diff} = \mu_{minuend} - $

$\mu_{subtrahend}$ and $\sigma_{diff} = \sqrt{\sigma_{minuend}^2 + \sigma_{subtrahend}^2}$, with the subtrahend always being the baseline

condition and the minuend representing either the rat encounter condition or the measurements

from the day after the simulated rat encounter. To determine the significance of the shifts, we used

the Mann-Whitney U-Test. The original thresholds in the figures were *=0.05, **=0.01,

***=0.001, and n.s. = not significant, which were then applied with a Bonferroni-correction

corresponding to the number of bins considered in each subplot.

For the summary data, multiple mice into one group, the means and standard deviations of

their individual bins were combined, under the assumption of independence between mice, with

the formulas $\mu = \frac{\sum_{i=1}^{n} \mu_i}{n}$ for the mean and $\sigma = \sqrt{\frac{\sum_{i=1}^{n} \sigma_i}{n}}$ for the standard deviation of the combined

bin. The p-values were averaged, weighted by the inverse of the variance of the individual bins

they originated from. This happened according to the formula $p_{group} = \frac{\sum_{i=1}^{n} \frac{1}{\sigma_i^2} \cdot p_i}{\sum_{i=1}^{n} \frac{1}{\sigma_i^2}}$, with n

corresponding to the individuals in the group. Due to the high number of single samples per bin,

the estimate of significance was limited to the ethologically relevant threshold of 10% of the 5^{th} to

95^{th} percentile span of each behavioral feature, across animals and conditions.

Data analysis for freely moving mice

All analyses were performed blind to the source of the mice. When shown, the unblinded

mouse ID follows the format "C{Cohort #}{Source}{Mouse #}," where the source is indicated as

C (Charles River), I (Internal), or J (Janvier). For example, the ID *C2J1* refers to the first mouse

from Janvier in cohort 2. No significant behavioral differences were observed among mice from

the three sources. Detailed categorizations by experiment, cohort, and source are presented in

Supplementary Figure 10.

Mice were classified as "fleeing" or "non-fleeing" based on the median of all loom-wise

ratios comparing the mean speed during the 5-second looming visual stimulus to the mean speed

in the subsequent 5 seconds. A threshold ratio of 2 was chosen as a conservative criterion for

identifying clear fleeing behavior.

The same threshold (a two-fold change) was used to conservatively identify mice that

exhibited significant changes in location preference in the odor-based experiments. Specifically,

this included avoidance or preference for the rat-odor side of the arena, as well as avoidance of the

region adjacent to the mesh in the live rat exposure condition. Both behaviors relate to avoidance

of rats as a defensive response. Mice falling below this threshold were classified as behaviorally

unchanged.

Data analysis Software

SLEAP

Two different SLEAP (Pereira et al. 2022) bottom-up models (version 1.3.3) were trained

to track the location of the mice in the verification experiments. The side view model for videos

recorded in the looming stimulus experiments was trained on 225 masked frames (to crop the

moving looming stimulus and reduce the complexity of the video). The top view model for videos

recorded in the rat odor and presence experiments was trained on 430 frames.

DeepLabCut

A series of DeepLabCut models (version 2.1; Mathis et al. 2018) were trained to track key

features of mice for behavioral analysis in the simulated foraging task. Separate models were

trained for each anatomical landmark: **Body key points** (nose, shoulder, tail base): trained on 220

labeled frames for one cycle of 1,030,000 iterations. **Nose key points** (nose ridge, nose base, nose

tip): trained for two cycles of 1,030,000 iterations each. The first cycle used 220 labeled frames;

the second used 337. **Eye key points** (eight equidistant points around the pupil, starting from the

12 o'clock position, plus the left and right corners of the eye): trained over three cycles of 1,030,000

iterations each. The training sets consisted of 260, 466, and 629 labeled frames, respectively. The

networks were trained across animals and sessions, when performance was lower in a video we

added more frames from that video to the training data.

                                          Results

   Simulated Foraging

This paradigm was designed to be easy for mice to perform but was also designed to

minimize stress for both rats and mice. Once mice were habituated to head restraint and to

obtaining reward by moving the treadmill (**Figure 1A,B,C**) they could perform 50-100 trials in a

day, obtaining a total reward of 0.5 ml in a ~30-45-minute session. When mice started a trial, the

lickspout was at a starting position ~30 cm from the mouse (**Video 1**, **Figure 1D**). A go cue, the

flashing of the virtual reality monitors, and a sound cue initiated the trial. A successful trial was

one in which mice moved the treadmill 28.5cm in 12 seconds, with the last 1.5 cm of movement

controlled automatically within the code. Two to four seconds after reward delivery, depending on

the start of licking, the lick spout and red tube over it were reset to their starting position.

Well trained, motivated mice started running on the tread mill immediately after licking

the reward, even before the lick spout moved back to starting position. As mice improved their

daily performance, adult female rats were habituated to being handled and to the apparatus. Once

a cohort of mice were stable in their performance for 3 successive days, reaching the threshold

criteria in their performance, the experimental data was collected over 3 consecutive days. On the

first day control data was collected, on the second day the rat was introduced into the behavioral

paradigm. The third day was a post-rat control session (**Figure 1E**). To maximize the potential for

interaction between the head-fixed mouse and restrained rat, we used only one setting, open, on

the disk that covered the mouth of the tube holding the rat. Note that in this setting mice could

almost touch the rat on each trial and mice could see, smell and hear the rat moving in the tube as

it came closer to the mouse. On each trial, with every mouse, rats could stick their nose out of the

tube. When rats were in this position, their nose was effectively just above the lickspout for mice

(**Video 1**). In the following plots the male mice are enumerated as mouse 1-4, and the female mice

as mouse 5-7.

Performance

Performance of each mouse was analyzed for three consecutive days: control / baseline

day, day with the rat, and a post-rat day. Surprisingly, the success rate of most mice (5/7 mice)

was unaffected (stayed above criterion for all three days) by the introduction of the rat (**Figure**

1F). Note that across the cohort of animals, performance was stable across days for the majority

of animals. Five of seven mice maintained success rates above criterion (>65%) on the baseline

day, during rat exposure, and on the post-rat day, with no systematic increase in trial failures or

missed rewards when the rat was present (**Figure 1F**). In contrast, two mice showed a marked

reduction in success rate during the rat session (>25% decrease relative to baseline), which

persisted into the post-rat day, indicating a stable, individual-specific effect rather than transient

trial-by-trial disruption. Importantly, the presence of the rat did not induce consistent hesitation at

trial onset or prolonged latencies to initiate treadmill movement across the cohort; instead, changes

in performance were confined to a small subset of animals. Next, we examined whether mice

changed their behavior when the rat was introduced.

Movement speed on treadmill

One measure of performance is the rate of success; another measure was the speed with

which mice moved the treadmill (**Figure 2; Video 2**). Mice could run or walk slowly and

consistently to cover 28.5 cm in 12 seconds. Most mice learned to move at a consistent speed of

around 0.2 m / s. This speed was mostly uniform and on average it stayed constant for most of the

trial (**Figure 2B**), but the speed decreased abruptly as mice stopped moving to lick the reward

spout (**Figure 2B**).

One issue that arose in monitoring and comparing the behavior of mice from day to day,

was whether the position of the mouse on the treadmill changed across days. It was possible that

mice were positioned at slightly different heights relative to the treadmill, and this effectively

changed the speed with which mice moved (**Figure 2B**). To examine this, we selected frames from

different time points during each session and examined them for any obvious differences in height

of the mouse relative to the treadmill (**Figure 2A; Supplementary Figure 9**). There were no

significant differences in the positioning of mice from day to day.

Next, we compared the speed of movement from control sessions (**Figure 2B**, green traces)

and sessions when the rat was present for individual mice. These data show that while the majority

of mice show no consistent significant changes in their movement speed in presence of the

potential predator (**Figure 2B**, blue traces), three out of seven mice showed significant ($p<0.05$,

binwise Mann-Whitney U (MWU) test , Bonferroni Correction) and consistent changes in their

movement speed throughout the session (**Figure 2C**). Their average speed was significantly

different at the onset of the trials when the rat was present. The two mice that decreased their

movement speed in the presence of the rat were slow on the treadmill from the beginning of the

trial. One mouse ran significantly faster in the presence of the rat. In the remaining mice, there

were no consistent changes in average speed (**Supplementary Figure 3, 4**). Thus, while the

presence of the rat altered locomotor speed in a subset of mice, these effects were directionally

heterogeneous (slowing in two mice, speeding in one) and did not reflect a uniform suppression

of movement or task engagement.

Eye movement and pupil diameter

To assess whether mice attend to the presence of the rat, we used DeeplabCut to track eye

movement and pupil diameter on the control day and on the day that the rat was introduced into

the tube (**Figure 3, Video 3**). We plotted the average position of the pupil in the horizontal axis

relative to the two corners of the eyes, over the course of the control session and in the presence

of the rat (**Figure 3A-C**). In five out of seven mice, there was a significant (p< 0.05, binwise

MWU, Bonferroni Correction) change in the horizontal position of the pupil in the presence of the

rat (**Figure 3B**, green traces); in the remaining 2 there were no consistent changes in eye position.

On average, the position of the eyes of five mice was significantly different (p < 0.05, binwise

Mann Whitney U Test, Bonferroni correction) during their reward phase compared to the control

days (**Supplementary Figure 5**). When the rat was present in the tube and the mice were directly

in front of the tube, on average, 3 mice positioned their eyes more in the direction of the rat (looked

right) and 2 looked away from the rat. These effects were significant and were evident throughout

the trial: across mice that showed significant effects, horizontal eye position shifted by an average

of ~1-3% of the half width of the eye relative to baseline. Two mice showed no significant change

in their eye position (**Figure 3C**, right panels).

Next, we examined pupil size (**Figure 3D-F; Video 4**). Pupil size changes with changes in

lighting, or changes in the parasympathetic or sympathetic system. To establish that light around

the face and head of the mouse remained consistent, we measured brightness and light intensity

around the eye and at another point on the head. There was no effect on the luminance / light levels

around the eyes or head when the rat was introduced. Our data suggest that across mice that showed

a significant reduction, pupil diameter decreased by ~10-13% during successful approaches, with

these changes remaining stable across trials within a session (Fig. 3; Supplementary Figs. 5–6).

But in the presence of the rat there were significant (p<0.05, binwise MWU, Bonferroni

correction) changes in pupil diameter in five out of seven mice (**Figure 3D, Supplementary**

Figure 6), especially when mice were close to the rat (Blue traces, **Figure 3E**). In these mice, the

pupil diameter was consistently smaller throughout the course of the trial and in four out of five of

these mice the change in pupil size persisted into the next session. The other 2 mice showed no

significant changes in their pupil diameter. Note that the reported significance values reflect

within-animal comparisons between baseline and rat sessions, with each mouse treated as an

independent observational unit. Taken together, these data show that mice detect and respond to

the predator, with pupil dynamics consistent with an altered attentional or autonomic state.

Posture and facial movements

To assess whether any other aspects of mouse posture or facial movement were affected

by the presence of the rat, we measured mouse posture, estimated by body-length, and facial

movements (**Figure 4A-C; Video 5, Supplementary Figure 7, 8**). In three out of seven mice there

was a significant change in the pose when the mouse was confronted by the rat (blue). Additionally,

in 2 other mice there was a significant increase in nasal movement speed in the presence of the rat

(**Figure 4D-F; Video 6**).

Taken together, these results show that although head-fixed mice do not express canonical

escape behaviors, the presence of a live rat reliably alters multiple behavioral and physiological

variables, including locomotor dynamics, eye position, pupil diameter, posture, and facial

movements (**Supplementary Figure 10**). These changes indicate that mice detect and respond to

the predator, but express this response through a constrained behavioral repertoire rather than

through overt flight or freezing. Note that although both male (n = 4) and female (n = 3) mice were

included in the head-fixed experiments, no consistent sex-specific differences were observed in

task performance, locomotor speed, eye movements, pupil responses, or posture; given the small

sample size, sex was not treated as an independent factor in statistical analyses.

Next, we considered whether head fixation, thirst, and the need to forage affected threat

perception or the mouse's behavioral response to the threat. At the same time, we also assessed

whether the mice in our colony were inherently less fearful. To address these issues, we used naive

freely moving mice, from our colony and from two other sources.

Effect of looming stimuli

In these experiments, we used 36 naive mice, 12 mice were from our animal facility, 12

newly acquired from Charles River, and 12 from Janvier. Mice obtained from our inhouse facility

showed no differences in behavior compared to mice obtained from external sources.

Mice were placed in an arena (**Figure 5A, B**), and when they entered a particular location
in the arena, the looming protocol was initiated. A dark shadow emerged, and grew larger, hovered
over the mouse, simulating a bird of prey swooping over the mouse. For each mouse, the looming
stimulus protocol was repeated 14 times. Post-hoc video analysis showed that none of the mice
froze in response to the looming stimulus; seven out of 36 mice ran into the shelter (**Figure 5C,
D**). The seven mice that ran into the shelter moved rapidly toward the shelter during the looming
stimulus (**Figure 5E, F**), the remaining 29 mice did not show any significant, consistent movement
toward the shelter. Overall, these results suggest that looming stimuli can reliably evoke fleeing in
mice, but this effect is only observed in ~19% (7 out of 36, Wilson 95% confidence interval of
~10–35%) of the mice.

Avoidance of rat odors or rats

Next, we examined whether freely moving mice (n = 36) reacted to rat odors (Figure 6A-
C). Mice were placed in an arena with a partition down the middle, a partition that mice could and
did traverse. Five minutes after putting mice in the arena, they were removed from the arena. A rat
was introduced on one side of the partition, a partition that rats could not traverse. Before mice
were returned to the arena, the rat was removed. Post-hoc analysis of the amount of time spent in
each portion of the arena, revealed that most mice either preferred the side that had contained the
rat (6 mice out of 36), or showed no clear preference for either side of the arena (29 mice out of
36) (Figure 6C).

We used the same arena but closed off the wire mesh partition that separated the arena into
two halves (**Figure 7A, B**). Once again naive mice were placed in the arena but were allowed to
explore only one half of the arena, then they were removed, and a rat was placed on the right side
of the arena. Mice were then returned to the arena. In these experiments, half the mice (n=18)
avoided being close to the wire mesh that separated the rat from the mice (**Figure 7C-E**). Of the
other half (n=18) fifteen mice showed no preference for one side and three mice preferred staying
close to the mesh. These data indicate that exposure to a live rat does not elicit a uniform, automatic
defensive response across mice. Instead, freely moving mice exhibit substantial inter-individual
variability, including avoidance, investigatory behavior, and neutral exploration, consistent with
context-dependent threat evaluation rather than a fixed innate response.

Discussion and conclusion

Defensive behaviors such as freezing, fleeing, or fighting are often treated as automatic responses to predator threat. Our results challenge this assumption by showing that predator exposure does not necessarily elicit an invariant defensive repertoire in mice, even under ethologically relevant conditions. Across both head-fixed and freely moving paradigms, mice frequently did not express stereotyped defensive actions yet showed clear evidence of threat engagement through changes in locomotion, gaze direction, pupil diameter, and posture. Together, these observations suggest that defensive behavior can be expressed flexibly and is not obligatorily triggered by predator cues.

In the head-fixed simulated foraging task, mice were physically unable to flee, and freezing would have resulted in trial failure and loss of reward. Under these constraints, most mice continued to perform the task despite close proximity to a live rat. Importantly, continued locomotion did not imply an absence of threat detection. Instead, individual mice exhibited reliable changes in multiple behavioral and physiological variables, including altered treadmill speed, systematic shifts in eye position, sustained pupil constriction, and changes in posture and facial movements. These measures indicate that mice detected and responded to the predator, but their response was restricted by head fixation and competing motivational demands of reward acquisition versus threat avoidance. Although the number of head-fixed animals was necessarily limited, each mouse contributed a large number of trials per condition, enabling sensitive within-animal comparisons, while constraining population-level generalizations.

This interpretation helps reconcile our findings with previous work. Vollmer et al. (2022) reported robust suppression of reward-seeking in head-fixed mice exposed to predator-related odors (e.g., TMT) under conditions in which animals were stationary, had ad libitum access to food and water, and reward-seeking was operationalized as discrete lever presses. In contrast, mice in our paradigm were water restricted, actively controlled their approach toward the stimulus, and were exposed to a live predator rather than an olfactory cue alone. Reward acquisition additionally required sustained locomotion followed by licking, rather than a single action. These differences in motivational context, threat modality, and action requirements are likely to strongly influence how defensive states interact with reward-seeking behavior, indicating that suppression of reward seeking under threat is context dependent rather than obligatory.

The freely moving experiments in this study provide an important complementary
reference point. Using established ethological assays—including looming stimuli, predator odor,
and live predator presence—we observed pronounced inter-individual variability and strong
context dependence, with defensive responses expressed in only a subset of animals. These
experiments therefore serve as a positive control, indicating that the absence of uniform freezing
or escape in the head-fixed paradigm does not reflect insensitivity to threat. To assess whether
head fixation, motivational state, or colony-specific factors contributed to this pattern (Burnett et
al. 2016; Verma et al. 2016), we tested freely moving, well-fed mice drawn from multiple sources.
Even under these conditions, looming stimuli elicited escape behavior in only a minority of
animals (Lenzi et al. 2022), predator odor did not produce consistent avoidance (Banik and Anand
2011), and exposure to a live rat resulted in avoidance in approximately half of the mice (Karli
1956; **Supplementary Figure 11**). Together, these findings mirror the heterogeneity observed in
the head-fixed paradigm and argue against a simple, hardwired mapping between predator cues
and defensive motor programs.

The observed pupil constriction in the presence of a predator was unexpected, as threat is
often associated with dilation. However, pupil size reflects a balance between sympathetic and
parasympathetic activity and is sensitive to attentional state. In this context, constriction may
reflect heightened visual focus or vigilance during close-range threat monitoring rather than
reduced arousal. This interpretation is consistent with the idea that mice adopt alternative
physiological strategies when classical defensive actions are unavailable or maladaptive.

As with many behavioral paradigms, responses to predator-related stimuli are shaped by
prior experience, social history, and internal state (Yilmaz and Meister 2013; Lenzi et al. 2022;
Wang 2020; Qi et al. 2025). Although we followed established protocols closely (Yilmaz and
Meister 2013; Banik and Anand 2011; Karli 1956), differences in housing, handling, and
individual experience are known to influence defensive expression. In earlier looming paradigms,
high rates of freezing or escape were reported, but these effects depended strongly on stimulus
parameters and arena structure, such as the availability of shelter. Defensive responses also vary
with threat history and repeated exposure, including rapid habituation or suppression across trials
(Yilmaz and Meister 2013; Lenzi et al. 2022; Qi et al. 2025). Together with prior work showing
experience- and history-dependent modulation of looming responses, our results indicate that

variability in defensive expression reflects flexible strategy selection rather than experimental
noise.

623   **Caveats and issues to consider**: In the present study, mice were socially isolated for
624 several days before testing and were drawn from multiple sources, yet defensive responses
625 remained limited and heterogeneous across a relatively large cohort (n = 36). Rather than treating
626 this variability as experimental noise, our results suggest it is a fundamental feature of defensive
627 behavior, reflecting flexible strategy selection shaped by context and experience rather than a
628 reflexive, invariant output.

629   Because the same cohort of mice was tested across paradigms, experience-dependent
630 carryover effects cannot be fully excluded. However, if such effects were present, prior exposure
631 to predator-related stimuli would be expected to enhance, rather than suppress, subsequent
632 defensive responses. The limited and heterogeneous defensive behaviors observed across tasks
633 therefore argue against sensitization-driven inflation of fear responses and support the conclusion
634 that defensive expression is strongly context dependent.

635   Note that in the head-fixed paradigm, the identity of the rat placed in the tube was rotated
across sessions to minimize systematic effects of individual rats. In the freely moving experiments,
rats were selected at random from a small set of animals (three rats in cohort 1 and two rats in
cohort 2). Differences in rat behavior could, in principle, influence the salience of the predator
stimulus and thus represent a potential source of variability. However, we observed no qualitative
or quantitative differences in mouse behavioral outcomes between the two freely moving cohorts
(**Figures 5–7C**), despite the use of different rats. This consistency argues against rat identity being
a major determinant of the observed patterns of defensive behavior and suggests that the reported
effects reflect mouse-specific and context-dependent factors rather than systematic differences in
predator behavior.

645   It is possible that some aspects of how we habituate, or house animals or how animal
646 caretakers look after mice reduce fear and stress or reduce the behavioral expression of fear in
647 mice (Gouveia and Hurst 2017; Furlong et al. 2016; Kallnik et al. 2007). It is also possible that
648 just as in rats, the sexual identity of mice affects how they respond in our behavioral paradigms
649 (Gruene et al. 2015).

650   A key limitation of head-fixed paradigms is the restriction of overt defensive actions such
as escape or freezing. Even though mice could have stopped walking forward, which would have

the effect of keeping the rat and the reward at distance, or could have walked backward to push

the rat away, they did not. Rather than viewing this as a failure of the model, we emphasize that

this constraint enables a complementary question: whether threat detection and defensive state

expression necessarily require these actions. Our findings suggest they do not. Even when classical

motor outputs are unavailable or maladaptive, mice engage alternative behavioral and

physiological strategies consistent with active threat assessment and coping.

Our study challenges simplified views of innate fear as a fixed behavioral program and

highlights the importance of considering how environment and task demands shape the expression

of defensive states.

Table of abbreviations

Abbreviation	Full Term
MWU	Mann-Whitney U Test

Acknowledgements

We thank the Charité Workshop for technical assistance, especially Alexander Schill, Jan-
Erik Ode and Daniel Deblitz. We also want to thank Melissa Long and the team of the ABPF for
their help with animal handling and room planning. Finally, we thank Laura Schwarz of the
Sainsbury Wellcome Centre and members of the Larkum lab for useful discussions about earlier
versions of this manuscript.

Data availability statement

The original contributions presented in the study are publicly available. This data can be
found here: http://doi.org/10.5281/zenodo.15264865. Extended data is available on request.

Code availability statement

The code used to generate the data shown in this study is publicly available. It can be found
here: https://github.com/Marti-Ritter/contextual-modulation-and-blunted-defensive-responses-to-
predators.

Ethics statement

All experiments were conducted in accordance with the guidelines of animal welfare of the
Charité Universitätsmedizin Berlin and the local authorities, the 'Landesamt für Gesundheit und
Soziales'.

Author Contributions

	Conceptualization	Data curation	Formal analysis	Funding acquisition	Investigation	Methodology	Project administration	Resources	Software	Supervision	Validation	Visualization	Writing – original draft	Writing – review & editing
MR	x	x	x		x	x	x		x	x	x	x	x	x
LC	x	x			x	x								x
LS	x	x			x	x								x
AH	x	x			x	x				x				x
SÖ		x			x						x			x
RB	x					x				x				x
AV	x	x			x	x		x						x
DS	x			x										x
PP	x			x										x
ML	x				x		x	x						x
RS	x			x		x	x	x		x			x	x

Funding

The following funding sources have supported this project: (1) Deutsche

Forschungsgemeinschaft (DFG), Grant Nos. 246731133, 250048060 and 267823436 to ML; (2)

DFG Project number 327654276 – SFB 1315 to ML; (3) European Commission Horizon 2020

Research And Innovation Program and Euratom Research and Training Program 2014–2018

(under grant agreement No. 670118 to ML); (4) Human Brain Project, EU Commission Grant

720270 (SGA1), 785907 (SGA2) and 945539 (SGA3) to ML; (5) Einstein Foundation Berlin EVF-

2017-363 to ML; Einstein Foundation Visiting Fellowship EVF-2019-508 to PP; Humboldt

Foundation Friedrich Wilhelm Bessel Research Award to PP.

Conflict of interest

The authors have declared no conflicts of interest.

Generative AI statement

The authors declare that no Generative AI was used in the creation of this manuscript.

References

Abs, Elisabeth, Rogier B. Poorthuis, Daniella Apelblat, et al. 2018. "Learning-Related Plasticity
in Dendrite-Targeting Layer 1 Interneurons." *Neuron* 100 (3): 684-699.e6.
https://doi.org/10.1016/j.neuron.2018.09.001.

Ahmad, Hammad, Khalid Rauf, Wahid Zada, et al. 2020. "Kaempferol Facilitated Extinction
Learning in Contextual Fear Conditioned Rats via Inhibition of Fatty-Acid Amide
Hydrolase." *Molecules* 25 (20): 4683. https://doi.org/10.3390/molecules25204683.

Banik, Avijit, and Akshay Anand. 2011. "Loss of Learning in Mice When Exposed to Rat Odor:
A Water Maze Study." *Behavioural Brain Research* 216 (1): 466–71.
https://doi.org/10.1016/j.bbr.2010.07.035.

Beckers, Tom, Angelos-Miltiadis Krypotos, Yannick Boddez, Marieke Effting, and Merel Kindt.
2013. "What's Wrong with Fear Conditioning?" *Biological Psychology* 92 (1): 90–96.
https://doi.org/10.1016/j.biopsycho.2011.12.015.

Blanchard, Robert J., and D. Caroline Blanchard. 1989. "Attack and Defense in Rodents as
Ethoexperimental Models for the Study of Emotion." *Progress in Neuro-*
Psychopharmacology and Biological Psychiatry 13 (January): S3–14.
https://doi.org/10.1016/0278-5846(89)90105-X.

Blanchard, Robert J., Mark A. Hebert, Pier Ferrari, et al. 1998. "Defensive Behaviors in Wild
and Laboratory (Swiss) Mice: The Mouse Defense Test Battery." *Physiology & Behavior*
65 (2): 201–9. https://doi.org/10.1016/S0031-9384(98)00012-2.

Burnett, C. Joseph, Chia Li, Emily Webber, et al. 2016. "Hunger-Driven Motivational State
Competition." *Neuron* 92 (1): 187–201. https://doi.org/10.1016/j.neuron.2016.08.032.

Dalmay, Tamas, Elisabeth Abs, Rogier B. Poorthuis, et al. 2019. "A Critical Role for Neocortical
Processing of Threat Memory." *Neuron* 104 (6): 1180-1194.e7.
https://doi.org/10.1016/j.neuron.2019.09.025.

Fanselow, Michael S., and Zachary T. Pennington. 2017. "The Danger of LeDoux and Pine's
Two-System Framework for Fear." *American Journal of Psychiatry* 174 (11): 1120–21.
https://doi.org/10.1176/appi.ajp.2017.17070818.

Farmer-Dougan, Valeri, Seshanand Chandrashekar, David Stutzman, Kari Bradham, and James
D. Dougan. 2005. "Fox Urine as an Aversive Stimulus: Modification of a Passive
Avoidance Task." *The Journal of General Psychology* 132 (3): 313–20.
https://doi.org/10.3200/GENP.132.3.313-320.

Furlong, Teri M., Rick Richardson, and Gavan P. McNally. 2016. "Habituation and Extinction of
Fear Recruit Overlapping Forebrain Structures." *Neurobiology of Learning and Memory*
128 (February): 7–16. https://doi.org/10.1016/j.nlm.2015.11.013.

Gibson, Eleanor J., and Richard D. Walk. 1960. *The "Visual Cliff."* April.

Gouveia, Kelly, and Jane L. Hurst. 2017. "Optimising Reliability of Mouse Performance in
737   Behavioural Testing: The Major Role of Non-Aversive Handling." *Scientific Reports* 7
738   (1): 44999. https://doi.org/10.1038/srep44999.

739   Gruene, Tina M., Katelyn Flick, Alexis Stefano, Stephen D. Shea, and Rebecca M. Shansky.
740   2015. "Sexually Divergent Expression of Active and Passive Conditioned Fear
741   Responses in Rats." *eLife* 4 (November): e11352. https://doi.org/10.7554/eLife.11352.

742   Iglewicz, Boris, and David C. Hoaglin. 1993. *How to Detect and Handle Outliers*. ASQC Basic
743   References in Quality Control, v. 16. ASQC Quality Press.

744   Kallnik, Magdalena, Ralf Elvert, Nicole Ehrhardt, et al. 2007. "Impact of IVC Housing on
745   Emotionality and Fear Learning in Male C3HeB/FeJ and C57BL/6J Mice." *Mammalian
746   Genome* 18 (3): 173–86. https://doi.org/10.1007/s00335-007-9002-z.

747   Karli, P. 1956. "The Norway Rat's Killing Response to the White Mouse: An Experimental
748   Analysis." *Behaviour* 10 (1/2): 81–103. https://doi.org/10.1163/156853956X00110.

749   LeDoux, Joseph E., and Richard Brown. 2017. "A Higher-Order Theory of Emotional
750   Consciousness." *Proceedings of the National Academy of Sciences* 114 (10): E2016–25.
751   https://doi.org/10.1073/pnas.1619316114.

752   Lenzi, Stephen C., Lee Cossell, Benjamin Grainger, Sarah F. Olesen, Tiago Branco, and Troy W.
753   Margrie. 2022. "Threat History Controls Flexible Escape Behavior in Mice." *Current
754   Biology* 32 (13): 2972-2979.e3. https://doi.org/10.1016/j.cub.2022.05.022.

755   Lovett-Barron, Matthew, Patrick Kaifosh, Mazen A. Kheirbek, et al. 2014. "Dendritic Inhibition
756   in the Hippocampus Supports Fear Learning." *Science* 343 (6173): 857–63.
757   https://doi.org/10.1126/science.1247485.

758   Maren, Stephen, K. Luan Phan, and Israel Liberzon. 2013. "The Contextual Brain: Implications
759   for Fear Conditioning, Extinction and Psychopathology." *Nature Reviews Neuroscience*
760   14 (6): 417–28. https://doi.org/10.1038/nrn3492.

761   Mathis, Alexander, Pranav Mamidanna, Kevin M. Cury, et al. 2018. "DeepLabCut: Markerless
762   Pose Estimation of User-Defined Body Parts with Deep Learning." *Nature Neuroscience*
763   21 (9): 1281–89. https://doi.org/10.1038/s41593-018-0209-y.

764   McDannald, Michael A. 2023. "Pavlovian Fear Conditioning Is More than You Think It Is."
765   Viewpoints. *Journal of Neuroscience* 43 (48): 8079–87.
766   https://doi.org/10.1523/JNEUROSCI.0256-23.2023.

767   Mobbs, Dean, Ralph Adolphs, Michael S. Fanselow, et al. 2019. "Viewpoints: Approaches to
768   Defining and Investigating Fear." *Nature Neuroscience* 22 (8): 1205–16.
769   https://doi.org/10.1038/s41593-019-0456-6.

Mongeau, Raymond, Gabriel A. Miller, Elizabeth Chiang, and David J. Anderson. 2003. "Neural
Correlates of Competing Fear Behaviors Evoked by an Innately Aversive Stimulus." *The*
Journal of Neuroscience 23 (9): 3855–68. https://doi.org/10.1523/JNEUROSCI.23-09-
03855.2003.

Mowrer, O. H., and R. R. Lamoreaux. 1946. "Fear as an Intervening Variable in Avoidance
Conditioning." *Journal of Comparative Psychology* 39 (1): 29–50.
https://doi.org/10.1037/h0060150.

Panksepp, Jaak. 1971. "Aggression Elicited by Electrical Stimulation of the Hypothalamus in
Albino Rats." *Physiology & Behavior* 6 (4): 321–29. https://doi.org/10.1016/0031-
9384(71)90163-6.

Pereira, Talmo D., Nathaniel Tabris, Arie Matsliah, et al. 2022. "SLEAP: A Deep Learning
System for Multi-Animal Pose Tracking." *Nature Methods* 19 (4): 486–95.
https://doi.org/10.1038/s41592-022-01426-1.

Phillips, R. G., and Joseph E. LeDoux. 1992. "Differential Contribution of Amygdala and
Hippocampus to Cued and Contextual Fear Conditioning." *Behavioral Neuroscience* 106
(2): 274–85. https://doi.org/10.1037//0735-7044.106.2.274.

Qi, Haoxuan, Alice Treloar, Greg J. Stuart, and Saba Gharaei. 2025. "Context Shapes Fear:
Habituation of Innate Defensive Behaviours Depends on Environmental Context."
Preprint, bioRxiv, August 21. https://doi.org/10.1101/2025.08.21.671638.

Schiff, William. 1965. "Perception of Impending Collision: A Study of Visually Directed
Avoidant Behavior." *Psychological Monographs: General and Applied* 79 (11): 1–26.
https://doi.org/10.1037/h0093887.

Silva, Bianca A., Cornelius T. Gross, and Johannes Gräff. 2016. "The Neural Circuits of Innate
Fear: Detection, Integration, Action, and Memorization." *Learning & Memory* 23 (10):
544–55. https://doi.org/10.1101/lm.042812.116.

Tovote, Philip, Jonathan Paul Fadok, and Andreas Lüthi. 2015. "Neuronal Circuits for Fear and
Anxiety." *Nature Reviews Neuroscience* 16 (6): 317–31. https://doi.org/10.1038/nrn3945.

Verma, Dilip, James Wood, Gilliard Lach, Herbert Herzog, Guenther Sperk, and Ramon Tasan.
2016. "Hunger Promotes Fear Extinction by Activation of an Amygdala Microcircuit."
Neuropsychopharmacology 41 (2): 431–39. https://doi.org/10.1038/npp.2015.163.

Vollmer, Kelsey M., Lisa M. Green, Roger I. Grant, et al. 2022. "An Opioid-Gated
Thalamoaccumbal Circuit for the Suppression of Reward Seeking in Mice." *Nature*
Communications 13 (1): 6865. https://doi.org/10.1038/s41467-022-34517-w.

Wang, Hao, ed. 2020. *Neural Circuits of Innate Behaviors*. Vol. 1284. Advances in Experimental
Medicine and Biology. Springer Singapore. https://doi.org/10.1007/978-981-15-7086-5.

Yilmaz, Melis, and Markus Meister. 2013. "Rapid Innate Defensive Responses of Mice to
Looming Visual Stimuli." *Current Biology* 23 (20): 2011–15.
https://doi.org/10.1016/j.cub.2013.08.015.

Figure 1. *Experimental design, setup, and performance in the presence of a rat.* **A)** Side
view of the mechanical components in the apparatus. The movement of a treadmill was tethered
to the movement of a lick spout and a large red tube. When mice walked forward, the lick tube
moved towards them, when mice moved backwards, the lick tube moved away. A flashing screen
(part of a virtual reality display) and a sound cue signaled the start of a trial. Mice had 12 seconds
to complete a trial, if they succeeded, reward was delivered within reach of the mouse, and a new
trial was initiated by resetting the position of the spout to its starting position away from the mouse.
B) Threat perception disc. The apparatus was designed to mimic four user defined interactive
conditions with the large red tube positioned over the lick spout. The full view condition was one
where the opening of the tube was completely open, and it was possible for a head-fixed mouse
and the rat to almost touch each other. The olfactory condition was one where odors could be
delivered but visual, tactile and other elements inside the tube were blocked. The visual condition
was one where mice could see inside the tube. The opaque condition blocked all cues from inside
the tube. Note that auditory cues were not filtered and that a fan at the end of the tube could be
activated to extract smells from the red tube. **C)** Training procedures. Habituation consisted of
head fixing mice on the treadmill and manually rewarding them when they moved forward
spontaneously. Following this brief, 1- to 2-day habituation period, the reward delivery was
automated. Reward was dispensed after mice had moved 28.5 centimeters. This movement was
sufficient to move the lick spout into a position where mice could lick the reward as it was
delivered. Over days the flashing light cue of the virtual reality screens, and the sound cue were
added as go-cues to begin the trial. The duration of the trial was shortened, giving mice 12 seconds
to reach the spout, or move at least 28.5 cm. **D)** Data acquisition. Once all mice of the cohort
reached criterion (65% success rate) baseline data was acquired. In the next session, a live rat was
placed inside the tube, and the experiment was repeated, with the full interaction between the
mouse and the rat. Finally, on a third day, reference data in the absence of the rat were acquired.
E) Performance data. Of the seven mice used for these data sets, five showed no change in their
performance in any of the three sessions. For the other two, performance was affected by the
presence of the live rat. For both these mice, the reduction in performance persisted through the
next day, post rat. Vector drawings of rat and mouse adapted from SciDraw (Branco and Costa
2020; Scidraw 2020).

Figure 2. *Movement speed in the presence of rat.* **A**) Representative images of mouse position from three sessions. The images show the position of a mouse at 3 different times on three different sessions. These kinds of images were used to assess whether the head-fixation height or position of the mouse were stable from day to day. Plots for each of the seven mice are in **Supplement-figure 8**. **B**) Speed of movement on the treadmill. The median and confidence interval of the speed of movement in baseline, control day, recordings (black trace) and the session with the rat present (blue) show the two outcomes of having the rat hovering over the lick spout. The left plot (trial phase) shows no change in velocity throughout the trial. In another mouse there was a momentary reduction in speed at the beginning of the trials (right panel, trial phase). When the mice were at the lick spout, they stopped moving (see reward phase). Significance was assessed with a Mann-Whitney U-test (*: $p<0.05$, **: $p<0.01$, ***: $p<0.001$). See methods for additional filtering applied to significant effects. **C**) Pie chart grouping mouse behavior. Mice could be divided into two main groups, one that showed significant changes in speed and another that showed no change in speed. One mouse significantly increased its speed during the approach to the tube with the rat, 4 mice showed no significant change of speed while approaching the rat, and 2 slowed down significantly.

Figure 3. *Pupil movement and pupil size in the presence of the rat.* A) Schematic showing the analysis of horizontal pupil position. The x-position was defined relative to the center of the pupil and the corners of the eyes. Movement of the eyes could be nasally oriented (toward the rat) or oriented to the back of the head away from the rat. **B)** Plots for a mouse that positioned its eyes toward the lick tube (nasally) as it moved toward the lick spout (left plot) and positioned its eyes to look forward toward the rat, when the mouse was stopped at lick spout (right plot). The second set of plots on the right show eye position for a mouse which showed no significant change in x-position while in front of the rat. **C**) Pie chart grouping mice. Five mice positioned their eyes significantly differently in the presence of the rat (n=5). Two mice showed no change. Three mice looked toward the tube holding the rat more often, and two looked away from the rat. **D**) Schematic of the pupil diameter analysis. The diameter was calculated based on points labeled by DeepLabCut. **E**) Example plots from two mice, one showing a significant reduction in pupil diameter in presence of the rat (black trace is control session, blue trace was with rat present) the other one showing no effect. **F**) Pie chart grouping mice. Mice could be divided into two groups,

mice that showed no change in pupil diameter and a second group of five mice that had a consistent

reduction in pupil diameter in the presence of the rat.

Figure 4. *Changes in posture and facial movement in the presence of a rat.* **A**) Schematic

showing analysis of posture using a tail to nose distance. **B**) Body-length diagrams for two mice.

The mouse on the left showed no significant change in posture when in front of the rat, the one on

the right was significantly more hunched up in the presence of the rat. **C**) Pie chart grouping mice.

Mice could be divided into three groups: 4 mice showed no change in body length in the presence

of the rat, and 2 mice were more hunched up at the lick tube in the presence of the rat, and one

elongated its body at the lick spout in the presence of the rat. **D**) Schematic showing the analysis

of facial movement / nose speed. The nose position was detected with DeepLabCut and normalized

to the distance between the corners of the eye, to account for mouse size and position. **E**) Facial

movement. The left plots show data from a mouse that did not significantly change the movement

speed of its nose even when it was directly in front of the rat. The right-side plots show movement

data from a mouse that increased the movement speed of its nose significantly, during the reward

phase. **F**) Pie chart grouping mice. Most mice showed no significant change in nose speed even

when they were directly in front of the rat.

Figure 5. *Effect of looming stimuli on the behavior of freely moving mice*. **A)** Experimental

protocol for the looming stimulus paradigm. **B)** Representative annotated video frame. Black bars

denote frame boundaries, and the white area marks the masked region used to isolate the tracking

area. The transformed arena outline is overlaid. **C)** Scatterplot showing the median shelter-relative

speed for each mouse across all loom presentations. The behavior classification threshold (speed

ratio = 2) is indicated by a dashed black line. Individual cohorts are marked with distinct symbols,

and behavioral groups ("fleeing" vs. "non-fleeing") are color-coded. The global distribution of all

average speeds is shown as a background contour plot. Seven of 36 mice met the criterion for

consistent flight behavior (see Methods). We avoided defining a "freezing" group based on these

parameters, as the trial-wise speed distribution did not show a clean separation as it did for the

"fleeing" animals. **D)** Strip plot of average speed ratios per mouse. The median ratio for each

animal is shown as a black horizontal bar. Mice were classified as "fleeing" if their median ratio

exceeded 2. Numbers in brackets next to mouse IDs indicate how many individual looms exceeded

the plot's y-axis limit. Mouse *C1I3* completed only seven looms, having remained in the shelter

after the seventh presentation. **E)** Time-aligned traces of shelter-relative X-position for all loom

events from mice in the "fleeing" group. The median and 95% confidence interval are shown as a
bold line and surrounding shaded region, respectively. **F)** Same as in E, but for loom events from
mice in the "non-fleeing" group. Vector illustrations of the rat and mouse in this and subsequent
figures were adapted from SciDraw (SciDraw, 2020a; Costa, 2020).

Figure 6. *Avoidance of rat odors.* **A)** Protocol of rat odor avoidance experiment. **B)** Same
as Fig. 5B, but for the rat odor avoidance experiment. **C)** Scatterplot mapping mouse-wise
preference for the clean area before and after the rat was introduced to the "rat odor" area. The
behavior class criteria are shown as dark black dashed lines, while the 2 cohorts of animals used
in the experiments are marked with different symbols. The resulting behavior groups are shown
with three different colors (avoiding the rat odor in blue, which did not occur); and no change in
preference (gold) and preferring the rat odor (green). Six out of 36 mice preferred the area with rat
odor after the rat was introduced. **D)** Overall distribution of positions of mice in the arena,
belonging to the group of mice "preferring" the rat odor, before the rat was introduced. **E)** Same
as D, but for the position distribution after the rat was introduced.

Figure 7. *Avoidance behavior in the presence of a live rat.* **A)** Protocol of rat presence
experiment. **B)** Same as Fig. 5B, but for the rat presence experiment. **C)** Same as Fig. 6B but for
the ratio of avoidance of the area close to the mesh without and with the rat present. All three
possible behavioral groups were observed (avoiding in blue, preferring in green and no change in
yellow). Eighteen out of 36 mice avoided the mesh when the rat was present, while 3 mice
preferred to explore the mesh when the rat was present. **D)** Same as Fig. 6D but applied to mice
that avoided the mesh once the rat was present, showing positions before the rat was present. **E)**
Same as D but showing the overall position distribution while the rat was present.

Figures

    Figure 1. Experimental design, setup, and performance in the presence of a rat.

Figure 2. Movement speed in the presence of rat.

Figure 3. Pupil movement and pupil size in the presence of the rat.

Figure 4. Changes in posture and facial movement in the presence of a rat.

Figure 5. Effect of looming stimuli on the behavior of freely moving mice

Figure 6. Avoidance of rat odors.

Figure 7. Avoidance behavior in the presence of a live rat.