A reversible feedback mechanism regulating mitochondrial heme synthesis

Iva Chitrakar[1], Alexis B. Roberson[1,2], Pedro H. Ayres-Galhardo[1], and Breann L. Brown[1,3*]

[1]Department of Biochemistry, Vanderbilt University School of Medicine, Nashville, Tennessee, USA, [2]Department of Chemistry, Fisk University, Nashville, Tennessee, USA, [3]Center for Structural Biology, Vanderbilt University School of Medicine, Nashville, Tennessee, USA

*For correspondence:

Breann L. Brown, Ph.D.

2215 Garland Ave

Nashville, TN 37232

615-343-1632

breann.brown@vanderbilt.edu

Abstract
Proper heme biosynthesis is essential for numerous cellular functions across nearly all life forms. In humans, dysregulated heme metabolism is linked to multiple blood diseases, neurodegeneration, cardiovascular disease, and metabolic disorders. Erythroid heme production begins with the rate-limiting enzyme Aminolevulinic Acid Synthase (ALAS2) in the mitochondrion. Although prior studies discuss the regulation of ALAS2 in the nucleus and cytoplasm, its modulation as a mature mitochondrial matrix enzyme remains poorly understood. We report that heme binds mature human ALAS2 with high affinity, acting as a reversible mixed inhibitor that reduces enzymatic activity. Structure-based modeling reveals two flexible regions of ALAS2 interact with heme, locking the enzyme in an inactive conformation and occluding the active site. Our work reveals a negative feedback mechanism for heme synthesis, offering insights into the spatial regulation of ALAS2 and the maturation of the essential heme cofactor.

Keywords
heme, enzyme inhibition, erythropoiesis, aminolevulinic acid, pyridoxal 5-phosphate, heme regulatory motif

Introduction
Our knowledge of cellular metabolism continues to expand with our understanding of the role of metabolites as cell signaling effectors (reviewed in [1]). Consequently, metabolic processes, which form the foundation of organismal biology, are studied in a new context to reveal critical insights into chemical and cellular signaling. Heme metabolism is central to multiple biological processes as it is a vital cofactor supporting oxygen transport, transcriptional regulation, and drug detoxification, among others[2-4]. Heme production must be tightly controlled due to the toxicity associated with heme deficiency or excessive amounts of porphyrin precursors, which produce reactive oxygen species and free radicals[5-8].

The enzyme that catalyzes the rate-limiting step for heme biosynthesis in many organisms is aminolevulinic acid synthase (ALAS)[9,10], a PLP-dependent homodimeric enzyme that mediates the condensation of glycine and succinyl-CoA to produce aminolevulinic acid[11,12]. Humans contain two ALAS isoforms, ALAS1 and ALAS2[13-15]. ALAS1 is a ubiquitously expressed housekeeping enzyme [13,14], and ALAS2 is the erythroid-specific isoform that governs the synthesis of 85-90% of the total body heme pool[5,16]. The high demand for heme to produce hemoglobin implies that erythroid heme synthesis is not tightly regulated. However, the fact that multiple blood disorders arise from dysfunctional heme biosynthesis indicates otherwise. Current clinical reports have identified over 95 human *ALAS2* mutations that result in the loss-of-function disease X-linked sideroblastic anemia (XLSA[17] or the gain-of-function disease X-linked protoporphyria (XLPP[18,19]. Compelling new evidence reveals that *ALAS2* is also expressed in non-erythroid cell[20], underscoring our critical need to understand ALAS2 enzyme regulation for metabolic processes beyond erythropoiesis.

Previous research primarily focused on the nuclear and cytosolic regulation of ALAS[21-31]. For example, heme binds directly to the ALAS mitochondrial targeting sequence, preventing protein translocation into the matri[21,22]. However, little is known about its enzymatic modulation in the mitochondrial matrix, which is the cellular compartment where heme biosynthesis is initiated and terminated. Human ALAS paralogs contain multiple putative heme-binding motifs in the mature enzyme, which lacks the mitochondrial targeting sequence. Various reports established that the interaction between heme and mitochondrial ALAS1 results in either LONP1 or CLPXP-mediated degradation[25,32,33]. Further, ALAS homologs from various α-proteobacteria are reported to be inactivated by heme binding[34,35]. Whether human ALAS2 is subject to negative regulation by heme in mitochondria remains to be fully explored. Due to the complex impacts of heme variation on effective erythropoiesis, gene transcription, and

several other cell signaling pathways, it is essential to establish *in vitro* principles of enzyme regulation that form the foundation of interpreting physiological outcomes.

Here, we identified a reversible mechanism by which heme inhibits its synthesis by affecting mature mitochondrial ALAS2 activity, challenging the notion that erythroid heme synthesis is an all-or-nothing process. This mechanism provides a response to heme stress that rapidly inactivates ALAS2, thereby decreasing heme synthesis. With our updated understanding of ALAS2 tissue expression, our study details a new mechanism of negative feedback regulation for heme biosynthesis, regulating erythropoiesis and potentially the maturation of hemoproteins, controlling numerous biological functions.

Results

Human ALAS2 sequence contains multiple heme-binding motifs

One challenge to identifying transient or regulatory heme-protein interactions is the diversity of interaction motifs and amino acid ligands. Although many heme-binding proteins use histidine residues as the axial ligand to coordinate the heme iron, other examples employ cysteine, methionine, tyrosine, or lysine residues[36]. The lower abundance of non-histidine motifs makes predicting these sites less accurate than other canonical motifs[37-39]. However, certain parameters have been identified to aid in prediction.

Previous studies identified a heme regulatory motif (HRM) that consists of a core cysteine-proline (CP) di-peptide in human ALAS1, with cysteine functioning as the axial ligand to the heme iron[21]. A multiple sequence alignment of vertebrate ALAS enzymes revealed the presence of five CP motifs conserved in both isoforms across homologs (**Fig. 1a, Supplemental Fig. 1**). The first two N-terminal HRMs reside in the mitochondrial target sequence, and heme binding to these sites negatively regulates ALAS mitochondrial import[21,22]. The remaining three motifs are dispersed throughout the N-terminal extension and catalytic core of ALAS (**Fig. 1b**).

In addition to the CP motif, a variation of a separate conserved heme-binding motif commonly found in cytochromes was also identified. This canonical CXXCH sequence, where X represents any amino acid, is found in the human ALAS2 C-terminal domain as CXXCR. Interestingly, the mouse homolog contains the classical CXXCH motif in the same position. In contrast, many other vertebrate ALAS enzymes retain the Arg residue in the terminal position (**Fig. 1a**). Collectively, ALAS2 contains two verified and four predicted HRMs per protomer, referred to as HRM1 through HRM6 in this paper.

Human ALAS2 binds heme in vitro

Due to the various motifs present in human ALAS2, we sought to determine if there is a role for heme regulating mature ALAS2 after it is processed in the mitochondrial matrix. Mature wildtype ALAS2 (WT, residues 54-587), which lacks the mitochondrial targeting sequence, was expressed and purified to homogeneity. Titration of WT ALAS2 into hemin (ferric chloride heme) resulted in robust binding exhibited by an exothermic binding curve (**Fig. 1c**). The ALAS2-heme dissociation constant was determined to be ~230 nM (**Table 1**). The binding stoichiometry was best fit to 4:1 (monomer equivalents), corresponding to two heme binding sites per ALAS2 protomer. Thus, heme binds mature ALAS2 tightly, and potentially at multiple sites.

Figure 1. Conserved heme regulatory motif in ALAS. (a) Consensus sequence logo generated based on the heme regulatory motifs (HRM) conserved across multiple ALAS homologs. Conserved motifs (pink box) include CP dipeptide and CXXC. (b) Domain map of human ALAS2 highlighting the experimentally confirmed HRMs in the mitochondrial target sequence (MTS) and predicted HRMs in dark and light blue, respectively. The ALAS N-terminal extension is colored purple and the C-terminal extension is colored orange. The motifs according to human ALAS2 residue numbers are HRM1: [11]CP[12]; HRM2: [38]CP[39]; HRM3: [70]CP[71]; HRM4: [338]CP[339]; HRM5: [471]CP[472]; and HRM6: [555]CXXC[558]. (c) Representative ITC thermograph (top) and binding isotherm (bottom) for the binding between hemin and WT human ALAS2.

Table 1. Isothermal Titration Calorimetry thermodynamic parameters

Construct	K_d (nM)	ΔH (kcal/mol)	ΔS (kcal/mol*K)	ΔG (kcal/mol)
WT	232 [170, 324]	-382 [-440, -342]	-1.25	-9.05
ΔN	589 [322, 1428]	-1006 [-1432, -810]	-3.35	-8.50
ΔC	279 [220, 360]	-544 [-604, -499	-1.80	-8.94
Cys-mut	610 [401, 1132]	-361 [-561, -280]	-1.18	-8.48

*95% confidence interval in brackets

Heme binds reversibly as a mixed inhibitor
Having established that there is robust heme binding to mature ALAS2, we sought to determine the functional impact of this interaction. The *in vitro* enzyme activity of ALAS2 was measured by monitoring the rate of ALA product release. WT ALAS2 activity decreased significantly with increasing hemin concentration, yielding an IC_{50} value of 18.7 ± 0.5 µM (**Fig. 2a, Table 2**). These studies were performed using the catalytically active holoenzyme that is covalently bound to the PLP cofactor. This covalent adduct is broken and reformed throughout the ALAS reaction cycle (**Supplemental Fig. 2**). To address the mechanism of heme inhibition, we generated a catalytically inactive form of ALAS2 by chemically cleaving the covalent bond between the endogenous PLP cofactor and a conserved active site Lys (apoALAS2). The apoenzyme is then reactivated with the addition of fresh cofactor. In the absence of heme, increasing concentrations of PLP restored enzymatic activity, with an activation constant (K_a) of 0.72 ± 0.08 µM (**Fig. 2b**). In the presence of 50 µM heme, PLP partially rescued activity with a 7 ± 1 µM EC_{50} (**Fig. 2b**). It is possible there is not complete restoration of maximal enzyme activity if apoALAS2 contains damaged cofactor that is unable to be exchanged for exogenous PLP. These data reveal that heme-mediated inhibition is reversible. Next, ALAS2 activity was monitored with increasing amounts of

PLP in the presence of various heme concentrations, up to 100-fold excess. The resulting data revealed significant differences in maximal enzyme velocity (V_{max}) and an increased PLP K_a, indicating that heme binds as a mixed inhibitor that binds both the free and cofactor-bound enzyme (**Fig. 2c**).

Figure 2. Heme reversibly inhibits the enzymatic activity of human ALAS2.
(a) The rate of ALA product released by WT ALAS2 measured as a function of increasing hemin concentration. (b). The rate of ALA product produced upon the addition of increasing concentrations of PLP cofactor into apo WT ALAS2 in either the absence (black diamonds) or the presence of 50 μM hemin (blue triangles). (c) Lineweaver Burk plots depicting the change in enzyme velocity in the presence of increasing concentrations of the hemin inhibitor (shades of blue).

Heme binding favors a catalytically inactive ALAS2 conformation
The crystal structure of human ALAS2 lacks the region containing HRM3 and portions of the C-terminal extension due to conformational flexibility and crystallographic disorder[40]. To determine how heme interacts with ALAS2 to inhibit enzyme activity, we used AlphaFold3 (AF3) to model heme bound to the mature ALAS2 homodimer[41] (**Fig. 3a-b, Supplemental Fig. 3**). The lowest-energy model resulting from the input of two mature ALAS chains and two heme b molecules proposes that one of the heme binding sites involves the coordination of two HRMs in the N-terminal and C-terminal extensions (**Fig. 3b-c**). In the absence of inhibitor, these regions are highly flexible, which allows for extensive rearrangement to coordinate heme simultaneously[42,43]. Although the heme-binding site is predicted with lower confidence, all of the models generated by AF3 depict heme coordinated between HRM3 and HRM6 (**Supplemental Fig. 3**). Notably, the manner of heme binding locks the C-terminal extension in a conformation that occludes the active site (**Fig. 3d**). This represents a similar inactive conformation identified in the crystal structure, indicating this orientation is energetically stable[40]. The confidence of the AlphaFold model is not high enough to indicate which of the three possible cysteine residues serve as the axial ligand to coordinate the heme iron center. Also, adding more than two heme molecules in the prediction led to spurious models, indicating the remaining heme binding sites are buried or require a more extensive conformational rearrangement that cannot be accurately modeled.

Heme potentially binds at multiple sites to inhibit ALAS2 activity
Multiple ALAS2 constructs with various HRM mutations were generated to interrogate the accuracy of the AlphaFold model and to identify additional heme regulatory sites. These included a truncation without HRM3 (ΔN, residues 75-587), a truncation without the C-terminal CXXC motif (ΔC, residues 54-547), and a construct where all cysteines in these two motifs were mutated to alanines (**Fig. 4a**, Cys-mut). All four constructs were purified as holoenzymes and confirmed to be catalytically active. To determine whether ALAS2 binds hemin when one or more of the HRMs are compromised, ITC was performed with each variant (**Fig. 4b, Table 1**). Titration of ΔN, ΔC, and Cys-mut into hemin resulted in an exothermic binding curve similar to WT. The WT enzyme exhibited the highest affinity for hemin, while disruption of the HRM at either terminus reduced affinity, with a smaller effect observed in the

absence of HRM6. Mutation of HRMs at both termini resulted in the lowest affinity for hemin. The ΔN construct lacking HRM3 exhibited an enthalpically driven binding mechanism and a decrease in disorder leading to less efficient binding. This implies that HRM3 plays a prominent role in heme binding compared to the other HRMs. However, other heme-binding sites appear to exist, as revealed by the residual heme binding to our HRM variant constructs.

Figure 3: AlphaFold3 model of the ALAS2:heme complex.
Two copies of mature ALAS2 (residues 54-587) were modeled without heme (a) and with two heme b molecules (b). The ALAS protomers are colored grey and light blue, the PLP cofactor is shown as yellow spheres, and the heme molecules are shown as blue spheres. The flexible N-terminal extension is colored purple, and the autoinhibitory C-terminal extension is shown in orange. (c) Magnified view of one heme-binding sites reveals heme coordinated between HRM3 (C70-P71) and the C-terminal HRM6 (residues 555-558) from the same chain. (d) Addition of the succinyl-CoA substrate (green spheres) reveals the heme-bound model adopts an inactive conformation where the C-terminal extension sterically interferes with substrate binding. The PLP cofactor and succinyl-CoA substrate were modeled based on superposition with *Rhodobacter capsulatus* ALAS (PDB 2BWO).

Compared to WT ALAS2, the basal activity of the Cys-mut construct decreased by approximately 45%. In contrast, the truncated constructs, ΔN and ΔC, showed activity increases of 5% and 70%, respectively (**Fig. 4c**). The large increase in basal activity of ALAS2 ΔC is expected as the deletion of the C-terminal extension is known to underlie the gain-of-function disorder XLPP[18]. Despite the changes in maximal enzyme velocity, the addition of heme resulted in significant inhibition for all constructs (**Fig. 4c**). Enzymatic activity for all mutant constructs decreased with increasing heme concentrations, but displayed higher IC_{50} values than WT (**Fig. 4c, Table 2**). Among the HRM variants, ALAS2 ΔN was the least sensitive to heme inhibition, exhibiting an IC_{50} value nearly 2-fold higher than WT. Both ΔC and Cys-mut constructs were also inhibited by heme to a lesser extent than WT, with IC_{50} values approximately 1.6 and 1.3 times higher than WT, respectively. There were also notable differences in residual enzyme activity under saturating heme concentrations. The WT, ΔN, and Cys-mut constructs retained approximately 11–13% of basal activity in the presence of 100-fold excess

heme. In contrast, ALAS2 ΔC retained ~25% of basal activity under the same conditions. These results indicate that perturbation of either the N-terminal or C-terminal heme regulatory motifs significantly reduces heme-mediated inhibition, but either motif is sufficient to facilitate ALAS2-heme interaction.

Figure 4. Heme binds human ALAS2 at multiple sites.
(a) Domain map of mature ALAS2 WT (colored as in Fig. 1) and HRM variants: WT (residues 54-587), ΔN (residues 75-587), ΔC (residues 54-547), and Cys-mut (C70A/C555A/C558A). (b) Representative ITC thermographs (top) and binding isotherms (bottom) for the interaction between hemin and ALAS2 ΔN (purple, left), ΔC (orange, middle) and Cys-mut (green, right). (c) The rate of ALA release was measured as a function of increasing hemin concentrations for WT (gray diamonds), ΔN (purple triangles), ΔC (orange circles), and Cys-mut (green squares) to monitor inhibition of holoenzyme. (d) The rate of ALA release was measured in the presence of either 50 μM hemin (WT, ΔC, Cys-mut) or 25 μM hemin (ΔN) and increasing concentrations of PLP to determine reactivation of apoenzyme.

The reversibility of heme inhibition of the HRM variants was monitored by titrating PLP into the apoenzymes preincubated with 50 µM heme (or 25 µM heme for ∆N). In the presence of excess PLP, the apoenzyme activity was restored by ~50% for WT and ∆C and to over 75% for Cys-mut and ∆N (**Fig. 4d**). Each ALAS2 variant exhibited EC_{50} values comparable to WT (**Table 2**). However, the maximal catalytic activity was restored to varying amounts as both constructs with HRM3 perturbations (∆N and Cys-mut) displayed the lowest recovery. Additionally, measuring activity in the presence of multiple heme concentrations supported heme interacting as an allosteric mixed inhibitor (**Supplemental Fig. 4**). Overall, these findings provide insight into the reversible interaction between heme and ALAS2, emphasizing its role in regulating enzymatic activity through allosteric effects.

Table 2. ALAS2-ligand dose-response values

Construct	IC_{50} (µM)*	EC_{50} (µM) †
WT	18.7 ± 0.5	7 ± 1
∆N	35.8 ± 0.4	4 ± 2
∆C	31.3 ± 0.8	9 ± 1
Cys-mut	24.8 ± 0.5	10 ± 4

*IC_{50} values were determined by titrating 0.1 µM–60 µM hemin into holoALAS2$_{54-587}$
†EC_{50} values were determined by measuring the activity of apoALAS2 variants in the presence of 50 µM hemin (or 25 µM hemin for ∆N) with increasing PLP titration.

Discussion
The heme biosynthetic enzyme ALAS2 serves as a critical node, governing the first and rate-limiting step of this essential metabolic pathway. Key differences in ALAS2, including tissue expression, protein interactions, and disease manifestation, necessitate examining this isoform separately from ALAS1. The interaction between heme and mature human ALAS2 has long been debated but lacked evidence. Here, we provide unprecedented insights into the intricate interaction between heme and the mature human ALAS2 isoform. Our comprehensive investigation reveals a novel and compelling molecular mechanism of action, wherein heme acts as a potent metabolic inhibitor, effectively decreasing the enzymatic activity of ALAS2. Although labile heme concentrations are difficult to measure directly, estimates vary from less than 1 µM[44] to greater than 20 µM in humans under non-pathogenic conditions[45]. This upper estimate is commensurate with the IC_{50} values we report for mitochondrial ALAS2-heme inhibition, connecting our *in vitro* research to a plausible cellular state. Critically, this unique mode of inhibition is specific to the enzyme as it localizes to the mitochondrial matrix, where little is known about how ALAS2 responds to chemical cues, but where it primarily operates.

Remarkably, heme binding to ALAS2 exhibits several distinctive traits, including redundancy, reversibility, and allostery. This study supports a model by which multiple motifs act in concert to form a single heme-binding site. Our *in vitro* reconstitution uncovered a process that would be otherwise difficult to identify in cells due to the pleiotropic effects of heme on transcriptional programs and erythropoiesis. The presence of multiple heme-binding sites within the enzyme likely serves as a fail-safe for feedback, ensuring the responsiveness of ALAS2 to heme even if one of the binding sites is compromised. For example, truncations of the ALAS2 C-terminal extension lead to the gain-of-function disease XLPP[18,19,46], which would affect the enzyme's ability to respond to heme stress if only one allosteric site existed. However, the presence of multiple non-equivalent heme binding sites across the ALAS2 homodimer makes their isolation and identification difficult, as compensatory changes may occur with enzyme mutagenesis. A redundant method for tuning ALAS2 activity in response to cellular heme levels provides a fitness advantage, and future studies aim to elucidate the unique allosteric properties exhibited by different HRMs.

Structural modeling and analysis reveal that heme functions as an allosteric effector, inducing a conformational change in ALAS2 that prevents substrate binding and, consequently, enzymatic activity. A recent report proposed that heme-bound ALAS2 recruits the CLPXP protease via an adaptor protein to trigger degradation[47]. Similarly, heme binding to the ALAS1 isoform also results in degradation[25,32,33]. Furthermore, our study indicates that heme binding may also influence the interactions of ALAS2 with proteins in the proposed heme synthesis metabolon, a complex of heme biosynthetic enzymes and other mitochondrial proteins that enhance metabolic efficiency[48-50]. This potential modulation of the metabolon would serve as an additional layer of regulation, further influencing the overall efficiency and dynamics of heme production.

Intriguingly, this heme-mediated inhibition discovered in our current work can coexist with a slower, irreversible means of heme-triggered ALAS2 degradation, providing the cell with a multifaceted mechanism to fine-tune heme production in response to various physiological demands (**Fig. 5**). Combined with previous studies of ALAS homologs in bacteria or other tissues[35,51-53], our work reveals a conserved role in heme-mediated inhibition that has evolved to meet organism and tissue-specific needs.

The tight control of heme biosynthesis is paramount, as heme plays a central role in a wide range of cellular processes, including oxygen transport, electron transport, and various metabolic systems such as iron homeostasis and lipid metabolis[54-60]. Given the pivotal role of heme in multiple processes, its dysregulation would also impact physiological functions dependent on other cofactors like FAD and cobalamin, exacerbating disease phenotypes such as anemia, neurotoxicity, oxidative stress, and metabolic dysfunction[63-65]. Therefore, by supporting hemoprotein maturation and integrating with key cellular pathways, the intricate regulation of heme production is crucial for maintaining cellular homeostasis in response to environmental and metabolic demands.

Figure 5. Model of heme-mediated negative feedback inhibition of mitochondrial ALAS2. Under basal conditions, the ALAS2 homodimer (gray ovals) binds PLP (yellow sphere) to subsequently produce heme (red spheres). In the case of toxic heme accumulation, heme binds to ALAS2, with potentially multiple outcomes. We discovered a mechanism where heme binding results in a conformational change that excludes substrate binding (purple box). Additionally, heme binding may recruit mitochondrial proteases, such as CLPXP, for enzyme degradation.

Experimental Methods
Protein expression and purification
Human ALAS2 54-587 constructs were cloned into a modified pET28b vector with a ULP1 protease-cleavable N-terminal hexahistidine-SUMO-tag (His$_6$-SUMO). The plasmids were transformed into *Escherichia coli* BL21-Codon Plus (DE3)-RIL cells (Agilent technologies) and grown in LB medium containing 25 µg/mL chloramphenicol and 50 µg/mL kanamycin at 37°C and 250 rpm. Expression was induced at an OD$_{600}$ of 0.6-0.8 with 0.5 mM IPTG for 4 hours at 22°C and 250 rpm. Cultures were harvested by centrifugation and the pelleted cells were stored at −80°C.

Cells were resuspended in lysis buffer containing 25 mM HEPES pH 8.0, 400 mM NaCl, 100 mM KCl, 2 mM MgCl$_2$, 20 mM imidazole, 10% glycerol, 1 mM DTT, and 20 µM PLP with EDTA-free protease inhibitor (Thermo Fisher Scientific). Cells were lysed by high-pressure homogenization and the lysate was cleared by centrifugation (at 4°C for 30 minutes, 30,000g). The clear lysate was incubated with Ni^{2+}-NTA agarose resin (Qiagen), pre-equilibrated in lysis buffer, for 1 hour at 4°C. After washing the resin with lysis buffer, the protein was eluted with lysis buffer containing 250 mM imidazole. The His-SUMO tag was cleaved using ULP1 overnight at 4°C while dialyzing into a buffer containing 25 mM HEPES pH 7.0, 100 mM KCl, 10% glycerol, 1 mM DTT, and 20 µM PLP. Tag-free protein was then passed through Ni^{2+}-NTA resin pre-equilibrated with dialysis buffer to remove any remaining tagged protein. The protein was further purified with gel-filtration chromatography (HiLoad 16/600 Superdex 200 pg) equilibrated with 25 mM HEPES pH 7, 150 mM KCl, 10% glycerol, and 0.5 mM tris(2-carboxyethyl)phosphine. The purified protein was concentrated using a centrifugal filter (Amicon) flash-frozen in liquid nitrogen, and stored at −80°C.

Site-directed Mutagenesis
Site-directed mutagenesis was performed on the mature ALAS2 (residues 54–587) construct to generate truncations and cysteine mutants targeting HRM3 and HRM6 (**Supplemental Table 1**). All constructs were verified with Sanger sequencing and were expressed and purified as described above.

Multiple Sequence Alignment
The amino acid sequences of human (Homo sapiens; P13196, P22557), bovine (Bos taurus; A6QLI6, Q3ZC31) rat (Rattus norvegicus; P13195, Q63147), mouse (Mus musculus; Q8VC19, P08680), and chicken (Gallus gallus: P07997, P18080) ALAS1 and 2 were retrieved from the UniProt database. The multiple sequence alignment was performed using Clustal Omega 1.2.4 with default parameters to identify conserved regions among the sequences. Heme regulatory motif 3 and 6 along with residues flanking the motifs were subsequently used to generate a sequence frequency plot using WebLogo 3.7.4. The alignment of all sequences was visualized using ESPript 3.0.24.

Isothermal Titration Calorimetry
Binding curves of the hemin-protein interaction were determined using NanoAffinity (TA instruments) at 25°C. Hemin was dissolved in 1 M NaOH and autoclaved to prepare a stock solution of 746 µM, ε_{385nm} = 58,440 M^{-1} cm^{-1}. The stock solution was diluted to 1.5 µM in titration buffer containing 25 mM HEPES pH 7, 150 mM KCl, 0.5 mM TCEP. Protein samples were dialyzed into the titration buffer and diluted to a final concentration of 40 µM. Fixed volumes of protein were injected at 200-second intervals into the cell containing hemin. An A+B hetero-association model was used to determine the thermodynamic parameters for each variant. Data were integrated and analyzed using NITPIC and SEDPHAT, and visualized using GUSSI[61-63].

Enzyme activity assay
A discontinuous colorimetric activity assay adapted from Shoolingin-Jordan et al. and Bailey et al. was used to measure the enzyme activity[40,64]. The reaction was initiated by adding 100 nM purified enzyme to 50 mM potassium phosphate pH 7.0, 10 mM MgCl$_2$, 1 mM DTT, 100 mM glycine and 300 µM

succinyl-CoA (175 µL total). The reaction was incubated at 37°C for 15 minutes and terminated with 100 µL of cold 10% trichloroacetic acid. The precipitated protein was removed by centrifugation for 5 minutes, 13,000g and 240 µL of the supernatant was mixed with 240 µL of 1 M sodium acetate pH 4.6. To derivatize ALA, 20 µL of acetylacetone was added to the mixture, and boiled at 100°C for 10 minutes. After the sample was cooled for 20 minutes, 100 µL aliquots were transferred into a clear 96-well plate in triplicate. An equal volume of Ehrlich's reagent (p-Dimethylaminobenzaldehyde, DMAB) was added to each well, and the absorbance was monitored at 553 nm using a microplate reader (Thermo Fisher Scientific). The peak absorbance value was converted to molar quantities of ALA with ε_{553}=of 60,400 M^{-1}cm^{-1}.

Inhibition of enzyme activity was determined by titrating hemin (0.1 µM–60 µM) into the substrate mixture before initiating the reaction upon enzyme addition. Hemin was dissolved in 100% DMSO, ε_{403nm}=170,000 M^{-1} cm^{-1}, and serially diluted in 100% DMSO to prepare stock solutions for each concentration. The final DMSO concentration in the reaction mixture was 2%. All experiments were performed with a minimum of three biological replicates, each with three technical replicates. Data were fitted using non-linear regression to generate an inhibitor dose-response curve (IC$_{50}$) or an agonist dose-response curve (EC$_{50}$), and statistical significance was determined using a one-way ANOVA with GraphPad Prism 10.

For inhibitor competition experiments, the apoenzymes were preincubated with hemin or PLP (0, 10, 25, 50 and 100 uM) for 15 minutes at 25°C in the dark prior to initiating the reaction. To restore the activity of hemin-apoenzyme and to inhibit the activity of PLP-apoenzyme, PLP or hemin (0, 1, 5, 10, 25, 50, 100 uM) were titrated, respectively. All experiments were performed with a minimum of three biological replicates, each with three technical replicates. Data were fit using non-linear regression to generate an agonist dose-response curve (EC$_{50}$) or an inhibitor dose-response curve (IC$_{50}$). Kinetic parameters (V$_{max}$ and K$_M$) were determined using a Michaelis-Menten model in GraphPad Prism 10 and used for generating Lineweaver Burk Plots.

In silico modeling
The mature ALAS2-heme model was generated using AlphaFold3 v3.0.0. Two copies of mature ALAS2 (54-587, Uniprot P22557), and two copies of heme b ligand were entered as the search query. Of the five models generated, the predicted structure with the highest scores was chosen to generate the heme-bound model. The structural figures were generated using ChimeraX.

Supplementary material
The supplemental material contains three figures and one table including entire ALAS enzyme multiple sequence alignment, AlphaFold models and corresponding pLDDT scores, Lineweaver Burk plots for HRM variants and, primer sequences used for site-directed mutagenesis.

Acknowledgements
This work was supported by the National Institute of General Medical Sciences of the National Institutes of Health grant DP2GM146255 (B.L.B.). We thank Lauren P. Jackson for the use of the NanoAffinity ITC instrument. The Vanderbilt Institute of Chemical Biology Synthesis Core provided the succinyl-Coenzyme A compound. Certain figures were created in BioRender.

Conflict of interest statement
The authors declare no conflict of interest.

Author Contributions
Iva Chitrakar: Conceptualization, methodology, investigation, validation, formal analysis, visualization, writing

Alexis B. Roberson: Investigation
Pedro H. Ayres Galhardo: Investigation
Breann L. Brown: Conceptualization, methodology, investigation, validation, formal analysis, visualization, writing, funding acquisition

ORCID
Iva Chitrakar: https://orcid.org/0000-0002-6359-1042
Alexis B. Roberson: https://orcid.org/0009-0001-3057-4350
Breann L. Brown: https://orcid.org/0000-0001-8109-4472

References
1. Baker, S.A. & Rutter, J. Metabolites as signalling molecules. *Nat Rev Mol Cell Biol* **24**, 355-374 (2023).
2. Dioum, E.M. et al. NPAS2: a gas-responsive transcription factor. *Science* **298**, 2385-7 (2002).
3. Freeman, S.L. et al. Heme binding to human CLOCK affects interactions with the E-box. *Proc Natl Acad Sci U S A* **116**, 19911-19916 (2019).
4. Yin, L. et al. Rev-erbalpha, a heme sensor that coordinates metabolic and circadian pathways. *Science* **318**, 1786-9 (2007).
5. Dailey, H.A. & Meissner, P.N. Erythroid heme biosynthesis and its disorders. *Cold Spring Harb Perspect Med* **3**, a011676 (2013).
6. Hamza, I. & Dailey, H.A. One ring to rule them all: trafficking of heme and heme synthesis intermediates in the metazoans. *Biochim Biophys Acta* **1823**, 1617-32 (2012).
7. Bechara, E.J.H., Ramos, L.D. & Stevani, C.V. 5-Aminolevulinic acid: A matter of life and caveats. *Journal of Photochemistry and Photobiology* **7**, 100036 (2021).
8. Chiabrando, D., Vinchi, F., Fiorito, V., Mercurio, S. & Tolosano, E. Heme in pathophysiology: a matter of scavenging, metabolism and trafficking across cell membranes. *Front Pharmacol* **5**, 61 (2014).
9. Hunter, G.A., Zhang, J. & Ferreira, G.C. Transient kinetic studies support refinements to the chemical and kinetic mechanisms of aminolevulinate synthase. *J Biol Chem* **282**, 23025-35 (2007).
10. Fratz, E.J. et al. Human Erythroid 5-Aminolevulinate Synthase Mutations Associated with X-Linked Protoporphyria Disrupt the Conformational Equilibrium and Enhance Product Release. *Biochemistry* **54**, 5617-31 (2015).
11. Gibson, K.D., Laver, W.G. & Neuberger, A. Initial stages in the biosynthesis of porphyrins. 2. The formation of delta-aminolaevulic acid from glycine and succinyl-coenzyme A by particles from chicken erythrocytes. *Biochem J* **70**, 71-81 (1958).
12. Kikuchi, G., Kumar, A., Talmage, P. & Shemin, D. The enzymatic synthesis of delta-aminolevulinic acid. *J Biol Chem* **233**, 1214-9 (1958).
13. Bishop, D.F. Two different genes encode delta-aminolevulinate synthase in humans: nucleotide sequences of cDNAs for the housekeeping and erythroid genes. *Nucleic Acids Res* **18**, 7187-8 (1990).
14. Bishop, D.F., Henderson, A.S. & Astrin, K.H. Human delta-aminolevulinate synthase: assignment of the housekeeping gene to 3p21 and the erythroid-specific gene to the X chromosome. *Genomics* **7**, 207-14 (1990).
15. Riddle, R.D., Yamamoto, M. & Engel, J.D. Expression of delta-aminolevulinate synthase in avian cells: separate genes encode erythroid-specific and nonspecific isozymes. *Proc Natl Acad Sci U S A* **86**, 792-6 (1989).
16. Besur, S., Hou, W., Schmeltzer, P. & Bonkovsky, H.L. Clinically important features of porphyrin and heme metabolism and the porphyrias. *Metabolites* **4**, 977-1006 (2014).

17. Stenson, P.D. et al. The Human Gene Mutation Database (HGMD((R))): optimizing its use in a clinical diagnostic or research setting. *Hum Genet* **139**, 1197-1207 (2020).
18. Whatley, S.D. et al. C-terminal deletions in the ALAS2 gene lead to gain of function and cause X-linked dominant protoporphyria without anemia or iron overload. *Am J Hum Genet* **83**, 408-14 (2008).
19. Balwani, M. Erythropoietic Protoporphyria and X-Linked Protoporphyria: pathophysiology, genetics, clinical manifestations, and management. *Mol Genet Metab* **128**, 298-303 (2019).
20. Lee, S., Lee, S., Desnick, R., Yasuda, M. & Lai, E.C. Noncanonical role of ALAS1 as a heme-independent inhibitor of small RNA-mediated silencing. *Science* **386**, 1427-1434 (2024).
21. Lathrop, J.T. & Timko, M.P. Regulation by heme of mitochondrial protein transport through a conserved amino acid motif. *Science* **259**, 522-5 (1993).
22. Munakata, H. et al. Role of the heme regulatory motif in the heme-mediated inhibition of mitochondrial import of 5-aminolevulinate synthase. *J Biochem* **136**, 233-8 (2004).
23. Srivastava, G. et al. Regulation of 5-aminolevulinate synthase mRNA in different rat tissues. *J Biol Chem* **263**, 5202-9 (1988).
24. Peoc'h, K. et al. Regulation and tissue-specific expression of delta-aminolevulinic acid synthases in non-syndromic sideroblastic anemias and porphyrias. *Mol Genet Metab* **128**, 190-197 (2019).
25. Nomura, K. et al. Heme-dependent recognition of 5-aminolevulinate synthase by the human mitochondrial molecular chaperone ClpX. *FEBS Lett* **595**, 3019-3029 (2021).
26. Tanimura, N. et al. Mechanism governing heme synthesis reveals a GATA factor/heme circuit that controls differentiation. *EMBO Rep* **17**, 249-65 (2016).
27. Cox, T.C., Bawden, M.J., Martin, A. & May, B.K. Human erythroid 5-aminolevulinate synthase: promoter analysis and identification of an iron-responsive element in the mRNA. *EMBO J* **10**, 1891-902 (1991).
28. Liu, J. et al. Long non-coding RNA-dependent mechanism to regulate heme biosynthesis and erythrocyte development. *Nat Commun* **9**, 4386 (2018).
29. Campagna, D.R. et al. X-linked sideroblastic anemia due to ALAS2 intron 1 enhancer element GATA-binding site mutations. *Am J Hematol* **89**, 315-9 (2014).
30. Kaneko, K. et al. Identification of a novel erythroid-specific enhancer for the ALAS2 gene and its loss-of-function mutation which is associated with congenital sideroblastic anemia. *Haematologica* **99**, 252-61 (2014).
31. Surinya, K.H., Cox, T.C. & May, B.K. Identification and characterization of a conserved erythroid-specific enhancer located in intron 8 of the human 5-aminolevulinate synthase 2 gene. *J Biol Chem* **273**, 16798-809 (1998).
32. Tian, Q. et al. Lon peptidase 1 (LONP1)-dependent breakdown of mitochondrial 5-aminolevulinic acid synthase protein by heme in human liver cells. *J Biol Chem* **286**, 26424-30 (2011).
33. Kubota, Y. et al. Novel Mechanisms for Heme-dependent Degradation of ALAS1 Protein as a Component of Negative Feedback Regulation of Heme Biosynthesis. *J Biol Chem* **291**, 20516-29 (2016).
34. Zhang, L. et al. Cloning of two 5-aminolevulinic acid synthase isozymes HemA and HemO from Rhodopseudomonas palustris with favorable characteristics for 5-aminolevulinic acid production. *Biotechnol Lett* **35**, 763-8 (2013).
35. Ikushiro, H. et al. Heme-dependent Inactivation of 5-Aminolevulinate Synthase from Caulobacter crescentus. *Sci Rep* **8**, 14228 (2018).
36. Li, T., Bonkovsky, H.L. & Guo, J.T. Structural analysis of heme proteins: implications for design and prediction. *BMC Struct Biol* **11**, 13 (2011).
37. Humayun, F. et al. A Computational Approach for Mapping Heme Biology in the Context of Hemolytic Disorders. *Front Bioeng Biotechnol* **8**, 74 (2020).

38.   Schubert, E. et al. Spectroscopic studies on peptides and proteins with cysteine-containing heme regulatory motifs (HRM). *J Inorg Biochem* **148**, 49-56 (2015).
39.   Rathod, D.C., Vaidya, S.M., Hopp, M.T., Kuhl, T. & Imhof, D. Shapes and Patterns of Heme-Binding Motifs in Mammalian Heme-Binding Proteins. *Biomolecules* **13**(2023).
40.   Bailey, H.J. et al. Human aminolevulinate synthase structure reveals a eukaryotic-specific autoinhibitory loop regulating substrate binding and product release. *Nat Commun* **11**, 2813 (2020).
41.   Abramson, J. et al. Accurate structure prediction of biomolecular interactions with AlphaFold 3. *Nature* **630**, 493-500 (2024).
42.   Na, I., DeForte, S., Stojanovski, B.M., Ferreira, G.C. & Uversky, V.N. Molecular dynamics analysis of the structural and dynamic properties of the functionally enhanced hepta-variant of mouse 5-aminolevulinate synthase. *J Biomol Struct Dyn* **36**, 152-165 (2018).
43.   Stojanovski, B.M., Hunter, G.A., Jahn, M., Jahn, D. & Ferreira, G.C. Unstable reaction intermediates and hysteresis during the catalytic cycle of 5-aminolevulinate synthase: implications from using pseudo and alternate substrates and a promiscuous enzyme variant. *J Biol Chem* **289**, 22915-22925 (2014).
44.   Sassa, S. Why heme needs to be degraded to iron, biliverdin IXalpha, and carbon monoxide? *Antioxid Redox Signal* **6**, 819-24 (2004).
45.   Aich, A., Freundlich, M. & Vekilov, P.G. The free heme concentration in healthy human erythrocytes. *Blood Cells Mol Dis* **55**, 402-9 (2015).
46.   Kadirvel, S. et al. The carboxyl-terminal region of erythroid-specific 5-aminolevulinate synthase acts as an intrinsic modifier for its catalytic activity and protein stability. *Exp Hematol* **40**, 477-86 e1 (2012).
47.   Cottle, T., Joh, L., Posner, C., DeCosta, A. & Kardon, J.R. An adaptor for feedback regulation of heme biosynthesis by the mitochondrial protease CLPXP. *bioRxiv* (2024).
48.   Medlock, A.E. et al. Identification of the Mitochondrial Heme Metabolism Complex. *PLoS One* **10**, e0135896 (2015).
49.   Grandchamp, B., Phung, N. & Nordmann, Y. The mitochondrial localization of coproporphyrinogen III oxidase. *Biochem J* **176**, 97-102 (1978).
50.   Ferreira, G.C., Andrew, T.L., Karr, S.W. & Dailey, H.A. Organization of the terminal two enzymes of the heme biosynthetic pathway. Orientation of protoporphyrinogen oxidase and evidence for a membrane complex. *J Biol Chem* **263**, 3835-9 (1988).
51.   Tan, Z. et al. Enhancing thermostability and removing hemin inhibition of Rhodopseudomonas palustris 5-aminolevulinic acid synthase by computer-aided rational design. *Biotechnol Lett* **41**, 181-191 (2019).
52.   He, G. et al. Construction of 5-aminolevulinic acid synthase variants by cysteine-targeted mutation to release heme inhibition. *J Biosci Bioeng* **134**, 416-423 (2022).
53.   Scholnick, P.L., Hammaker, L.E. & Marver, H.S. Soluble -aminolevulinic acid synthetase of rat liver. II. Studies related to the mechanism of enzyme action and hemin inhibition. *J Biol Chem* **247**, 4132-7 (1972).
54.   Dohi, Y. et al. Bach1 inhibits oxidative stress-induced cellular senescence by impeding p53 function on chromatin. *Nat Struct Mol Biol* **15**, 1246-54 (2008).
55.   Igarashi, K. & Sun, J. The heme-Bach1 pathway in the regulation of oxidative stress response and erythroid differentiation. *Antioxid Redox Signal* **8**, 107-18 (2006).
56.   Zenke-Kawasaki, Y. et al. Heme induces ubiquitination and degradation of the transcription factor Bach1. *Mol Cell Biol* **27**, 6962-71 (2007).
57.   Keel, S.B. et al. A heme export protein is required for red blood cell differentiation and iron homeostasis. *Science* **319**, 825-8 (2008).
58.   Miller, M.A. et al. CO recombination in cytochrome c peroxidase: effect of the local heme environment on CO binding explored through site-directed mutagenesis. *Biochemistry* **29**, 1777-91 (1990).

59.   Vinchi, F. et al. Heme exporter FLVCR1a regulates heme synthesis and degradation and controls activity of cytochromes P450. *Gastroenterology* **146**, 1325-38 (2014).
60.   Zhuang, J. et al. Evaluating the roles of the heme a side chains in cytochrome c oxidase using designed heme proteins. *Biochemistry* **45**, 12530-8 (2006).
61.   Keller, S. et al. High-precision isothermal titration calorimetry with automated peak-shape analysis. *Anal Chem* **84**, 5066-73 (2012).
62.   Zhao, H., Piszczek, G. & Schuck, P. SEDPHAT--a platform for global ITC analysis and global multi-method analysis of molecular interactions. *Methods* **76**, 137-148 (2015).
63.   Brautigam, C.A. Calculations and Publication-Quality Illustrations for Analytical Ultracentrifugation Data. *Methods Enzymol* **562**, 109-33 (2015).
64.   Shoolingin-Jordan, P.M., LeLean, J.E. & Lloyd, A.J. Continuous coupled assay for 5-aminolevulinate synthase. *Methods Enzymol* **281**, 309-16 (1997).