1
2
3
4
5
6 **Species-specific oculomotor control tolerances predict saccadic suppression**
7 **strength in macaques and zebrafish**
8
9
10

11 Giulia Soto[1,#], Florian A. Dehmelt[1,#], Matthias P. Baumann[1,2,#], Ibrahim Tunç[1,2,3,4], Yue
12 Yu[1,2], Tatiana Malevich[1,2], Ziad M. Hafed[1,2*], Aristides B. Arrenberg[1,*]
13
14
15
16
[1] Werner Reichardt Centre for Integrative Neuroscience, University of Tübingen,
Tübingen, Germany
[2] Hertie Institute for Clinical Brain Research, University of Tübingen, Tübingen,
Germany
[3] Graduate Training Centre of Neuroscience, University of Tübingen, Tübingen,
Germany
[4] Current address: Computational Systems Neuroscience, Institute of Zoology,
University of Cologne, Cologne, Germany
* corresponding authors: aristides.arrenberg@uni-tuebingen.de, and
ziad.m.hafed@cin.uni-tuebingen.de
[#] contributed equally
28

Summary

Active sensing necessarily requires integrating information about both self-generated

movements as well as past, present, and future afferent inputs[1,2]. However, such

information is inherently variable and relies, at least in part, on uncertain extrapolations.

Here, using saccadic suppression[3,4] of visual sensitivity as a classic example of sensory-

motor integration, we show that suppression strength in two drastically different species,

macaque monkeys and zebrafish larvae, may be a direct outcome of efficient state

estimation in the presence of uncontrollable sensory and motor variability. Bayesian

estimator models[5] suggest that optimal saccadic suppression should rely not just on the

sensory-motor information being processed, but also on the time-dependent magnitude

of unexplained variability in the nervous system encoding it. In both macaques and

zebrafish larvae, and using matched visual stimulation regimes across the species, we

experimentally measured saccadic suppression strength in the superior colliculus (SC) of

the monkeys and the homologous optic tectum (OT) of the fish. We also experimentally

quantified additive and multiplicative noise components in the motor systems of both

species, and we furthermore estimated noise in the sensory systems. We found that inter-

species differences in noise levels and their theoretically predicted impacts on visual

sensitivity are qualitatively consistent with our experimentally observed differences in

saccadic suppression strength between the fish and the monkeys. Because sensory and

motor noise levels can reflect the amounts of available neural resources committed to a

given task, our results strongly underscore the value of incorporating computational

resource limits in investigating performance differences that have evolved in homologous

brain areas.

Results and Discussion

When moving through the world, our brains constantly reconcile potentially conflicting information arising from sensory inputs in a changing environment and from the internal signals of our own body movements. One striking example of this non-stationary relationship between sensory and motor information is the active vision phenomenon of saccadic suppression: a brief reduction in visual sensitivity around the time of saccadic eye movements[3,4,6–10] (Figure 1A).

Saccadic suppression is a highly conserved phenomenon across a wide range of species, from zebrafish larvae[11] to rodents[9,10] and primates/humans[3,4,7,9,12,13]. However, the neural resources in these different species are very drastically different (e.g. Figures 1B-C), providing us with an ideal opportunity to understand how computational limits in different brain sizes act to quantitatively reshape what are otherwise qualitatively identical phenomena in multiple species.

Starting from a theoretical foundation, we implemented a computational model allowing parameterization of different sensory and motor noise contributions to saccadic suppression strength. Then, using matched visual stimulation paradigms (Methods), we experimentally measured saccadic suppression strength in homologous brain structures within macaque monkeys and zebrafish larvae, two species at opposite ends of the evolutionary spectrum. Our results revealed a particularly important effect of sensory noise in dictating saccadic suppression strength in the two species.

Saccadic suppression as efficient sensory-motor estimation

Eye position control can be optimized in a closed-loop system, and a recent study suggested that saccadic suppression might be caused by efficient sensory-motor estimation across rapid eye movements[5]. According to this framework (Figure 1D), a Bayesian estimator dynamically balances predictions from both sensory and motor information according to their ever-changing reliability, particularly across saccades. Uncertainty arises explicitly from four distinct sources: the sensory input (additive sensory

noise, σ_t), the dynamics of the eye plant (additive motor noise, ξ_t), the control signal itself

(multiplicative motor noise, ε_t), and the relay delay of sensory signals, δt, from the retina

to higher computing areas. Taking these variable sources of noise into account,

specifically around the time of rapid eye movements, sensory signals are very unreliable

in estimating the eye plant's instantaneous state. Thus, to optimize state estimation during

saccades, sensory signals should be discounted and motor signals preferred.

Figure 1. Saccadic suppression as efficient sensory-motor estimation.

(A) In a suppression window around the time of a saccadic eye movement, visual perception is reduced.

(B) Horizontal eye position during a saccadic eye movement of macaques (orange trace) and zebrafish (blue trace). The zebrafish saccade is executed slower and less accurately than the macaque saccade.

(C) Schematic linking neuronal resources to accuracy of sensory estimation in macaque and zebrafish. Macaque superior colliculus (SC) contains ~100 times as many neurons as the homologous zebrafish optic tectum (OT). This allows macaques to sample their significantly smaller visual field with many small-size receptive fields. Whereas zebrafish have to fill a large visual field (here centered around the area centralis) with fewer larger receptive fields. Thus, macaques can estimate the position of a visual stimulus (red dot) more precisely than zebrafish.

(D) Schematic illustration of Bayesian state estimation for closed-loop control of eye movements (adapted from Crevecoeur and Kording[5]). To estimate the current position x of a visual target, its previous position is relayed as visual input y with delay δt and visual noise σ_t, then combined with motor activity from the corollary discharge. This extrapolation with sensory information is weighted against a purely feedforward motor extrapolation via Kálmán filter K. Eye position is moved towards this estimated target by motor control signal u with multiplicative noise ε_t, additive noise ξ_t, and a scaling factor α. A_t and B_t capture the dynamics of the eye plant, t is time, dt the time step. Changes in K correspond to saccadic suppression or enhancement. See Methods for the model and notations.).

The above mechanistic explanation of saccadic suppression relies on anatomical structures that combine sensory and motor signals. In mammals, this is found in the superior colliculus (SC), and in lower vertebrates such as zebrafish, in the optic tectum (OT)[14–18] (Figure 1C). Studies in the macaque SC and zebrafish OT have shown that saccadic suppression can indeed be observed in neurons within these structures[7,11,16,19,20], despite their large evolutionary distance from one another. Having said that, while the SC and OT are anatomically homologous, they vary vastly in size, and therefore in their ability to contribute neural resources to the optimization task required by saccadic suppression. For example, we estimated, based on brain volume and the total number of neurons in each organism, that the macaque SC has ~100 times as many neurons as the zebrafish OT (Figure 1C, Methods). Concomitantly, observations of eye position during saccadic eye movements in zebrafish and macaques (Figure 1B) revealed less accurate saccadic eye movement execution in zebrafish, indicating less precise motor control. Further, when comparing sensory accuracy (Figure 1C), we noted that zebrafish have a much larger visual field compared to macaques, and with fewer neurons and larger receptive fields[21–23], this visual field is much more coarsely sampled in

123   zebrafish. In summary, the zebrafish oculomotor system may have to rely on much more

124   uncertain visual and motor information than that of macaques, particularly peri-

125   saccadically. As we show below, this has direct implications on the strength of saccadic

126   suppression in the two species.

128 Zebrafish motor signals are much noisier than those of macaques during saccadic eye
129 movements

To assess the effect of noise levels on theoretical predictions of efficient sensory

estimation, we experimentally quantified motor noise before, during, and after saccades

in macaque monkeys and zebrafish (Figure 2). We recorded saccade traces (see

Methods, "Noise estimation") in both species (Figures 2A and C) and estimated noise

parameters for use in the model (Figures 2B, 2D). First, eye traces were split into pre-

saccadic, saccadic, and post-saccadic sections (Figures 2B, 2D). Pre- and post-saccadic

sections were used to determine additive motor noise, ξ, since it is modeled as

independent of the motor control signal for saccades (see Methods, "Additive motor

noise"). This represents the instability of eye position in the absence of a motor control

signal. To quantify multiplicative motor noise, ε, we fit a sigmoidal curve to the saccadic

section of the eye traces (Figures 2B and D). From the time derivative of the sigmoidal fit,

we derived the control signal u(t). A piecewise-constant approximation of u(t), combined

with the previously determined additive motor noise, ξ, was then used to estimate ε (see

Methods, "Multiplicative motor noise"). This represents the variability of the saccadic

motor control signal and the resulting change of eye position. We thus determined that

zebrafish motor control is much noisier than that of macaque monkeys, with the additive

motor noise being about 10 times higher in zebrafish, and the multiplicative motor noise

being nearly 40 times higher (Figure 2E). In our subsequent simulations, both types of

motor noise were coupled, and always covaried.

Figure 2. Zebrafish motor signals are much noisier than those of macaques during saccadic eye movements.

(A) Normalized eye position during saccadic eye movements in macaques. Grey lines indicate individual trials, colored line indicates average eye position and shaded envelope indicates standard deviation.

(B) Example macaque saccade with sigmoidal fit (dashed line). Deviations from the fit (RMS) are considered motor noise (for detailed description see STAR Methods).

(C and D) Same as (A) and (B) for zebrafish.

(E-F) Standard deviations of the estimated multiplicative (ϵ) motor noise, representing the saccadic motor control signal, and the estimated additive (ξ) motor noise, representing pre- and post-saccadic variability of eye positions. Boxplot shows standard deviations of multiplicative motor noise estimates for multiple velocity bins for macaque and zebrafish during saccadic eye movement (see STAR Methods). Central line indicates median, box encloses 25th to 75th percentile and whiskers extend to minimum and maximum of data; circles show outliers. (F) Violin plot shows distribution of additive motor noise estimates pooled across all pre- and post-saccadic stationary phases for macaque and zebrafish; error bars indicate the standard deviation.

On the assumption that the precision of sensory information increases with the number of available neurons for tiling visual space, the sensory noise parameter σ for both species was estimated from the number of neurons in their respective visual brain areas (see Methods, "Estimation of neuron count", "Sensory noise"). With about 1 million neurons in the SC and 10,000 neurons in OT, we expected the sensory noise to be 10 times larger

194   in zebrafish compared to macaques. We were now in a position to generate predictions
195   on saccadic suppression strength across the two species.

Suppression strength differences across species are consistent with their noise levels

We first identified model parameters to reproduce the curve predicted in the original
study[5]. Then, relative to this baseline, we ran further simulations in which we increased
the respective noise parameters by factors of up to 37.5 times (Figure 3A). This range
included the ratios observed between macaque and zebrafish noise parameters. Higher
sensory noise indeed reduced the predicted optimum of saccadic suppression strength,
consistent with the notion that if sensory information is always unreliable, then saccades
have less effect on the trade-off involved in efficient state estimation (Figure 3B). Because
absolute contributions from different noise sources vary over time, the effect of motor
noise was not symmetric to that of sensory noise: Multiplicative motor noise matters only
during saccade execution, accumulates over its course and may require a post-saccadic
correction; but if high levels of additive motor noise make the eye position unstable, the
(inevitably delayed) sensory information may be unable to provide such a correction.
When set to levels observed in zebrafish, the two coupled types of motor noise noticeably
affected the time course of post-saccadic corrections, predicting post-saccadic
enhancement at lower noise levels only (Figure 3C; see below and Fig. 4 for what we
mean by post-saccadic enhancement). In summary, if zebrafish have higher sensory and
motor noise than macaque monkeys, model results predict much stronger saccadic
suppression (and post-saccadic enhancement) in monkey than in fish.

To test for such cross-species differences in saccadic suppression strength, we recorded
neural activity in the SC of macaque monkeys and the OT of zebrafish. The visual
stimulation regime for the fish was described recently[11], and it involved saccades across
textured backgrounds; here we picked one of the background textures tested in the earlier
study (low spatial frequency), and we directly compared it with observations from the
macaque monkey data. In such data, the monkeys generated horizontal saccades

(rightward or leftward across a coarse texture (Methods). The texture was generated

similarly to the fish experiments, with the only difference being that the spatial frequency

ranges were matched to the receptive field sizes expected in the monkeys (Methods); this

is a similar approach to our earlier multi-species comparisons of saccadic suppression[9,10].

Figure 3. Bayesian model predicts differences in saccadic suppression across the range of noise parameters observed in macaque and zebrafish

(A) Optimal time course of saccadic suppression predicted by the model. All solid black traces show the first element of the optimal Kálmán matrix, vertical dashed lines show saccade onset. Different combinations of sensory and motor noise require different saccadic suppression. The thick trace on the upper left was obtained for macaque-like noise parameters, and the thick trace on the bottom right for zebrafish-like values. All noise parameters shown here (e.g., "10x") are relative to macaque values ("1x").

(B) Increased sensory noise reduces the depth of saccadic suppression across motor noise regimes, but in the parameter range investigated, hardly affects its overall shape. Curves are superpositions of those in the columns of (A), with shades of green indicating different magnitudes of sensory noise for the same motor noise.

(C) Increased motor noise discourages post-saccadic enhancement, and this effect is robust across
sensory noise regimes. Curves are superpositions of those in the rows of (A), with shades of magenta
indicating different magnitudes of motor noise for the same sensory noise.

Figures 4A and B schematically illustrate the timing of visual stimulus presentation relative

to saccade onset in both species. Example neurons from the SC and OT are shown in

Figures 4C and D, respectively. In both cases, responses to the flash were markedly

reduced when the stimulus appeared shortly after a saccade, indicating clear saccadic

suppression at the level of single units in both primates and fish. To compare suppression

strength across the population, we computed cumulative distribution functions (CDFs) of

suppression indices (Methods) for all recorded neurons (Figure 4E). The distributions

revealed a clear species difference: saccadic suppression in the macaque SC was

generally significantly stronger (statistical results are detailed in the legend of Fig. 4). and

more prevalent than in the zebrafish OT, consistent with the model prediction (Fig. 3) that

higher sensory noise in zebrafish reduces the benefit of discounting sensory input during

saccades. Thus, while saccadic suppression is qualitatively similar in zebrafish[11] and

macaque monkeys[7,16], the quantitative differences between the species can be easily

reconciled by an optimization principle related to efficient state estimation in the presence

of noise.

Finally, in both species, we experimentally tested visual sensitivity at multiple times

relative to saccades (Methods). This allowed us to further compare the empirical data to

the predictions of our model (Figures 4F and 4G). Specifically, for both species, the model

reproduced the fact that there is always an extended time period in which visual sensitivity

is reduced around saccades, regardless of suppression strength. Importantly, besides

correctly predicting that saccadic suppression would be stronger in monkeys than in fish

– driven solely by species-specific noise levels estimated independently (cf. Figure 2), the

model also predicted that post-saccadic enhancement (Fig. 4F and data from Chen et al.[16]) is more likely in monkeys than in fish.

Figure 4. Saccadic suppression in the monkey superior colliculus and the zebrafish optic tectum.

(A) Example eye trace from a monkey with the timing of the baseline flash and a test flash ~50 ms after saccade onset, the saccade is highlighted in grey.

(B) Example eye movement trace from a zebrafish, illustrating baseline and test flash timing. Note: only a test flash 20 ms after saccade onset is shown here; in panels (D) and (E), test flashes at 20, 100, and 250 ms were pooled.

(C) Example neuron from monkey SC: visual response (normalized firing rate) to the baseline flash (black) and to the test flash (orange).

(D) Example neuron from zebrafish tectum: visual response (normalized calcium signal) to baseline flash (black) and pooled test flashes (blue).

(E) Cumulative distribution functions of suppression strength across neurons. Example neurons from (C)
and (D) are highlighted. 83% of SC neurons (orange) show suppression at 50 ms post-saccade; 68% of
tectal neurons (blue) show suppression across the three tested post-saccadic times.

(F) Time course of suppression in monkey SC. Faint lines indicate data from Bellet et al. 2017; crosses
show the current data from all neurons (dashed line) and the significantly suppressed ones (solid line),
closely matching model predictions. Inset: model prediction for low sensory and motor noise, as in Figure
3A, top left.

(G) Suppression time course for zebrafish tectum neurons, analogous to (F). Inset: model prediction for
high sensory and motor noise, as in Figure 3A, bottom right.

In our view, the alignment between modeling and data in our study suggests that efficient

sensory-motor estimation under species-specific noise constraints provides a unifying

account of saccadic suppression strength across vertebrates. We suggest that theoretical

considerations of the implications of computational resource limits, like we did here, will

also be valuable for our understanding of a wide range of further multi-modal, integrative

brain processes.

Methods

Animal Care

All non-human primate and zebrafish experiments were evaluated by ethics committees
and approved by the Tübingen regional government. For macaques we recorded SC
activity from two adult, male rhesus macaque monkeys (A and F), aged 13 to 14 y and
weighing 9 to 13 kg. For zebrafish we recorded calcium activity larval zebrafish 5-7 days
post fertilization (dpf) with pan-neuronal GCaMP6f expression and homozygous for the
mitfa mutation (*Tg(HuC:H2B-GCaMP6f)jf7; nacre -/-*)[24]. For eye position recordings we
used 5-6 dpf larvae *nacre -/-*.   Larvae were raised in E3 solution at 28°C on a 14/10
light/dark cycle.

Noise Estimation

Macaque eye position recordings

The eye data used for the noise estimation is a subset of the data published by Soto et
al.[25]. It comes from the delayed saccade task, in which the monkey fixated a central spot
while a peripheral target (1-degree-diameter circle) appeared and remained visible. After
a variable delay (500–1,000 ms), the fixation spot disappeared, cueing a saccade to the
target. We used 1,445 trials involving four oblique saccade directions with varying
amplitudes. Eye data acquisition and saccade detection were performed similarly to the
procedures described below for the current monkeys' task.

Zebrafish eye position recordings

Eye position of larval zebrafish was recorded at 1 kHz with a high-speed camera (iN8-S1
IDT, Integrated Design Tools, Inc., Pasadena, CA, USA) mounted on a stereomicroscope.
Spontaneous saccades were detected using a motion trigger and 1000 frames recorded,
including 200 frames pre-trigger. Recordings were manually screened for false triggers
caused by non-saccadic motion within the frame and, if necessary, discarded. Eye
position was extracted post-hoc using the ZebEyeTrack virtual machine[26].

Additive motor noise

Before analysis, zebrafish trials were discarded which did not meet the following criteria: both eyes move along the same direction, saccade amplitude is greater than 5° for both eyes, and traces show no strong fluctuations for either eye. The remaining traces were visually inspected, and 5 further traces clearly showing no saccade were manually discarded. In each trial, saccade onset and offset are detected with a velocity criterion matching the variability of the eye trace[27], and trials split into pre-saccadic, saccadic, and post-saccadic periods accordingly. Additive motor noise is estimated from pre- and post-saccadic data only by fitting one affine function each to the pre- and post-saccadic traces of each trial. Deviations between best fits and raw data are interpreted as noise. We compute them for each time point, pool across trials, then compute their standard deviation to estimate ξ.

Multiplicative motor noise

Multiplicative motor noise is associated with the control signal, and thus estimated from saccadic periods under the assumption that the (previously estimated) additive motor noise remains stationary. We fit a sigmoid function to the saccadic part of each eye trace using a soft ramp[28]. As larvae occasionally show pre-saccadic drift as well as large saccadic overshoots or undershoots, and these are not the object of the present study, such trials are excluded based on fit score. The exclusion threshold is high enough to ensure that both high-noise and low-noise trials with sigmoidal saccades are included, and only significantly distorted saccades excluded. For each trial, we then computed the time derivative of the best fit to estimate control signal u, which is pooled over all eye traces. To facilitate estimation of ε, we convert u into a piecewise-constant function by dividing saccades into time bins of 100 values, computing an average $\bar{u}$ per bin, and discretizing it. Across trials, eye positions with the same discretized $\bar{u}$ are pooled; the standard deviation of their distance from best fits provides one estimate of multiplicative noise per value of $\bar{u}$. As we observed no characteristic differences between estimates obtained under high or low control signals, we averaged them to obtain ε. The arbitrary

347   scaling parameter α was kept at 0.06 throughout, as this allowed us to reproduce the
348   original model results[5].

Estimation of neuron count within brain areas

Our estimation of sensory noise depended on the difference in number of neurons in
primate superior colliculus versus zebrafish optic tectum. The superior colliculus of
macaques has an estimated volume of 15 mm^3 (cf. Figure 3D from Chen et al. [29], we
used the raw data of the SC surface reconstruction). The average macaque brain has a
volume of 90.000 - 100.000 mm^3 and an estimated number of 6 x 10^9 neurons[30,31].
Assuming the neuronal density to be roughly homogeneous, we multiplied the total
neuron number by the fraction of SC volume relative to whole brain volume, roughly
totaling 10^6 neurons. Note, however, that the homogeneous density assumption is a
simplification, since subcortical structures have been reported to have lower neuron
density than cortical structures[31]. 7-day-old zebrafish contain about 100,000 neurons[32]
and due to the large volume of the OT, we estimate that it contains about 10,000 neurons.
Based on our estimated neuron count and assuming, for the sake of argument, a uniform
two-dimensional tiling of visual space, the 100-fold difference in neuron count translates
into a difference of visual resolution on the order of σ = $\sqrt{100}$ = 10.

Sensory Estimation Model

To model the effect of noise on efficient sensory estimation, we reconstructed a closed-
loop control model that had previously been used to explain the existence and time course
of saccadic suppression[5]. This abstract model treats sensory and motor information as
simple 4D variables (position and velocity of eye and target, respectively), without any
reference to specific neuronal implementations. In addition to our own, the original
authors kindly provided unpublished MATLAB code. We manually identified parameters
to reproduce the originally published time course and to improve the stability of
simulations across a wider range of noise levels. Final parameters (all unitless as per the
model) included $\xi \in$ [0.01, 0.375], σ $\in$ [10^{-6}, 10^{-5}], scaling factor α = 0.06, sensory delay
= 0.1, initial fixation time = 0.5, movement time = 0.05, stabilization time = 0.25, cost
function tolerance = 0.01, initial target position = -10, final target position = 20, discrete

time step = 0.02. Because inter-species differences were on a comparable order of
magnitude in both ξ (37.5x) and ε (10x), and because the effect of multiplicative motor
noise ε was relatively minor, we covaried the two types of motor noise in all simulations
reported here, with $\varepsilon = \xi$. Because of the model's high degree of abstraction, we cannot
directly substitute our experimentally determined noise values. Therefore, we chose an
approach, where we first reproduced the curve from the original modeling study[5] and then
multiplied the used noise levels to investigate the effects on the saccadic suppression
curve (Figure 3).

Macaque Electrophysiology

Experimental setup

The experimental setup closely followed the procedures outlined in our previous study[33].
In brief, experiments were conducted in a darkened room with the animal positioned
approximately 72 cm away from a calibrated and gamma-linearized CRT monitor
(covering ~30° horizontally and ~23° vertically). Stimulus presentation and data collection
were managed using a custom-modified version of PLDAPS[34], integrated with the
Psychophysics Toolbox[35] and interfaced with an OmniPlex data acquisition system
(Plexon, Inc.).

Animal preparation for behavioral training and electrophysiological recording was carried
out as described in earlier work[33]. Briefly, each animal was surgically implanted with a
head post for head stabilization and a scleral search coil in one eye[36] to enable high-
precision eye tracking via electromagnetic induction[37]. A recording chamber was also
implanted along the midline, tilted 38° posterior to vertical to target the superior colliculus.
We used linear multi-electrode arrays (V-Probes, Plexon Inc.) with either 20 or 24
channels and 50 μm inter-electrode spacing to record neuronal activity. We minimized
sampling bias by offline spike sorting.

Behavioral task

In each trial, monkeys performed a visually guided saccade task against a background of
either a fine or a coarse texture. For each texture condition, one of 80 pre-generated
images (80 fine, 80 coarse) was randomly selected and displayed as the background.

Each trial began with the presentation of a fixation spot, located 7.56° to the left or right
of the screen center. After the monkey acquired and maintained fixation on this spot, it
disappeared and simultaneously reappeared at the center of the screen. The monkey
was trained to perform a visually guided saccade to the new (central) fixation location and
to maintain gaze there for a set duration (500 ms), after which a juice reward was
delivered.

During the saccade, when the eye position reached within 2.5° of the new fixation
location, a probe flash was triggered. This probe, a square of size 1° × 1°, consisted of a
brief luminance change (either an increase or decrease) in the pixel values of the texture
at the receptive field (RF) location of the recorded superior colliculus (SC) neurons. The
RFs had been previously mapped using a separate task.

The flash could occur at one of three randomly chosen onset times relative to the eye
crossing the 2.5° threshold: either immediately (0 ms), or with a delay of 100 ms or 200
ms. Shortly before the initial saccade onset in each trial, we calculated the fixation error
and adjusted the probe position accordingly to ensure accurate alignment with the
intended RF location.

Data Analysis

Eye position data were recorded at 1 kHz using scleral search coils, and neuronal activity
in the superior colliculus (SC) was acquired via laminar probes. Saccades were detected
offline using a previously described analysis toolbox[38], and the detected events were
manually verified to ensure accuracy.

Spike sorting was performed offline using the Kilosort Toolbox[39], followed by manual
refinement with the phy software. Subsequent analyses focused on firing rates across

experimental conditions. Firing rates were estimated by convolving spike trains with a

Gaussian kernel (σ = 10 ms).

Only SC neurons that exhibited a significant visual response relative to baseline activity

(*t*-test) were included in the analysis. To quantify suppression strength, we employed an

analysis approach similar to that used in prior work[9].

First, to isolate the neural response associated with the saccade itself, we aligned spiking

activity to the offline-detected saccade onset and averaged responses across the portion

of the trial influenced by the saccade but not by the visual probe. This saccade-related

response was computed separately for each neuron, saccade direction, and background

texture condition.

Next, we aligned the neural activity to the onset of the probe flash and determined the

temporal offset between saccade onset and probe onset for each trial. The saccade-

related response was then temporally shifted by this offset and subtracted from the

individual trial activity, isolating the probe-evoked component of the response.

In this study, we averaged across trials in which the probe flash was either an increase

or a decrease in local luminance. Analysis was restricted to trials in which a coarse texture

served as the background.

To quantify the degree of suppression, we computed a suppression index for each neuron

by dividing the average evoked response in the 0 ms delay condition by the average

response in the 200 ms delay condition (baseline). The evoked response was defined as

the mean firing rate in the window from 50 to 200 ms after probe onset.

Zebrafish *in-vivo* calcium imaging

Experimental setup

Larval zebrafish were embedded in 1.6% low-melting agarose with the agarose removed

around the eyes to allow for free eye movements. The fish were then placed onto a

custom-built stage and submerged in E3 within a spherical stimulation arena, previously

described[11]. Visual stimuli were projected onto the surface of the stimulation arena at 60

Hz refresh rate at 625 nm.

To measure eye position, the fish were illuminated using a custom-built IR-LED ring and

the eyes were recorded with an infrared-filtered camera (Basler ace 2 R, a2A1920-

160umBAS, Basler AG, Ahrensburg, Germany) at 100 Hz. Fish were allowed to perform

spontaneous saccades. Eye position was extracted online and instantaneous velocity

computed. Saccades were detected, using a velocity threshold of at least 200°/s. To

optimize saccade detection, the actual velocity threshold was adjusted for each fish to

ensure ideal saccade detection and to avoid noise interference. To generate and display

visual stimuli, to extract eye position and velocity and to trigger visual stimuli in a closed-

loop manner we used *VxPy* (Version 0.1.4).

To measure calcium activity in tectal neurons we used a moveable objective two-photon

setup (Sutter Instruments, Novato, CA, USA) with an acquisition rate of 2.18 fps, an image

size of 512 x 512 pixels and a resolution of 0.88 μm x 0.88 μm per pixel. GCaMP6f was

excited at 920 nm and emitted light was collected using a 20x/1.0 Zeiss water immersion

objective.

Visual stimuli

Binary gaussian blur textures were displayed onto the spherical stimulation arena, as

previously described[11]. In response to spontaneous saccades after a delay period of 20,

100, 250, 500, 1000, 2000, or 4000 ms a probe flash consisting of a whole-field luminance

decrease with a cosine shape, lasting 500 ms at 0.5 Hz and an amplitude of 50%

luminance change was displayed. To obtain a baseline response to the probe flash, it

was also shown 8 s after saccade detection to ensure no temporal connection between

saccade and probe flash.

Data Analysis

Calcium recordings were registered and segmented into ROIs using suite2p[40]. Further

analysis of registered ROIs was performed with custom-written Python (PyCharm,

JetBrains, Prague, Czech Republic) and Matlab (The MathWorks Inc., Natick, USA) code.

Fluorescence traces of individual ROIs were normalized to $\Delta F/F_0$, where F_0 was a 120

second sliding median window. ROIs that showed responses to the probe flash were

identified using regression-based analysis and only those ROIs were considered for

further analysis. For each ROI, saccade related responses were quantified, using control

conditions, where no probe flash was shown. Said saccade-related response was

subtracted in all other conditions during and shortly after the time of the saccade. Peak

$\Delta F/F_0$ responses to probe flashes were quantified and averaged for each delay condition.

For each ROI a suppression index for each delay was computed as in the macaque data.

Amongst all probe flash-responsive ROIs, truly suppressed ROIs were identified with a

permutation test, described previously[11].

References

495   1.   Wurtz, R. H. Neuronal mechanisms of visual stability. *Vision Res* **48**, 2070–2089 (2008).

496   2.   Sommer, M. A. & Wurtz, R. H. Brain circuits for the internal monitoring of movements.

497   *Annu Rev Neurosci* **31**, 317 (2008).

498   3.   Zuber, B. L. & Stark, L. Saccadic suppression: Elevation of visual threshold associated

499   with saccadic eye movements. *Exp Neurol* **16**, 65–79 (1966).

500   4.   Beeler, G. W. Visual threshold changes resulting from spontaneous saccadic eye

501   movements. *Vision Res* **7**, 769–775 (1967).

502   5.   Crevecoeur, F. & Kording, K. P. Saccadic suppression as a perceptual consequence of

503   efficient sensorimotor estimation. *Elife* **6**, (2017).

504   6.   Bremmer, F., Kubischik, M., Hoffmann, K. P. & Krekelberg, B. Neural dynamics of

505   saccadic suppression. *Journal of Neuroscience* **29**, 12374–12383 (2009).

506   7.   Hafed, Z. M. & Krauzlis, R. J. Microsaccadic suppression of visual bursts in the primate

507   superior colliculus. *Journal of Neuroscience* **30**, 9542–9547 (2010).

508   8.   Burr, D. C., Morrone, M. C. & Ross, J. Selective suppression of the magnocellular visual

509   pathway during saccadic eye movements. *Nature 1994 371:6497* **371**, 511–513 (1994).

510   9.   Idrees, S., Baumann, M. P., Franke, F., Münch, T. A. & Hafed, Z. M. Perceptual saccadic

511   suppression starts in the retina. *Nat Commun* **11**, 1–19 (2020).

512   10.   Idrees, S. *et al.* Suppression without inhibition: how retinal computation contributes to

513   saccadic suppression. *Communications Biology 2022 5:1* **5**, 1–23 (2022).

514   11.   Soto, G. *et al.* Saccadic Suppression Enhances Saliency of Prey-Like Stimuli in the Optic

515   Tectum. *bioRxiv* 2025.02.21.639462 (2025) doi:10.1101/2025.02.21.639462.

516   12.   Jing Chen, X., Valsecchi, M. & Gegenfurtner, K. R. Saccadic suppression measured by

517   steady-state visual evoked potentials. *J Neurophysiol* **122**, 251–258 (2019).

518   13.   Zhang, Y., Valsecchi, M., Gegenfurtner, K. R. & Chen, J. The execution of saccadic eye

519   movements suppresses visual processing of both color and luminance in the early visual

520   cortex of humans. *J Neurophysiol* (2024) doi:10.1152/JN.00419.2023.

521   14.   Liu, X. *et al.* The Superior Colliculus: Cell Types, Connectivity, and Behavior. *Neurosci*

522   *Bull* **38**, 1519 (2022).

523   15.   Cang, J., Savier, E., Barchini, J. & Liu, X. Visual function, organization, and development

524   of the mouse superior colliculus. *Annu Rev Vis Sci* **4**, 239–262 (2018).

525   16.   Chen, C. Y. & Hafed, Z. M. A neural locus for spatial-frequency specific saccadic
526   suppression in visual-motor neurons of the primate superior colliculus. *J Neurophysiol*
527   **117**, 1657–1673 (2017).

528   17.   Wurtz, R. H. & Albano, J. E. Visual-motor function of the primate superior colliculus. *Annu*
529   *Rev Neurosci* **3**, 189–226 (1980).

530   18.   Isa, T., Marquez-Legorreta, E., Grillner, S. & Scott, E. K. The tectum/superior colliculus
531   as the vertebrate solution for spatial sensory integration and action. *Current Biology* **31**,
532   R741–R762 (2021).

533   19.   Chen, C. Y., Ignashchenkova, A., Thier, P. & Hafed, Z. M. Neuronal response gain
534   enhancement prior to microsaccades. *Current Biology* **25**, 2065–2074 (2015).

535   20.   Wu, W. & Hafed, Z. M. Stronger premicrosaccadic sensitivity enhancement for dark
536   contrasts in the primate superior colliculus. *Scientific Reports 2025 15:1* **15**, 1–17 (2025).

537   21.   Easter, S. S. & Nicola, G. N. The development of vision in the zebrafish (Danio rerio).
538   *Dev Biol* **180**, 646–663 (1996).

539   22.   Rolls, E. T. & Cowey, A. Topography of the retina and striate cortex and its relationship to
540   visual acuity in rhesus monkeys and squirrel monkeys. *Exp Brain Res* **10**, 298–310
541   (1970).

542   23.   Schwartz, E. L. Spatial mapping in the primate sensory projection: Analytic structure and
543   relevance to perception. *Biol Cybern* **25**, 181–194 (1977).

544   24.   Dunn, T. W. *et al.* Brain-wide mapping of neural activity controlling zebrafish exploratory
545   locomotion. *Elife* **5**, (2016).

546   25.   Hafed, Z. M. & Chen, C. Y. Sharper, Stronger, Faster Upper Visual Field Representation
547   in Primate Superior Colliculus. *Current Biology* **26**, 1647–1658 (2016).

548   26.   Dehmelt, F. A., Von Daranyi, A., Leyden, C. & Arrenberg, A. B. Evoking and tracking
549   zebrafish eye movement in multiple larvae with ZebEyeTrack. *Nat Protoc* **13**, 1539–1568
550   (2018).

551   27.   Nyström, M. & Holmqvist, K. An adaptive algorithm for fixation, saccade, and glissade
552   detection in eyetracking data. *Behav Res Methods* **42**, 188–204 (2010).

553   28.   Dai, W., Selesnick, I., Rizzo, J. R., Rucker, J. & Hudson, T. A parametric model for
554   saccadic eye movement. *2016 IEEE Signal Processing in Medicine and Biology*
555   *Symposium, SPMB 2016 - Proceedings* (2017) doi:10.1109/SPMB.2016.7846860.

556   29.   Chen, C. Y., Hoffmann, K. P., Distler, C. & Hafed, Z. M. The Foveal Visual
557   Representation of the Primate Superior Colliculus. *Current Biology* **29**, 2109-2119.e7
558   (2019).

30. Scott, J. A. *et al.* Longitudinal analysis of the developing rhesus monkey brain using
magnetic resonance imaging: birth to adulthood. *Brain Struct Funct* **221**, 2847–2871
(2016).

31. Herculano-Houzel, S., Collins, C. E., Wong, P. & Kaas, J. H. Cellular scaling rules for
primate brains. *Proc Natl Acad Sci U S A* **104**, 3562–3567 (2007).

32. Sumbre, G. & de Polavieja, G. G. The world according to zebrafish: how neural circuits
generate behavior. *Front Neural Circuits* **8**, 91 (2014).

33. Bogadhi, A. R. & Hafed, Z. M. Express detection of visual objects by primate superior
colliculus neurons. *Sci Rep* **13**, 1–15 (2023).

34. Eastman, K. M., Huk, A. C. & Galloway, G. J. PLDAPS: A Hardware Architecture and
Software Toolbox for Neurophysiology Requiring Complex Visual Stimuli and Online
Behavioral Control. *Front Neuroinform* **6**, 14243 (2012).

35. Brainard, D. H. The Psychophysics Toolbox. *Spat Vis* **10**, 433–436 (1997).

36. Judge, S. J., Richmond, B. J. & Chu, F. C. Implantation of magnetic search coils for
measurement of eye position: An improved method. *Vision Res* **20**, 535–538 (1980).

37. Fuchs, A. F. & Robinson, D. A. A method for measuring horizontal and vertical eye
movement chronically in the monkey. *J Appl Physiol* **21**, 1068–1070 (1966).

38. Chen, C. Y. & Hafed, Z. M. Postmicrosaccadic Enhancement of Slow Eye Movements.
Journal of Neuroscience **33**, 5375–5386 (2013).

39. Pachitariu, M., Steinmetz, N. A., Kadir, S. N., Carandini, M. & Harris, K. D. Fast and
accurate spike sorting of high-channel count probes with KiloSort. *Adv Neural Inf Process*
Syst **29**, (2016).

40. Pachitariu, M. *et al.* Suite2p: beyond 10,000 neurons with standard two-photon
microscopy. *bioRxiv* 061507 (2017) doi:10.1101/061507.

Acknowledgements

This work was funded by the Deutsche Forschungsgemeinschaft (DFG; German Research Foundation) through the Special Priority Programme: "SPP 2205 Evolutionary optimization of neural processing" (430158665; AR 1076/1-2, HA 6749/3-2, AR 1076/1-1), by the Major Research Instrumentation programme of the DFG (INST 37/1254-1 FUGG), and by the Human Frontier Science Program (HFSP) Young Investigator grant RGY0079. We're grateful to Frédéric Crèvecoeur of UC Louvain, Belgium, for sharing code and advice on how to approximate their original model results.

Author Contributions

G.S. performed experiments in zebrafish optic tectum, analyzed zebrafish calcium data, generated figures, and wrote the manuscript. F.A.D. reproduced the saccadic suppression model, generated figures, and wrote the manuscript. M.P.B. performed macaque experiments, analyzed macaque eye movement data, analyzed macaque electrophysiology data, generated figures, and wrote the manuscript. I.T. analyzed zebrafish and macaque eye trace data and generated figures. Y.Y. performed macaque experiments. T.M. performed macaque experiments and analyzed macaque electrophysiology data. Z.M.H. and A.B.A. designed the overall study, acquired funding and wrote the manuscript.

Competing Interests

The authors declare no competing interests.