Alveolar epithelial cell plasticity and injury memory in human pulmonary fibrosis

Authors:

Taylor S Adams[1,*], Jonas C Schupp[1,2,3,4,*], Agshin Balayev[1], Johad Khoury[1], Aurélien Justet[1,5], Fadi Nikola[1], Laurens J De Sadeleer[6,7], Juan Cala-Garcia[1,8], Marta Zapata-Ortega[6], Panayiotis V Benos[9], John E McDonough[1,6,10,11], Farida Ahangari[1], Melanie Königshoff[12], Jun Ding[13,14,15,16], Robert J Homer[17], Ivan Rosas[8], Xiting Yan[1,18], Bart M Vanaudenaerde[6], Wim A Wuyts[6,7], Naftali Kaminski[1]

*Both authors contributed equally to this work.

Affiliations:

[1] Section of Pulmonary, Critical Care and Sleep Medicine, Yale School of Medicine, New Haven, CT, USA

[2] Department of Respiratory Medicine and Infectious Diseases, Hannover Medical School, Hannover, Germany

[3] Biomedical Research in Endstage and Obstructive Lung Disease Hannover (BREATH), German Center for Lung Research (DZL), Hannover, Germany

[4] Department of Clinical Airway Research, Fraunhofer Institute for Toxicology and Experimental Medicine (ITEM), Hannover, Germany

[5] Department of Respiratory Medicine, University Hospital of Caen, France; Unicaen, CEA, CNRS, UMR 6030, ISTCT Group, GIP Cyceron, Université de Normandie, Caen, France

[6] Laboratory of Respiratory Diseases and Thoracic Surgery (BREATHE), Department of Chronic Diseases and Metabolism, KU Leuven, Leuven, Belgium

[7] Unit for Interstitial Lung Diseases, Department of Pneumology, University Hospitals Leuven, Leuven, Belgium

[8] Section of Pulmonary, Critical Care and Sleep Medicine, Baylor College of Medicine

[9] Department of Epidemiology, University of Florida, Florida, USA

[10] Department of Medicine McMaster University Hamilton, Ontario, Canada.

[11] Firestone Institute for Respiratory Health St. Joseph's Healthcare Hamilton Hamilton, Ontario, Canada

[12] Department of Medicine, University of Pittsburgh, Pittsburgh, Pennsylvania, USA

[13] Department of Medicine, Faculty of Medicine and Health Sciences, McGill University, Montreal, Quebec, Canada

38 [14] Meakins-Christie Laboratories, Translational Research in Respiratory Diseases Program, Research Institute of the McGill University Health Centre, Montreal, Quebec, Canada
40 [15] School of Computer Science, McGill University, Montreal, Quebec, Canada
41 [16] Mila - Quebec AI Institute, Montreal, Quebec, Canada
42 [17] Department of Pathology, Yale School of Medicine, New Haven, CT, USA
43 [18] Department of Biostatistics, Yale School of Public Health, New Haven, CT, USA
44
45

46 **Corresponding author:**

47 Naftali Kaminski. Boehringer-Ingelheim Endowed Professor of Internal Medicine. Chief of

48 Pulmonary, Critical Care and Sleep Medicine. Yale School of Medicine. 300 Cedar Street, TAC–

49 441 South. P.O. Box 208057. New Haven, CT 06520-8057. Tel: 203-7854162, Fax: 203-7853826,

50 E-Mail: naftali.kaminski@yale.edu

Abstract

Acute and repetitive lung epithelial injury can lead to irreversible and even progressive pulmonary fibrosis; Idiopathic pulmonary fibrosis (IPF) is a fatal disease and quintessential example of this phenomenon. The composition of epithelial cells in human pulmonary fibrosis – irrespective of disease etiology – is marked by the presence of Aberrant Basaloid cells: an abnormal cell phenotype with pro-fibrotic and senescent features, localized to the surface of fibrotic lesions. Despite their relevance to human pulmonary fibrosis, the exotic molecular profile of Aberrant Basaloid cells has obscured their etiology, preventing insights into how or why these cells emerge with fibrosis. Here we identify cellular intermediaries between Aberrant Basaloid and normal alveolar epithelial cells in human IPF tissue. We track the emergence of Aberrant Basaloid cells from alveolar epithelial cells *ex vivo* and uncover a role for similar cells in epithelial regeneration under normal conditions. Lastly, we characterize the epigenetic changes that distinguish Aberrant Basaloid cells from their progenitors and identify hallmarks of AP-1 injury memory retention. This study elucidates the phenomenon of maladaptive epithelial plasticity and regeneration in pulmonary fibrosis and re-contextualizes therapeutic strategies for epithelial dysfunction.

Main

The alveoli of the lung are the primary interface between the atmosphere and our circulation. Each alveolus is structurally optimized to maximize two critical functions: gas exchange and pulmonary compliance. On the alveolar surface, these tasks are delegated to two distinct types of epithelial cells: alveolar epithelial type I (ATI) cells enable gas-diffusion, while alveolar type II (ATII) cells produce surfactants that reduce surface tension[1,2]. Environmental exposure carries the risk of injury and ATII serve as the primary stem cells for alveolar epithelial regeneration. While alveolar epithelial injury and regeneration are a normal part of life, repeated injury can potentially lead to progressive and irreversible fibrosis[3]. Understanding both functional and maladaptive responses to alveolar epithelial regeneration remain critical topics in pulmonary medicine[4,5].

Single cell RNA sequencing (scRNAseq) studies enabled the discovery of Aberrant Basaloid cells in tissue from patients with end-stage human lung fibrosis[6,7]. Aberrant Basaloid cells lack hallmarks of ATII and ATI cells and are characterized by a pronounced epithelial mesenchymal transition (EMT) signal (CDH2, VIM, COL1A1) and some overlapping features with airway basal cells (KRT17, TP63, LAMB3). Notably, this Aberrant Basaloid cell phenotype is found across many diseases including interstitial lung diseases like idiopathic pulmonary fibrosis (IPF)[6,7] and systemic sclerosis[8], COPD[7] and COVID-19 induced fibrosis[9,10]. A concurrent discovery was made in mice of Transitional Krt8+ cells: a cell phenotype which emerges from ATII following pulmonary injury and persist until resolution before becoming ATI cells[11–16]. Murine Transitional Krt8+ cells are a conserved phenotype following genotoxic, endotoxic and mechanical forms of lung injury: they lack hallmarks of ATII and ATI cells and are characterized by expression of Cldn4, Krt8, Krt19, Sfn and Fn1[11–16]. Despite their situational likeness, murine Transitional Krt8+ cells lack most defining features of human disease Aberrant Basaloid cells[17]; this lack of apparent molecular similarity raises questions regarding the degree of cellular orthology between the two epithelial phenotypes. Unlike Transitional Krt8+ cells, the cellular origin of Aberrant Basaloid cells in human disease remains unknown. More importantly, it remains unclear whether Aberrant Basaloid cells play a normal role in human epithelial injury repair or whether their extreme expression profile and persistence through end-stage fibrosis are indicative of an abnormal phenomenon.

In this study, we perform single nuclei RNA sequencing (snRNAseq) on differentially fibrotic regions in human IPF lungs and identify intermediate states between Aberrant Basaloid cells and

98 other epithelial cell types. We longitudinally profile a model of Aberrant Basaloid development

99 and uncover a role for these cells in normal regeneration. Lastly, we dissect the epigenetic profile

100 of Aberrant Basaloid cells and describe what makes them fundamentally different from normal

101 epithelial cells.

Results

Tissue specimens of differentially fibrotic IPF and control lungs were prepared and evaluated as

described by McDonough *et al.*[18]. Explanted IPF lungs or donor lungs declined for transplantation

were inflated, frozen and CT scanned. Small cores of tissue were isolated and subject to microCT

assessment of surface density: a proxy of localized fibrosis (Fig. 1A). snRNAseq was then

performed on 54 control and IPF cores with variable levels of fibrosis, and cell types were classified

into major lineages (Figures 1A, S1A). To provide further context, epithelial nuclei were integrated

with epithelial cell data from two previously published scRNAseq datasets of lung disease (Fig.

1A; S1A)[6,7] before cell types were collectively re-classified (Fig. 1B, Supp. Table S1).

A "Missing Link" Between Aberrant Basaloid and Alveolar Epithelial Cells

The unbiased cell representation afforded by snRNAseq (Figure S1A) enabled us to identify a

previously overlooked population of epithelial cells in IPF, herein referred to as Alveolar

Intermediate (AT*i*) cells (Figure 1B). Analyzing the intersection of disease-associated cell marker

genes with otherwise distinct markers from control epithelial cell types, we observe AT*i* cells to

share some features with ATI (NTM, GPC5), ATII (SERPINA1, SFTA2) and Aberrant Basaloid

cells (Fig. 1C,D). The distinct combination of positive and negative features characterizing AT*i*

cells (Fig. 1D) indicate that these are a unique population of cells straddling the phenotypic

boundary of otherwise distinct cell states in health and disease. AT*i* cells share only a subset of

features with Aberrant Basaloid cells (Fig. 1D): Overlapping features include DNA damage-related

genes MDM2 and GDF15; elevated levels of TGFβ-activating integrin subunit ITGB6 and distinct

expression of TGFβ signaling genes TGFBR2, SMAD4 and MIR100HG - a host gene for micro-

RNA let-7a; hedgehog signaling repressor PTCHD4; the Angiotensin-1 converting protease CPA6,

and CTSE – a metastatic hallmark of pancreatic ductal adenocarcinoma and Barretts'

Esophagus[19,20]. In contrast, AT*i* cells lack basaloid features TP63 and KRT17; senescence

hallmarks CDKN2A and CDKN2B; TGFβ-inducible NOX4, EPHB2, CASC15 and TGFBI;

extracellular matrix (ECM) remodeling genes COL1A1, COL5A1, PRSS2, SPINK1 ADAMTS9,

MMP7; the epithelial-mesenchymal transition (EMT) hallmark CDH2 and mutation-inducing

transposase PGBD5 (Figure 1D).

AT*i* and Aberrant Basaloid cells with the same features are also observed in publicly available

datasets of end-stage COVID-19 lungs[10,21,22] (Supp. Fig. S2A, Supp. Table S8) extending their

relevance from chronic to acute etiologies of pulmonary fibrosis.

Composition of Abnormal Epithelial Cells in IPF

The median percent of epithelial cells classified as Aberrant Basaloid are similar among IPF

samples across all 3 datasets, ranging from 2.72% to 2.91% (Figure 1E). AT*i* cells are significantly

enriched in IPF samples vs controls in snRNAseq (Wilcox pVal=2.165E-5) and in Adams *et al.*

(Wilcox pVal=0.043), though not in Habermann *et al.* (Wilcox pVal=0.22; Fig. 1E). The proportion

of AT*i* cells is significantly higher in IPF snRNAseq samples (median 3.87%) when compared to

IPF samples from either scRNAseq datasets (1.32%, p=0.0015 in Adams *et al.*; 2.17%, p=0.038 in

Habermann *et al.*; Figure 1E), suggesting that cell dissociation survivor bias negatively impacted

the recovery of these cells in previous studies[23].

We explored the relationship between tissue surface density, a proxy for fibrosis, and the percent

of abnormal epithelial cells. The decline in surface density across IPF tissues is significantly

associated with a decline in the proportion of ATII cells (p=5.34e-4) alongside an increase in large

airway-associated secretory (p=0.013) and ciliated cells (p=0.023) (Figure 1F; Supp. Table S2).

Neither Aberrant Basaloid nor AT*i* cell proportions had significant relationships with tissue surface

density in IPF (Fig. 1F, Supp. Table S2). This suggests that the emergence of these cells pre-dates

the development of pathophysiologic characteristics of fibrosis and it is thus a potential target of

anti-fibrotic drugs.

Assessing co-occurrence patterns across all lung cores, the most positive compositional correlates

of surface density are unsurprisingly ATI and ATII cells (Figure 1G). Notably, the strongest

abundance correlates of AT*i* cells are Aberrant Basaloid cells (Spearman's rho=0.813) followed by

epithelial cells in cell cycle (Spearman's rho=0.442; Figure 1G), indicating spatial associations

with both senescent and proliferating epithelial cells.

The Continuum of States Between ATI, ATII and Aberrant Basaloid Cells

After evaluating disease-relevant cell types in discrete terms, we assessed their phenotypic connectivity to healthy cell types to model continuous trajectories. Common epithelial cell types were re-clustered into granular sub-populations; inter-connectivity between sub-populations was evaluated with the lineage reconstruction technique "partition-based graph abstraction" (PAGA)[24]. Using an edge confidence threshold > 0.2, all healthy cell types remain connected with a plausible lineage topology, and Aberrant Basaloid cells are exclusively connected to the graph via IPF AT*i* cells towards either ATI or ATII cells (Fig. 2A). To avoid spurious connections in trajectory analysis, low confidence edge pruning from the PAGA graph was extended to connections in the original nuclei-level graph, and a force-directed graph layout of pruned nuclei similarities was built to complement pseudotime analysis (Fig. 2A). Lacking certainty in the orientation of trajectories, we chose to model independent trajectories from either ATII or ATI nuclei towards disease-restricted Aberrant Basaloid nuclei. Generalized additive models (GAMs) of genes were fit to changes in pseudotime across nuclei from each trajectory and organized pseudotemporally by expression peak.

Of the two trajectories, nuclei along the trajectory from ATII towards Aberrant Basaloid features a more gradual pseudotime mapping and more detailed model of pseudotemporal gene dynamics[25] (Fig. 2B). While markers of ATII cell identity (SFTPC, LAMP3, PDE4D, ERBB4) decline, "Early Peak" features emerge including Let-7a host gene MIR100HG, unconventional myosin MYO16 and the peptidase CTSE (Fig. 2B,C). "Early Peak" features then decline with the loss of cell identity and rise of "Late Peak" genes – including: GDF15, FGF13, ITGAV and CPA6 – before specific Aberrant Basaloid hallmarks emerge (TP63, CDH2, CASC15, CDKN2A; Fig. 2B,C). Trajectory analysis of ATI towards Aberrant Basaloid revealed a similar – though less detailed – sequence of gene expression patterns. ATI features (RTKN2, NCKAP5, CAV1) decline while "Early Peak" features (CTSE, MYO16, STK32A, NTM, GPC5) rise prior to "Late Peak" features (GDF15, CPA6), followed by peaks of Aberrant Basaloid hallmarks (CDH2, TP63, CDKN2A; Fig. 2C,D). Though it remains unclear whether these trajectories represent real lineage fates, this analysis demonstrates that AT*i* cells are a diverse population occupying a branching continuum of phenotypes between ATI, ATII and IPF Aberrant Basaloid cells.

Immunofluorescent microscopy of control and IPF tissues confirmed the presence of both SFTPC+ ATII cells and AGER+ ATI cells with "Early Peak" AT*i* marker CTSE in affected regions of IPF

lungs (Fig. 3B) but not in control lungs (Fig. 3A). Despite "Late Peak" features like CPA6

transcriptionally emerging with the loss of cell-type markers (Fig. 2B,D), we observed many

SFTPC+/CPA6+ and AGER+/CPA6+ cells in IPF lungs, as well as in hyper-keratinized SFTPC-

/AGER-/CPA6+ cells (Fig. 3D), confirming the presence of AT*i* populations in IPF lungs.

Longitudinal Assessment of *ex vivo* Fibrosis

To study the emergence of Aberrant Basaloid cells in real time and compare *ex vivo* outcomes to

in vivo disease, we took advantage of a recent observation that Aberrant Basaloid cells arise *ex vivo*

from disease-free human precision cut lung slices (PCLS) following 5 days of exposure to a

"fibrotic cocktail" (FC: PDGF-αβ, TGF-β, LPA, and TNFα)[26–28]. We longitudinally assessed this

model with snRNAseq (Fig. 4A,B) and observed that both cocktail and control PCLS undergo a

decline in epithelial cells over the first 48 hours, before gradually rising (Fig. 4C, Supp. Table S3,

Supp. Table S4). Concurrent with this gradual rise is a dramatic increase in epithelial cell

proliferation: 3-8% of epithelial cells from 72 hours onwards are undergoing mitosis (Fig. 4C).

Hereby, FC exposure significantly reduces the rate of epithelial mitosis between 72 and 120 hours

(t-test pVal= 0.010607), from an average of 6.3% in controls to 3.8% with FC (Fig. 4C).

The phenotypic landscape of PCLS alveolar epithelial cells is characterized by many distinct

phenotypic substates, associated with different points in time (Fig. 4D,E). All substates could be

classified under *in vivo* conventions based on the conservation of marker genes: CAV1, COL4A1,

COL4A2 in ATI and PDE4D, ACOXL and ABCC4 in ATII cells (Fig. 4E, Supp. Fig. S3A). The

majority of different substates are associated with samples from the first 48 hours, reflecting a

period of dynamic behavior with characteristic expression patterns. Early timepoint ATII subtypes

exhibit features of stress (HSPA1A, STIP1) and metabolic change (ALDH1L2, GK, KYNU) (Fig.

4E, Supp. Fig. S3E). ATI substates enriched in the first 48 hours exhibit hallmarks of stress (FOS,

JUN, HSPA1A), cell cycle arrest (CDKN1C, CCN2, ID2), endoplasmic reticular stress (ERO1B,

CLPTM1L, PLA2G4) and cytoskeleton regulation (FSCN1, SAMSN1) (Fig. 4E, Supp. Fig. S3B).

PCLS epithelial cells from 72 hours onwards are primarily classified into just three phenotypes:

"ATIend", "ATIIend" and Aberrant Basaloid – suggesting that some degree of homeostasis has

been achieved. Of interest, Aberrant Basaloid cells were detected in both control and FC PCLS as

early as hour 24 onwards. Aberrant Basaloid cells are significantly more common in FC samples

from 72 hours onwards vs control (tTest pVal=0.004385), increasing from an average 1.25% of

epithelial cells in control PCLS to 30.4% following FC treatment (Fig. 4E).

A detailed description of temporal ATI and ATII gene expression dynamics is provided in the

supplement (Supp. Discussion; Figs. S3, S4). These results depict a biphasic response of alveolar

epithelial cells to PCLS experiment conditions where 48 hours of injury response and acclimation

are followed by several days of phenotypic stability and tenacious regeneration.

Comparing control, fibrotic cocktail and IPF Aberrant Basaloid cells

To model the etiology of Aberrant Basaloid cells in PCLS, graph connectivity and pseudotime

analysis was performed as described earlier. Similar to *in vivo* results, the strongest connectivities

to PCLS Aberrant Basaloid cells are substates of both ATI and ATII cells (Supp. Fig. S3B). Unlike

the *in vivo* results, the AT*i* gene signature was not particularly representative of any cell population

detected in PCLS, with the prominent exception of FRMD5 and MIR100HG in the PCLS

pseudotime trajectory from ATII towards Aberrant Basaloid cells (Supp. Fig. S3E).

Comparing markers of PCLS "ATIend", "ATIIend" and Aberrant Basaloid to their *in vivo* IPF

equivalents, we find most *in vivo* ATI and ATII hallmarks are conserved *ex vivo* (Fig. 4F, Supp.

Table S5). Although only 17.5% of the *in vivo* Aberrant Basaloid signal overlaps with ex vivo

variants, this overlap encapsulates 54.6% of their defining ex vivo features (Fig. 4F). Control PCLS

Aberrant Basaloid cells share many features with FC and *in vivo* variants, including KRT17,

ADAMTS9, SLC35F3, PTGS2, HMGA2, FOSL2 and elevated levels of KRT7 and ITGAV (Fig.

4F). However, unlike FC PCLS Aberrant Basaloid cells, control PCLS variants notably lack core

disease hallmarks of aberrancy (CDH2, COL1A1, SOX4, SERPINE1, EPHB2) and basal-likeness

(TP63) (Fig. 4F). Though Aberrant Basaloid-like cells are observed in control PCLS, FC

stimulation induces a more complete molecular representation of Aberrant basaloid cells described

in IPF.

Among proliferating epithelial cells, ATII were the most commonly detected, though SFTPB+

Secretory, Aberrant Basaloid and even ATI cells with mitotic hallmarks are found. Strikingly, 30%

of all control Aberrant Basaloid cells are observed undergoing mitosis compared to just 0.5% of

FC stimulated variants (Fig. 4G); for reference, 7% of all ATII cells and 1.2% of ATI cells are

observed mitotic from 24 hours onwards. From hour 72 onwards, FC induced a dramatic decline

in the percent of mitotic Aberrant Basaloid (tTest pVal=0.04077) from a per-sample average of

32.6% in controls to just 0.49% (Fig. 4G). FC did not significantly affect the percent of ATII cells

undergoing mitosis over this time (tTest pVal=0.6304), indicating that the overall decline in

epithelial proliferation induced by FC arose not from direct effects on ATII proliferation, but rather
their displacement with senescent Aberrant Basaloid cells.

FC stimulation thus confers a more pathologically relevant presentation of Aberrant Basaloid cells
in PCLS in terms of abundance, molecular signature and cell cycle arrest. The observations that
control PCLS aberrant basaloid cells are rare yet frequently found dividing, suggest they are
normally a transient phenotype associated with homoeostatic repair response. This contrasts with
abundant yet rarely dividing Aberrant basaloid Cells found in FC-stimulated PCLS, patterns
indicative of a change towards persistence and senescence. Beyond highlighting FC's ability to
emulate the disease phenotype of Aberrant Basaloid cells, these results place a spotlight on control
PCLS Aberrant Basaloid cells, understanding their role and plasticity may be key to distinguishing
normal repair from maladaptive outcomes.

Chromatin Accessibility Patterns of Aberrant Basaloid Cells and ATi cells
To explore the epigenetic landscape of epithelial cells in lung fibrosis, we subjected three control
and seven IPF lung samples to snMultiOme analysis, where each nucleus is simultaneously profiled
with snRNAseq and snATAC-seq chemistry (Fig. 5A). Cell types were classified by transcriptional
identity, and ATAC fragments from each cell type per sample were pooled for pseudo bulk peak
calling using the R package ArchR[29]; fragments from merged peaks were used in downstream
analyses (Fig. 5A). UMAP embeddings of gene expression and chromatin-accessibility show
remarkable topological similarities among epithelial cells (Fig. 5B,C). ATI, ATII and Aberrant
Basaloid cells occupy unique regions of both RNA and ATAC feature-space. AT*i* straddle the
feature-space between these cell types in both UMAP embeddings, however their ATAC profile
was far less distinct (Supp. Table S6). The tight co-localization of Aberrant Basaloid cells in
unsupervised ATAC-space indicates a divergence from normal epithelial cell chromatin
accessibility.

We compared differentially expressed genes to differentially accessible genes using ArchR's Gene
Score[29] (Supp. Table S6). Across epithelial cell types, distinct marker genes often have
correspondingly distinct accessibility patterns – MIR205HG in basal cells, FOXJ1 in ciliated,
LAMP3 and SFTPC in ATII, RTKN2 and NCKAP5 in ATI (Fig. 5D). Likewise, many uniquely
expressed genes in Aberrant Basaloid cells are associated with uniquely accessible chromatin,
including TGF-B signaling mediators SKIL, FSTL1, LTBP1 and HMGA2, as well as EMT

hallmarks CDH2, COL6A1 and COL18A1 (Fig. 5D). AT*i* cells have no significantly distinct

ATAC Gene Scores, but – alongside Aberrant Basaloid cells – they share ATI features like KRT7,

NTM and collagen IV and show elevated accessibility of pathological features like CPA6 and the

cytoskeleton regulators FGD6 and TMEM132D (Fig. 5D). Collectively, Aberrant Basaloid cells

are both epigenetically and transcriptionally distinct from normal lung epithelial cells. The

chromatin accessibility profile of AT*i* cells is less categorically distinct, consistent with their

diverse intermediate transcriptional states.

AP-1 Transcription Factor Accessibility is a Defining Feature of Aberrant Basaloid

Cells

We explored patterns of transcription factor (TF) motif enrichment across differentially accessible

peaks using the TF database CIS-BP[30]. TF motifs with the largest contributions to variance in the

data are primarily associated with specific epithelial cell types (Fig. 5E). The top TF families

associated with ATII cells are C/EBP, NKX-2 and FOXA1/2; in ATI they are TEAD, KLF, GATA

(Supp. Table S7) – observations consistent with previous studies[31–33]. Differentially accessible

motif positions from these families have cis-regulatory links to over 60% of their respective

transcriptional features (Supp. Fig. S4A,B).

TF motifs for JUND, FOSL2, SMARCC1, JUNB, FOSL1, FOS, JUN, BACH2, BACH1, FOSB

and JDP2 are outliers in terms of their contribution to overall variance in the data (Fig. 5E, left)

and in terms of the significance of their enrichment among Aberrant Basaloid relative to other cells

(pVal<1E-20; Figure 5SE, right). Further examination of the top 15 significantly enriched TF

motifs among Aberrant Basaloid cells revealed that they all shared a common sequence: TGA G/C

TCA (Fig. 5F, Supp. Table S7) – the consensus sequence for AP-1 – we subsequently focused on

this particular sequence motif.

Though 12.5% of all merged peaks contain one or more AP-1 consensus sequences, they are found

in over 50% of all differentially accessible peaks in Aberrant Basaloid cells (hypergeometric test

p-value < 1e-50; Fig. 5G). Cell-level AP-1 enrichment z-scores are profoundly elevated in Aberrant

Basaloid compared to other cell types (Wilcoxon test pVal=2.2e-16; Fig. 5I), where the median

cell's AP-1 enrichment zScore is 5.29; Alveolar Intermediate cells have the second highest median

zScore of 1.09. Footprint analysis of accessible chromatin surrounding AP-1 sequences reveals a

prominent dip at the binding site among Aberrant Basaloid cells (Fig. 5I), indicating AP-1 TF

binding. AP-1 TF motif sequences are found in the majority of differentially accessible chromatin
in Aberrant Basaloid cells, these cells have the highest breadth of accessible AP-1 binding sites of
any cell in the lung and these sites are actively occupied.

Discussion

This study provides an unprecedented view of human alveolar epithelial plasticity and unearths
important context behind the identity of Aberrant Basaloid cells and their role in regeneration and
disease. Analyzing differentially affected IPF tissues, we observe similar frequencies of aberrant
basaloid cells across disease tissues, irrespective of their level of fibrosis. We uncover AT*i* cells: a
diverse spectrum of cell phenotypes between Aberrant Basaloid cells and ATII and ATI cells – we
verify their presence across multiple cohorts and diseases, and localize them in fibrotic tissues.
Tracking the emergence of these cells in human PCLS, we observe that fibrotic cocktail stimulation
reduces the overall rate of epithelial cell mitosis while increasing the abundance of Aberrant
Basaloid cells and imparting them with a molecular profile more aligned with human pulmonary
fibrosis. Finally, we establish that IPF Aberrant Basaloid cells have a common epigenetic profile
which is distinct from all other lung cell types, characterized by extreme levels of AP-1 family TF
chromatin accessibility and binding site occupation.

Upon their discovery, the exotic molecular profile of Aberrant Basaloid cells precluded the ability
to infer where these cells originated, a limitation that undercut their translational impact[6,7,34]. Not
only was it unclear which cells should be prophylactically targeted to prevent Aberrant Basaloid
emergence, but interventions aimed at targeting the behavior of Aberrant Basaloid cells lacked a
phenotypic compass to orient the desired outcome. The discovery of AT*i* cells in IPF and fibrotic
COVID-19 lungs represents the first example of a "missing link" between these pathological cells
and normal epithelial lung cell phenotypes in human lung disease. In both *in vivo* and *ex vivo*
models of fibrosis, we observe phenotypic intermediate states between Aberrant Basaloid cells and
both ATI and ATII cells – however the PCLS intermediate phenotypes are largely different from
the AT*i* cells found in human disease. In addition, studies of cells exposed to fibrosis-associated
signaling molecules have shown that human – but not mouse - ATII[35] and Basal cells[36] are each
capable of achieving an Aberrant Basaloid feature set. Overall, this suggests that the Aberrant
Basaloid cell phenotype is an example of cellular convergence: a singular state that different cell
types acquire upon injury.

A key limitation of this study is its inability to provide evidence regarding the orientation of

changes between these cell types, beyond the assumption that Aberrant Basaloid cells emerge from

normal epithelial cells. It is further possible that AT*i* are not transitioning at all. In multiple murine

models of fibrosis, it is well established that Transitional Krt8+ cells temporarily emerge from ATII

before becoming ATI cells [11–14]; human disease Aberrant Basaloid cells could reflect malfunction

in this process. Perhaps the better analogue to transient murine Transitional Krt8+ cells are the

human control PCLS Aberrant Basaloid cells, serving conceptually similar roles in epithelial

alveolar regeneration – a form of cellular homoplasy.

Genome-wide chromatin accessibility profiling reveals that IPF Aberrant Basaloid cells share a

common epigenetic profile distinct from any other cell type in the lung, forming a unique valley in

Waddington's epigenetic landscape model[37]. Their defining molecular hallmarks are often in

regions of the genome uniquely accessible to them, denoting a direct relationship between their

functional profile and epigenetic programming.

A defining property of the IPF Aberrant Basaloid epigenome is the degree of AP-1 TF accessibility

on display. Gene expression of AP-1 is widely associated with cellular responses to injury and

stress, including in lung epithelial cells following cigarette smoke exposure[38], viral[39] or bacterial

infection[40], or in the stress response of adjusting to the PCLS environment (Supp. Fig. S4B).

However, FOSL2 is the only AP-1 family gene we observe constitutively expressed in either IPF

or *ex vivo* Aberrant Basaloid variants, although Footprint analysis indicates that TFs are occupying

these binding sites. It is conceivable that these changes arose from transcriptional AP-1 responses

to injuries which occurred long before the fibrotic lungs described herein were profiled. Although

the AP-1 motif has been among many TF motifs enriched in murine Transitional Krt8+ cells[11,12]

and aged human AT2 cells[41], this is the first time that genome-wide AP-1 epigenetics are implicated

in epithelial cell dysfunction of human non-cancerous lung disease.

Similar AP-1 epigenetics are central to the first recognized instance of innate memory in non-

immune cells: skin epithelial stem cells with a history of injury exposure maintained enhanced AP-

1 chromatin accessibility and sustained AP-1 binding[42,43]. Retaining this inflammation memory

allows cells to respond to future injuries faster and more robustly[44]. Maladaptive outcomes of AP-

1 injury memory have been implicated in several diseases where chronic injury leads to epithelial

stem cell metaplasia: from benign cases like nasal polyps in chronic rhinosinusitis to precancerous

conditions like Barrett's Esophagus and pancreatic intra-epithelial neoplasia[19,20]. Our study

describes the pulmonary disease rendition of AP-1 epigenetic associated dysfunction: where abnormal Aberrant Basaloid cells retain the memory of injuries responsible for triggering lung fibrosis.

In summary, our study provides detailed evidence that alveolar epithelial cell plasticity and regeneration play a significant role in human lung disease. Our findings establish maladaptive epithelial plasticity and regeneration as key mechanistic events in the shift of alveolar epithelial cells from injured innocent bystanders to potential drivers of disease, as proposed by Pardo and Selman nearly 20 years ago[3]. Beyond their pathogenetic significance, these observations highlight the restoration of normal epithelial differentiation as a key therapeutic aim in pulmonary fibrosis.

References

1. Travaglini, K. J. *et al.* A molecular cell atlas of the human lung from single-cell RNA sequencing. *Nature* **587**, 619–625 (2020).

2. Wang, Y., Wang, L., Ma, S., Cheng, L. & Yu, G. Repair and regeneration of the alveolar epithelium in lung injury. *FASEB J* **38**, e23612 (2024).

3. Selman, M. & Pardo, A. Role of epithelial cells in idiopathic pulmonary fibrosis: from innocent targets to serial killers. *Proc Am Thorac Soc* **3**, 364–372 (2006).

4. Jones, D. L. *et al.* An injury-induced mesenchymal-epithelial cell niche coordinates regenerative responses in the lung. *Science* **386**, eado5561 (2024).

5. Parimon, T. *et al.* Senescence of alveolar epithelial progenitor cells: a critical driver of lung fibrosis. *Am J Physiol Cell Physiol* **325**, C483–C495 (2023).

6. Habermann, A. C. *et al.* Single-cell RNA sequencing reveals profibrotic roles of distinct epithelial and mesenchymal lineages in pulmonary fibrosis. *Science Advances* **6**, eaba1972 (2020).

7. Adams, T. S. *et al.* Single-cell RNA-seq reveals ectopic and aberrant lung-resident cell

populations in idiopathic pulmonary fibrosis. *Science Advances* **6**, eaba1983 (2020).

8. Valenzi, E. *et al.* Disparate Interferon Signaling and Shared Aberrant Basaloid Cells in

Single-Cell Profiling of Idiopathic Pulmonary Fibrosis and Systemic Sclerosis-Associated

Interstitial Lung Disease. *Front Immunol* **12**, 595811 (2021).

9. Narasimhan, H. *et al.* An aberrant immune-epithelial progenitor niche drives viral lung

sequelae. *Nature* **634**, 961–969 (2024).

10. Bharat, A. *et al.* Lung transplantation for patients with severe COVID-19. *Sci Transl Med* **12**,

eabe4282 (2020).

11. Choi, J. *et al.* Inflammatory Signals Induce AT2 Cell-Derived Damage-Associated Transient

Progenitors that Mediate Alveolar Regeneration. *Cell Stem Cell* **27**, 366-382.e7 (2020).

12. Kobayashi, Y. *et al.* Persistence of a regeneration-associated, transitional alveolar epithelial

cell state in pulmonary fibrosis. *Nat Cell Biol* **22**, 934–946 (2020).

13. Jiang, P. *et al.* Ineffectual Type 2-to-Type 1 Alveolar Epithelial Cell Differentiation in

Idiopathic Pulmonary Fibrosis: Persistence of the KRT8hi Transitional State. *Am J Respir*

Crit Care Med **201**, 1443–1447 (2020).

14. Strunz, M. *et al.* Alveolar regeneration through a Krt8+ transitional stem cell state that

persists in human lung fibrosis. *Nat Commun* **11**, 3559 (2020).

15. Niethamer, T. K. *et al.* Longitudinal single-cell profiles of lung regeneration after viral

infection reveal persistent injury-associated cell states. *Cell Stem Cell* **32**, 302-321.e6 (2025).

16. Konkimalla, A. *et al.* Transitional cell states sculpt tissue topology during lung regeneration.

Cell Stem Cell **30**, 1486-1502.e9 (2023).

17. Xie, T. *et al.* Abnormal respiratory progenitors in fibrotic lung injury. *Stem Cell Res Ther* **13**,

64 (2022).

18. McDonough, J. E. *et al.* Transcriptional regulatory model of fibrosis progression in the

human lung. *JCI Insight* **4**, 131597 (2019).

19. Fisher, O. M. *et al.* High Expression of Cathepsin E in Tissues but Not Blood of Patients with

Barrett's Esophagus and Adenocarcinoma. *Ann Surg Oncol* **22**, 2431–2438 (2015).

20. Azuma, T. *et al.* Expression of cathepsin E in pancreas: a possible tumor marker for pancreas,

a preliminary report. *Int J Cancer* **67**, 492–497 (1996).

21. Delorey, T. M. *et al.* COVID-19 tissue atlases reveal SARS-CoV-2 pathology and cellular

targets. *Nature* **595**, 107–113 (2021).

22. Melms, J. C. *et al.* A molecular single-cell lung atlas of lethal COVID-19. *Nature* **595**, 114–

119 (2021).

23. Zheng, Y. *et al.* A deep generative model for deciphering cellular dynamics and in silico drug

discovery in complex diseases. *Nature Biomedical Engineering* (2025) doi:10.1038/s41551-

025-01423-7.

24. Wolf, F. A. *et al.* PAGA: graph abstraction reconciles clustering with trajectory inference

through a topology preserving map of single cells. *Genome Biol* **20**, 59 (2019).

25. Setty, M. *et al.* Characterization of cell fate probabilities in single-cell data with Palantir. *Nat*

Biotechnol **37**, 451–460 (2019).

26. Lang, N. J. *et al.* Ex vivo tissue perturbations coupled to single-cell RNA-seq reveal

multilineage cell circuit dynamics in human lung fibrogenesis. *Sci Transl Med* **15**, eadh0908

(2023).

27. Alsafadi, H. N. *et al.* An ex vivo model to induce early fibrosis-like changes in human

precision-cut lung slices. *American Journal of Physiology-Lung Cellular and Molecular*

Physiology **312**, L896–L902 (2017).

28. Lehmann, M. *et al.* Precision Cut Lung Slices: Emerging Tools for Preclinical and

Translational Lung Research. An Official American Thoracic Society Workshop Report. *Am*

J Respir Cell Mol Biol **72**, 16–31 (2024).

29. Granja, J. M. *et al.* ArchR is a scalable software package for integrative single-cell chromatin

accessibility analysis. *Nat Genet* **53**, 403–411 (2021).

30. Weirauch, M. T. *et al.* Determination and inference of eukaryotic transcription factor

sequence specificity. *Cell* **158**, 1431–1443 (2014).

31. Hassan, D. & Chen, J. CEBPA restricts alveolar type 2 cell plasticity during development and

injury-repair. *bioRxiv* 2023.10.10.561625 (2023) doi:10.1101/2023.10.10.561625.

32. Penkala, I. J. *et al.* Age-dependent alveolar epithelial plasticity orchestrates lung homeostasis

and regeneration. *Cell Stem Cell* **28**, 1775-1789.e5 (2021).

33. Zepp, J. A. *et al.* Genomic, epigenomic, and biophysical cues controlling the emergence of

the lung alveolus. *Science* **371**, eabc3172 (2021).

34. Adegunsoye, A. *et al.* Genetics and Genomics of Pulmonary Fibrosis: Charting the Molecular

Landscape and Shaping Precision Medicine. *Am J Respir Crit Care Med* **210**, 401–423

(2024).

35. Kathiriya, J. J. *et al.* Human alveolar type 2 epithelium transdifferentiates into metaplastic

KRT5+ basal cells. *Nat Cell Biol* **24**, 10–23 (2022).

36. Uthaya Kumar, D. B. *et al.* Bronchial epithelium epithelial-mesenchymal plasticity forms

aberrant basaloid-like cells in vitro. *Am J Physiol Lung Cell Mol Physiol* **322**, L822–L841

(2022).

37. *The Strategy of the Genes: A Discussion of Some Aspects of Theoretical Biology*. (Routledge,

Abingdon, Oxon New York, New York, 2014).

38. Zhong, C.-Y., Zhou, Y.-M., Douglas, G. C., Witschi, H. & Pinkerton, K. E. MAPK/AP-1

signal pathway in tobacco smoke-induced cell proliferation and squamous metaplasia in the

lungs of rats. *Carcinogenesis* **26**, 2187–2195 (2005).

39. Ludwig, S. *et al.* Influenza virus-induced AP-1-dependent gene expression requires activation

of the JNK signaling pathway. *J Biol Chem* **276**, 10990–10998 (2001).

40. Rajasekaran, S. *et al.* Visualization of Fra-1/AP-1 activation during LPS-induced

inflammatory lung injury using fluorescence optical imaging. *Am J Physiol Lung Cell Mol*

Physiol **309**, L414-424 (2015).

41. Liu, X. *et al.* Transcriptomics Analysis Identifies the Decline in the Alveolar Type II Stem

Cell Niche in Aged Human Lungs. *Am J Respir Cell Mol Biol* **71**, 229–241 (2024).

42. Naik, S. *et al.* Inflammatory memory sensitizes skin epithelial stem cells to tissue damage.

Nature **550**, 475–480 (2017).

43. Larsen, S. B. *et al.* Establishment, maintenance, and recall of inflammatory memory. *Cell*

Stem Cell **28**, 1758-1774.e8 (2021).

44. Gonzales, K. A. U. *et al.* Stem cells expand potency and alter tissue fitness by accumulating

diverse epigenetic memories. *Science* **374**, eabh2444 (2021).

45. Van den Berge, K. *et al.* Trajectory-based differential expression analysis for single-cell

sequencing data. *Nat Commun* **11**, 1201 (2020).

Data availability

491   Raw sequencing data generated for this study was deposited at Gene Expression Omnibus (GEO)

492   under the accession number GSE286182.

Acknowledgments

493   We are grateful to all patients and control subjects who participated in this study. Special thanks to

494   Mei Zhong from the Yale Stem Cell Center genomics core facility who performed sequencing, and

495   Joerg Niklaus from the Yale West Campus Imaging Core for provided microscope training and

496   guidance.

Authors Contribution

NK supervised the study. NK and JCS conceptualized the study and acquired funding. JM

performed the micro-CT. JCS, AB and TSA performed snRNAseq and analysis. TSA, AB and AJ

performed snMultiOme and analysis. TSA performed immunofluorescent microscopy

experiments. JK and AJ performed the PCLS time course experiments. AB, TA and JCS analyzed

the COVID19 scRNAseq data. LDS, BMV, MZO and WAW assembled the IPF cohort and

supervised the biobanking. IR, JCG provided healthy lung tissue. LDS, JD, FN, JCG, BVP, FA,

RJH, IR, XY, BMV, WAW provided critical interpretation, review, and commentary on data and

the manuscript. The manuscript was drafted by TSA, JCS and NK and reviewed and edited by all

authors.

Funding

This project was supported by NIH grants R01HL127349, R01HL141852, U01HL145567 to NK,

and a Three Lakes Foundations grant to NK and IOR; NIH grants R01HL159805, R01DK130294

to PVB; NIH grant R01LM014087 to XY; by the by DOD grant HT9425-23-1-0126 and

Boehringer Ingelheim discovery award grant to JK; DOD grant W81XWH-19-1-0131, the Else

Kröner-Fresenius Foundation (2023_EKCS.18) and the German Center for Lung research (FKZ

82DZL002B1, FKZ 82DZL002C1 & FKZ 82DZLT82C1) to JCS. JCS was supported by

CORE100Pilot Advanced Clinician Scientist Program of Hannover Medical School funded by

Else-Kröner-Fresenius Foundation (EKFS, 2020_EKSP.78) and the Ministry of Science and

Culture of Lower Saxony. BMV and WAW are funded by the KULeuven research fund

(C16/19/005) supporting the biobanking and lung core imaging.

Competing interests

NK reports consulting to Boehringer Ingelheim, Pliant, GSK, Three Lake Partners, Merck, Astra-Zeneca, RohBar, BMS, Galapagos, Chiesi, Sofinnova, Fibrogen, reports Equity in Pliant and grants from Astra Zeneca and BMS. NK has IP on novel biomarkers and therapeutics in IPF licensed to Biotech. JCS served as a consultant to Boehringer Ingelheim, Merck/MSD, GSK, AOP Health, and received lecture honoraria from Boehringer Ingelheim, GSK, Kinevant. JCS has IP on basal cell-targeted therapies in IPF.

Figure Legends and Tables

Figure 1: Study design and epithelial cell atlas of interstitial lung disease.
A) Overview of tissue procurement and experiment design. B) UMAPs of epithelial cells from integrated analysis of multiple datasets. C) Euler diagram of intersecting conserved gene expression markers between normal and disease-associated cell types. D) Heatmap of corresponding features. Each column represents the average normalized gene expression of one patient. Within each dataset, values for each gene are min-max normalized to a common range. E) Boxplots of cell type abundances normalized to the total number of epithelial cells per subject. F) Left: Stacked bar plots of epithelial cell compositions per core in descending rank order of tissue surface density. Right: Linear regression models of changes in epithelial cell proportions as a function of surface density. G) Heatmap of spearman distances among cell-type proportions and surface density, across lung cores.

Figure 2: Continuous changes between healthy and disease alveolar epithelial cells
A) From left to right: UMAPs of epithelial nuclei labelled by cell type, Louvain cluster and disease; PAGA graph of Louvain clusters with edges colored by confidence levels above (black) or below (red) the threshold of 0.2; pruned PAGA graph in Force Atlas 2 (FA2) layout; FA2 layout of nuclei with corresponding edge pruning labelled by Louvain cluster, disease and cell type. B) Upper left: FA2 layout of pseudotime trajectory from ATII to Aberrant Basaloid. Upper right: Generalized additive model fits of genes changing as a function of pseudotime. Lower: heatmap of corresponding genes; each column is one nucleus randomly sampled from fixed intervals in pseudotime, gaps correspond to pseudotime intervals with missing representation. C) FA2 layouts colored by gene expression. D) Pseudotime trajectory from ATI to Aberrant Basaloid, represented in the same manner as Fig 2B.

Figure 3: Immunofluorescent Imaging of AT*i* Cells in IPF

A) CTSE expression in control alveolar epithelial cells. B) CTSE expression in IPF alveolar epithelial cells. C) CPA6 expression in control alveolar epithelial cells. D) CPA6 expression in IPF alveolar epithelial cells.

Figure 4: Longitudinal snRNAseq Assessment of PCLS

A) Overview of experiment design. B) UMAP of snRNAseq data labeled by condition, timepoint and cell type. C) Upper: Stacked bar plots of cell type composition. Middle: Line plot of cell type makeup changes as a function of time. Lower: Line plot of the percent of mitotic cells per cell type as a function of time. D) UMAP of epithelial cells labeled by condition, timepoint and cell subtype. E) Heatmap of cell subtype markers and corresponding percent makeup values. The average normalized gene expression value for each sample and subtype are normalized between 0 and 1 across samples. The corresponding percent makeup value is shown in a bar graph above. F) Heatmap of intersecting hallmarks between *ex vivo* and *in vivo* cells. G) Above: UMAPs of epithelial cells in cell cycle labeled by cell type, condition or timepoint. Middle) Heatmap of mitosis marker genes (above) and cell type marker genes for cells undergoing mitosis in cell cycle (CC) or not undergoing mitosis (nonCC). Below: Line plot of the proportion of cells found in cell cycle for each epithelial cell type.

Figure 5: snMultiOme Analysis of IPF and Control Lung Tissue

A) Overview of analysis strategy. B) UMAP of snRNAseq data labeled by cell type. C) UMAP of corresponding snATAC-seq data colored by cell type. D) Heatmap of epithelial gene expression markers (blue) with elevated chromatin accessibility geneScores (red). Average gene expression per sample and average cell type geneScore values for each gene are independently normalized between 0 and 1. E) Left: The top transcription factor (TF) motifs ranked by global contributions to variance; TFs associated with a particular cell type are highlighted. Right: The top TF motifs enriched in Aberrant Basaloid cells compared to other cell types. F) Above: Sequence logos of the top TF motifs in Aberrant Basaloid cells. Below: Sequence logo for AP-1 motif. G) Euler plot showing intersections of all merged peaks, merged peaks significantly enriched in Aberrant Basaloid cells and merged peaks containing at least one AP-1 motif sequence. H) Density plots of AP-1 enrichment z-scores per cell, organized by cell type. I) Footprint analysis of AP-1 transcription factor occupation across epithelial cell types.