Title: Essentialome-Wide Multigenerational Imaging Reveals Mechanistic Origins of Cell Growth Laws

Authors: Daniel S. Eaton[1][†], Carlos Sánchez[1], Luis Gutiérrez-López[1,4], Jacob Q. Shenker[1], Youlian Goulev[1], Brianna R. Watson[2], Véronique Henriot[1,3], Ethan C. Garner[4], Quincey A. Justman[1], Jeffrey R. Moffitt[2,5,†], Johan Paulsson[1,†]

[1]Department of Systems Biology, Harvard Medical School, Boston, MA 02115, USA
[2]Program in Cellular and Molecular Medicine, Boston Children's Hospital, Boston, MA 02115, USA
[3]Institut Curie, Université PSL, CNRS UMR3348, Orsay, France
[4]Department of Molecular and Cellular Biology, Harvard University, Boston, MA 02138, USA
[5]Department of Microbiology, Harvard Medical School, Boston, MA 02115, USA

[†]To whom correspondence should be addressed: danscotteaton@gmail.com, Jeffrey.Moffitt@childrens.harvard.edu, Johan_Paulsson@hms.harvard.edu

Optical Pooled Screening (OPS) enables precise measurement of pooled genetic libraries in single, highly multiplexed microscopy experiments,[1–4] linking image-based phenotypes to genotypes at genome-scale. However, no method has combined large-scale OPS with multi-generational time-lapse imaging, which is especially important for studying single-cell dynamics and physiology in microbes. Here we develop MARLIN, a technique for multigenerational time-lapse OPS in a microfluidic device and leverage it to map live-cell imaging data to genetic perturbations in a ~130,000-strain mismatch-CRISPRi library targeting all essential genes in *E. coli*. The resulting dataset comprises nearly a hundred million cell cycles and 1.6 billion cells, representing an average of 33 different knockdowns per gene, and provides quantitative measures of morphology and growth for each essential gene with respect to dose. We clustered our multidimensional time-lapse phenotypes to reveal exceptional correlations across co-functional genes, allowing us to systematically discover roles for essential genes of unknown function, as well as implicate RNase E as required for replication initiation. Further, we assayed dose-dependent correlations between growth and cell size for single genes, demonstrating shared regulation of size with ribosome synthesis, while also identifying limitations in existing growth control models. We reconcile these models with our data by positing SpoT as a sensor of translation elongation, which we validate through several lines of genetic and biochemical evidence. Finally, we show that a simple proteome-allocation model – where (p)ppGpp-control only passively limits division protein synthesis – explains all our size-growth scaling behaviors while also quantitatively linking cell size to ribosome abundance. Looking forward, we anticipate that MARLIN will be an essential tool for high-throughput genotype to phenotype mapping in microbes.

Main

Time-lapse microscopy is a powerful tool for cell biology, but is often throughput-limited – finely tracking a few labeled components in a handful of genetic backgrounds, but only managing genome-level screens with great effort[5,6]. Optical Pooled Screening (OPS) promises to change that by leveraging randomly pooled rather than structurally arrayed cell libraries, and then jointly phenotyping and genotyping every cell under the microscope. The randomization reduces batch

artifacts when comparing cells, and, by eliminating the need to keep custody of cells in parallel
compartments, approaches the physical limit of experimental miniaturization where each
individual cell provides a biological replicate. However, current OPS also suffers several
limitations. By pooling cells together, it has been difficult to culture and track them over multiple
generations of growth, and OPS has therefore focused on single snapshots or short-term dynamics
in constant environments. OPS is also challenging[7] for bacteria and yeast, because of their hard-
to-penetrate membranes and low per cell nucleic acid abundance. Microbes would otherwise be
exceptionally well suited for multigenerational OPS, thanks to their small physical footprint, short
generation times and relative ease of building large, pooled libraries.

Here we present an optical screening platform that combines and extends the key benefits of both
arrayed and pooled screens. Specifically, we achieve large-scale OPS for microbes in a
microfluidic cell culture format based on so-called "mother machine"[2,8–18] designs. Cells are
loaded as random pools that self-organize into isogenic lineages by single-file growth in trenches,
while diffusively fed from continuously replenished media that also serves to evacuate surplus
offspring (Fig. 1b). This allows for tracking growth over hundreds of consecutive generations[13,14]
while maintaining exceptionally high cell densities throughout – in local environments that can be
finely controlled in time and space[15,17] – while also minimizing measurement artifacts[15], allowing
for retrieval of cells of interest[16], and many other features[9,18,2]. Early attempts to expand OPS to
this format used Fluorescent In situ Hybridization (FISH) to optically read out RNA barcodes at
the end of the experiment.[2,18] While these were limited to 40 strains per experiment, they laid a
foundation that we expand on here, to consider >120,000 strains per experiment, made possible by
systematic optimizations:

First, we phenotype hundreds of millions of *Escherichia coli* cell divisions per day, organized into
over a million parallel lineages in a dense geometry that enables image acquisition for about
~100,000 lineages per minute (Fig. 1b). The resulting multi-terabyte data volumes are segmented
and analyzed using custom software[19] for parallel computation on HPC systems (Supplementary
Methods).

Second, we link the observed phenotypes to genetically encoded barcodes by optically reading out
the latter under the microscope. Traditional DNA barcodes of sequence length n can in principle
separate between 4^n variants, but in practice less since the number of nucleotide differences
between barcodes (the Hamming distance) must be large compared to the average number of errors
inherited from sequencing. Here we instead use an expression-based strategy with n FISH rounds,
using labeled probes against unique RNA targets and asking if each target is expressed ("1") or
not expressed ("0") in a given cell lineage. By using three spectrally distinct probes
simultaneously, we in principle separate between 8^n variants in n rounds of FISH, but in practice
less since we ensure that the number of different FISH targets between barcodes (here termed the
Herring distance) is much larger than the number of FISH errors. We have heavily optimized this
process, placing our barcodes under strong, inducible expression post-phenotyping, extensively
screening FISH probes for minimal adsorption into the device polymer, and adapting MERFISH
chemistry[1,20,21] to speed up each probe cycle – typically running 10 rounds of three-color imaging
in 14 hours (Methods). To test accuracy, we assayed a pooled library of 120,000 different strains,
each expressing a unique 30-bit barcode together with an intact or inactivated copy of a *gfp* gene,
with fluorescence providing a ground truth for the *gfp* genotype (Extended Data Fig. 2a,b). This
revealed that our FISH barcode read-outs were >99.8% accurate, in a process that is easily
automated and costs around $200 per run (Supplementary Methods).

Finally, we link the barcodes to the genetic regions of interest by sequencing both elements on the
same read. This requires a challenging combination of long-read sequencing (>1kb) and a low
error rate to capture even single point mutations. However, our barcodes help here as well: with
typical Herring distances of 3-4 from the nearest barcode, and single-bit Hamming distances of
~10, the whole-barcode Hamming distances are generally >30, providing a clear barcode
consensus for every read. Grouping the reads by the barcodes allows us to pinpoint other mutations
with great accuracy, e.g. detecting the 2bp substitution distinguishing our *gfp* variants with an
accuracy of 99.8% in as few as 20 grouped reads from an R10.3.1 Oxford Nanopore run, with
higher accuracy likely possible with recent sequencing chemistries (Fig. 1d and Supplementary
Methods).

Taken together, this 'big FISH' platform – MARLIN for Multiplexed Assignment of RNA-
barcoded LINeages – allows us to precisely track over a hundred thousand strains at the cellular
and subcellular levels, each across thousands of individual cells and for dozens of generations of
growth in controllable environments, while precisely calling point mutations in a genetic region of
interest for each cell. We believe that MARLIN can revolutionize screening in microbes, of
anything from perturbation libraries to systematic protein mutants or ensembles of genetic circuits.

Systematic Measurement of Cell Physiology in an Essentialome-wide Knockdown Library

In our first application of MARLIN, we chose to investigate the contribution of essential genes to
cell growth and morphology in *Escherichia coli*. Because loss of essential function leads to
inviable cells, we designed our CRISPRi library to include systematically "mismatched" sgRNAs
with varying complementarity, and therefore activity, on their targets, manifesting as non-lethal
hypomorphic phenotypes.[22–24] We further measure phenotypes from the onset of knockdown to
circumvent the need for long-term cultivation, which also minimizes selection for compensatory
mutations in our uninduced pre-cultures.

Specifically, we designed a CRISPRi library of around 134,000 strains, each with a unique
barcode, covering 29,738 different sgRNAs to target 585 essential genes (Fig. 1e,f). Each barcode
corresponds to an initial library transformant and so each sgRNA is comprised of on average 4-5
separate clonal lineages, mitigating artifacts from random mutations in lineage founders. We then
performed four replicate perturbation experiments, imaging growing lineages for roughly 16
consecutive generations while inducing CRISPRi after 3-4 generations to deplete essential gene
products. Technically we could image many more generations, but 16 was chosen for biological
reasons, to genotype before all cells in a lineage died from essential gene depletion. In total, ~2
million lineages passed quality controls over the four replicates. This aggregate MARLIN dataset
contained at least 200 lineages for 581 of 585 essential gene targets, with ~3,400 lineages observed
on average per essential gene (Fig. 2b and Extended Data Fig. 1g).

To capture the global effects of knocking down essential genes, we extracted six phenotypic
parameters (Fig. 2a and Methods) from each of the 1.6 billion cell detections included in the
MARLIN dataset – cell length (L), width (W), interdivision time (τ), instantaneous single-cell
growth rate (λ), error in division septum placement normalized by cell length (L_s) and pRpsL
fluorescence reporter intensity (I_{rpsL}) as a measure of ribosome synthesis regulation. As a control,
we imaged growing cells in the absence of CRISPRi for over 40 generations, showing that all
measured parameters exhibit stable means and variances over the course of imaging (Extended
Data Fig. 3a).

Following CRISPRi induction, our library variants exhibited great phenotypic diversity in every measured parameter across the library (Fig. 2d and Extended Data Fig. 4), producing a gradual spectrum of physiological impacts at different knockdown strengths (Fig. 2f). Quantities were reproducible on repeated measurements of the same sgRNA across regions of the same microfluidic chip, in replicate experiments (Extended Data Fig. 3d) and in select isolates moved into individual non-barcoded strains (Extended Data Fig. 3e). We also verified that well-known biology is captured by MARLIN across genes. For example, members of the Divisome complex, responsible for carrying out cell division, produced significantly increased cell length upon knockdown (Fig. 2e, blue dots). Characteristic phenotypes were also obtained for depletions targeting the Elongasome, the MinCDE system and the ParCE system (Extended Data Fig. 4b-d).

For a subset of sgRNAs, we further used a DNA marker and higher imaging resolution, to better distinguish division and replication defects. Specifically, we selected one or two sgRNAs per gene with a strong effect on cell physiology and introduced these into a strain bearing a HU-mCherry fluorescent fusion protein which localizes to the nucleoid (Fig. 2g,h). This allowed us to quantify both the ratio of the nucleoid to cell area (NC ratio) as well as the number of nucleoids per cell (Fig. 2i). A reduced NC ratio appears to serve as a proxy for the failure to replicate, as exhibited by a DnaA knockdown (Extended Data Fig. 5a). Further, an elevation in the number of nucleoids indicates division failure, as inhibition of division results in the formation of long, poly-nucleated cell filaments, as seen in FtsN knockdowns (Extended Data Fig. 5b). Together our libraries link comprehensive measurements of physiology and replication status to perturbations for essential function, laying the foundation for mechanistic insights into the molecular basis of these traits.

Clustering Timeseries Phenotypes Reveals Unknown Physiological Functions of Many Genes

Though *E. coli* is arguably the most thoroughly characterized model organism, almost a third of its genes[25] – including up to 20% of essential genes[26]– encode proteins with unknown function. Homology-based approaches have predicted molecular-level properties for some of these proteins, such as enzymatic activity, but rarely identify the cellular process or phenotypes associated with them.

We therefore asked whether disruptions to the expression of functionally related genes tend to group into similar phenotypes that can be identified by clustering. By quantifying single-cell phenotypes dynamically, separating between initial responses to perturbations and secondary epistatic effects, we reasoned that our approach could result in well-defined clusters for highly specific molecular functions. We could then identify specific functions for poorly characterized genes by examining annotations for those genes they co-cluster with.

To that end we performed multiparametric clustering to systematically classify CRISPRi knockdowns according to their measured phenotypes.[2,6,27] To retain information encoded in phenotype dynamics over the course of knockdown, we aggregated measurements for each sgRNA into a multi-dimensional timeseries of Z-scores corresponding to each measured parameter (Fig. 3a and Supplemental Methods). We then generated clusters and a 2-dimensional UMAP based on a distance metric (softDTW)[28] between those timeseries (Fig. 3b). Following this, each cluster was assessed for statistical stability, measured by comparing cluster membership across jackknife resamples of the same data (Extended Data Fig. 6c). Finally, we collapsed redundant information from different sgRNAs targeting the same gene (Fig. 3e), which usually appear in the same cluster. As an initial validation of this approach, we found that members of multi-protein complexes

reproduced phenotypes observed in previous studies and clustered together with their complex
partners (Fig. 3c,d).[29–31]

We focused on a set of 36 clusters with strong phenotypes to build a global map of physiology
under partial loss of essential function. Strikingly, though these clusters were defined agnostically,
based purely on measured phenotypes, we observed an exceptional level of co-clustering for
functionally related genes (Fig. 3c and Extended Data Fig. 6d). Integrating gene association, we
were able to divide these clusters into five functional supergroups: division, replication, envelope,
r-protein and tRNA synthetase. Excluding genes with off-target effects (Supplementary Methods),
100% of Divisome genes were assigned to the division supergroup, 100% of Replisome gene were
assigned to the replication supergroup, and 100% of Elongasome genes were assigned to the
envelope supergroup. Meanwhile, 93% of sgRNAs targeting r-proteins appeared in the r-protein
supergroup and 76% of sgRNAs targeting tRNA synthetases appeared in the tRNA synthetase
supergroup. Each supergroup further contained multiple clusters, and these in turn correlated with
more specific physiological sub-functions within the supergroup. All five supergroups also
contained a minority of other gene depletions, most of which were previously known to be
implicated in the supergroup's associated function, e.g. *minE* which acts as an indirect activator of
division.[31,32] This strongly validates our measurements and analysis, and positions us to annotate
genes of unknown function. Leveraging these phenotypic clusters, we associate 13 poorly
characterized genes with distinct physiological functions, as well as discover a new molecular
factor involved in replication initiation control.

Aside from our multi-gene clusters, five gene depletions in the whole data set did not cluster with
others, but were highly reproducible across redundant sgRNAs targeting the same gene, indicating
gene-unique phenotypes that could be interesting for follow-up analyses (Extended Data Fig. 6e).
Similarly, a few rare exceptions of genes in the same molecular pathways reliably associated with
different clusters (Extended Data Fig. 6d). For instance, knockdowns targeting fatty acid or lipid
A synthesis enzymes appear in a variety of clusters (Extended Data Fig. 6d) and with a diversity
of size scaling behaviors (Extended Data Fig. 7b-d). This suggests that metabolic intermediates in
these pathways possess underappreciated activities distinct from those of end products
(Supplementary Note S1), e.g. as regulatory players.

The r-protein and tRNA synthetase supergroups were each composed of highly interconnected
clusters (Extended Data Fig. 8a,b), with relatively low statistical stability (Extended Data Fig. 6c),
reflecting titration of individual genes which populate a phenotypic continuum across these groups
(Extended Data Fig. 6a). Nonetheless, many clusters were still enriched for more specifically co-
functional genes (Extended Data Fig. 8c), suggesting that finer physiological distinctions could be
made in the future with deeper phenotyping. We also observed that the knockdowns targeting the
tRNA synthetases for histidine and leucine (*hisS* and *leuS*) had unique phenotypes and occupied
single-gene clusters (Extended Data Fig. 6e), demonstrating that single synthetase depletions can
exert physiological consequences that are distinct from the general trend. In addition to tRNA
synthetases (Extended Data Fig. 8b, orange), we found genes involved in amino acid and fatty acid
synthesis, as well as glycolysis, within the tRNA synthetase supergroup. We also found four genes
of unknown function within this group – *ysaA*, *ybcF*, *ylbF*, and *yfjT* (Extended Data Fig. 8d). *ybcF*
and *ylbF* are consecutive genes on a putative catabolic operon,[33] suggesting that this
uncharacterized metabolism contributes significantly to overall growth.

The remaining three supergroups spanning 13 curated clusters were smaller and more statistically
stable, allowing us to further increase the confidence of our functional assignments, by

systematically analyzing sgRNAs for evidence of off-target effects using both fitness and cluster
phenotypes (Supplementary Methods). The envelope synthesis supergroup includes six distinct
and highly stable sub-clusters all involving changes in cell width (Extended Data Fig. 7a, non-
underlined). Based on the genes in each group, and consistent with previous data,[30,34] increased
width was associated with outer membrane (OM) or cell wall defects, while decreased width was
associated with increased OM stiffness (Supplementary Note S1). Most interestingly, factors
implicated in modulating OM composition were found in both wide (*lpxABDH, lptAG, yejM*) and
narrow (*waaC, gmhA, rfaE, ftsH*) clusters, implicating the OM as a major determinant of cell width
homeostasis. Within the envelope synthesis supergroup, the only gene of unknown function *ykgE*
co-clustered with envelope synthesis genes in the high width "Envelope II" cluster shown in
Extended Data Fig. 7a, suggesting a role in maintaining envelope integrity.

The cell division supergroup included three distinct and highly stable sub-clusters, characterized
by an increase in cell length, without any change in growth (Fig. 4a and Extended Data Fig. 7a,
Underlined and Unbolded), consistent with previous findings for partially division-defective
cells.[35] Integrating our nucleoid imaging data, we confirmed that the "Division I" cluster shown in
Fig. 4a exhibited clear division-specific defects, with filamentous, polynucleate phenotypes (Fig.
4a,b). The genes of unknown function among this group (*yagH, ybcN, ycgB* and *yecM*) similarly
exhibit this phenotype (Fig. 4a), implying impaired division but no replication defect. The other
clusters, while division defective, exhibited more complex nucleoid and morphological
phenotypes (Extended Data Fig. 7a). As discussed in Supplementary Note S1, these clusters likely
impair division indirectly through either reduced lipid synthesis ("Division II" cluster) or
dysregulated division localization ("Septum Error" cluster), and include several genes of unknown
function (*yeiH, yfaU*, and *yhcN*).

The DNA replication supergroup contained four distinct sub-clusters (Fig. 4c and Extended Data
Fig. 7a, Underlined and Bold). Like the division-defective clusters, this group exhibited an increase
in cell length, but were instead distinguished by a decreased growth rate, consistent with recent
reports.[36] Remarkably, our replication-associated subclusters were primarily distinguished from
each other by cell width, and each associated with a specific function in replication (Fig. 4c and
Extended Data Fig. 7a, Underlined and Bold). Based on the genes in each of the four clusters, we
successfully associated these groups with either replication initiation, dTTP synthesis or the
Replisome itself. Broadly, members of these clusters are either directly involved in replication, or
indirectly involved through impacts on transcription or transcription termination that have
previously been shown to affect replication (Supplementary Note S1). We found that genes with a
direct role in replication exhibit a decrease in nucleoid extent, consistent with their function, while
those with an indirect role through transcription exhibit an increase in extent, which has indeed
been observed when treating cells with transcription-inhibiting drugs.[37,38] Among these indirect
hits was the only gene of unknown function in this supergroup *yejF*, though its role remains
ambiguous since its sgRNAs are split evenly between replication and envelope-related clusters
(Extended Data Fig. 7a).

Interestingly, our replication initiation-related clusters suggested a previously unappreciated
player in this textbook molecular process. Replication initiation has been extensively characterized
both *in vivo* and *in vitro*, with three core factors, DnaA, DnaB and DnaC, participating in opening
the replication bubble at the origin of replication (*oriC*).[39] Strikingly, we detected one replication
initiation cluster consisting of only these three proteins as well as one other well-studied gene: the
single-strand endoribonuclease RNase E, encoded by *rne*, which is not known to play a role in this

process (Fig. 4c,d). Interestingly, RNase E does not co-cluster with either of the 3' to 5' ssRNA
exonucleases RNase II (*rnb*) and PNPase (*pnp*) (Fig. 4e), indicating that its phenotype does not
result from generic loss of mRNA degrading activity. To test whether RNase E activity is required
for replication initiation, we used the fact that halting initiation reduces ploidy near *oriC*.
Specifically, using a temperature-sensitive RNase E allele, we confirmed that the relative *oriC*
copy number dropped within a few minutes of RNase E inactivation, measured by deep sequencing
(Fig. 4f and Extended Data Fig. 9b), while a replication runout assay (Fig. 4h, *bottom* and Extended
Data Fig. 9c-e) indicated immediate loss of initiation activity. RNase E is thus required for the
initiation of DNA replication *in vivo*, demonstrating that our approach can reveal new essential
functions for even the most well-studied genes involved in the most central processes in *E. coli*.

Growth Control in *E. coli* Depends on Translation Elongation

When cells are perturbed, whether externally by changing conditions or internally by genetic
modifications, their phenotypic responses can follow distinct quantitative patterns. For example,
the classic 'nutrient growth law' of Schaechter, Maaløe, and Kjeldgaard (SMK) shows that cell
size approximately scales exponentially with growth rate for a wide range of external changes in
nutrient levels.[40] Ribosomes in turn obey a famous linear growth law where faster growth produces
proportionally more ribosome expression[41]. Both laws are enforced by the regulatory action of the
small molecule alarmone (p)ppGpp, which represses both size and ribosome synthesis.[42,43] Our
methods above allow us to more comprehensively explore such phenotypic patterns by comparing
growth and size for tens of thousands of internal perturbations arising from graded loss of gene
expression. Strikingly, this reveals three distinct variants of each growth law. Here we describe
these phenotypically, explain how they arise from two distinct translation-mediated mechanisms
for (p)ppGpp control, and propose a unified mechanistic and quantitative explanation for all
variants.

First, many gene knockdowns (65 of 185) with a growth defect (<0.9 doublings/hour) conform to
classical SMK scaling, including tRNA synthetases, central metabolic enzymes and fatty acid
biosynthesis genes (Fig. 5a-d, Extended Data Fig. 10c). This supports our understanding of how
translation affects (p)ppGpp: under conditions of low tRNA charge, there are more elongating
ribosomes with uncharged tRNAs in their A-site,[44,45] where the enzyme RelA can bind to stimulate
its synthesis of (p)ppGpp. In a separately cloned knockdown of the tRNA synthetase PheT, we
used LC-MS/MS to directly confirm that (p)ppGpp indeed was strongly elevated (Fig. 6b) and
measured the RNA/Protein ratio to confirm that ribosomes were correspondingly downregulated
(Fig. 5f).

Second, 90 genes in our dataset – including 41 of the 45 strongly growth-inhibiting ribosomal
protein (r-protein) knockdowns (Fig. 5c,d, purple bars) – instead *inverted* the SMK growth law
and produced negative scaling between cell size and growth rate. This seems to support the same
tRNA charge explanation as above: knockdowns of individual r-proteins often compromise
ribosome elongation[46] and decrease tRNA consumption, which increases tRNA charge while
reducing RelA activity and (p)ppGpp levels. We also observed the accompanying inversion in
ribosome scaling with increased RNA/Protein at lower growth rates (Fig. 6a), and found that
ectopic expression of a hyperactive (p)ppGpp synthase (RelA*) largely restores size and ribosome
content to wildtype levels (Fig. 6a and Extended Data Fig. 5g). However, we then identified a third
scaling class where knockdowns targeting the few remaining r-proteins, located in the ribosome
stalk, as well as translation initiators *infA* and *infB*, form a group that exhibit flat scaling with *no*

discernable correlation between size and growth (Fig. 5d, Extended Data Fig. 10b). Both (p)ppGpp and ribosome levels in an *infA* knockdown are indeed similar to wildtype (Fig. 5f).

This is an important discrepancy: by our expectations, knocking down initiation factors should have mimicked r-protein knockdowns and similarly reduced the consumption of charged tRNA. The fact that there is a distinct difference suggests that the inverse SMK scaling for r-protein knockdowns is *not* in fact due to altered tRNA charging in general, but to a more specific step. We therefore analyzed knockdowns of the EF-G protein – specifically targeting EF-G-mediated ribosome translocation. The results closely mirrored the effects of knocking down r-proteins: decreased growth, decreased (p)ppGpp, increased size and an increased RNA/Protein ratio (Fig. 5e,f, 6b)[47–49]. This further shows that initiation and translocation differentially affect (p)ppGpp levels, suggesting additional control mechanisms beyond RelA. Next we explore those mechanisms.

Ribosomes and Cell Size Respond to Translation Elongation Defects via SpoT and (p)ppGpp

A recent mathematical analysis[47] of how (p)ppGpp levels quantitatively scale with elongation rate suggested that translating ribosomes play a direct regulatory role, not just in (p)ppGpp production via RelA, but also by making the (p)ppGpp degradation rate proportional to the number of ribosomes carrying a charged A-site tRNA i.e. that are primed to translocate. Such a mechanism would diminish (p)ppGpp in response to reductions in translocation rate, but will likely not respond to initiation defects – which should only weakly impact the translocation rate in rich conditions i.e. where this rate is already maximized – as we observed both for the libraries and in the targeted experiments focused specifically on pure initiation and translocation knockdowns. This prompted us to look for such a translocation-specific regulator.

The only known (p)ppGpp regulator in *E. coli*, apart from RelA, is the bifunctional enzyme SpoT which both constitutively synthesizes and degrades (p)ppGpp[50]. We therefore hypothesized that SpoT provides the translocation-specific (p)ppGpp hydrolysis responsible for the inverse SMK scaling under inhibited translation. To test this, we sought to compare CRISPRi libraries with and without *spoT*. While the single *ΔspoT* knockout does not grow, due to the stabilization and accumulation of lethal amounts of (p)ppGpp synthesized by RelA[51], the *spoT* deletion can be made in a *ΔrelA* strain. We therefore compared growth and size patterns across three CRISPRi libraries – the WT above as well as both the double knock-out *ΔrelA ΔspoT* and the single knockout *ΔrelA*.

In the *ΔrelA ΔspoT* background, which have no (p)ppGpp[15], both the SMK scaling for tRNA synthetase knockdowns and the inverse SMK scaling for r-protein knockdowns collapse to flat lines, like the initiation factor knockdowns. This confirms that the scaling laws depend on (p)ppGpp. We then considered the single *ΔrelA* knockout where cells have the SpoT activity added back, which should restore both its constitutive (p)ppGpp production activity[51] and its putative translocation-dependent hydrolysis. Strikingly, we find that adding back SpoT completely restores the inverse SMK scaling under r-protein knockdown observed in the wildtype (Fig. 6d). We also confirm that this response is specific to translocation defects, by demonstrating similar effects for an EF-G knockdown in *ΔrelA* strains which produce a wildtype-like increase in ribosome levels and reduction in (p)ppGpp (Fig. 6e). Similar results hold for initiation defects, where again as in wildtype cells, size and ribosome levels are unaffected in a *ΔrelA* background (Fig. 6d,e).

In contrast to wildtype, however, in the *ΔrelA* strain the tRNA synthetase knockdowns *do not* follow SMK scaling, but rather *inverse* SMK scaling (Fig. 6d). Similarly, an inverted (p)ppGpp response to single amino acid auxotrophy has been observed previously[50], which has not been

accounted for thus far. This, we argue, further supports a SpoT-dependent translocation response,
as these knockdowns are predicted to impede translocation due to ribosome queuing upstream of
codons with undercharged cognate tRNAs (see Supplementary Theory). Consistent with this
explanation, knockdowns which inhibit the synthesis of multiple amino-acids, and should thus not
produce excessive queueing, do not exhibit this inverse scaling in *ΔrelA* (Fig. 6d). Therefore, RelA
appears to mask what would otherwise be an inverted, SpoT-mediated response to single amino
acid depletion in WT – showing that our proposed mechanism also clarifies previously unintuitive
results. These data show that *spoT* is necessary for the reduction of (p)ppGpp under reduced
translocation, while *relA* is required for standard SMK scaling under reduced tRNA synthetase
activity.

Finally, we show that a combined RelA/SpoT model explains the remarkable linearity of
ribosome-growth scaling, under varying nutrient quality, elongation inhibitors and translation
initiation defects (Fig. 5f).[41,52,53] Previously, linear ribosome-growth scaling was theoretically
suggested to arise as a consequence of maximizing overall translation flux[53] through control of
ribosomes. We show that even without maximizing translation flux, the same behavior also
emerges from the proposed (p)ppGpp control model – assuming simple, first-order kinetics in
(p)ppGpp production and degradation. Such a model produces positive-linear ribosome scaling
under reduced tRNA charge (Fig. 6c, orange line) – previously derived elsewhere[47], negative-
linear scaling under reduced elongation (Fig. 6c, purple line) and invariance under reduced
translation initiation (Fig. 6c, red line) (see Theory Box).

Briefly, the previous derivation[47] of positive-linear ribosome scaling under reduced tRNA charge
follows from the inverse scaling of the (p)ppGpp concentration g with the elongation rate ε under
these conditions. In particular, $g \propto (\varepsilon_{Max}/\varepsilon) - 1$ where ε_{Max} is the maximum elongation rate, which
was subsequently empirically validated. Continuing, the authors assume (p)ppGpp control of
ribosome levels R is given by a simple inverse function $R \propto 1/g$, while the growth rate λ itself
depends on the product of ribosome levels with the elongation rate $\lambda = \varepsilon R$. To capture the
accumulation of inactive "hibernating" ribosomes H at high g, the authors inserted a corrective
term – resulting in $\lambda = \varepsilon(R-H)$. The measured scaling of hibernating ribosomes with (p)ppGpp[47]
subsequently motivated the form $H \propto g$. Solving this system of equations for $\lambda(R)$ results in a
complex form. However, under the simplifying, empirically verified assumption that the minimum
elongation rate ε_0 is half ε_{Max}, the authors obtained the observed positive-linear scaling $\lambda \propto - R_0$
$+R$.

Following a similar approach, we show that linear model behavior can also be obtained for the
reduced translocation and initiation regimes. Beginning with reduced translocation, we note that
for this perturbation, the (p)ppGpp concentration g instead scales positive-linearly with ε. Here,
we model (p)ppGpp control of ribosome levels R by a simple monovalent repressor function $R \propto$
$k/(k+g)$, while $\lambda = \varepsilon(R-H)$ still holds. However, we instead treat hibernating ribosomes as a fixed
value H_0, motivated by the observed invariance in this value except at low growth rates.[54] We
subsequently note that H_0 and the minimum ribosome expression level R_0 take similar values –
allowing us to use the simplifying assumption $H_0 = R_0$. Along with the proportionality between
elongation rates and (p)ppGpp levels, this allows us to obtain $g \propto \lambda/(R-R_0)$. Finally, we may
rearrange the ribosome repressor function to get $g \propto (R_{max}-R)/(R-R_0)$. Since both ribosome levels
and growth rate are scaled to (p)ppGpp levels by the same inverse dependence on $R-R_0$, it follows
that growth rate and ribosome levels obey a negative-linear relationship $\lambda \propto R_{max} - R$ as observed.[52]

The remaining invariant scaling under initiation factor depletion (Fig. 6c, red line) proceeds simply
from the expectation that neither dwelling nor translocation times will change. That is, under the
rich growth conditions examined here, ternary complex diffusion should be limiting for the
dwelling step[55] and increases in tRNA charge from lower translation flux should not impact
dwelling. Similarly, the translocation rate is likely near-maximum under these conditions[54] and so
increases from i.e. reduced ribosome queueing are unlikely. As such, ribosome levels should
remain fixed with respect to growth under translation initiation defects, as we observe (Fig. 5f).
Thus, our model for the combined RelA-SpoT system largely accounts for the remarkably robust
linearity between ribosomes and growth across diverse translation perturbations.

The Nutrient Growth Law is Negative Hyperbolic and Emerges from Optimal Protein Synthesis
408
409

Above we showed how translation-mediated (p)ppGpp control creates the linear scaling of
ribosomes with growth rate, for all three classes of perturbations observed. Next we seek to explain
the nutrient size growth law. Because cell volume responds to growth in a (p)ppGpp-dependent
manner, and (p)ppGpp directly controls transcription of many genes, it is tempting to hypothesize
that some key division protein is under direct transcriptional control of (p)ppGpp. However, to our
knowledge no such candidates have been identified, and most known division proteins are instead
demonstratively *not* regulated by (p)ppGpp.[56] Here we propose that not only the classical SMK
growth law, where volume seems to depend exponentially on growth, but all three variants
observed above, precisely reflect this *absence* of direct (p)ppGpp control of division proteins.

Specifically, our model postulates that cells on average divide after accumulating enough of some
unspecified set of 'division proteins' that are not directly controlled by (p)ppGpp, whether
structurally involved in division or in some indirect regulatory process. This assumption is
supported by the observation that most division-related promoters are indeed not regulated by
(p)ppGpp[56,57] and is consistent both with *E. coli*'s behavior as a division 'adder'[58,10] where cells
correct size deviations regressively by adding constant mass regardless of deviations in birth size,
and the fact that changes in the expression of division proteins correlate with changes in cell size.[59]
In the simplest model, the average volume V of cells is then inversely proportional to the fraction
or 'sector' ϕ_{NC} of the proteome not controlled by (p)ppGpp, which in turn is complementary to the
fraction $\phi_C = 1 - \phi_{NC}$ that *is* controlled by (p)ppGpp, i.e., $V \propto 1/(1-\phi_C)$.

We further postulate that the sector ϕ_C controlled by (p)ppGpp varies linearly with the ribosome
sector, i.e., $\phi_C = a_1\phi_R + b_1$. This is roughly true simply because ribosomes constitute a large part
of this sector. However, linearity also follows exactly if (p)ppGpp controls all (p)ppGpp-regulated
promoters by the same factor, which seems likely since it regulates expression by binding RNA
polymerase far from the DNA binding domains, minimizing promoter-specific effects (see Theory
Box). We experimentally validate that, across many growth conditions, the (p)ppGpp-repressed
fraction of expression varies linearly with the ribosomal protein fraction, at both the protein and
mRNA levels (Extended Data Fig. 11).[57,60,61] This predicts a volume-ribosome relationship of the
form $V \propto (1-c\phi_R)^{-1}$ where c is a positive constant, as also directly supported by our measurements
across different knockdowns (Fig. 7e, white dots). Finally, we connect the ribosome sector ϕ_R to
growth rate λ by using the linear ribosome growth law above, $\phi_R = a_2\lambda + b_2$ with different slopes
under defects for tRNA charging, translocation and initiation, a form we and others have
experimentally verified (Fig. 5f, data pending for *fusA*).[53]

These simple and experimentally supported assumptions predict a negative *hyperbolic* relationship between volume V and growth rate λ, i.e., on the form $V \sim (1 - k\lambda)^{-1}$ rather than the classic exponential $V \sim e^{k\lambda}$. For the SMK perturbation class, the hyperbolic curve strikingly fits data at least as well as the classical exponential curve, even though the exponential was chosen based on fitting (Fig. 7c) and the hyperbolic is predicted from underlying mechanisms. In fact, the hyperbolic function quantitatively captures all three perturbation classes (Fig. 7d), with class-specific *size scaling* constants strictly proportional to their underlying *ribosome scaling* constants, consistent with our mechanistic expectations (Extended Data Fig. 12b). Together, our evidence shows that, despite describing different phenotypes and taking different quantitative forms, both the nutrient and ribosome growth laws arise from (p)ppGpp-mediated regulation of ribosome synthesis.

For maximal generality we refrain from proposing a specific model for the inverse proportionality between division volume and division protein levels. However, we do note that the assembly of the division-mediating Z-ring, which requires sufficient division protein to span the perimeter of the mid-cell cross-section, could mediate such a proportionality. Indeed, we observe that division volume exhibits a positive, roughly linear dependence on the mid-cell perimeter for the knockdowns that alter cell width (Extended Data Fig. 12a), and that applying a perimeter-based correction to our size-growth scaling model also results in a high-quality fit (Extended Data Fig. 12c).

It has been a long-standing mystery why so many bacteria follow the nutrient growth law of approximately exponential size-growth scaling.[10,40,62] Our results show that the exponential is similar to the hyperbolic, which arises naturally from simple criteria that could be satisfied in many organisms. Specifically, the linear relation between ribosome levels and growth rate, the complementary linear decrease in protein sectors not co-regulated with ribosomes, and the lack of strong co-regulation between division and ribosomes, may all hold broadly. Many other cell processes could similarly respond to changes in conditions in a seemingly controlled and coordinated manner, simply by *not* being directly regulated by (p)ppGpp or its equivalents. In fact, properties that stay *constant* across conditions are the ones more likely to be under (p)ppGpp control. For example, the initiation volume per origin of replication in *E. coli* changes vary little between environments, requiring levels of the key regulator DnaA to be maintained at a constant level. Indeed DnaA is repressed by (p)ppGpp.[57]

Evolutionarily, all these quantitative patterns, from division 'adders' to nutrient or ribosome growth laws, could just be spandrels – byproducts of selection on other traits, such as maximal ribosome efficiency. However, cell size matters for many reasons, including the surface-to-volume ratio needed to support nutrient intake, stochastic low-number effects, or transport rates across the cell. If other size-growth patterns would be beneficial, they could also easily be achieved by e.g. adding or removing (p)ppGpp control elements to a few genes. We therefore view the simple passive control above as an elegant and highly efficient way for cells to ensure baseline size adaptation to different conditions.

Conclusion

Our MARLIN approach combines multi-generational imaging of bacteria in stable, micro-controlled environments followed by precise genotyping of every cell – at a scale of close to a billion cells per day, across a million separate lineages each of which can in principle be a separate

strain. In doing so, MARLIN combines the comprehensiveness of genomic screening with the
quantitative measurements demanded by biophysics and physiology.

Looking forward, combining MARLIN with deep mutational scanning of RNAs or proteins
enables far deeper and more precise phenotypic readouts for library variants than possible with
e.g. sort-seq, including single-cell heterogeneity, dynamics and extremely subtle differences in
expression. For many biomolecules, such as DNA-binding proteins like activators and repressors,
precise quantification and dynamics are essential to understanding the full sequence-function
landscape. MARLIN allows us to consider millions of DNA-binding protein variants,
systematically connecting binding kinetics to amino acid substitutions.

Expanding out into gene networks, we have seen that MARLIN is uniquely suited to characterizing
the quantitative rules emerging from circuits coordinating fundamental cellular processes.
Extending this with process-specific reporters for division[63], replication[64] and various subcellular
structures[65,66] will uncover additional rules and their molecular implementation. Moreover,
MARLIN is also useful for studying synthetic cellular circuits, allowing us to consider not only
the individual effects of simple building blocks like promoters and ribosome binding sites, but also
far more complex emergent dynamics of networks. Behaviors like bistability, oscillations and
excitable dynamics can now be deeply analyzed not just for a handful of systems, but for hundreds
of thousands of circuits.

At the broadest cell scale, MARLIN's combination of immense throughput with phenotypic depth
is greatly enabling for physiology, where gene networks interact with each other, biophysical
constraints, and the environment to govern global cellular behaviors. By testing hundreds of
thousands of variants, we remain agnostic to mechanisms and allow our data to guide holistic,
parsimonious physiological models. Moreover, expanding our clustering approach with deeper
phenotyping – using more imaging modalities, more reporters and more conditions – we may
further functionally annotate the ~1,200 genes[67] that still have unknown functions in *E. coli*.
Finally, using MARLIN to pool commonly-held structured libraries – such as the ASKA, Keio,
and various GFP reporter libraries – would enable their routine measurement in combination with
different genetic disruptions, reporters and extracellular environments, multiplying our
understanding of how major cellular systems interlock. The low cost of running these experiments,
combined with the ease of building, sharding and storing pooled libraries, further promises to
democratize access.

We believe MARLIN will serve as a tremendous quantitative tool for understanding both the
internal mechanics of microbes as well as the synthetic systems we engineer into them. The
measurements enabled by MARLIN, coupled with the recent explosion of computational
approaches for genomics and model identification, represent a huge step towards the goal of
modeling emergent cellular behavior from the cell's constituent parts.

Theory Box

(p)ppGpp Control of Ribosomes

It has previously been shown that ribosome concentration shows positive-linear scaling with growth rate, under varying nutrient conditions and negative-linear scaling under varying concentrations of the elongation-targeting antibiotic chloramphenicol.[52] In addition, our data suggests that ribosome levels do not vary under growth defects incurred by inhibiting translation initiation. Here, we show how all three of these behaviors may be derived from the (p)ppGpp regulatory model discussed in the main text.

Under this model, (p)ppGpp synthesis is hypothesized to occur in proportion to the concentration of uncharged ribosomes R_U competent to bind an uncharged tRNA and stimulate RelA, while (p)ppGpp degradation occurs in proportion to the concentration of charged ribosomes R_C with a charged A-site tRNA awaiting translation. The steady state concentration of (p)ppGpp g is thus the ratio of these concentrations, scaled by a constant k equal to the ratio of rate constants associated with synthesis and degradation:

$$g = k \cdot \frac{R_U}{R_C} = k \cdot \frac{\tau_D}{\tau_T}$$

During elongation, ribosomes switch from the uncharged to charged state with an average dwell time τ_D, followed by switching back into the uncharged state once again after an average translocation time τ_T. At equilibrium, elongating ribosomes obey detailed balance and satisfy the condition $R_U \tau_D^{-1} = R_C \tau_T^{-1}$, meaning that we may freely substitute ribosome concentrations with the kinetic rates of dwelling and translocation; this form of g will thus allow us to connect (p)ppGpp levels to translation kinetics.

Previously, the behavior of this model was analyzed under variable dwelling time, which was associated with conditions of varying nutrient quality and tRNA charging.[47] The form of g under variable dwell time was given as a function of the overall translation elongation rate $\varepsilon = (\tau_D + \tau_T)^{-1}$, with the maximum elongation rate $\varepsilon_{max} = \tau_T^{-1}$ associated with the limit $\tau_D \rightarrow 0$:

$$g_{\text{Dwell}}(R, \varepsilon) = k \cdot \left(\frac{\varepsilon_{max}}{\varepsilon} - 1 \right)$$

We extend this model to varying initiation and translocation rates by considering model behavior when tRNA charging is already high – i.e. under a reference condition of rapid growth. Under these conditions, dwell time will be primarily limited by the diffusion of ternary complexes[55] and only weakly dependent on tRNA charge. Therefore, despite the expected increase in tRNA charge under initiation defects, the resulting change in dwell time will be small while the translocation time will be unaffected. As such, (p)ppGpp levels would be expected to remain constant under initiation knockdown:

$$g_{\text{Init}}(R, \varepsilon) = g_{\text{ref}}$$

Finally, modeling translocation defects by varying τ_T under a fixed dwelling time τ_D^0, we arrive at the following relationship between (p)ppGpp levels and the elongation rate:

$$g_{\text{Elong}}(R, \varepsilon) = c \cdot \varepsilon \tau_D^0$$

Previously, it was shown that the relationship between ribosome levels R and growth rate λ, under variable dwell time (i.e. obeying g_{Dwell}) could be obtained by satisfying a system of equations including the (p)ppGpp repressive function on ribosomes $R(g)$ and the dependence of growth rate on the ribosome concentration and elongation rates $\lambda(R, \varepsilon)$. We adopt a similar approach here to model $R(\lambda)$ under variable initiation and translocation. Motivated by the empirical scaling of (p)ppGpp with ribosome levels[48] we assume it acts as a simple monovalent repressor of ribosome synthesis:

$$R(g) = \frac{k}{k + g} \cdot (R_{max} - R_0) + R_0$$

569 The growth rate is proportional to the product of the number of active ribosomes $(R - R_{inact})$ engaged in
570 translation and the elongation rate:

$$\lambda(g) = a \cdot \varepsilon(g) \cdot (R(g) - R_{inact})$$

572 In the simplifying case where $R_{inact} = R_0$ – which appears to be satisfied over most of the range of growth
573 rates examined – solving this system of equations yields $R(\lambda)$ under both varying initiation and
574 translocation. Combining this with the previously derived form under varying dwelling time results in the
575 following three functions:

$$R_{Dwell}(\lambda) = R_0 + \varepsilon_{max}^{-1}\lambda$$
$$R_{Init}(\lambda) = R_{ref}$$
$$R_{Elong}(\lambda) = R_{max} - \frac{c\tau_D}{ak}\lambda$$

576
577
578

Scaling of Unregulated Proteins

We hypothesize that the set of proteins whose activities are limiting for division are only passively regulated
by (p)ppGpp, influenced primarily through competition with (p)ppGpp-regulated transcripts for the
translation machinery. To formalize how (p)ppGpp regulation will impact this set of proteins and modulate
size, we use a conventional proteome allocation framework.[52,68] In this framework, the cell's total proteome
is partitioned into functionally distinct "sectors," each occupying a fraction ϕ_i of the total proteome. Each
such sector is further divided into a basal ϕ_i^0 and an additional $\Delta\phi_i$ allocation capturing any changes from
this reference. We will specify the following two coarse-grained proteome sectors: a (p)ppGpp repressed
sector ϕ_C and an unregulated sector ϕ_{NC} – which we hypothesize contains the set of limiting division
proteins assigned to a sub-sector ϕ_D. Due to the finite size of the proteome, additional ϕ_C will come at the
expense of ϕ_{NC} with each sharing a fixed 'flexible' portion of the proteome $\Delta\phi^{max}$ according to $\Delta\phi_C +$
$\Delta\phi_{NC} = \Delta\phi^{max}$.

To link the fraction of unregulated proteins ϕ_{NC} directly to the growth rate, we first introduce the following
relationship between the ribosome fraction ϕ_R – which depends on growth as described previously – and
the overall fraction of genes repressed by (p)ppGpp ϕ_C:

$$\Delta\phi_C(\lambda) = f_R^{-1}\Delta\phi_R(\lambda)$$

595 Or equivalently:

$$\phi_C(\lambda) = f_R^{-1}\Delta\phi_R(\lambda) + \phi_C^0 = a_1\phi_R(\lambda) + b_1$$

In other words, we assume that the change in the ϕ_C sector is linearly proportional to the change in the ϕ_R
sector, according to a constant f_R^{-1}. We may obtain this condition if we assume that (p)ppGpp regulated
genes are controlled according to functions of the same form as the ribosome repression function $R(g)$
introduced earlier – with the (p)ppGpp binding constant k held constant i.e. assuming (p)ppGpp affinity for
RNAP is not strongly impacted by promoter sequence. Combining this result with the constraint of a finite
proteome produces the following dependence of ϕ_{NC} on growth rate:

$$\Delta\phi_{NC}(\lambda) = \Delta\phi^{max} - f_R^{-1}\Delta\phi_R(\lambda)$$

604 We may specify $\Delta\phi_R(\lambda)$ according to the linear relationship of ribosome concentration and growth rate
605 $R(\lambda)$ under tRNA charge depletion and elongation inhibition, described in the previous section:

$$\phi_{NC}^{Dwell}(\lambda) = \phi_{NC}^0 + \Delta\phi^{max} - f_R^{-1}\beta_{tRNA} \cdot \lambda$$
$$\phi_{NC}^{Init}(\lambda) = \phi_{NC}^0 + \Delta\phi^{max} - f_R^{-1}\Delta\phi_R^{Ref}$$
$$\phi_{NC}^{Elong}(\lambda) = \phi_{NC}^0 - f_R^{-1}\beta_{Elong} \cdot \lambda$$

606
607
608

609 where each β refers to the slope of the corresponding ribosome fraction-growth relationship $\phi_R(\lambda)$. In
610 general, all three equations correspond to a linear form:

611
$$\phi_{NC}^{Class}(\lambda) = a_2^{Class} \cdot \lambda + b_2^{Class}$$

612 with perturbation-specific slopes and intercepts.

Negative Hyperbolic Size-Growth Scaling

614 Finally, to link the abundance of division proteins to cell size, we will need a rough model for cell division.
615 To this end, we will require that the division volume V_D scales inversely with the concentration of division
616 proteins ϕ_D – which being a subset of the unregulated sector ϕ_{NC} is equivalent to:

617
$$V_D \propto 1/\phi_{NC} = 1/(1 - \phi_C)$$

Note that this criterion is generally obtained for the division accumulator family of size control models –
while replication-driven models exhibit a more complex dependence of V_D on the abundance of the
replication-initiating protein and the growth rate. This condition will hold for the average cell volume V,
which is proportional to V_D for cells growing at steady state – with the exact proportionality set by the
conditions of observation e.g. a sample from liquid culture vs trapped lineages (Supplementary Theory).

We integrate our proteome model with this size-concentration relation – substituting $\Delta\phi_C$ with $f_R^{-1}\Delta\phi_R$ –
to arrive at the following negative hyperbolic function describing the explicit dependence of cell size on
ribosome concentration:

626
$$\frac{V}{V_0} = \left(1 - \theta \cdot \frac{\Delta\phi_R}{\phi_R^{max}}\right)^{-1} \qquad \theta = \frac{f_R^{-1}\phi_R^{max}}{\phi_{NC}^{max}} \qquad \phi_i^{max} = \phi_i^0 + \Delta\phi^{max}$$

which is equivalent to the relation $V \propto (1-c\phi_R)^{-1}$ from the main text. Qualitatively, the parameter θ serves
to convert a given percentage change in the ribosome sector, normalized to its maximum size, into an
equivalent percentage change in the unregulated sector. This is then converted into a change in volume,
with the normalization condition that the minimum volume is achieved when the unregulated sector is
maximized (ϕ_{NC}^{max}). Applying the condition-dependent forms of $\Delta\phi_R(\lambda)$ from the previous section to this
equation, we arrive at a distinct linear ribosome-growth relationship for each translation perturbation class:

633
$$\frac{V_{Dwell}}{V_0}(\lambda) = \left(1 - \theta \cdot \frac{\beta_{Dwell}}{\phi_R^{max}} \cdot \lambda\right)^{-1} = (1 - \kappa_{Dwell} \cdot \lambda)^{-1}$$

634
$$\frac{V_{Init}}{V_0}(\lambda) = \left(1 - \theta \cdot \frac{\Delta\phi_R^{Ref}}{\phi_R^{max}}\right)^{-1} = \frac{V_{Dwell}}{V_0}(\lambda^{Ref})$$

635
$$\frac{V_{Elong}}{V_0}(\lambda) = \left(\frac{\phi_{NC}^0}{\phi_{NC}^{max}} - \theta \cdot \frac{\beta_{Elong}}{\phi_R^{max}} \cdot \lambda\right)^{-1} = \left(\alpha_{Elong} - \kappa_{Elong} \cdot \lambda\right)^{-1}$$

636 Examining the specific case of tRNA synthetase knockdowns – which exhibit positive growth-size scalings
637 of the SMK type – we can see directly how this classic law emerges naturally from our integrated model:

638
$$V_{Dwell} = V_0 exp(\kappa_{Dwell} \cdot \lambda) exp\left(\frac{1}{2}(\kappa_{Dwell} \cdot \lambda)^2 + O(\lambda^3)\right)$$

639 In other words, our model form can be approximated as exponential in growth rate when the change in $\Delta\phi_R$
640 over conditions is small (e.g. $\kappa_{Dwell} \cdot \lambda$ is small). From this we also see that the deviation from an equivalent
641 exponential fit will be second order in the growth rate, reflecting an increased relative curvature in the
642 negative hyperbolic model.

Fig. 1: MARLIN - Multiplexed Assignment of RNA-barcoded LINeages. a, MARLIN libraries are constructed through combinatorial ligation of MERFISH barcodes to genetic variants and subsequently bottlenecked to ~10^5 CFUs. During an experiment, libraries are loaded as a pool and cultivated on our high-throughput mother machine. **b,** In a single imaging run, our device enables continuous imaging of >1 million bacterial lineages every 10 minutes, enabling high precision measurement of growth dynamics at scale. **c,** As an endpoint, cells are fixed, and barcodes read using FISH. Dye signal is subsequently eliminated by a disulfide reduction allowing multiple cycles of sequential probing. Decoded barcode sequences are mapped back to their associated library members' time-lapse data using a sequencing-established lookup table. **d,** The barcode-library variant lookup table is established using nanopore sequencing of our pooled plasmid library. Reads are robustly grouped by FISH barcode in the presence of sequencing-associated error, which is subsequently corrected by consensus averaging for each group. **e,** Our library is designed according to a mismatch-CRISPRi strategy where gene expression is titrated between 0 and 10 mismatches between sgRNA and target. We included mismatch series for up to 5 PAM sites per gene, depending on the number available in the gene. **f,** Our essentialome-wide CRISPRi screen was designed targeting 585 essential genes in *E. coli* with 29738 sgRNAs, derived a previous set of matched sgRNAs[26], which was barcoded and introduced into MG1655 *E. coli* containing a genomically integrated pRpsL-mKate2Hyb fluorescent reporter.

Fig. 2: Essentialome-wide Mismatch-CRISPRi with Cell Cycle Measurements. a, Top, our CRISPRi library was monitored for over 2 hours of normal growth followed by 8 hours of CRISPRi induction, before being genotyped. Bottom, kymograph of an example lineage from our library, arrows show identified division events. Top-left zoom-in shows that we may quantify length (L), width (W) and pRpsL reporter intensity (I_{rpsL}) from single snapshots. Bottom-right zoom-in shows our terminal or "mother" cell growing over time, with its cell volume over time indicated on the graph above. From these data, we may measure both the instantaneous growth rate (λ) and the interdivision time (τ), as well as the absolute error in septum placement (L_S). **b,** Histograms of the number of lineages or cell cycles observed per gene target across all library replicates. **c,** Top, single-cell length distribution for sgRNA variants targeting the indicated genes, between 5-8 hours of CRISPRi induction. Bottom, bootstrap estimator distributions for the length averaged within single lineages (left) and the median of these average lengths, across trenches containing the same sgRNA (right). **d,** Top, log histograms of sgRNA variant average phenotypes, between 5-8 hours of CRISPRi induction. Black line indicates the mean of control sgRNA mean phenotypes. Bottom, representative kymographs of lineages containing sgRNA variants with the indicated mean phenotypes. **e,**

Volcano plot of cell lengths measured between 5-8 hours post-induction, for each sgRNA in the library. The horizontal line corresponds to FDR=0.05 and the vertical lines indicate two standard deviations around the distribution of control sgRNAs. Sample lineages are shown of representative and control strains, after 8 hours of CRISPRi induction. **f,** Length vs growth rate plot of a single sgRNA targeting ValS over time. Inset indicates the growth-length scaling of representative ValS sgRNAs after 8 hours of CRISPRi induction. **g,** Length vs growth rate plots for sgRNAs targeting the indicated gene after 8 hours of CRISPRi induction. The white circle indicates the mean length and growth rate of control strains, while the orange dots indicate sgRNAs used in the HU-mCherry reporter library. **h,** Left, length vs growth rate plot for sgRNAs targeting DnaB after 8 hours of CRISPRi induction. A strong sgRNA is selected from the set of DnaB-targeting sgRNAs and pooled with representative sgRNAs targeting all other genes in the library. Right, these sgRNAs were introduced into a strain bearing chromosomally integrated HU-mCherry and pRpsL-mVenus markers. **i,** Left, kymograph of an example lineage from our HU-mCherry library. Top-left zoom-in shows that we may quantify the nucleoid to cell area ratio (NC) and the number of nucleoids (N_n) in each cell. Right, joint distributions of cell and nucleoid area for single cells with 1 or 2 nucleoids, prior to CRISPRi induction.

Fig. 3: Clustering Reveals Phenotypic Signatures of Essential Processes. a, Left, each parameter, measured over time, is converted to a 20-point timeseries using gaussian kernel regression on the data from each trench. These timeseries are converted to Z-scores (Methods) and averaged over trenches belonging to the same sgRNA variant to produce an average timeseries. Right, we subsequently filter out variants with weak phenotypes by excluding those which do not exhibit a Z-score > 1 for any timepoint, along any dimension (grey area). **b,** Conversion of Z-score cutoffs to real values shows where the cutoffs for length and growth rate lie for exemplary genes with size and growth phenotypes. Distances between timeseries data for sgRNA variants are established using a softDTW distance metric, which produces a sgRNA vs sgRNA distance matrix that is then used to generate a 2-dimensional UMAP and clusters based on the Leiden clustering algorithm. **c,** Top, red dots correspond to sgRNAs targeting genes belonging to the indicated class, mapped onto a UMAP representation of each sgRNA variant. Bottom, endpoint values of Length, Width, and Septum Error for each sgRNA. Blue represents values below those of control sgRNAs and red represents values above the controls, with scale given by the right-side bar. **d,** Representative lineage kymographs from our HU-mCherry library, from the indicated classes in Fig. 3c. **e,** Left, subset of clusters associated with a division/replication-like defect, indicated by color. Heatmap indicates average phenotypic quantities after 8 hours of CRISPRi induction for each sgRNA, scale in Fig. 3f. Right, sgRNA

variants composing each of our clusters of interest are aggregated into single genes by taking the average of each phenotypic value. Single genes are assigned to the cluster in which the plurality of their targeting sgRNAs can be found, with two clusters assigned in the case of ties. **f,** Top, schematic of quantified variables, see Fig. 2 for details. Bottom, linear scale for variables in Fig. 3e, with red indicating quantities above the EV controls while blue indicates quantities below the controls.

Fig. 4: Clusters Implicate Unknown Genes in Division, Replication and Envelope Integrity. a, Heatmap of cluster phenotypes for the "Division I" cluster. Arrows indicate genes mentioned in the text. The leftmost column color denotes cluster membership, right columns display the average value of each quantity at the end of the experiment, for each gene. Same scale as in Fig. 3e. Entries beginning with O denote polycistronic operons, where the subsequent gene name indicates the last gene in a consecutive block all belonging to the indicated cluster. See Extended Data Fig. 7 for all curated clusters. **b,** Representative kymographs of genes from the "Division I" cluster, with the cytoplasm in magenta and nucleoid in green, as in the diagram in Fig. 3f. Arrows indicate clear manifestations of the cluster-associated phenotype. Left, kymograph of a *ftsN* knockdown, canonically associated with division. Right, kymograph of a *yagH* knockdown, not previously characterized as division-defective. **c,** Heatmap of cluster phenotypes for the "Replication Initiation I" cluster, same as in Fig. 4a. **d,** Representative kymographs of genes from the "Replication Initiation I" cluster, same as in Fig. 4c. Left, kymograph of a *dnaA* knockdown, canonically associated with replication initiation. Right, kymograph of a *rne* knockdown, not previously characterized as initiation-defective. **e,** Top, colored dots correspond to sgRNAs targeting the indicated genes involved in RNA degradation, mapped onto a UMAP representation of each sgRNA variant. Bottom, heatmap of gene phenotypes for *rnb* and *pnp*. **f,** Marker frequencies based on deep gRNA sequencing for 100kb bins across the *E. coli* genome, with the reference coordinate set to *oriC*. Frequencies normalized to reference measurements from non-replicating cells grown to stationary phase in MBM+0.2% Succinate. Measurements taken after shifting either rne^{WT} (black) or *rne-1(ts)* (red) cells from 30 °C to 43 °C for 2 hours, with shaded area indicating $\pm 2\sigma_{SEM}$ based on three biological replicates. **g,** CFU/mL over time for

rne^{WT} (triangles) and *rne-1(ts)* (circles) cells during a shift from 30 °C to 43 °C occurring at 80 minutes (red
shading). The vertical dotted line indicates 30 minutes post-shift, the time at which post-treatment samples
were taken for replication runout, dividing 1-2 times in this period. **h,** Flow cytometric measurement of
PicoGreen-stained DNA before (left) and after (right) replication runout for cells grown first at 30 °C and
shifted to 43 °C for 30 minutes, in strains bearing either *rne^{WT}* (black) or *rne-1(ts)* (red) alleles. *rne-1(ts)*
cells drop from 16 to 4-8 oriC copies after 30 minutes at 43 °C, indicating that initiation immediately halts
on temperature shift and the original 16 oriC have simply partitioned into either 2 or 4 descendent cells.

787
788

Fig. 5: Three Size-Growth Scalings Emerge from Translation Knockdowns. a, Length vs growth rate
for each sgRNA variant averaged between 5-8 hours after CRISPRi induction. Dashed line a guide for the
log-linear decrease in length with growth rate. **b,** Top, select $\log_2$ length vs growth rate linear fits to groups
of sgRNAs targeting single genes. Bottom, histogram of the $\log_2$ length vs growth rate slope for genes
exhibiting at least one sgRNA with a growth rate less than 0.9 doublings per hour. Colors correspond to
slopes less than -0.2 (purple), greater than 0.2 (orange) and between -0.2 and 0.2 (red). **c,** Histograms of
$\log_2$ length vs growth rate slope for indicated gene groups. **d,** Length vs growth rate averaged between 5
and 8 hours after CRISPRi induction for variants corresponding to ribosomal proteins/rRNAs (purple
circles), tRNA synthetases (orange circles), ribosome stalk proteins L7/L12 and L10 (black outlined purple
circles) and initiation factors (black outlined red circles). Initiation factor InfC was omitted due to likely
disruption of downstream gene expression. Note that ribosome stalk knockdowns exhibit exceptionally
weak size scaling compared to other r-proteins (see Fig. 6c). Triangles correspond to points for which
individual isolates were cloned in E, including 50S subunit proteins RplQ and RplA (purple), the 30S
subunit protein RpsO (pink) and an EV control sgRNA (blue). **e,** Left, simplified kinetic model of
translation, consisting of initiation, elongation and termination. Elongation kinetics are coarse-grained into
an uncharged state (empty A-site) and a charged state (charged tRNA in the A-site). Colors indicate gene
functions targeted in the corresponding knockdown strains. Right, small circles indicate average length vs
average exponential growth rate for the indicated genes after 5-8 hours of CRISPRi induction (pheT, infA,
rplL) in lDE20. Inverted triangles indicate the same values for a strain with FusA under AHL induction
(P_{AHL}-fusA), 5-8 hours after stepping down from an AHL concentration of 1 μM to 0.2 μM, 0.16 μM, 0.13
μM, 0.1 μM, 0.07 μM, 0.05 μM, or 0.03 μM. Large circles represent the same values for indicated CRISPRi
strains (DE428, DE830, DE831, and DE832) induced by aTc or the AHL induction strain (DE828) stepped
down to 0 μM AHL. **f,** Average RNA/Protein ratio (n=5-6) after 6 hours of CRISPRi induction or removal
of AHL vs average exponential growth rate from Fig. 5e. Large circles indicate the same strains and
knockdown conditions as in the large circles in Fig. 5f and Fig. 6b. Error bars denote $2\sigma_{SEM}$ range.

Fig. 6: (p)ppGpp Degradation by Translocating Ribosomes Reconciles Diverse Translation Phenotypes. a, RNA/Protein ratio after 6 hours CRISPRi depletion of indicated genes. WT denotes the EV sgRNA control. Unhatched bars indicate cells uninduced for the hyperactive (p)ppGpp synthase RelA* and hatched lines indicate +25uM IPTG to induce RelA* expression, resulting in ectopic synthesis of (p)ppGpp. Error bars denote $2\sigma_{SEM}$ range. Bars indicated with "*" exhibit a significant increase in RNA/Protein relative to WT ($p<0.05$). Dotted line indicates the maximum RNA/Protein ratio (~0.89) reported in a previous study limiting protein synthesis by chloramphenicol.[69] **b,** Left, same schematic as in Fig. 5e. Right, RNA/Protein and (p)ppGpp measurements of the indicated sgRNA knockdowns, P_{AHL}-fusA and an EV sgRNA control (WT), in a $relA^+ spoT^+$ background. All measurements made after 6 hours of CRISPRi induction or removal of AHL. Error bars denote $2\sigma_{SEM}$ range. **c,** Left, illustration of a two-step model of elongation. Production of (p)ppGpp in proportion to uncharged or "dwelling" ribosomes (R_D) is mediated by RelA, as previously described.[47] Degradation of (p)ppGpp promoted by charged or "translocating" ribosomes (R_T), in a SpoT-dependent manner, is proposed to reconcile our observations. Constitutive (p)ppGpp production by SpoT is assumed insignificant in WT cells, but considered as the only productive term in $\Delta relA\ spoT^+$. Right, model-predicted changes in (p)ppGpp, cell length, or RNA/Protein vs growth rate under increased dwell time (orange), increased translocation time (purple) or reduced translation initiation (red). Note the prediction for cell length is qualitative, since the (p)ppGpp to length function is not modeled. The model also accounts for RNA/Protein and size invariance under $rplL$ knockdown, which interferes with initiation,

translocation and charged tRNA capture (Supplementary Theory).[70] **d,** Average length vs average
exponential growth rate for translation-related variants after 5-8 hours of CRISPRi induction in $relA^+ spoT^+$,
$\Delta relA\ spoT^+$, $\Delta relA\ \Delta spoT$ backgrounds. Ribosomal proteins are purple, tRNA synthetases are orange,
glycolysis enzymes are blue, and initiation factors are red. **e,** RNA/Protein and (p)ppGpp measurements of
the same knockdowns as in Fig. 6a, in a $\Delta relA\ spoT^+$ background. Empty circles indicate values in $relA^+$
$spoT^+$.

891
Fig. 7: Size Scaling Law Emerges from Proteome Distortion by Ribosome Expression. a, Illustration
of our size scaling model. The proteome is divided into two coarse-grained sectors, ϕ_P containing proteins
repressed by (p)ppGpp and ϕ_{NP} containing proteins that are not directly regulated by (p)ppGpp. ϕ_P obeys
the same regulation as ribosomal proteins and therefore the proteome fraction made up of ribosomes ϕ_R is
a subset of this fraction. Division is assumed to occur when a subset of proteins in ϕ_{NP}, with a fractional
abundance of ϕ_D, reach a threshold abundance D*. **b,** Model-derived scaling of the ribosome proteome
fraction ϕ_R (left) and cell volume (right) with growth rate, under decreasing translocation rate (purple),
initiation rate (red) and ternary complex capture rate (orange). **c,** Similar measurements as in Fig. 5e, for
pheT knockdowns, with cell volume estimated from length and width assuming the cell has a
spherocylindrical shape. Lines correspond to fits to the negative hyperbolic and exponential (i.e. SMK) size
models. **d,** Same measurements as in Fig. 7d, for *pheT* (orange), *infA* (red) and *fusA*/EF-G (purple)
knockdowns. Lines correspond to a three parameter, joint fit of the negative hyperbolic to *pheT* and *fusA*
knockdowns, with the trendline for *infA* given by the point of intersection. 95% confidence intervals were
determined by residual bootstrapping. **e,** Datapoints correspond to the empirical RNA/Protein vs volume
relationship, derived from the data in Figs. 5e-f. Shaded area corresponds to the 95% confidence interval
of the model-predicted relationship, given the fit in Fig. 7e.

Sequencing

Imaging

Extended Data Fig. 1: Sequencing Error Rates and Sampling Distributions for MARLIN Experiments. a, Error rate (per bp) in the random 15-bp UMI vs read depth, for synthetically down-sampled data. lDE11 (small *gfp* variant library) variants with read depth >200 were sequenced and their consensus UMIs taken as ground truth. Reads assigned to these variants were randomly downsampled to a fixed read depth, re-run through our consensus calling algorithm and compared to the high depth ground truth to estimate the error rate. **b,** Error rate in calling GFPmut2 vs DarkGFP vs read depth, for synthetically down-sampled data generated as in A. **c,** Log scale histogram of the read depth distribution for each lDE15 (large *gfp* variant library) barcode, with a depth cut-off of 15 reads. Grey bars are below the depth threshold and do not contribute to the codebook, while blue bars above the threshold do. **d,** Bar chart showing the number of sequenced barcodes assigned to the GFPmut2 and DarkGFP variants. **e,** Log scale histogram of the read depth distribution for each lDE20 (mismatch-CRISPRi library) barcode, with a depth cut-off of 15 reads. **f,** Histogram of the number of MARLIN barcodes associated with each lDE20 sgRNA sequence. **g,** Histograms of the number of lineages measured for each designed sgRNA in lDE20. **h,** Same histogram for lDE26, lDE28 and lDE30 (HU-mCherry libraries). **i,** Histogram of the number of sgRNAs measured per gene target, for sgRNAs passing the quality threshold (>7 lineages, $CV_{SEM} < 0.2$) for inclusion in our growth-length scaling analysis (Fig. 3-4).

Extended Data Fig. 2: Barcode Error Rate Measurement. a, A barcoded library (lDE15) of two
fluorescent protein variants, GFP (i.e. GFPmut2) and DarkGFP, was measured by MARLIN to estimate
our barcode error rates. **b,** During imaging of lDE15, we measured the fluorescent signal from both a
constitutively expressed fluorescent protein encoded on the chromosome (pRpsL-mKate2Hyb) and the
constitutively expressed GFP variant expressed on the lDE15 plasmid, every 15 minutes for 6 hours. **c,**
Histogram of the average GFP/mKate2 intensity ratio for each lineage in the experiment. We called the
lineage as being GFP positive if this ratio was above 1 (dotted line). **d,** Decomposition of total experimental
error into contributions from library sequencing and MERFISH readout. Note that the FISH contribution is
halved since barcode assignment errors will predict an incorrect GFP variant roughly half the time. **e,**
Estimating error contributions by increasing error robustness of barcodes. Computationally filtering lDE15
data for sub-libraries composed of well-separated barcodes with hamming distances of up to four should
eliminate the FISH contribution to error. The marginal difference of ~0.09 indicates that the FISH error
contribution is ~0.18%. See Supplementary Methods for error estimation details.

Extended Data Fig. 3: Phenotypic Quantities in MARLIN-CRISPRi Experiments are Highly Reproducible. a, Mean and standard deviation estimates for physiological parameters over time for 26 hours of growth of DE32. Values are computed for each 2-hour time window. **b,** Kernel density estimates for distributions of measured physiological parameters without correction (blue line), with trench-wise baseline rescaling (orange line) and with both rescaling and filtering based on growth characteristics (green line). See Supplementary Methods for details. **c,** Mean values of physiological parameters within each FOV across the chip using uncorrected or rescaled and filtered values. **d,** Top, relationship between cell length, width or mKate2 intensity, for each sgRNA in replicate lDE20 experiments. Points correspond to average values in the period from 5 to 8 hours post-induction, for the same sgRNA in each experiment. Bottom, relationship between cell length, width or mKate2 intensity, for each replicate sgRNA in the top or bottom half of the microfluidic. **e,** Distributions of measured physiological parameters in the period from 5 to 8 hours post-induction, for the indicated CRISPRi knockdowns. Distributions correspond to either values determined from the library experiment (right, grey) or from corresponding clonal isolates (left, green). Dotted lines indicate the quartiles of the distributions.

Extended Data Fig. 4: Volcano Plots of 6 Physiological Quantities After CRISPRi Knockdown. a,
Sample lineages of EV control strains, after 8 hours of CRISPRi induction in our mismatch-CRISPRi
library (lDE20). **b,** Volcano plot of cell lengths measured between 5- and 8-hours post-induction, for each
sgRNA in lDE20. The horizontal line corresponds to FDR=0.05 and the vertical lines indicate three standard
deviations above and below the distribution of control sgRNAs. Sample lineages are shown of
representative strains, imaging the cytoplasmic pRpsL-mKate2Hyb marker. **c-g,** Similar volcano plots also
included for cell width (**c**), septum placement error (**d**), growth rate (**e**), interdivision time (**f**), and mKate2
intensity (**g**). Consistent with previous reports, depletions targeting members of the cell wall synthesizing
Elongasome (*mrdAB*, *mreBCD*, *rodZ*) exhibit an increase in width of >0.1 μm relative to EV controls.[30,71,72]
Also similar to previous reports[31,32], depletions targeting the division localization factor *minC* and nucleoid
partitioning *parCE* system exhibit an increase in septum placement error of >10% relative to EV controls.

Extended Data Fig. 5: Nucleoid Compaction is Associated with Translation Elongation Defects, without RelA Activation. a, Volcano plot of the nucleoid to cell area (NC) ratio, measured between 5 and 8 hours post-induction, for each sgRNA in our HU-mCherry library (lDE26). The horizontal line corresponds to FDR=0.05 and the vertical lines indicate two standard deviations around the distribution of control sgRNAs. Sample lineages are shown of representative strains, where indicated. The cytoplasmic pRpsL-mVenus marker is shown in magenta and the nucleoid-localized HU-mCherry marker is shown in green. **b,** Volcano plot of the number of nucleoids in the cell. **c,** Relationship of NC ratio to growth rate, for all sgRNAs in lDE26. Indicated dots are genes coding for tRNA synthetases (tRNA Synth), translation initiation factors (Initiation), the 50S ribosomal subunit (50S) or the 30S ribosomal subunit (30S). **d,** Same measurement as in panel c, for each sgRNA in our *ΔrelA* library (lDE28). **e,** Same measurement as in panel c, for each sgRNA in our *ΔrelA ΔspoT* library (lDE30). **f,** Same measurement as in panel c. Indicated dots are genes coding for the Sec translocation complex (Sec) or the signal recognition particle (SRP). **g,** Cell length and NC ratio, as measured by DAPI staining, in CRISPRi strains depleted for the indicated gene products, WT denoting the EV sgRNA control. Top, snapshots of the indicated isolates imaged by phase contrast (greyscale) and for fluorescent DAPI stain (blue). *ppGpp^{WT}* indicates cells uninduced for the

hyperactive (p)ppGpp synthase RelA* and $ppGpp^{++}$ indicates cell cultured with +25uM IPTG to induce
RelA* expression, resulting in ectopic synthesis of (p)ppGpp. Bottom, length and NC ratio measurements
of the same strains. Error bars denote $2\sigma_{SEM}$ range. **h,** Same measurements as panel g. *ΔrelA* indicates the
knockdown was performed in a *ΔrelA* background, while $relA^{+}$ indicates an intact locus. pAHL-fusA
denotes the AHL induction strain introduced in Fig. 3e, in which *fusA* was depleted according to the same
protocol as in Fig. 3f. Note the difference between this strain and *fusA*, which likely owes to disrupted
expression of the downstream *tufA* gene.

1000

1001

1002

1003

1004

1005

1006

1007

1008

1009

1010

1011

1012

1013

1014

1015

1016

1017

Extended Data Fig. 6: Clustering and Phenotype Distributions in MARLIN-CRISPRi Library. **a,** Distribution of measured parameters in our mismatch-CRISPRi library (lDE20). Colors on UMAPs indicate whether sgRNA variants on the UMAP take an average parameter value above (red), below (blue) or at the value of the sgRNA EV controls (white). **b,** Clusters determined by the Leiden community detection algorithm[73], indicated by color, with a resolution of 2. **c,** Cluster robustness at different resolutions, determined from 10% jackknife resampling of Leiden clusters and expressed as either the average Jaccard score or the average recall score between a cluster and the most similar cluster in each resampling. **d,** Distribution of different gene groups across lDE20 UMAP, indicated in red. **e,** Top, distribution of different genes or operons across lDE20 UMAP, indicated in red. Points outlined in black correspond to Leiden clusters containing primarily the indicated gene or operon. The percentage corresponds to the fraction of sgRNAs targeting the indicated gene/operon that appear in the selected cluster. Bottom, the average phenotype of the indicated cluster, using the same scaling as in panel a.

1031
Extended Data Fig. 7: Extended Cluster Heatmaps and Select Scaling Phenotypes for Genes Involved
in Fatty Acid, Phospholipid and LPS Synthesis. **a,** Heatmap of cluster phenotypes. Underlined cluster
names indicate the division supergroup, bold names indicate the replication supergroup, and plain names
indicate the envelope supergroup. Arrows indicate genes mentioned in the text. The leftmost column color
denotes cluster membership, right columns display the average value of each quantity at the end of the
experiment, for each gene. Same scale as in Fig. 5e. Entries beginning with O denote polycistronic operons,
where the subsequent gene name indicates the last gene in a consecutive block all belonging to the indicated
cluster. **b-c,** Semi-log plots of average growth rate vs length, measured between 5- and 8-hours post-
induction, for sgRNAs in our mismatch-CRISPRi library (lDE20) targeting the indicated genes (red)
involved in phospholipid and fatty acid synthesis (grey). **d,** Semi-log plots of average growth rate vs width,
measured between 5- and 8-hours post-induction, for sgRNAs in our mismatch-CRISPRi library (lDE20)
targeting the indicated genes (red) involved in LPS synthesis (grey). **e.** Representative kymographs of *minC*
and *lolE* knockdowns, with the cytoplasm in magenta and nucleoid in green, as in the diagram in Fig. 5f.
Top, knockdown, arrows indicating asymmetric divisions. Bottom, kymograph of a *lolE* knockdown,

arrows indicating extracellular HU-mCherry resulting from lysis. Insets highlight instances of cytoplasmic
invagination and plasmolysis.

1048

1049

1050

1051

1052

1053

1054

1055

1056

1057

1058

1059

1060

1061

1062

1063

1064

1065

1066

1067

1068

1069

1070

1071

1072

Extended Data Fig. 8: Ribosomal Protein and tRNA Synthetase Clusters are Highly Interconnected.
a, Left, heatmap indicates the percentage of each cluster composed of either ribosomal proteins (purple) or
tRNA synthetases (orange), with each row corresponding to a cluster in our mismatch-CRISPRi library
(lDE20). Right, heatmap values indicate statistical connectivity between clusters using the PAGA
connectivity metric[74]. Rows are matched to the same clusters as in the left heatmap. Boxed cluster groups
indicated highly inter-connected groups enriched for either ribosomal proteins (purple) or tRNA synthetases
(orange). **b,** Highlighted UMAP indicating the inter-connected cluster groups corresponding to the boxes
in panel a, with the same color scheme. **c,** Distribution of different gene groups across lDE20 UMAP,
indicated in red. **d,** Distribution of different genes of unknown function across lDE20 UMAP, indicated in
red.

1083

1084

1085

1086

1087

1088

Extended Data Fig. 9: *rne-1(ts)* Cells Rapidly Lose Initiation Activity at a Restrictive Temperature.

a, Representative lineages depleted for *rne* after 8 hours of CRISPRi induction in our HU-mCherry library

(lDE26). The cytoplasmic pRpsL-mVenus marker is shown in magenta and the nucleoid-localized HU-

mCherry marker is shown in green. **b,** Marker frequencies based on deep gRNA sequencing for 100kb bins

across the *E. coli* genome, with the reference coordinate set to *oriC*. Frequencies normalized to reference

measurements from non-replicating cells grown to stationary phase in MBM+0.2% Succinate at 37 °C.

Measurements before (left) and after (right) shifting either *rne^WT* (black) or *rne-1(ts)* (red) cells from 30 °C

to 43 °C for 2 hours, with shaded area indicating ±2σSEM based on three biological replicates. **c,** Triplicate

flow cytometric measurement of PicoGreen-stained DNA after replication runout for *rne^WT* cells grown to

stationary phase in MBM+0.2% Succinate at 37 °C, used to determine fluorescence intensity of a single

chromosome. Chromosome equivalents reported as various multiples (2,4,8,16) of this value. **d,** Same as

panel c, for *rne^WT* cells before (0 mins) and at various times after a shift from 30 °C to 43 °C. **e,** Same as

panel d, for *rne-1(ts)* cells.

1102

1103

1104

1105

1106

Extended Data Fig. 10: Growth-Size Scaling Analysis for Translation-Related Knockdowns. **a,** Left,
histograms of log2 length vs growth rate slope for indicated gene groups including translation termination
and initiation factors. Same color scheme for slopes as in Fig. 3b. Right, select log2 length vs growth rate
linear fits to groups of sgRNAs targeting single genes belonging to the same gene class. **b,** Same as in panel
a, with indicated gene groups including tRNA synthetases and lipid synthesis factors. **c,** Same as in panel
a, with indicated gene groups including ribosomal proteins and ribosome biogenesis factors.

1113

1114

1115

1116

1117

1118

1119

1120

1121

1122

1123

1124

1125

Extended Data Fig. 11: Ribosomal Protein Expression Varies Linearly with the (p)ppGpp Regulon.
a, Measured ribosomal protein transcriptome fraction vs (p)ppGpp-repressed transcriptome fraction, under
limitation of glucose import by PtsG titration (orange points) or translation inhibition by sub-lethal
chloramphenicol (blue points). Transcriptome data from Balakrishnan and Mori et al. 2022.[61] List of
(p)ppGpp repressed promoters from Sanchez-Vazquez et al. 2019.[57] **b,** Measured ribosomal proteome
fraction vs (p)ppGpp-repressed proteome fraction, with the same conditions as in panel a. The ribosome
fraction of the variable component of the (p)ppGpp regulon f_R – defined in the Theory Box – is measured
as the slope of this relationship. Proteome data from Mori et al. 2021.[60] **c-d,** Same plots as panels a-b, but
for genes belonging to proteome sectors – identified in Mori et al. 2021[60] – which are co-regulated with
ribosomal proteins (i.e. the A,S,U,C sectors), which are collectively referred to as the (p)ppGpp-repressed
or "P-sector".

1137

1138

1139

1140

1141
Extended Data Fig. 12: Strict Proportionality of Ribosome and Volume Scaling Constants. **a,** Single-
cell measurements of division volume vs mid-cell perimeter at division, between 6-8 hours post-induction
in lDE26. Points are only shown for library variants targeting members of the Rod complex (*mrdA*, *mrdB*,
mreB, mreC, mreD, rodZ) or UDP-GlcNAc synthesis (*glmS*, *glmM*, *glmU*), which primarily impact cell
width through reduced cell wall synthesis. **b,** Measured ribosome-growth scaling constants β vs volume-
growth scaling constants κ, for *pheT* (orange), *infA* (red), *rplL* (light purple) and *fusA* (dark purple)
knockdown series. Derived from scaling data in Fig. 5e,f. See Theory Box for definitions of β and κ. **c,**
Same fit of volume vs growth rate as in Fig. 7d (left) as well as equivalent fits where volume is normalized
by the measured mid-cell perimeter (center) or the mid-cell cross section (right).

1151

1152

1153

1154

1155

1156

1157

References:

1. Emanuel, G., Moffitt, J. R. & Zhuang, X. High-throughput, image-based screening of pooled genetic-variant libraries. *Nat. Methods* **14**, 1159–1162 (2017).

2. Camsund, D. *et al.* Time-resolved imaging-based CRISPRi screening. *Nat. Methods* **17**, 86–92 (2020).

3. Feldman, D. *et al.* Optical Pooled Screens in Human Cells. *Cell* **179**, 787-799.e17 (2019).

4. Funk, L. *et al.* The phenotypic landscape of essential human genes. *Cell* **185**, 4634-4653.e22 (2022).

5. Juillot, D. *et al.* A High-Content Microscopy Screening Identifies New Genes Involved in Cell Width Control in Bacillus subtilis. *mSystems* **6**, e01017-21 (2021).

6. Campos, M. *et al.* Genomewide phenotypic analysis of growth, cell morphogenesis, and cell cycle events in *Escherichia coli*. *Mol. Syst. Biol.* **14**, e7573 (2018).

7. Daims, H., Stoecker, K. & Wagner, M. Fluorescence in situ hybridization for the detection of prokaryotes. in *Molecular Microbial Ecology* (Taylor & Francis, 2005).

8. Wang, P. *et al.* Robust Growth of *Escherichia coli*. *Curr. Biol.* **20**, 1099–1103 (2010).

9. Baltekin, Ö., Boucharin, A., Tano, E., Andersson, D. I. & Elf, J. Antibiotic susceptibility testing in less than 30 min using direct single-cell imaging. *Proc. Natl. Acad. Sci.* **114**, 9170–9175 (2017).

10. Taheri-Araghi, S. *et al.* Cell-Size Control and Homeostasis in Bacteria. *Curr. Biol.* **25**, 385–391 (2015).

11. Si, F. *et al.* Mechanistic Origin of Cell-Size Control and Homeostasis in Bacteria. *Curr. Biol.* **29**, 1760-1770.e7 (2019).

12. Potvin-Trottier, L., Lord, N. D., Vinnicombe, G. & Paulsson, J. Synchronous long-term oscillations in a synthetic gene circuit. *Nature* **538**, 514–517 (2016).

13. Norman, T. M., Lord, N. D., Paulsson, J. & Losick, R. Memory and modularity in cell-fate decision making. *Nature* **503**, 481–486 (2013).

14. Lord, N. D. *et al.* Stochastic antagonism between two proteins governs a bacterial cell fate switch. *Science* **366**, 116–120 (2019).

15. Bakshi, S. *et al.* Tracking bacterial lineages in complex and dynamic environments with applications

for growth control and persistence. *Nat. Microbiol.* **6**, 783–791 (2021).

16. Luro, S., Potvin-Trottier, L., Okumus, B. & Paulsson, J. Isolating live cells after high-throughput, long-

term, time-lapse microscopy. *Nat. Methods* **17**, 93–100 (2020).

17. Kaiser, M. *et al.* Monitoring single-cell gene regulation under dynamically controllable conditions

with integrated microfluidics and software. *Nat. Commun.* **9**, 212 (2018).

18. Lawson, M. J. *et al.* In situ genotyping of a pooled strain library after characterizing complex

phenotypes. *Mol. Syst. Biol.* **13**, 947 (2017).

19. paulssonlab/TrenchRipper. Paulsson Lab at HMS (2024).

20. Chen, K. H., Boettiger, A. N., Moffitt, J. R., Wang, S. & Zhuang, X. Spatially resolved, highly

multiplexed RNA profiling in single cells. *Science* **348**, aaa6090 (2015).

21. Moffitt, J. R. *et al.* High-throughput single-cell gene-expression profiling with multiplexed error-

robust fluorescence in situ hybridization. *Proc. Natl. Acad. Sci.* **113**, 11046–11051 (2016).

22. Vigouroux, A., Oldewurtel, E., Cui, L., Bikard, D. & van Teeffelen, S. Tuning dCas9's ability to block

transcription enables robust, noiseless knockdown of bacterial genes. *Mol. Syst. Biol.* **14**, e7899

(2018).

23. Hawkins, J. S. *et al.* Mismatch-CRISPRi Reveals the Co-varying Expression-Fitness Relationships of

Essential Genes in Escherichia coli and Bacillus subtilis. *Cell Syst.* **11**, 523-535.e9 (2020).

24. Mathis, A. D., Otto, R. M. & Reynolds, K. A. A simplified strategy for titrating gene expression reveals

new relationships between genotype, environment, and bacterial growth. *Nucleic Acids Res.* **49**, e6

(2021).

25. Goodall, E. C. A. *et al.* The Essential Genome of Escherichia coli K-12. *mBio* **9**, e02096-17 (2018).

26. Calvo-Villamañán, A. *et al.* On-target activity predictions enable improved CRISPR–dCas9 screens in

bacteria. *Nucleic Acids Res.* **48**, e64 (2020).

27. Govers, S. K., Campos, M., Tyagi, B., Laloux, G. & Jacobs-Wagner, C. Apparent simplicity and

emergent robustness in bacterial cell cycle control. 2023.01.16.524295 Preprint at

https://doi.org/10.1101/2023.01.16.524295 (2023).

28. Cuturi, M. & Blondel, M. Soft-DTW: a Differentiable Loss Function for Time-Series. Preprint at

https://doi.org/10.48550/arXiv.1703.01541 (2018).

29. Yang, X. *et al.* A two-track model for the spatiotemporal coordination of bacterial septal cell wall

synthesis revealed by single-molecule imaging of FtsW. *Nat. Microbiol.* **6**, 584–593 (2021).

30. Kruse, T., Bork-Jensen, J. & Gerdes, K. The morphogenetic MreBCD proteins of Escherichia coli form

an essential membrane-bound complex. *Mol. Microbiol.* **55**, 78–89 (2005).

31. Kato, J. *et al.* New topoisomerase essential for chromosome segregation in E. coli. *Cell* **63**, 393–404

(1990).

32. Boer, P. A. J. de, Crossley, R. E. & Rothfield, L. I. A division inhibitor and a topological specificity

factor coded for by the minicell locus determine proper placement of the division septum in E. coli.

Cell **56**, 641–649 (1989).

33. Smith, A. A. T., Belda, E., Viari, A., Medigue, C. & Vallenet, D. The CanOE Strategy: Integrating

Genomic and Metabolic Contexts across Multiple Prokaryote Genomes to Find Candidate Genes for

Orphan Enzymes. *PLOS Comput. Biol.* **8**, e1002540 (2012).

34. Fivenson, E. M. *et al.* A role for the Gram-negative outer membrane in bacterial shape

determination. *Proc. Natl. Acad. Sci.* **120**, e2301987120 (2023).

35. Zheng, H. *et al.* Interrogating the Escherichia coli cell cycle by cell dimension perturbations. *Proc.*

Natl. Acad. Sci. **113**, 15000–15005 (2016).

36. Mäkelä, J. *et al.* Genome concentration limits cell growth and modulates proteome composition in

Escherichia coli. *eLife* **13**, (2024).

37. Dworsky, P. & Schaechter, M. Effect of Rifampin on the Structure and Membrane Attachment of the

Nucleoid of Escherichia coli. *J. Bacteriol.* **116**, 1364–1374 (1973).

38. Bakshi, S., Choi, H., Mondal, J. & Weisshaar, J. C. Time-dependent effects of transcription- and

translation-halting drugs on the spatial distributions of the Escherichia coli chromosome and

ribosomes. *Mol. Microbiol.* **94**, 871–887 (2014).

39. Mott, M. L. & Berger, J. M. DNA replication initiation: mechanisms and regulation in bacteria. *Nat.*

Rev. Microbiol. **5**, 343–354 (2007).

40. Schaechter, M., MaalØe, O. & Kjeldgaard, N. O. Dependency on Medium and Temperature of Cell

Size and Chemical Composition during Balanced Growth of Salmonella typhimurium. *Microbiology*

19, 592–606 (1958).

41. Neidhardt, F. C. & Magasanik, B. Studies on the role of ribonucleic acid in the growth of bacteria.

Biochim. Biophys. Acta **42**, 99–116 (1960).

42. Potrykus, K., Murphy, H., Philippe, N. & Cashel, M. ppGpp is the major source of growth rate control

in E. coli. *Environ. Microbiol.* **13**, 563–575 (2011).

43. Fernández-Coll, L. *et al.* The Absence of (p)ppGpp Renders Initiation of Escherichia coli

Chromosomal DNA Synthesis Independent of Growth Rates. *mBio* **11**, 10.1128/mbio.03223-19

(2020).

44. Haseltine, W. A. & Block, R. Synthesis of Guanosine Tetra- and Pentaphosphate Requires the

Presence of a Codon-Specific, Uncharged Transfer Ribonucleic Acid in the Acceptor Site of

Ribosomes. *Proc. Natl. Acad. Sci.* **70**, 1564–1568 (1973).

45. Wendrich, T. M., Blaha, G., Wilson, D. N., Marahiel, M. A. & Nierhaus, K. H. Dissection of the

Mechanism for the Stringent Factor RelA. *Mol. Cell* **10**, 779–788 (2002).

46. Subramanian, A. R. & Dabbs, E. R. Functional Studies on Ribosomes Lacking Protein L1 from Mutant

Escherichia coli. *Eur. J. Biochem.* **112**, 425–430 (1980).

47. Wu, C. *et al.* Cellular perception of growth rate and the mechanistic origin of bacterial growth law.

Proc. Natl. Acad. Sci. **119**, e2201585119 (2022).

48. Ryals, J., Little, R. & Bremer, H. Control of rRNA and tRNA syntheses in Escherichia coli by guanosine

tetraphosphate. *J. Bacteriol.* **151**, 1261–1268 (1982).

49. Chure, G. & Cremer, J. An optimal regulation of fluxes dictates microbial growth in and out of steady

state. *eLife* **12**, e84878 (2023).

50. Murray, D. K. & Bremer, H. Control ofspoT-dependent ppGpp Synthesis and Degradation

inEscherichia coli. *J. Mol. Biol.* **259**, 41–57 (1996).

51. Xiao, H. *et al.* Residual guanosine 3′,5′-bispyrophosphate synthetic activity of relA null mutants can

be eliminated by spoT null mutations. *J. Biol. Chem.* **266**, 5980–5990 (1991).

52. Scott, M., Gunderson, C. W., Mateescu, E. M., Zhang, Z. & Hwa, T. Interdependence of Cell Growth

and Gene Expression: Origins and Consequences. *Science* **330**, 1099–1102 (2010).

53. Scott, M., Klumpp, S., Mateescu, E. M. & Hwa, T. Emergence of robust growth laws from optimal

regulation of ribosome synthesis. *Mol. Syst. Biol.* **10**, 747 (2014).

54. Dai, X. *et al.* Reduction of translating ribosomes enables Escherichia coli to maintain elongation

rates during slow growth. *Nat. Microbiol.* **2**, 1–9 (2016).

55. Klumpp, S., Scott, M., Pedersen, S. & Hwa, T. Molecular crowding limits translation and cell growth.

Proc. Natl. Acad. Sci. **110**, 16754–16759 (2013).

56. Anderson, S. E., Vadia, S. E., McKelvy, J. & Levin, P. A. The transcription factor DksA exerts opposing

effects on cell division depending on the presence of ppGpp. *mBio* **14**, e02425-23 (2023).

57. Sanchez-Vazquez, P., Dewey, C. N., Kitten, N., Ross, W. & Gourse, R. L. Genome-wide effects on

Escherichia coli transcription from ppGpp binding to its two sites on RNA polymerase. *Proc. Natl.*

Acad. Sci. **116**, 8310–8319 (2019).

58. Campos, M. *et al.* A constant size extension drives bacterial cell size homeostasis. *Cell* **159**, 1433–

1446 (2014).

59. Männik, J., Kar, P., Amarasinghe, C., Amir, A. & Männik, J. Determining the rate-limiting processes

for cell division in Escherichia coli. *Nat. Commun.* **15**, 9948 (2024).

60. Mori, M. *et al.* From coarse to fine: the absolute Escherichia coli proteome under diverse growth

conditions. *Mol. Syst. Biol.* **17**, 1–23 (2021).

61. Balakrishnan, R. *et al.* Principles of gene regulation quantitatively connect DNA to RNA and proteins

in bacteria. *Science* **378**, eabk2066 (2022).

62. Sargent, M. G. Control of cell length in Bacillus subtilis. *J. Bacteriol.* **123**, 7–19 (1975).

63. Moore, D. A., Whatley, Z. N., Joshi, C. P., Osawa, M. & Erickson, H. P. Probing for Binding Regions of

the FtsZ Protein Surface through Site-Directed Insertions: Discovery of Fully Functional FtsZ-

Fluorescent Proteins. *J. Bacteriol.* **199**, 10.1128/jb.00553-16 (2016).

64. Reyes-Lamothe, R., Sherratt, D. J. & Leake, M. C. Stoichiometry and Architecture of Active DNA

Replication Machinery in Escherichia coli. *Science* **328**, 498–501 (2010).

65. Marceau, A. H. *et al.* Structure of the SSB–DNA polymerase III interface and its role in DNA

replication. *EMBO J.* **30**, 4236–4247 (2011).

66. Gray, W. T. *et al.* Nucleoid Size Scaling and Intracellular Organization of Translation across Bacteria.

Cell **177**, 1632-1648.e20 (2019).

67. Karp, P. D. *et al.* The EcoCyc Database (2023). *EcoSal Plus* **11**, eesp-0002-2023 (2023).

68. Hui, S. *et al.* Quantitative proteomic analysis reveals a simple strategy of global resource allocation

in bacteria. *Mol. Syst. Biol.* **11**, 784 (2015).

69. Si, F. *et al.* Invariance of Initiation Mass and Predictability of Cell Size in Escherichia coli. *Curr. Biol.*

CB **27**, 1278–1287 (2017).

70. Helgstrand, M. *et al.* The Ribosomal Stalk Binds to Translation Factors IF2, EF-Tu, EF-G and RF3 *via* a

Conserved Region of the L12 C-terminal Domain. *J. Mol. Biol.* **365**, 468–479 (2007).

71. Shiomi, D., Sakai, M. & Niki, H. Determination of bacterial rod shape by a novel cytoskeletal

membrane protein. *EMBO J.* **27**, 3081–3091 (2008).

72. Tamaki, S., Matsuzawa, H. & Matsuhashi, M. Cluster of mrdA and mrdB genes responsible for the

rod shape and mecillinam sensitivity of Escherichia coli. *J. Bacteriol.* **141**, 52–57 (1980).

73. Traag, V. A., Waltman, L. & van Eck, N. J. From Louvain to Leiden: guaranteeing well-connected

communities. *Sci. Rep.* **9**, 5233 (2019).

74. Wolf, F. A. *et al.* PAGA: graph abstraction reconciles clustering with trajectory inference through a

topology preserving map of single cells. *Genome Biol.* **20**, 59 (2019).

Methods

Barcode Assembly

Our barcode library consists of a set of plasmids, with each plasmid containing a DNA sequence encoding an RNA barcode. Each barcode contains a set of 30 unique binding sites representing a 30 bit word. Each binding site is 20 bp long and consists of one of two possible sequences, "0" or "1". If the site consists of sequence "1", it will be complementary to a cognate FISH probe for that site, while if it consists of sequence "0" it will not bind that cognate probe. Site sequences were designed to maximize hybridization rates and minimize crosstalk as described previously for MERFISH secondary probes.[1,21]

Barcodes were generated by cycled ligation assembly (CLA) of pools of 3-bit blocks, each consisting of all 2^3 possible block sequences.[75] Both strands of dsDNA coding for each 3-bit block were synthesized as single stranded oligos and pooled to a final concentration of 50 nM for each oligo. These oligos were then phosphorylated by T4 PNK (44 µL oligo pool, 5 µL T4 ligase buffer, 2 µL PNK (NEB, M0201S)). Following this, the phosphorylated oligo blocks were assembled, in proper order, using a mix of "Scaffold Oligonucleotide Connectors" or SOCs, which are unphosphorylated oligos designed with scaffold sequences complementary to the 3 and 5 prime ends of consecutive blocks (4 µL phosphorylation mix, 2 µL 100 nM SOC mix, 2 µL Ampligase buffer, 2 µL 5 µM/µL Ampligase, 10 water). Thermocycling protocol was 95 °C for 2 min, then 60 cycles of 95 °C for 10 s, 60 °C for 30s and 66 °C for 60 s.

To generate barcodes for insertion into plasmids, we diluted the CLA reaction five-fold and amplified full length barcodes using a 100 µL Phusion polymerase (NEB, M0531S0) reaction with 2 µL of diluted CLR template, for 20 cycles, using primers oDE311 and oDE201. An additional 10 µL of 100 µm oDE201 primer (10X) was added to the reaction, which was followed by a single additional thermocycle. Notably, oDE311 contained a random 20bp N-mer sequence such that the final barcode library also included a (20-mer) unique molecular identifier (UMI). 5 µL of this reaction was inspected on an agarose gel for products of the expected length. If non-specific bands were present, reactions would be repeated with fewer cycles. Acceptable products were purified (Monarch PCR & DNA Cleanup Kit, T1030S) and inserted into a oDE202 and oDE253 primer-amplified pDE47 backbone using isothermal assembly (NEB NEBuilder HiFi DNA Assembly Master Mix, E2621L). Assembled barcode plasmids contained a pBR322 origin, a kanamycin resistance gene, and a T7 promoter controlling barcode expression. These assemblies were then purified (Monarch PCR & DNA Cleanup Kit, T1030S), eluted in 6 µL water of which 2 µL was transformed by electroporation into *E. coli* (DH5G Gold Electrocompetent *E.coli*, Genlantis or NEB 10-beta Electrocompetent *E. coli*). 1 mL SOC was added after electroporation and cells were recovered at 37 °C with shaking for 1 hour. The SOC culture was then diluted in 50 mL of LB broth with 50 µg/mL Kanamycin and cultured at 37 °C 220 rpm in a shaking incubator overnight. In parallel, samples of the SOC culture were dilution plated on LB plates with 50 µg/mL Kanamycin to check for adequate transformation efficiency (>10 million CFU/mL). The following day, DNA from the liquid culture was extracted (GeneJET Plasmid Midiprep Kit, K0482) producing the barcode library.

sgRNA Cloning and Library Assembly

To create a mismatch-CRISPRi library targeting essential genes, we began by selecting all sgRNA sequences in a previous genome-wide CRISPRi screen in *E. coli*.[26] We considered as essential all genes which exhibited a log fold change < -5.57 at the final timepoint in the original

screen or were annotated as essential in a separate TraDIS screen.[25] For each initial sgRNA, the
first member of a mismatch series was generated by mutating the 5' base pair of the targeting
sequence. The second member was then made by further mutating this single-mismatch sequence
at the base pair just 3' to the original mutation. This was continued up to 10 consecutive mutations
to generate a total of 11 sgRNAs per target site. 40 control sequences following the design rules
of the original CRISPRi screen[26] and containing less than 8bp of complementarity to any sites in
the genome were also included, as well as a handful of sgRNAs from the previous screen targeting
non-essential genes.

sgRNA targeting sequences were ordered as an oligo pool from Twist, resuspended to a
final (total) concentration of 100 nM in TE buffer. The pool was amplified using a 50 µL Phusion
polymerase (NEB, M0531S0) reaction with 1 µL of a 1:250 dilution of the 100 nM oligo pool and
oDE418/oDE433 primers, for 8 cycles. The PCR product was purified (Monarch PCR & DNA
Cleanup Kit, T1030S) and subsequently digested using BbsI-HF (NEB, R3539S) for 30 mins at
37 °C. 1 µL of 5M NaCl to condition the buffer for a subsequent digestion with BsmBI-v2 (NEB,
R0739S) for 30 mins at 55 °C. After heat inactivation at 80 °C for 20 mins, the digested product
was purified with a low molecular weight protocol (Monarch PCR & DNA Cleanup Kit, T1030S).

The sgRNA plasmid backbone (pDE93) contains a constitutive sgRNA expression
cassette, a pBR322 origin and a Kanamycin resistance gene. This plasmid was digested with BbsI-
HF (NEB, R3539S) and CIP treated (NEB, M0525S) to prevent backbone relegation at 37 °C for
1 hour. After heat inactivation at 80 °C for 20 mins, the digested product was run on an agarose
gel and the band at the expected backbone size gel purified (Monarch DNA Gel Extraction Kit,
T1020L). The digested sgRNA library was then ligated into the plasmid backbone using a 50-100
µM T4 DNA ligase reaction (NEB, M0202S) at RT for 30 mins. The reaction was then purified
(Monarch PCR & DNA Cleanup Kit, T1030S) and eluted in 7 µL of water, of which 2 µL was
transformed into electrocompetent *E. coli*, recovered in SOC, cultured overnight, and the library
was extracted according to the same protocol as with the barcode library.

Barcoding Library

To assemble our variant libraries with barcodes, we digested both our variant library,
barcode plasmid library and a destination vector with BsaI-HFv2 (NEB, R3733S) for 1 hour at 37
°C. The digests were then run on an agarose gel and bands were gel purified corresponding to the
cassettes of the two libraries and the backbone of the destination plasmid (p15A ori, Chlor-R).
Finally, the two libraries were combinatorially ligated using a 40 µL T4 DNA ligase reaction at
RT for 30 mins, purified and eluted in 10 µL. 1 µL of this elution was transformed into
electrocompetent *E. coli* with the relevant genetic background and recovered in SOC according to
the same protocol as with the barcode library.

To test transformation efficiency, the recovery culture was dilution plated on LB plates
supplemented with 25 µg/mL Chloramphenicol, cultured overnight at 37 °C and plating efficiency
(CFU/mL recovery mix) measured the next day. Transformation and recovery were then repeated,
followed by dilution and plating of the recovery mixture with a target of ~10000 CFU/plate, onto
20-30 plates. An additional dilution series was also performed on this day to measure the efficiency
of this transformation. Plates were cultured overnight at 37 °C and the library bottlenecked to the
desired complexity by solubilizing colonies on a number of plates corresponding to the desired
CFUs, in EZRDM (Teknova, M2105) supplemented with 25 µg/mL Chloramphenicol. The OD600
of the resulting culture was measured and further diluted in media to desired OD600 of 5. Finally,

the diluted culture was diluted 1:1 in 50% glycerol, separated into 100 µL aliquots and stored at -
80 °C for storage. The residual culture plasmid was extracted (GeneJET Plasmid Midiprep Kit,
K0482) for later use in sequencing of the barcode to variant map.

Library Sequencing

According to our procedure, barcodes are randomly ligated to variants, so we use
sequencing to establish the lookup table from each barcode to its corresponding variant. Since the
length of the variant-barcode cassette is long (>700bp), we used Nanopore long read sequencing
to acquire reads spanning the entire construct and leveraged our barcodes to bin reads by library
members.

Library plasmid cassettes were first amplified in a 400 µL Phusion polymerase PCR with
primers oDE154 and oDE201 for 12-15 cycles, depending on how many cycles were needed to
yield sufficient material. PCR products were then purified by AMPure bead cleanup (Beckman
Coulter, A63881). Amplicons were then prepared for sequencing using a standard protocol from
Oxford Nanopore Technologies (ONT) which included FFPE DNA Repair (NEB, M6630S), End
Repair/dA-Tailing (NEB, E7546S), and ligation of adaptors from the ONT SQK-LSK109 ligation
sequencing kit. Prepared amplicons were then run on a ONT MinION with either R9.4.3 or R10.3.1
pores. If necessary to achieve adequate depth, multiple sequencing runs were conducted on a single
library and the reads pooled.

Raw reads were processed using high accuracy calling (HAC) with the Guppy basecaller.
Following this, reads were aligned to a graph genome reference (gfa) specifying the combinatorial
structure of our barcodes using GraphAligner.[76] The alignment, corresponding to a traversal
through this barcode graph, provided the barcode contained in each read, allowing us to
subsequently group reads based on barcode. Grouped reads were then aligned to a reference using
Minimap2[77], also specified by the graph alignment, and finally used to generate a barcode-variant
consensus using Medaka. This analysis was parallelized using the Snakemake[78] module in Python
(https://github.com/paulssonlab/marlin-nanopore), enabling us to scale this approach to large
libraries.

MARLIN Phenotyping

Prior to experiment, 100 µL library aliquots of *E. coli* were inoculated into 5 mL cultures
of EZRDM with 25 µg/mL Chloramphenicol and grown at 37 °C 220 rpm on a shaking incubator
to saturation (6-8 hours). Prior to loading cells, the microfluidic was primed by injecting ~200 µL
of faCellatite BIOFLOAT™ solution (faCellitate, F202005) and allowed it to adsorb at RT for 30
mins. 3 mL of saturated library culture was concentrated 10X by spinning in a microcentrifuge at
4000 rcf for 1 min. Supernatant was then collected to leave a residual ~300 µL of liquid before
resuspension of the pellet. The concentrated culture was then injected into the microfluidic to load
the library. To ensure maximum loading, the microfluidic was spin loaded using a custom
microfuge adaptor, at 650 rcf for 30s, loading the cells into chambers in the downward orientation.
The orientation of the device in the adaptor was then reversed and spun at 650 rcf for 30 s, to load
chambers in the upward orientation. Once loaded, the device is connected to tubing at the inlet
carrying growth medium driven upstream of the device by a peristaltic pump (Longer Precision
Pump Co. Ltd, T60-S2&WX10-14), while the outlet is connected to a waste line. The device and
feeding lines were then transferred to the microscope for imaging.

Imaging was conducted on a Nikon Ti-2 microscope, with the entire stage enclosed in an incubator (Okolabs) to maintain a constant temperature of 30 °C (lDE20, lDE28 and lDE30) or 37 °C (lDE15). The device was imaged using either a Nikon Plan Apo Lambda 20x Ph2 DM air objective with 0.75 NA (lDE15 and lDE20) or a Nikon Plan Apo Lambda 40x Ph2 DM air objective with 0.95 NA (lDE26, lDE28 and lDE30). All images were acquired using a high pixel density (4.5 μm x 4.5 μm pixel size) sCMOS camera (Photometrics Iris 9) to enable adequate sampling of the 20x objective point spread function. We used an LED lamp (CoolLED pE-100) as a white-light source for phase-contrast imaging and a Spectra III mixed LED/Laser light source (Lumencor) for fluorescence imaging. For most experiments, our light path consisted of a multi-band dichroic (409/493/573/652/759 Brightline, Semrock) and a variable emission filter (519/26, 595/33, or no filter), mounted on an emission filter wheel for fast multi-channel imaging (Finger Lakes Instruments HS-625). For imaging the mKate2Hyb marker in lDE20, we used an alternative configuration with a multi-band dichroic (89403, Chroma) and an 642/80 emission filter.

Before imaging, cells in the microfluidic were allowed to proliferate on-scope for a period of 10-14 hours, under constant flow of media. The phenotyping portion of the experiment then proceeded, with all microscope control and acquisition handled by the Nikon ND Elements software. Depending on the phenotype being measured, different numbers of fields of view and image cadences were used. Imaging of GFPmut2 and mKate2Hyb in lDE15 was conducted every 40 mins for a period of 5 hours and 20 mins, over a grid of 855 fields of view (FOVs). Imaging of mKate2Hyb in lDE20 was conducted every 10 mins for a period of 10 hours, over 629 FOVs. Finally, imaging of mVenus and HU-mCherry in lDE26, lDE28 and lDE30 was conducted every 10 mins for a period of 10 hours, over 325 FOVs. In all cases, the Nikon perfect focus system (PFS) was engaged throughout the experiment to maintain the sample in the focal plane. In all 10 hour CRISPRi experiments, aTc (Sigma, #37919) was added to the flask containing the input growth medium to a final concentration of 0.2 μM at the 2 hour mark, in order to induce dCas9 expression.

MARLIN Genotyping

Genotyping FISH probes complementary to the MERFISH barcode were ordered from IDT, with the probe DNA conjugated at the 5' end with a six carbon (C6) linker to a di-thiol group followed by another C6 linker conjugated to a dye. Depending on the experiment, we used probe sets utilizing three of the following five dyes: Alexa Fluor 488, Alexa Fluor 555, Alexa Flour 647, Cy5 and Alexa Fluor 750. Probes were received as lyophilized pellets, resuspended to 100 μM using nuclease-free water, split into a number of strip tubes, and re-lyophilized in 2 nmol aliquots. The dry aliquots were stored at -80C for long term storage. 100 μM stocks were prepared from these aliquots by adding 20 μL of nuclease-free water and kept at -20C for less than six months or until exhaustion. Working stocks of probe combinations to be used in the same imaging cycle were prepared by adding 6.66 μL each of three 100 μM probe stocks to 180 μL of readout probe buffer (TE buffer with 0.1% Triton X-100) to a final concentration of 3.33 μM per probe. Working stocks were stored at -20C for less than two months or until exhaustion. Each round of FISH probing uses 15 μL of this working stalk in a 5 mL hybridization buffer (10% Ethylene Carbonate, 0.125% Triton X-100, 40 U/mL Murine RNAse inhibitor and 20 nM DNA probe in pH 7.0 2X SSC).

Genotyping was conducted immediately after the conclusion of phenotype imaging for each library experiment. Cells in the device were induced for barcode expression for a period of ~75 mins by the addition of Cuminic acid to a final concentration of 100 μM in the input growth media. During this time, we performed setup of our automated liquid handling system (see

extended methods), which enabled automated selection of FISH reagents in conjunction with image acquisition. We controlled this system, in conjunction with the microscope, from a Jupyter notebook making use of both a custom Python module for this purpose (https://github.com/paulssonlab/marlin) and the Python Micromanager API.[79] During setup, all liquid lines in this system were cleaned using DI water and inspected for adequate flow. Once barcode induction was concluded, we loaded the liquid handler with our FISH reagents and replaced the growth medium line with the liquid handler line, just upstream of the device inlet. The liquid handler proceeded to perform fixation and FISH imaging over the next 16-20 hours.

During FISH genotyping, we first inject a 4:1:1:4 solution of methanol, acetone, 20X SSC and nuclease-free water (50% MeAc) into the device, followed immediately by a 4:1 solution of methanol and acetone (MeAc) to fix the cells. We allow fixation to proceed for 45 mins, then inject 50% MeAc, followed by the hybridization solution containing the first probe set. After allowing probes to hybridize for 30 mins, we inject an oxygen-scavenging imaging buffer solution consisting of 3mM PCA, 0.03% Trolox, ~0.18 U/mL rPCO, and Trolox-Quinone (>30 μM, prepared according to a previously published protocol[80]) in pH 7.0 2X SSC. Imaging buffer is flowed for an additional 5 mins to clear any bubbles from the device and then FISH imaging proceeds. During each round of imaging, all FOVs phenotyped previously are imaged in phase-contrast for lineage-detection purposes and three additional fluorescence channels, corresponding to each probe dye. After all channels are imaged, we inject a cleavage buffer (33.3 mL TCEP in pH 7.0 2x SSC) containing a reducing agent to break the di-thiol bond linking dyes to DNA probes, eliminating the fluorescence signal from this round of imaging. To continue genotyping, we repeat this protocol from the hybridization step, for the next set of probes. We repeat 10 such cycles, reading out 3 DNA probes in each, for a total of 30 probe readouts, completing the experiment.

Mother Machine Isolate Imaging

Mother machine experiments with single isolates were performed using the same imaging setup as described in "MARLIN Imaging". Starter cultures for strains to be loaded into the device were generally prepared 12-16 hours in advance, in the same manner as described in "Strains and Growth Conditions". All CRISPRi validation strains were imaged according to the same schedule and under the same conditions as described for lDE20.

Strains and Growth Conditions

Cells were initially inoculated into 5 mL EZ Rich Defined Medium (EZRDM) with appropriate antibiotics in a test tube from a single colony on an agar plate, struck within 2 weeks of the experiment. Cells were then cultured for 14-16 hrs at 37 °C in a shaking incubator (220 rpm), until saturation. Cells were then diluted 10^{-7} into 5 mL EZRDM with appropriate antibiotics, as a pre-culture, and incubated for an additional 10-12 hrs at 30 °C in a shaking incubator (220 rpm). We then measured OD600 for the pre-cultures and proceeded to dilute each into 25 mL or 5 mL of inductive media (EZRDM + 0.2 μM aTc) with or without 1000 μM IPTG, depending on the experiment. Cultures with OD600 over 0.5 were discarded. Induced cultures were subsequently incubated for 6 hrs at 30 °C in a shaking incubator (220 rpm). Cultures with OD600 between 0.1 and 0.4 at this time were further processed to measure the RNA/Protein ratio (see "RNA/Protein Measurement") and the (p)ppGpp concentration (see "ppGpp/GTP Sample Harvest Protocol").

To prepare fixed samples for agar pad imaging the culture protocol was similar, with starter cultures and pre-cultures prepared in the same manner. Pre-cultures were then diluted in 2.5 mL of inductive media (EZRDM + 0.4 μM aTc) and incubated at 30 °C in a shaking incubator (220

rpm) for 6 hours. The OD600 of the cultures were measured and the cultures processed if their OD600 was between 0.1 and 0.4. Subsequently, 360 µL of 32% PFA was added to each tube, which was then placed back into the incubator for a total of 30 minutes. Cells were then collected by centrifugation of the fixed culture at 4000 g for 5 mins, at RT in 10 mL falcon tubes, removing the supernatant and resuspending the pellet in 500 µL of PBS, this centrifugation step was repeated with a resuspension volume of 500 µL and then once more with a final resuspension volume of 250 µL. Samples were then stored at 4 °C until staining and imaging.

Methods References

75. Roth, T. L., Milenkovic, L. & Scott, M. P. A Rapid and Simple Method for DNA Engineering Using Cycled Ligation Assembly. *PLOS ONE* **9**, e107329 (2014).

76. Rautiainen, M. & Marschall, T. GraphAligner: rapid and versatile sequence-to-graph alignment. *Genome Biol.* **21**, 253 (2020).

77. Li, H. Minimap2: pairwise alignment for nucleotide sequences. *Bioinformatics* **34**, 3094–3100 (2018).

78. Köster, J. & Rahmann, S. Snakemake—a scalable bioinformatics workflow engine. *Bioinformatics* **28**, 2520–2522 (2012).

79. Edelstein, A. D. *et al.* Advanced methods of microscope control using µManager software. *J. Biol. Methods* **1**, e10 (2014).

80. Cordes, T., Vogelsang, J. & Tinnefeld, P. On the Mechanism of Trolox as Antiblinking and Antibleaching Reagent. *J. Am. Chem. Soc.* **131**, 5018–5019 (2009).

Data availability:

Raw sequencing reads for Nanopore and NGS libraries have been deposited in the NCBI BioProject database under accession number PRJNA1205775. Processed data will be deposited on Zenodo (https://doi.org/10.5281/zenodo.14537796) upon submission. An archived release of the TrenchRipper module version used for this publication is also available on Zenodo at https://doi.org/10.5281/zenodo.14552982. Other source code for image analysis, sequencing analysis, and figure generation are available on GitHub (https://github.com/paulssonlab/2025_Eaton_optical-pooled-screen). Raw or processed image data are available upon request.

Acknowledgements:

We are grateful for the gifts of bacterial strains and plasmids from the labs of Richard Murray, David Bikard, Christopher Voigt, Stanley Qi, Kathleen Matthews, Drew Endy, Keith Shearwin, Greg Bokinsky,

Philippe Cluzel, Donald Court, James Keck, Michael Baym, and Thomas Bernhardt. Mass spectrometry measurements and consultation were performed at the Harvard Center for Mass Spectrometry. Fluid handling systems were built at the Research Instrumentation core at Harvard Medical School. All Computational work was performed on the O2 cluster supported by the Research Computing Group at Harvard Medical School. D.S.E. is grateful for formal and informal feedback on this work from Andrew Murray, Noah Olsman, Gene-Wei Li, Rebecca Ward, Thomas Bernhardt, James Taggart, Julius Palme, and Maximillian Marin. This work was supported by National Institute of Health grant R01GM081563, Defense Advanced Research Projects Agency grant W911NF-19-2-0018, and Advanced Research Projects Agency for Health grant 1AY2AX000005-01. D.S.E., C.S., L.G.L., and J.Q.S. acknowledge support from the Systems, Synthetic and Quantitative Biology PhD program training award (T32GM135014). D.S.E. was supported in part by the NSF-Simons Center for Mathematical and Statistical Analysis Biology at Harvard (award number #1764269), and the Harvard Quantitative Biology Initiative.

Author Contributions: D.S.E., J.E.M. and J.P. conceived of and designed the study. D.S.E., C.S., and L.G.L. carried out the principal experiments and data analysis. J.Q.S. and Y.G. designed and fabricated the microfluidic master molds. B.R.W. assisted with automation of liquid handling. D.S.E., C.S., L.G.L. and V.H. engineered the genetic strains made for this study. J.E.M., J.P. and E.C.G. supervised the study. D.S.E., Q.A.J., J.E.M. and J.P. wrote the paper.

Competing Interest Declaration: The authors declare no competing interests.

Corresponding Author Line: Correspondence to Daniel S. Eaton, Jeffrey R. Moffitt, or Johan Paulsson.