Neural crest induction requires SALL4-mediated BAF recruitment to lineage specific enhancers

1 Martina Demurtas[1], Samantha M. Barnada[2], Emma van Domselaar[1], Zoe H. Mitchell[1],

2 Laura Deelen[1], and Marco Trizzino[1,*]

4 [1]Department of Life Sciences, Imperial College London, London, UK

5 [2] Department of Biochemistry and Molecular Biology, Thomas Jefferson University, Philadelphia, PA, USA

6 *Lead contact: m.trizzino@imperial.ac.uk

ABSTRACT

Neural crest induction begins early during neural plate formation, requiring precise transcriptional

control to activate lineage-specific enhancers. Here, we demonstrate that SALL4, a transcription

factor associated with syndromes featuring craniofacial anomalies, plays a critical role in early

cranial neural crest (CNCC) specification. Using *SALL4*-het-KO human iPSCs to model clinical

haploinsufficiency, we show that SALL4 directly recruits BAF to CNCC-lineage specific enhancers

at the neuroectodermal stage, specifically when neural crest gene expression is induced at the

neural plate border. Without functional SALL4, BAF is not loaded at chromatin, leaving CNCC

enhancers inaccessible. Consequently, the cells cannot undergo proper CNCC induction and

specification due to persistent enhancer repression, despite normal neuroectodermal and neural

plate progression. Moreover, by performing SALL4 isoform-specific depletion, we demonstrate

that the SALL4A is the isoform essential for CNCC induction and specification, and that SALL4B

cannot compensate for SALL4A loss in this developmental process.

In summary, our findings reveal SALL4 as essential regulator of BAF-dependent enhancer

activation during early stages of neural crest development, providing molecular insights into

SALL4-associated craniofacial anomalies.

KEYWORDS

26 SALL4, DPF2, SALL4A, BAF, neural crest, enhancers, chromatin accessibility

INTRODUCTION

Transcription factors and chromatin remodelers play crucial roles in cell fate determination (Gopinathan and Diekwisch, 2022; Islam et al., 2021; Kim and Shendure, 2019; Li, 2002; Voss and Hager, 2014; Zheng et al., 2021). Among these, the C2H2 zinc-finger SALL4 is involved in a wide range of developmental processes (Tatetsu et al., 2016). SALL4 is highly expressed in pluripotent cells and throughout embryonic development, gradually decreasing after birth to become undetectable in adult tissues (Xiong et al., 2015).

In humans, *SALL4* produces two main isoforms: SALL4A (1053 aa) and SALL4B (617 aa), which have been proposed to play distinct roles in ESCs. Namely, Rao et al. (2010) suggested that SALL4B has function limited to pluripotency regulation in ESCs, while SALL4A is primarily active during development. Although *SALL4* is expressed as early as the two-cell stage, its role in maintaining pluripotency remains debated. Some studies suggest that SALL4 cooperates with core pluripotency factors to sustain stem cell identity (Yang, 2018; Zhang et al., 2006). In contrast, others propose that SALL4 plays a more marginal role in pluripotency regulation (Miller et al., 2016; Wang et al., 2014; Yuri et al., 2009).

During early mouse development, *Sall4* is broadly expressed throughout the embryo, however, as gastrulation progresses, its expression becomes more posteriorly restricted, localizing primarily to the developing tail bud and limb buds (Chen et al., 2022; Tahara et al., 2019). Here, it regulates cell fate commitment of neuromesodermal progenitors by suppressing neural genes in mesodermal cells while simultaneously enhancing mesodermal gene expression (Pappas et al., 2024). Additionally, *Sall4* remains expressed in mouse craniofacial regions until E9.5, including the developing frontonasal process and the mandibular and maxillary arches (Tahara et al., 2019). These craniofacial structures originate from cranial neural crest cells (CNCCs), a subpopulation of the neural crest cells that gives rise to the cartilage and bones of the face and anterior skull, as well as cranial neurons and glia (Dash and Trainor, 2020; Milet and Monsoro-Burq, 2012; Pla and Monsoro-Burq, 2018; Roth et al., 2021; Szabó et al., 2018). Consistent with this, individuals with *SALL4* haploinsufficiency exhibit a range of phenotypes, including distinctive facial features such as epicanthal folds, widely spaced eyes, and a depressed and flattened nasal bridge, along with cranial nerve-related conditions and underdevelopment of one side of the face (Kekunnaya and Negalur, 2017). Many of these distinctive features are classic indicators of impaired CNCC development. *SALL4*-associated phenotypes include Duane anomaly, a congenital strabismus resulting from abnormal oculomotor nerve innervation (Kekunnaya and Negalur, 2017), and sensorineural hearing loss (Terhal et al., 2006), but also upper limb anomalies (Duane-radial ray syndrome) and, in some cases, renal anomalies.

Together, these findings suggest that SALL4 may play a critical role in CNCC development, which

has not yet been investigated.

Previous studies have also suggested that SALL4 interacts directly with chromatin remodeling

complexes. For example, Sall4 is an interactor of the NuRD complex, in mESCs and during

development (Bode et al., 2016; Liu et al., 2018b; Lu et al., 2009; Miller et al., 2016; Wang et al.,

2023). Additionally, recent mass-spectrometry experiments conducted by our group in early

stages of human iPSC-to-CNCC differentiation, and specifically at the neuroectodermal stage,

revealed that SALL4 interacts with the BAF complex at this developmental stage (Pagliaroli et al.,

2021). Since this was the first report of a potential direct SALL4-BAF interaction, here we set out

to investigate the scope of this interaction and determine the role of SALL4 during CNCC

development.

RESULTS

SALL4 interacts with the BAF complex

CNCCs can be generated *in vitro* using human induced pluripotent stem cells (hiPSCs). This

protocol is two weeks long, and it first goes through a neuroectoderm intermediate (Bajpai et al.,

2010; Barnada et al., 2024; Mitchell et al., 2025; Pagliaroli et al., 2021; Prescott et al., 2015). In

our previous work (Pagliaroli et al., 2021), mass spectrometry (IP-MS) conducted at early stages

of iPSC-to-CNCC differentiation (Day-5, corresponding to neuroectoderm), revealed that SALL4

interacts with BAF at this developmental stage. Since this was the first report of a SALL4-BAF

interaction, we performed a co-immunoprecipitation (co-IP) at Day-5 to validate it, pulling-down

ARID1B, which is the only ARID1 subunit active and incorporated in BAF at this stage (Pagliaroli

et al., 2021), and blotting for SALL4. This co-IP confirmed that ARID1B interacts with both SALL4

isoforms at this time-point (Fig. 1A).

BAF is a multi-subunit assembly composed of a dozen subunits, which are differentially

incorporated in a tissue-specific manner (Hodges et al., 2016). Based on which subunit is present

in each complex, in mammals these can be classified into canonical BAF, polybromo-associated

BAF, and non-canonical BAF (Zhang et al., 2021). While our mass-spec and co-IP experiments

supported the interaction between SALL4 and canonical BAF (ARID1B is not incorporated in other

BAF configurations), neither of these two approaches has the resolution to identify the exact BAF

subunit that mediates the interaction with SALL4. Therefore, we performed an *in silico* analysis

using AlphaFold3 (Jumper et al., 2021). We tested various BAF subunit combinations along with

four histones and an AT-rich DNA sequence which has been empirically shown to be recognized

by SALL4 (5'-GGAAATATTACC-3'; PDB: 8A4I) and SALL4A. The output was statistically

significant, with 0.45 predicted template modeling (pTM) score, and 0.47 interface predicted

template modeling (ipTM) score (Figs. S1A, B). Through this approach, AlphaFold3 predicted a

direct interaction between DPF2, a broadly expressed BAF subunit, and the zinc-fingers 5 and 6

(ZFC3) and 7 (ZFC4) of SALL4A (Figs. 1B, C, D; Figs. S1A, B; Table 1). *DPF2* is a member of

the *BAF45* gene family, which also includes *PHF10*, *DPF1*, and *DPF3*. We were able to rule out

its paralogues based on their distinct expression patterns, incorporation into specific BAF

complexes, and previous reports that BAF45 subunits are not functionally redundant (Zhang et

al., 2019a). In particular, PHF10 is exclusively associated with the pBAF, which does not

incorporate ARID1B (Krasteva et al., 2017), while DPF1 and DPF3 are incorporated in BAF only

in neurons (Lessard et al., 2007). Moreover, we compared the gene expression of each *DPF*

paralogue at both Day-0 and Day-5 and observed no significant difference in expression between

SALL4-WT and *SALL4*-het-KO for any of them (Figs. 1E, F).

This prediction was particularly interesting given that DPF2 deletion has been linked to early

neural ectoderm differentiation in embryonic stem cells (ESCs) (Zhang et al., 2019a). Notably, of

the two main SALL4 isoforms (SALL4A and SALL4B), SALL4B lacks the portion of exon 2 that

contains ZFC3. This prompted us to explore whether the SALL4-DPF2 interaction could be

isoform-specific. To address this, we reanalysed the interactions in AlphaFold3, incorporating

SALL4B to the same BAF subunits as before. To ensure biological relevance, we first excluded

two of the five models that did not exhibit interactions with the AT-rich DNA sequence, as SALL4

is expected to bind DNA for its function. Among the remaining three models, we observed that

SALL4B interacted with DPF2, but unlike the SALL4A isoform, its interaction was not domain

specific. Instead, the predicted interactions for SALL4B were characterized by annotated

disordered regions and appeared to be largely driven by a non-specific, zinc-finger-to-zinc-finger

association (Figs. S1C, D; Table S2), with no particular domain-specific interactions identified.

To accurately interpret our AlphaFold3 predictions, we examined two key confidence metrics: the

Predicted Aligned Error (PAE), which assesses the reliability of the relative positioning of residues

within the structure, and the Predicted Local Distance Difference Test (pLDDT), which provides a

per-residue confidence score for the model (Table S1). Our SALL4-DPF2 contact analysis

revealed 394 atomic interactions (center-to-center distance ≤ 8Å) involving 139 residues (Fig. 1D

and Table S1). Encouragingly, the pLDDT values were highest in known functional domains

involved in the interaction, particularly ZF5, ZF6, and ZF7 in SALL4A, as well as the PHD1 and

PHD2 domains in DPF2, reinforcing the reliability of these predicted structures. Consistently, our

model also aligned closely with experimentally resolved structures at 3 Å resolution available for

the BAF base module (Fig. 1E). To refine our analysis, we focused on residues where at least

half of their atomic interactions had a pLDDT > 65 (70 $\pm$ 10%) and that were annotated as part of

a functional domain, ensuring that only higher-confidence and potentially functional interactions

were considered. As a result, ZF7 was excluded from further analysis (Table S1).

In summary, our experiments and analyses confirmed that SALL4 interacts with BAF during early

stages of human CNCC specification, and predicted that this interaction might be mediated by

the BAF subunit DPF2 through ZFC3.

Generation of a *SALL4*-haploinsufficient human iPSC lines

Our experiments so far revealed that SALL4 interacts with BAF at early stages of CNCC

specification, and specifically at the neuroectodermal stage. To investigate the scope of this

interaction, we generated *SALL4* haploinsufficient iPSC lines (hereafter *SALL4*-het-KO; Fig. 2A).

We opted for heterozygous KO of *SALL4* as this recapitulates the human *SALL4* syndromes,

which are all caused by *SALL4* haploinsufficient variants (Al-Baradie et al., 2002; Rgen Kohlhase

et al., 2002; Terhal et al., 2006). Two different CRISPR/Cas9 clones were generated and used

as biological replicates. Specifically, the CRISPR-KO was generated by introducing a 11 bp (clone

C4) deletion and a 19 bp (clone F1) deletion, respectively (Fig. S2A), both within the 3'-terminus

of exon 1 of *SALL4*. This is a conserved region across both *SALL4* isoforms that lacks functionally

significant domains. Therefore, both *SALL4* isoforms were equally impacted by the CRISPR

deletions.

SALL4 immunoblot and immunofluorescence assays confirmed that SALL4 protein expression

was significantly reduced in the *SALL4*-het-KO iPSCs compared to controls (*SALL4*-WT; Figs.

2B, C; Fig. S2B, C). We noted that SALL4A protein expression in the *SALL4*-het-KO line was

significantly less than the expected 50% of the wild-type, suggesting potential negative feedback

mechanisms triggered by the reduced SALL4 levels, possibly due to the preferentially

homodimeric nature of this protein. On the other hand, SALL4B protein was barely detectable in

both *SALL4*-WT and *SALL4*-het-KO conditions (Fig. 2B).

In contrast, at the gene expression level we did not observed significant differences in *SALL4*

expression between *SALL4*-WT and *SALL4*-het-KO, for any isoform (Fig. 2D). This may reflect

autoregulatory mechanisms, where reduced SALL4 protein levels in the knockout trigger

compensatory transcriptional upregulation of the gene which however does not result in increased

protein level because of nonsense-mediated decay or other post-transcriptional regulatory

mechanisms.

Next, we investigated whether other *SALL* paralogs (*SALL1*, *SALL2*, *SALL3*) could compensate

for SALL4 loss. First, we investigated their mRNA expression (Fig. 2E) and found *SALL1* as the

only paralog significantly upregulated in *SALL4*-het-KO (Fig. 2E). SALL1 upregulation was ever

greater at protein level (Fig. 2F), indicating a potential compensation in pluripotent cells by post-

transcriptional regulation.

Importantly, *SALL4*-het-KO iPSCs remained viable and exhibited hallmarks iPSC morphology

(Fig. S2D). Furthermore, these cells showed no significant difference in growth rate compared to

controls (Fig. 2G).

Decreased SALL4 expression does not affect the stem cell identity of the iPSCs

Given the debated role of SALL4 in maintaining pluripotency, we investigated whether *SALL4*-

het-KO impacted iPSC stem cell identity. Protein levels of the pluripotency transcription factors

NANOG and OCT4 remained unaltered in the *SALL4*-het-KO cells relative to controls (Fig. 3A).

Similarly, the levels of the pluripotency surface markers TRA-1-60-R and SSEA4 were also not

significantly affected by decreased SALL4 (Fig. 3B; Fig. S4A). Moreover, we examined the

transcriptome of the iPSCs and identified only 114 differentially expressed genes in *SALL4*-het-

KO cells relative to controls (FDR < 0.05; Fig. 3C and Table 3). Notably, canonical pluripotency

markers, such as *POU5F1, NANOG, SOX2, KLF4,* and *MYC*, were not significantly differentially

expressed (Fig. 3C and Table S3). Consistent with this, gene ontology analysis confirmed that

neither up- (Fig. S3A) or downregulated (Fig. S3B) genes were enriched for pluripotency-

associated terms. Finally, by performing tri-lineage differentiation we confirmed that SALL4

depletion did not impact the ability of the *SALL4*-het-KO cells to successfully differentiate into all

three of the germ layers (Fig. S3C).

Together, these findings suggest that *SALL4* haploinsufficiency did not affect IPSC stemness

and self-renewal, possibly as a result of SALL1-mediated compensation.

SALL4 relocates to CNCC developmental enhancers upon CNCC induction cues

After establishing that *SALL4*-het-KO did not affect iPSC stemness, we differentiated the iPSCs

into CNCCs using the same aforementioned protocol (Bajpai et al., 2010; Barnada et al., 2024;

Mitchell et al., 2025; Pagliaroli et al., 2021; Prescott et al., 2015) (Fig. 4A). Since the SALL4-BAF

interaction was detected at Day-5 (Pagliaroli et al., 2021), corresponding to neuroectodermal

spheres, we performed CUT&RUN to profile SALL4 binding in *SALL4*-WT cells in iPSCs and at

Day-5 of differentiation. We performed two SALL4 CUT&RUN replicates per timepoint, and only

retained replicated peaks in each timepoint (FDR < 0.05). With this approach, we identified

comparable amounts of SALL4-bound genomic sites at Day-0 and Day-5 (29,088 and 32,707,

replicated peaks, respectively, FDR < 0.05; Fig. 4B). Further analysis of these sites indicated that

approximately 20% of the Day-0 SALL4 peaks were unique to iPSCs and not detected at Day-5

(hereafter SALL4 Day-0 specific peaks; n = 6,114; Fig. 4B, 4C, left). On the other hand, ~50% of

the replicated peaks detected at Day-5 were exclusive of this developmental stage and were not

detected in iPSCs (hereafter SALL4 Day-5-specific peaks, n = 17,137; Fig. 4B and 4C). Finally, a

subset of ~15,000 sites remained bound by SALL4 at both time points.

These data indicate that, upon neural crest differentiation cues, there is a major shift in SALL4

binding, resulting in the relocation of this transcription factor at ~17,000 previously unbound

regions.

To characterize the scope of SALL4 relocation, we analyzed the positional distribution of SALL4

peaks by associating each peak with its nearest transcription start site (TSS), identifying 4,362

genes at Day-0 and 6,861 genes at Day-5 (Table S4). Consistent with SALL4's established role

as an enhancer-binding transcription factor, 87% of SALL4-bound regions at Day-0 were

located >1 kb from the nearest TSS, suggesting these represent putative enhancers. This

proportion increased substantially to 97% at Day-5 (Fig. 4D).

GO terms enriched in the "Day-5 specific" genes included pathways associated with neural crest

development, such as neural tube development, face development, and cardiac chamber

development (Fig. 4E; FDR < 0.05).

To further characterise the SALL4 bound elements at Day-0 and Day-5, we performed sequence-

based computational motif analysis. At Day-0, motifs associated with core pluripotency

transcription factors such as OCT4, NANOG, and SOX2 did not rank among the top 20 enriched

sequences, further supporting that SALL4 may not be essential for maintaining pluripotency in

human cells (Fig. S5A).

On the other hand, in addition to general enhancer activator factors (AP-1, TEAD), the Day-5

specific SALL4 peaks were enriched for many CNCC signature factors, including SOX9, TFAP2A,

TWIST1/2, SNAI1, NR2F1/2, TCF4/12 and several others (Fig. 4F). Moreover, neuroectodermal

transcription factors, such as ZIC2/3 were also enriched. Additionally, *de novo* motif analysis in

iPSCs revealed enrichment for the CTCF-like motifs (Fig. S5B), while at Day-5 there was

enrichment for enhancer activators TEADS/ETS and AP-1 (Fig. S5C). Finally, AT-rich motifs

resembling the SALL4 annotated motif were found only in a small minority of peaks in iPSCs (Fig.

S5B). This might suggest that SALL4 binding at the CNCC enhancers is not dictated by

recognition of its cognate motif, and possibly mediated by other lineage-specific transcription

factors.

In summary, our data so far indicate that SALL4 relocates to CNCC developmental enhancers

upon differentiation cues.

Chromatin accessibility at the CNCC enhancers is SALL4 dependent

Given that we demonstrated that SALL4 interacts with BAF at early stages of CNCC specification, we reasoned that SALL4 binding at the CNCC developmental enhancers could be necessary for their chromatin activation. We profiled chromatin accessibility by ATAC-seq at Day-0 and Day-5 in both *SALL4*-WT and *SALL4*-het-KO conditions (two biological replicates per timepoint per condition, only peaks present in both replicates were considered; FDR < 0.05). The number of ATAC-seq replicated peaks identified in *SALL4*-WT condition was comparable between Day-0 (61,660 peaks; FDR < 0.05) and Day-5 (58,375 peaks; FDR < 0.05; Fig. 5A). Next, we specifically investigated the chromatin state of the 17,137 genomic sites that gained SALL4 binding at Day-5, and found that SALL4 binding at these sites correlates with a marked increase in chromatin accessibility compared to Day-0 (Fig. 5B). Strikingly, the accessibility at these regions was significantly attenuated in *SALL4*-het-KO relative to *SALL4*-WT (Fig. 5C). This suggests that, in response to differentiation signals, SALL4 relocates to CNCC developmental enhancers, which subsequently become activated through chromatin opening, which is SALL4 dependent.

Motif analysis of regions that lost accessibility in *SALL4*-het-KO cells (n = 1,649; Fig. 5A) showed again enrichment for CNCC factors such as ZIC2, ZEB1/2, and SNAIL, along with BMP signalling factors such as SMADs, and enhancer activators such as AP-1 (Fig. 5D). On the other hand, regions aberrantly accessible in the *SALL4*-het-KO cells (n = 3,460; Fig. 5A) showed enrichment for pluripotency factors, CTCF, and neural plate markers (Fig. 5E).

SALL4 recruits the BAF complex at CNCC developmental enhancers to increase chromatin accessibility

Since SALL4 binding at the CNCC enhancers is necessary for chromatin opening, we investigated whether the SALL4-bound developmental enhancers were also occupied by BAF, and if BAF binding at these sites could be SALL4 dependent. We performed ChIP-seq for BRG1 (SMARCA4), which is the only catalytic BAF subunit active at this sage, in *SALL4*-WT cells at Day-5. We found that BRG1 consistently binds the same genomic sites as SALL4, supporting the idea that these factors interact and co-occupy regulatory elements during early differentiation (Fig. 6A).

To assess whether BRG1 binding depends on SALL4, we performed ChIP-seq for BRG1 in both *SALL4*-WT and *SALL4*-het-KO cells at Day-5. In *SALL4*-WT, we detected 32,833 replicated BRG1 peaks (FDR < 0.05), as opposed to only 14,116 peaks in *SALL4*-het-KO. This suggests that SALL4 loss triggered widespread loss of BAF from chromatin. We focused on BRG1 peaks

lost in *SALL4*-het-KO cells and found that these sites were largely not bound by SALL4 at Day-0,

but they gained SALL4 binding at Day-5 (Figs. 6B–D). As expected, loss of SALL4 and BRG1

from these sites led to marked decrease of chromatin accessibility (Fig. 6E). Moreover, we

explored the identity of some of these putative enhacers by confirming that these overlapped with

some of those reported on databases such as VISTA Enhancer Browser (Kosicki et al., 2025) and

GeneHancer (Fishilevich et al., 2017; Fig. 6F). Overall, these experiments support that SALL4 is

required to recruit BAF at CNCC enhancers.

To further confirm that these regions are enhancers, we leveraged publicly available H3K4me1

and H3K4me3 ChIP-seq datasets generated using a comparable protocol at our Day-5 equivalent

(Prescott et al., 2015; Rada-Iglesias et al., 2011). Moreover, we performed ChIP-seq for H3K27ac

at Day-5, Day-11 and Day-14, roughly corresponding to CNCC-induction, early CNCC, late CNCC

stages. This analysis revealed that nearly all the sites bound by SALL4 and BRG1 are marked by

the enhancer mark H3K4me1, and only a small subset of them has the promoter mark H3K4me3

(Fig. S6A). Moreover, nearly all these sites have the active enhancer mark H3K27ac at Day-5,

and the acetylation levels progressively increase from CNCC induction to CNCC stage,

supporting that these are *bona fide* CNCC enhancers that increase in activation after SALL4

binding (Figs. S6B, C)

While a large portion of BRG1 peaks was lost in the absence of SALL4, we also identified a subset

of novel BRG1 peaks that were absent in *SALL4*-WT cells but gained in the *SALL4*-het-KO cells.

These BRG1 peaks gained in the *SALL4*-het-KO cells were associated with genes linked to

neurogenesis and nervous system development (Fig. S7A and Table S4). In contrast, the genes

located near BRG1 peaks lost in the *SALL4*-het-KO cells were involved in embryonic organ

development, epithelial differentiation, and cell migration processes (Fig. S7B and Table S4).

Together, these results suggest that SALL4 is a regulator of BAF recruitment to CNCC enhancers,

ensuring their accessibility during differentiation. In the absence of SALL4, BRG1 fails to localize

to these sites and is partially redirected to neurodevelopmental enhancers, leading to a

misallocation of chromatin remodeling, and to a shift towards default neural differentiation at the

expense of CNCC induction and specification.

SALL4-het-KO cells are not able to exit the neural ectoderm stage and to induce the cranial

neural crest fate

Our data thus far indicate that SALL4 relocates to CNCC enhancers as early as the

neuroectodermal stage, recruiting BAF to these regulatory elements for chromatin activation.

Next, we investigated the functional consequences of SALL4 loss by generating CNCCs using

SALL4-WT and *SALL4*-het-KO cells. As expected, *SALL4*-WT CNCCs expressed the signature

CNCC markers at both the mRNA (Fig. 7A) and protein (Figs. 7B and S4B) level, while *SALL4*-

het-KO failed to induce these genes (Fig. 7A). This was further supported by RNA-seq conducted

at Day-14, which identified 1,106 genes differentially expressed in *SALL4*-het-KO CNCCs relative

to *SALL4*-WT (FDR < 0.05; Fig. 7C and Table S3). This was in sharp contrast with the small

number of genes differentially expressed at the iPSC stage (n = 114; Fig. 3C and Table S3) and

at Day-5 (n = 170; Fig. S7C; Table S3).

Of the 1,106 genes differentially expressed at Day-14, ~60% (n = 659) were downregulated while

the remaining were upregulated (Fig. 7C). Canonical CNCC markers such as *TFAP2A*, *NR2F1*,

and *SOX9* were significantly downregulated in *SALL4*-het-KO cells. Additionally, genes

associated with neural plate border, early neural crest induction and epithelial-to-mesenchymal

transition (EMT) were also downregulated (Fig. 7D, F). In line with this, gene ontology analysis

revealed that pathways associated with face development, autonomic nervous system,

gastrulation, and regulation of morphogenesis were all downregulated (Fig. 7D). The 447

upregulated genes were instead involved in cell regulatory pathways (Fig. 7E).

As aforementioned, genes associated with EMT were dysregulated. The process of EMT requires

a switch in cadherin expression where class-1 cadherins must be downregulated, while class-2

cadherins must be upregulated. Related to this, our RNA-seq data revealed that this cadherin

switch did not occur in the differentiating *SALL4*-het-KO iPSCs, suggesting that the cells remained

in an epithelial state. This was further corroborated by the upregulation of several epithelial genes

in the differentiating *SALL4*-het-KO cells, including *EpCAM*, *CDH1*, and *OVOL2* (Fig. 7F). The

latter is involved in maintaining epithelial lineages by repressing the expression of the EMT gene

ZEB1, which was downregulated in the *SALL4*-het-KO condition (Haensel et al., 2019; Liu et al.,

2018a).

In summary, our data indicate that CNCC specification failed in *SALL4*-het-KO conditions, as

reflected by the large number of differentially expressed genes. On the other hand, the

transcriptomes of iPSCs and neuroectodermal cells (i.e. Day-5) were only modestly affected by

SALL4 loss, suggesting that SALL4 may not be essential for neuroectoderm formation, while it is

essential for CNCC induction and specification.

To investigate this further, we systematically examined the expression of a panel of stage-specific

marker genes at Day-0, Day-5 and Day-14 (Fig. S8A–D) including core pluripotency genes

(*POU5F1* and *NANOG*), neural plate border specifiers (*ZIC1, GBX2, MSX1/2, PAX3/7, GATA2/3*),

CNCC specification markers (*SNAI1/2, SOX9, TFAP2A*), and post-migratory CNCC markers

(*SOX5, SOX6, SOX10*). In *SALL4*-WT cells, pluripotency markers exhibited a progressive and

expected downregulation as differentiation advanced from Day-0 to Day-14. In *SALL4*-het-KO
cells, a similar decline in pluripotency markers was observed between Day-0 and Day-5, but
instead of continuing to decrease, these markers remained similarly expressed through Day-14
(Fig. S8A). Despite this, the two conditions showed similar trends, confirming that decreased
levels of SALL4 did not significantly impact pluripotency markers expression during exit from
pluripotency. Similarly, no significant difference was observed in the expression of neural plate
border specifiers (Fig. S8B), suggesting that SALL4 is not required for human neuroectoderm
specification. Conversely, CNCC specification markers and post-migratory CNCC markers all
increased in expression after Day-5 in *SALL4*-WT cells, while the SALL4-depleted cells failed to
maintain a comparable increase in expression (Fig. S8C, D).
In summary, *SALL4*-het-KO cells are competent to generate neuroectoderm-like cells but fail to
induce the neural crest lineage.
351

SALL4A but not SALL4B is necessary for CNCC specification
Given previous reports suggesting distinct roles for the two SALL4 isoforms in ESCs (Rao et al.,
2010), and our prediction that the interaction between SALL4 and DPF2 may be isoform-specific,
we investigated whether CNCC differentiation exhibits preferential dependence on a particular
SALL4 isoform.
Time-course mRNA expression in *SALL4*-WT cells revealed that *SALL4B* is consistently lowly
expressed throughout differentiation, whereas *SALL4A* expression increases upon differentiation
cues, peaking around Day-5 and then declining over time (Figs. S9A, B). On the other hand, the
paralog SALL1 is expressed at low levels in iPSCs and barely detectable throughout the entire
differentiation protocol (Fig. S9C).
To investigate potential isoform-specific requirements for CNCC development, we selectively
degraded SALL4A while preserving SALL4B expression. To achieve this, we treated cells with
pomalidomide, a thalidomide derivative that selectively degrades SALL4A but not SALL4B (Anh
Vu et al., 2023; Liu et al., 2024; Fig. 8A).
To test the effects of SALL4A degradation, we differentiated *SALL4*-WT iPSCs into CNCCs. Cells
were treated with either DMSO (control) or 5 µM pomalidomide (5 µM POM) every 24 hours (Fig.
8B). To confirm consistent degradation of SALL4A throughout the differentiation process, we
performed immunoblotting at key time points (Day-5 and Day-14). As expected, SALL4A was
successfully degraded, while SALL4B expression remained intact (Fig. 8C). Importantly,
pomalidomide degrades several proteins beyond SALL4, and its protein degradation activity is
typically cell-type specific (Sievers et al., 2018). Two of the known pomalidomide targets have

suggested functions in neural crest development. These are ARID2 (Bajpai et al., 2010;
Yamamoto et al., 2020) and ZMYM2 (Jourdeuil et al., 2025; Renneville et al., 2021). Crucially,
immunoblot analysis for these two proteins performed at Day 5 following treatment with 5 µM
pomalidomide revealed that the levels of both ARID2 and ZMYM2 remained unchanged
compared to the DMSO control (Figs. S9D, E), further supporting that pomalidomide affinity for
its targets is cell-type specific, as previously suggested (Bajpai et al., 2010; Yamamoto et al.,
2020). This control experiments allowed to rule out potential off-target effects caused by unwanted
degradation of one of these two proteins. The other known pomalidomide targets, including
IKAROS, AIOLOS, PLZF and ZNFX1 have no established contribution to neural crest
specification and in fact they are not highly expressed in our system.
Next, we assessed the downstream effects of SALL4A degradation by performing RT-qPCR and
immunofluorescence for key neural crest markers, such as *TFAP2A*, *NR2F1*, and *SOX9* (Figs.
8D, E), and EMT markers VIM and EpCAM (Fig. 8F). Degradation of SALL4A recapitulated what
we previously observed in *SALL4*-het-KO cells. Specifically, the cells treated with pomalidomide
failed to differentiate successfully into CNCCs despite the persistent expression of SALL4B.
Taken together, our results highlight a specific and essential role for SALL4A in orchestrating
chromatin dynamics during CNCC specification and suggest that SALL4B might function in a
different regulatory capacity rather than acting as a compensatory factor for SALL4A.

391
392

DISCUSSION
Mutations in *SALL4* are associated with congenital syndromes which frequently include distinct
craniofacial anomalies(Kohlhase et al., 2005a; Kohlhase et al., 2005b; Miller, 1991), suggesting
disrupted CNCC development.
Notably, at the murine embryonic stage E9.5, Sall4 expression persists within the frontonasal
prominence (Tahara et al., 2019), implying a potential role for SALL4 in craniofacial development.
Given the limited existing knowledge about SALL4 function in this context, we utilized a
heterozygous *SALL4* knockout hiPSC line to investigate its role CNCC specification.
At early stages of CNCC induction, SALL4 relocates to enhancers of genes involved in CNCC
formation, and SALL4 binding at these regulatory elements is required for their activation. In
mESCs, approximately 10% of the total Sall4 protein, and primarily the Sall4A isoform, is
incorporated into the NuRD complex (Bode et al., 2016; Miller et al., 2016). Here, we identify and
characterize a novel interaction between SALL4 and the BAF complex, predicted to be mediated
specifically through the SALL4A isoform and the BAF subunit DPF2. Interestingly, DPF2 and

SALL4 exhibit notable functional parallels. Similar to SALL4, DPF2 occupancy undergoes
significant repositioning during the transition from pluripotency toward early neural commitment
in mouse embryoid bodies (Zhang et al., 2019b). Although the role of DPF2 in CNCCs has not
been directly investigated, its early recruitment to neuronal enhancers suggests a potential
involvement in neuroectodermal differentiation and lineage commitment of neuroectoderm-
derived cells, including CNCCs. Moreover, genomic sequences bound by DPF2 in mouse
neuroectoderm are AT-rich, resembling the preferred binding motif of SALL4 (Kong et al., 2021;
Nazim et al., 2024; Pantier et al., 2021). It is important to note that AlphaFold3 is built on deep
learning rather than reinforcement learning. Consequently, the absence of more comprehensive
experimental SALL4 data could significantly affect the reliability of the proposed SALL4–DPF2
interaction model. Additionally, AlphaFold 5,000-token limit per prediction poses a constraint
when modeling large multi-subunit complexes such as BAF. Fortunately, BAF has been
extensively characterized, and our predicted model match experimentally resolved structures at
3 Å resolution.

Our study demonstrated that while CNCC induction fails without SALL4, early neuroectodermal
commitment and neural plate border identity remain unaffected. Our findings demonstrate that,
during the earliest stages of CNCC specification, SALL4 occupies CNCC-associated enhancers
and recruits BAF. Together, these observations support previous studies that proposed that
neural crest cell fate commitment is initiated well before overt gene expression changes occur
(Basch et al., 2006; Patthey et al., 2008; Stuhlmiller and García-Castro, 2012). This finding is
further supported by clinical observations. In fact, while phenotypes arising from *SALL4*
haploinsufficiency affect many neural crest derivatives, phenotypes implicating defects in
neurodevelopment are exceptionally rare (Borozdin et al., 2004; Ma et al., 2022) and usually
associated with dysregulation of potassium channels (Kodytková et al., 2023). In the rare cases
of intellectual disability reported, such phenotypes were attributed to broader deletions on
chromosome 20q13.13-q13.2 rather than *SALL4* loss alone (Borozdin et al., 2004).

In the absence of SALL4, after two weeks of differentiation the cells acquire an aberrant fate,
expressing a mix of pluripotency genes (suggesting lack of cell differentiation), and neuronal
genes (suggesting derailed cell fate). This indicates that most cells safely reach the
neuroectodermal stage but fail to go beyond that, while some of the cells simply do not
differentiate or regress back to pluripotency.

Interestingly, only a small minority of the CNCC enhancers bound by SALL4 has the AT-rich motif
that is thought to be recognized by SALL4 (Kong et al., 2021; Pantier et al., 2021). This suggests
that SALL4 might bind these enhancers in cooperation with other lineage-specific factors. SALL1

also recognizes AT-rich motifs, but similar to SALL4 it also binds unspecifically. Since SALL1 is upregulated in the *SALL4*-het-KO cells, it might possible that some of the sites associated with a gain of BRG1 upon loss of SALL4 could be associated with SALL1 being upregulated and hijacking BAF recruitment to distinct sites.

Limitations

SALL4 was recently identified as a direct target of thalidomide (Donovan KA, 2018), a drug widely prescribed to pregnant women in the late 1950s to alleviate morning sickness. The subsequent alarming rise in births of children with congenital anomalies, including limb malformations, facial abnormalities such as inner ear defects, ocular motility issues, and facial nerve palsy, as well as kidney and heart defects, led to the identification of thalidomide as the causative agent. However, the molecular mechanism by which thalidomide induced these developmental defects remained elusive until nearly six decades later. Building on earlier findings that thalidomide and its derivatives bind to cereblon (CRBN), Matyskiela et al. demonstrated that SALL4 is recruited to the CRL4–CRBN E3 ubiquitin ligase complex following thalidomide-induced ubiquitination, resulting in its proteasomal degradation (Matyskiela et al., 2018). More recently, research demonstrated that only the SALL4A isoform, but not SALL4B, is susceptible to degradation by immunomodulatory imide drugs such as pomalidomide (Anh Vu et al., 2023). Our data suggests that SALL4 role in CNCC specification might be isoform-specific and these findings might contribute to shed light on the birth malformations caused by thalidomide. However, while we ensured that known pomalidomide targets important for CNCC developments were not degraded in our system, we cannot rule out degradation of previously uncharacterized and unknown pomalidomide targets with key roles in CNCC specification. While highly unlikely, this is an important limitation of our study, and therefore the findings of our pomalidomide experiments, should be taken with caution.

ACKNOWLEDGMENTS

We thank the Facility for Imaging by Light Microscopy (FILM) at Imperial College London for providing access to instrumentation and technical support, which is part-supported by funding from the Wellcome Trust (grant 104931/Z/14/Z) and BBSRC (grant BB/L015129/1). We thank Prof. Brian Hendrich (University of Cambridge) and Dr. Andria Koulle (Imperial College London) for insightful discussion of the data and of early versions of the manuscript.

FUNDING

For this work, MT was funded by the Mathers Foundation and by Biotechnology and Biological Sciences Research Council (BBSRC, grant BB/Y000854/1).

COMPETING INTERESTS

None declared.

AUTHOR CONTRIBUTIONS

MT designed the project. MD performed most of the experiments. SMB, EVD, ZHM, and LD contributed to some of the experiments. MD and MT analyzed the data and wrote the manuscript. All the authors read and approved the manuscript.

FIGURE LEGENDS

Figure 1. SALL4A interacts with the BAF complex through DPF2.

(A) Immunoblot of nuclear extract (5% input), co-immunoprecipitations (co-IP) of ARID1B, and IgG (negative control) from *SALL4*-WT cells at Day-5.

(B) AlphaFold structure model showing the predicted interactions between SALL4A (red), DPF2 (dark blue), and other BAF components, including SMARCA4, SMARCB1, and ACTL6A.

(C) SALL4A protein sequence and visual representation of output model of SALL4A (model 0). C2H2 zinc-fingers annotations are highlighted in dark purple and numbered. Interactions between SALL4A and DPF2 through the ZFs 4, 5 and 6 (ZFC3) and 7 (ZFC4) are highlighted in light purple. Made with BioRender.com.

(D) Zoomed view of SALL4 (pink) and DPF2 (purple) interacting surfaces. Highlighted are the functional domains involved in the predicted interaction. In red, contacts predicted by AlphaFold(Jumper et al., 2021).

(E) Box plot displaying expression (TPMs) of the four members of the DPF protein family in iPSCs (n = 2). Differences between conditions were assessed using unpaired student's t-test.

(F) Box plot displaying expression (TPMs) of the four members of the DPF protein at Day-5 (n = 3). Differences between conditions were assessed using unpaired student's t-test.

Figure 2. Generation of a *SALL4*-haploinsufficient human iPSC line

(A) Visual representation of the CRISPR/Cas9 approach used on for *SALL4*-het-KO. The star represents the exact location of the deletion. Sanger sequencing confirming the 11-bp deletion in one allele of *SALL4* is displayed (clone F1). Made with BioRender.com.

(B) Immunoblot for SALL4 in *SALL4*-WT and *SALL4*-het-KO (clone F1) iPSC lines.

(C) Bar plots showing densitometry analysis of SALL4A and SALL4B expression in WT and *SALL4*-het-KO iPSC lines from immunoblot assay.

(D) Bar plots showing RNA expression in TPMs for SALL4A and SALL4B in WT and *SALL4*-het-KO iPSCs. Individual biological replicates (n = 2) are displayed as circles. Statistical significance was assessed using unpaired student's t-test. Error bars display standard error.

(E) Bar plots displaying expression TPMs of all *SALL* paralogues in *SALL4*-WT and *SALL4*-het-KO iPSCs. Individual biological replicates (n = 2) are displayed as circles. Statistical significance was assessed using unpaired student's t-test. Error bars display standard error of the mean.

(F) Immunoblot for SALL1 in *SALL4*-WT and *SALL4*-het-KO (clone F1) iPSC lines.

(G) Growth curve comparing *SALL4*-WT with *SALL4*-het-KO iPSC lines. N = 2 biologically independent experiments. Error bars display standard error.

Figure 3. Stem cell identity is unaffected by SALL4 loss

(A) (Left) Immunofluorescence labeling quantifying the expression of the canonical pluripotency markers OCT4 and NANOG in *SALL4*-WT and *SALL4*-het-KO (clone F1) iPSCs. Scale bar: 100 μm. (Right) Corrected Total Cell Fluorescence (CTCF) values were used to quantify and compare markers expression (OCT4 n=34; NANOG n=37). Error bars display standard error.

(B) Flow cytometry quantifying the expression of surface markers for pluripotency (TRA-60-1-R and SSEA-4).

(C) Volcano plot of genes differentially expressed in *SALL4*-het-KO relative to *SALL4*-WT in iPSCs (Day-0). Blue dots (n = 50) represent downregulated genes with p-adj < 0.05 and log2FoldChange < -1.5. Purple dots (n = 63) represent upregulated genes with p-adj < 0.05 and log2FoldChange > 1.5.

Figure 4. SALL4 relocates to CNCC enhancers to induce chromatin accessibility

(A) (Bottom) Schematic representation of the iPSC-to-CNCC differentiation protocol. The experimental timeline shows in vitro differentiation of iPSCs into CNCCs via a neuroectodermal-like stage. Immunofluorescence labeling quantifying the expression of the canonical stage-specific markers (iPSCs: OCT4; neuroectoderm: OTX2; CNCCs: SOX9) showing ratio of cells successfully undergoing stage-specific specifications during CNCC generation. Scale bar: 100 μm.

(B) Diagram showing the number of regions bound by SALL4 uniquely at Day-0 ("Day-0 specific"; n = 6,114), at Day-5 ("Day-5 specific"; n = 17,137), and regions that maintain SALL4 binding at both time points ("shared", n = 15,570).

(C) Heatmaps showing the distribution of SALL4 binding at Day-0 and Day-5 across Day-0 specific (left) and Day-5 specific (right) binding regions. CUT&RUN signal intensity was plotted across ±2 kb regions centered on peak summits. Rows represent individual peaks ranked by signal intensity.

(D) Genomic distribution of SALL4 binding sites relative to the nearest TSS. Peaks are categorized by distance into six bins: <1 kb, 1–3 kb, 3–5 kb, 5–10 kb, 10–100 kb, and >100 kb. Peaks located at >1kb from the nearest TSS were classified as putative enhancers. Percentages of peaks for each bin are represented by bars to the left (downstream) or right (upstream) of the TSS.

(E) Gene ontology analysis of downregulated nearest genes to Day-5 specific SALL4-bound regions as determined using WebGestalt (Liao et al., 2019) over-representation pathway analysis, using affinity propagation as parameter for redundancy removal (FDR < 0.05).

(F) Table of motifs enriched in Day-5 specific SALL4-bound regions as determined using HOMER (Heinz et al., 2010).

Figure 5. Chromatin accessibility at the CNCC enhancers is SALL4 dependent

(A) Venn diagram showing chromatin accessibility in *SALL4*-WT (ATAC-seq peaks n = 58,375) and *SALL4*-het-KO (n = 60,186). It also depicts the number of regions uniquely accessible in *SALL4*-WT ("Sites losing accessibility in KO"; n = 1,649), in *SALL4*-het-KO ("Sites gaining accessibility in KO"; n = 3,460), and regions that remain accessible in both conditions ("Sites maintaining accessibility in WT and KO", n = 60,186).

(B) Heatmaps showing ATAC-seq signals at Day-0 (left) and Day-5 (right), at Day-5 specific SALL4 CUT&RUN peaks. Signal intensity was plotted across ±2 kb regions centered on peak summits. Rows represent individual peaks ranked by signal intensity.

(C) Average profiles showing ATAC-seq signal at Day 5-specific SALL4 CUT&RUN peaks (n =
17,137) in *SALL4*-WT and *SALL4*-het-KO cells in two distinct biological replicates.
(D) Table of motifs enriched in regions gaining or maintaining accessibility at Day-5 in *SALL4*-het-
KO regions as determined using HOMER (Heinz et al., 2010).
(E) Table of motifs enriched in regions losing accessibility at Day-5 in *SALL4*-het-KO regions as
determined using HOMER (Heinz et al., 2010).
577

Figure 6. SALL4 recruits BAF to CNCC enhancers.
(A) Heatmap showing BRG1 ChIP-seq signal at the genomic sites bound by SALL4 in WT cells
at Day-5 (n = 17,137). Signal intensity was plotted across ± 2 kb regions centered on peak
summits. Rows represent individual peaks ranked by signal intensity.
(B) Venn diagram showing peaks bound by SALL4 uniquely at Day-5 and that lose accessibility
in *SALL4*-het-KO.
(C) Heatmap and average profiles show BRG1 ChIP-seq signal at Day-5 at the BRG1 peaks lost
in *SALL4*-het-KO (clone F1; n = 13,505). Signal intensity was plotted across ±2 kb regions
centered on peak summits. Rows represent individual peaks ranked by signal intensity.
(D) Heatmap and average profiles show SALL4 CUT&RUN signal at the genomic sites that lose
BRG1 in *SALL4*-het-KO cells at Day-0 and Day-5 (n = 13,505).
(E) Heatmaps and average profile show ATAC-seq signal at the genomic sites that lose BRG1 in
SALL4-het-KO cells (clone F1; n = 13,505). Signal intensity was plotted across ±2 kb regions
centered on peak summits. Rows represent individual peaks ranked by signal intensity.
(F) Two representative examples of enhancers bound by SALL4 at Day 5 that lose BRG1 co-
occupancy and chromatin accessibility in the *SALL4*-het-KO (clone F1) cells. Both correspond to
ChIP-seq peaks annotated as enhancer regions, as reported in GeneHancer (left) and VISTA
Enhancer Browser (right).
596

Figure 7. CNCC induction and specification are impaired in *SALL4*-het-KO.
(A) Bar plot displaying relative mRNA expression of canonical CNCC markers in *SALL4*-WT and
SALL4-het-KO cells at Day-14 (n = 2). Statistical significance was assessed using unpaired
student's t-test. 18S rRNA was used to normalize.
(B) Flow cytometry for the CNCC surface marker CD99 in *SALL4*-WT and *SALL4*-het-KO (clone
F1) CNCCs at Day-14.
(C) Volcano plot of genes differentially expressed in *SALL4*-het-KO relative to *SALL4*-WT in
CNCCs (Day-14). Blue dots (n= 659) represent downregulated genes with p-adj < 0.05 and

log2FoldChange < -1.5. Purple dots (n= 447) represent upregulated genes with p-adj < 0.05 and

log2FoldChange > 1.5.

(D) and (E) GO term enrichment for upregulated (D) and downregulated (E) genes in *SALL4*-het-

KO cells (FDR < 0.05).

(F) Heatmap displaying the expression of canonical epithelial-to-mesenchymal transition (EMT)

markers in *SALL4*-WT and *SALL4*-het-KO cells (Day-14).

Figure 8. SALL4A is the only SALL4 isoform necessary for CNCC specification

(A) Schematic of pomalidomide-mediated ubiquitin ligation. Ub, ubiquitin. POM, pomalidomide.

CRBN, DDB1, Cul4 and Rbx1 are subunits of CRL4/CRBN E3 ubiquitin ligase.

(B) Schematic representation of pomalidomide treatment during iPSC-to-CNCC differentiation

protocol.

(C) Time-course immunoblot for SALL4 in *SALL4*-WT at Day-5 and Day-14 after daily treatment

with DMSO (control) or 5μM pomalidomide (5μM POM).

(D) Box plot displaying the relative mRNA expression of canonical CNCC markers (SOX9,

TFAP2A, and NR2F1) in *SALL4*-WT at Day-14 treated with 5μM pomalidomide or DMSO (control)

daily for 14 days. Individual biological replicates (n = 2) are displayed as circles. Statistical

significance was assessed using unpaired student's t-test. Error bars display standard error.

(E) Immunofluorescence labeling quantifying the expression of the canonical CNCC markers

SOX9 and TFAP2A in *SALL4*-WT treated with 5μM pomalidomide or DMSO (control) daily for 14

days. Scale bar: 100 μm.

(F) Immunofluorescence labeling quantifying the expression of the canonical EMT markers

EpCAM and VIM in *SALL4*-WT treated with 5uM pomalidomide or DMSO (control) daily for 14

days. Scale bar: 100 μm.

REFERENCES

Al-Baradie, R., Yamada, K., St Hilaire, C., Chan, W.-M., Andrews, C., McIntosh, N., Nakano, M., Jean
Martonyi, E., Raymond, W. R., Okumura, S., et al. (2002). *Duane Radial Ray Syndrome (Okihiro*
Syndrome) Maps to 20q13 and Results from Mutations in SALL4, a New Member of the SAL Family.

Anh Vu, K. L., Kumari, K., Hui Liu, B., Gao, C., Li, F., Ping Tang, J., Maddalo, D., Auld, D. S., Casalena, D.
E., Tian, X., et al. (2023). SALL4B, not targeted by IMiD, is important for SALL4-mediated
tumorigenesis.

Bajpai, R., Chen, D. A., Rada-Iglesias, A., Zhang, J., Xiong, Y., Helms, J., Chang, C. P., Zhao, Y., Swigut, T.
and Wysocka, J. (2010). CHD7 cooperates with PBAF to control multipotent neural crest formation.
Nature **463**, 958–962.

Barnada, S. M., Giner de Gracia, A., Morenilla-Palao, C., López-Cascales, M. T., Scopa, C., Waltrich, F.
J., Mikkers, H. M. M., Cicardi, M. E., Karlin, J., Trotti, D., et al. (2024). ARID1A-BAF coordinates
ZIC2 genomic occupancy for epithelial-to-mesenchymal transition in cranial neural crest
specification. *The American Journal of Human Genetics*.

Basch, M. L., Bronner-Fraser, M. and García-Castro, M. I. (2006). Specification of the neural crest occurs
during gastrulation and requires Pax7. *Nature* **441**, 218–222.

Bode, D., Yu, L., Tate, P., Pardo, M. and Choudhary, J. (2016). Characterization of two distinct
nucleosome remodeling and deacetylase (NuRD) complex assemblies in embryonic stem cells.
Molecular and Cellular Proteomics **15**, 878–891.

Borozdin, W., Boehm, D., Leipoldt, M., Wilhelm, C., Reardon, W., Clayton-Smith, J., Becker, K.,
Mühlendyck, H., Winter, R., Giray, O., et al. (2004). SALL4 deletions are a common cause of
Okihiro and acro-renal-ocular syndromes and confirm haploinsufficiency as the pathogenic
mechanism. *J Med Genet* **41**,.

Chen, K. Q., Anderson, A., Kawakami, H., Kim, J., Barrett, J. and Kawakami, Y. (2022). Normal
embryonic development and neonatal digit regeneration in mice overexpressing a stem cell factor,
Sall4. *PLoS One* **17**,.

Dash, S. and Trainor, P. A. (2020). The development, patterning and evolution of neural crest cell
differentiation into cartilage and bone. *Bone* **137**,.

Donovan KA, A. J. N. R. Y. J. F. E. B. B. E. B. F. E. (2018). Thalidomide promotes degradation of SALL4, a
transcription factor implicated in Duane Radial Ray syndrome. *Elife*.

Fishilevich, S., Nudel, R., Rappaport, N., Hadar, R., Plaschkes, I., Stein, T. I., Rosen, N., Kohn, A., Twik,
M., Safran, M., et al. (2017). GeneHancer: Genome-wide integration of enhancers and target
genes in GeneCards. *Database* **2017**,.

Gopinathan, G. and Diekwisch, T. G. H. (2022). Epigenetics and Early Development. *J Dev Biol* **10**,.

Haensel, D., Sun, P., MacLean, A. L., Ma, X., Zhou, Y., Stemmler, M. P., Brabletz, S., Berx, G., Plikus, M.
V, Nie, Q., et al. (2019). An Ovol2-Zeb1 transcriptional circuit regulates epithelial directional
migration and proliferation. *EMBO Rep* **20**,.

Heinz, S., Benner, C., Spann, N., Bertolino, E., Lin, Y. C., Laslo, P., Cheng, J. X., Murre, C., Singh, H. and
Glass, C. K. (2010). Simple Combinations of Lineage-Determining Transcription Factors Prime cis-
Regulatory Elements Required for Macrophage and B Cell Identities. *Mol Cell* **38**, 576–589.

Herbreteau, G., Airaud, F., Pierre-Noël, E., Vallée, A., Bézieau, S., Théoleyre, S., Blons, H., Garinet, S.
and Denis, M. G. (2021). MEM: An algorithm for the reliable detection of microsatellite instability
(MSI) on a small NGS panel in colorectal cancer. *Cancers (Basel)* **13**,.

Hodges, C., Kirkland, J. G. and Crabtree, G. R. (2016). The many roles of BAF (mSWI/SNF) and PBAF
complexes in cancer. *Cold Spring Harb Perspect Med* **6**,.

Islam, Z., Ali, A. M., Naik, A., Eldaw, M., Decock, J. and Kolatkar, P. R. (2021). Transcription Factors: The
Fulcrum Between Cell Development and Carcinogenesis. *Front Oncol* **11**,.

Jourdeuil, K., Neilson, K. M., Tavares, A. L. P. and Moody, S. A. (2025). Zmym2 Alters Expression of Key
Craniofacial Genes. *Genesis (United States)* **63**,.

Jumper, J., Evans, R., Pritzel, A., Green, T., Figurnov, M., Ronneberger, O., Tunyasuvunakool, K., Bates,
R., Žídek, A., Potapenko, A., et al. (2021). Highly accurate protein structure prediction with
AlphaFold. *Nature* **596**, 583–589.

Kekunnaya, R. and Negalur, M. (2017). Duane retraction syndrome: Causes, effects and management
strategies. *Clinical Ophthalmology* **11**, 1917–1930.

Kim, S. and Shendure, J. (2019). Mechanisms of Interplay between Transcription Factors and the 3D
Genome. *Mol Cell* **76**, 306–319.

Kodytková, A., Dušátková, P., Anne Amaratunga, S., Plachý, L., Pruhová, Š. and Lebl, J. (2023).
Integrative role of the sall4 gene: From thalidomide embryopathy to genetic defects of the upper
limb, internal organs, cerebral midline, and pituitary. *Horm Res Paediatr* **97**, 106–112.

Kohlhase, J., Chitayat, D., Kotzot, D., Ceylaner, S., Froster, U. G., Fuchs, S., Montgomery, T. and Rösler,
B. (2005a). SALL4 mutations in Okihiro syndrome (Duane-radial ray syndrome), acro-renal-ocular
syndrome, and related disorders. *Hum Mutat* **26**, 176–183.

Kohlhase, J., Chitayat, D., Kotzot, D., Ceylaner, S., Froster, U. G., Fuchs, S., Montgomery, T. and Rösler,
B. (2005b). SALL4 mutations in Okihiro syndrome (Duane-radial ray syndrome), acro-renal-ocular
syndrome, and related disorders. *Hum Mutat* **26**, 176–183.

Kong, N. R., Bassal, M. A., Tan, H. K., Kurland, J. V., Yong, K. J., Young, J. J., Yang, Y., Li, F., Lee, J. D.,
Liu, Y., et al. (2021). Zinc Finger Protein SALL4 Functions through an AT-Rich Motif to Regulate
Gene Expression. *Cell Rep* **34**,.

Kosicki, M., Baltoumas, F. A., Kelman, G., Boverhof, J., Ong, Y., Cook, L. E., Dickel, D. E., Pavlopoulos,
G. A., Pennacchio, L. A. and Visel, A. (2025). VISTA Enhancer browser: an updated database of
tissue-specific developmental enhancers. *Nucleic Acids Res* **53**, D324–D330.

Krasteva, V., Crabtree, G. R. and Lessard, J. A. (2017). The BAF45a/PHF10 subunit of SWI/SNF-like
chromatin remodeling complexes is essential for hematopoietic stem cell maintenance. *Exp*
Hematol **48**, 58-71.e15.

Lessard, J., Wu, J. I., Ranish, J. A., Wan, M., Winslow, M. M., Staahl, B. T., Wu, H., Aebersold, R., Graef,
I. A. and Crabtree, G. R. (2007). An Essential Switch in Subunit Composition of a Chromatin
Remodeling Complex during Neural Development. *Neuron* **55**, 201–215.

Li, E. (2002). Chromatin modification and epigenetic reprogramming in mammalian development. *Nat*
Rev Genet **3**, 662–673.

Liao, Y., Wang, J., Jaehnig, E. J., Shi, Z. and Zhang, B. (2019). WebGestalt 2019: gene set analysis toolkit
with revamped UIs and APIs. *Nucleic Acids Res* **47**, W199–W205.

Liu, J., Wu, Q., Wang, Y., Wei, Y., Wu, H., Duan, L., Zhang, Q. and Wu, Y. (2018a). Ovol2 induces
mesenchymal–epithelial transition via targeting ZEB1 in osteosarcoma. *Onco Targets Ther* **11**,
2963–2973.

Liu, B. H., Jobichen, C., Chia, C. S. B., Chan, T. H. M., Tang, J. P., Chung, T. X. Y., Li, J., Poulsen, A., Hung,
A. W., Koh-Stenta, X., et al. (2018b). Targeting cancer addiction for SALL4 by shifting its
transcriptome with a pharmacologic peptide. *Proc Natl Acad Sci U S A* **115**, E7119–E7128.

Liu, B. H., Liu, M., Radhakrishnan, S., Jaladanki, C. K., Gao, C., Tang, J. P., Kumari, K., Go, M. L., Vu, K. A.
L., Seo, H.-S., et al. (2024). Targeting transcription factors through an IMiD independent zinc finger
domain.

Love, M. I., Huber, W. and Anders, S. (2014). Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2. *Genome Biol* **15**,.

Lu, J., Jeong, H., Kong, N., Yang, Y., Carroll, J., Luo, H. R., Silberstein, L. E., Ma, Y. and Chai, L. (2009).
Stem cell factor SALL4 represses the transcriptions of PTEN and SALL1 through an epigenetic
repressor complex. *PLoS One* **4**,.

Ma, X., Huang, R., Li, G., Zhang, T. and Ma, J. (2022). A de novo mutation of SALL4 in a Chinese family
with Okihiro syndrome. *Mol Med Rep* **25**,.

Matyskiela, M. E., Couto, S., Zheng, X., Lu, G., Hui, J., Stamp, K., Drew, C., Ren, Y., Wang, M.,
Carpenter, A., et al. (2018). SALL4 mediates teratogenicity as a thalidomide-dependent cereblon
substrate. *Nat Chem Biol* **14**, 981–987.

Milet, C. and Monsoro-Burq, A. H. (2012). Neural crest induction at the neural plate border in
vertebrates. *Dev Biol* **366**, 22–33.

Miller, M. T. (1991). Thalidomide embryopathy: a model for the study of congenital incomitant
horizontal strabismus. *Transaction of the American Ophtalmological Society* 623–674.

Miller, A., Ralser, M., Kloet, S. L., Loos, R., Nishinakamura, R., Bertone, P., Vermeulen, M. and
Hendrich, B. (2016). Sall4 controls differentiation of pluripotent cells independently of the
nucleosome remodelling and deacetylation (NuRD) complex. *Development (Cambridge)* **143**, 3074–
3084.

Mitchell, Z. H., den Hoed, J., Claassen, W., Demurtas, M., Deelen, L., Campeau, P. M., Liu, K., Fisher, S.
E. and Trizzino, M. (2025). CHD3 regulates BMP signalling response during cranial neural crest cell
specification.

Nazim, M., Lin, C. H., Feng, A. C., Xiao, W., Yeom, K. H., Li, M., Daly, A. E., Tan, X., Vu, H., Ernst, J., et al.
(2024). Alternative splicing of a chromatin modifier alters the transcriptional regulatory programs
of stem cell maintenance and neuronal differentiation. *Cell Stem Cell* **31**, 754-771.e6.

Pagliaroli, L., Porazzi, P., Curtis, A. T., Scopa, C., Mikkers, H. M. M., Freund, C., Daxinger, L., Deliard, S.,
Welsh, S. A., Offley, S., et al. (2021). Inability to switch from ARID1A-BAF to ARID1B-BAF impairs

exit from pluripotency and commitment towards neural crest formation in ARID1B-related
neurodevelopmental disorders. *Nat Commun* **12**,.

Pantier, R., Chhatbar, K., Quante, T., Skourti-Stathaki, K., Cholewa-Waclaw, J., Alston, G., Alexander-
Howden, B., Lee, H. Y., Cook, A. G., Spruijt, C. G., et al. (2021). SALL4 controls cell fate in response
to DNA base composition. *Mol Cell* **81**, 845-858.e8.

Pappas, M. P., Kawakami, H., Corcoran, D., Chen, K. Q., Scott, E. P., Wong, J., Gearhart, M. D.,
Nishinakamura, R., Nakagawa, Y. and Kawakami, Y. (2024). Sall4 regulates posterior trunk
mesoderm development by promoting mesodermal gene expression and repressing neural genes
in the mesoderm. *Development (Cambridge)* **151**,.

Patthey, C., Gunhaga, L. and Edlund, T. (2008). Early development of the central and peripheral nervous
systems is coordinated by Wnt and BMP signals. *PLoS One* **3**,.

Pla, P. and Monsoro-Burq, A. H. (2018). The neural border: Induction, specification and maturation of
the territory that generates neural crest cells. *Dev Biol* **444**, S36–S46.

Polit, L., Kerdivel, G., Gregoricchio, S., Esposito, M., Guillouf, C. and Boeva, V. (2021). CHIPIN: ChIP-seq
inter-sample normalization based on signal invariance across transcriptionally constant genes. *BMC*
Bioinformatics **22**,.

Prescott, S. L., Srinivasan, R., Marchetto, M. C., Grishina, I., Narvaiza, I., Selleri, L., Gage, F. H., Swigut,
T. and Wysocka, J. (2015). Enhancer Divergence and cis-Regulatory Evolution in the Human and
Chimp Neural Crest. *Cell* **163**, 68–83.

Quinlan, A. R. and Hall, I. M. (2010). BEDTools: A flexible suite of utilities for comparing genomic
features. *Bioinformatics* **26**, 841–842.

Rada-Iglesias, A., Bajpai, R., Swigut, T., Brugmann, S. A., Flynn, R. A. and Wysocka, J. (2011). A unique
chromatin signature uncovers early developmental enhancers in humans. *Nature* **470**, 279–285.

Ramírez, F., Ryan, D. P., Grüning, B., Bhardwaj, V., Kilpert, F., Richter, A. S., Heyne, S., Dündar, F. and
Manke, T. (2016). deepTools2: a next generation web server for deep-sequencing data analysis.
Nucleic Acids Res **44**, W160–W165.

Rao, S., Zhen, S., Roumiantsev, S., McDonald, L. T., Yuan, G.-C. and Orkin, S. H. (2010). Differential
Roles of Sall4 Isoforms in Embryonic Stem Cell Pluripotency. *Mol Cell Biol* **30**, 5364–5380.

Renneville, A., Gasser, J. A., Grinshpun, D. E., Jean Beltran, P. M., Udeshi, N. D., Matyskiela, M. E.,
Clayton, T., McConkey, M., Viswanathan, K., Tepper, A., et al. (2021). Avadomide Induces
Degradation of ZMYM2 Fusion Oncoproteins in Hematologic Malignancies. *Blood Cancer Discov* **2**,
251–265.

Rgen Kohlhase, J., Heinrich, M., Schubert, L., Liebers, M., Kispert, A., Laccone, F., Turnpenny, P.,
Winter, R. M. and Reardon, W. (2002). Okihiro syndrome is caused by SALL4 mutations. *Oxford*
University Press **11**, 2979–2987.

Roth, D. M., Bayona, F., Baddam, P. and Graf, D. (2021). Craniofacial Development: Neural Crest in
Molecular Embryology. *Head Neck Pathol* **15**,.

Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibisch, S., Rueden,
C., Saalfeld, S., Schmid, B., et al. (2012). Fiji: An open-source platform for biological-image analysis.
Nat Methods **9**, 676–682.

Sievers, Q. L., Petzold, G., Bunker, R. D., Renneville, A., Słabicki, M., Liddicoat, B. J., Abdulrahman, W.,
Mikkelsen, T., Ebert, B. L. and Thomä, N. H. (2018). Defining the human C2H2 zinc finger degrome
targeted by thalidomide analogs through CRBN. *Science (1979)* **362**,.

Stuhlmiller, T. J. and García-Castro, M. I. (2012). FGF/MAPK signaling is required in the gastrula epiblast
for avian neural crest induction. *Development* **139**, 289–300.

Szabó, A., Szabó, S. and Mayor, R. (2018). Mechanisms of Neural Crest Migration.

Tahara, N., Kawakami, H., Chen, K. Q., Anderson, A., Peterson, M. Y., Gong, W., Shah, P., Hayashi, S.,
Nishinakamura, R., Nakagawa, Y., et al. (2019). Sall4 regulates neuromesodermal progenitors and
their descendants during body elongation in mouse embryos. *Development (Cambridge)* **146**,.

Tatetsu, H., Kong, N. R., Chong, G., Amabile, G., Tenen, D. G. and Chai, L. (2016). SALL4, the missing link
between stem cells, development and cancer. *Gene* **584**, 111–119.

Terhal, P., Rösler, B. and Kohlhase, J. (2006). A family with features overlapping Okihiro syndrome,
hemifacial microsomia and isolated Duane anomaly caused by a novel SALL4 mutation. *Am J Med*
Genet **140 A**, 222–226.

Voss, T. C. and Hager, G. L. (2014). Dynamic regulation of transcriptional states by chromatin and
transcription factors. *Nat Rev Genet* **15**, 69–81.

Wang, F., Zhao, W., Kong, N., Cui, W. and Chai, L. (2014). The next new target in leukemia: The
embryonic stem cell gene SALL4. *Mol Cell Oncol* **1**,.

Wang, B., Li, C., Ming, J., Wu, L., Fang, S., Huang, Y., Lin, L., Liu, H., Kuang, J., Zhao, C., et al. (2023). The
NuRD complex cooperates with SALL4 to orchestrate reprogramming. *Nat Commun* **14**,.

Xiong, J., Todorova, D., Su, N. Y., Kim, J., Lee, P. J., Shen, Z., Briggs, S. P. and Xu, Y. (2015). Stemness
factor Sall4 is required for DNA damage response in embryonic stem cells. *Journal of Cell Biology*
208, 513–520.

Yamamoto, J., Suwa, T., Murase, Y., Tateno, S., Mizutome, H., Asatsuma-Okumura, T., Shimizu, N.,
Kishi, T., Momose, S., Kizaki, M., et al. (2020). ARID2 is a pomalidomide-dependent CRL4CRBN
substrate in multiple myeloma cells. *Nat Chem Biol* **16**, 1208–1217.

Yang, J. (2018). SALL4 as a transcriptional and epigenetic regulator in normal and leukemic
hematopoiesis. *Biomark Res* **6**,.

Yuri, S., Fujimura, S., Nimura, K., Takeda, N., Toyooka, Y., Fujimura, Y. I., Aburatani, H., Ura, K., Koseki,
H., Niwa, H., et al. (2009). Sall4 is essential for stabilization, but not for pluripotency, of embryonic
stem cells by repressing aberrant trophectoderm gene expression. *Stem Cells* **27**, 796–805.

Zhang, J., Tam, W. L., Tong, G. Q., Wu, Q., Chan, H. Y., Soh, B. S., Lou, Y., Yang, J., Ma, Y., Chai, L., et al.
(2006). Sall4 modulates embryonic stem cell pluripotency and early embryonic development by the
transcriptional regulation of Pou5f1. *Nat Cell Biol* **8**, 1114–1123.

Zhang, Y., Liu, T., Meyer, C. A., Eeckhoute, J., Johnson, D. S., Bernstein, B. E., Nussbaum, C., Myers, R.
M., Brown, M., Li, W., et al. (2008). Model-based analysis of ChIP-Seq (MACS). *Genome Biol* **9**,.

Zhang, W., Chronis, C., Chen, X., Zhang, H., Spalinskas, R., Pardo, M., Chen, L., Wu, G., Zhu, Z., Yu, Y., et
al. (2019a). The BAF and PRC2 Complex Subunits Dpf2 and Eed Antagonistically Converge on Tbx3
to Control ESC Differentiation. *Cell Stem Cell* **24**, 138-152.e8.

Zhang, Z., Zhan, X., Kim, B. and Wu, J. (2019b). A proteomic approach identifies SAFB-like transcription
modulator (SLTM) as a bidirectional regulator of GLI family zinc finger transcription factors. *Journal*
of Biological Chemistry **294**, 5549–5561.

Zhang, H., Wang, X., Li, J., Shi, R. and Ye, Y. (2021). BAF Complex in Embryonic Stem Cells and Early
Embryonic Development. *Stem Cells Int* **2021**,.

Zheng, Z. H., Sam, T. W., Zeng, Y., Chu, J. J. H. and Loh, Y. H. (2021). Chromatin Regulation in
Development: Current Understanding and Approaches. *Stem Cells Int* **2021**,.

832
833

834 **METHODS**

835 **REAGENT TABLE**

REAGENT	SOURCE	IDENTIFIER
Antibodies		
Brg1 (D1Q7F)	Cell Signaling Technology	AB_2728743
CD99 Antibody, anti-human, Vio® Bright B515, REAfinity™	Miltenyi Biotec	AB_2784331
GAPDH (D16H11) XP Rabbit mAb	Cell Signaling Technology	AB_10622025
Human Nanog Antibody	R&D Systems	AB_355097
Human/Mouse Oct-3/4 Antibody	R&D Systems	AB_354975
SALL1 Polyclonal Antibody	Thermo Fisher Scientific	AB_2900498
Sall4 (EE-30)	Santa Cruz Biotechnology	AB_1129262
Sall4 (D16H12) Rabbit mAb	Cell Signaling Technology	AB_10949321
APC anti-human SSEA-4	Biolegend	AB_2616818
PE anti-human TRA-1-60-R	Biolegend	AB_1279447
Rabbit IgG, monoclonal [EPR25A] Antibody	Abcam	AB_2687931
AP2 alpha Monoclonal Antibody (3B5)	Thermo Fisher Scientific	AB_2199412
CD326 (EpCAM) Monoclonal Antibody (1B7), eBioscience	Thermo Fisher Scientific	AB_795876
Anti-SOX9 antibody	Abcam	AB_2715497
Vimentin Polyclonal Antibody	Thermo Fisher Scientific	AB_2216267
PAX6 antibody	Abcam	AB_305110
ARID1B antibody	Abcam	AB_2243092
ARID2 Polyclonal Antibody	Bethyl Laboratories	AB_1731041
ZMYM2 Polyclonal Antibody	Thermo Fisher Scientific	AB_11152210
IRDye 800CW Goat anti-Rabbit IgG	LI-COR Biosciences	AB_621843

Goat polyclonal Secondary Antibody to Mouse (IRDye® 800CW)	LI-COR Biosciences	AB_10859231
Donkey anti-Rabbit IgG (H+L) Highly Cross-Adsorbed Secondary Antibody, Alexa Fluor™ Plus 488	Thermo Fisher Scientific	AB_2762833
Donkey anti-Mouse IgG (H+L) Highly Cross-Adsorbed Secondary Antibody, Alexa Fluor™ Plus 647	Thermo Fisher Scientific	AB_2762830
Goat anti-Chicken IgY (H+L) Secondary Antibody, Alexa Fluor™ 488	Thermo Fisher Scientific	AB_2534096
Critical commercial assays		
ATAC-Seq Kit	Active Motif	53150
Human Pluripotent Stem Cell Functional Identification Kit	R&D Systems	SC027B
Monarch® Total RNA Miniprep Kit	New England Biolabs	T2010S
Maxima First Strand cDNA Synthesis Kit for RT-qPCR	Thermo Fisher	K1641
NEBNext® Ultra™ II Directional RNA Library Prep Kit for Illumina®	New England Biolabs	E7760L
CUT&RUN Assay Kit	Cell Signaling Technology	86652
NEBNext® Ultra™ II DNA Library Prep Kit for Illumina®	Cell Signaling Technology	E7645L
Experimental models: Cell lines		
PENN123i-SV20	iPSC Core of the University of Pennsylvania	CVCL_EL23
PENN123i-SV20 *SALL4*-het-KO iPSC	Synthego	
Oligonucleotides		

18S	5'-ATACATGCCGACGGGCGCTG-3'	5'-AGGGCTGACCGGGTTGGTT-3'
TFAP2A	5'-GCCTCTCGCTCCTCAGCTCC-3'	5'-CGTTGGCAGCTTTACGTCTCCC-3'
NR2F1	5'-ATCGTGCTGTTCACGTCAGA-3'	5'-GCTCCTCACGTACTCCTCCA-3'
SOX9	5'-GTACCCGCACTTGCACAAC-3'	5'-TCTCGCTCTCGTTCAGAAGTC-3'
NANOG	5'-ATGCCTCACACGGAGACTGT-3'	5'-AAGTGGGTTGTTTGCCTTTG-3'
POU5F1	5'-TCGAGAACCGAGTGAGAGG-3'	5'-GAACCACACTCGGACCACA-3'
MSX1	5'-GAAGATGCGCTCGTCAAAGC-3'	5'-CTTACGGTTCGTCTTGTGTTTG-3'
PAX3	5'-ACAACGCCTGACGTGGAG-3'	5'-ATTTACTTCTCAGGATGCGGCT-3'
TWIST1	5'-GCCAGGTACATCGACTTCCTCT-3'	5'-TCCATCCTCCAGACCGAGAAGG-3'
Software and algorithms		
AlphaFold3	https://alphafoldserver.com/	SCR_025454
TrimGalore!	https://www.bioinformatics.babraham.ac.uk/projects/trim_galore/	SCR_011847

FlowJo Software version 10.9		SCR_008520
BEDTools		SCR_006646
Deeptools		SCR_016366
R Studio version 4.4.3		
GraphPad Prism 10.2.3	http://www.graphpad.com/faq/viewfaq.cfm?faq=1362	SCR_002798
DESeq2		SCR_015687

METHOD DETAILS

AlphaFold3 structure prediction

FASTA files for each BAF subunits and SALL4 isoforms were collected from the "Sequence & Isoforms" section of Uniprot (www.uniprot.org). The AT-rich DNA sequences was collected from an empirically reported crystal structure of SALL4 zinc finger cluster 4 bound to DNA on RCSB PDB (www.rcsb.org, 8A4I – DOI: https://doi.org/10.2210/pdb8A4I/pdb). FASTA sequences are listed in Table 2. AlphaFold outputs were downloaded and .cif files were analysed using UCSF ChimeraX-1.9 (https://www.rbvi.ucsf.edu/chimerax). Contacts were found using default parameters and center-center distance ≤ 8 Å.

Human iPSC culture

Wild-type iPSC lines were obtained from the iPSC Core of the University of Pennsylvania (Control line-1: SV20 line, male, age 43). iPSC lines were cultured in feeder-free, serum-free mTeSR Plus medium (STEMCELL Technologies) with 1% Penicillin-Streptomycin (5,000 U/mL; Gibco #15140122) on tissue culture plates coated overnight with Geltrex™ LDEV-Free hESC-qualified Reduced Growth Factor Basement Membrane Matrix (Fisher-Scientific). Cells were passaged ~1:8 at 60-80% confluency using 0.5mM EDTA to create 50-200 cell clusters.

CRISPR/Cas9-mediated SALL4 knock-out

The heterozygous *SALL4* knock-out in the human SV20 iPSC line was generated via CRISPR/Cas9 gene-editing by the biotech company Synthego. Specifically, two clones were used. In both clones Cas9 targeted the first exon of the *SALL4* gene in the established SV20 iPSC line producing an 11-base deletion (c.230delGGGCCGCATCT; clone C4, named KO1 in the manuscript) and a 19-base deletion (c.del226GCTGGGGCCGCATCT; clone F1, named KO2 in

the manuscript) respectively, which in both cases generated a premature stop codon. The two
clones were used as biological replicates in all experiments.

Trilineage differentiation
iPSCs were differentiated into the three germ layers (endoderm, ectoderm and mesoderm) using
the Human Pluripotent Stem Cell Functional Identification Kit (R&D Systems, SC027B). The
expression of relevant markers was then assessed using immunofluorescence with 10 ug/ml of
antibodies provided in the kit (Goat anti-human SOX17, Goat anti-human OTX2 and, Goat anti-
human TBXT).

iPSC-to-CNCC differentiation
All iPSC lines were differentiated into CNCCs as previously described (Prescott et al., 2015).
When iPS cells reached ~60% confluence in Geltrex-coated 6-well plates, cells were dissociated
with accutase in small suspension clusters and were treated with CNCC differentiation media (1:1
Neurobasal medium/D-MEM F-12 medium (Invitrogen), 0.5× B-27 supplement with Vitamin A
(50× stock, Gibco, #17504044), 0.5× N-2 supplement (100× stock, Gibco; #17502001), 20 ng/ml
human FGF (Gibco), 20 ng/ml recombinant EGF (Sigma-Aldrich), 5 µg/ml bovine insulin (Sigma-
Aldrich) and 1× Glutamax-I supplement (100× stock, Invitrogen). Medium (2 ml) was changed
every other day. At Day-7 of differentiation, adhering neurospheres and early migrating CNCCs
were transitioned to CNCC early maintenance media until Day-14 (1:1 Neurobasal medium/D-
MEM F-12 medium (Invitrogen), 0.5× B-27 supplement with Vitamin A (50× stock, Invitrogen),
0.5× N-2 supplement (100× stock, Invitrogen), 20 ng/ml bFGF (Biolegend), 20 ng/ml EGF (Sigma-
Aldrich), 1 mg/ml bovine serum albumin, serum replacement grade (Gemini Bio-Products # 700-
104 P) and 1× Glutamax-I supplement (100× stock, Invitrogen). Cell medium (2 ml) was changed
every other day. Differentiating cells were passaged with Accutase Cell Detachment Solution
(Stemcell Technologies; #07920) for all the subsequent passages when cells reached 80%
confluence, plated in a fresh, Geltrex-coated 6-well plates. After passaging, cells were treated
with 0.05% ROCK-inhibitor (Y27632; Stemcell Technologies) for 24 hours.

Real-time quantitative polymerase chain reaction (RT-qPCR)
Cells were harvested as single cells, and the total RNA was extracted using Monarch® Total RNA
Miniprep Kit (New England Biolabs) following manufacturer's instructions. 600ng of total RNA was
used to synthesize cDNA using Maxima First Strand cDNA Synthesis Kit for RT-qPCR (K1641,
Thermo Fisher). For each RT-qPCR reaction, 15 ng of cDNA was used with 0.1 µM of each primer,

10 μL of PowerUp™ SYBR™ Green Master Mix (Applied Biosystems) in a final volume of 20 μl.

Samples were analyzed with CFX Connect Real-Time PCR Detection System (Bio-Rad) using

the following thermal cycling parameters: 3 min at 95 °C, followed by 40 cycles of 10 s at 95 °C,

20 s at 63 °C followed by 30 s at 72 °C. Each sample was run in triplicate. 18 S rRNA was used

as normalizer.

Nuclear Fractionation and Co-immunoprecipitation

Cells were washed twice with ice-cold PBS, scraped, and collected by centrifugation at 10,000 ×

g for 1 minute. Pellets were resuspended in 500 μL of Buffer A (10mM HEPES pH7.9, 1.5mM

$MgCl_2$, 10mM KCl, 1mM PMSF, 0.5mM DTT, protease inhibitor cocktail) and cytoplasmic fractions

were obtained by incubating the cells on ice for 15 minutes, adding of 0.05% NP-40, followed by

centrifugation at 17,000 × g for 10 minutes at 4 °C. The resulting pellet was gently washed with

Buffer A and resuspended in Buffer C (10mM HEPES pH7.9, 12.5% glycerol, 0.75mM $MgCl_2$,

10mM KCl, 0.1mM EDTA, 400mM NaCl, 1mM PMSF, 0.5mM DTT, protease inhibitor cocktail).

Samples were incubated at 4 °C with shaking at 3,400 rpm for 1 hour, with vortexing every 10

minutes. Nuclear fractions were collected by centrifugation at 17,000 × g for 10 minutes at 4 °C.

For co-immunoprecipitation (co-IP), 25 μL of Dynabeads™ Protein A (ThermoFisher, Cat#

10002D) were washed twice with PBS containing 0.1% Tween-20 (PBS-T), resuspended in 1 mL

of PBS-T, and incubated with 2 μg of the appropriate antibody for 1 hour at 4 °C with rotation.

Bead-antibody complexes were washed twice with PBS-T, resuspended in 125 μL PBS-T, and

incubated with 100 μg of nuclear extract for 3 hours at 4 °C with rotation. Beads were then washed

five times with Buffer C and eluted in 25 μL of Buffer C supplemented with 4.1 μL of 6× reducing

Laemmli SDS sample buffer (ThermoFisher Scientific). Eluates and 5% input controls (nuclear

extract diluted 1:10 in PBS and mixed with 1:6 volume of 6× Laemmli buffer) were boiled for 5

minutes prior to SDS-PAGE. For Western blotting, 20 μL of each sample were loaded as

described below.

Western Blot

For each condition, cells were harvested and lysed with cold radioimmunoprecipitation assay

buffer (Pierce™ RIPA buffer, Thermo Fisher) with Pierce™ Protease Inhibitor Mini Tablets, EDTA-

free (Thermo Fisher). After centrifugation at 4°C at 16,000x g for 10 minutes, the total extracted

protein was quantified using Pierce™ BCA Protein Assay Kit, following manufacturer's

indications. Thirty micrograms of total protein were loaded in Novex WedgeWell 4–20% Tris-

Glycine Gel (Invitrogen) and separated through gel electrophoresis (SDS–PAGE) Tris-Glycine-

SDS buffer (Invitrogen). The proteins were then transferred to nitrocellulose membrane (Thermo
Fisher Scientific) and incubated for one hour in Intercept® (PBS) Blocking Buffer (Licor). After
blocking, membranes were incubated overnight at 4°C (while rocking) in suitable antibody diluted
in blocking buffer and 0.2% Tween-20, at recommended concentrations. Membranes were
washed in 0.1% PBS-T before and after being incubated for one hour in the appropriate secondary
antibody (1:15,000 dilution). The antibodies were then visualized using Odyssey XF Imaging
System (Licor) or ChemiDocTM MP Imaging System and imaged with Image Studio Software
(Licor) or Fiji(Schindelin et al., 2012).

Flow Cytometry analysis of surface markers
To obtain a single-cell suspension for flow cytometry analysis, cells were treated with Accutase
for 5 min at 37°C. Cells were then washed with cold phosphate-buffered saline (PBS)-2% fetal
bovine serum (FBS) and live cells were counted. 1×10^6 cells/condition were resuspended in
100 μL PBS-2% FBS and stained. For pluripotency evaluation, 4 μl of the respective antibodies
were used. For analysis of differentiation, 2 μl of the respective antibodies were used. Cells were
incubated for 15 min on ice and protected from light, before transferring them into FACS tubes
containing an additional 300 μL PBS-2% FBS. Flow cytometry data were acquired using
NovoCyte Penteon Flow Cytometer Systems 5 Lasers and analyzed with FlowJo Software version
10.9.

Immunofluorescence
Cells were passaged onto Geltrex-coated coverslips and fixed at required confluence or time-
point. In detail, cells were incubated in 4% PFA at 37°C for 15 minutes, followed by three 2-minute
washes in 1× PBS. For permeabilization, cells were treated with 0.1% (v/v) Triton X-100 (T8787,
Sigma-Aldrich) at room temperature for 10 minutes before three 2-minute washes in 1× PBS +
0.1% (v/v) Tween 20 and a 1-hour blocking upon 10% Donkey serum (D9663, Sigma-Aldrich) at
room temperature. Cells were incubated with corresponding PBS-diluted primary antibodies
protected from direct light at 4°C overnight. After three 5-minute washes with 1× PBS + 0.1%
Tween 20, cells were incubated for 1 hour in PBS-diluted secondary antibodies and 15 minutes
with DAPI (1:5,000; 422801, BioLegend). Cells were then washed three times with 1× PBS +
0.1% Tween for 5 minutes and once with 1× PBS. Coverslips were then fetched and mounted
onto slides using Fluorescence Mounting Medium (S3023, Agilent Technologies). Slides were
imaged using confocal microscope (LSM780, Zeiss). The intensity of fluorescent signal was
measured using Corrected Total Cell Fluorescence (CTCF) using Image J. Briefly, for each cell,

a region of interest (ROI) was drawn around the cell boundary, and the Integrated Density (sum of all pixel intensity values within the ROI), area of ROI (total number of pixels contained in the ROI), and the mean background fluorescence (average pixel intensity measured from a background region of the image lacking fluorescent signal) were measured.

The corrected fluorescence intensity was then calculated as:

$$CTCF = Integrated\ Density - (Area\ of\ ROI\ \times\ Mean\ Background\ Fluorescence)$$

For colony-forming cells (e.g., iPSCs), we analyzed segmented nuclei located within a section along the horizontal diameter of each colony, encompassing approximately 30–40 cells. This approach allowed us to capture differences between the colony center and edges while enabling comparisons across colonies of similar size.

RNA-seq

Total RNA was extracted using Monarch® Total RNA Miniprep Kit (New England Biolabs, T2010S) according to the manufacturer's instructions. RNA was quantified using NanoDrop 2000 UV Visible Spectrophotometer, while the RNA integrity was checked on Tapestation 2200 (Agilent). Only samples with RIN value above 8.0 were used for transcriptome analysis. RNA libraries were prepared using 1 µg of total RNA input using NEBNext® Poly(A) mRNA Magnetic Isolation Module, NEBNext® UltraTM II Directional RNA Library Prep Kit for Illumina® according to the manufacturer's instructions (New England Biolabs).

RNA-sequencing analysis

After removing the adapters with TrimGalore! (https://www.bioinformatics.babraham.ac.uk/projects/trim_galore/) , Kallisto was used to quantify reads mapping to each gene. Differential gene expression levels were determined using DESeq2(Love et al., 2014). All statistical analyses were performed using the latest versions of BEDTools(Quinlan and Hall, 2010), deepTools(Ramírez et al., 2016), R Studio (version 4.4.3; Trophy Case), and GraphPad Prism software version 10.2.3 (http://www.graphpad.com/faq/viewfaq.cfm?faq=1362; GraphPad, San Diego, CA).

CUT&RUN

For each condition, 1x 10^5 cells were processed using CUT&RUN Assay Kit #86652 (Cell Signaling Technology). Briefly, each sample was resuspended in Concanavalin A Beads and treated with an Antibody Binding Buffer and the suitable antibody. After the antibody incubation, each reaction underwent a step of pAG-MNase Enzyme binding, and subsequent DNA Digestion

and diffusion. The diffused DNA was then purified using Phenol/Chloroform Extraction and
Ethanol Precipitation. The purified DNA was quantified using QuantiFluor dsDNA System, using
the ONE dsDNA protocol (Promega). Libraries for sequencing were prepared using NEBNext
Ultra II DNA Library Prep Kit for Illumina (New England BioLabs) and the libraries size distribution
was assessed through automated electrophoresis using the Agilent 2200 Tapestation System
(Agilent Technology).

1005

ChIP-seq

1006
All samples from different conditions were processed together to prevent batch effects.
Approximately 15 million cells were double cross-linked with 2mM disuccinimidyl glutarate (DSG)
for 20 minutes followed by 1% formaldehyde for 10 minutes at room temperature, quenched with
125 mM glycine, harvested, and washed twice with 1X PBS. The fixed cell pellet was resuspended
in ChIP lysis buffer (150 mM NaCl, 1% Triton X-100, 0.7% SDS, 500 µM DTT, 10 mM Tris-HCl, 5
mM EDTA) and chromatin was sheared to an average length of 200–900 base-pairs, using a
Covaris S220 Ultrasonicator at 5% duty factor for 7 minutes for 3 times. The chromatin lysate was
diluted with SDS-free ChIP lysis buffer. 10µg of antibody was used for BRG1. The antibody was
added to at least 5µg of sonicated chromatin along with Dynabeads Protein G magnetic beads
(Invitrogen) and incubated with rotation at 4°C overnight. The beads were washed twice with each
of the following buffers: Mixed Micelle Buffer (150 mM NaCl, 1% Triton X- 100, 0.2% SDS, 20 mM
Tris-HCl, 5 mM EDTA, 65% sucrose), Buffer 200 (200 mM NaCl, 1% Triton X-100, 0.1% sodium
deoxycholate, 25 mM HEPES, 10 mM Tris-HCl, 1 mM EDTA), LiCl detergent wash (250 mM LiCl,
0.5% sodium deoxycholate, 0.5% NP-40, 10 mM Tris-HCl, 1 mM EDTA) and a final wash was
performed with cold 0.1X TE. Finally, beads were resuspended in 1X TE containing 1% SDS and
incubated at 65°C for 10 min to elute immunocomplexes. The elution was repeated twice, and the
samples were incubated overnight at 65°C to reverse cross-linking, along with the input (5% of
the starting material). The DNA was digested with 0.5 mg/ml Proteinase K for 3 hours at 55°C
and then purified using the ChIP DNA Clean & Concentrator kit (Zymo) and quantified with QUBIT.
Barcoded libraries were made with NEBNext Ultra II DNA Library Prep Kit for Illumina) using
NEBNext Multiplex Oligos Dual Index Primers for Illumina (New England BioLabs) and sequenced
on NovaSeq X Plus instrument (Illumina).

1029

ATAC-seq

1030
One hundred thousand cells/condition were harvested by enzymatic dissociation using
StemPro™ Accutase™ (Thermo Fisher Scientific). Each cell pellet was processed using ATAC-

Seq Kit (Active Motif), following the manufacturer's directions. Briefly, the fresh cell pellets were

washed in 1X PBS and lysate in ATAC Lysis Buffer. Each lysate underwent a process of

tagmentation and purification, before being amplified through PCR, using the following thermal

cycler parameters: 72°C for 5 min, 98°C for 30 sec, followed by 10 cycles of 98°C for 10 sec,

63°C for 30 sec, 72°C for 1 min. Each PCR reaction was cleaned up using SPRI beads (1.2X of

the sample volume). The Agilent 2200 Tapestation System (Agilent Technologies) was used as

quality control tool before performing next generation sequencing (NGS).

1040

Bioinformatics analysis of CHIP-seq, ATAC-seq and CUT&RUN data

Adapters were removed with TrimGalore!

(https://www.bioinformatics.babraham.ac.uk/projects/trim_galore/), the sequences were aligned

to the reference hg19, using Burrows-Wheeler Alignment tool, with the MEM

algorithm(Herbreteau et al., 2021). Uniquely mapping aligned reads were filtered based on

mapping quality (MAPQ > 10) to restrict our analysis to higher quality and uniquely mapped reads,

and PCR duplicates were removed. MACS2(Zhang et al., 2008) was used to call peaks using the

default parameters at 5% FDR.

1049

1050 **Motif analysis**

1051 BED files for the regions of interest were produced using BEDTools(Quinlan and Hall, 2010). Motif

1052 analysis was performed using HOMER(Heinz et al., 2010). Shuffled input sequences were used

1053 as background. E-values < 0.001 were used as a threshold for significance.

1054

QUANTIFICATION AND STATISTICAL ANALYSIS

RT-qPCR and image quantification data were analyzed using Prism software version 10.2.3

(http://www.graphpad.com/faq/viewfaq.cfm?faq=1362; GraphPad, San Diego, CA). Data is

represented as mean ± standard error of mean (SEM). GO enrichment analysis was performed

using Webgestalt 2019(Liao et al., 2019). All statistical analyses were performed using the latest

versions of BEDTools(Quinlan and Hall, 2010), Deeptools(Ramírez et al., 2016), R version 4.4.3

(Throphy Case; https://www.R-project.org/), and Prism software version 10.2.3

(http://www.graphpad.com/faq/viewfaq.cfm?faq=1362; GraphPad, San Diego, CA). In particular,

Bigwigs were generated with Deeptools and normalized by sequencing depth using the same

software, and upper-quantile normalized using CHIPIN(Polit et al., 2021).

A

B

C

D

E   **Day-0**   **F**   **Day-5**

A ATAC-seq peaks at Day 5

Accessible sites in WT
(n = 58,375)

Sites losing accessibility in KO
(n = 1,649)

Sites maintaining accessibility in WT and KO
(n = 60,186)

Sites gaining accessibility in KO
(n = 3,460)

Accessible sites in KO
(n = 70,040)

B ATAC-seq

Day-0
Day-5

Day-0   Day-5

Day-5 specific SALL4-bound regions (n=17,137)

Peak Summit (± 2Kbs)

C ATAC-seq at SALL4 Day-5 specific regions

Replicate 1
WT
KO

Peak Summit (± 2Kbs)

Replicate 2
WT
KO

Peak Summit (± 2Kbs)

D

Gaining accessibility in KO Day 5 motifs		
MOTIF #	**Motif Consensus**	**p-value**
51	CTCF BORIS	$1e^{-270}$
256	POU5F1/SOX2/ NANOG POU2F3 (epidermis) POU3F1 (nervous s.) POU2F2 (metastases)	$1e^{-103}$
410 (8%)	ZIC3 ZIC2 NANOG	$1e^{-84}$

E

Losing accessibility in KO Day 5 motifs			
MOTIF #	**Motif Consensus**	**TF**	**p-value**
410 (11%)		ZIC3 ZIC2 NANOG	$1e^{-59}$
92		ERG ETS	$1e^{-58}$
402		ZEB1 ZEB2 SNAIL1	$1e^{-50}$
407		SMAD4 SMAD2 P53	$1e^{-47}$
107		AP-1 JUN FOS ATF3	$1e^{-37}$

A

Very high (pIDDT > 90) Confident (90 > pIDDT > 70) Low (70 > pIDDT > 50) Very low (pIDDT < 50)

B

SALL4A

ARID1B C-terminus

SMARCA4 (BRG1)

ACTL6A

SMARCAB1

DPF2

DNA-Nucleosomes

SALL4A

Aligned Residue

Scored Residue

C

SALL4B

ARID1B C-terminus

SMARCA4 (BRG1)

ACTL6A

SMARCAB1

DPF2

DNA-Nucleosomes

SALL4B

Aligned Residue

Scored Residue

0 5 10 15 20 25 30
Expected Position Error (Ångströms)

D

SALL4B   1   7 8

Model 1
Model 3
Model 4

E

PREDICTED

SALL4
DPF2
SMARCA4
SMARCB1
ACTL6
ARID1B

EM - 3Å resolution

SMARCA4
DPF2
SMARCB
ACTL6
ARID1A

Supplementary Figure 1.

(A) (Left) Visual representation of the confidence score pLDDT (predicted Local Distance Difference Test) for each residue of the AlphaFold-predicted complex, with color gradients indicating prediction confidence (blue: very high confidence, yellow: low confidence, orange/red: very low confidence). (Right) Close-up view of the SALL4A-DPF2 interface highlighting key interacting residues in red.

(B) AlphaFold predicted aligned error (PAE) plot for model 0 of SALL4A. The plot shows the expected positional error in Ångströms for each residue pair.

(C) AlphaFold predicted aligned error (PAE) plot for model 0 of SALL4B. The plot shows the expected positional error in Ångströms for each residue pair.

(D) SALL4B protein sequences. C2H2 zinc fingers annotations are highlighted in dark purple and numbered. Visual representation of output models of SALL4B (models 1, 3, and 4). Interactions between SALL4B and DPF2 are highlighted in pink, light purple, and light blue. Made with BioRender.com.

(E) Comparison between the predicted BAF-SALL4A in silico structure and an experimentally identified structure of the human BAF Base module (PDB: 6LTH).

A

CLONE F1

Status ?	Guide Target ?	PAM Sequence ?	Indel % ?	Model Fit (R²) ?	Knockout-Score ?	Nuclease ?
✅ Succeeded	GTTTGCAGATGCGGCCCCAG	CGG	51	0.97	51	

POWERED BY EditCo ICE

RELATIVE CONTRIBUTION OF EACH SEQUENCE (NORMALIZED)

INDEL	CONTRIBUTION ▾	SEQUENCE
-19	51%	AGCTCCCCCGCCGCGGGCGCC----- ------------------ACTCCGGGGTCTGCTGCTGCGGCTGCTGCTCGCCC
+ 0	46%	AGCTCCCCCGCCGCGGGCGCCGCTG GGGCCGCATCTGCAAACTCCGGGGTCTGCTGCTGCGGCTGCTGCTCGCCC

CLONE C4

Status ?	Guide Target ?	PAM Sequence ?	Indel % ?	Model Fit (R²) ?	Knockout-Score ?	Nuclease ?
✅ Succeeded	GTTTGCAGATGCGGCCCCAG	CGG	52	0.98	52	

POWERED BY EditCo ICE

RELATIVE CONTRIBUTION OF EACH SEQUENCE (NORMALIZED)

INDEL	CONTRIBUTION ▾	SEQUENCE
-11	52%	AGCTCCCCCGCCGCGGGCGCCGCTG ----------GCAAACTCCGGGGTCTGCTGCTGCGGCTGCTGCTCGCCC
+ 0	46%	AGCTCCCCCGCCGCGGGCGCCGCTG GGGCCGCATCTGCAAACTCCGGGGTCTGCTGCTGCGGCTGCTGCTCGCCC

B

	DAPI	SALL4
WT		
SALL4-het-KO		

C

SALL4 immunofluorescence

WT
KO

CTCF

D

WT   *SALL4*-het-KO

Supplementary Figure 2.

(A) Sanger sequencing reports of the two iPSC *SALL4*-het-KO clones used in this study: clone F1 (19-bp deletion, top) and clone C4 (11-bp deletion, bottom). The data were analyzed using the ICE (Inference of CRISPR Edits) tool to quantify CRISPR-Cas9 editing efficiency. Editing resulted in an overall indel frequency of 51% and 52%, respectively. The sequence traces show the relative contribution of each allele, normalized across the population, with deleted bases indicated by dashed lines. Cut sites are represented by black vertical dotted lines, and the wild-type sequence is marked by a "+" symbol on the far left.

(B) Immunofluorescence labeling quantifying the expression of SALL4 in WT and clone F1 iPSC lines. Scale bar: 50 μm.

(C) Box plot showing Corrected Total Cell Fluorescence (CTCF) values quantifying and comparing SALL4 expression in WT and SALL4-het-KO (clone F1) in immunofluorescence assay (sample n= 30; background n= 5). Error bars display standard error of the mean.

(D) Phase-contrast microscopic images showing the morphology of *SALL4*-WT and *SALL4*-het-KO (clone F1) iPSC colonies at 4x (top) and 10x (bottom).

A Upregulated in *SALL4*-het-KO

- establishment of tissue polarity
- midbrain development
- regulation of animal organ morphogenesis
- cytoplasmic translation
- morphogenesis of a branching structure
- protein processing
- stem cell differentiation
- epithelial cell proliferation
- gland development
- synapse organization

Enrichment Ratio (0, 10, 20, 30)

B Downregulated in *SALL4*-het-KO

- **membrane raft organization**
- regulation of epithelial cell differentiation
- **multi-organism reproductive process**
- **multi-multicellular organism process**
- **organic hydroxy compound transport**
- carbohydrate homeostasis
- **hormone transport**
- epidermis development
- negative regulation of transport
- **signal release**

Enrichment Ratio (0, 10, 20, 30, 40, 50)

C

	WT	*SALL4*-het-KO
Ectoderm	OTX2 / OTX2/DAPI	OTX2 / OTX2/DAPI
Mesoderm	TBXT / TBXT /DAPI	TBXT / TBXT /DAPI
Endoderm	SOX17 / SOX17/DAPI	SOX17 / SOX17 /DAPI

Scale bar: 100 µm

Supplementary Figure 3.

(A) Top 10 GO term enrichment for upregulated genes in *SALL4*-het-KO iPSCs determined using WebGestalt[71] over-representation pathway analysis, using affinity propagation as parameter for redundancy removal. All terms are non-significant (FDR > 0.05).

(B) Top 10 GO term enrichment for downregulated genes in *SALL4*-het-KO iPSCs determined using WebGestalt[71] over-representation pathway analysis, using affinity propagation as parameter for redundancy removal. In bold terms with FDR < 0.05.

(C) Immunofluorescence for ectodermal (OTX2), mesodermal (TBXT), and endodermal (SOX17) markers in *SALL4*-WT and *SALL4*-het-KO (clone F1) after trilineage differentiation. Scale bar: 100 µm.

A. GATING STRATEGY FIG. 3B

Supplementary Figure 4.

Gating strategy for flow cytometry data, as reported in (A) figure 3B and (B) figure 7B.

A

	SALL4 CUT&RUN Day-0 specific motifs		
MOTIF #	**Motif Consensus**		**p-value**
51		CTCF BORIS	$1e^{-270}$
260		OCT:OCT	$1e^{-106}$

B

Day-0 specific SALL4 binding sites – *de novo* motifs				
Motif Consensus	**% of Target**	**% of Background**	**p-value**	**Closest match**
	20.17%	2.23%	$1e^{-486}$	CTCF BORIS
	4.63%	0.40%	$1e^{-118}$	
	2.60%	0.17%	$1e^{-73}$	SALL4 (ZFC4) ARID/HMG

C

Day-5 specific SALL4 binding sites – *de novo* motifs				
Motif Consensus	**% of Target**	**% of Background**	**p-value**	**Closest match**
	51.16%	17.48%	$1e^{-1613}$	TEAD
	34.74%	10.61%	$1e^{-1108}$	AP-1
	26.53%	9.00%	$1e^{-698}$	NANOG

Supplementary Figure 5.

(A) Table of motifs enriched in Day-0 specific SALL4-bound regions as determined using HOMER[68].

(B) Table of de novo motifs enriched in Day-0 specific SALL4-bound regions as determined using HOMER[68].

(C) Table of de novo motifs enriched in Day-5 specific SALL4-bound regions as determined using HOMER[68].

A

hNECs

H3K4me1 · H3K4me3

Sites losing BRG1 in *SALL4*-het-KO cells (n = 13,505)

PROMOTERS

ENHANCERS

Peak Summit (± 5Kbs)

B

H3K27ac (*SALL4*-WT)

genes

— Day-5
— Day-11
— Day-14

Mean density/tag

Peak Summit (± 5Kbs)
Sites losing BRG1 in *SALL4*-het-KO cells (n = 13,505)

C

H3K27ac (*SALL4*-WT)

Day-5 · Day-11 · Day-14

Sites losing BRG1 in *SALL4*-het-KO cells (n = 13,505)

Peak Summit (± 5Kbs)

Supplementary Figure 6.

(A) ChIP-seq signal intensity for H3K4me1 and H3K4me3 plotted across ±5 kb of sites losing BRG1 in SALL4-het-KO cells regions (n= 13,505) in human neuroectoderm-like cells (hNECs), centered on peak summits. Rows represent individual peaks ranked by signal intensity.

(B) Average profiles and (C) heatmaps showing H3K27ac ChIP-seq signal genomic sites plotted across ±5 kb of sites losing BRG1 in *SALL4*-het-KO cells regions (n= 13,505) at Day-5, Day-11, and Day-14 of the iPSC-to-CNCC specification. Centered on peak summits.

A

B

C

Supplementary Figure 7.

(A) GO term enrichment for genes near BRG1 ChIP-seq peaks gained in *SALL4*-het-KO cells. GO terms were determined using WebGestalt[1] over-representation pathway analysis, using affinity propagation as parameter for redundancy removal (FDR<0.05).

(B) GO term enrichment for genes near BRG1 ChIP-seq peaks lost in *SALL4*-het-KO cells. GO terms were determined using WebGestalt[1] over-representation pathway analysis, using affinity propagation as parameter for redundancy removal (FDR<0.05).

(C) Volcano plot of genes differentially expressed in *SALL4*-het-KO relative to *SALL4*-WT in neuroectoderm-like cells (Day-5). Blue dots (n= 126) represent downregulated genes with p-adj < 0.05 and log2FoldChange < -1.5. Purple dots (n= 44) represent upregulated genes with p-adj < 0.05 and log2FoldChange > 1.5.

A

Pluripotency Markers

NANOG

POU5F1

B

Neural Plate Border Markers

MSX1

PAX3

C

CNCC Specification Markers

SOX9

TFAP2A

TWIST1

D

Supplementary Figure 8.

Trajectory of stage-specific markers at Day-0 (n=2), Day-5 (n=2), and Day-14 (n=2). Displayed are the relative mRNA expression for:

(A) pluripotency markers (POU5F1 and NANOG),

(B) neural plate border markers (MSX1, PAX3),

(C) CNCC specification markers (SOX9, TFAP2A).

Highlighted are the approximate stages in which each marker is conventionally expressed in *SALL4*-WT.

(D) Expression (Log2[TPMs]) of neuronal-associated markers at Day-14.

Supplementary Figure 9.

(A) Trajectory of SALL4A and SALL4B RNA expression (TPMs) at significant iPSC-to-CNCC specification time-points: Day-0 (n=2); Day-5 (n=3); Day-14 (n=2).

(B) Time-course immunoblot for SALL4 in SALL4-WT at Day-0, Day-5, and Day-14. GAPDH was used as a loading control.

(C) Time-course immunoblot for SALL4 in SALL4-WT at Day-0, Day-5, and Day-14. GAPDH was used as a loading control.

(D) Immunoblot for ARID2 in SALL4-WT at Day-5 after 24-hour treatment with DMSO (control) or 5μM pomalidomide (5μM POM). GAPDH was used as a loading control.

(E) Immunoblot for ZMYM2 in SALL4-WT at Day-5 after 24-hour treatment with DMSO (control) or 5μM pomalidomide (5μM POM). GAPDH was used as a loading control.