Neural crest induction requires SALL4-mediated BAF recruitment to lineage specific enhancers

Martina Demurtas[1], Samantha M. Barnada[2], Emma van Domselaar[1], Zoe H. Mitchell[1],

Laura Deelen[1], and Marco Trizzino[1,*]

[1]Department of Life Sciences, Imperial College London, London, UK

[2] Department of Biochemistry and Molecular Biology, Thomas Jefferson University, Philadelphia, PA, USA

*Lead contact: m.trizzino@imperial.ac.uk

SUMMARY

Neural crest induction begins early during neural plate formation, requiring precise transcriptional

control to activate lineage-specific enhancers. Here, we demonstrate that SALL4, a transcription

factor strongly expressed in cranial neural crest cells (CNCCs) and associated with syndromes

featuring craniofacial anomalies, plays a critical role in this early regulatory process. Using a

SALL4-haploinsufficient human iPSC model that recapitulates clinical haploinsufficiency, we

show that the SALL4A isoform directly interacts with the BAF complex subunit DPF2 through its

zinc-finger-3 domain. This interaction enables SALL4-mediated recruitment of BAF to CNCC-

specific enhancers during early neural plate stages. Without functional SALL4, BAF fails to load

at chromatin, leaving CNCC enhancers inaccessible despite normal neuroectodermal

progression. Consequently, cells cannot undergo proper CNCC induction and specification due

to persistent enhancer repression. Our findings reveal SALL4 as an essential regulator of BAF-

dependent enhancer activation during early neural crest development, providing molecular

insights into SALL4-associated craniofacial anomalies.

KEYWORDS

SALL4, DPF2, SALL4A, BAF, neural crest, enhancers, chromatin accessibility

INTRODUCTION

Embryonic development is a tightly regulated process that depends on the ability of pluripotent embryonic stem cells (ESCs) to progressively acquire specialized fates and functions[1]. This enables their differentiation into a diverse range of somatic lineages, ultimately giving rise to a complex, fully integrated organism from a single cell[2]. These diverse cell identities are established through epigenetic mechanisms that regulate the spatiotemporal expression of lineage-specific gene regulatory networks, primarily by modulating chromatin accessibility. Chromatin states range from open to closed, shaped by a combination of epigenetic modifications, including histone methylation and acetylation, incorporation of histone variants, DNA methylation, and the activity of chromatin remodeling complexes. These large, multi-subunit complexes utilize ATP hydrolysis to reposition nucleosomes, dynamically regulating access to genes and their regulatory elements[3–6].

Transcription factors also play a crucial role in cell fate determination, acting as molecular switches that trigger mechanisms that ultimately activate or repress genes as a result to their binding to specific regulatory DNA sequences within *cis*-regulatory elements. They influence gene expression by recruiting or inhibiting the transcriptional machinery and regulating 3D chromatin organization[3,4,6–9]. Among these, the embryonic C2H2 zinc-finger transcription factor SALL4 is involved in a wide range of developmental processes[10]. SALL4 is highly expressed in pluripotent cells and throughout embryonic development, gradually decreasing after birth to become undetectable in adult tissues[11].

In humans, the SALL4 gene produces two main isoforms through alternative splicing: SALL4A (1053 aa) and SALL4B (617 aa), which have been proposed to play distinct roles in ESCs. Namely, Rao et al.[12] suggested that SALL4B has function limited to pluripotency regulation in ESCs, while SALL4A is the isoform primarily active during development. Although SALL4 is expressed as early as the two-cell stage, its role in maintaining pluripotency remains debated. Some studies suggest that SALL4 cooperates with core pluripotency transcription factors, such as OCT4 and NANOG, to sustain stem cell identity[13,14]. In contrast, other studies propose that SALL4 plays a more marginal role in directly regulating the pluripotency gene regulatory network[15–17].

During early mouse development, *Sall4* is broadly expressed throughout the embryo, however, as gastrulation progresses, its expression becomes more posteriorly restricted, localizing primarily to the developing tail bud and limb buds[18,19]. Here, it regulates the cell fate commitment of neuromesodermal progenitors by suppressing the neural gene regulatory network in mesodermal cells while simultaneously enhancing mesodermal gene expression[20]. Additionally,

Sall4 remains expressed in mouse craniofacial regions until the E9.5 stage, including the
developing frontonasal process as well as the mandibular and maxillary arches[18]. These
craniofacial structures originate from cranial neural crest cells (CNCCs), a subpopulation of neural
crest cells that migrate dorsolaterally to give rise to the cartilage and bones of the face and anterior
skull, as well as cranial neurons and glia. Consistent with this, individuals with *SALL4*
haploinsufficiency exhibit a range of phenotypes, including distinctive facial features such as
epicanthal folds, widely spaced eyes, and a depressed nasal bridge, along with cranial nerve-
related conditions. These include Duane anomaly, a congenital strabismus resulting from
abnormal oculomotor nerve innervation[21], and sensorineural hearing loss[22].

Together, these findings suggest that SALL4 may play a critical role in CNCC induction,
specification and differentiation. However, its involvement in this developmental process has not
yet been investigated. Previous studies have also suggested that SALL4 interacts directly with
chromatin remodeling complexes. For example, several studies in mice have identified Sall4 as
an interactor of the NuRD chromatin remodeling complex, in both embryonic cells and various
developmental processes[15,23–26]. Additionally, recent mass-spectrometry experiments conducted
by our group in early stages of human iPSC-to-CNCC differentiation, and specifically at the
neuroectodermal stage, revealed that SALL4 interacts with the BAF chromatin remodeling
complex during this developmental process[27].

Since this was the first report of a potential direct SALL4-BAF interaction, here we set out to
investigate the scope of this interaction, and to determine the role of SALL4 during cranial neural
crest cell specification. Given that human SALL4-associated developmental syndromes result
from *SALL4* haploinsufficiency, we generated a *SALL4* heterozygous human iPSC line using
CRISPR-Cas9-mediated deletion. With this approach, we demonstrate that, at early stages of
CNCC specification, SALL4 is required for the recruitment of the BAF complex to open the
chromatin to enhancers essential for CNCC fate. In the absence of SALL4, cells progress
normally through the neuroectodermal stage but fail to specify into CNCCs, as the lineage-specific
enhancers remain repressed. Finally, using *in silico* approaches, we predict that SALL4A isoform
directly interacts with the BAF subunit DPF2. Specifically, this interaction occurs between the
zinc-finger-cluster-3 (ZFC3) domain of SALL4A and the plant homeodomains (PHDs) of DPF2.

RESULTS

SALL4 interacts with the BAF complex

Cranial neural crest cells can be generate *in vitro* starting from human pluripotent stem cells
(hiPSCs). This protocol is approximately two weeks long, and it first goes through a

neuroectoderm intermediate [27–31]. In our previous work[27], mass spectrometry (IP-MS) conducted

at early stages of the iPSC-to-CNCC differentiation protocol, and specifically Day-5, which

corresponds to the neuroectodermal stage, revealed that SALL4 interacts with the BAF chromatin

remodeling complex in this developmental cell population. Since this was the first report of a

SALL4-BAF interaction, we performed a co-immunoprecipitation (co-IP) at Day-5 to validate it.

Specifically, we performed pull-down of ARID1B, which is the only ARID1 subunit active and

incorporated in BAF at this stage [27], and blotted for SALL4. This co-IP confirmed that ARID1B

interacts with both SALL4 isoforms at this time-point (Fig. 1A).

The BAF chromatin remodeling complex is a multi-subunit assembly composed of a dozen

components, which are differentially incorporated in a tissue-specific manner[32]. Based on which

subunit is present in each complex, in mammals these can be classified into canonical BAF,

polybromo-associated BAF, and non-canonical BAF[33]. While our mass-spec and co-IP

experiments support the interaction between SALL4 and the canonical BAF (ARID1B is not

incorporated in the other BAF configurations), neither of these two approaches has the resolution

to identify the exact BAF subunit that mediates the interaction with SALL4. Therefore, we

performed an *in silico* analysis using AlphaFold3[34]. We tested various BAF subunit combinations

along with four histones and an AT-rich DNA sequence which has been empirically shown to be

recognized by SALL4 (5'-GGAAATATTACC-3', PBD: 8A4I). The output was statistically

significant, with a 0.45 predicted template modeling (pTM) score and 0.47 interface predicted

template modeling (ipTM) score. Through this approach, AlphaFold3 predicted a direct interaction

between DPF2, a broadly expressed BAF subunit, and the zinc-fingers 5 and 6 (ZFC3) and 7

(ZFC4) of SALL4 (Figs. 1B, 1C, 1D; Supplementary Table 1). *DPF2* is a member of the *BAF45*

gene family, which also includes *PHF10*, *DPF1*, and *DPF3*. We were able to rule out its

paralogues based on their distinct expression patterns, incorporation into specific BAF

complexes, and previous reports that BAF45 subunits are not functionally redundant[35]. In

particular, PHF10 is exclusively associated with the pBAF complex, which does not incorporate

ARID1B[36], while DPF1 and DPF3 are specifically incorporated into the neuron-specific nBAF

complex[37] and are expressed at very low levels in both our iPSCs and neuroectoderm-like cells

at Day-5 (Fig. 1E).

This finding was particularly interesting given that DPF2 deletion has been linked to early neural

ectoderm differentiation in embryonic stem cells (ESCs)[35]. DPF2 exists in two isoforms, resulting

from alternative splicing of exon 7 and are spliced in a tissue-specific and developmental-stage-

specific manner[38]. DPF2-L is not alternatively spliced and is expressed in ESCs, while DPF2-S is

expressed in neurons. Interestingly, both isoforms are expressed in neural stem cells. Notably, of

the two main SALL4 isoforms (SALL4A and SALL4B), SALL4B lacks the portion of exon 2 that

contains ZFC3. This prompted us to explore whether the SALL4-DPF2 interaction could be

isoform-specific. To address this, we reanalysed the interactions in AlphaFold3, incorporating

SALL4B to the same BAF subunits as before. To ensure biological relevance, we first excluded

two of the five models that did not exhibit interactions with the AT-rich DNA sequence, as SALL4

is expected to bind DNA for its function. Among the remaining three models, we observed that

SALL4B interacted with DPF2, but unlike the SALL4A isoform, its interaction was not domain

specific. Instead, the predicted interactions for SALL4B were characterized by annotated

disordered regions and appeared to be largely driven by a non-specific, zinc-finger-to-zinc-finger

association (Fig. 1C; Supplementary Table 2), with no particular domain-specific interactions

identified. This pattern may be the result of the tendency of AlphaFold3 learning algorithm to

associate similar domains, such as zinc fingers, based on their predicted functional specificity

rather than spatial conformations. These findings suggest that, during neural ectoderm

specification, SALL4 may associate with BAF in an isoform-dependent manner.

To accurately interpret our AlphaFold3 predictions, we examined two key confidence metrics: the

Predicted Aligned Error (PAE), which assesses the reliability of the relative positioning of residues

within the structure, and the Predicted Local Distance Difference Test (pLDDT), which provides a

per-residue confidence score for the model. Our SALL4-DPF2 contact analysis revealed 394

atomic interactions (center-to-center distance ≤ 8Å) involving 139 residues (Fig. 1B, right;

Supplementary Table 1). Encouragingly, the pLDDT values were highest in known functional

domains involved in the interaction, particularly ZF5, ZF6, and ZF7 in SALL4A, as well as the

PHD1 and PHD2 domains in DPF2, reinforcing the reliability of these predicted structures. To

refine our analysis, we focused on residues where at least half of their atomic interactions had a

pLDDT > 65 (70 $\pm$ 10%) and that were annotated as part of a functional domain, ensuring that

only higher-confidence and potentially functional interactions were considered. As a result, ZF7

was excluded from further analysis (Supplementary Table 1). In summary, our experiments and

analyses confirm that SALL4 interacts with BAF during early stages of human CNCC

specification, and predict that this interaction is mediated by the BAF subunit DPF2.

Generation of a *SALL4*-haploinsufficient human iPSC line

Our experiments so far have revealed that SALL4 interacts with the BAF complex at early stages

of CNCC specification, and specifically at the neuroectodermal stage (Day-5 of our iPSC-to-

CNCC differentiation protocol). To investigate the scope of the SALL4-BAF interaction during this

developmental process, we generated a *SALL4* haploinsufficient iPSC line (hereafter *SALL4*-het-

KO; Fig. 2A). We opted for a heterozygous KO of *SALL4* as this recapitulates the human *SALL4*

syndromes, which are all caused by *SALL4* haploinsufficient genetic variants[22,39,40]. The CRISPR-

KO was generated by introducing a 11 bp deletion within the 3'-terminus of exon 1 of *SALL4*. This

is a conserved region across both *SALL4* isoforms that lacks functionally significant domains.

SALL4 immunoblot confirmed that SALL4 protein expression was significantly reduced in the

SALL4-het-KO iPSCs compared to the unedited wild-type line (*SALL4*-WT; Fig. 2B). We noted

that SALL4 expression in the *SALL4*-het-KO line was significantly less than the expected 50% of

the wild-type, suggesting potential negative feedback mechanisms triggered by the reduced

SALL4 levels, possibly due to the preferentially homodimeric nature of this protein.

Next, we investigated whether other SALL paralogs (*SALL1*, *SALL2*, *SALL3*) could compensate

for the loss of SALL4. Thus, we investigated their expression at gene level (Fig. 2C) and found

that, while *SALL2* and *SALL3* were not differentially expressed at the RNA level, *SALL1* was

instead significantly upregulated in the *SALL4*-het-KO cell line, at both RNA and protein level

(Figs. 2C and 2D), indicating a potential compensation in pluripotent cells.

Importantly, *SALL4*-het-KO iPSCs remain viable and exhibit hallmark iPSC morphology, including

a high nucleus-to-cytoplasm ratio and growth in compact colonies with well-defined borders (Fig.

2E). Furthermore, these cells show no significant difference in growth rate compared to their

isogenic controls (Fig. 2F).

Decreased SALL4 expression does not affect the stem cell identity of the iPSCs

Given the debated role of SALL4 in maintaining pluripotency, we investigated whether *SALL4*-

het-KO impacted iPSC stem cell identity. Thus, we examined the protein levels of the canonical

pluripotency transcription factors NANOG and OCT4 and found that they remained unaltered in

the *SALL4*-het-KO cells relative to the *SALL4*-WT control (Fig. 3A). Similarly, the levels of the

pluripotency surface markers TRA-1-60-R and SSEA4 were also not significantly affected by

decreased SALL4 expression (Fig. 3B). Moreover, we examined the transcriptome of the iPSCs

using bulk RNA-seq and identified only 114 differentially expressed genes in *SALL4*-het-KO cells

relative to the *SALL4*-WT control (FDR < 0.05; Fig. 3C). Gene Ontology analysis revealed that

these genes were not enriched for pluripotency (Fig. 3D and 3E). Finally, by performing tri-lineage

differentiation we confirmed that decreased SALL4 expression levels did not impact the ability of

the *SALL4*-het-KO cells to successfully differentiate into all three of the germ layers

(Supplementary Fig. 1A).

Together, these findings suggest that *SALL4* haploinsufficiency does not affect the stem cell

identity and self-renewal of human iPSCs, possibly as a result of SALL1-mediated compensation.

SALL4 relocates to CNCC developmental enhancers upon CNCC induction cues

After establishing that *SALL4*-het-KO did not affect iPSC stem cell identity, we differentiated the iPSCs into CNCCs using the same aforementioned protocol[27–31] (Fig. 4A). Since the SALL4-BAF interaction was detected at Day-5 of this protocol[27] (neuroectodermal stage), we profiled SALL4 binding in *SALL4*-WT cells by performing CUT&RUN at the iPSC stage and at Day-5 of differentiation. We performed two SALL4 CUT&RUN replicates per timepoint, and only retained replicated peaks in each timepoint (FDR < 0.05). With this approach, we identified comparable amounts of SALL4-bound genomic sites at Day-0 and Day-5 (29,088 and 32,707, replicated peaks, respectively, FDR < 0.05). Further analysis of these sites indicated that approximately 20% of the Day-0 SALL4 peaks were unique to the iPSC stage and not detected at Day-5 (hereafter SALL4 Day-0 specific peaks; n=6,114; Fig. 4B, left). On the other hand, ~50% of the peaks detected at Day-5 were exclusive of this developmental stage and were not detected at Day-0 (hereafter SALL4 Day-5-specific peaks, n= 17,137; Fig. 4B, right). Finally, a subset of ~15,000 sites remained bound by SALL4 throughout exit from pluripotency to neural ectoderm cell fate commitment (i.e. peaks conserved and replicated between Day-0 and Day-5). Together, this data indicate that, upon differentiation cues, there is a major shift in SALL4 binding, resulting in the relocation of this transcription factor at ~17,000 previously unbound regions.

To understand the scope of SALL4's relocation, we analysed the positional distribution of the SALL4 peaks by associating each peak with the nearest transcription start site (TSS). Consistent with SALL4's role as an enhancer-binding transcription factor, 86% of SALL4-bound regions at Day-0 were located >1Kb from the nearest TSS (n= 3,729), indicating they could be putative enhancers. This proportion increased even further to 97% (n= 6,860) at Day-5, further emphasizing the preferential binding of SALL4 at enhancer regions (Fig. 4C). To investigate the scope of the newly acquired SALL4 Day-5 peaks, we first looked at the nearest genes. GO terms enriched in the "Day-5 specific" genes included pathways associated with neural crest development, such as neural tube development, face development, cardiac chamber development, and cell-cell adhesion mediated by cadherin (Fig. 4D; FDR < 0.05, clustered using affinity propagation redundancy removal parameter).

To further characterise the SALL4 bound enhancers and promoters at Day-0 and Day-5, we performed sequence-based computational motif analysis. At Day-0, motifs associated with core pluripotency transcription factors such as OCT4, NANOG, and SOX2 did not rank among the top 20 enriched sequences, further supporting the notion that SALL4 may not be essential for maintaining pluripotency in human cells (Supplementary Figure 2A). Moreover, Day-0 specific

SALL4 peaks were enriched for the enhancer box (E-box) motif (Supplementary Figure 2A). The

E-box is usually bound by homo- or heterodimers of transcription factors containing a basic Helix-

Loop-Helix (bHLH) or a Zn-finger domain to regulate the expression of genes associated to cell

lineage commitment, especially in the context of skeletal muscle, cardiovascular and neuronal

tissue[41]. On the other hand, in addition to general enhancer activator factors (AP-1, TEAD), the

Day-5 specific SALL4 peaks were enriched for many CNCC signature factors, including SOX9,

TFAP2A, TWIST1/2, SNAI1, NR2F1/2, TCF4/12 and several others (Fig. 4E). Moreover,

neuroectodermal transcription factors, such as ZIC2/3 were also enriched. In summary, this

analysis suggests that, as early as the neuroectodermal stage, SALL4 relocates to enhancers of

genes associated with craniofacial development.

Chromatin accessibility at the CNCC enhancers is SALL4 dependent

Given that we demonstrated that SALL4 interacts with BAF at early stages of CNCC specification,

we reasoned that SALL4 binding at the CNCC developmental enhancers could be necessary for

their chromatin activation. Thus, we profiled chromatin accessibility by performing ATAC-seq at

Day-0 and Day-5 in both *SALL4*-WT and *SALL4*-het-KO conditions (two replicates per timepoint

per condition, only peaks present in both replicates were considered; FDR < 0.05). The number

of ATAC-seq replicated peaks identified in *SALL4*-WT condition was comparable between Day-0

(61,660 peaks; FDR < 0.05) and Day-5 (58,375 peaks; FDR < 0.05). Next, we specifically

investigated the chromatin state of the 17,137 genomic sites that gain SALL4 binding at Day-5

and found that SALL4 binding at these sites correlates with a marked increase in chromatin

accessibility compared to Day-0 (Fig. 4F). Strikingly, the accessibility at these regions was

significantly attenuated in *SALL4*-het-KO relative to *SALL4*-WT (Fig. 4G). Overall, these findings

suggest that, in response to differentiation signals, SALL4 relocates to CNCC developmental

enhancers, which subsequently become activated through chromatin opening, which is SALL4

dependent.

SALL4 recruits the BAF complex at CNCC developmental enhancers to increase chromatin

accessibility

Since SALL4 binding at the CNCC enhancers is necessary for chromatin opening, we sought to

determine whether the SALL4-bound developmental enhancers were also occupied by BAF, and

if BAF binding at these sites could be SALL4 dependent. We performed ChIP-seq for BRG1

(SMARCA4), the only catalytic subunit of BAF active at this time point, in *SALL4*-WT cells at Day-

5 of the iPSC-to-CNCC differentiation protocol. This experiment revealed that BRG1 consistently

binds the same genomic sites as SALL4, supporting the idea that these factors interact and co-
occupy regulatory elements during early differentiation (Fig. 5A).
To assess whether BRG1 binding depends on SALL4, we performed ChIP-seq for BRG1 in both
SALL4-WT and *SALL4*-het-KO cells at Day-5. In *SALL4*-WT cells, we detected 32,833 replicated
BRG1 peaks (FDR < 0.05), as opposed to only 14,116 peaks in *SALL4*-het-KO cells. This
suggested that SALL4 loss leads to a widespread loss of BAF from chromatin. We thus focused
on the BRG1 peaks that were lost in *SALL4*-het-KO cells and found that these sites were largely
not bound by SALL4 at Day-0, but they gained SALL4 binding at Day-5 (Figs. 5B and 5C). As
expected, loss of SALL4 and BRG1 from these sites led to marked decrease of chromatin
accessibility (Fig. 5D). Overall, these experiments support that SALL4 is required to recruit the
BAF complex at CNCC enhancers. Consistent with this, genes near BRG1 peaks lost in the
SALL4-het-KO cells are involved in embryonic organ development, epithelial differentiation, and
cell migration processes (Fig. 5E).
While a large portion of BRG1 peaks was lost in the absence of SALL4, we also identified a subset
of novel BRG1 peaks that were absent in *SALL4*-WT cells but gained in the *SALL4*-het-KO cells.
These BRG1 peaks gained in the *SALL4*-het-KO cells were associated with genes linked to
neurogenesis and nervous system development (Fig. 5F). These findings indicate that, in the
absence of SALL4, BRG1 is relocated from CNCC enhancers to neurodevelopmental enhancers,
shifting chromatin remodeling activity away from CNCC specification.
Together, these results suggest that SALL4 is a key regulator of BAF complex recruitment to
CNCC-specific enhancers, ensuring their accessibility during differentiation. In the absence of
SALL4, BRG1 fails to localize to these sites and is instead partially redirected to
neurodevelopmental enhancers, leading to a misallocation of chromatin remodeling activity,
potentially leading to a shift towards neural differentiation at the expense of CNCC induction and
specification.
296
SALL4-het-KO cells are not able to exit the neural ectoderm stage and to induce the cranial
neural crest fate
Our data thus far indicate that SALL4 relocates to CNCC enhancers as early as the
neuroectodermal stage, recruiting BAF to these regulatory elements for chromatin activation.
Next, we investigated the functional consequences of SALL4 loss by generating CNCCs using
SALL4-WT and *SALL4*-het-KO cells. As expected, the *SALL4*-WT CNCCs expressed the
signature CNCC markers at both the RNA (Fig. 6A) and protein (Fig. 6B) level, while *SALL4*-het-
KO showed a significant downregulation of these genes (Fig. 6A). This was further supported by

RNA-seq conducted at Day-14, which identified 1,106 differentially expressed genes in *SALL4*-het-KO CNCCs relative to *SALL4*-WT (FDR < 0.05; Fig. 6C). This is in sharp contrast with the small number of genes differentially expressed at the iPSC stage (n= 114; Fig. 3C) and at Day-5 (n= 170; Supplementary Fig. 2B).

Of the 1,106 genes differentially expressed at Day-14, ~60% (n= 659) were downregulated while the remaining were upregulated (Fig. 6C). Remarkably, canonical CNCC markers such as TFAP2A, NR2F1, and SOX9 were significantly downregulated in *SALL4*-het-KO cells. Additionally, genes associated with neural plate border, early neural crest induction and epithelial-to-mesenchymal transition (EMT) were also downregulated (*MSX1*, *GBX2*, *SNAI2*, *SOX5*, *ZEB1/2*; Fig. 6C). In line with these findings, gene ontology analysis revealed that pathways associated with face development, autonomic nervous system, gastrulation, and regulation of morphogenesis were all downregulated (Fig. 6D). The 447 upregulated genes were instead involved in regulatory pathways, especially vesicle-mediated transport of inorganic substances.

As aforementioned, genes associated with EMT were also dysregulated. The process of EMT requires a switch in cadherin expression where E-cadherin (*CDH1*) must be downregulated, while N-cadherin (*CDH2*) must be upregulated. Remarkably, our RNA-seq data revealed that this cadherin switch did not occur in the differentiating *SALL4*-het-KO iPSCs, suggesting that the cells remained in an epithelial state. This was further corroborated by the upregulation of several epithelial genes in the differentiating *SALL4*-het-KO cells, including *EpCAM*, *OTX2*, *CDH1*, and *OVOL2* (Fig. 6E). The latter is involved in maintaining epithelial lineages by repressing the expression of the EMT gene *ZEB1*, which was downregulated in the *SALL4*-het-KO condition[42,43]. In summary, our data indicate that CNCC specification failed in *SALL4*-het-KO conditions, as reflected by the large number of differentially expressed genes. On the other hand, the transcriptome of the iPSCs and of the neuroectodermal cells (i.e. Day-5) were only modestly affected by SALL4 loss, suggesting that SALL4 may not be essential for neuroectoderm formation, while it is essential for CNCC induction and specification.

To investigate this further, we systematically examined the expression of a panel of stage-specific marker genes at Day-0, Day-5 and Day-14 (Fig. 6F) including core pluripotency genes (*POU5F1* and *NANOG*), neural plate border specifiers (*ZIC1, GBX2, MSX1/2, PAX3/7, DLX5/6, GATA2/3*), CNCC specification markers (*SNAI1/2, SOX9, TFAP2A*), and post-migratory CNCC markers (*SOX5, SOX6, SOX10*). In *SALL4*-WT cells, pluripotency markers exhibited a progressive and expected downregulation as differentiation advanced from Day-0 to Day-14. In *SALL4*-het-KO cells, a similar decline in pluripotency markers was observed between Day-0 and Day-5, but instead of continuing to decrease, these markers remained similarly expressed through Day-14

(Fig. 6F). Despite this, the two conditions showed similar trends, confirming that decreased levels
of SALL4 dId not significantly impact pluripotency markers expression during exit from
pluripotency. Similarly, no significant difference was observed in the expression of neural plate
border specifiers (Fig. 6G), suggesting that SALL4 is not required for neuroectoderm specification.
Conversely, CNCC specification markers and post-migratory CNCC markers all increased in
expression after Day-5 in *SALL4*-WT cells, while the SALL4-depleted cells failed to maintain a
comparable increase in expression (Fig. 6H and 6I).
In summary, *SALL4*-het-KO cells are competent to generate neuroectoderm-like cells (Day-5) but
fail to induce the neural crest lineage (Day-14).
348
DISCUSSION
During embryonic development, SALL4 plays a critical role in forming major anatomical
structures, including the heart, limbs, cerebral midline, and pituitary gland[44]. Correspondingly,
mutations in the haploinsufficient gene *SALL4* are associated with congenital syndromes
characterized by malformations in these structures. These syndromes also frequently include
distinct craniofacial anomalies, such as epicanthal folds, hypertelorism, a depressed nasal bridge,
and hemifacial microsomia[45–47], all of which are hallmark features of disrupted cranial neural crest
cell development.
Notably, at the murine embryonic stage E9.5, Sall4 expression persists within the frontonasal
prominence, which is the most anterior embryonic region[18], implying a potential role for SALL4 in
craniofacial development. Given the limited existing knowledge about SALL4 function in this
context, we utilized a heterozygous *SALL4* knockout human induced pluripotent stem cell (iPSC)
line to investigate its role during cranial neural crest cell (CNCC) specification. Although SALL4
expression begins as early as the 2-cell embryonic stage, its role in maintaining pluripotency
remains debated. Some studies indicate that SALL4 cooperates closely with key pluripotency
factors such as OCT4 and NANOG to sustain stem cell identity[13,14]. Others, however, report that
SALL4 has only limited involvement within pluripotency regulatory networks[15–17]. Our findings
align with the latter hypothesis, demonstrating that decreased expression of SALL4 does not
compromise iPSC identity, possibly due to compensatory mechanisms involving its paralogue,
SALL1.
Using CUT&RUN, we observed that at early stages of CNCC induction, SALL4 relocates to
enhancers of genes involved in CNCC formation, and that SALL4 binding at these regulatory
elements is required for their activation. In humans, SALL4 exists in two major isoforms generated
by alternative splicing: SALL4A (1,053 amino acids), which retains all eight zinc-finger motifs, and

SALL4B (617 amino acids), which contains a partially spliced exon 2. Rao et al.[12] proposed distinct functional roles for these isoforms in murine ESCs, demonstrating that SALL4A homodimers localize predominantly to enhancers of developmental genes, whereas pluripotency-associated enhancers are occupied either by heterodimers comprising both isoforms or by SALL4B alone. In mouse ESCs, approximately 10% of the total Sall4 protein, and primarily the Sall4A isoform, is incorporated into the NuRD complex [15,24]. Here, we identify and characterize a novel interaction between SALL4 and the BAF complex, predicted to be mediated specifically through the SALL4A isoform and the BAF subunit DPF2. Furthermore, we show that upon receiving CNCC induction cues, SALL4 and BAF co-localize to enhancers of genes involved in CNCC specification to prime and activate them, with BAF recruitment being SALL4-dependent.

Interestingly, DPF2 and SALL4 exhibit notable functional parallels. Similar to SALL4, DPF2 occupancy undergoes significant repositioning during the transition from pluripotency toward early neural commitment in mouse embryoid bodies [48]. Although the role of DPF2 in CNCCs has not been directly investigated, its early recruitment to neuronal enhancers suggests a potential involvement in neuroectodermal differentiation and the subsequent lineage commitment of neuroectoderm-derived cells, including CNCCs. Moreover, genomic sequences bound by DPF2 in mouse neuroectodermal cells are AT-rich, resembling the preferred binding motif of SALL4[38,49,50].

Our study demonstrated that while cranial neural crest cell induction and specification fail in the absence of SALL4, early neuroectodermal commitment remains unaffected (Fig. 7). This finding is further supported by clinical observations. In fact, phenotypes arising from SALL4 haploinsufficiency affect many neural crest derivatives, including craniofacial structures and peripheral neurons. In contrast, phenotypes implicating defects in neural ectoderm or neurogenesis are exceptionally rare[51,52] and usually associated with dysregulation of potassium channels[44]. In the rare cases of intellectual disability reported, such phenotypes were attributed to broader deletions on chromosome 20q13.13-q13.2 rather than *SALL4* loss alone[51].

SALL4 was recently identified as a direct target of thalidomide[53], a drug widely prescribed to pregnant women in the late 1950s to alleviate morning sickness. The subsequent alarming rise in births of children with congenital anomalies, including limb malformations (phocomelia), facial abnormalities such as inner ear defects, ocular motility issues, and facial nerve palsy, as well as kidney and heart defects, led to the identification of thalidomide as the causative agent. However, the molecular mechanism by which thalidomide induced these developmental defects remained elusive until nearly six decades after the drug's withdrawal from the market. Building on earlier findings that thalidomide and its derivatives bind to cereblon (CRBN), Matyskiela et al.

demonstrated that SALL4 is recruited to the CRL4–CRBN E3 ubiquitin ligase complex following thalidomide-induced ubiquitination, resulting in its proteasomal degradation[54]. More recently, research demonstrated that only the SALL4A isoform, but not SALL4B, is susceptible to degradation by immunomodulatory imide drugs such as thalidomide and its derivative pomalidomide[55]. In this context, our findings might contribute to shed light on the birth malformations caused by thalidomide in the 1960s, potentially revealing some of the underlying mechanisms.

RESOURCE AVAILABILITY

Data availability

RNA-seq, ATAC-seq, CUT&RUN-seq, and ChIP-seq data have been deposited in the Gene Expression Omnibus (GEO) under accession code GEO: GSE293821 and are publicly available at https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE293821.

Lead contact

Further information and requests for resources and reagents should be directed to and will be fulfilled by the lead contact, Marco Trizzino (m.trizzino@imperial.ac.uk).

ACKNOWLEDGMENTS

We thank the Facility for Imaging by Light Microscopy (FILM) at Imperial College London for providing access to instrumentation and technical support, which is part-supported by funding from the Wellcome Trust (grant 104931/Z/14/Z) and BBSRC (grant BB/L015129/1). We also thank the Genomics Facility at The Wistar Institute for the Next Generation Sequencing support. For this work, MT was funded by the Mathers Foundation. We thank Prof. Brian Hendrich (University of Cambridge) and Dr. Andria Koulle (Imperial College London) for insightful discussion of the data and of early versions of the manuscript.

AUTHOR CONTRIBUTIONS

MT designed the project. MD performed most of the experiments. SMB, EVD, ZHM, and LD contributed to some of the experiments. MD and MT analyzed the data and wrote the manuscript. All the authors read and approved the manuscript.

DECLARATION OF INTERESTS

Authors declare no competing interests.

SUPPLEMENTAL INFORMATION

Document S1. Supplemental figures

Table S1. List of AlphaFold predicted contacts between SALL4A and DPF2.

Table S2. List of AlphaFold predicted contacts between SALL4B and DPF2.

FIGURE TITLES AND LEGENDS

Figure 1. SALL4A interacts with the BAF complex through DPF2.

(A) Immunoblot of nuclear protein (5% input) and co-immunoprecipitations (co-IP) of ARID1B and IgG (negative control) from *SALL4*-WT cells at Day-5.

(B) (Left) AlphaFold structure model showing the predicted interactions between SALL4A (red), DPF2 (dark blue), and other BAF components, including SMARCA4, SMARCB1, and ACTL6A. (Center) Visual representation of the confidence score (predicted Local Distance Difference Test - pLDDT) for each residue of the AlphaFold-predicted complex, with color gradients indicating prediction confidence (blue: very high confidence, yellow: low confidence, orange/red: very low confidence). (Right) Close-up view of the SALL4A-DPF2 interface highlighting key interacting residues in red.

(C) SALL4A and SALL4B protein sequences. C2H2 zinc fingers annotations are highlighted in dark purple and numbered. Visual representation of output models of SALL4A (model 0) and SALL4B (models 1, 3, and 4). Predicted interactions between SALL4 isoforms and DPF2 are highlighted in different colors in each model. SALL4A is predicted to interact with DPF2 through the ZFs 4, 5 and 6 (ZFC3) and 7 (ZFC4), light purple. Made with BioRender.com.

(D) Zoomed view of SALL4 (pink) and DPF2 (purple) interacting surfaces. Highlighted are the functional domains involved in the predicted interaction (ZF6 and ZF7 of SALL4 and PHD1 and PHD2 of DPF2). In red, contacts predicted by AlphaFold[34].

(E) Violin plots displaying expression (log2[TPM]) of the four members of the DPF protein family (*PHF10*, *DPF1*, and *DPF3*) in iPSCs (left), neuroectodermal-like cells (Day-5 – center), and in CNCCs (Day-14 – right). Differences between conditions were assessed using unpaired student's t-test.

Figure 2. Generation of a *SALL4*-haploinsufficient human iPSC line

(A) Visual representation of the CRISPR/Cas9 approach used on the *SALL4* gene model. The star represents the exact location of the deletion generated by Cas9. Sanger sequencing confirming the 19 bp deletion in one allele of *SALL4* is displayed. Made with BioRender.com.

(B) Immunoblot for SALL4 in *SALL4*-WT and *SALL4*-het-KO iPSC lines. GAPDH was used as a loading control.

(C) Violin plots displaying expression (log2[TPM]) of the *SALL4* paralogues (*SALL1*, *SALL2*, *SALL3*) in *SALL4*-WT and *SALL4*-het-KO iPSCs. Statistical significance was assessed using unpaired student's t-test.

(D) Immunoblot for SALL1 in *SALL4*-WT and *SALL4*-het-KO iPSC lines. GAPDH was used as a

loading control.

(E) Growth curve comparing *SALL4*-WT with *SALL4*-het-KO iPSC lines. N = 2 biologically

independent experiments. Error bars display standard error of the mean.

(F) Morphology of *SALL4*-WT and *SALL4*-het-KO iPSC colonies at 4x (top) and 10x (bottom).

Figure 3. Stem cell identity is unaffected by SALL4 loss

(A) (Left) Immunofluorescence labeling quantifying the expression of the canonical pluripotency

markers OCT4 and NANOG in the two iPSC lines. Scale bar: 50 μm. (Right) Corrected Total Cell

Fluorescence (CTCF) values were used to quantify and compare markers expression.

(B) Flow cytometry quantifying the expression of surface markers for pluripotency (TRA-60-1-R

and SSEA-4).

(C) Volcano plot of genes differentially expressed in *SALL4*-het-KO relative to *SALL4*-WT in

iPSCs (Day-0). Blue dots (n= 50) represent downregulated genes with p-adj < 0.05 and

log2FoldChange < -1.5. Purple dots (n= 63) represent upregulated genes with p-adj < 0.05 and

log2FoldChange > 1.5.

(D) Top 10 GO term enrichment for downregulated genes in *SALL4*-het-KO iPSCs determined

using WebGestalt[56] over-representation pathway analysis, using affinity propagation as

parameter for redundancy removal. In bold terms with FDR < 0.05).

(E) Top 10 GO term enrichment for upregulated genes in *SALL4*-het-KO iPSCs determined using

WebGestalt[56] over-representation pathway analysis, using affinity propagation as parameter for

redundancy removal. All terms are non-significant (FDR > 0.05).

Figure 4. SALL4 relocates to CNCC enhancers to induce chromatin accessibility

(A) Schematic representation of the iPSC-to-CNCC differentiation protocol. The experimental

timeline shows in vitro differentiation of iPSCs into cranial neural crest cells (CNCCs) via a

neuroectodermal-like stage. Cells were maintained in pluripotency self-renewing conditions until

Day-0, transitioned into differentiation medium starting at Day-0 and further to maintenance

medium at ~ Day-7/8. Key analyses were performed at Day-0, Day-5, and Day-14.

(B) Heatmaps showing the distribution of SALL4 binding at Day-0 and Day-5 across Day-0

specific (left) and Day-5 specific (right) binding regions.

(C) Genomic distribution of SALL4 binding sites relative to the nearest transcription start site

(TSS). Peaks are categorized by distance into six bins: <1 kb, 1–3 kb, 3–5 kb, 5–10 kb, 10–100

kb, and >100 kb. Peaks located at >1kb from the nearest TSS were classified as putative

enhancers.

(D) Gene ontology analysis of Day-5 specific SALL4-bound regions as determined using

WebGestalt[56] over-representation pathway analysis, using affinity propagation as parameter for

redundancy removal (FDR<0.05).

(E) Table of motifs enriched in Day-5 specific SALL4-bound regions as determined using

HOMER[57].

(F) Heatmaps showing ATAC-seq signals at Day-0 (left) and Day-5 (right), at Day-5 specific

SALL4 CUT&RUN peaks. Each row represents an individual peak. Signal intensity is color-coded,

with higher intensity indicating higher chromatin accessibility.

(G) Average profiles showing ATAC-seq signal at Day 5-specific SALL4 CUT&RUN peaks in

SALL4-WT and *SALL4*-het-KO cells in two distinct biological replicates.

Figure 5. SALL4 recruits BAF to CNCC enhancers.

(A) Heatmap and average profiles show BRG1 ChIP-seq signal at the SALL4 Day-5 specific

CUT&RUN peaks (n = 17,137).

(B) Heatmap and average profiles show SALL4 CUT&RUN signal at the genomic sites that lose

BRG1 in *SALL4*-het-KO cells at Day-0 and Day-5 (n = 13,505).

(C) Heatmap and average profiles show BRG1 ChIP-seq signal at the genomic sites that lose

BRG1 in *SALL4*-het-KO cells at day 5 (n = 13,505).

(D) ATAC-seq signal at the genomic sites that lose BRG1 in *SALL4*-het-KO cells (n = 13,505).

(E) and (F) GO term enrichment for genes near BRG1 ChIP-seq peaks lost (E) and gained (F) in

SALL4-het-KO cells. GO terms were determined using WebGestalt[56] over-representation

pathway analysis, using affinity propagation as parameter for redundancy removal (FDR<0.05).

Figure 6. CNCC induction and specification are impaired in *SALL4*-het-KO.

(A) Box plot displaying relative mRNA expression of canonical CNCC markers in *SALL4*-WT and

SALL4-het-KO cells at Day-14. Statistical significance was assessed using unpaired student's t-

test. 18S rRNA was used to normalize samples.

(B) Flow cytometry for the CNCC surface marker CD99 in *SALL4*-WT and *SALL4*-het-KO CNCCs

at Day-14.

(C) Volcano plot of genes differentially expressed in *SALL4*-het-KO relative to *SALL4*-WT in

CNCCs (Day-14). Blue dots (n= 659) represent downregulated genes with p-adj < 0.05 and

log2FoldChange < -1.5. Purple dots (n= 447) represent upregulated genes with p-adj < 0.05 and

log2FoldChange > 1.5.

(D) GO term enrichment for downregulated genes in *SALL4*-het-KO cells determined using

WebGestalt[56] over-representation pathway analysis, using affinity propagation as parameter for

redundancy removal (FDR < 0.05).

(E) Heatmap displaying the expression of canonical epithelial-to-mesenchymal transition (EMT)

markers in *SALL4*-WT and *SALL4*-het-KO.

(F), (G), (H), and (I). Trajectory of stage-specific markers at Day-0, Day-5, and Day-14. Displayed

are the medians of normalized expression (log2[TPMs]) for (F) pluripotency (*POU5F1* and

NANOG), (G) neural plate border (NPB; *ZIC1, GBX2, MSX1, MSX2, PAX3, PAX7, DLX5, DLX6,*

GATA2, GATA3), (H) CNCC specification (*SNAI1, SNAI2, SOX9, TFAP2A*), and (I) post-migratory

CNCC (*SOX5, SOX6, SOX10*) markers at each time-point. Dotted lines represent simple linear

regression.

Figure 7. Working model.

In wild-type conditions, SALL4 recruits BAF at neural crest induction enhancers. This activates

lineage specific gene expression, allowing for NCC induction and specification. In the absence of

SALL4, the NCC induction enhancers remain repressed because BAF is not loaded at chromatin,

and the NCC gene program is not launched, and NCC induction is aborted.

REFERENCES

1.   Jia, Y. kang, Yu, Y., and Guan, L. (2024). Advances in understanding the regulation of pluripotency fate transition in embryonic stem cells. Preprint at Frontiers Media SA, https://doi.org/10.3389/fcell.2024.1494398 https://doi.org/10.3389/fcell.2024.1494398.

2.   Fagan, M.B. (2017). Stem Cell Lineages: Between Cell and Organism. Philosophy, Theory, and Practice in Biology *9*. https://doi.org/10.3998/ptb.6959004.0009.006.

3.   Zheng, Z.H., Sam, T.W., Zeng, Y., Chu, J.J.H., and Loh, Y.H. (2021). Chromatin Regulation in Development: Current Understanding and Approaches. Preprint at Hindawi Limited, https://doi.org/10.1155/2021/8817581 https://doi.org/10.1155/2021/8817581.

4.   Gopinathan, G., and Diekwisch, T.G.H. (2022). Epigenetics and Early Development. Preprint at MDPI, https://doi.org/10.3390/jdb10020026 https://doi.org/10.3390/jdb10020026.

5.   Moore, L.D., Le, T., and Fan, G. (2013). DNA methylation and its basic function. Preprint, https://doi.org/10.1038/npp.2012.112 https://doi.org/10.1038/npp.2012.112.

6.   Li, E. (2002). Chromatin modification and epigenetic reprogramming in mammalian development. Preprint, https://doi.org/10.1038/nrg887 https://doi.org/10.1038/nrg887.

7.   Islam, Z., Ali, A.M., Naik, A., Eldaw, M., Decock, J., and Kolatkar, P.R. (2021). Transcription Factors: The Fulcrum Between Cell Development and Carcinogenesis. Preprint at Frontiers Media S.A., https://doi.org/10.3389/fonc.2021.681377 https://doi.org/10.3389/fonc.2021.681377.

8.   Kim, S., and Shendure, J. (2019). Mechanisms of Interplay between Transcription Factors and the 3D Genome. Preprint at Cell Press, https://doi.org/10.1016/j.molcel.2019.08.010 https://doi.org/10.1016/j.molcel.2019.08.010.

9.   Voss, T.C., and Hager, G.L. (2014). Dynamic regulation of transcriptional states by chromatin and transcription factors. Preprint, https://doi.org/10.1038/nrg3623 https://doi.org/10.1038/nrg3623.

10. Tatetsu, H., Kong, N.R., Chong, G., Amabile, G., Tenen, D.G., and Chai, L. (2016). SALL4, the missing link between stem cells, development and cancer. Preprint at Elsevier B.V., https://doi.org/10.1016/j.gene.2016.02.019 https://doi.org/10.1016/j.gene.2016.02.019.

11. Xiong, J., Todorova, D., Su, N.Y., Kim, J., Lee, P.J., Shen, Z., Briggs, S.P., and Xu, Y. (2015). Stemness factor Sall4 is required for DNA damage response in embryonic stem cells. Journal of Cell Biology *208*, 513–520. https://doi.org/10.1083/jcb.201408106.

12. Rao, S., Zhen, S., Roumiantsev, S., McDonald, L.T., Yuan, G.-C., and Orkin, S.H. (2010). Differential Roles of Sall4 Isoforms in Embryonic Stem Cell Pluripotency. Mol Cell Biol *30*, 5364–5380. https://doi.org/10.1128/mcb.00419-10.

13. Zhang, J., Tam, W.L., Tong, G.Q., Wu, Q., Chan, H.Y., Soh, B.S., Lou, Y., Yang, J., Ma, Y., Chai, L., et al. (2006). Sall4 modulates embryonic stem cell pluripotency and early

embryonic development by the transcriptional regulation of Pou5f1. Nat Cell Biol *8*, 1114–
1123. https://doi.org/10.1038/ncb1481.

14. Yang, J. (2018). SALL4 as a transcriptional and epigenetic regulator in normal and
leukemic hematopoiesis. Preprint at BioMed Central Ltd., https://doi.org/10.1186/s40364-
017-0115-6 https://doi.org/10.1186/s40364-017-0115-6.

15. Miller, A., Ralser, M., Kloet, S.L., Loos, R., Nishinakamura, R., Bertone, P., Vermeulen, M.,
and Hendrich, B. (2016). Sall4 controls differentiation of pluripotent cells independently of
the nucleosome remodelling and deacetylation (NuRD) complex. Development
(Cambridge) *143*, 3074–3084. https://doi.org/10.1242/dev.139113.

16. Wang, F., Zhao, W., Kong, N., Cui, W., and Chai, L. (2014). The next new target in
leukemia: The embryonic stem cell gene SALL4. Mol Cell Oncol *1*.
https://doi.org/10.4161/23723548.2014.969169.

17. Yuri, S., Fujimura, S., Nimura, K., Takeda, N., Toyooka, Y., Fujimura, Y.I., Aburatani, H.,
Ura, K., Koseki, H., Niwa, H., et al. (2009). Sall4 is essential for stabilization, but not for
pluripotency, of embryonic stem cells by repressing aberrant trophectoderm gene
expression. Stem Cells *27*, 796–805. https://doi.org/10.1002/stem.14.

18. Tahara, N., Kawakami, H., Chen, K.Q., Anderson, A., Peterson, M.Y., Gong, W., Shah, P.,
Hayashi, S., Nishinakamura, R., Nakagawa, Y., et al. (2019). Sall4 regulates
neuromesodermal progenitors and their descendants during body elongation in mouse
embryos. Development (Cambridge) *146*. https://doi.org/10.1242/DEV.177659.

19. Chen, K.Q., Anderson, A., Kawakami, H., Kim, J., Barrett, J., and Kawakami, Y. (2022).
Normal embryonic development and neonatal digit regeneration in mice overexpressing a
stem cell factor, Sall4. PLoS One *17*. https://doi.org/10.1371/journal.pone.0267273.

20. Pappas, M.P., Kawakami, H., Corcoran, D., Chen, K.Q., Scott, E.P., Wong, J., Gearhart,
M.D., Nishinakamura, R., Nakagawa, Y., and Kawakami, Y. (2024). Sall4 regulates
posterior trunk mesoderm development by promoting mesodermal gene expression and
repressing neural genes in the mesoderm. Development (Cambridge) *151*.
https://doi.org/10.1242/dev.202649.

21. Kekunnaya, R., and Negalur, M. (2017). Duane retraction syndrome: Causes, effects and
management strategies. Preprint at Dove Medical Press Ltd,
https://doi.org/10.2147/OPTH.S127481 https://doi.org/10.2147/OPTH.S127481.

22. Terhal, P., Rösler, B., and Kohlhase, J. (2006). A family with features overlapping Okihiro
syndrome, hemifacial microsomia and isolated Duane anomaly caused by a novel SALL4
mutation. Am J Med Genet *140 A*, 222–226. https://doi.org/10.1002/ajmg.a.31060.

23. Lu, J., Jeong, H., Kong, N., Yang, Y., Carroll, J., Luo, H.R., Silberstein, L.E., Ma, Y., and
Chai, L. (2009). Stem cell factor SALL4 represses the transcriptions of PTEN and SALL1
through an epigenetic repressor complex. PLoS One *4*.
https://doi.org/10.1371/journal.pone.0005577.

24. Bode, D., Yu, L., Tate, P., Pardo, M., and Choudhary, J. (2016). Characterization of two
distinct nucleosome remodeling and deacetylase (NuRD) complex assemblies in
embryonic stem cells. Molecular and Cellular Proteomics *15*, 878–891.
https://doi.org/10.1074/mcp.M115.053207.

25. Liu, B.H., Jobichen, C., Chia, C.S.B., Chan, T.H.M., Tang, J.P., Chung, T.X.Y., Li, J.,
Poulsen, A., Hung, A.W., Koh-Stenta, X., et al. (2018). Targeting cancer addiction for
SALL4 by shifting its transcriptome with a pharmacologic peptide. Proc Natl Acad Sci U S
A *115*, E7119–E7128. https://doi.org/10.1073/pnas.1801253115.

26. Wang, B., Li, C., Ming, J., Wu, L., Fang, S., Huang, Y., Lin, L., Liu, H., Kuang, J., Zhao, C.,
et al. (2023). The NuRD complex cooperates with SALL4 to orchestrate reprogramming.
Nat Commun *14*. https://doi.org/10.1038/s41467-023-38543-0.

27. Pagliaroli, L., Porazzi, P., Curtis, A.T., Scopa, C., Mikkers, H.M.M., Freund, C., Daxinger,
L., Deliard, S., Welsh, S.A., Offley, S., et al. (2021). Inability to switch from ARID1A-BAF
to ARID1B-BAF impairs exit from pluripotency and commitment towards neural crest
formation in ARID1B-related neurodevelopmental disorders. Nat Commun *12*.
https://doi.org/10.1038/s41467-021-26810-x.

28. Bajpai, R., Chen, D.A., Rada-Iglesias, A., Zhang, J., Xiong, Y., Helms, J., Chang, C.P.,
Zhao, Y., Swigut, T., and Wysocka, J. (2010). CHD7 cooperates with PBAF to control
multipotent neural crest formation. Nature *463*, 958–962.
https://doi.org/10.1038/nature08733.

29. Barnada, S.M., Giner de Gracia, A., Morenilla-Palao, C., López-Cascales, M.T., Scopa, C.,
Waltrich, F.J., Mikkers, H.M.M., Cicardi, M.E., Karlin, J., Trotti, D., et al. (2024). ARID1A-
BAF coordinates ZIC2 genomic occupancy for epithelial-to-mesenchymal transition in
cranial neural crest specification. The American Journal of Human Genetics.
https://doi.org/10.1016/j.ajhg.2024.07.022.

30. Mitchell, Z.H., den Hoed, J., Claassen, W., Demurtas, M., Deelen, L., Campeau, P.M., Liu,
K., Fisher, S.E., and Trizzino, M. (2025). CHD3 regulates BMP signalling response during
cranial neural crest cell specification. Preprint, https://doi.org/10.1101/2025.02.14.638260
https://doi.org/10.1101/2025.02.14.638260.

31. Prescott, S.L., Srinivasan, R., Marchetto, M.C., Grishina, I., Narvaiza, I., Selleri, L., Gage,
F.H., Swigut, T., and Wysocka, J. (2015). Enhancer Divergence and cis-Regulatory
Evolution in the Human and Chimp Neural Crest. Cell *163*, 68–83.
https://doi.org/10.1016/j.cell.2015.08.036.

32. Hodges, C., Kirkland, J.G., and Crabtree, G.R. (2016). The many roles of BAF (mSWI/SNF)
and PBAF complexes in cancer. Cold Spring Harb Perspect Med *6*.
https://doi.org/10.1101/cshperspect.a026930.

33. Zhang, H., Wang, X., Li, J., Shi, R., and Ye, Y. (2021). BAF Complex in Embryonic Stem
Cells and Early Embryonic Development. Preprint at Hindawi Limited,
https://doi.org/10.1155/2021/6668866 https://doi.org/10.1155/2021/6668866.

34.   Jumper, J., Evans, R., Pritzel, A., Green, T., Figurnov, M., Ronneberger, O.,
684   Tunyasuvunakool, K., Bates, R., Žídek, A., Potapenko, A., et al. (2021). Highly accurate
685   protein   structure   prediction   with   AlphaFold.   Nature   *596*,   583–589.
686   https://doi.org/10.1038/s41586-021-03819-2.

687   35.   Zhang, W., Chronis, C., Chen, X., Zhang, H., Spalinskas, R., Pardo, M., Chen, L., Wu, G.,
688   Zhu, Z., Yu, Y., et al. (2019). The BAF and PRC2 Complex Subunits Dpf2 and Eed
689   Antagonistically Converge on Tbx3 to Control ESC Differentiation. Cell Stem Cell *24*, 138-
690   152.e8. https://doi.org/10.1016/j.stem.2018.12.001.

691   36.   Krasteva, V., Crabtree, G.R., and Lessard, J.A. (2017). The BAF45a/PHF10 subunit of
692   SWI/SNF-like chromatin remodeling complexes is essential for hematopoietic stem cell
693   maintenance. Exp Hematol *48*, 58-71.e15. https://doi.org/10.1016/j.exphem.2016.11.008.

694   37.   Lessard, J., Wu, J.I., Ranish, J.A., Wan, M., Winslow, M.M., Staahl, B.T., Wu, H.,
695   Aebersold, R., Graef, I.A., and Crabtree, G.R. (2007). An Essential Switch in Subunit
696   Composition of a Chromatin Remodeling Complex during Neural Development. Neuron *55*,
697   201–215. https://doi.org/10.1016/j.neuron.2007.06.019.

698   38.   Nazim, M., Lin, C.H., Feng, A.C., Xiao, W., Yeom, K.H., Li, M., Daly, A.E., Tan, X., Vu, H.,
699   Ernst, J., et al. (2024). Alternative splicing of a chromatin modifier alters the transcriptional
700   regulatory programs of stem cell maintenance and neuronal differentiation. Cell Stem Cell
701   *31*, 754-771.e6. https://doi.org/10.1016/j.stem.2024.04.001.

702   39.   Rgen Kohlhase, J., Heinrich, M., Schubert, L., Liebers, M., Kispert, A., Laccone, F.,
703   Turnpenny, P., Winter, R.M., and Reardon, W. (2002). Okihiro syndrome is caused by
704   SALL4   mutations.   Oxford   University   Press   *11*,   2979–2987.
705   https://doi.org/https://doi.org/10.1093/hmg/11.23.2979.

706   40.   Al-Baradie, R., Yamada, K., St Hilaire, C., Chan, W.-M., Andrews, C., McIntosh, N.,
707   Nakano, M., Jean Martonyi, E., Raymond, W.R., Okumura, S., et al. (2002). Duane Radial
708   Ray Syndrome (Okihiro Syndrome) Maps to 20q13 and Results from Mutations in SALL4,
709   a New Member of the SAL Family.

710   41.   Pan, Y., van der Watt, P.J., and Kay, S.A. (2023). E-box binding transcription factors in
711   cancer. Preprint at Frontiers Media SA,   https://doi.org/10.3389/fonc.2023.1223208
712   https://doi.org/10.3389/fonc.2023.1223208.

713   42.   Liu, J., Wu, Q., Wang, Y., Wei, Y., Wu, H., Duan, L., Zhang, Q., and Wu, Y. (2018). Ovol2
714   induces mesenchymal–epithelial transition via targeting ZEB1 in osteosarcoma. Onco
715   Targets Ther *11*, 2963–2973. https://doi.org/10.2147/OTT.S157119.

716   43.   Haensel, D., Sun, P., MacLean, A.L., Ma, X., Zhou, Y., Stemmler, M.P., Brabletz, S., Berx,
717   G., Plikus, M. V, Nie, Q., et al. (2019). An Ovol2-Zeb1 transcriptional circuit regulates
718   epithelial   directional   migration   and   proliferation.   EMBO   Rep   *20*.
719   https://doi.org/10.15252/embr.201846273.

720   44.   Kodytková, A., Dušátková, P., Anne Amaratunga, S., Plachý, L., Pruhová, Š., and Lebl, J.
721   (2023). Integrative role of the sall4 gene: From thalidomide embryopathy to genetic defects

of the upper limb, internal organs, cerebral midline, and pituitary. Preprint at S. Karger AG,
https://doi.org/10.1159/000531452 https://doi.org/10.1159/000531452.

45. Miller, M.T. (1991). Thalidomide embryopathy: a model for the study of congenital
incomitant horizontal strabismus. Transaction of the American Ophtalmological Society,
623–674.

46. Kohlhase, J., Chitayat, D., Kotzot, D., Ceylaner, S., Froster, U.G., Fuchs, S., Montgomery,
T., and Rösler, B. (2005). SALL4 mutations in Okihiro syndrome (Duane-radial ray
syndrome), acro-renal-ocular syndrome, and related disorders. Preprint,
https://doi.org/10.1002/humu.20215 https://doi.org/10.1002/humu.20215.

47. Kohlhase, J., Chitayat, D., Kotzot, D., Ceylaner, S., Froster, U.G., Fuchs, S., Montgomery,
T., and Rösler, B. (2005). SALL4 mutations in Okihiro syndrome (Duane-radial ray
syndrome), acro-renal-ocular syndrome, and related disorders. Preprint,
https://doi.org/10.1002/humu.20215 https://doi.org/10.1002/humu.20215.

48. Zhang, Z., Zhan, X., Kim, B., and Wu, J. (2019). A proteomic approach identifies SAFB-
like transcription modulator (SLTM) as a bidirectional regulator of GLI family zinc finger
transcription factors. Journal of Biological Chemistry *294*, 5549–5561.
https://doi.org/10.1074/jbc.RA118.007018.

49. Kong, N.R., Bassal, M.A., Tan, H.K., Kurland, J. V., Yong, K.J., Young, J.J., Yang, Y., Li,
F., Lee, J.D., Liu, Y., et al. (2021). Zinc Finger Protein SALL4 Functions through an AT-
Rich Motif to Regulate Gene Expression. Cell Rep *34*.
https://doi.org/10.1016/j.celrep.2020.108574.

50. Pantier, R., Chhatbar, K., Quante, T., Skourti-Stathaki, K., Cholewa-Waclaw, J., Alston, G.,
Alexander-Howden, B., Lee, H.Y., Cook, A.G., Spruijt, C.G., et al. (2021). SALL4 controls
cell fate in response to DNA base composition. Mol Cell *81*, 845-858.e8.
https://doi.org/10.1016/j.molcel.2020.11.046.

51. Borozdin, W., Boehm, D., Leipoldt, M., Wilhelm, C., Reardon, W., Clayton-Smith, J.,
Becker, K., Mühlendyck, H., Winter, R., Giray, O., et al. (2004). SALL4 deletions are a
common cause of Okihiro and acro-renal-ocular syndromes and confirm haploinsufficiency
as the pathogenic mechanism. J Med Genet *41*. https://doi.org/10.1136/jmg.2004.019901.

52. Ma, X., Huang, R., Li, G., Zhang, T., and Ma, J. (2022). A de novo mutation of SALL4 in a
Chinese family with Okihiro syndrome. Mol Med Rep *25*.
https://doi.org/10.3892/mmr.2022.12647.

53. Donovan KA, A.J.N.R.Y.J.F.E.B.B.E.B.F.E. (2018). Thalidomide promotes degradation of
SALL4, a transcription factor implicated in Duane Radial Ray syndrome. Preprint,
https://doi.org/10.7554/eLife.38430.001 https://doi.org/10.7554/eLife.38430.001.

54. Matyskiela, M.E., Couto, S., Zheng, X., Lu, G., Hui, J., Stamp, K., Drew, C., Ren, Y., Wang,
M., Carpenter, A., et al. (2018). SALL4 mediates teratogenicity as a thalidomide-dependent
cereblon substrate. Nat Chem Biol *14*, 981–987. https://doi.org/10.1038/s41589-018-0129-
x.

55. Anh Vu, K.L., Kumari, K., Hui Liu, B., Gao, C., Li, F., Ping Tang, J., Maddalo, D., Auld,
D.S., Casalena, D.E., Tian, X., et al. (2023). SALL4B, not targeted by IMiD, is important for
SALL4-mediated tumorigenesis. https://doi.org/10.1101/2023.07.07.548071.

56. Liao, Y., Wang, J., Jaehnig, E.J., Shi, Z., and Zhang, B. (2019). WebGestalt 2019: gene
set analysis toolkit with revamped UIs and APIs. Nucleic Acids Res *47*, W199–W205.
https://doi.org/10.1093/nar/gkz401.

57. Heinz, S., Benner, C., Spann, N., Bertolino, E., Lin, Y.C., Laslo, P., Cheng, J.X., Murre, C.,
Singh, H., and Glass, C.K. (2010). Simple Combinations of Lineage-Determining
Transcription Factors Prime cis-Regulatory Elements Required for Macrophage and B Cell
Identities. Mol Cell *38*, 576–589. https://doi.org/10.1016/j.molcel.2010.05.004.

58. Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T.,
Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B., et al. (2012). Fiji: An open-source
platform for biological-image analysis. Preprint, https://doi.org/10.1038/nmeth.2019
https://doi.org/10.1038/nmeth.2019.

59. Love, M.I., Huber, W., and Anders, S. (2014). Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. Genome Biol *15*.
https://doi.org/10.1186/s13059-014-0550-8.

60. Quinlan, A.R., and Hall, I.M. (2010). BEDTools: A flexible suite of utilities for comparing
genomic features. Bioinformatics *26*, 841–842.
https://doi.org/10.1093/bioinformatics/btq033.

61. Ramírez, F., Ryan, D.P., Grüning, B., Bhardwaj, V., Kilpert, F., Richter, A.S., Heyne, S.,
Dündar, F., and Manke, T. (2016). deepTools2: a next generation web server for deep-
sequencing data analysis. Nucleic Acids Res *44*, W160–W165.
https://doi.org/10.1093/NAR/GKW257.

62. Herbreteau, G., Airaud, F., Pierre-Noël, E., Vallée, A., Bézieau, S., Théoleyre, S., Blons,
H., Garinet, S., and Denis, M.G. (2021). MEM: An algorithm for the reliable detection of
microsatellite instability (MSI) on a small NGS panel in colorectal cancer. Cancers (Basel)
13. https://doi.org/10.3390/cancers13164203.

63. Zhang, Y., Liu, T., Meyer, C.A., Eeckhoute, J., Johnson, D.S., Bernstein, B.E., Nussbaum,
C., Myers, R.M., Brown, M., Li, W., et al. (2008). Model-based analysis of ChIP-Seq
(MACS). Genome Biol *9*. https://doi.org/10.1186/gb-2008-9-9-r137.

64. Polit, L., Kerdivel, G., Gregoricchio, S., Esposito, M., Guillouf, C., and Boeva, V. (2021).
CHIPIN: ChIP-seq inter-sample normalization based on signal invariance across
transcriptionally constant genes. BMC Bioinformatics *22*. https://doi.org/10.1186/s12859-
021-04320-3.

METHODS

KEY RESOURCES TABLE

REAGENT OR RESOURCE	SOURCE	IDENTIFIER
Antibodies		
Brg1 (D1Q7F)	Cell Signaling Technology	AB_2728743
CD99 Antibody, anti-human, Vio® Bright B515, REAfinity™	Miltenyi Biotec	AB_2784331
GAPDH (D16H11) XP Rabbit mAb	Cell Signaling Technology	AB_10622025
Human Nanog Antibody	R&D Systems	AB_355097
Human/Mouse Oct-3/4 Antibody	R&D Systems	AB_354975
SALL1 Polyclonal Antibody	Thermo Fisher Scientific	AB_2900498
Sall4 (EE-30)	Santa Cruz Biotechnology	AB_1129262
Sall4 (D16H12) Rabbit mAb	Cell Signaling Technology	AB_10949321
APC anti-human SSEA-4	Biolegend	AB_2616818
PE anti-human TRA-1-60-R	Biolegend	AB_1279447
Rabbit IgG, monoclonal [EPR25A] Antibody	Abcam	AB_2687931
AP2 alpha Monoclonal Antibody (3B5)	Thermo Fisher Scientific	AB_2199412
CD326 (EpCAM) Monoclonal Antibody (1B7), eBioscience	Thermo Fisher Scientific	AB_795876
Anti-SOX9 antibody	Abcam	AB_2715497
Vimentin Polyclonal Antibody	Thermo Fisher Scientific	AB_2216267
ARID1B antibody	Abcam	AB_2243092
IRDye 800CW Goat anti-Rabbit IgG	LI-COR Biosciences	AB_621843
Goat polyclonal Secondary Antibody to Mouse (IRDye® 800CW)	LI-COR Biosciences	AB_10859231

Donkey anti-Rabbit IgG (H+L) Highly Cross-Adsorbed Secondary Antibody, Alexa Fluor™ Plus 488	Thermo Fisher Scientific	AB_2762833
Donkey anti-Mouse IgG (H+L) Highly Cross-Adsorbed Secondary Antibody, Alexa Fluor™ Plus 647	Thermo Fisher Scientific	AB_2762830
Goat anti-Chicken IgY (H+L) Secondary Antibody, Alexa Fluor™ 488	Thermo Fisher Scientific	AB_2534096
Critical commercial assays		
ATAC-Seq Kit	Active Motif	53150
Human Pluripotent Stem Cell Functional Identification Kit	R&D Systems	SC027B
Monarch® Total RNA Miniprep Kit	New England Biolabs	T2010S
Maxima First Strand cDNA Synthesis Kit for RT-qPCR	Thermo Fisher	K1641
NEBNext® Ultra™ II Directional RNA Library Prep Kit for Illumina®	New England Biolabs	E7760L
CUT&RUN Assay Kit	Cell Signaling Technology	86652
NEBNext® Ultra™ II DNA Library Prep Kit for Illumina®	Cell Signaling Technology	E7645L
Experimental models: Cell lines		
PENN123i-SV20	iPSC Core of the University of Pennsylvania	CVCL_EL23
PENN123i-SV20 *SALL4*-het-KO iPSC	Synthego	
Oligonucleotides		
18S	5'-ATACATGCCGACGGGCGCTG-3'	5'-AGGGCTGACCGGGTTGGTT-3'
TFAP2A	5'-GCCTCTCGCTCCTCAGCTCC-3'	5'-CGTTGGCAGCTTTACGTCTCCC-3'
NR2F1	5'-ATCGTGCTGTTCACGTCAGA-3'	5'-GCTCCTCACGTACTCCTCCA-3'

SOX9	5'-GTACCCGCACTTGCACAAC-3'	5'-TCTCFCTCTCGTTCAGAAGTC-3'
SALL1	5'-CCCAGTGTGTGTCCTGTAGT-3'	5'-CCACCTTCACCAACCCTTTG-3'
SALL2	5'-TAGGTGTTGTGTAGGCTGGG-3'	5'-CTCACTTCCATTCCCACCCT-3'
SALL3	5'-GCGCTGCGTAAAGATGAGAA-3'	5'-GGCTACCCTGTCTGTTGAGT-3'
SALL4	5'-GGCATCCAGAGACAGACCTT-3'	5'-GGTCCTCGAGCAGATCTTGT-3'
Software and algorithms		
AlphaFold3	https://alphafoldserver.com/	SCR_025454
TrimGalore!	https://www.bioinformatics.babraham.ac.uk/projects/trim_galore/	SCR_011847
FlowJo Software version 10.9		SCR_008520
BEDTools		SCR_006646
Deeptools		SCR_016366
R Studio version 4.4.3		
GraphPad Prism 10.2.3	http://www.graphpad.com/faq/viewfaq.cfm?faq=1362	SCR_002798
DESeq2		SCR_015687

METHOD DETAILS

AlphaFold3 structure prediction

FASTA files for each BAF subunits and SALL4 isoforms were collected from the "Sequence &

Isoforms" section of Uniprot (www.uniprot.org). The AT-rich DNA sequences was collected from

an empirically reported crystal structure of SALL4 zinc finger cluster 4 bound to DNA on RCSB

PDB (www.rcsb.org, 8A4I – DOI: https://doi.org/10.2210/pdb8A4I/pdb). FASTA sequences are

listed in Supplementary Table 2. AlphaFold outputs were downloaded and .cif files were analysed

using UCSF ChimeraX-1.9 (https://www.rbvi.ucsf.edu/chimerax). Contacts were found using

default parameters and center-center distance ≤ 8 Å.

Human iPSC culture

Wild-type iPSC lines were obtained from the iPSC Core of the University of Pennsylvania (Control

line-1: SV20 line, male, age 43). iPSC lines were cultured in feeder-free, serum-free mTeSR Plus

medium (STEMCELL Technologies) with 1% Penicillin-Streptomycin (5,000 U/mL; Gibco

#15140122) on tissue culture plates coated overnight with Geltrex™ LDEV-Free hESC-qualified

Reduced Growth Factor Basement Membrane Matrix (Fisher-Scientific). Cells were passaged

~1:8 at 60-80% confluency using 0.5mM EDTA to create 50-200 cell clusters.

CRISPR/Cas9-mediated SALL4 knock-out

822   The heterozygous *SALL4* knock-out in the human SV20 iPSC line was generated via

823   CRISPR/Cas9 gene-editing by the biotech company Synthego. Specifically, Cas9 targeted the

824   first exon of the *SALL4* gene in the established SV20 iPSC line producing an 11-base deletion

825   (c.230delGGGCCGCATCT) which generated a premature stop codon.

Trilineage differentiation

828   iPSCs were differentiated into the three germ layers (endoderm, ectoderm and mesoderm) using

829   the Human Pluripotent Stem Cell Functional Identification Kit (R&D Systems, SC027B). The

830   expression of relevant markers was then assessed using immunofluorescence with 10 ug/ml of

831   antibodies provided in the kit (Goat anti-human SOX17, Goat anti-human OTX2 and, Goat anti-

832   human Brachyury).

iPSC-to-CNCC differentiation

All iPSC lines were differentiated into CNCCs as previously described (Prescott et al., 2015).

Briefly, iPSCs at 60% confluence were treated with CNCC differentiation media (1:1 Neurobasal

medium/D-MEM F-12 medium (Invitrogen), 0.5× B-27 supplement with Vitamin A (50× stock,

Gibco, #17504044), 0.5× N-2 supplement (100× stock, Gibco; #17502001), 20 ng/ml human FGF

(Gibco), 20 ng/ml recombinant EGF (Sigma-Aldrich), 5 µg/ml bovine insulin (Sigma-Aldrich) and

1× Glutamax-I supplement (100× stock, Invitrogen). Medium (2 ml) was changed every other day.

Around day 7-9 of differentiations, cells were transitioned to CNCC early maintenance media (1:1

Neurobasal medium/D-MEM F-12 medium (Invitrogen), 0.5× B-27 supplement with Vitamin A

(50× stock, Invitrogen), 0.5× N-2 supplement (100× stock, Invitrogen), 20 ng/ml bFGF

(Biolegend), 20 ng/ml EGF (Sigma-Aldrich), 1 mg/ml bovine serum albumin, serum replacement

grade (Gemini Bio-Products # 700-104 P) and 1× Glutamax-I supplement (100× stock,

Invitrogen). Differentiating cells were passaged ~1:6 with 0.5mM EDTA or Accutase Cell

Detachment Solution (Stemcell Technologies; #07920), plated in a fresh, Geltrex-coated 6-well

plates. After passaging, cells were treated with 0.05% ROCK-inhibitor (Y27632; Stemcell

Technologies) for 24 hours.

Real-time quantitative polymerase chain reaction (RT-qPCR)

Cells were harvested as single cells, and the total RNA was extracted using Monarch® Total RNA Miniprep Kit (New England Biolabs) following manufacturer's instructions. 600ng of total RNA was used to synthesize cDNA using Maxima First Strand cDNA Synthesis Kit for RT-qPCR (K1641, Thermo Fisher). For each RT-qPCR reaction, 15 ng of cDNA was used with 0.1 μM of each primer, 10 μL of PowerUp™ SYBR™ Green Master Mix (Applied Biosystems) in a final volume of 20 μl. Samples were analyzed with CFX Connect Real-Time PCR Detection System (Bio-Rad) using the following thermal cycling parameters: 3 min at 95 °C, followed by 40 cycles of 10 s at 95 °C, 20 s at 63 °C followed by 30 s at 72 °C. Each sample was run in triplicate. 18 S rRNA was used as normalizer.

Nuclear Fractionation and Co-immunoprecipitation

Cells were washed twice with ice-cold PBS, scraped, and collected by centrifugation at 10,000 × g for 1 minute. Pellets were resuspended in 500 μL of Buffer A (10mM HEPES, 1.5mM Mg2Cl,10mM KCl, 1mM EDTA, 0.5mM DTT) and cytoplasmic fractions were obtained by incubating the cells on ice for 15 minutes, adding of 0.05% NP-40, followed by centrifugation at 17,000 × g for 10 minutes at 4 °C. The resulting pellet was gently washed with Buffer A and resuspended in Buffer C (10mM HEPES, 1.5mM Mg2Cl, 300mM NaCl, 0.2mM EDTA, 0.5mM DTT, 26% glycerol). Samples were incubated at 4 °C with shaking at 3,400 rpm for 1 hour, with vortexing every 10 minutes. Nuclear fractions were collected by centrifugation at 17,000 × g for 10 minutes at 4 °C.

For co-immunoprecipitation (co-IP), 25 μL of Dynabeads™ Protein A (ThermoFisher, Cat# 10002D) were washed twice with PBS containing 0.1% Tween-20 (PBS-T), resuspended in 1 mL of PBS-T, and incubated with 2 μg of the appropriate antibody for 1 hour at 4 °C with rotation. Bead-antibody complexes were washed twice with PBS-T, resuspended in 125 μL PBS-T, and incubated with 100 μg of nuclear extract for 3 hours at 4 °C with rotation. Beads were then washed five times with Buffer C and eluted in 25 μL of Buffer C supplemented with 4.1 μL of 6× reducing Laemmli SDS sample buffer (ThermoFisher Scientific). Eluates and 5% input controls (nuclear extract diluted 1:10 in PBS and mixed with 1:6 volume of 6× Laemmli buffer) were boiled for 5 minutes prior to SDS-PAGE. For Western blotting, 20 μL of each sample were loaded as described below.

Western Blot

For each condition, cells were harvested and lysed with cold radioimmunoprecipitation assay
buffer (Pierce™ RIPA buffer, Thermo Fisher) with Pierce™ Protease Inhibitor Mini Tablets, EDTA-
free (Thermo Fisher). After centrifugation at 4°C at 16,000x g for 10 minutes, the total extracted
protein was quantified using Pierce™ BCA Protein Assay Kit, following manufacturer's
indications. Thirty micrograms of total protein were loaded in Novex WedgeWell 4–20% Tris-
Glycine Gel (Invitrogen) and separated through gel electrophoresis (SDS–PAGE) Tris-Glycine-
SDS buffer (Invitrogen). The proteins were then transferred to nitrocellulose membrane (Thermo
Fisher Scientific) and incubated for one hour in Intercept® (PBS) Blocking Buffer (Licor). After
blocking, membranes were incubated overnight at 4°C (while rocking) in suitable antibody diluted
in blocking buffer and 0.2% Tween-20, at recommended concentrations. Membranes were
washed in 0.1% PBS-T before and after being incubated for one hour in the appropriate secondary
antibody (1:15,000 dilution). The antibodies were then visualized using Odyssey XF Imaging
System (Licor) or ChemiDocTM MP Imaging System and imaged with Image Studio Software
(Licor) or Fiji[58].

Flow Cytometry analysis of surface markers
To obtain a single-cell suspension for flow cytometry analysis, cells were treated with Accutase
for 5 min at 37°C. Cells were then washed with cold phosphate-buffered saline (PBS)-2% fetal
bovine serum (FBS) and live cells were counted. 1×10^6 cells/condition were resuspended in
100 μL PBS-2% FBS and stained. For pluripotency evaluation, 4 μl of the respective antibodies
were used. For analysis of differentiation, 2 μl of the respective antibodies were used. Cells were
incubated for 15 min on ice and protected from light, before transferring them into FACS tubes
containing an additional 300 μL PBS-2% FBS. Flow cytometry data were acquired using
NovoCyte Penteon Flow Cytometer Systems 5 Lasers and analyzed with FlowJo Software version
10.9.

Immunofluorescence
Cells were passaged onto Geltrex-coated coverslips and fixed at required confluence or time-
point. In detail, cells were incubated in 4% PFA at 37°C for 15 minutes, followed by three 2-minute
washes in 1× PBS. For permeabilization, cells were treated with 0.1% (v/v) Triton X-100 (T8787,
Sigma-Aldrich) at room temperature for 10 minutes before three 2-minute washes in 1× PBS +
0.1% (v/v) Tween 20 and a 1-hour blocking upon 10% Donkey serum (D9663, Sigma-Aldrich) at
room temperature. Cells were incubated with corresponding PBS-diluted primary antibodies
protected from direct light at 4°C overnight. After three 5-minute washes with 1× PBS + 0.1%

Tween 20, cells were incubated for 1 hour in PBS-diluted secondary antibodies and 15 minutes
with DAPI (1:5,000; 422801, BioLegend). Cells were then washed three times with 1× PBS +
0.1% Tween for 5 minutes and once with 1× PBS. Coverslips were then fetched and mounted
onto slides using Fluorescence Mounting Medium (S3023, Agilent Technologies). Slides were
imaged using confocal microscope (LSM780, Zeiss).

RNA-seq
Total RNA was extracted using Monarch® Total RNA Miniprep Kit (New England Biolabs,
T2010S) according to the manufacturer's instructions. RNA was quantified using NanoDrop 2000
UV Visible Spectrophotometer, while the RNA integrity was checked on Tapestation 2200
(Agilent). Only samples with RIN value above 8.0 were used for transcriptome analysis. RNA
libraries were prepared using 1 µg of total RNA input using NEBNext® Poly(A) mRNA Magnetic
Isolation Module, NEBNext® UltraTM II Directional RNA Library Prep Kit for Illumina® according
to the manufacturer's instructions (New England Biolabs).

RNA-sequencing analysis
After   removing   the   adapters   with   TrimGalore!
(https://www.bioinformatics.babraham.ac.uk/projects/trim_galore/) , Kallisto was used to quantify
reads mapping to each gene. Differential gene expression levels were determined using
DESeq2[59]. All statistical analyses were performed using the latest versions of BEDTools[60],
deepTools[61], R Studio (version 4.4.3; Trophy Case), and GraphPad Prism software version 10.2.3
(http://www.graphpad.com/faq/viewfaq.cfm?faq=1362; GraphPad, San Diego, CA).

CUT&RUN-seq
For each condition, 1x 10^5 cells were processed using CUT&RUN Assay Kit #86652 (Cell
Signaling Technology). Briefly, each sample was resuspended in Concanavalin A Beads and
treated with an Antibody Binding Buffer and the suitable antibody. After the antibody incubation,
each reaction underwent a step of pAG-MNase Enzyme binding, and subsequent DNA Digestion
and diffusion. The diffused DNA was then purified using Phenol/Chloroform Extraction and
Ethanol Precipitation. The purified DNA was quantified using QuantiFluor dsDNA System, using
the ONE dsDNA protocol (Promega). Libraries for sequencing were prepared using NEBNext
Ultra II DNA Library Prep Kit for Illumina (New England BioLabs) and the libraries size distribution
was assessed through automated electrophoresis using the Agilent 2200 Tapestation System
(Agilent Technology).

ChIP-seq

All samples from different conditions were processed together to prevent batch effects. Approximately 15 million cells were double cross-linked with 2mM disuccinimidyl glutarate (DSG) for 20 minutes followed by 1% formaldehyde for 10 minutes at room temperature, quenched with 125 mM glycine, harvested, and washed twice with 1X PBS. The fixed cell pellet was resuspended in ChIP lysis buffer (150 mM NaCl, 1% Triton X-100, 0.7% SDS, 500 μM DTT, 10 mM Tris-HCl, 5 mM EDTA) and chromatin was sheared to an average length of 200–900 base-pairs, using a Covaris S220 Ultrasonicator at 5% duty factor for 7 minutes for 3 times. The chromatin lysate was diluted with SDS-free ChIP lysis buffer. 10μg of antibody was used for BRG1. The antibody was added to at least 5μg of sonicated chromatin along with Dynabeads Protein G magnetic beads (Invitrogen) and incubated with rotation at 4°C overnight. The beads were washed twice with each of the following buffers: Mixed Micelle Buffer (150 mM NaCl, 1% Triton X- 100, 0.2% SDS, 20 mM Tris-HCl, 5 mM EDTA, 65% sucrose), Buffer 200 (200 mM NaCl, 1% Triton X-100, 0.1% sodium deoxycholate, 25 mM HEPES, 10 mM Tris-HCl, 1 mM EDTA), LiCl detergent wash (250 mM LiCl, 0.5% sodium deoxycholate, 0.5% NP-40, 10 mM Tris-HCl, 1 mM EDTA) and a final wash was performed with cold 0.1X TE. Finally, beads were resuspended in 1X TE containing 1% SDS and incubated at 65°C for 10 min to elute immunocomplexes. The elution was repeated twice, and the samples were incubated overnight at 65°C to reverse cross-linking, along with the input (5% of the starting material). The DNA was digested with 0.5 mg/ml Proteinase K for 3 hours at 55°C and then purified using the ChIP DNA Clean & Concentrator kit (Zymo) and quantified with QUBIT. Barcoded libraries were made with NEBNext Ultra II DNA Library Prep Kit for Illumina) using NEBNext Multiplex Oligos Dual Index Primers for Illumina (New England BioLabs) and sequenced on NovaSeq X Plus instrument (Illumina).

ATAC-seq

One hundred thousand cells/condition were harvested by enzymatic dissociation using StemPro™ Accutase™ (Thermo Fisher Scientific). Each cell pellet was processed using ATAC-Seq Kit (Active Motif), following the manufacturer's directions. Briefly, the fresh cell pellets were washed in 1X PBS and lysate in ATAC Lysis Buffer. Each lysate underwent a process of tagmentation and purification, before being amplified through PCR, using the following thermal cycler parameters: 72°C for 5 min, 98°C for 30 sec, followed by 10 cycles of 98°C for 10 sec, 63°C for 30 sec, 72°C for 1 min. Each PCR reaction was cleaned up using SPRI beads (1.2X of

the sample volume). The Agilent 2200 Tapestation System (Agilent Technologies) was used as

quality control tool before performing next generation sequencing (NGS).

Bioinformatics analysis of CHIP-seq, ATAC-seq and CUT&RUN data

Adapters   were   removed   with   TrimGalore!

(https://www.bioinformatics.babraham.ac.uk/projects/trim_galore/), the sequences were aligned

to the reference hg19, using Burrows-Wheeler Alignment tool, with the MEM algorithm[62]. Uniquely

mapping aligned reads were filtered based on mapping quality (MAPQ > 10) to restrict our

analysis to higher quality and uniquely mapped reads, and PCR duplicates were removed.

MACS2[63] was used to call peaks using the default parameters at 5% FDR.

Motif analysis

BED files for the regions of interest were produced using BEDTools[60]. Motif analysis was

performed using HOMER[57]. Shuffled input sequences were used as background. E-values <

0.001 were used as a threshold for significance.

1000

QUANTIFICATION AND STATISTICAL ANALYSIS

RT-qPCR and image quantification data were analyzed using Prism software version 10.2.3

(http://www.graphpad.com/faq/viewfaq.cfm?faq=1362; GraphPad, San Diego, CA). Data is

represented as mean ± standard error of mean (SEM). GO enrichment analysis was performed

using Webgestalt 2019[56]. All statistical analyses were performed using the latest versions of

BEDTools[60], Deeptools[61], R version 4.4.3 (Throphy Case; https://www.R-project.org/), and Prism

software version 10.2.3 (http://www.graphpad.com/faq/viewfaq.cfm?faq=1362; GraphPad, San

Diego, CA). In particular, Bigwigs were generated with Deeptools and normalized by sequencing

depth using the same software, and upper-quantile normalized using CHIPIN[64].

1010

1011

FIG 1

A

5% INPUT   IP:ARID1B   IP:IgG

SALL4A ——————————— 165 kDa

SALL4B ——————————— 95 kDa

α-SALL4

B

SALL4A
SMARCA4
DPF2
SMARCB1
ACTL6A
ARID1B

SALL4-DPF2 contacts

Very high (pIDDT > 90)   Confident (90 > pIDDT > 70)   Low (70 > pIDDT > 50)   Very low (pIDDT < 50)

C

ZFC1   ZFC2   ZFC3   ZFC4

SALL4A   1   2 3   4 5 6   7 8

SALL4B   1   7 8

Model 1
Model 3
Model 4

D

ZF6   ZF5

PHD1   PHD2

E

Day-0   **Day-5**   **Day-14**

Log2(TPM)

PHF10   DPF1   DPF3

0.580861   0.580861   0.580861

0.880699   0.880699   0.880699

0.182706   0.182706   0.182706

WT
SALL4-het-KO

A

WT

CONTROL SAMPLE 197 TO 262 BP

SALL4-het-KO

EDITED SAMPLE 198 TO 263 BP

TRUNCATED SALL4

B

α-SALL4 — 165 kDa

α-GAPDH — 37 kDa

D

α-SALL1 — 140 kDa

α-GAPDH — 37 kDa

C

Log2(TPM)

0.002659   0.602814   0.602814

SALL1   SALL2   SALL3

WT
SALL4-het-KO

E

WT   *SALL4*-het-KO

F

Growth Curve

Log2(Mln Alive Cells/ml)

Days

WT
SALL4-het-KO

FIG 3

A

WT / *SALL4*-het-KO

DAPI/NANOG

DAPI/OCT4

IF quantification - NANOG

IF quantification - OCT4

B

TRA-60-1-R (PE)

SSEA-4 (APC)

C

Differentially Expressed Genes – Day-0

50
downregulated

63
upregulated

NANOG

POU5F1

total = 17112 variables

Top 10 Gene Ontology Terms – Day-0

D   Downregulated in *SALL4*-het-KO

- **membrane raft organization**
- regulation of epithelial cell differentiation
- **multi-organism reproductive process**
- **multi-multicellular organism process**
- **organic hydroxy compound transport**
- carbohydrate homeostasis
- **hormone transport**
- epidermis development
- negative regulation of transport
- **signal release**

Enrichment Ratio

E   Upregulated in *SALL4*-het-KO

- establishment of tissue polarity
- midbrain development
- regulation of animal organ morphogenesis
- cytoplasmic translation
- morphogenesis of a branching structure
- protein processing
- stem cell differentiation
- epithelial cell proliferation
- gland development
- synapse organization

Enrichment Ratio

FIG4

A

B

SALL4 CUT&RUN

C

SALL4 stage-specific binding distribution

D

Top 10 Gene Ontology Terms
Day-5 specific

F

ATAC-seq

E

SALL4 CUT&RUN Day-5 specific motifs		
TF	**Motif Consensus**	**p-value**
TEAD		$1e^{-1356}$
JUN/FOS		$1e^{-938}$
SOX9		$1e^{-817}$
ZIC		$1e^{-498}$
TCF/TWIST		$1e^{-132}$
TFAP2A		$1e^{-90}$

G

ATAC-seq at SALL4 Day 5-specific regions

A CNCC Markers – Day-14

B CNCC – Day-14

C Differentially Expressed Genes – Day-14

659 downregulated

447 upregulated

D Downregulated Genes in *SALL4*-het-KO Day-14

face development
artery development
regulation of animal organ morphogenesis
cardiac chamber development
autonomic nervous system development
exocrine system development
gastrulation
extracellular structure organization
tissue migration
mesenchyme development

E EMT markers - CNCC

F Pluripotency Markers

G Neural Plate Border Markers

H CNCC Specification Markers

I Post-migratory CNCC Markers

FIG 7

PLURIPOTENT CELLS

NEUROECTODERMAL
CELLS

CRANIAL NEURAL CREST
CELLS

Supplementary Figures

A

Supplementary Figure 1.
A. Immunofluorescence for ectodermal (OTX2), mesodermal (Brachyury), and endodermal (SOX17) markers in *SALL4*-WT and *SALL4*-het-KO after trilineage differentiation. Scale bar: 100 μm.

A

SALL4 CUT&RUN Day-0 specific motifs		
Transcription Factor	**Motif Consensus**	**p-value**
CTCF/BORIS

 | $1e^{-461}$ |
bHLH		$1e^{-106}$
OCT:OCT (MORE)		$1e^{-103}$
ASCL1/2		$1e^{-67}$
OCT4-NANOG-SOX2		$1e^{-36}$

B

Differentially Expressed Genes – Day-5

Supplementary Figure 2.
(A) Table of motifs enriched in Day-0 specific SALL4-bound regions as determined using HOMER[63].
(B) Volcano plot of genes differentially expressed in *SALL4*-het-KO relative to *SALL4*-WT in neuroectodermal-like cells (Day-5). Blue dots (n= 126) represent downregulated genes with p-adj < 0.05 and log2FoldChange < -1.5. Purple dots (n= 44) represent upregulated genes with p-adj < 0.05 and log2FoldChange > 1.5.