Cell-free RNA reveals host and microbial correlates of broadly neutralizing

antibody development against HIV

Mark Kowarsky[1], Mercedes Dalman[2,3], Mira N Moufarrej[4], Jennifer Okamoto[5], Yike Xie[2,3],

Norma F Neff[5], Salim S Abdool Karim[6,7], Nigel Garrett[6,8,9], Penny L Moore[6,10,11,12,†], Joan

Camunas-Soler [2,3,13,†], Stephen R Quake[4,5,†]

[1]Department of Physics, Stanford University, Stanford, CA

[2]Department of Medical Biochemistry and Cell Biology, Institute of Biomedicine, University of Gothenburg,

Sweden

[3]Wallenberg Centre for Molecular and Translational Medicine, Sahlgrenska Academy, University of

Gothenburg, Sweden

[4]Department of Bioengineering, Stanford University, Stanford, CA

[5]Chan Zuckerberg Biohub, San Francisco, CA

[6]Centre for the AIDS Programme of Research in South Africa (CAPRISA), University of KwaZulu Natal,

Durban, South Africa

[7]Department of Epidemiology, Columbia University, New York, NY 10032, USA

[8] Desmond Tutu HIV Centre, University of Cape Town, Cape Town, South Africa

[9] Discipline of Public Health Medicine, School of Nursing and Public Health, University of KwaZulu-Natal,

Durban, South Africa

[10] SA MRC Antibody Immunity Research Unit, Faculty of Health Sciences, University of the Witwatersrand,

Johannesburg, South Africa

[11] Centre for HIV and STIs, National Institute for Communicable Diseases (NICD) of the National Health

Laboratory Service (NHLS), Johannesburg, South Africa.

[12] Wits Infectious Diseases and Oncology Research Institute, Faculty of Health Sciences, University of the

Witwatersrand, Johannesburg, South Africa

[13] Science for Life Laboratory, Institute of Biomedicine, University of Gothenburg, Gothenburg, Sweden

†To whom correspondence should be addressed: steve@quake-lab.org, pennym@nicd.ac.za,

joan.camunas@gu.se

Abstract:

A small number of people living with HIV (PLWH) develop broadly neutralizing antibodies (bNAbs) targeting multiple HIV strains. Although several viral and immune factors contribute to bNAb development, the genetic and environmental factors driving this response remain largely unknown. We performed combined cell-free DNA (cfDNA) and cell-free RNA (cfRNA) sequencing in 42 plasma samples from a longitudinal cohort of 14 PLWH (7 who develop bNAbs and 7 matched controls). This approach enabled us to non-invasively monitor the host transcriptome, viral genetic variation, and microbiome composition during HIV infection, and to identify molecular correlates of bNAb development. We find that development of bNAbs is associated with a transcriptomic signature of early immune activation characterized by elevated levels of MHC class I antigen presentation genes. This signature is independent of viral load or CD4 count and declines over time. In addition to host features, we recovered sufficient viral reads to reconstruct HIV consensus sequences, supporting the utility of cfRNA for viral genotyping. Finally, we also identified an enrichment of several microbial taxa in bNAb producers and increased levels of GB virus C (GBV-C), a non-pathogenic lymphotropic virus. Our findings suggest a distinct early immune activation profile in PLWH who develop bNAbs. More broadly, we show that combined cfDNA/cfRNA sequencing can reveal relationships between a protective immunogenic response to HIV infection, the host immune system, and microbiome, highlighting its potential for biomarker discovery in future vaccine and therapeutic studies.

Author Summary

A subset of people living with HIV develop broadly neutralizing antibodies (bNAbs), a phenomenon that remains poorly understood but relevant to vaccine development. We applied

combined cell-free RNA and DNA sequencing to 42 longitudinal plasma samples (7 bNAb producers, 7 controls) to monitor the host transcriptome, viral variation, and microbiome composition during bNAb development. Early in infection, bNAb producers showed an MHC class I-linked immune activation signature that declined over time, alongside changes in commensal microbial sequences and increased levels of GB virus C in cfRNA. These findings support the use of cfDNA/cfRNA sequencing for biomarker discovery in HIV research.

Introduction

HIV has infected over 36 million people worldwide, reaching epidemic proportions in Eastern and Southern Africa with an estimated adult prevalence close to 7% (1). Antiretroviral therapy (ART) has greatly improved outcomes, allowing HIV to be a manageable condition throughout life. However, global access to ART is difficult to achieve and also requires lifelong daily adherence to a pill taking regime. Moreover, despite effective treatment, transmission of HIV remains common in vulnerable populations, and people living with HIV (PLWH) are at increased risk of aging-related and chronic diseases (2).

Although virtually all PLWH develop strain-specific autologous responses (3–6), few develop broadly neutralizing antibodies (bNAbs), which can neutralize multiple strains of the virus (7), and when administered passively can prevent acquisition of susceptible HIV strains and suppress viral replication in PLWH (8). There is great interest in understanding the drivers of bNAbs, as this could inform the development of an effective vaccine. Efforts to design an effective HIV vaccine have increasingly focused on strategies to induce bNAbs by guiding B-cell affinity maturation, with studies demonstrating promising immunization approaches through the rational design of immunogens (9–11). Despite these advances, which specific cellular processes give rise to bNAbs during infection remains incompletely described. So far,

there is evidence that viral and host genetic factors might play a role (12–16), and that a

combination of high viral titres and reduced CD4 counts early in infection is associated with

the development of bNAbs in some individuals (17,18). Additionally, changes in frequency of

certain immune subpopulations, such as T follicular helper cells (Tfh) might also influence

affinity maturation towards bNAb production (19–23). Moreover, there is some evidence that

bNAbs might arise from pre-existing memory B cells initially stimulated by bacterial antigens

-likely from the gut microbiome- which later cross-react with the HIV glycan shield and

develop neutralizing capabilities (24,25). Although microbial antigen cross-reactivity has also

been shown to lead to non-protective immune responses in HIV vaccine trials (26), a better

understanding of the interplay between the host immune response, microbiome and a protective

bNAb response could aid in vaccine development.

Sequencing of plasma cell-free DNA (cfDNA) has become an established tool in clinical

medicine, with applications ranging from non-invasive prenatal testing and oncology to organ

transplant monitoring (27,28). In contrast, plasma cell-free RNA (cfRNA) provides an

emerging complementary approach to measure dynamic changes in the host transcriptome (29).

Because transcripts released into circulation can originate from many different cell types and

tissues (30), cfRNA enables non-invasive monitoring of systemic physiological and immune

states (31,32). To date, cfRNA sequencing has been used in diverse biomedical settings,

including the monitoring of placental and pregnancy complications for maternal-fetal health

(33–36), to follow bone marrow reconstitution after stem cell transplantation (37), and to

characterize host immune responses to tuberculosis infection (38). However, integrative

analyses that capture host, viral, and microbial nucleic acids simultaneously remain

uncommon, and cfRNA sequencing approaches have been rarely applied to monitor retroviral

infections such as HIV. Here, we conducted a pilot study to investigate whether combined

cfDNA and cfRNA sequencing of plasma could reveal molecular features associated with the

development of bNAbs in a longitudinal cohort of PLWH.

Results

Study design, sequencing overview and viral analysis

We analyzed 42 plasma samples from 14 PLWH from the CAPRISA studies, of whom 7

developed bNAbs and 7 did not. For each participant, plasma was collected at approximately

6 months, 1 year, and 3 years post-HIV acquisition and prior to ART (Figure 1A). At the 6-

month time point, both groups had comparable CD4 counts and viral loads (Figure S1). The

extent of bNAb development was assessed by measuring plasma neutralization breadth against

a panel of 18 Env-pseudotyped HIV viruses, summarized as percentage breadth, a continuous

measure of bNAb development (Figure 1B, Table S1, Methods).

We performed combined cfDNA and cfRNA sequencing in each of the 42 samples (Methods).

After quality control and filtering, we obtained enough reads to sample the host transcriptome

(80% of total reads for cfRNA) and microbiome (0.2% and 20% reads of cfDNA and cfRNA

respectively) (Figure S2). Sequencing depth before and after sample QC processing (cleaning)

is similar for both cfDNA and cfRNA, with comparable metrics between bNAb producers and

non-producers (Figure S3). Similarly, cell-type deconvolution of the host transcriptome

showed comparable contributions of the most abundant cell types (e.g. red blood cells, immune

cells, platelets) across both groups (Figure S4). We detected HIV reads in the cfRNA assay for

all samples, with coverage sufficient to genotype the majority strain in 38 out of 42 samples

(Figure S5A,B). For high coverage samples, the nucleotide identity of HIV clustered within

individuals based on previous single-genome amplified envelope sequences (Figure 2A) except

for one mislabeled sample that was excluded from further analyses (Methods).

We validated the accuracy of cfRNA-derived consensus sequences by comparing them to

previously published envelope sequences from the same CAPRISA participants (Table S2),

observing consistent clustering and strain concordance across time points and sequencing

methods (Figure 2B, S5C). By comparing differences in consensus sequences within

individuals (Figure S5D), we found that most mutations occur in the *env* gene, which encodes

the surface protein responsible for host cell binding. The Env protein is the sole target of bNAbs

and is known to be the most rapidly evolving part of the HIV genome (39–41). Using cfRNA

we estimated that the dominant viral strain in each individual accumulates approximately 10

mutations per year (Figure S5E), in line with other estimates from virion or PBMC-derived

studies (42,43). Altogether, these results show that cfRNA is a quantitative tool to perform

genotype analysis of viral pathogens and to track strain-specific mutational dynamics over

time.

Analysis of read lengths and mapping start and end positions indicates that HIV-derived reads

in plasma cfRNA (cfHIV) predominantly consist of short RNA fragments (200-500 bases) and

do not show systematic positional biases along the HIV genome (Figure S6). This is consistent

with cfHIV being largely derived from fragmented cell-free viral RNA rather than libraries

dominated by intact full-length viral genomes. In addition, to assess the quality of HIV genome

recovery from cfRNA, we performed a subsampling analysis showing that approximately

50,000 cfHIV reads are sufficient to recover >90% of the genome at ≥50X depth (Figure S7).

In this cohort, this level of coverage was achieved for most samples (83%).

Host transcriptome correlates of cfHIV and clinical parameters

To characterize host responses to HIV infection, we measured the correlation between the host

transcriptome and the total abundance of cfHIV, as well as with clinical parameters such as

viral load and CD4 counts. Spearman correlation of cfHIV, viral load and CD4 counts to the

human transcriptome revealed several genes associated with these measurements (Figure 3A,

Figure S8A-B). Analysis of genes significantly correlated to cfHIV revealed 19 genes typically

associated with host immune responses to viral infection (FDR<0.05), whereas only 1 for each

of viral load and CD4 count, respectively (Figure 3B). CD4 count levels showed an association

with *IL4*, a cytokine inducing differentiation of naive T helper cells. Although the relation

between cytokine profile and HIV progression is controversial, increased secretion of IL4 in

relation to IFN-gamma is a known hallmark of switching from Th1 to Th2 lymphokine

secretion in PLWH (44). Our observation of IL4 being increased in low CD4 samples is

compatible with an increase of IL-4 levels upon disease progression and T helper cell decrease.

Top correlates of cfHIV include: *CCL5*, also known as RANTES, an HIV suppression factor

released by $CD8^+$ T cells, *TFRC* (CD71), a marker of T-cell activation, and *CCL2*, a chemokine

that attracts immune cells to sites of infection (Figure 3C, Figure S8C). These genes do not

show a differential behaviour between bNAb producers and controls, indicating that they

represent shared responses to HIV infection. This shows that cfRNA sequencing captures

shared host transcriptional responses associated with peripheral viral burden during HIV

infection.

Development of bNAbs is associated with a transcriptional phenotype of immune activation

We next compared the transcriptome of bNAb producers and non-producers. We performed

differential gene abundance analysis finding 256 genes elevated in bNAbs producers

(FDR<0.05, fold-change>1, Figure 4A, Methods). This variation could not be attributed to

environmental factors such as site of collection, that showed little to no correlation to these

transcriptomic signatures. Remarkably, 52% (n=131) of elevated genes in bNAb producers are

immune-related (primarily associated with adaptive immunity), compared with only 14%

(n=39) of the 281 decreased genes. Among the elevated genes, 25 of them are associated with

production of RANTES (CCL5), a chemokine involved in homing and migration of effector

and memory T cells. Although CCL5 itself is not enriched in bNAb producers in our dataset,

elevated RANTES, along with other inflammatory markers, have previously been linked to the

development of bNAbs in spontaneous controllers of HIV (45).

A pathway analysis confirmed these results and further revealed an enrichment of molecular

functions related to an immunologic response in bNAb producers (Figure S9A). Terms such as

receptor activity, IgG and peptide antigen binding, cytokine signaling and NF-kB signaling

were enriched in bNAb positive samples. Analyses of the genes decreased in the bNAb

producers showed lower enrichment scores and did not point to the existence of other host

immunoregulatory effects (Figure S9B). Overall, host cfRNA sequencing provides evidence

for an increased immune activation in participants who developed bNAbs, despite having

similar CD4 counts and viral loads to non-bNAb producers.

To investigate this immune activation in more detail, we performed pseudotime analysis on the

two groups, using plasma neutralization breadth to define the temporal progression (Figure 4B,

Methods). Pseudotime is a trajectory-inference approach that orders samples by relative

biological state rather than chronological time, allowing comparison across individuals at

different stages of bNAb development. We found that the immunologic gene signature is

elevated only early in breadth development in bNAb producers (when only a limited number

of viral strains are neutralized) and later converges to levels comparable to non-bNAb

producers as neutralization breadth increases. This aligns with previous studies showing that

factors such as CD4 levels inversely correlate with bNAb development only early in infection

(4). However, in our study, both groups had similar CD4 counts at the first time point. Focusing

on the genes significantly correlated with breadth in both groups (Figure S9C-D, FDR<0.05)

revealed almost the entirety of the MHC class I antigen presentation pathway was

anticorrelated with the development of breadth (i.e. pseudotime) and sometimes differentially

altered or correlated with the level of cfHIV measured (Figure 4C). Hence, MHC class I antigen

genes are highly elevated early in infection in participants who go on to develop bNAbs, and

their abundance decrease as breadth develops, becoming comparable to that of participants who

do not develop bNAbs. MHC class I antigen presentation is present in almost all cells of the

body, presents antigens to be killed by cytotoxic T cells (CD8) but is not generally associated

with antigen presenting cells or the production of antibodies by B cells (4). Taken together,

these results indicate that early immune activation associated with MHC class I antigen

presentation, independent of viral load or CD4 count, is a relevant factor in the development

of bNAbs.

Microbiome associated with bNAbs and the development of breadth

Development of bNAbs has been hypothesized to be triggered by antigen cross-reactivity

between HIV and bacterial components present in the gut microbiome (24,25,46,47). To test

whether a differentiated microbiota could be observed in participants who developed bNAbs,

we mapped all non-human reads (both cfDNA and cfRNA) to a database of bacterial and viral

genomes. After discarding reads that mapped to potential reagent contaminants (Figure S10,

Methods) (48), we identified 249 taxa in cfDNA and 4366 taxa in cfRNA that could be

confidently attributed to the participants microbiome. We combined data from all participants

to assemble a phylogenetic tree of the human microbiome during HIV infection as measured

from circulating nucleic acids (Figure S11). This accounted for ~70% of non-human reads in

both cfDNA and cfRNA, and spanned several areas of the tree of life including archaea,

bacteria and both DNA and RNA viruses. Among other features, we identified the presence of

several DNA and RNA viruses such as Anelloviruses, Flaviviruses, Herpesviruses indicative

of concomitant viral infections for several participants. In cfRNA, we found that ribosomal

reads encompass 37% of the nonhost microbiome reads, and we also identified a moderate

fraction of reads (~20%) that could not be attributed to specific species and mapped to

uncultured bacteria.

We then compared the microbiome of both groups (bNAb producers/controls) finding 19

genera that were differentially abundant between both groups (FDR<0.05, $|\log_2 FC|>1$, Figure

5A). Among genera enriched in bNAb producers, we identified several taxa that include

opportunistic pathogens and are commonly associated with mucosal surfaces and the

gastrointestinal tract, including Enterobacter, Serratia and Leclercia (Figure S12). By

performing a temporal analysis, we found that the detection of these pathogens does not differ

across timepoints and does not precede development of breadth (Figure S13). Overall, and

despite the relatively small sample size of this study, our data shows the existence of differences

in the microbiota compositions in participants who develop bNAbs from those who do not.

Additionally, metagenomic analysis of cfRNA revealed the presence of GB virus C (GBV-C,

previously known as Hepatitis G) in several participants (Fig. 5B). GBV-C is a flavivirus that

infects lymphocytes, has no known pathology, and has been associated with improved survival

in people with HIV infection (49,50). Given the phylogenetic relationship between GBV-C and

hepatitis C virus (HCV), we also examined reads mapping to HCV. However, we did not find

evidence of HCV signal, whereas reads mapping to GBV-C showed broad coverage across the

viral genome (Figure S13). In our study, all bNAb producers had non-zero levels of GBV-C (p

< 0.05, one-sided Fisher exact test) and showed higher odds of elevated GBV-C levels (Figure

5B). Differential gene abundance analysis on samples with high levels of GBV-C (defined as

>1/2000 nonhost reads) revealed 42 host genes that are also elevated in these participants.

Among top candidates we identified several genes of the MHC class II antigen presentation

complex (*HLA-DQA1*, *HLA-DQB1*, *HLA-DRB1*), an HIV suppressive factor involved in

inflammatory signaling (*CCL3*), several proinflammatory cytokines (*IL6*, *EBI3*), and

immunoglobulin receptors (*KIR3DL2*). Notably, increased expression of inhibitory KIRs,

including KIR3DL2, has been associated with bNAb development, suggesting a potential link

between natural killer cell maturation and the regulation of antibody production (51). These

findings suggest that high levels of GBV-C might be associated with an increased activity of

antigen presenting cells.

Discussion

In this work, we demonstrate simultaneous monitoring of viral infection and host response

through sequencing of circulating nucleic acids (DNA and RNA) in plasma samples of PLWH

prior to ART. We show that cfRNA can be used to characterize circulating HIV viral strains

and, at the same time, to profile the host transcriptome, allowing characterization of immune

activity during HIV infection. Analysis of fragment length and genome coverage suggests that

HIV-derived cfRNA likely reflects fragmented viral RNA circulating in plasma rather than

being dominated by intact full-length viral genomes. Subsampling analyses further shows that

near-complete genome coverages (>90%) at moderate depths (≥50X) can be recovered from

cfRNA, supporting the feasibility of viral genome analyses.

As cfRNA contains transcripts released from diverse tissues and cell types across the body

(30), it offers a systemic view of immune processes that may be otherwise difficult to capture.

We find that overall cfHIV levels—defined as HIV-derived fragments in cfRNA—correlate

with abundance of host genes involved in immune responses to viral infections, including

CCL5, TFRC, and CCL2, consistent with responses to peripheral viral burden. For instance,

CCL5 encodes RANTES, a chemokine with HIV-suppressive activity that can block CCR5-

mediated viral entry, while TFRC (CD71) and CCL2 (MCP-1) are involved in pathways

commonly perturbed during infection (52–54). However, when comparing participants who

develop bNAbs with those who did not, we observed a distinct and orthogonal transcriptomic

signature suggestive of an enhanced immune activation early in infection that was specific to

bNAb producers. This response is independent of viral load and CD4 counts, and is enriched

in MHC class I antigen presentation genes, RANTES-associated pathways (though not CCL5

itself), and other immune activation pathways.

Previous studies have highlighted the central role of lymph nodes in shaping humoral responses

during HIV infection, yet their direct study remains challenging (23,55,56). Cell-free RNA

may provide a non-invasive window into immune activity occurring in tissue compartments,

including lymphoid tissues where Tfh-B cell interactions drive affinity maturation. Hence, it is

plausible that the immune-related cfRNA transcripts we report here reflects immune activity

within lymph nodes, consistent with previous evidence that cfRNA can capture tissue-resident

hematopoietic activity such as bone marrow progenitor transcripts (37). However, we

emphasize that cfRNA represents an integrated signal from multiple sources, and we cannot

exclude the possibility that increased cfRNA levels are due to contributions of circulating

immune cells or other tissues.

Analysis of non-human reads further allowed us to assess the microbiome and virome,

revealing a reduced set of taxa whose presence or absence is associated with the development

of bNAbs. Microbial sequences detected in plasma cfRNA/cfDNA most likely reflect leakage

of microbial products into circulation due to barrier dysfunction and dysbiosis, a process well

described in chronic HIV infection (57–60). While gut microbiota has been speculated to

increase antigen cross-reactivity, our data cannot establish a direct role in bNAb development.

Surprisingly, we found that participants who developed bNAbs consistently had higher levels

of GBV-C. GBV-C is a non-pathogenic lymphotropic flavivirus that has been associated with

improved outcomes in PLWH (49), with some studies suggesting this may reflect an indirect

rather than a causal association (50). Mechanistically, co-infection with GBV-C has been

reported to inhibit HIV replication and modulate cytokine release, co-receptor expression, and

T-cell homeostasis (50,61,62). In our study, GBV-C abundance correlated with increased

abundance of proinflammatory cytokines and adaptative immune genes, raising the possibility

that co-infection influences the host immune milieu in a way that could facilitate bNAb

development. While causality cannot be inferred from our dataset, understanding such

modulatory viral co-infections early after HIV acquisition could inform strategies to mimic

beneficial immune environments in the context of HIV vaccine design.

Our study represents a small pilot investigation (42 samples, 14 PLWH) in a well-characterized

cohort of young women from South Africa (63), followed from prior to HIV acquisition into

acute infection and before initiation of ART. Biobanked samples from such cohorts are an

invaluable resource for studies of factors associated with the development of breadth, as earlier

ARV treatment regimens are now standard of care. Although this provides a relatively

homogeneous and well-controlled population for an initial discovery study, the small sample

size limits generalizability and statistical power. For example, associations between viral load

or CD4 counts and bNAb induction reported in other studies may not be detectable here due to

the limit size of this pilot cohort. Expanded studies within the CAPRISA cohort or in other

settings, and with finer temporal resolution, will be important to validate the findings reported

here and their relationship to bNAb development. Despite these limitations, this work

illustrates the potential of cfRNA/cfDNA sequencing as a non-invasive approach to interrogate

host, viral, and microbial processes simultaneously from a single plasma sample. This approach

could be of use to support biomarker discovery in vaccine studies and help identify biological

contexts that favor the induction of bNAbs.

Materials and Methods

Ethics Statement

336 The CAPRISA 002 Acute Infection study was reviewed and approved by the research ethics

337 committees of the University of KwaZulu-Natal (E013/04), the University of Cape Town

338 (025/2004), and the University of the Witwatersrand (MM040202). All participants provided

339 written informed consent for screening, enrolment and specimen storage.

Participants

342 The study utilized biobanked plasma specimen of PLWH with known neutralization status

343 enrolled in the CAPRISA cohorts. The CAPRISA 002 Acute Infection cohort was established

344 in 2004 and has been following women from the early/acute stage of infection (63). We

345 included participants from two CAPRISA clinical sites, Ethekwini in Durban (site 1) and

346 Vulindlela (site 2), and accounted for potential site-specific effects in all downstream analyses.

Neutralization measurements

Neutralization breadth was measured using a standardized pseudovirus assay that included 18

Env-pseudotyped HIV viruses (Table S1) (64,65). Briefly, HIV pseudoviruses were produced

by co-transfecting HEK293T cells with each HIV-1 Env plasmid and a pSG3 backbone. Serial

dilutions of heat-inactivated plasma were preincubated with each pseudovirus, followed by the

addition of TZM-bl cells. After 48h, infection was quantified by luciferase read-out, and titers

reported as the reciprocal dilution where 50% of the virus is neutralized (IC_{50}). Breadth was

defined as the percentage of pseudoviruses neutralized at an $IC_{50} \geq 1:45$.

Participants were classified as bNAb producers if they achieved $\geq 20\%$ breadth within year of

infection or $\geq 40\%$ breadth against the 18-pseudovirus panel at any three years post-infection.

Isolation of cfDNA and cfRNA and library preparation

Blood samples were collected in sodium citrate tubes and plasma extracted according to the

standard clinical centrifugation protocol of the HIV/AIDS Network Coordination, and stored

at -80 C. Each plasma sample was thawed once and split into two aliquots for cfDNA/cfRNA

extraction (0.4-1 mL per aliquot). To minimize confounding effects from hemolysis and

platelet-derived RNA, plasma samples were visually inspected prior to processing for signs of

hemolysis (e.g. pink discoloration), which was not visible for any sample. Immediately before

nucleic acid extraction, plasma samples were subjected to a final centrifugation step (1 min at

16,000 rpm) to remove any residual debris. cfDNA was extracted using the Qiagen Circulating

Nucleic Acid kit. cfRNA was extracted using the Plasma/Serum Circulating RNA and

Exosomal Purification kit (Norgen, cat 42800), followed by DNA digestion using Baseline-

ZERO DNase (Epicentre) and then cleaned up using the RNA Clean and Concentrator-5 kit

(Zymo) (32,66). cfDNA/cfRNA libraries were prepared using the automated Ovation SP

Ultralow Library Systems (Mondrian ST) and the SMARTer Stranded Total RNA Pico Input

Mammalian kit (Clontech) respectively (32,66). A negative control (1 mL PBS, Gibco) was

processed in parallel for every batch to control for potential contaminants (67).

Sequencing methods

Paired end Illumina sequencing (2×75 bp) was performed on the Nextseq 500 for cfDNA and

on the Novaseq 6000 for cfRNA. This resulted in a median of 69 million (cfDNA) and 109

million (cfRNA) reads per sample (Figure S3).

Sequencing Analysis

Sequencing reads from cfRNA and cfDNA were processed using a shared initial pipeline. First,

low quality bases and adapter sequences were trimmed by Trimmomatic v0.38 (68), and

overlapping read pairs were merged using FLASH v1.2.11 (69). Merged reads were then

aligned to the UniVec database using bowtie2 to remove reads mapping to common vectors,

linkers and control sequences (70). These processing QC steps removed only a fraction of reads

in cfRNA and cfDNA libraries from most plasma samples (Figure S2-S3). Higher cleaning

rates were observed in cfDNA negative controls (~50%) and in three cfRNA plasma samples

(~25%), that may reflect lower sample input or quality. Overall, no differences in QC

processing were observed between bNAb producers and non-producers.

The remaining reads were first aligned to the human genome (GRCh38, Ensembl) using

bowtie2, retaining unmapped reads, which were subsequently passed through a second

homology filtering by BLASTN against the NCBI nt database restricted to taxid 9606, to obtain

non-human reads for microbial and viral analysis. Taxonomic classification of non-human

reads was performed by BLASTN (NCBI BLAST+ v2.8) against the NCBI nt database, using

curated taxid lists limited to viral, bacterial and fungal references (66,67). BLAST hits were

filtered by requiring ≥80% nucleotide identity and ≥80% per-read query coverage, also de-

duplicating counts to the best-supported accession to avoid counting the same read twice across

hits. Subsequently, per-accession counts and coverage profiles were generated and aggregated

at taxonomic ranks for downstream analysis.

For cfRNA, reads were additionally aligned to the human genome using STAR v.25.4a (71)

and duplicate-filtered using Picard v2.18.2,and gene abundance was quantified using HTSeq

(72). Downstream analysis of gene abundance data is detailed in the following section.

For HIV-specific analysis, non-human cfRNA reads were re-aligned to the HIV-1 group M

consensus reference (CON-C.fa) using bowtie2. Consensus sequences were generated with

samtools mpileup and bcftools consensus (73), followed by multiple sequence alignment using

MAFFT v7 (74). A maximum-likelihood phylogenetic tree was constructed using IQ-TREE

(75), and visualized with Bio.Phylo (76), with metadata-based label coloring and shaded

backgrounds highlighting predefined sample clusters. Previously published HIV reference

sequences from the same CAPRISA participants were obtained from the Los Alamos National

Laboratory (LANL) HIV Sequence Database (Table S2). Sample identity was primarily

validated by assessing HIV sequence consistency across all the three time-points per participant

(Fig. 2 and S3). As an orthogonal check, we performed HLA-B typing on cfRNA using

seq2HLA (v2.3, 4-digit resolution) (77), which confirmed exclusion of one sample (CAP200,

time point 1), where a mismatch was detected in both the HIV sequence analysis and HLA-B

typing.

To control for potential environmental contamination in both cfDNA and cfRNA data, six

negative controls were used to create a database of background environmental taxa that were

blacklisted in downstream analysis. This filtering preserved 72% of microbial reads in cfRNA

and 17% in cfDNA. In addition, non-template control barcodes (n=8) were spiked into the

cfRNA libraries to estimate sample cross-contamination, which was found to be <0.5% and

compatible with index-hopping during sequencing. This threshold was used to define the

detection limit for each individual microbial taxon.

Differential Gene Abundance Analysis

Differential Gene Abundance Analysis was performed using edgeR with a generalized linear

model and quasi-likelihood F-tests. Site of sample collection was included as a covariate in the

model to account for potential confounding effects. All p-values were corrected for multiple

hypothesis testing using the Benjamini-Hochberg method, and results are reported as false

discovery rate (FDR). For downstream analysis, we selected genes with FDR < 0.05 and

absolute fold change > 1. Associations between gene abundance and clinical metadata (e.g.

viral load, CD4+ counts) were measured using nonparametric Spearman correlation unless

otherwise stated (SciPy package, Python). Differential abundance of microbial taxa was also

assessed in edgeR using generalized linear models, with the same thresholds and covariate

adjustments. Overall, no significant site-associated differences were observed in either the

transcriptomic or microbial enrichment analyses.

Pseudotime analysis

Pseudotime analysis was used as an alternative approach to investigate factors associated with

the development of neutralization breadth, given the limited number of longitudinal samples

and heterogeneity in bNAb development across individuals. Samples were ordered according

to the number of pseudoviruses neutralized by each plasma sample, which was used as a proxy

for progression along the bNAb development continuum. This trajectory-inference approach

allows us to account for differences in the pace of bNAb development across individuals.

Spearman's rank correlation analyses were then performed against this pseudotime variable to

evaluate changes both between bNAb producers and non-producers and along the continuous

trajectory of breadth progression.

GBV-C analysis

GBV-C reads were normalized to the total number of non-human reads per sample and scaled

by a factor of 1,000 (GBV-C reads per 1,000 non-human reads), analogous to the normalization

used for cfHIV. Differences in GBV-C abundance between bNAb producers and non-

producers were assessed using a logistic mixed-effects model (statsmodels package, Python),

where we model GBV-C abundance as a binary outcome and included participant as a random

effect to account for repeated sampling across time points.

Acknowledgments

This work was supported by the funding from the Chan Zuckerberg Biohub to S.R.Q, the Bill

& Melinda Gates Foundation (OPP1113682) and the Global Health Vaccine Accelerator

Platform (GH-VAP-SI-ID-14) to S.R.Q and P.L.M. We thank the Centre for the AIDS

Programme of Research In South Africa (CAPRISA) for sample procurement, Dr Mashudu

Madzivhandila for neutralization data, and MD Gerlinde Obermoser (Stanford Center for

Human Systems Immunology) for project coordination. P.L.M. is supported by South African

Medical Research Council (MRC) SHIP program and the South African Research Chairs

Initiative of the Department of Science and Technology and the NRF (Grant No 98341). J.C.-

S. is supported by the Knut and Alice Wallenberg Foundation (Wallenberg Molecular Medicine

Fellow), the Swedish Research Council (2021-05109) and the Erling Perssons Stiftelse

(Swedish Foundations' Starting Grant). M.N.M. was supported by the Stanford Bio-X Bowes

Fellowship. Finally, we are indebted to the study participants for their generous support of

scientific research.

Data availability

The datasets generated and analyzed in the study are available in the NCBI Gene Expression

Omnibus (GEO) and Sequence Read Archive (SRA) with accession number GSE313190. The

code to reproduce the analyses in this study is publicly available at

https://github.com/CamunasLab/cfHIV

Bibliography

1. UNAIDS Report 2017. Ending AIDS. Progress towards the 90-90-90 targets. (Global Aids Update 2017.). Joint United Nations Programme on HIV/AIDS (UNAIDS); 2017.

2. Deeks SG, Overbaugh J, Phillips A, Buchbinder S. HIV infection. Nature reviews Disease primers. 2015;1:15035.

3. Moore PL, Gray ES, Choge IA, Ranchobe N, Mlisana K, Abdool Karim SS, et al. The C3-V4 Region Is a Major Target of Autologous Neutralizing Antibodies in Human Immunodeficiency Virus Type 1 Subtype C Infection. J Virol. 2008;82:1860–9.

4. Gray ES, Moore PL, Choge IA, Decker JM, Bibollet-Ruche F, Li H, et al. Neutralizing Antibody Responses in Acute Human Immunodeficiency Virus Type 1 Subtype C Infection. J Virol. 2007;81:6187–96.

5. Li B, Decker JM, Johnson RW, Bibollet-Ruche F, Wei X, Mulenga J, et al. Evidence for Potent Autologous Neutralizing Antibody Titers and Compact Envelopes in Early Infection with Subtype C Human Immunodeficiency Virus Type 1. J Virol. 2006;80:5211–8.

6. Richman DD, Wrin T, Little SJ, Petropoulos CJ. Rapid evolution of the neutralizing antibody response to HIV type 1 infection. Proc Natl Acad Sci USA. 2003;100:4144–9.

7. Landais E, Huang X, Havenar-Daughton C, Murrell B, Price MA, Wickramasinghe L, et al. Broadly Neutralizing Antibody Responses in a Large Longitudinal Sub-Saharan HIV Primary Infection Cohort. PLoS Pathog. 2016;12:e1005369.

8. Corey L, Gilbert PB, Juraska M, Montefiori DC, Morris L, Karuna ST, et al. Two Randomized Trials of Neutralizing Antibodies to Prevent HIV-1 Acquisition. N Engl J Med. 2021;384:1003–14.

9. Caniels TG, Prabhakaran M, Ozorowski G, MacPhee KJ, Wu W, Van Der Straten K, et al. Precise targeting of HIV broadly neutralizing antibody precursors in humans. Science. 2025;eadv5572.

10. Willis JR, Prabhakaran M, Muthui M, Naidoo A, Sincomb T, Wu W, et al. Vaccination with mRNA-encoded nanoparticles drives early maturation of HIV bnAb precursors in humans. Science. 2025;eadr8382.

11. Leggat DJ, Cohen KW, Willis JR, Fulp WJ, deCamp AC, Kalyuzhniy O, et al. Vaccination induces HIV broadly neutralizing antibody precursors in humans. Science. 2022;378:eadd6502.

12. The Swiss HIV Cohort Study, Rusert P, Kouyos RD, Kadelka C, Ebner H, Schanz M, et al. Determinants of HIV-1 broadly neutralizing antibody induction. Nat Med. 2016;22:1260–7.

13. Moody MA, Pedroza-Pacheco I, Vandergrift NA, Chui C, Lloyd KE, Parks R, et al. Immune perturbations in HIV-1-infected individuals who make broadly neutralizing antibodies. Science Immunology. 2016;1:aag0851.

14. Haynes BF, Shaw GM, Korber B, Kelsoe G, Sodroski J, Hahn BH, et al. HIV-Host Interactions: Implications for Vaccine Design. Cell Host & Microbe. 2016;19:292–303.

15. Roskin KM, Jackson KJL, Lee J-Y, Hoh RA, Joshi SA, Hwang K-K, et al. Aberrant B cell repertoire selection associated with HIV neutralizing antibody breadth. Nature Immunology. 2020;21:199–209.

16. Landais E, Moore PL. Development of broadly neutralizing antibodies in HIV-1 infected elite neutralizers. Retrovirology. 2018;15:61.

17. Euler Z, van Gils MJ, Bunnik EM, Phung P, Schweighardt B, Wrin T, et al. Cross-reactive neutralizing humoral immunity does not protect from HIV type 1 disease progression. J Infect Dis. 2010;201:1045–53.

18. Gray ES, Madiga MC, Hermanus T, Moore PL, Wibmer CK, Tumba NL, et al. The Neutralization Breadth of HIV-1 Develops Incrementally over Four Years and Is Associated with CD4+ T Cell Decline and High Viral Load during Acute Infection. J Virol. 2011;85:4828–40.

19. Cottrell CA, Hu X, Lee JH, Skog P, Luo S, Flynn CT, et al. Heterologous prime-boost vaccination drives early maturation of HIV broadly neutralizing antibody precursors in humanized mice. Sci Transl Med. 2024;16:eadn0223.

20. Cohen K, Altfeld M, Alter G, Stamatatos L. Early preservation of CXCR5+ PD-1+ helper T cells and B cell activation predict the breadth of neutralizing antibody responses in chronic HIV-1 infection. J Virol. 2014;88:13310–21.

21. Locci M, Havenar-Daughton C, Landais E, Wu J, Kroenke MA, Arlehamn CL, et al. Human Circulating PD-1+CXCR3−CXCR5+ Memory Tfh Cells Are Highly Functional and Correlate with Broadly Neutralizing HIV Antibody Responses. Immunity. 2013;39:758–69.

22. Perreau M, Savoye A-L, De Crignis E, Corpataux J-M, Cubas R, Haddad EK, et al. Follicular helper T cells serve as the major CD4 T cell compartment for HIV-1 infection, replication, and production. J Exp Med. 2013;210:143–56.

23. Crotty S. T Follicular Helper Cell Biology: A Decade of Discovery and Diseases. Immunity. 2019;50:1132–48.

24. Liao H-X, Chen X, Munshaw S, Zhang R, Marshall DJ, Vandergrift N, et al. Initial antibodies binding to HIV-1 gp41 in acutely infected subjects are polyreactive and highly mutated. Journal of Experimental Medicine. 2011;208:2237–49.

25. Trama AM, Moody MA, Alam SM, Jaeger FH, Lockwood B, Parks R, et al. HIV-1 Envelope gp41 Antibodies Can Originate from Terminal Ileum B Cells that Share Cross-Reactivity with Commensal Bacteria. Cell Host & Microbe. 2014;16:215–26.

26.   Williams WB, Liao H-X, Moody MA, Kepler TB, Alam SM, Gao F, et al. Diversion of HIV-1
vaccine–induced immunity by gp41-microbiota cross-reactive antibodies. Science.
2015;349:aab1253.

27.   Loy C, Ahmann L, De Vlaminck I, Gu W. Liquid Biopsy Based on Cell-Free DNA and RNA.
Annual Review of Biomedical Engineering. 2024;26:169–95.

28.   Moufarrej MN, Bianchi DW, Shaw GM, Stevenson DK, Quake SR. Noninvasive Prenatal
Testing Using Circulating DNA and RNA: Advances, Challenges, and Possibilities. Annu Rev
Biomed Data Sci. 2023;6:397–418.

29.   Koh W, Pan W, Gawad C, Fan HC, Kerchner GA, Wyss-Coray T, et al. Noninvasive in vivo
monitoring of tissue-specific global gene expression in humans. Proc Natl Acad Sci USA.
2014;111:7361–6.

30.   Vorperian SK, Moufarrej MN, Tabula Sapiens Consortium, Overall Project Direction and
Coordination, Jones RC, Karkanias J, et al. Cell types of origin of the cell-free transcriptome.
Nat Biotechnol. 2022;40:855–61.

31.   Loy CJ, Servellita V, Sotomayor-Gonzalez A, Bliss A, Lenz JS, Belcher E, et al. Plasma cell-
free RNA signatures of inflammatory syndromes in children. Proc Natl Acad Sci USA.
2024;121:e2403897121.

32.   Pan W, Ngo TTM, Camunas-Soler J, Song C-X, Kowarsky M, Blumenfeld YJ, et al.
Simultaneously Monitoring Immune Response and Microbial Infections during Pregnancy
through Plasma cfRNA Sequencing. Clinical Chemistry. 2017;63:1695–704.

33.   Camunas-Soler J, Gee EPS, Reddy M, Mi JD, Thao M, Brundage T, et al. Predictive RNA
profiles for early and very early spontaneous preterm birth. American Journal of Obstetrics and
Gynecology. 2022;227:72.e1-72.e16.

34.   Ngo TTM, Moufarrej MN, Rasmussen M-LH, Camunas-Soler J, Pan W, Okamoto J, et al.
Noninvasive blood tests for fetal development predict gestational age and preterm delivery.
Science. 2018;360:1133–6.

35.   Rasmussen M, Reddy M, Nolan R, Camunas-Soler J, Khodursky A, Scheller NM, et al. RNA
profiles reveal signatures of future health and disease in pregnancy. Nature. 2022;601:422–7.

36.   Moufarrej MN, Vorperian SK, Wong RJ, Campos AA, Quaintance CC, Sit RV, et al. Early
prediction of preeclampsia in pregnancy with cell-free RNA. Nature. 2022;602:689–94.

37.   Ibarra A, Zhuang J, Zhao Y, Salathia NS, Huang V, Acosta AD, et al. Non-invasive
characterization of human bone marrow stimulation and reconstitution by cell-free messenger
RNA sequencing. Nat Commun. 2020;11:400.

38.   Chang A, Loy CJ, Eweis-LaBolle D, Lenz JS, Steadman A, Andgrama A, et al. Circulating cell-
free RNA in blood as a host response biomarker for detection of tuberculosis. Nat Commun.
2024;15:4949.

39.   Burton DR, Stanfield RL, Wilson IA. Antibody vs. HIV in a clash of evolutionary titans.
Proceedings of the National Academy of Sciences. 2005;102:14943–8.

40.   Haddox HK, Dingens AS, Bloom JD. Experimental Estimation of the Effects of All Amino-
Acid Mutations to HIV's Envelope Protein on Viral Replication in Cell Culture. Swanstrom R,
editor. PLoS Pathog. 2016;12:e1006114.

41.   Burton DR, Mascola JR. Antibody responses to envelope glycoproteins in HIV-1 infection. Nat
594   Immunol. 2015;16:571–6.

595   42.   Cuevas JM, Geller R, Garijo R, López-Aldeguer J, Sanjuán R. Extremely High Mutation Rate of
596   HIV-1 In Vivo. Rowland-Jones SL, editor. PLoS Biol. 2015;13:e1002251.

597   43.   Zanini F, Puller V, Brodin J, Albert J, Neher RA. In vivo mutation rates and the landscape of
598   fitness costs of HIV-1. Virus Evolution. 2017;3.

599   44.   Meyaard L, Hovenkamp E, Keet IP, Hooibrink B, de Jong IH, Otto SA, et al. Single cell
600   analysis of IL-4 and IFN-gamma production by T cells from HIV-infected individuals:
601   decreased IFN-gamma in the presence of preserved IL-4 production. Journal of Immunology.
602   1996;157:2712–8.

603   45.   Dugast A-S, Arnold K, Lofano G, Moore S, Hoffner M, Simek M, et al. Virus-driven
604   Inflammation Is Associated With the Development of bNAbs in Spontaneous Controllers of
605   HIV. Clinical Infectious Diseases. 2017;64:1098–104.

606   46.   Mascola JR, Haynes BF. HIV-1 neutralizing antibodies: understanding nature's pathways.
607   Immunological Reviews. 2013;254:225–44.

608   47.   Moir S, Fauci AS. B-cell responses to HIV infection. Immunological Reviews. 2017;275:33–48.

609   48.   Salter SJ, Cox MJ, Turek EM, Calus ST, Cookson WO, Moffatt MF, et al. Reagent and
610   laboratory contamination can critically impact sequence-based microbiome analyses. BMC
611   Biology. 2014;12:87.

612   49.   Xiang J, Wünschmann S, Diekema DJ, Klinzman D, Patrick KD, George SL, et al. Effect of
613   Coinfection with GB Virus C on Survival among Patients with HIV Infection. N Engl J Med.
614   2001;345:707–14.

615   50.   Schwarze-Zander C, Blackard JT, Rockstroh JK. Role of GB virus C in modulating HIV
616   disease. Expert Review of Anti-Infective Therapy. 2012;10:563–72.

617   51.   Bradley T, Peppa D, Pedroza-Pacheco I, Li D, Cain DW, Henao R, et al. RAB11FIP5
618   Expression and Altered Natural Killer Cell Function Are Associated with Induction of HIV
619   Broadly Neutralizing Antibody Responses. Cell. 2018;175:387-399.e17.

620   52.   Mengozzi M, De Filippi C, Transidico P, Biswas P, Cota M, Ghezzi S, et al. Human
621   Immunodeficiency Virus Replication Induces Monocyte Chemotactic Protein-1 in Human
622   Macrophages and U937 Promonocytic Cells. Blood. 1999;93:1851–7.

623   53.   Packard TA, Schwarzer R, Herzig E, Rao D, Luo X, Egedal JH, et al. CCL2: a Chemokine
624   Potentially Promoting Early Seeding of the Latent HIV Reservoir. Moscona A, editor. mBio.
625   2022;13:e01891-22.

626   54.   Madrid R, Janvier K, Hitchin D, Day J, Coleman S, Noviello C, et al. Nef-induced Alteration of
627   the Early/Recycling Endosomal Compartment Correlates with Enhancement of HIV-1
628   Infectivity. Journal of Biological Chemistry. 2005;280:5032–44.

629   55.   Cirelli KM, Carnathan DG, Nogal B, Martin JT, Rodriguez OL, Upadhyay AA, et al. Slow
630   Delivery Immunization Enhances HIV Neutralizing Antibody and Germinal Center Responses
631   via Modulation of Immunodominance. Cell. 2019;177:1153-1171.e28.

632   56.   Lindqvist M, Van Lunzen J, Soghoian DZ, Kuhl BD, Ranasinghe S, Kranias G, et al. Expansion
633   of HIV-specific T follicular helper cells in chronic HIV infection. J Clin Invest. 2012;122:3271–
634   80.

635   57.   Alzahrani J, Hussain T, Simar D, Palchaudhuri R, Abdel-Mohsen M, Crowe SM, et al.
636   Inflammatory and immunometabolic consequences of gut dysfunction in HIV: Parallels with
637   IBD and implications for reservoir persistence and non-AIDS comorbidities. eBioMedicine.
638   2019;46:522–31.

639   58.   Das Adhikari U, Froehle LM, Pipkin AN, Baharlou H, Linder AH, Shah P, et al.
640   Immunometabolic defects of CD8+ T cells disrupt gut barrier integrity in people with HIV. Cell.
641   2025;S0092867425009791.

642   59.   Brenchley JM, Price DA, Schacker TW, Asher TE, Silvestri G, Rao S, et al. Microbial
643   translocation is a cause of systemic immune activation in chronic HIV infection. Nat Med.
644   2006;12:1365–71.

645   60.   Nazli A, Chan O, Dobson-Belaire WN, Ouellet M, Tremblay MJ, Gray-Owen SD, et al.
646   Exposure to HIV-1 Directly Impairs Mucosal Epithelial Barrier Integrity Allowing Microbial
647   Translocation. Hope TJ, editor. PLoS Pathog. 2010;6:e1000852.

648   61.   Stapleton JT. Human Pegivirus Type 1: A Common Human Virus That Is Beneficial in
649   Immune-Mediated Disease? Front Immunol. 2022;13:887760.

650   62.   Acchioni C, Sandini S, Acchioni M, Sgarbanti M. Co-Infections and Superinfections between
651   HIV-1 and Other Human Viruses at the Cellular Level. Pathogens. 2024;13:349.

652   63.   Van Loggerenberg F, Mlisana K, Williamson C, Auld SC, Morris L, Gray CM, et al.
653   Establishing a Cohort at High Risk of HIV Infection in South Africa: Challenges and
654   Experiences of the CAPRISA 002 Acute Infection Study. Ugarte-Gil CA, editor. PLoS ONE.
655   2008;3:e1954.

656   64.   Montefiori DC. Evaluating Neutralizing Antibodies Against HIV, SIV, and SHIV in Luciferase
657   Reporter Gene Assays. CP in Immunology . 2005;64: Chapter 12:12.11.1-12.11.17.

658   65.   Moyo-Gwete T, Scheepers C, Makhado Z, Kgagudi P, Mzindle NB, Ziki R, et al. Enhanced
659   neutralization potency of an identical HIV neutralizing antibody expressed as different isotypes
660   is achieved through genetically distinct mechanisms. Sci Rep. 2022;12:16473.

661   66.   De Vlaminck I, Khush KK, Strehl C, Kohli B, Luikart H, Neff NF, et al. Temporal response of
662   the human virome to immunosuppression and antiviral therapy. Cell. 2013;155:1178–87.

663   67.   Kowarsky M, Camunas-Soler J, Kertesz M, De Vlaminck I, Koh W, Pan W, et al. Numerous
664   uncharacterized and highly divergent microbes which colonize humans are revealed by
665   circulating cell-free DNA. Proc Natl Acad Sci USA. 2017;114:9623–8.

666   68.   Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina sequence data.
667   Bioinformatics. 2014;30:2114–20.

668   69.   Magoc T, Salzberg SL. FLASH: fast length adjustment of short reads to improve genome
669   assemblies. Bioinformatics. 2011;27:2957–63.

670   70.   Langmead B, Salzberg SL. Fast gapped-read alignment with Bowtie 2. Nat Methods.
671   2012;9:357–9.

71. Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, et al. STAR: ultrafast universal
RNA-seq aligner. Bioinformatics. 2013;29:15–21.

72. Putri GH, Anders S, Pyl PT, Pimanda JE, Zanini F. Analysing high-throughput sequencing data
in Python with HTSeq 2.0. Boeva V, editor. Bioinformatics. 2022;38:2943–5.

73. Danecek P, Bonfield JK, Liddle J, Marshall J, Ohan V, Pollard MO, et al. Twelve years of
SAMtools and BCFtools. GigaScience. 2021;10:giab008.

74. Katoh K, Standley DM. MAFFT Multiple Sequence Alignment Software Version 7:
Improvements in Performance and Usability. Molecular Biology and Evolution. 2013;30:772–
80.

75. Minh BQ, Schmidt HA, Chernomor O, Schrempf D, Woodhams MD, Von Haeseler A, et al.
Corrigendum to: IQ-TREE 2: New Models and Efficient Methods for Phylogenetic Inference in
the Genomic Era. Molecular Biology and Evolution. 2020;37:2461–2461.

76. Talevich E, Invergo BM, Cock PJ, Chapman BA. Bio.Phylo: A unified toolkit for processing,
analyzing and visualizing phylogenetic trees in Biopython. BMC Bioinformatics. 2012;13:209.

77. Boegel S, Löwer M, Schäfer M, Bukur T, De Graaf J, Boisguérin V, et al. HLA typing from
RNA-Seq sequence reads. Genome Med. 2012;4:102.

Figure 1: Study design and development of neutralization breadth. A) Longitudinal

trajectories of neutralization breadth in bNAb producers (red) and controls (blue). Time zero

corresponds to the estimated HIV infection point, with breadth defined as the ability of plasma

antibodies to neutralize a cross-clade panel of 18 env-pseudotyped viruses (Table S2). B)

Heatmap of serum neutralization titers (IC50) against the 18-virus panel, grouped by

individual, with time points sorted from first (top) to last (bottom). Colors represent log10-

transformed IC50 neutralization titers, with higher values (yellow) indicating stronger

neutralization potency and black indicating values below the limit of detection.

Figure 2: HIV consensus sequences from cfRNA. A) Genomic similarity matrix of consensus

701 HIV sequences obtained from cfRNA, showing clustering within individuals. B) Maximum-

702 likelihood phylogeny of cfRNA-derived HIV consensus sequences (colored labels) with

703 previously published envelope sequences (bold, CAPRISA). Shaded boxes highlight

704 individual-level clustering. With the exception of one sample, cfRNA consensus sequences

705 consistently cluster with the corresponding envelope sequences.

Figure 3: Correlation between cfHIV levels and host transcriptome. A) Number of genes

significantly correlated with cfHIV counts, viral load (VL) and CD4 counts (Spearman's rank

correlation, see Methods). B) Subset of genes from panel A previously described as involved

in host immune responses to viral infection. C) Example correlations between cfHIV counts

and three HIV-associated genes. Spearman correlation coefficients indicated. No partitioning

is observed between bNAb producers (red) and controls (blue).

Figure 4: Gene level differences in cfRNA between bNAb producers and controls. A)

Volcano plot of differentially abundant genes. Genes with a fold change > 2 and FDR < 0.05

are highlighted (red = increased, blue = decreased in bNAb producers). Labelled points are

genes involved in host immune responses to viral infection. B) Violin plots of the distribution

of elevated genes in bNAb producers, stratified by group and neutralization breadth (number

of viral strains neutralized), used here as a proxy for progression along bNAb development. C)

Genes in the MHC class I antigen presentation pathway. Most show negative correlations with

breadth, and some are also differentially abundant between bNAb producers and controls.

Correlations with breadth were assessed using Spearman's rank correlation (Spearman's ρ

indicated next to each gene, see Methods). * correlated with breadth at FDR <0.05, †

differentially abundant between bNAb producers and controls at FDR<0.05, ‡ correlated with

cfHIV at FDR <0.05.

Figure 5: Microbial differences and GBV-C enrichment in bNAb producers. A) Volcano

plot of differentially abundant microbial genera between bNAb producers and non-producers,

colored by phylum (FDR < 0.05, Benjamini-Hochberg). Positive log2 fold change indicates

enrichment in bNAb producers. B) Scatterplots showing cfGBV-C abundance in samples

partitioned by time and study group (left, bNAbs producers, right, controls). The dashed line

indicates the threshold used to define elevated cfGBV-C levels. Using a logistic mixed-effects

model with participant as a random effect, bNAb producers showed higher odds of elevated

GBV-C levels (odds ratio [OR]: 6.4, 95% CI: 2.3-17.9). In some participants, GBV-C read

counts exceeded those of HIV up to 100-fold.