An Adaptive Visuomotor Transformation Reservoir Embedded in the Vertebrate Brain

Yu Qian[1,2,4,*], Sha Li[1,3,4], Ming-Chuan Chen[1], Xu-Fei Du[1,*] & Jiu-Lin Du[1,2,3,*]

[1] Institute of Neuroscience, Laboratory of Brain Cognition and Brain-Inspired Intelligence Technology, Center for Excellence in Brain Science and Intelligence Technology, Chinese Academy of Sciences, 320 Yue-Yang Road, Shanghai 200031, China

[2] University of Chinese Academy of Sciences, 19A Yu-Quan Road, Beijing 100049, China

[3] School of Life Science and Technology, ShanghaiTech University, 319 Yue-Yang Road, Shanghai 200031, China

[4] These authors contributed equally.

[*] Correspondence E-mail: forestdu@ion.ac.cn; xufeidu@ion.ac.cn; yqian@ion.ac.cn

Abstract

Adaptive sensorimotor transformation is essential for animal survival in dynamic environments. The optic tectum (OT), homologous to the mammalian superior colliculus, serves as a central hub for visuomotor processing. To elucidate how OT circuits discriminate visual inputs and generate relevant behavior outputs, we leveraged the mesoscopic connectome of zebrafish OT to build a biophysically constrained reservoir computing network that models visuomotor transformations. *In silico* lesion revealed that the accuracy and robustness of sensorimotor outputs are governed by different combinations of tectal interneuron (TIN) subtypes. Furthermore, specific lamina-projecting serotoninergic subsystems distinctly bias network output toward either escape or orienting behaviors via regulating TIN activities. Our findings establish a mesoscopic connectome-based reservoir for vertebrate sensorimotor processing, where TINs control accuracy and robustness, highlighting TINs as a tunable gate and serotoninergic neurons as a context-dependent modulator of behavioral flexibility.

Introduction

Biological neural networks are the result of billions of years of evolutionary "hyperparameter tuning", optimizing the structure and function across generations. Unlike artificial neural networks that requires extensive data and iterative training, biological neural networks are endowed with innate computational capabilities. Even from birth, animals can perform complex, survival-critical behaviors such as threat avoidance, reward seeking with minimal experience[1–4]. These remarkable capabilities stem from hardwired circuit architectures whose core computational principles remain largely undeciphered. Understanding these intrinsic network structures is fundamental to uncovering the biological substrate of intelligence[5].

Vision is a primary sensory modality for animal to interpret dynamic environments. The superior colliculus (SC), an evolutionarily conserved sensorimotor hub[6], that rapidly converts retinotopic visual inputs into appropriate behaviors[7–10]. Prior studies have focused primarily on feedforward input-output pathways, providing limited insight into the local computational mechanisms embedded within the sensorimotor hub[11–15].

Larval zebrafish (*Danio rerio*) provide a unique model system for investigating these questions. Their compact and evolutionarily conserved sensorimotor hub-OT(the homologs of SC), optical transparency, and in particular a complete mesoscopic connectome mapped at single-cell resolution enable a level of structural access unmatched in vertebrate systems[16–20]. Yet despite these advantages, experimental interrogation of tectal circuit logic remains challenging. The extreme diversity and density of tectal neuron types make it physically intractable to selectively manipulate defined subnetworks or to systematically trace causal relationships between structure and function[21,22]. These technical constraints have limited our ability to uncover mechanisms about how neural architecture gives rise to appropriate visuomotor transformations.

Computational modeling offers a complementary approach to overcome these challenges. In particular, biologically constrained spiking neural networks (SNNs) grounded in empirical connectome data provide a powerful platform for mechanistic dissection of circuit dynamics[23]. Unlike abstract artificial networks, these models preserve biologically realistic connectivity patterns, neuron types, and temporal dynamics, allowing precise and scalable *in silico* manipulation of network components. This permits causal exploration of how specific subcircuits contribute to emergent population dynamics and behavior, especially in cases where experimental interventions are not feasible[24]. Furthermore, the reservoir computing framework provides a theoretical bridge between structural connectivity and dynamic computation. Reservoirs with fixed, recurrent connectivity, resembling biological networks, have been shown to perform complex temporal tasks such as memory, pattern separation, and dynamic input mapping[25,26].Importantly, biological reservoirs are not static in function. Rather, their dynamic regimes can be modulated by internal states and neuromodulatory influences, enabling a single circuit to flexibly switch between different computational modes[27]. This balance of innate structure and flexible dynamics is a defining feature of adaptive neural systems, yet remains underexplored in large-scale, biologically realistic models[28].

In this study, we leveraged these insights to reconstruct a biophysically constrained reservoir model using over 2,000 single-cell reconstructions of retinal ganglion cells (RGCs) and tectal neurons (TNs) in larval zebrafish. We implemented this architecture as a SNN to simulate visuomotor transformation and validated its emergent dynamics with *in vivo* calcium imaging data. *In silico* lesion experiments revealed that specific tectal interneuron (TIN) types are critical for preserving visuomotor accuracy and robustness. Then we demonstrated that this tectal-inspired

reservoir outperforms artificial recurrent neural networks in chaotic time-series prediction tasks,
such as the Lorenz attractor, underscoring the computational advantages of biologically grounded
architectures. Furthermore, we identified lamina-specific neuromodulation by serotoninergic (5-
HT) subsystems, which gate tectal network dynamics to enable flexible sensorimotor output.
Together, these results highlight a paradigm shift in computational neuroscience – moving beyond
abstract models toward biologically realistic simulations that reveal how structural motifs and
neuromodulation interact to produce adaptive neural computations.

Results

Identification of tectofugal pathways underlying escape and orienting behaviors

To investigate the intrinsic properties of the tectal network, we focused on two well-characterized
premotor targets involved in distinct behaviors as the output of the tectal network: the Mauthner
cell (M-cell), which drives rapid escape responses[29–31], and the nucleus of the medial longitudinal
fasciculus (nMLF), which is implicated in orienting and postural control[15,32,33](Fig. 1a).
Leveraging the mesoscopic connectome of larval zebrafish[16], we systematically analyzed the
structural connectivity of the OT with these downstream premotor nuclei.

TNs were classified into two principal types: TINs (also known as periventricular
interneurons, PVINs) and tectal projection neurons (TPNs, also referred to as periventricular
projection neurons, PVPNs) (Fig. 1b). TINs are largely confined to local intratectal circuits and
include both excitatory and inhibitory subtypes (Extended Data Fig. 1b). Inhibitory TINs are
thought to mediate lateral inhibition and contribute to visual feature extraction and contrast
enhancement. In contrast, TPNs serve as the primary output neurons of the tectum. These neurons
are predominantly excitatory and extend long-range projections to downstream premotor and
motor-associated brainstem regions (Extended Data Fig. 1a). In addition to TINs and TPNs, two
other cell types with somata located within the neuropil (Np) of the OT were identified. The first
is the superficial interneuron (SIN), an inhibitory neuron that modulates synaptic input in the
superficial tectal layers and is implicated in filtering sensory input[34,35]. The second is the
superficial projection neuron (SPN), which typically sends descending projections to the
contralateral hemisphere (Extended Data Fig. 1g).

To map input–output relations at synaptic resolution, we used sparse labeling of TNs via
UAS:sypb-APEX2-TagRFP plasmid injections to tag presynaptic release sites, enabling

morphological reconstruction and polarity inference. Axons and dendrites were algorithmically classified using a machine learning-based pipeline (Methods; Extended Data Fig. 1c). We further inferred putative synaptic connectivity using a columnar distance-based proximity algorithm, assigning connection weights based on the number of closely apposed axon–dendrite contact points falling below a defined distance threshold (Methods; Extended Data Fig. 1d).

Using this approach, we identified two distinct subtypes of TPNs based on their preferential connectivity to premotor targets. One subtype projects to the caudal dendritic compartment of the M-cell, which we term TPN-E (Escape-type TPNs), and is thus likely involved in initiating escape responses[29–31] (Fig. 1c, left). The other subtype preferentially innervates the nMLF, which we refer to as TPN-O (Orientation-type TPNs), likely contributing to orienting and posture-related visuomotor transformations[15,32,33] (Fig. 1c, right).

Retinotectal laminar and topographic organization subdivides the tectal neuropil into microdomains

To examine the spatial organization of retinal inputs onto tectal projection neurons (TPNs), we sparsely labeled individual retinal ganglion cells (RGCs) by microinjecting the plasmid *UAS:tdTomato-CAAX* into *Tg(Huc:H2b-GCaMP6s);Ki(isl2b:GAL4FF)* embryos. This strategy enabled single-cell resolution reconstructions of RGC morphologies, which we registered to a standardized brain template to produce a population-level retinotectal projection map (Extended Data Fig. 2). Among the 464 reconstructed RGCs, the vast majority projected exclusively to the contralateral OT, while a small minority targeted the contralateral intermediate hypothalamus (Extended Data Fig. 2b, c). Within the OT, RGC axon terminals exhibited precise laminar specificity, forming stereotyped arborization patterns across the tectal Np (Fig. 1d). Most RGCs terminated in superficial layers—specifically, in sublayers 1–6 of the stratum fibrosum et griseum superficiale (SFGS)—whereas a smaller proportion arborized in deeper strata, including the stratum opticum (SO), stratum griseum centrale (SGC), and stratum album centrale (SAC) (Extended Data Fig. 2f, g).

Consistent with established principles of retinotopy[15,36,37], the spatial positions of RGC somata within the retina predicted their tectal termination zones. Accordingly, we subdivided the tectal neuropil into four functional microdomains: anterior, posterior, medial, and lateral Np (Fig. 1e). Before entering the neuropil, RGC axons diverged into distinct arborization fields (AFs),

forming presynaptic terminals within specific laminar compartments. Based on branch morphologies and terminal distributions, we identified nine discrete arborization fields (AF1–AF9), each associated with a subset of retinotectal laminae (Extended Data Fig. 2d, e). Notably, two minor RGC branches with established behavioral functions were observed. One traversed AF7 to reach the SO layer, previously associated with high-resolution visual processing during prey detection[17,38]. The other projected via AF9 to the deeper SFGS5–6, SGC, or SAC layers—regions implicated in the encoding of global luminance changes, such as flashes, dimming, or looming stimuli[39,40] (Extended Data Fig. 2f, g). The angular size of stimuli conveyed through this latter pathway is a known determinant for initiating visually evoked escape behaviors.

Based on the six laminar strata and four topographic orientations projected from RGCs onto the Np, we subdivided the entire tectal Np into 24 distinct subdomains, each representing a unique retinotectal convergence zone (Figure 1f).

Simplified parallel visuomotor transformation pathways reveal functional segregation of escape and orienting circuits

Leveraging the mesoscopic connectome, the visuomotor transformation architecture in the OT was modeled based on anatomical assumptions: RGCs project axons into the neuropil (Np), TINs extend both dendrites and axons within the Np, and TPNs contribute primarily dendritic processes to receive processed information (Extended Data Fig. 1c). Analysis of dendritic distributions for the two identified TPN subtypes, TPN-E and TPN-O, across the 24 previously defined Np subdomains revealed subtype-specific spatial segregation.

Several subdomains contained dendrites from only one TPN subtype. For example, the SO-A subdomain was innervated exclusively by TPN-O, whereas SGC-P and SAC/SPV-P subdomains—together referred to as the SGC/SAC-P domain—were selectively targeted by TPN-E dendrites (Fig. 1g, h). Tracing upstream connectivity identified two distinct RGC subtypes with restricted projection fields: one terminating in SGC/SAC-P, the other in SO-A (Fig. 1i, j). Synaptic connectivity analysis confirmed this segregation: RGCs projecting to SGC/SAC-P were connected almost exclusively to TPN-E, while those projecting to SO-A selectively innervated TPN-O (Fig. 1k).

Downstream connectivity further supported this parallel organization. TPN-E projects selectively to the M-cell, whereas TPN-O targets the nMLF (Fig. 1l). Together, these connections

delineate two anatomically and functionally distinct visuomotor transformation pathways. This modular and topographically organized circuit structure suggests that discrete tectal microcircuits transform specific retinal inputs into behaviorally relevant outputs. Furthermore, the observed anterior–posterior segregation is consistent with prior findings[15,39,41–44] that posterior OT domains are preferentially involved in threat-evoked escape responses, whereas anterior regions support prey detection and orienting, reflecting the evolutionary compartmentalization of visuomotor processing in the vertebrate midbrain.

Simulation of tectal visuomotor transformation using SNN

The OT contains diverse TNs with distinct morphologies and distributions, forming a recurrent network capable of complex sensory integration and motor output selection. To investigate the computational roles of different TN subtypes in visuomotor transformation, a SNN was implemented using leaky integrate-and-fire (LIF) neuron models (see Methods).

To assess the validity of the model, initial simulations focused on the simplified parallel visuomotor transformation pathways described earlier. Brief current pulses (25 mV, 100 ms) were injected into either SO-A-projecting RGCs or SGC/SAC-P-projecting RGCs (Fig. 2a). Selective activation of the SO-A RGC population elicited strong spiking responses in TPN-O neurons, but failed to activate TPN-E (Fig. 2b). Conversely, stimulation of SGC/SAC-P RGCs robustly activated TPN-E while leaving TPN-O largely unresponsive (Fig. 2c).

This input specificity was preserved at the premotor level. Stimulation of the SGC/SAC-P pathway led to reliable activation of the M-cell. In contrast, SO-A pathway activation selectively recruited nMLF neurons (Fig. 2d). These simulations confirm that the modeled SNN recapitulates key anatomical and functional features of parallel tectal circuits and reliably transforms visual input patterns into subtype-specific motor-related outputs. This framework provides a mechanistic foundation for dissecting circuit-level computations underlying visuomotor behaviors in the vertebrate midbrain.

***In silico* ablation reveals TIN types contributing to visuomotor transformation accuracy**

Natural visual scenes contain rich spatiotemporal dynamics, requiring coordinated input from diverse RGC subtypes tuned to specific visual features. To provide realistic input to the tectal model, in vivo calcium imaging was performed under monocular visual stimulation. Zebrafish

larvae were immobilized in low-melting-point agarose, and stimuli were projected onto a screen contralateral to the recorded OT (Fig. 2f). Three behaviorally relevant stimuli were used: a small moving dot (SMD) representing prey-like motion, a neutral large moving dot (BMD), and a looming stimulus associated with threat detection. Calcium activity in the contralateral tectal Np was recorded across multiple layers and converted to estimate firing rates (see Methods). These time-varying inputs were fed into the SNN, which incorporated excitatory and inhibitory TINs forming a recurrent reservoir within the Np (Fig. 2g).

Under intact conditions, the model exhibited accurate and stimulus-specific visuomotor transformation without the need for training. SMD selectively activated TPN-O, while looming stimuli robustly engaged TPN-E, consistent with their respective downstream targets—nMLF for orienting and M-cell for escape behavior. However, in silico ablation of the TIN reservoir profoundly disrupted these dynamics. Network activity became pathologically elevated, and both TPN-O and TPN-E were coactivated in response to all stimulus types, indicating a loss of stimulus discrimination (Fig. 2h).

To quantify this disruption, incorrect TPN activations—e.g., TPN-E activation during SMD or TPN-O during looming—were classified as computational errors. Following TIN ablation, error rates significantly increased, reflecting a marked decline in the accuracy of visuomotor transformation (Fig. 2i). To dissect the contributions of individual TIN subtypes, targeted *in silico* ablations were performed by selectively zeroing connection weights from specific TIN morphologies in the connectome matrix (Fig. 2j). Among the various TIN subtypes (Extended Fig. 1h–j), the most profound deficits in visuomotor accuracy occurred when inhibitory bistratified TINs targeting both SFGS1-2 and SFGS5-6 (I_SFGS1-2_SFGS5-6; 21.1% accuracy reduction on average) were ablated, followed by the inhibitory non-stratified TINs (I_ns; 14.9% accuracy reduction on average) (Fig. 2k).

These findings demonstrate that TINs are essential for constraining tectal network activity, stabilizing recurrent dynamics, and ensuring population-level selectivity. Morphologically distinct inhibitory TINs play critical roles in shaping stimulus-specific outputs, enabling accurate visuomotor transformations from naturalistic inputs. This highlights the functional importance of intratectal processing in mediating behaviorally relevant decisions.

***In silico* ablation reveals TIN types contributing to network robustness**

Robustness to sensory noise is a hallmark of biological neural networks, enabling reliable behavior in unpredictable environments. To investigate the role of specific TIN subtypes in promoting robustness, a combination of behavioral and imaging experiments was performed using noisy visual stimuli. First, behavioral responses were recorded in freely swimming zebrafish exposed to looming stimuli embedded with varying levels of spatiotemporal noise (Fig. 3a). Escape probabilities were quantified using infrared video tracking. Subsequently, the same fish were immobilized for two-photon calcium imaging, allowing direct measurement of RGC activity in response to the same noisy stimuli (Fig. 3b). Using SGC-layer-projecting RGCs (SGC-RGCs) as a representative population, calcium signals revealed a gradual reduction in signal-to-noise ratio as stimulus noise density increased (Fig. 3b, c).

These empirical input patterns were then used to drive the computational SNN model. Looming stimuli with increasing noise levels were fed into the network, and escape probabilities were inferred from M-cell activity (Fig. 3d). Both the real zebrafish and the intact model exhibited notable robustness, maintaining high escape probabilities at low noise levels ($\rho < 0.5$). However, TIN network reshuffling or complete removal significantly impaired both response accuracy and robustness (Fig. 3e). To pinpoint which TIN subtypes are critical for maintaining robustness, targeted in silico ablation was performed at a representative noise level ($\rho = 0.5$). Among all TIN morphologies tested, the inhibitory non-stratified subtype (I_ns) exhibited the strongest impact, with a mean normalized reduction in escape probability of 13.15%. This was followed by bistratified inhibitory TINs projecting exclusively to the SGC and SAC layers (I_SGC_SAC), which produced an 11.23% average reduction in escape probability (Fig. 3f)..

These results demonstrate that distinct TIN morphotypes contribute differentially to the network's resilience against noisy input. Specifically, I_ns and I_SGC_SAC neurons appear to play stronger roles in maintaining decision fidelity under suboptimal sensory conditions, highlighting the functional diversity of intratectal interneurons in promoting behavioral robustness.

TIN reservoir exhibits robust generalization on Lorenz sequence denoising

To evaluate whether the robustness observed in the TIN reservoir extends beyond biologically inspired tasks and generalizes to machine learning domains, we employed the Lorenz chaotic time series—a standard benchmark in nonlinear dynamics and temporal prediction. This system is highly sensitive to initial conditions, such that even small perturbations can result in large

deviations over time, making it an ideal testbed for assessing noise tolerance and predictive stability. Noisy variants of the Lorenz sequence were generated by adding Gaussian noise of varying standard deviations (SD) and were used as input to the TIN reservoir model. In this setup, only the output weights (from reservoir to readout) were trained, while all internal weights remained fixed—analogous to standard reservoir computing frameworks (Fig. 3g). The goal was to reconstruct the clean Lorenz trajectory from the corrupted input.

Despite the high sensitivity of the chaotic sequence, the TIN reservoir consistently demonstrated strong denoising capabilities across noise levels. The output closely tracked the ground truth trajectory, even under moderate perturbations (Fig. 3h). Quantitative comparisons revealed that under low-noise conditions (SD ≤ 0.1), the TIN reservoir outperformed conventional recurrent neural networks (RNN), long short-term memory networks (LSTM), and gated recurrent units (GRU) of equivalent size in terms of predictive accuracy (Fig. 3i). Additionally, the TIN reservoir exhibited lower prediction error variance, further highlighting its noise robustness (Fig. 3j).

These results indicate that the structured recurrent dynamics of the TIN reservoir confer not only biological plausibility but also computational advantages in machine learning contexts. Specifically, its inherent ability to filter noise and preserve temporal information makes it a promising candidate for robust time-series modeling in both neuroscience and artificial intelligence applications.

Discrete serotonergic subsystems differentially modulate tectal output pathways to endow behavioral flexibility

In large-scale neural networks such as OT, which contains over 20,000 neurons and occupies more than one-fifth of the larval zebrafish brain volume. Behavioral fine-tuning through slow synaptic plasticity mechanisms (e.g., STDP, Hebbian learning, LTP, or LTD) is metabolically costly and temporally inefficient. Instead, fast-acting neuromodulatory systems have evolved to flexibly reconfigure circuit dynamics in a context-dependent manner[45–47], enabling adaptive behavioral outputs with minimal rewiring. Among these, the serotonergic (5-HT) system is a phylogenetically conserved monoaminergic pathway implicated in emotions, sleep-wake, learning and memory, reward and punishment modulation [48–54].In mammals, serotonergic projections directly innervate the superior colliculus[55,56], suggesting the candidate role in SC computation.

To investigate whether the 5-HT system directly shape tectal (SC) computations, we leveraged a comprehensive anatomical atlas and identified 5-HT neurons projecting to the OT neuropil(Fig. 4a, left). Serotonergic axons exhibited pronounced laminar specificity, targeting both superficial layers (SO) and deeper layers (SFGS 5–6 and SGC)—layers previously associated with distinct visual feature processing[37] (Fig. 4a, right). Based on axonal projection patterns, we identified two subtypes of 5-HT neurons: SO-projecting (SO-type) type, which arborizing in superficial layer (Fig. 4b), and SGC-projecting (SGC-type) type targeting deeper layers (Fig. 4c). In addition to their distinct axonal projection patterns, SO-type and SGC-type 5-HT neurons also exhibit significant differences in dendritic architecture. Notably, the dendrites of SGC-type neurons extend deep into the intermediate hypothalamus (Fig. 4c, bottom), suggesting potential regulation by internal states such as hunger or satiety. In vivo calcium imaging during light-dark transitions revealed divergent sensory tuning: SO-type neurons exhibited sustained responses to continuous illumination, consistent with a role in contextual background modulation, whereas SGC-type neurons showed transient responses to luminance changes, suggesting sensitivity to dynamic visual cues (Fig. 4d).

To examine the physiological impact of serotonergic modulation on tectal microcircuits, we performed whole-cell recordings of TINs. Bath application of 10 µM 5-HT significantly increased spontaneous firing rates (Fig. 4e, f), indicating that 5-HT can increase neuronal excitability. Thus, we computationally lowered the firing thresholds of TNs with dendrites in either the SO or SGC layers, to mimicking selective activation of serotonergic subtypes (Fig. 4g) in the spiking neural network (SNN) model. Simulation results revealed that activating the SO-type selectively enhanced TPN-O responses (Fig. 4i, top), whereas SGC-type preferentially increased TPN-E activity (Fig. 4i, bottom) with BMD stimulus input. However, the BMD stimulus that typically engages both output pathways (Fig. 4h). These findings suggest that laminar-selective serotonergic subsystems can dynamically bias tectal output based on contextual demands.

To test the behavioral relevance of this mechanism, we selectively ablated SGC-type 5-HT neurons using targeted two-photon laser ablation in (Fig. 4j). This manipulation significantly impaired looming-evoked escape behaviors in free-swimming zebrafish (Fig. 4k), confirming the critical role of SGC-type serotonergic modulation in threat-driven visuomotor transformation. Collectively, these results demonstrate that discrete serotonergic subsystems, defined by laminar innervation and functional tuning, differentially gate tectal output pathways. Such architecture

enables rapid and flexible reconfiguration of visuomotor responses based on internal states and sensory context, providing an efficient solution for behavioral adaptation in dynamic environments.

Discussion

This study establishes the zebrafish optic tectum (OT) as a biologically grounded reservoir computing network in which the morphological diversity of tectal interneurons (TINs) enables robust visuomotor transformations, and lamina-specific serotonergic modulation dynamically reconfigures output pathways to support behavioral flexibility. By integrating a large-scale mesoscopic connectome with a spiking neural network (SNN) constrained by single-cell morphology and synaptic topography, we provide a mechanistic account of how innate midbrain circuits perform flexible sensorimotor computations without the need for learning.

Our results reveal two principal mechanisms underlying this computation. First, the diversity of stratified inhibitory and excitatory TINs forms a heterogeneous recurrent network that expands the separability of sensory inputs, echoing the high-dimensional projection principle in artificial reservoir networks. This suggests that evolutionary diversification of interneuron morphology serves as a structural solution for enhancing temporal coding and stimulus discrimination[57,58]. Second, we show that serotonergic subsystems projecting to distinct tectal layers act as laminar-specific gain controllers: activating SO-projecting 5-HT neurons biases the network toward orienting behavior, while SGC-projecting 5-HT neurons enhance escape-related outputs. This demonstrates that a conserved anatomical feature—laminar targeting—can be repurposed to implement a form of neuromodulatory multiplexing, allowing the same hardwired circuit to flexibly switch behavioral modes depending on internal state or environmental context[59,60].

In contrast to prior tectal models that emphasized feedforward pathways or abstract dynamics, our reservoir model captures both the structural and functional complexity of the OT, enabling causal dissection of recurrent computations that are otherwise inaccessible to experimental perturbation. Moreover, the model generalizes to artificial tasks—such as Lorenz sequence denoising—outperforming conventional recurrent neural networks in noise-robust prediction. This underscores the computational advantage of biologically evolved network architectures, even outside their native domain[61]. Importantly, our findings also suggest a framework for reinterpreting tectal circuit motifs in terms of computational primitives. For instance, the parallel TPN-E and TPN-O pathways resemble a hardwired "output layer," while the TIN reservoir transforms

incoming RGC input into a high-dimensional population code that is selectively decoded based on behavioral context. This circuit design enables a form of real-time sensory selection, where behavior-specific outputs are recruited according to both sensory input and neuromodulatory state.

Despite these advances, several key questions remain. One outstanding issue is whether neuromodulators beyond serotonin, such as dopamine and norepinephrine, exhibit similarly stratified targeting in the OT and how these systems interact to coordinate flexible sensorimotor control[62,63]. Another question concerns the role of synaptic plasticity in shaping or refining tectal computations: while our model operates with fixed connectivity, experience-dependent modifications may adjust reservoir dynamics over longer timescales[64]. Additionally, it remains unknown how this tectal reservoir interacts with other midbrain and hindbrain structures to coordinate distributed sensorimotor transformations—a question that will require multi-region imaging and perturbation in freely behaving animals.

Finally, our work raises intriguing developmental questions: How do such precise morphologies and laminar targeting rules arise during circuit assembly? Do different TIN subtypes follow distinct genetic programs or activity-dependent cues? Addressing these issues could provide fundamental insights into how evolution sculpts flexible yet robust neural systems through constrained developmental rules.

In sum, our study offers a biologically grounded computational framework that unifies structural connectivity, interneuron diversity, and neuromodulation into a coherent model of real-time sensorimotor processing. These findings underscore that biological structure is not merely descriptive—it is fundamentally computational. By embracing this principle, we may uncover generalizable algorithms from the brain's architecture to inspire more robust and adaptable artificial intelligence.

Materials and methods

Zebrafish husbandry and preparation

Zebrafish larvae were maintained under standard conditions at 28.5°C with a 14 h light/10 h dark photoperiod. Larvae were kept in Hank's solution until 2 dpf, after which they were transferred to facility water. Beginning at 5 dpf, larvae were fed rotifers once daily. All experiments were conducted using larvae aged 6–7 dpf.

For in vivo imaging experiments, larvae were embedded in 2.0% low-melting-point agarose (Sigma) dissolved in facility water. 5 minutes post-embedding, facility water was added to the recording chamber. For whole-cell recordings, facility water was replaced with an external solution containing (in mM): 134 NaCl, 2.9 KCl, 2.1 CaCl$_2$·2H$_2$O, 1.2 MgCl$_2$·6H$_2$O, 10 HEPES, and 10 D-glucose (pH 7.8, 290 mOsm).

Zebrafish lines

Transgenic zebrafish larvae were in a nacre background[65]. Fishline *Tg(-7atoh7:GAL4-VP16)s1992t* (in abbreviation: *Tg(-7atoh7:GAL4-VP16)*)[66], *Tg(UAS:GCaMP6s)nkUAShspzGCaMP6s13a* (*Tg(UAS:GCaMP6s)*)[67], *Tg2(elavl3:GCaMP6s)a13203* (*Tg2(elavl3:GCaMP6s)*)[68] were described in previous studies, respectively.

Tg(dbh:GAL4FF,5×UAS-hsp:Rno.Trpv1-TagRFP)ion88d (*Tg(dbh:GAL4FF,UAS:Trpv1-TagRFP)*) was constructed by Dr. Chang-Mei Zhang at Jiu-Lin Du's laboratory with Tol2 system. *Ki(tph2:GAL4FF,myl7:EGFP)ion127d* (*Ki(tph2:GAL4FF,myl7:EGFP)*)line was constructed by Dr. Fu-Nin Li at Jiu-Lin Du's laboratory with a CRISPR–Cas9-mediated knock-in approach. *Tg(elavl3:Hsa.H2B-GCaMP8f)ion1m* (*Tg(elavl3:Hsa.H2B-GCaMP8f)*) was constructed by Yu Mu's Laboratory.

SNN construction

In this study, we constructed a biologically plausible SNN using BrainPy, a flexible and high-performance simulation framework for spiking dynamics and neural computation[69]. The neuronal units are modeled using the LIF neuron with an absolute refractory period, capturing key temporal dynamics observed in vertebrate neurons.

395   The subthreshold dynamics of membrane potential $V(t)$ are governed by:

396
$$\tau \frac{dV(t)}{dt} = -\left(V(t) - V_{rest}\right) + R_m I_{syn}(t), \text{for } t \notin \left[t_{spike}, t_{spike} + \tau_{ref}\right]$$

Where $V(t)$ is the membrane potential, V_{rest} is the resting potential, V_{reset} is the reset potential

after a spike, V_{th} is the spike threshold, τ is the membrane time constant, τ_{ref} is the refractory

period, R_m is the membrane resistance, $I_{syn}(t)$ is the total synaptic current input.

When $V(t) \geq V_{th}$, the neuron emits a spike and resets to V_{reset}, remaining inactive for the duration

of τ_{ref}. The detailed parameterization for each modeled neuron type is listed in Supplementary

Table 1.

Synaptic inputs are modeled using single-exponential decay conductance-based synapses, a

commonly used approximation in SNN modeling that assumes rapid neurotransmitter release,

diffusion, and receptor dynamics. We included two types of synapses: AMPA (excitatory)

synapses with a short decay time constant; GABA (inhibitory) synapses with a longer decay time

constant and negative reversal potential. The conductance $g(t)$ of a synapse after a presynaptic

spike at time t_k evolves as:

409
$$g(t) = \bar{g} \sum_k \exp\left(-\frac{t - t_k}{\tau_s}\right) H\left(t - t_k\right)$$

410   Where $\bar{g}$ is the maximal synaptic conductance, τ_s is the synaptic decay time constant

411   ($\tau_{AMPA} = 5$ ms, $\tau_{GABA} = 10$ ms), t_k is the time of the k-th presynaptic spike, V_{th} is the spike

412   threshold, $H\left(t - t_k\right)$ is the Heaviside step function.

413   The corresponding differential form is:

414
$$\frac{dg(t)}{dt} = -\frac{g(t)}{\tau_s} + \bar{g} \sum_k \delta\left(t - t_k\right)$$

415   The total synaptic current into the postsynaptic neuron is:

416
$$I_{syn}(t) = \sum_i g_i(t) \cdot \left(E_i - V(t)\right)$$

Where E_i is the synaptic reversal potential ($E_{AMPA} = 0$ mV, $E_{GABA} = -80$ mV), and $g_i(t)$ is the

time-varying conductance of the i-th synapse.

The number of excitatory and inhibitory neurons in the network is scaled according to the

anatomically derived neuron counts from each zebrafish brain region. To simulate *in silico*

knockout of a specific neuron type, we set the connection probability between the targeted neurons

and all other neurons to zero:

423
$$P_{\text{conn}}(i,j) = \{0, \quad \text{if } i \in \mathcal{K} \qquad \mathcal{K} \qquad), \quad \text{otherwise}]$$

where $P_{\text{conn}}(i,j)$ is the connection probability between presynaptic neuron i and postsynaptic

neuron j, $P_{\text{base}}(i,j)$ is the baseline (non-zero) probability determined by the connectome or

anatomical prior, and $\mathcal{K}$ is the set of neurons targeted for knockout.

Calcium fluorescence signals to spike trains

To simulate the effect of RGCs' activity, we convert experimentally recorded calcium fluorescence

signals into Poisson-distributed spike trains, where the normalized calcium signal determines the

instantaneous firing rate. These spike trains are then used as input to downstream neurons in the

model:

432
$$s_i(t) \sim \text{Poisson}(\lambda_i(t)), \quad \lambda_i(t) = \lambda_{\text{max}} \cdot \frac{C_i(t)}{C_{\text{max}}}$$

where $s_i(t)$ is the binary spike train (0 or 1) for upstream neuron i at time t, $\lambda_i(t)$ is the time-

varying firing rate, $C_i(t)$ is the calcium signal (e.g., $\Delta F/F$) of neuron i at time t, C_{max} is the

maximum value of $C_i(t)$ over time for normalization, λ_{max} is the maximum firing rate.

TIN reservoir construction and comparative modeling

We constructed a biologically inspired reservoir network using zebrafish TINs, where the

connection weights were determined by the estimated number of potential synapses between

neurons based on anatomical connectivity. Excitatory neurons were assigned positive weights,

while inhibitory neurons were assigned negative weights. To ensure the reservoir dynamics remain

in the echo state regime, we constrained the spectral radius $\rho(W)$ of the recurrent weight matrix

W to approximately 1:

443
$$\rho(W) \approx 1$$

For assessing the accuracy and robustness of visual motion transformations, the network is
configured with 6 input nodes and 2 output nodes. For benchmarking using the Lorenz attractor
sequence, both the input and output dimensions are adjusted to 3, and the system is converted into
a standard Echo State Network (ESN) structure for direct comparison with traditional recurrent
neural networks.

The ESN update rule is given by:

$$\mathbf{x}(t+1) = f\left(W_{in}\mathbf{u}(t) + W\mathbf{x}(t) + \mathbf{b}\right), \quad \mathbf{y}(t) = W_{out}\mathbf{x}(t)$$

where $\mathbf{x}(t)$ is the reservoir state vector, $\mathbf{u}(t)$ is the input vector, W_{in}, W, W_{out} are the input,
recurrent, and output weight matrices, respectively. $\mathbf{b}$ is the bias term, $f(\cdot)$ is a nonlinear tanh
activation function.

The Lorenz system used as a benchmark task is defined by the following set of differential
equations:

456
$$\begin{cases} \dfrac{dx}{dt} = \sigma(y-x) \\ \dfrac{dy}{dt} = x(\rho-z) - y \\ \dfrac{dz}{dt} = xy - \beta z \end{cases}$$

457 with standard parameters $\sigma = 10$, $\rho = 28$, and $\beta = 8/3$, yielding a chaotic trajectory.
458

In the visuomotor transformation task, the reservoir performs feedforward processing without
weight updates. However, in the Lorenz task, output weights W_{out} are trained using ridge
regression (L2-regularized linear regression):

$$W_{out} = \mathbf{Y}_{target} \mathbf{X}^\top \left(\mathbf{X}\mathbf{X}^\top + \alpha I\right)^{-1}$$

where $\mathbf{X}$ is the matrix of reservoir states, $\mathbf{Y}_{target}$ is the target output, α is the regularization
coefficient, I is the identity matrix. For benchmarking, we compared our model with several
standard recurrent architectures:

Vanilla RNN:

$$\mathbf{h}_t = \tanh\left(W_{hh}\mathbf{h}_{t-1} + W_{xh}\mathbf{x}_t + \mathbf{b}_h\right), \quad \mathbf{y}_t = W_{hy}\mathbf{h}_t + \mathbf{b}_y$$

LSTM (Long Short-Term Memory):

$$\mathbf{f}_t = \sigma\left(W_f \mathbf{x}_t + U_f \mathbf{h}_{t-1} + \mathbf{b}_f\right)$$

$$\mathbf{i}_t = \sigma\left(W_i \mathbf{x}_t + U_i \mathbf{h}_{t-1} + \mathbf{b}_i\right)$$

$$\mathbf{o}_t = \sigma\left(W_o \mathbf{x}_t + U_o \mathbf{h}_{t-1} + \mathbf{b}_o\right)$$

$$\tilde{} \quad W_c \mathbf{x}_t + U_c \mathbf{h}_{t-1} + \mathbf{b}_c\right)$$

$$\mathbf{c}_t = \mathbf{f}_t \odot \qquad \odot$$

$$\mathbf{h}_t = \mathbf{o}_t \odot \qquad)$$

GRU (Gated Recurrent Unit):

$$\mathbf{z}_t = \sigma\left(W_z \mathbf{x}_t + U_z \mathbf{h}_{t-1} + \mathbf{b}_z\right)$$

$$\mathbf{r}_t = \sigma\left(W_r \mathbf{x}_t + U_r \mathbf{h}_{t-1} + \mathbf{b}_r\right)$$

$$\mathbf{h}_t = \tanh\left(W_h \mathbf{x}_t + U_h \left(\mathbf{r}_t \odot \qquad \mathbf{'}_h\right)\right)$$

$$\mathbf{h}_t = \left(1 - \mathbf{z}_t\right) \odot \qquad \odot$$

To evaluate the robustness of the reservoir to noisy inputs, we added Gaussian noise to the Lorenz time series. The noise level was controlled by the SD parameter. The noisy input $\mathbf{x}(t)$ is defined as:

$$\mathbf{x}(t) = \mathbf{x}(t) + \mathcal{N}$$

where $\mathbf{x}(t)$ is the original Lorenz time series at time t, $\mathcal{N}$ represents zero-mean Gaussian noise with variance σ^2, σ is the standard deviation regulating the noise level.

The dataset was split as follows: The first 15% of the sequence was used as a warm-up period to allow the reservoir dynamics to stabilize; the next 70% was used as the training set; the final 15% was reserved for testing the model's prediction accuracy.

Finally, to quantify model performance, we employed the Lorenz chaotic time series to evaluate the predictive capability of each reservoir configuration. The Mean Squared Error (MSE) between the predicted time series and the ground truth was used as the primary evaluation metric. The MSE is defined as:

$$\mathrm{MSE} = \frac{1}{T} \sum_{t=1}^{T} \left\| \mathbf{y}_{\mathrm{pred}}(t) - \mathbf{y}_{\mathrm{true}}(t) \right\|^2$$

where T is the total number of time steps, $\mathbf{y}_{\mathrm{pred}}(t)$ is the predicted output at time t, $\mathbf{y}_{\mathrm{true}}(t)$ is the ground truth at time t, $\|\cdot\|$ denotes the Euclidean norm.

***In vivo* calcium imaging**

Calcium imaging experiments were usually performed under room temperature on an Olympus

Fluoview FVMPE-RS Multiphoton Laser Scanning Microscope with a 20X/1.0 NA water-

immersion objective, with 4-5 Hz frame rate (Galvano scanning mode). *Tg(-7atoh7:GAL4-*

VP16);Tg(UAS:GCaMP6s)   for   RGC   calcium   imaging.

Ki(tph2:GAL4FF,myl7:EGFP);Tg(UAS:GCaMP6s) for serotoninergic fiber calcium imaging.

Calcium imaging paired with whole-cell recording experiment was performed under room

temperature on an Olympus Fluoview FV3000 Confocal Laser Scanning Microscope with a

20X/1.0 NA water-immersion objective (Resonant scanning mode), with 30-40 Hz frame rate

(resonant scanning mode). *Tg(elavl3:Hsa.H2B-GCaMP8f)* for Calcium imaging paired with

whole-cell recording experiment. Images with severe drifts were abandoned.

Calcium activity analysis

Images were aligned using the MultiStackReg plugin in ImageJ. Regions of interest (ROIs) were

manually defined based on morphology and regional signal activity. Ca^{2+} signals were extracted

by the mean gray value of ROI. $\Delta F/F_0$ as computed using a custom MATLAB script, where F0

was calculated by the mean filter of Ca^{2+} signals with the function ordfilt2. Response peak

amplitude was defined as the maximum $\Delta F/F0$ within 5 s of stimulus onset, minus the baseline

average in the 10 s preceding stimulation. The area under the curve (AUC) was quantified over

the 10 s post-stimulus window. Calcium transients exceeding baseline mean ± 2 SD were

considered significant events.

Visual stimulation

All stimuli were presented monocularly using a mini-projector positioned lateral to the larva, with

a red filter applied to minimize imaging interference.

-Looming stimulus: A black dot expanded at a constant rate on a bright background until

filling the visual field over ~3.7 s [70]; 5 trials/fish with 120 s intervals.

Moving-dot stimuli: A black dot moved across a white background. Small and large dots

subtended 4° and 16°, respectively, with the same moving velocity; 5 trials/fish with 120 s intervals.

Light–dark stimulation: 1 min of bright illumination (1.23 mW) followed by 1 min of

darkness (0.167 mW), repeated three times for a total of 6 min.

***In vivo* electrophysiological recording**

Larvae were paralyzed with 100 μg/ml α-bungarotoxin (Sigma) for 10–15 min, then embedded in

2% low-melt agarose. In whole-cell recording experiments, micropipettes were filled with internal

solution, which consisted of (in mM): 110 K-gluconate, 6 NaCl, 2 MgCl2, 2 CaCl2, 10 HEPES,

and 10 EGTA (pH 7.41, 285 mOsm). The calculated chloride reversal potential (E_Cl) was ∼−60

mV. In whole-cell recording TNs paired calcium response in *Tg(elavl3:Hsa.H2B-GCaMP8f)*,

EGTA was replaced with phosphocreatine to avoid calcium chelation. In vivo whole-cell

recording of TNs with bath 5-HT solution (dissolved in DMSO, 10mM for stock solution) was

performed in wild-type.

For in vivo whole-cell recording of TNs, Following paralysis, a small incision was made at

the skin between the tectum and hindbrain using a glass pipette (~1 μm tip). After filled with the

internal solution, the recording micropipette (20 - 25 MΩ in resistance, 1-mm tip opening) was

advanced into the brain through the dissected hole and slowly approached to the TNs with a

persistent positive pressure to keep tip clean. After the contact of micropipette tip with TNs

membrane, giga-ohm seal was formed by removing the positive pressure and applying a slight

negative pressure. Whole-cell recording was achieved by delivering a few brief electrical zaps (50

μs to 2 ms) to break the cell membrane beneath the micropipette tip. Recordings were made with

a patch-clamp amplifier (MultiClamp 700B, Axon Instruments), and signals were filtered at 5 kHz

and sampled at 10 kHz by using Clampex 10.2 (Molecular Devices).

Behavioral assay

Looming-evoked escape responses were captured using a high-speed, infrared-sensitive camera

(Redlake Motionscope M3). Each larva was tested in five trials. Escape probability was defined as

the number of escape responses across trials. Body posture was quantified using skeletonization

algorithms, followed by an angular threshold to detect C-start escape responses[71]. All automated

detections were confirmed manually.

Two-photon ablation of SGC-type 5-HT neurons

Two-photon ablations were performed on an Olympus FVMPE-RS multiphoton microscope.

SGC-type 5-HT neurons were ablated based on both visual response properties and TagRFP

expressed in these neurons in *Tg2(elavl3:GCaMP6s);Ki(tph2:GAL4FF,myl7:EGFP);*

Tg(dbh:GAL4FF,UAS:Trpv1-TagRFP) larvae. Neurons were ablated by focused 800-nm laser

scanning (~2.92 mW) on the soma (~1 μm radius) for 10–12 ms at a constant angular speed. Cellular fragmentation could be observed after successful ablation. Each target was sequentially ablated by manually positioning the ROI on its soma. After ablation, larvae were recovered in facility water for 18–24 h before post-ablation testing.

Acknowledgments

We thank Chao-ming Wang, Tian-qiu Zhang, Si Wu and Yu Mu for their kind support. We appreciate the technical assistance from Hong-Li Wan, Hui Zhang and Qian Hu. We acknowledge the assistance with neuronal morphology registration on the brain template from the Facility of Mapping Brain-wide Mesoscale Connectome at CEBSIT (Institute of Neuroscience) for technical support. This work was supported by National Science and Technology Innovation 2030 Major Program of the Ministry of Science and Technology (2021ZD0204500 and 2021ZD0204502), Science Fund for Creative Research Groups (32321003) and Key Joint Funds for Cooperation (U22B200571) of the National Natural Science Foundation of China, Shanghai Municipal Science and Technology Major Project (18JC1410100 and 2018SHZDZX05), and Key Research Program of Frontier Sciences (QYZDYSSW-SMC028) and Strategic Priority Research Program (XDB32010200) of Chinese Academy of Sciences.

Author contributions

Conceptualization: Y.Q., J.D.; Code writing and brain simulation: Y.Q.; Methodology: Q.Y.; Investigation: Y.Q., S.L., J.D.; Data acquisition and pre-processing of wet experiments (including neuron morphology imaging and reconstruction, calcium imaging, and electrophysiology): S.L., Q.Y., M.C., X.D.; Funding acquisition: J.D.; Project administration: Y.Q., S.L., J.D.; Supervision: J.D., X.D.; Manuscript writing: Y.Q., S.L., J.D., X.D.

Data availability

Relevant data and codes have been released on GitHub: https://github.com/yqian-CEBSIT/Reservoir-of-Zebrafish-Optic-tectum.

Competing interests

The authors declare no competing financial interests.

References

1. Zador, A. M. A critique of pure learning and what artificial neural networks can learn from
animal brains. *Nat. Commun.* **10**, 3770 (2019).

2. Spelke, E. S. & Kinzler, K. D. Core knowledge. *Dev. Sci.* **10**, 89–96 (2007).

3. Hassabis, D., Kumaran, D., Summerfield, C. & Botvinick, M. Neuroscience-Inspired Artificial
Intelligence. *Neuron* **95**, 245–258 (2017).

4. He, L. *et al.* Network model with internal complexity bridges artificial intelligence and
neuroscience. *Nat. Comput. Sci.* **4**, 584–599 (2024).

5. Luo, L. Architectures of neuronal circuits. *Science* **373**, eabg7285 (2021).

6. Isa, T., Marquez-Legorreta, E., Grillner, S. & Scott, E. K. The tectum/superior colliculus as
the vertebrate solution for spatial sensory integration and action. *Curr. Biol.* **31**, R741–R762
(2021).

7. Basso, M. A. & May, P. J. Circuits for Action and Cognition: A View from the Superior
Colliculus. *Annu. Rev. Vis. Sci.* **3**, 197–226 (2017).

8. Kardamakis, A. A., Pérez-Fernández, J. & Grillner, S. Spatiotemporal interplay between
multisensory excitation and recruited inhibition in the lamprey optic tectum. *eLife* **5**, e16472
(2016).

9. Kardamakis, A. A., Saitoh, K. & Grillner, S. Tectal microcircuit generating visual selection
commands on gaze-controlling neurons. *Proc. Natl. Acad. Sci.* **112**, (2015).

10. Suzuki, D. G., Pérez-Fernández, J., Wibble, T., Kardamakis, A. A. & Grillner, S. The role of
the optic tectum for visually evoked orienting and evasive movements. *Proc. Natl. Acad. Sci.*
116, 15272–15281 (2019).

11. Naumann, R. K. & Laurent, G. Function and Evolution of the Reptilian Cerebral Cortex. in

Evolutionary Neuroscience 213–245 (Elsevier, 2020). doi:10.1016/B978-0-12-820584-

6.00009-X.

12. Wylie, D. R. W., Gutierrez-Ibanez, C., Pakan, J. M. P. & Iwaniuk, A. N. The optic tectum of

birds: Mapping our way to understanding visual processing. *Can. J. Exp. Psychol. Rev. Can.*

Psychol. Expérimentale **63**, 328–338 (2009).

13. Cang, J., Savier, E., Barchini, J. & Liu, X. Visual Function, Organization, and Development

of the Mouse Superior Colliculus. *Annu. Rev. Vis. Sci.* **4**, 239–262 (2018).

14. Robles, E., Laurell, E. & Baier, H. The Retinal Projectome Reveals Brain-Area-Specific

Visual Representations Generated by Ganglion Cell Diversity. *Curr. Biol.* **24**, 2085–2096

(2014).

15. Helmbrecht, T. O., Dal Maschio, M., Donovan, J. C., Koutsouli, S. & Baier, H. Topography

of a Visuomotor Transformation. *Neuron* **100**, 1429-1445.e4 (2018).

16. Du, X. *et al.* Central nervous system atlas of larval zebrafish constructed using the morphology

of single excitatory and inhibitory neurons. *bioRxiv* 2025.06.06.658008 (2025)

doi:10.1101/2025.06.06.658008.

17. Förster, D. *et al.* Retinotectal circuitry of larval zebrafish is adapted to detection and pursuit

of prey. *eLife* **9**, e58596 (2020).

18. Simmich, J., Staykov, E. & Scott, E. Zebrafish as an appealing model for optogenetic studies.

in *Progress in Brain Research* vol. 196 145–162 (Elsevier, 2012).

19. Vanwalleghem, G. C., Ahrens, M. B. & Scott, E. K. Integrative whole-brain neuroscience in

larval zebrafish. *Curr. Opin. Neurobiol.* **50**, 136–145 (2018).

20. Scott, E. K. The cellular architecture of the larval zebrafish tectum, as revealed by Gal4

enhancer trap lines. *Front. Neural Circuits* **3**, (2009).

21. Alivisatos, A. P. *et al.* The Brain Activity Map Project and the Challenge of Functional
Connectomics. *Neuron* **74**, 970–974 (2012).

22. Bauer, A. Q. *et al.* Effective Connectivity Measured Using Optogenetically Evoked
Hemodynamic Signals Exhibits Topography Distinct from Resting State Functional
Connectivity in the Mouse. *Cereb. Cortex* **28**, 370–386 (2018).

23. Sun, Z., Cutsuridis, V., Caiafa, C. F. & Solé-Casals, J. Brain Simulation and Spiking Neural
Networks. *Cogn. Comput.* **15**, 1103–1105 (2023).

24. Gerstner, W., Sprekeler, H. & Deco, G. Theory and Simulation in Neuroscience. *Science* **338**,
60–65 (2012).

25. Suárez, L. E. *et al.* Connectome-based reservoir computing with the conn2res toolbox. *Nat.*
Commun. **15**, 656 (2024).

26. Damicelli, F., Hilgetag, C. C. & Goulas, A. Brain connectivity meets reservoir computing.
PLOS Comput. Biol. **18**, e1010639 (2022).

27. Cazettes, F. *et al.* A reservoir of foraging decision variables in the mouse brain. *Nat. Neurosci.*
26, 840–849 (2023).

28. Shine, J. M. Neuromodulatory control of complex adaptive dynamics in the brain. *Interface*
Focus **13**, 20220079 (2023).

29. Temizer, I., Donovan, J. C., Baier, H. & Semmelhack, J. L. A Visual Pathway for Looming-
Evoked Escape in Larval Zebrafish. *Curr. Biol.* **25**, 1823–1834 (2015).

30. Kohashi, T. & Oda, Y. Initiation of Mauthner- or Non-Mauthner-Mediated Fast Escape
Evoked by Different Modes of Sensory Input. *J. Neurosci.* **28**, 10641–10653 (2008).

31. Korn, H. & Faber, D. S. The Mauthner Cell Half a Century Later: A Neurobiological Model
for Decision-Making? *Neuron* **47**, 13–28 (2005).

32. Thiele, T. R., Donovan, J. C. & Baier, H. Descending Control of Swim Posture by a Midbrain Nucleus in Zebrafish. *Neuron* **83**, 679–691 (2014).

33. Severi, K. E. *et al.* Neural Control and Modulation of Swimming Speed in the Larval Zebrafish. *Neuron* **83**, 692–707 (2014).

34. Baier, H. & Scott, E. K. The Visual Systems of Zebrafish. *Annu. Rev. Neurosci.* **47**, 255–276 (2024).

35. Abbas, F. & Meyer, M. P. Fish Vision: Size Selectivity in the Zebrafish Retinotectal Pathway. *Curr. Biol.* **24**, R1048–R1050 (2014).

36. Kunst, M. *et al.* A Cellular-Resolution Atlas of the Larval Zebrafish Brain. *Neuron* **103**, 21-38.e5 (2019).

37. Baier, H. & Scott, E. K. The Visual Systems of Zebrafish. *Annu. Rev. Neurosci.* **47**, 255–276 (2024).

38. Semmelhack, J. L. *et al.* A dedicated visual pathway for prey detection in larval zebrafish. *eLife* **3**, e04878 (2014).

39. Temizer, I., Donovan, J. C., Baier, H. & Semmelhack, J. L. A Visual Pathway for Looming-Evoked Escape in Larval Zebrafish. *Curr. Biol.* **25**, 1823–1834 (2015).

40. Dunn, T. W. *et al.* Neural Circuits Underlying Visually Evoked Escapes in Larval Zebrafish. *Neuron* **89**, 613–628 (2016).

41. Zhao, P. *et al.* The visuomotor transformations underlying target-directed behavior. *Proc. Natl. Acad. Sci.* **122**, e2416215122 (2025).

42. Bianco, I. H. & Engert, F. Visuomotor Transformations Underlying Hunting Behavior in Zebrafish. *Curr. Biol.* **25**, 831–846 (2015).

43. Fernandes, A. M. *et al.* Neural circuitry for stimulus selection in the zebrafish visual system. *Neuron* **109**, 805-822.e6 (2021).

44. Preuss, S. J., Trivedi, C. A., vom Berg-Maurer, C. M., Ryu, S. & Bollmann, J. H. Classification
of Object Size in Retinotectal Microcircuits. *Curr. Biol.* **24**, 2376–2385 (2014).

45. Bessonova, Y. & Raman, B. Serotonergic amplification of odor-evoked neural responses maps
flexibly onto behavioral outcomes.

46. Cohn, R., Morantte, I. & Ruta, V. Coordinated and Compartmentalized Neuromodulation
Shapes Sensory Processing in Drosophila. *Cell* **163**, 1742–1755 (2015).

47. Aston-Jones, G., Rajkowski, J. & Cohen, J. Role of locus coeruleus in attention and behavioral
flexibility. *Biol. Psychiatry* **46**, 1309–1320 (1999).

48. Marcinkiewcz, C. A. *et al.* Serotonin engages an anxiety and fear-promoting circuit in the
extended amygdala. *Nature* **537**, 97–101 (2016).

49. Wang, Y. *et al.* Dorsal Raphe Serotonergic Neurons-Ventral Tegmental Area Neural Pathway
Promotes Wake From Sleep. *CNS Neurosci. Ther.* **30**, e70141 (2024).

50. Oikonomou, G. *et al.* The Serotonergic Raphe Promote Sleep in Zebrafish and Mice. *Neuron*
103, 686-701.e8 (2019).

51. Harkin, E. F., Grossman, C. D., Cohen, J. Y., Béïque, J.-C. & Naud, R. A prospective code for
value in the serotonin system. *Nature* (2025) doi:10.1038/s41586-025-08731-7.

52. Kawashima, T., Zwart, M. F., Yang, C.-T., Mensh, B. D. & Ahrens, M. B. The Serotonergic
System Tracks the Outcomes of Actions to Mediate Short-Term Motor Learning. *Cell* **167**,
933-946.e20 (2016).

53. Cohen, J. Y., Amoroso, M. W. & Uchida, N. Serotonergic neurons signal reward and
punishment on multiple timescales. *eLife* **4**, e06346 (2015).

54. Ren, J. *et al.* Anatomically Defined and Functionally Distinct Dorsal Raphe Serotonin Sub-
systems. *Cell* **175**, 472-487.e20 (2018).

55. Muzerelle, A. Conditional anterograde tracing reveals distinct targeting of individual serotonin

cell groups (B5–B9) to the forebrain and brainstem. *Brain Struct Funct* (2016).

56. Ren, J. *et al.* Single-cell transcriptomes and whole-brain projections of serotonin neurons in

the mouse dorsal and median raphe nuclei. *eLife* **8**, e49424 (2019).

57. Buonomano, D. V. & Maass, W. State-dependent computations: spatiotemporal processing in

cortical networks. *Nat. Rev. Neurosci.* **10**, 113–125 (2009).

58. Maass, W., Natschläger, T. & Markram, H. Real-Time Computing Without Stable States: A

New Framework for Neural Computation Based on Perturbations. *Neural Comput.* **14**, 2531–

2560 (2002).

59. Dayan, P. Twenty-Five Lessons from Computational Neuromodulation. *Neuron* **76**, 240–256

(2012).

60. Disney, A. A., Aoki, C. & Hawken, M. J. Gain Modulation by Nicotine in Macaque V1.

Neuron **56**, 701–713 (2007).

61. Winnubst, J. *et al.* Reconstruction of 1,000 Projection Neurons Reveals New Cell Types and

Organization of Long-Range Connectivity in the Mouse Brain. *Cell* **179**, 268-281.e13 (2019).

62. Morilak, D. A. *et al.* Role of brain norepinephrine in the behavioral response to stress. *Prog.*

Neuropsychopharmacol. Biol. Psychiatry **29**, 1214–1224 (2005).

63. Devilbiss, D. M. & Waterhouse, B. D. Phasic and Tonic Patterns of Locus Coeruleus Output

Differentially Modulate Sensory Network Function in the Awake Rat. *J. Neurophysiol.* **105**,

69–87 (2011).

64. Zenke, F., Agnes, E. J. & Gerstner, W. Diverse synaptic plasticity mechanisms orchestrated to

form and retrieve memories in spiking neural networks. *Nat. Commun.* **6**, 6922 (2015).

65. Lister, J. A., Robertson, C. P., Lepage, T., Johnson, S. L. & Raible, D. W. nacre encodes a

zebrafish microphthalmia-related protein that regulates neural-crest-derived pigment cell fate.

Development **126**, 3757–3767 (1999).

66. Del Bene, F. *et al.* Filtering of Visual Information in the Tectum by an Identified Neural Circuit.

Science **330**, 669–673 (2010).

67. Muto, A. *et al.* Activation of the hypothalamic feeding centre upon visual prey detection. *Nat.*

Commun. **8**, 15029 (2017).

68. Kim, D. H. *et al.* Pan-neuronal calcium imaging with cellular resolution in freely swimming

zebrafish. *Nat. Methods* **14**, 1107–1114 (2017).

69. Wang, C. *et al.* BrainPy, a flexible, integrative, efficient, and extensible framework for

general-purpose brain dynamics programming. *eLife* **12**, e86365 (2023).

70. Yao, Y. *et al.* Visual Cue-Discriminative Dopaminergic Control of Visuomotor

Transformation and Behavior Selection. *Neuron* **89**, 598–612 (2016).

71. Dunn, T. W. *et al.* Neural Circuits Underlying Visually Evoked Escapes in Larval Zebrafish.

Neuron **89**, 613–628 (2016).

Figures

Fig. 1. Anatomical substrates for transforming visual information into escape and orienting behaviors in the zebrafish brain.

a, Two distinct descending pathways in zebrafish mediating orienting and escape behaviors, driven by premotor neurons in the nucleus of the nMLF and M-cells. Schematic representation (left) and mesoscale connectome structural data demonstrating single-neuron resolution across both pathways (right).

b, Schematic illustration of the anatomical organization of TINs and TPNs in the OT.

c, Two distinct subtypes of TPNs exhibiting selective connectivity with either the nucleus of the nMLF or M-cells. Schematic organization of TPN-E subtype exclusively connecting to M-cells (left) and TPN-O subtype showing specific connectivity with nMLF (right).

d, Lamina-specific retinotectal projections establish six distinct Np strata through retinal RGCs. Stratification progresses from superficial to deep layers: stratum opticum (SO), superficial fibrosum et griseum superficiale sublayers 1-2 (SFGS1-2), SFGS3-4, SFGS5-6, stratum griseum centrale (SGC), and stratum album centrale/stratum periventriculare (SAC/SPV), with individual RGCs projecting exclusively to single laminae.

e, Topographic organization of retinotectal projections demonstrating spatial inversion along four axes. Anterior-posterior axis: RGCs in the anterior retina projected to posterior Np regions, while posterior retinal RGCs innervated anterior Np regions; Dorsoventral axis: Dorsal retinal RGCs targeted medial Np domains, whereas ventral retinal RGCs terminated in lateral Np domains.

f, Schematic of visuomotor transformation circuitry. Combinatorial integration of the six-layered stratification (from panel d) and four-column topographic organization (from panel e) within the Np generated 24 functional microdomains. TPN subtypes TPN-E and TPN-O received processed visual signals from distinct Np microdomains, subsequently relaying information to M-cells and the nMLF nucleus, respectively, eliciting escape and orienting behavioral responses.

g,h Dendritic arborization heatmaps of TPN subtypes TPN-E (g) and TPN-O (h) within the neuropil (Np). Dendritic distribution was quantified by normalizing total branch length per neuronal cluster to maximum projection density (0-1 scale).

i,j Selective connectivity patterns between RGC subtypes and TPNs. (i) SGC/SAC-P RGCs exhibited preferential connectivity with TPN-E neurons, whereas SO-A RGCs showed stronger anatomical coupling with TPN-O neurons. (j) Coronal view orthogonal to the laminar axis demonstrated enhanced synaptic appositions between SGC/SAC-P RGC terminals and TPN-E dendrites, and between SO-A RGC terminals and TPN-O dendrites, within their respective Np microdomains.

k,l Connection specificity across neuronal populations. **(**k) SGC/SAC-P RGCs exhibited selective connectivity with TPN-E (connection probability: 7.9%), while SO-A RGCs showed preferential coupling with TPN-O (5.6%). Residual RGCs projecting to other Np microdomains demonstrated no significant connectivity bias. (l) TPN-E neurons formed high-probability synapses with M-cells

(45.7%) but not with the nucleus of the nMLF (0.0%), whereas TPN-O neurons preferentially

connected to nMLF (25.4%) over M-cells (0.0%, $p < 0.001$, χ^2 test).

Fig. 2. Computational dissection and morphotype-specific manipulation of visuomotor

transformation in the TIN reservoir network.

a, Schematic of selective retinal ganglion cell (RGC) activation via 100-ms intracellular current

injection (25 mV): SO-A RGCs (top) and SGC/SAC-P RGCs (bottom).

b,c Activation of SO-A RGCs evoked robust firing in TPN-O neurons (b), whereas SGC/SAC-P

RGC stimulation selectively activated TPN-E without eliciting TPN-O responses (c).

d, Mesoscale connectome-based simulations predicting premotor nucleus engagement probability. SGC/SAC-P RGC activation triggered M-cell responses with 99% probability, while SO-A RGC stimulation activated nMLF in 90% of iterations (trials=100)**.**

e, Experimental setup for visual stimulation (projector) and simultaneous two-photon calcium imaging in head-fixed zebrafish.

f, Laminar-specific calcium dynamics in OT Np evoked by three visual stimuli: Small Moving Dot (SMD), Big Moving Dot (BMD), and Looming threat**.**

g, Visuomotor transformation network architecture: Visual inputs processed through E-TIN/I-TIN reservoir with readouts via TPN-E (escape) and TPN-O (orienting) nodes.

h, Firing patterns of TPN-E and TPN-O under SMD (top) /Looming (bottom) stimuli with intact (left) vs. TIN-depleted (right) reservoir (normalized spike rates).

i, TIN-mediated behavioral accuracy enhancement. Correct responses: nMLF activation without M-cell firing during SMD (left); M-cell responses without high-frequency nMLF activity during Looming (right) (Accuracy: 96.7% $\pm$ 1.8% with TINs vs. 8.3% $\pm$ 1.2% without; $p < 0.001$).

j, *In silico* ablation paradigm targeting morphologically defined TIN subtypes in the reservoir network.

k, Top eight TIN subtypes whose ablation maximally impaired visuomotor accuracy. Nomenclature: [E/I][layers][stratification] (e.g., I_SFGS1-2_SFGS5-6: GABAergic bsTINs restricted to SFGS1-2 and SFGS5-6 layers; ns: non-stratified).

Fig. 3. Neural mechanisms of noise robustness in the visuomotor transformation reservoir

a, Freely swimming assay: visual stimulation with synchronized infrared behavioral recording in zebrafish.

b, Noise-dependent retinal encoding: Top: Looming stimuli with increasing noise density (ρ). Bottom: Corresponding calcium responses of SGC-layer RGCs in OT Np showing signal-to-noise ratio (SNR) degradation (gray lines: SEM, n=5 fish).

c, Quantified noise susceptibility: Inverse correlation between noise density (ρ) and SNR of SGC-RGC responses.

d, TIN reservoir robustness test: Escape probability estimation via M-cell activation to noisy looming stimuli.

e, Noise resilience of behavior.

f, Critical TIN subtypes for robustness: Top 8 morphotypes sustaining escape probability under

noise ($\rho=0.1$).

g, Chaotic signal processing paradigm: Lorenz attractor inputs with additive Gaussian noise

processed through TIN reservoir network.

h, Noise filtering capacity: Denoised Lorenz outputs (red) reconstructed from noisy inputs (blue)

at SD from 0.01 to 1.0.

i, Comparative denoising performance: Mean squared error (MSE) of reconstructed Lorenz signals

across network architectures (TIN reservoir vs shuffled TIN, RNN, LSTM, GRU). TIN achieved

minimal MSE.

j, 3D morphological reconstruction: Lorenz attractor morphology recovery under SD=0.1 noise.

TIN reservoir (red) outperformed artificial networks in trajectory preservation.

Fig. 4. 5-HT systems endow the TIN reservoir with flexibility for visual processing.

a, Spatial distribution of 5-HT across 24 OT Np microdomains.

b,c, Serotonergic neurons are divided into two types based on innervated in neuropil: SO-Type axons concentrated in superficial layers (b), while SGC-Type axons dominated deeper strata (c) (z-score normalized axonal density).

d, Light-response dichotomy in 5-HT subtypes: SO-Type 5-HTergic neurons exhibited sustained ON responses (tonic firing during illumination), whereas SGC-Type neurons showed transient ON-OFF responses (phasic bursts at light transitions).

e, In vitro modulation assay: Whole-cell patch-clamp recording during 5-HT bath application (10 μM, 5 min) in TNs.

f,g (f) Spontaneous firing increased post-5-HT and (g) Proposed mechanism: 5-HT lowers

activation thresholds of layer-specific TINs (schematic).

h,i Pathway-specific gain control: Under BMD stimuli, (h) control: balanced TPN-E/O firing; (i)

SO-Type 5-HT photo-stimulation biased responses to TPN-O (orienting), while SGC-Type 5-HT

enhanced TPN-E (escape).

j, Optogenetic lesion of SGC-Type 5-HT neurons identified by photo-response screening.

k, SGC-Type 5-HT lesions reduced looming-triggered escape probability.

Extended Data Figures

Extended Data Fig. 1. Multidimensional characterization of TNs.

a,b Excitatory-inhibitory (E-I) anatomical organization of (a) TPN and (b) TIN subtypes.

Including 348 excitatory TPNs (E-TPNs), 57 inhibitory TPNs (I-TPNs), 180 excitatory TINs (E-

TINs), and 845 inhibitory TINs (I-TINs) from zebrafish mesoscopic connectome.

c, Axo-dendritic morphological differentiation of TINs and TPNs. Synaptic release sites were

labeled by crossing with *Tg(UAS:sypb-APEX2-TagRFP)* zebrafish, where presynaptic

compartments (TagRFP+ puncta) defined axonal territories, and dendritic domains were identified

by the absence of TagRFP signal.

d, Schematic of neuronal connectivity quantification. Putative connections were defined by

axodendritic appositions within ≤2 voxel distances (3 um isotropic resolution). Connection

weights (edge thickness) were proportionally scaled to the number of contiguous voxel appositions

between axonal terminals and dendritic arbors.

e,f Excitatory-inhibitory neurite distribution in (c) TPNs and (d) TINs. Heatmap of normalized total neurite length across OT Np microdomains. All neurite lengths were normalized to the maximum value per subtype (0–1 scale).

g, Excitatory projection neurons (E-SPN) and inhibitory superficial interneurons (I-SIN) with soma located in the neuropil.

h, Laminar classification of OT neurons by stratification and projection. Upper: Extratectal projection neurons - monostratified TPN, msTPNs; bistratified TPN, bsTPN; tristratified TPN, tsTPN; non-stratified TPN, nsTPNs; excitatory superficial projection neuron, E-SPN. Lower: tectal interneurons with stratification - monostratified TIN, msTIN; bistratified TIN, bsTIN; tristratified TIN, tsTIN; non-stratified TINs nsTIN; inhibitory superficial interneuron, I-SIN.

i,j, Pie charts quantifying morphological diversity of (g) excitatory and (h) inhibitory TNs. Excitatory subtypes: msE-TIN (4.93%), bsE-TIN (16.14%), tsE-TIN (3.59%), nsE-TIN (7.77%), msE-TPN (9.87%), bsE-TPN (29.00%), tsE-TPN (23.02%), nsE-TPN (0.45%), E-SPN (3.29%), E-other types (1.94%) (n=561 cells). Inhibitory subtypes: msI-TIN (12.49%), bsI-TIN (38.09%), tsI-TIN (22.16%), nsI-TIN (15.19%), I-TPN (5.93%), I-SIN (1.66%), I-other types (4.47%) (n=961 cells).

Extended Data Fig. 2. Topographic projections of RGCs.

a, Schematic diagram illustrating the workflow for acquiring single-neuron resolution structural data in the zebrafish mesoscopic connectome (Details see Methods).

b,c, Axonal projection targets of RGCs. (b) OT-dominant projections: 99.35% of RGCs innervated contralateral OT Np microdomains. (c) Non-canonical hypothalamic projections: A small subset of RGCs (0.65%) exhibited contralateral projections to the intermediate hypothalamus (iHyp).

d-f, Retinotectal axonal arborization patterns. (d) Delineated AFs in the OT Np (upper) and two distinct axonal trajectories - RGC bundles traversing AF7 to target the SO, and those passing through AF9 projecting to the SAC/SPV (lower). (e) Coronal view of RGC laminar-specific projection architecture within OT Np. (f) Schematic model illustrating retinotopic-laminar convergence principles.

g, Population-wide RGC projection density heatmap across OT microdomains. Axonal arborization patterns were visualized by normalizing total axonal length projections to maximum intensity (0–1 scale).

887   **h,** Quantitative mapping of AF-specific projection distributions. Percentages of RGCs innervating

888   each AF that projected to specific OT Np microdomains. For every AF, the cumulative projection

889   weights along the white dashed lines to six Np layers (stratification axis) and four spatial columns

890   (topography axis) summed to unity.

Extended Data Fig. 3. Multidimensional characterization of TNs.

a,b Excitatory-inhibitory (E-I) anatomical organization of (a) TIN and (b) TPN subtypes.

c,d Excitatory-inhibitory neurite distribution in (c) TINs and (d) TPNs. Heatmap of normalized total neurite length across OT Np microdomains. All neurite lengths were normalized to the maximum value per subtype (0–1 scale).

e, Morphological segregation of excitatory projection neurons (E-SPN) and inhibitory superficial interneurons (I-SIN).

f, Laminar classification of OT neurons by stratification and projection. Upper: tectal interneurons with stratification - monostratified TIN, msTIN; bistratified TIN, bsTIN; tristratified TIN, tsTIN; non-stratified TINs nsTIN; inhibitory superficial interneuron, I-SIN. Lower: Extratectal projection neurons - monostratified TPN, msTPNs; bistratified TPN, bsTPN; tristratified TPN, tsTPN; non-stratified TPN, nsTPNs; excitatory superficial projection neuron, E-SPN.

g,h, Pie charts quantifying morphological diversity of (g) excitatory and (h) inhibitory TNs. Excitatory subtypes: msE-TIN (4.93%), bsE-TIN (16.14%), tsE-TIN (3.59%), nsE-TIN (7.77%), msE-TPN (9.87%), bsE-TPN (29.00%), tsE-TPN (23.02%), nsE-TPN (0.45%), E-SPN (3.29%),

E-other types (1.94%) (n=561 cells). Inhibitory subtypes: msI-TIN (12.49%), bsI-TIN (38.09%), tsI-TIN (22.16%), nsI-TIN (15.19%), I-TPN (5.93%), I-SIN (1.66%), I-other types (4.47%) (n=961 cells).

i, Axo-dendritic morphological differentiation of TINs and TPNs. Synaptic release sites were labeled by crossing with *Tg(UAS:sypb-APEX2-TagRFP)* zebrafish, where presynaptic compartments (TagRFP$^+$ puncta) defined axonal territories, and dendritic domains were identified by the absence of TagRFP signal.

j, Schematic of neuronal connectivity quantification. Putative connections were defined by axodendritic appositions within ≤2 voxel distances (3 um isotropic resolution). Connection weights (edge thickness) were proportionally scaled to the number of contiguous voxel appositions between axonal terminals and dendritic arbors.

Supplementary Table 1. Parameter ranges for LIF neuron models of key cell types.

	RGCs	All TNs	M-cell	nMLF neurons
τ_m (membrane time constant, ms)	10-20	15–30	15–30	15–25
V_{rest} (resting potential, mV)	-65	-65	-65 to -70	-65
V_{th} (threshold, mV)	-50 to -55	-50 to -55	-50	-50 to -55
V_{reset} (resetting potential, mV)	-65 to -70	-65 to -70	-65 to -70	-65 to -70
R_m (membrane resistance, MΩ)	150–300	150–250	100	150–250
τ_{ref} (refractory period, ms)	2–3	2–5	2–3	2–5