Structural basis of RNA-guided transcription by a dCas12f-σE-RNAP complex

Renjian Xiao[1,5], Florian T. Hoffmann[3,5], Dan Xie[1], Tanner Wiegand[3,4], Adriana I. Palmieri[3], Samuel H. Sternberg[3,4†], Leifu Chang[1,2†]

[1]Department of Biological Sciences, Purdue University, 915 W. State Street, West Lafayette, IN 47907, USA. [2]Purdue Institute for Cancer Research, Purdue University, 915 W. State Street, West Lafayette, IN 47907, USA. [3]Department of Biochemistry and Molecular Biophysics, Columbia University, New York, NY, USA. [4]Howard Hughes Medical Institute, Columbia University, New York, NY, USA. [5]These authors contributed equally.

[†]Corresponding authors

ABSTRACT

RNA-guided proteins have emerged as critical transcriptional regulators in both natural and engineered biological systems by modulating RNA polymerase (RNAP) and its associated factors[1-5]. In bacteria, diverse clades of repurposed TnpB and CRISPR-associated proteins repress gene expression by blocking transcription initiation or elongation, enabling non-canonical modes of regulatory control and adaptive immunity[1,6,7]. Intriguingly, a distinct class of nuclease-dead Cas12f homologs (dCas12f) instead activates gene expression through its association with unique extracytoplasmic function sigma factors (σ^E)[8], though the molecular basis has remained elusive. Here we reveal a novel mode of RNA-guided transcription initiation by determining cryo-electron microscopy structures of the dCas12f-σ^E system from *Flagellimonas taeanensis*. We captured multiple conformational and compositional states, including the DNA-bound dCas12f-σ^E-RNAP holoenzyme complex, revealing how RNA-guided DNA binding leads to σ^E-RNAP recruitment and nascent mRNA synthesis at a precisely defined distance downstream of the R-loop. Rather than following the classical paradigm of σ^E-dependent promoter recognition, these studies show that recognition of the -35 element is largely supplanted by CRISPR-Cas targeting, while the melted -10 element is stabilized through unusual stacking interactions rather than insertion into the typical recognition pocket. Collectively, this work provides high-resolution insights into an unexpected mechanism of RNA-guided transcription, expanding our understanding of bacterial gene regulation and opening new avenues for programmable transcriptional control.

INTRODUCTION

Bacterial transcription is catalyzed by a conserved RNA polymerase (RNAP) core enzyme comprising four distinct subunits: α, β, β', and ω. RNAP exhibits only non-specific interactions with DNA[9,10], whereas sequence-specific binding of promoter motifs during transcription initiation relies critically on a dedicated sigma (σ) factor that comprises a key part of the σ-RNAP holoenzyme complex[11,12]. Bacteria typically encode one RNAP core enzyme but multiple σ factors, each of which recognizes a distinct promoter sequence, thus allowing adaptation to diverse cellular conditions[13,14]. The primary Group 1 σ factor in *E. coli*, σ^{70}, is responsible for transcribing essential housekeeping genes and contains four conserved domains ($\sigma1$–$\sigma4$) that bind the –35 and –10 elements in standard σ^{70} promoters[15]. Group 2 and 3 σ factors play more specialized regulatory roles, including controlling genes involved in stationary phase, flagellar biosynthesis, and heat shock, and share many of the same conserved regions as σ^{70}. Group 4 σ factors, also known as extracytoplasmic function (ECF) σ^E, are structurally more compact and retain only the σ_2 and σ_4 domains connected by a $\sigma_{2/4}$ linker[16]. Intriguingly, whereas *E. coli* encodes only a single homolog, σ^E factors are the largest and most diverse group in other bacterial taxa[13,17], where they canonically respond to diverse environmental cues via interactions with their cognate anti-sigma factors that localize to the cellular membrane[14,15,18-20].

Recent genomic analyses uncovered a subset of ECF σ^E factors encoded adjacent to atypical Cas12f-like homologs that lack the expected association with CRISPR arrays[21]. These unusual σ^E proteins feature a C-terminal extension resembling a helix-turn-helix (HTH) domain predicted to interact with Cas12f, and the presence of conserved Cas12f mutations in the catalytic RuvC domain, as well as the absence of C-terminal regions critical for DNA cleavage, further suggesting a novel function beyond adaptive immunity[21]. In particular, the strong co-occurrence between σ^E and nuclease-dead Cas12f (dCas12f) implied the possibility that guide RNA (gRNA) molecules might be involved in directing a new mode of transcriptional activation through direct interactions with RNAP. Alongside recently described systems that employ repurposed Cas9, Cas12, or TnpB-like homologs for transcriptional repression[1,6,7,22-24], a natural mechanism of RNA-guided transcription initiation would expand the known roles of programmable gene regulation while also complementing existing technologies for CRISPR interference (CRISPRi) and CRISPR activation (CRISPRa) applications[2-5].

Here, we present the biochemical and structural basis of RNA-guided transcription initiation by the dCas12f-σ^E, whose cellular activities are detailed in a companion article[8]. Focusing on homologs from *Flagellimonas taeanensis* that regulate SusC/SusD-family outer membrane proteins[8] **(Fig. 1a,b)**, we resolved multiple cryo-EM structures including dCas12f-gRNA, DNA-bound dCas12f-gRNA, and the DNA-bound dCas12f-σ^E-RNAP holoenzyme complex. Our structures elegantly reveal how recognition of a simple 5'-G target-adjacent motif (TAM) leads to R-loop formation, σ^E-RNAP recruitment, and a tightly defined transcription start site (TSS) ~46-bp downstream, without obligate recognition of typical promoter elements. These findings provide new insights into diverse σ factor function and inform future strategies for synthetic, promoter-independent gene activation tools.

RESULTS

Specificity and biochemical basis of dCas12f-σ^E coupling

We focused on a *F. taeanensis* (*Fta*) strain that encodes one dCas12f-σ^E system alongside 26 additional σ^E homologs (*rpoE* gene) and 3 σ^70 homologs **(Fig. 1a, Extended Data Fig. 1)**. Based on their phylogenetic distribution, we selected nine representatives for functional testing in *E. coli*, including the dCas12f-associated σ^E, and developed a fluorescence-based reporter assay to monitor activity[8], in which the native gRNA-matching target site is cloned upstream of RFP **(Fig. 1b)**. The resulting data revealed that only the cognate, dCas12f-associated σ^E enabled RNA-guided transcription above background levels, whereas all other tested σ factors showed baseline activity comparable to the no-σ control **(Fig. 1c)**. These data are consistent with a functional coupling that is exclusive to dCas12f and its genomically proximal *rpoE* gene product.

We next separately purified the *Fta* dCas12f-gRNA binary complex and the *Fta* RNAP core enzyme (α, β, β', ω) using a heterologous *E. coli* expression system. Analysis by SDS-PAGE revealed an absence of contaminating *E. coli* RNAP components, suggesting that the purified products predominantly corresponded to homogeneous *Fta* RNAP **(Fig. 1d)**. While *Fta* σ^E alone proved recalcitrant to purification, co-expression with RNAP yielded a stable σ^E-RNAP complex **(Fig. 1d)**. Similarly, while we were unable to isolate a homogenous complex from cells co-expressing σ^E with dCas12f-gRNA **(Extended Data Fig. 2a–c)**, size-exclusion chromatography indicated that a stable dCas12f-σ^E-RNAP supercomplex specifically formed in the presence of target DNA **(Extended Data Fig. 2d,e)**. This latter result suggests that substrate DNA binding and R-loop formation triggers RNAP holoenzyme recruitment.

Structure of dCas12f-gRNA binary complex

To elucidate the molecular basis of RNA-guided transcription, we first determined a 3.28 Å cryo-EM structure of the dCas12f-gRNA complex **(Fig. 1e, Extended Data Fig. 3, and Extended Data Table 1)**. As with canonical Cas12f effectors such as *Un*Cas12f[25,26] and *As*Cas12f[27,28], *Fta* dCas12f assembles as an asymmetric dimer with a single gRNA **(Fig. 1e)**.

dCas12f consists of an N-terminal region (1–214), further divided into the REC (recognition) and WED (wedge) domains, and a C-terminal RuvC domain (223–365) connected by a linker **(Fig. 1f)**. The REC domain forms the dimerization interface, and consistent with previous nomenclature[25-28], we designate the subunit that extensively interacts with the gRNA and binds the TAM as dCas12f.1, and the other as Cas12f.2. Domains within both subunits share identical folds, but the N- and C-terminal regions adopt different orientations due to linker variations. A key motif within the RuvC domain is the lid motif, a region between the β4 and β5 strands of the RuvC domain spanning residues 325–346 **(Fig. 1g)**, which undergoes major conformational changes upon target DNA recognition, as discussed below.

The gRNA is naturally encoded downstream of *dcas12f* as a single-guide transcript[8], in contrast to the natural dual-guide RNA comprising crRNA and tracrRNA used by canonical CRISPR-Cas12f nuclease systems[29,30]. The gRNA consists of a core structure with four based-paired regions (designated P1–P4) and a 14-nt 3' guide region, which strikingly adopts an unexpected stem-loop structure P5 **(Fig.**

Figure 1 | Biochemical characterization of the dCas12f-σ^E system and structure of dCas12f-gRNA. a, Genomic organization of the dCas12f-σ^E operon in *F. taeanensis* (left), and the gRNA-matching DNA target site upstream of *susC/susD* membrane transport genes (right). **b,** Schematic showing base-pairing between the gRNA and target DNA, which includes native mismatches at positions 11–12. The downstream transcription start site (TSS) is indicated. **c,** OD-normalized RFP fluorescence from cellular transcriptional reporter assays using the indicated *Fta* σ^E homologs. dCas12f-gRNA complexes were presented in all conditions, and contained a targeting (T) or non-targeting (NT) guide. Data are shown as mean ± s.d. for *n* = 3 biologically independent samples. **d,** SDS-PAGE and urea-PAGE analysis of purified dCas12f-gRNA, RNAP, and σ^E-RNAP complexes. **e,** Cryo-EM density map of the dCas12f-gRNA complex at 3.28 Å. **f,** Domain organization of dCas12f.1 and dCas12f.2, showing the REC, WED, RuvC, and Lid motif domains. **g,** Structures of the Lid motif in 'Loop' (left) and 'Helical' (right) conformations. **h,** Cartoon representation of the experimental gRNA structure. **i,** Secondary structure diagram of the gRNA, highlighting key structural elements. Stem-loops are labeled P1–P5.

1h,i, Extended Data Fig. 3g). This differs markedly from other Cas12 nucleases, where the seed sequence typically maintains a pre-ordered conformation to facilitate target DNA hybridization, with the rest of the guide region remaining flexible[26,31-34]. Conventional CRISPR-Cas systems rely on diversified guides that

comprise unique, foreign DNA-derived spacers within the CRISPR array, whereas the dCas12f-σ^E system has evolved an inflexible guide region that targets a specific, conserved genomic site upstream of *susCD* for gene regulation[8]. This reduced variability may thus favor the formation of the P5 stem-loop structure, which contains base pairs between residues 77–79 and 87–89 **(Fig. 1h,i)**. This region must unwind to enable RNA-DNA heteroduplex formation, likely imposing an energetic barrier that enhances target specificity.

Structure of the DNA-bound dCas12f-gRNA complex

To investigate the mechanism of RNA-guided DNA binding, we next investigated the ternary complex comprising dCas12f, gRNA, and target DNA **(Fig. 2a,b, Extended Data Fig. 4, and Extended Data Table 1)**. Overall, the ternary complex resembles *As*Cas12f-gRNA-DNA and *Un*Cas12f-gRNA-DNA complexes[25-27], but with minimized protein and gRNA constituents **(Extended Data Fig. 5)**. The TAM duplex binds within a groove formed by the REC and WED domains of dCas12f.1, which make extensive contacts with the phosphate backbone of the non-target strand **(Fig. 2c,d)**, and alanine substitutions of key residues involved in this interaction (Y89 and R90) reduce RNA-guided transcription by roughly 50% **(Fig. 2e)**. Strand separation is facilitated by H110 and M85, which stack against a G-C pair at position −1, and recognition of this minimal 5'-G TAM is mediated by a hydrogen bond between the S106 main-chain carbonyl group and the C(−1) base of the target strand **(Fig. 2f)**. Additionally, Q144 forms a hydrogen bond with A(−4) of the non-target strand **(Fig. 2c,d)**. While the TAM binding site in Cas12f.2 is accessible, no DNA is bound to this site, consistent with the paradigm in which TAM (or PAM) interactions are transient and only stabilized upon RNA-DNA hybrid formation[35] **(Fig. 2a,b)**.

Hybridization of the guide RNA and target DNA occurs subsequent to TAM recognition, and 3D classification of the ternary complex revealed two distinct conformational states. The majority of particles (~88%) adopt a partial R-loop conformation, with only a minor subset (~12%) favoring a complete R-loop **(Fig. 2g,h, Extended Data Fig. 4f)**. The partial R-loop structure features a 7-bp RNA-DNA heteroduplex, and although RNA and DNA densities at positions 8 and 9 are observed, they deviate from A-form duplex geometry and fail to form base pairs **(Fig. 2g,i, Extended Data Fig. 4g–j)**. In the full R-loop structure, base pairs at positions 7–9 reorganize to adopt canonical A-form structure, and the duplex extends to position 14, corroborating accompanying *in vivo* data from RIP-seq and transcriptional activity assays[8] **(Extended Data Fig. 4h,j)**. A twist in the target DNA forms a wobble base pair at position 10, potentially accommodating mismatches at positions 11 and 12 **(Extended Data Fig. 4j)**, and densities for DNA and RNA are undetectable beyond position 14.

The transition between the partial and full R-loop states involves coordinated conformational changes in the RuvC domain, particularly in the lid motif. In the partial R-loop state, where we refer to the motif as Lid^{Loop}, only the C-terminal portion of the lid motif (residues 337–346) adopts a defined structure, while the N-terminal region (residues 325–336) is disordered **(Fig. 2g)**. Two consecutive tyrosines (Y344 and Y345) are positioned beneath the target DNA strand and sterically block position 8 from hybridizing with the gRNA in an A-form configuration. Upon RNA-DNA heteroduplex extension beyond position 7

in the full R-loop state, both tyrosines rearrange into an extended α-helical structure formed by residues 347–358. In the full R-loop state, the entire lid motif (referred to as Lid^Helical) becomes ordered, forming stable helical structures that interact with the target DNA **(Fig. 2h,i)**. These structures therefore reveal a likely mechanism whereby the lid motif functions as a sensor for R-loop formation. In canonical Cas12f nucleases, the lid motif guides the DNA substrate into the RuvC catalytic center, positioning it for cleavage[26] **(Fig. 2j)**, whereas for dCas12f-σ^E systems, we hypothesized that it would instead play a critical role in σ^E recruitment **(Fig. 2k)**.

Figure 2 | Structure of the dCas12f-gRNA-target DNA complex. **a,** Cryo-EM density map of dCas12f-gRNA-DNA in a partial R-loop state. **b,** Cryo-EM density map of dCas12f-gRNA-DNA in a full R-loop state. **c,** Interaction between the TAM duplex and dCas12f.1. **d,** Schematic of interactions between the TAM duplex and dCas12f.1. **e,** OD-normalized RFP fluorescence from cellular transcriptional reporter assays using the indicated dCas12f mutants that perturb TAM duplex recognition. Data are shown as mean ± s.d. for *n* = 3 biologically independent samples. **f,** Recognition of the 5'-G TAM, highlighting hydrogen bonding between the base of C(−1) and the backbone carbonyl oxygen of S106. **g,** Lid motif conformation in the partial R-loop state. **h,** Lid motif conformation in the full R-loop state. Insets in **g** and **h** show focused views of key residues in the Lid motif undergoing conformational rearrangement, along with the corresponding cryo-EM density. **i,** Structural superimposition of RNA-DNA duplex in partial and full R-loop states. **j,** Lid motif conformation in *Un*Cas12f-gRNA-DNA complex for comparison. **k,** Lid motif conformation in the dCas12f-gRNA-DNA-σ^E-RNAP complex.

Overall structure of the DNA-bound dCas12f-σ^E-RNAP holoenzyme

To directly resolve the mechanism of σ^E and RNAP recruitment during transcription initiation, we next determined the structure of the fully assembled dCas12f-σ^E-RNAP nucleoprotein complex comprising dCas12f-gRNA, a 99-bp target DNA, σ^E, and *Fta* RNAP. The dCas12f-σ^E-RNAP complex was refined to an average resolution of 3.30 Å from 11,161 particles, and 3D classification also allowed us to determine the apo RNAP structure to 2.95 Å from 171,804 particles **(Extended Data Fig. 6, Supplementary Table. 1)**. Optimized DNA constructs incorporating mismatches to favor DNA bubble formation within the RNAP core improved the overall resolution to 2.88 Å **(Extended Data Fig. 7, Extended Data Table 1)**, and focused refinement using a mask around the dCas12f-gRNA region yielded a local resolution of 3.34 Å **(Extended Data Fig. 7e)**. Overall, the final map showed excellent density, allowing model building for nearly all protein subunits of the complex **(Extended Data Fig. 8)**. However, the α subunit C-terminal domains, known to interact with the UP element of canonical promoter DNA[36,37], are not resolved in the map, indicating their inherent flexibility. and the lineage-specific insertions in the β and β' subunits, which likely interact with transcriptional factors[12,38,39], are visible but at lower resolution **(Extended Data Fig. 8a,b)**.

The DNA substrate can be divided into six regions: the TAM duplex, R-loop, transition region, bridge DNA, transcription bubble, and downstream DNA **(Fig. 3a)**. The 6-bp TAM duplex and 14-bp RNA-DNA heteroduplex within the R-loop are well-resolved, although density for the non-target DNA strand is largely absent **(Extended Data Fig. 8j, 9a,b)**. The bridge DNA — which spans between dCas12f and the RNAP core — adopts a B-form conformation with well-resolved major and minor grooves **(Fig. 3b–d)**, but weak density between the R-loop and bridge DNA suggests structural flexibility in this transition region, preventing precise sequence tracing **(Extended Data Fig. 8j)**. Within the transcription bubble itself, we modeled a 13-nt sequence for the template strand, whereas only the first 3 nt are visible for the non-template strand **(Extended Data Fig. 8j)**; this observation aligns with the typical transcription bubble length of 12–14 bp observed in homologous bacterial RNAP structures[40-45]. Finally, the downstream DNA is modeled as a 17-bp B-form duplex.

To provide a mechanistic framework for transcription initiation, we modeled a 4-nt transition region (positions 15–18) based on structural constraints and cryo-EM densities from the unsharpened map **(Extended Data Fig. 8j)**. This allowed confident tracing of the template DNA strand sequence up to position 67 in the downstream DNA, yielding a model that aligns well with the observed density and is biologically plausible, as it places position 46, the most frequently observed TSS from RNA-sequencing data[8], within the RNAP catalytic center. We note, however, that minor shifts in conformation may occur given the inherent flexibility of DNA[46], and this could readily explain the observation of context-dependent TSSs located 45–48 bp downstream of the TAM site[8]. Collectively, these findings suggest that our structure represents a biologically relevant state of RNA-guided transcription driven by the dCas12f-σ^E system.

The final structural model reveals a pronounced bipartite architecture, consisting of a well-ordered RNAP core region bound to the σ_2 domain of σ^E, and a smaller dCas12f-gRNA region that engages with

the σ$_4$ and C-terminal domains (CTD) of σ^E **(Fig. 3a–e)**. These two regions are physically connected by three distinct structural elements: the 16-bp bridge DNA **(Fig. 3b–d)**, the elongated σ^E α-helix that links its RNAP-bound σ$_2$ domain to its dCas12f-bound σ$_4$ domain **(Fig. 3b–d)**, and electrostatic interactions between the dCas12f.1 RuvC domain and the RNAP ω subunit **(Fig. 3f)**.

Structure of the *Fta* RNAP core complex

Figure 3 | Structures of the DNA-bound dCas12f-σ^E-RNAP complex and apo RNAP. a, Schematic showing the target DNA and gRNA in the DNA-bound dCas12f-σ^E-RNAP structure. The target DNA features two DNA bubbles, one mediated by the dCas12f-gRNA R-loop and the other mediated by the RNAP transcription bubble; the TSS is located at position 46 of the template strand. **b,** Two representative 2D class averages of cryo-EM images of the DNA-bound dCas12f-σ^E-RNAP complex. **c,** Cryo-EM density map of the DNA-bound dCas12f-σ^E-RNAP complex. **d,** Cartoon representation of the atomic model. **e,** Schematic of the DNA-bound dCas12f-σ^E-RNAP structure. **f,** Contact between dCas12f.1 and the RNAP ω subunit, showing the cryo-EM map displayed as a surface representation (top) and the surface potential of dCas12f.1 and the ω subunit (bottom). **g,** Cryo-EM map of apo RNAP. **h,** Atomic model of apo RNAP.

RNAP and σ factors canonically function together for faithful promoter recognition[47-49], but in dCas12f-σ[E] systems, RNAP also promotes key interactions between DNA-bound dCas12f and σ[E]. Overall, the core subunits closely resemble other bacterial RNAP structures, such as those from *E. coli* and *M. tuberculosis*[40,50] **(Extended Data Fig. 10)**, with the characteristic 'crab-claw' shape formed by the large β and β' subunits **(Fig. 4a–e)**. These two subunits form a structural platform that is stabilized by peripheral α subunits (α1 and α2) and the ω subunit, while also coordinating σ[E] recruitment.

The β and β' subunits define key structural features critical for transcription, including the clamp, the DNA-binding cleft, the entry funnel for incoming nucleoside triphosphates (NTPs), and the transcription bubble stabilization in coordination with σ[E]. The clamp domain, formed by the β-clamp and β'-clamp regions, undergoes significant conformational rearrangement upon σ[E] and target DNA binding, which stabilizes the downstream DNA together with the β'-jaw **(Figs. 4f and 3g,h)**. The β' subunit contains a

Figure 4 | Structure of *Fta* RNAP core subunits. a, Domain organization of the RNAP core subunits. **b,** Structure of the β' subunit with each domain color-coded. **c,** Structure of β' in the context of σ[E] and DNA. **d,** Structure of the β subunits with each domain color-coded. **e,** Structure of β in the context of σ[E] and DNA. **f,** Structure of the Clamp domain alone (left) and in the context of σ[E] and DNA (right); the conformational change between both states is indicated. **g,** Structure of the 'Cleft' formed by the β-cleft, β'-cleft, and β-protrusion. **h,** View of the 'Funnel' structure formed by the β'-pore, β'-funnel, and β'-cleft domains.

characteristic coiled-coil motif (β'-CC) and zinc-finger domain (ZnF) within the clamp domain, both of which engage in critical interactions with σ^E **(Fig. 4f)**. The cleft formed by β and β' encloses the DNA bubble and harbors the catalytic center in the β' subunit **(Fig. 4g)**, which is responsible for RNA synthesis. Adjacent to the catalytic center, the positively charged bridge helix (BH) plays a crucial role in catalysis and RNA-DNA translocation[51]. The incoming NTPs access the active site through a 'funnel' structure that is predominantly formed by the β'-pore and β'-funnel domains **(Fig. 4c,h)**, highlighting the conserved mechanism of substrate entry and catalysis across bacteria, archaea, and eukaryotes (RNAP II) [40,50,52,53]. Additionally, the β-lobe and β-external domains, positioned near the downstream DNA, and the β-flap domain, proximal to the bridge DNA, are essential for complex stability **(Fig. 4e)**. Notably, the β flat tip helix (β-FTH) plays a crucial role in engaging the σ_4 domain of σ^E (ref. [54]), as detailed below.

While the α-NTD structure in *Fta* RNAP is highly similar to *E. coli* RNAP, the ω subunit exhibits notable differences, consistent with its chaperone-like role in RNAP assembly[55,56] **(Extended Data Fig. 10d)**. As the smallest RNAP core subunit, ω primarily interacts with the β subunit for proper β subunit folding[56-59]. Compared to *E. coli* ω, *Fta* ω contains a negatively charged insertion (residues 48–82) that adopts a helix-loop-helix structure, which appears conserved in genomes containing dCas12f-σ^Esystems **(Extended Data Fig. 10e,f)** and engages the basic RuvC domain of Cas12f.1 **(Fig. 3f)**. A DALI search identified a similar ω subunit insertion in *Porphyromonas gingivalis* RNAP (PDB: 8DKC) [60] **(Extended Data Fig. 10f)**, but dCas12f is notably absent from the genome. *P. gingivalis*, like *F. taeanensis*, belongs to the bacterial phylum *Bacteroidetes*, thus suggesting that the conserved structural insertion reflects a shared phylogenetic origin and was co-opted for dCas12f interactions.

Molecular basis of σ^E–dCas12f interactions

σ^E is recruited to target-bound dCas12f via both protein–protein and protein-DNA interactions **(Fig. 5a,b)** that provide critical insights into RNA-guided transcription. The σ^E CTD, which is unique to dCas12f-associated σ^E and absent in other σ^E paralogs[8], binds directly to the RuvC domain of dCas12f.1, complementing its partially truncated structure **(Fig. 5c)**. The interface centers on a hydrophobic core formed by σ^E residues F225 and F243, and hydrophobic residues in the dCas12f.1 RuvC domain. The interaction is further strengthened by hydrogen bonding interactions, including those mediated by H240 and R218 **(Fig. 5c)**. Deletion of residues 201–265 in the σ^E CTD, or alanine substitutions in this region (R218A and H240A/F243A), led to a striking reduction in transcriptional activity (~50%) compared to wild type **(Fig. 5d)**.

The σ_4 domain acts as a linchpin, contacting dCas12f, the target DNA, and the β-FTH **(Fig. 5b,e,f,g)**. Two hydrophobic residues, I184 and L137, form key contacts with the dCas12f.1 Lid^Helical motif **(Fig. 5f)**. Substitution of L137 with alanine or lysine significantly reduced transcription activity in cellular assays, and the L137A/I184A double mutation almost completely abolished transcriptional activity **(Fig. 5h)**. In addition to this key contact, the σ_4 domain engages with the target DNA and dCas12f.2 simultaneously via residues H144 and Y152 of the latter's lid motif **(Fig. 5g)**. The WT activity of H144A and Y152A mutants, however, suggests that this interaction is ancillary rather than critical. Notably, the σ_4 domain engages

with the target DNA through minor groove contacts **(Fig. 5b**, **Extended Data Fig. 9f)**, contrasting with its canonical role in recognizing −35 promoter elements via major groove interactions in typical σ factors[40,61]. One exception is the σ[I] (SigI) factor, which recognizes the −35 region through both the major groove and A-tract minor groove, likely enhancing binding stability via increased interface area[62]. Minor groove recognition follows a similar pattern, with a positively charged histidine inserting into the groove and positively charged residues interacting with the phosphate backbone, commonly found in protein-DNA interactions[63]. This altered DNA binding mode suggests that σ_4 primarily stabilizes the complex rather than conferring sequence specificity, allowing dCas12f-gRNA to provide target recognition function.

Molecular basis of RNAP-σ[E] interactions

Figure 5 | Interactions between dCas12f and σ[E]. a, Structure of σ[E] in cartoon representation, with the remainder of the DNA-bound dCas12f-RNAP complex shown in transparency, for clarity. The σ[E] domain organization is shown at the top. **b,** Overall view showing the interactions between σ[E] and other components, with magnified inset regions indicated with dotted lines. **c,** Interactions between the σ[E] CTD and dCas12f.1. **d,** OD-normalized RFP fluorescence from cellular transcriptional reporter assays using the indicated σ[E] CTD mutants. **e,** Interactions between the σ^E_4 domain and β-flip-tip-helix (β-FTH). **f,** Interactions between the σ^E_4 domain and dCas12f.1. **g,** The σ^E_4 domain is encircled by β-FTH, the lid motif of dCas12f.2, and DNA. **h,** OD-normalized RFP fluorescence from cellular transcriptional reporter assays using the indicated σ^E_4 mutants. Data in **d** and **h** are shown as mean ± s.d. for *n* = 3 biologically independent samples.

The σ$_2$ domain plays key roles in transcription initiation, particularly in DNA unwinding and strand separation at the −10 region[13]. In our structure, R76 packs against the last base-paired position (G-C pair at position −12 from the TSS), facilitating strand separation **(Fig. 6a,b, Extended Data Fig. 9c,d)**. Upon DNA melting, both strands are stabilized by distinct protein interactions to prevent re-annealing: the non-template strand is primarily stabilized by the σ$_2$ domain **(Fig. 6a,b)**, while the template strand is stabilized through interactions with both σ^E and the RNAP β-protrusion **(Extended Data Fig. 9c,d)**. Overall, multiple key residues contribute to maintaining a melted DNA state, though single mutations did not reduce transcriptional activity, suggesting that individual perturbations are tolerated under these experimental conditions **(Extended Data Fig. 9g)**.

The first 3–4 nucleotides of the non-template strand typically contribute to σ factor sequence specificity in the −10 region (i.e. the TATA motif in primary σ factors)[40-42,64-67]. In *E. coli* σ^E, a specificity loop between two helices forms a binding pocket that recognizes the flipped-out nucleotide[40] **(Fig. 6c)**. In our structure, however, no obvious binding pocket is present for recognizing a flipped nucleotide, and H65 instead packs against C(−10) and G(−9) **(Fig. 6b)**, likely stabilizing the melted bases and preventing

Figure 6 | Target DNA interactions and comparison with canonical σ^E factors. a, Interactions between *Fta* σ^E and the −10 region of DNA; the cryo-EM map is shown in mesh. **b,** Interactions between *Fta* σ^E and the −10 region of DNA, with the σ^E surface displayed. **c,** Interactions between the *Eco* σ^E and the −10 region of DNA, shown as in **b**, highlighting the pocket formed by the specificity loop. **d,** Interactions between *Fta* σ^E and the target DNA. **e,** Interactions between *Eco* σ^E and the target DNA. **f,** Superimposed structure comparing dCas12f-associated *Fta* σ^E (yellow) and non-dCas12f-associated *Eco* σ^E (brown).

reannealing without sequence specificity. Consistently, perturbation of the −10 region had minimal effect on RNA-guided transcription[8], suggesting that nucleobase-dependent interactions at this site are not essential.

The $\sigma_{2/4}$ linker extends into the catalytic cleft, functioning similarly to the primary $\sigma_{3.2}$ domain ($\sigma_{3/4}$ linker) of σ^{70} (ref. [64,68]) by pre-organizing the template strand for subsequent *de novo* RNA synthesis, as is also observed in homologous ECF sigma factors[40-42]. In our structure, Y105 from the linker packs against G(−4) of the template strand **(Extended Data Fig. 9e)**.

In comparing our *Fta* dCas12f-σ^E-RNAP structure with the *E. coli* σ^E-RNAP complex, notable differences emerge. Whereas the σ_4 domain typically recognizes the −35 region through major groove contacts[40,61], in our structure it instead engages with the minor groove **(Fig. 6d–f)**. Additionally, the bridge DNA (corresponding to the upstream DNA in the *E. coli* σ^E-RNAP complex) adopts a distinct angle by rotating away from the RNAP β'-ZnF **(Fig. 6d–f, Extended Data Fig. 10a-c)**, thus deviating from the typical configuration observed in standard transcription initiation complexes[42,43,65,68,69]. The bridge DNA region is notably A-T rich, with a continuous A-T sequence spanning positions 21–28. The A-T rich bridge may act as a flexible hinge, allowing the DNA to adopt a conformation favorable for interaction with the dCas12f-σ^E-RNAP complex.

DISCUSSION

Altogether, our results reveal an unprecedented mechanism of transcription initiation in dCas12f-σ^E systems, in which protein-DNA promoter recognition is replaced by RNA-DNA complementarity, enabling a flexible and programmable mode of defining transcription start sites (TSS). The dCas12f-gRNA complex binds upstream of the TSS at a position analogous to the −35 promoter region in canonical bacterial transcription[41,69,70], leading to subsequent recruitment of the σ^E-RNAP complex. The σ_4 domain, which is usually responsible for −35 sequence recognition, plays a distinct role by serving as a structural adaptor that bridges dCas12f and RNAP through critical engagement of the minor groove of the bridge DNA. Meanwhile, the σ_2 domain retains its conventional role in core RNAP interactions and DNA melting at the −10 region, though the sequence specificity enforced is substantially reduced. Together, these structural adaptations exemplify striking co-evolution between RNA-guided, nuclease-dead Cas12f proteins and extracytoplasmic function sigma factors, that enabled a novel mode of transcriptional regulation.

Our structural analysis identified three key interaction points that establish the dCas12f-σ^E-RNAP complex: a bridge DNA-mediated connection, direct interactions between dCas12f and σ^E_4 that are further stabilized through the minor groove DNA contacts, and direct dCas12f-RNAP core interactions through the ω subunit. This latter interaction may represent a key feature to accommodate RNA-guided transcription.

The σ^E C-terminal domain (CTD) extension is a unique feature of dCas12f-associated systems that distinguishes it from canonical σ factors **(Extended Data Fig. 1)**. It binds the truncated, nuclease-inactive RuvC domain of dCas12f, likely enhancing the affinity between the RNA-guided effector and the σ^E-RNAP complex. Given its positioning, we speculate that this critical σ^E-Cas12f.1 interaction functions

as an initial docking mechanism, facilitating the recruitment of σ^E-RNAP to the bound target DNA. This protein-protein interaction may provide a more efficient means of transcriptional activation compared to conventional sigma factor-mediated promoter search mechanisms[71-74]. Our data also highlight the conformational coupling between R-loop formation and transcription complex assembly. Specifically, conformational changes of the dCas12f lid motif function as a molecular switch that ensures transcription initiation occurs exclusively at targeted sites, likely enhances both specificity and regulatory control in cellular contexts.

In conclusion, our work establishes a structural framework for RNA-guided transcription, elucidating key interactions between an RNA-guided protein and the σ^E-RNAP holoenzyme complex. These insights provide new opportunities for developing programmable transcriptional regulators for precise gene expression control.

METHODS

Molecular constructs. The initial plasmid construct (pSL6726) that encodes *F. taeanensis* dCas12f, gRNA, and σ^E (*rpoE* gene) was synthesized by GenScript[8]. *dcas12f* and *rpoE* genes were individually subcloned into the bacterial expression vector pET28-MKH8SUMO (Addgene #79526). The His$_8$-tag in the *rpoE* vector was subsequently removed for co-expression with *Fta* RNAP. For co-expression with dCas12f, the gRNA was subcloned into a pUC19 vector. Genes encoding *Fta* RNAP subunits were also synthesized by GenScript and subsequently cloned into the pACYCDuet vector (*rpoA_rpoB* in MCS-1 with a His$_6$-tag on *rpoA*; *rpoC_rpoZ* in MCS-2).

Mutagenesis. Single amino acid mutations were introduced using the QuikChange site-directed mutagenesis method. Multiple amino acid mutations were generated by ligating inverse PCR-amplified backbone with mutation-bearing DNA oligonucleotides using the In-Fusion Cloning Kit (Clontech). All mutations were verified by Sanger sequencing.

Protein expression and purification. dCas12f and gRNA were co-expressed in Rosetta(DE3)pLysS cells (Novagen #70956), while σ^E and RNAP were co-expressed in BL21-AI cells (Thermofisher # C607003). Cells were cultured in Terrific Broth (TB) to an OD$_{600}$ of 0.6. Protein expression was induced with 0.3 mM IPTG, and for BL21-AI cells, 0.2% arabinose was added. Cultures were incubated overnight at 16 °C. Cells were harvested and resuspended in lysis buffer (25 mM Tris-HCl pH 7.6, 500 mM NaCl, 5% glycerol) supplemented with 1 mM PMSF and 5 mM β-mercaptoethanol, then lysed by sonication. The lysate was clarified by centrifugation, and the supernatant was applied to Ni-NTA resin. After extensive washing with lysis buffer containing 30 mM imidazole, target proteins were eluted with lysis buffer containing 250 mM imidazole.

The His-SUMO tag from dCas12f and the SUMO tag from σ^E were removed by overnight digestion with SUMO protease at 4 °C. The dCas12f-gRNA eluate was diluted in buffer containing 25 mM Tris-HCl pH 7.6, 125 mM NaCl, and 5% glycerol, then loaded onto a Heparin column (Cytiva) and eluted using a linear NaCl gradient (0.1 to 1 M). The σ^E-RNAP eluate was diluted in the same buffer but purified using an HP Q column (Cytiva). Following concentration, proteins were further purified by size exclusion chromatography (SEC) using a Superose 6 Increase 10/300 GL column (Cytiva) in equilibration buffer containing 25 mM Tris-HCl pH 7.6, 250 mM NaCl, 5% glycerol, and 2 mM DTT. Purified fractions were concentrated and stored at -80 °C.

The *E. coli* RNAP–σ^{70} holoenzyme (NEB, M0551S) was purchased as a control for detecting chimeric *Fta-Eco* RNAP formation.

Complex assembly. The dCas12f-gRNA-target DNA ternary complex was reconstituted by incubating dCas12f-gRNA with a 60-bp target DNA **(Supplementary Table 1)** at a 1:1.3 ratio for 30 minutes at 30 °C. The mixture was then subjected to SEC using a Superose 6 column equilibrated in buffer containing

25 mM Tris-HCl pH 7.6, 150 mM NaCl, 2 mM DTT, and 5 mM $MgCl_2$.

For assembly of the dCas12f-gRNA-target DNA-σ^E-RNAP complex, a longer 99-bp target DNA was used, consisting of the original 60-bp sequence extended by 39 bp at the 3' end (**Supplementary Table 1**). The ternary complex was assembled following the same procedure described above, then incubated with σ^E-RNAP overnight at 4 °C. The final complex was purified by SEC using the same equilibration buffer.

Electron microscopy. Aliquots of 3 μL dCas12f-gRNA binary complex (1 mg/mL), dCas12f-gRNA-target DNA ternary complex (1 mg/mL), or dCas12f-gRNA-target DNA-σ^E-RNAP complex (3 mg/mL) were applied to glow-discharged Quantifoil holey carbon girds (R1.2/1.3, 300mesh). The grids were blotted for 2.5 seconds and plunged into liquid ethane using a ThermoFisher Scientific Mark IV Vitrobot. Cryo-EM data were collected with a Titan Krios G4 microscope (FEI) operated at 300 kV and images were collected using EPU at a nominal magnification of 105,000x (resulting in a calibrated physical pixel size of 0.822 Å/pixel) with a defocus range of 0.8–2.0 μm. Three images per hole were collected using EPU with the beam-image shift method. Images were recorded on a K3 electron direct detector in super-resolution mode at the end of a GIF-Quantum energy filter operated with a slit width of 20 eV. A dose rate of 20 electrons per pixel per second and an exposure time of 2.01 seconds were used, generating 40 movie frames with a total dose of ~59.2 electrons per $Å^2$.

For the dCas12f-gRNA binary complex, we collected 4,149 micrographs. After screening based on CTF-estimated resolution and manual inspection, 3,982 micrographs were selected for further analysis. For the dCas12f-gRNA-DNA ternary complex, 2,521 micrographs were selected from a total of 2,652 collected images.

For the dCas12f-gRNA-DNA-σ^E-RNAP complex with 99-bp native DNA, we collected 7,601 micrographs and selected 7,289 for further analysis. We then modified this complex by introducing a 15-bp mismatch region in the native DNA to facilitate RNAP open complex formation. For this modified complex, we collected 8,378 micrographs and selected 7,656 for further analysis.

Image processing. Movie frames were initially imported to cryoSPARC[75] live for on-the-fly processing during data collection. Movie frames were aligned by patch motion correction with a binning factor of 2. Contrast transfer function (CTF) parameters were estimated using Patch CTF. A few thousand particles were auto-picked without a template to generate 2D averages for subsequent template-based auto-picking. The auto-picked and extracted particles were screened by 2D classification to exclude false and bad particles that fall into 2D averages with poor features. A subset of the particles (100,000 particles) was used to generate initial models in cryoSPARC. All screened particles were subjected to heterogenous refinement using the initial models. Particles in each 3D class were subject to homogenous refinement.

Selected particles were used to generate a Topaz model for automated particle picking[76] from all micrographs. The picked particles underwent sequential processing in cryoSPARC: 2D classification, *ab initio* reconstruction (2–4 classes), heterogeneous refinement, and Non-uniform Refinement. Results were

manually inspected after each step to determine whether to rerun with modifications (e.g., adjusting class numbers) or proceed to the next step. In the final Non-uniform Refinement, we applied per-particle CTF correction. Local Refinement was performed when needed, such as for the Cas12f region in the dCas12f-gRNA-DNA-σ^E-RNAP complex. Resolution was estimated using the FSC = 0.143 criterion. Maps were post-processed using DeepEMhancer[77] to reduce noise levels and used for model building and refinement. For the dCas12f-gRNA-DNA-σ^E-RNAP complex, a composite map was generated by combining two maps (one from general refinement and one from local refinement around the Cas12f region) using the *'vop maximum'* command in Chimera[78], which was then used for model refinement.

Model building and refinement. AlphaFold3 models[79] were used as initial models. The individual structures were manually fitted onto respective maps as a rigid-body in UCSF Chimera[78] and manually adjusted in COOT[80]. The model was then fitted in COOT with all-molecule self-restrains to maintain H-bonds and stacking interactions between bases. Finally, refinement of the structure models against corresponding maps were performed using *phenix.real_space_refine* tool in Phenix[81].

Sequence analysis. Sequence alignments were performed using Clustal Omega[82], and alignment diagrams were plotted using ESPript[83]. The secondary structure of gRNA **(Fig. 2d)** was visualized by Forna (force-directed RNA)[84] and manually adjusted according to the cryo-EM structure.

Structural analysis and visualization. Figures were generated using PyMOL, UCSF Chimera[78] and ChimeraX[85]. To calculate the RMSD of two structures, the *align* command in PyMOL was used.

RFP fluorescence assay. RFP assays were performed as described in Hoffmann *et al.*[8]. Briefly, *E. coli* BL21(*DE3*) cells encoding *sfGFP* in the chromosome were co-transformed with a pTarget plasmid, encoding *mRFP1* downstream of a 500 bp sequence cloned from the native *Fta* strain that includes dCas12f-gRNA target site, and a plasmid encoding *Fta*RNAP. In another round of transformations, a pEffector plasmid encoding the native *Fta* gRNA and *Fta* dCas12f-σ^E protein components (or mutants thereof) from the native strain (GenBank accession GCA_003584105.1) was delivered. *E. coli* cultures were grown by picking three individual colonies per sample and inoculating 5 ml of LB in 24-well plates, or 0.5 ml of LB in 96-well plates. Expression of *Fta* RNAP was induced by adding IPTG to a final concentration of 0.01 mM. Cell cultures were grown at 28 °C for 20-24 h. Then, 200 μl from each well was transferred to a 96-well optical bottom plate (Thermo Scientific) to take endpoint measurements for OD_{600} mRFP fluorescence using a Synergy Neo2 microplate reader (Biotek).

F. taeanensis **σ phylogeny**. The amino acid sequence for *E. coli* K-12 MG1655 (genomic accession NC_000913.3) σ^E (protein accession: WP_001295364.1) and σ^{70} (protein accession WP_000437376.1) was used to perform a BLASTp search in the *F. taenensis* genome (genomic accession GCA_003584105.1) to identify all *Fta*-encoded σ factors. To ensure that all σ^D and σ^E-encoding genes were discovered, anoth-

er BLASTp search using an *Fta*-encoded σ^D (protein accession RIV53857.1) and σ^E (protein accession RIV50554.1) was performed, respectively. The resulting 30 σ^D and σ^E sequences were used to build a Hidden Markov Model (HMM) using the hmmbuild command. The output file was used to perform an hmmsearch against the *Fta* proteome which yielded 46 hits, of which 34 were unique. Four sequences with non-σ annotations were manually excluded. MAFFT (LINSI option) was used to generate a multiple sequence alignment of the *Fta* σ sequences. A phylogenetic tree was generated using the following command: iqtree2 [alignment.fasta] -nt AUTO -m WAG -alrt 1000 -bb 1000 -abayes. The unrooted phylogenetic tree was generated in iTOL (online). σ homologs were chosen for gene synthesis to reflect the phylogenetic diversity of *Fta* σ factors.

Quantification and statistical analysis. Detailed statistical information for the experiments is provided in the figure legends. Statistical validation for the final structural models was performed using Phenix[81].

Data availability. Cryo-EM reconstructions have been deposited in the Electron Microscopy Data Bank under the accession numbers EMD-49165, EMD-49173, EMD-49174, EMD-49175, and EMD-49176. Coordinates for atomic models have been deposited in the Protein Data Bank under the accession numbers 9N9C, 9N9M, 9N9O, 9N9P, and 9N9Q.

ACKNOWLEDGMENTS

We thank Frank Vago and Thomas Klose for assistance in cryo-EM data collection, and Steven Wilson for computation. This work made use of the Purdue Cryo-EM Facility. L.C. was supported by the NIH grant R01GM138675 and the NSF CAREER Award 2339799. S.H.S. was supported by NSF Faculty Early Career Development Program (CAREER) Award 2239685, a Pew Biomedical Scholarship, an Irma T. Hirschl Career Scientist Award, the Howard Hughes Medical Institute Investigator Program, and a generous startup package from the Columbia University Irving Medical Center Dean's Office and the Vagelos Precision Medicine Fund.

AUTHOR CONTRIBUTIONS

R.X. prepared the samples for cryo-EM and performed all biochemical assays. F.T.H and A.I.P. performed the cellular transcription assays for protein mutants. T.W. performed bioinformatics analyses. D.X. contributed to molecular cloning and protein purification. R.X. and L.C. collected cryo-EM data. L.C. and R.X. processed cryo-EM data and built the structural models. S.H.S. contributed to data analysis, interpretation, and manuscript editing. All authors participated in data analysis and contributed to the preparation of the manuscript. L.C. and S.H.S. supervised the research.

COMPETING INTERESTS

S.H.S. is a co-founder and scientific advisor to Dahlia Biosciences, a scientific advisor to CrisprBits and Prime Medicine, and an equity holder in Dahlia Biosciences and CrisprBits. S.H.S., F.T.H., and T.W. are inventors on patents related to CRISPR-Cas-like systems and uses thereof. The other authors declare no competing interests.

Correspondence and requests for materials should be addressed to L.C. (lchang18@purdue.edu.) and S.H.S. (shsternberg@gmail.com).

EXTENDED DATA

Extended Data Figure 1 | Bioinformatics and structural analysis of all 30 σ factors in the *F. taeanensis* genome.

Extended Data Figure 2 | Biochemical analysis of the dCas12f-σ^E system.

Extended Data Figure 3 | Cryo-EM of the *Fta* dCas12f-gRNA complex.

Extended Data Figure 4 | Cryo-EM of the *Fta* dCas12f-gRNA-DNA complex.

Extended Data Figure 5 | Structural comparison of *Fta* dCas12f, *As*Cas12f, and *Un*Cas12f.

Extended Data Figure 6 | Cryo-EM structure of the dCas12f-σ^E-RNAP complex bound to 99-bp fully matched dsDNA.

Extended Data Figure 7 | Cryo-EM structure of the dCas12f-σ^E-RNAP complex bound to a 99-bp dsDNA containing 15-bp mismatches.

Extended Data Figure 8 | Model-to-map fitting for each subunit of the DNA-bound dCas12f-σ^E-RNAP complex.

Extended Data Figure 9 | Interaction between target DNA and protein subunits in the dCas12f-σ^E-RNAP complex.

Extended Data Figure 10 | Structural comparison of DNA-bound *Fta* dCas12f-σ^E-RNAP and *E. col*i σ^E-RNAP complexes.

Extended Data Table 1 | Cryo-EM data collection, refinement and validation statistics.

SUPPLEMENTARY TABLE

Supplementary Table 1 | Sequence of RNA and DNA substrates used in this study.

SUPPLEMENTARY FIGURE

Supplementary Figure 1 | Source data for electrophoretic separation experiments.

REFERENCES

Wiegand, T., Hoffmann, F. T., Walker, M. W. G., Tang, S., Richard, E., Le, H. C., Meers, C. & Sternberg, S. H. TnpB homologues exapted from transposons are RNA-guided transcription factors. *Nature* **631**, 439-448, doi:10.1038/s41586-024-07598-4 (2024).

Qi, L. S., Larson, M. H., Gilbert, L. A., Doudna, J. A., Weissman, J. S., Arkin, A. P. & Lim, W. A. Repurposing CRISPR as an RNA-guided platform for sequence-specific control of gene expression. *Cell* **152**, 1173-1183, doi:10.1016/j.cell.2013.02.022 (2013).

Gilbert, L. A., Horlbeck, M. A., Adamson, B., Villalta, J. E., Chen, Y., Whitehead, E. H., Guimaraes, C., Panning, B., Ploegh, H. L., Bassik, M. C., Qi, L. S., Kampmann, M. & Weissman, J. S. Genome-Scale CRISPR-Mediated Control of Gene Repression and Activation. *Cell* **159**, 647-661, doi:10.1016/j.cell.2014.09.029 (2014).

Bikard, D., Jiang, W., Samai, P., Hochschild, A., Zhang, F. & Marraffini, L. A. Programmable repression and activation of bacterial gene expression using an engineered CRISPR-Cas system. *Nucleic Acids Res* **41**, 7429-7437, doi:10.1093/nar/gkt520 (2013).

Zalatan, J. G., Lee, M. E., Almeida, R., Gilbert, L. A., Whitehead, E. H., La Russa, M., Tsai, J. C., Weissman, J. S., Dueber, J. E., Qi, L. S. & Lim, W. A. Engineering complex synthetic transcriptional programs with CRISPR RNA scaffolds. *Cell* **160**, 339-350, doi:10.1016/j.cell.2014.11.052 (2015).

Workman, R. E., Pammi, T., Nguyen, B. T. K., Graeff, L. W., Smith, E., Sebald, S. M., Stoltzfus, M. J., Euler, C. W. & Modell, J. W. A natural single-guide RNA repurposes Cas9 to autoregulate CRISPR-Cas expression. *Cell* **184**, 675-688 e619, doi:10.1016/j.cell.2020.12.017 (2021).

Wu, W. Y., Mohanraju, P., Liao, C., Adiego-Perez, B., Creutzburg, S. C. A., Makarova, K. S., Keessen, K., Lindeboom, T. A., Khan, T. S., Prinsen, S., Joosten, R., Yan, W. X., Migur, A., Laffeber, C., Scott, D. A., Lebbink, J. H. G., Koonin, E. V., Beisel, C. L. & van der Oost, J. The miniature CRISPR-Cas12m effector binds DNA to block transcription. *Mol Cell* **82**, 4487-4502 e4487, doi:10.1016/j.molcel.2022.11.003 (2022).

Hoffmann, F. T., Wiegand, T., Palmieri, A. I., Glass-Klaiber, J., Xiao, R., Tang, S., Le, H. C., Meers, C., Lampe, G. D., Chang, L. & Sternberg, S. H. Exapted CRISPR-Cas12f homologs drive RNA-guided transcription. *Submitted* (2025).

Burgess, R. R., Travers, A. A., Dunn, J. J. & Bautz, E. K. Factor stimulating transcription by RNA polymerase. *Nature* **221**, 43-46, doi:10.1038/221043a0 (1969).

Ross, W., Gosink, K. K., Salomon, J., Igarashi, K., Zou, C., Ishihama, A., Severinov, K. & Gourse, R. L. A third recognition element in bacterial promoters: DNA binding by the alpha subunit of RNA polymerase. *Science* **262**, 1407-1413, doi:10.1126/science.8248780 (1993).

Lee, D. J., Minchin, S. D. & Busby, S. J. Activating transcription in bacteria. *Annu Rev Microbiol* **66**, 125-152, doi:10.1146/annurev-micro-092611-150012 (2012).

Chen, J., Boyaci, H. & Campbell, E. A. Diverse and unified mechanisms of transcription initiation in bacteria. *Nat Rev Microbiol* **19**, 95-109, doi:10.1038/s41579-020-00450-2 (2021).

Feklistov, A., Sharon, B. D., Darst, S. A. & Gross, C. A. Bacterial sigma factors: a historical, structural, and genomic perspective. *Annu Rev Microbiol* **68**, 357-376, doi:10.1146/annurev-mi-

cro-092412-155737 (2014).

14   Paget, M. S. Bacterial Sigma Factors and Anti-Sigma Factors: Structure, Function and Distribu-
      tion. *Biomolecules* **5**, 1245-1265, doi:10.3390/biom5031245 (2015).

15   Paget, M. S. & Helmann, J. D. The sigma70 family of sigma factors. *Genome Biol* **4**, 203,
      doi:10.1186/gb-2003-4-1-203 (2003).

16   Campbell, E. A., Tupy, J. L., Gruber, T. M., Wang, S., Sharp, M. M., Gross, C. A. & Darst, S. A.
      Crystal structure of Escherichia coli sigmaE with the cytoplasmic domain of its anti-sigma RseA.
      Mol Cell **11**, 1067-1078, doi:10.1016/s1097-2765(03)00148-5 (2003).

17   Casas-Pastor, D., Muller, R. R., Jaenicke, S., Brinkrolf, K., Becker, A., Buttner, M. J., Gross,
      C. A., Mascher, T., Goesmann, A. & Fritz, G. Expansion and re-classification of the extracyto-
      plasmic function (ECF) sigma factor family. *Nucleic Acids Res* **49**, 986-1005, doi:10.1093/nar/
      gkaa1229 (2021).

18   Mascher, T. Past, Present, and Future of Extracytoplasmic Function sigma Factors: Distribution
      and Regulatory Diversity of the Third Pillar of Bacterial Signal Transduction. *Annu Rev Microbi-
      ol* **77**, 625-644, doi:10.1146/annurev-micro-032221-024032 (2023).

19   Rowley, G., Spector, M., Kormanec, J. & Roberts, M. Pushing the envelope: extracytoplasmic
      stress responses in bacterial pathogens. *Nat Rev Microbiol* **4**, 383-394, doi:10.1038/nrmicro1394
      (2006).

20   Lonetto, M. A., Brown, K. L., Rudd, K. E. & Buttner, M. J. Analysis of the Streptomyces co-
      elicolor sigE gene reveals the existence of a subfamily of eubacterial RNA polymerase sigma
      factors involved in the regulation of extracytoplasmic functions. *Proc Natl Acad Sci U S A* **91**,
      7573-7577, doi:10.1073/pnas.91.16.7573 (1994).

21   Altae-Tran, H., Shmakov, S. A., Makarova, K. S., Wolf, Y. I., Kannan, S., Zhang, F. & Koonin,
      E. V. Diversity, evolution, and classification of the RNA-guided nucleases TnpB and Cas12. *Proc
      Natl Acad Sci U S A* **120**, e2308224120, doi:10.1073/pnas.2308224120 (2023).

22   Sampson, T. R., Saroj, S. D., Llewellyn, A. C., Tzeng, Y. L. & Weiss, D. S. A CRISPR/Cas sys-
      tem mediates bacterial innate immune evasion and virulence. *Nature* **497**, 254-257, doi:10.1038/
      nature12048 (2013).

23   Ratner, H. K., Escalera-Maurer, A., Le Rhun, A., Jaggavarapu, S., Wozniak, J. E., Crispell, E. K.,
      Charpentier, E. & Weiss, D. S. Catalytically Active Cas9 Mediates Transcriptional Interference
      to Facilitate Bacterial Virulence. *Mol Cell* **75**, 498-510 e495, doi:10.1016/j.molcel.2019.05.029
      (2019).

24   Huang, C. J., Adler, B. A. & Doudna, J. A. A naturally DNase-free CRISPR-Cas12c enzyme
      silences gene expression. *Mol Cell* **82**, 2148-2160 e2144, doi:10.1016/j.molcel.2022.04.020
      (2022).

25   Takeda, S. N., Nakagawa, R., Okazaki, S., Hirano, H., Kobayashi, K., Kusakizako, T., Nishiza-
      wa, T., Yamashita, K., Nishimasu, H. & Nureki, O. Structure of the miniature type V-F CRIS-
      PR-Cas effector enzyme. *Mol Cell* **81**, 558-570 e553, doi:10.1016/j.molcel.2020.11.035 (2021).

26   Xiao, R., Li, Z., Wang, S., Han, R. & Chang, L. Structural basis for substrate recognition and
      cleavage by the dimerization-dependent CRISPR-Cas12f nuclease. *Nucleic Acids Res* **49**, 4120-

4128, doi:10.1093/nar/gkab179 (2021).

Hino, T., Omura, S. N., Nakagawa, R., Togashi, T., Takeda, S. N., Hiramoto, T., Tasaka, S., Hirano, H., Tokuyama, T., Uosaki, H., Ishiguro, S., Kagieva, M., Yamano, H., Ozaki, Y., Motoo-ka, D., Mori, H., Kirita, Y., Kise, Y., Itoh, Y., Matoba, S., Aburatani, H., Yachie, N., Karvelis, T., Siksnys, V., Ohmori, T., Hoshino, A. & Nureki, O. An AsCas12f-based compact genome-editing tool derived by deep mutational scanning and structural analysis. *Cell* **186**, 4920-4935 e4923, doi:10.1016/j.cell.2023.08.031 (2023).

Wu, T., Liu, C., Zou, S., Lyu, R., Yang, B., Yan, H., Zhao, M. & Tang, W. An engineered hyper-compact CRISPR-Cas12f system with boosted gene-editing activity. *Nat Chem Biol* **19**, 1384-1393, doi:10.1038/s41589-023-01380-9 (2023).

Harrington, L. B., Burstein, D., Chen, J. S., Paez-Espino, D., Ma, E., Witte, I. P., Cofsky, J. C., Kyrpides, N. C., Banfield, J. F. & Doudna, J. A. Programmed DNA destruction by miniature CRISPR-Cas14 enzymes. *Science* **362**, 839-842, doi:10.1126/science.aav4294 (2018).

Bigelyte, G., Young, J. K., Karvelis, T., Budre, K., Zedaveinyte, R., Djukanovic, V., Van Ginkel, E., Paulraj, S., Gasior, S., Jones, S., Feigenbutz, L., Clair, G. S., Barone, P., Bohn, J., Acharya, A., Zastrow-Hayes, G., Henkel-Heinecke, S., Silanskas, A., Seidel, R. & Siksnys, V. Miniature type V-F CRISPR-Cas nucleases enable targeted DNA modification in cells. *Nat Commun* **12**, 6191, doi:10.1038/s41467-021-26469-4 (2021).

Yang, H., Gao, P., Rajashankar, K. R. & Patel, D. J. PAM-Dependent Target DNA Recognition and Cleavage by C2c1 CRISPR-Cas Endonuclease. *Cell* **167**, 1814-1828 e1812, doi:10.1016/j.cell.2016.11.053 (2016).

Swarts, D. C., van der Oost, J. & Jinek, M. Structural Basis for Guide RNA Processing and Seed-Dependent DNA Targeting by CRISPR-Cas12a. *Mol Cell* **66**, 221-233 e224, doi:10.1016/j.molcel.2017.03.016 (2017).

Liu, L., Chen, P., Wang, M., Li, X., Wang, J., Yin, M. & Wang, Y. C2c1-sgRNA Complex Structure Reveals RNA-Guided DNA Cleavage Mechanism. *Mol Cell* **65**, 310-322, doi:10.1016/j.molcel.2016.11.040 (2017).

Zhang, H., Li, Z., Xiao, R. & Chang, L. Mechanisms for target recognition and cleavage by the Cas12i RNA-guided endonuclease. *Nat Struct Mol Biol* **27**, 1069-1076, doi:10.1038/s41594-020-0499-0 (2020).

Sternberg, S. H., Redding, S., Jinek, M., Greene, E. C. & Doudna, J. A. DNA interrogation by the CRISPR RNA-guided endonuclease Cas9. *Nature* **507**, 62-67, doi:10.1038/nature13011 (2014).

Liu, B., Hong, C., Huang, R. K., Yu, Z. & Steitz, T. A. Structural basis of bacterial transcription activation. *Science* **358**, 947-951, doi:10.1126/science.aao1923 (2017).

Kompaniiets, D., Wang, D., Yang, Y., Hu, Y. & Liu, B. Structure and molecular mechanism of bacterial transcription activation. *Trends Microbiol* **32**, 379-397, doi:10.1016/j.tim.2023.10.001 (2024).

Chen, J., Gopalkrishnan, S., Chiu, C., Chen, A. Y., Campbell, E. A., Gourse, R. L., Ross, W. & Darst, S. A. E. coli TraR allosterically regulates transcription initiation by altering RNA polymerase conformation. *Elife* **8**, doi:10.7554/eLife.49375 (2019).

Bae, B., Chen, J., Davis, E., Leon, K., Darst, S. A. & Campbell, E. A. CarD uses a minor groove wedge mechanism to stabilize the RNA polymerase open promoter complex. *Elife* **4**, doi:10.7554/eLife.08505 (2015).

Fang, C., Li, L., Shen, L., Shi, J., Wang, S., Feng, Y. & Zhang, Y. Structures and mechanism of transcription initiation by bacterial ECF factors. *Nucleic Acids Res* **47**, 7094-7104, doi:10.1093/nar/gkz470 (2019).

Lin, W., Mandal, S., Degen, D., Cho, M. S., Feng, Y., Das, K. & Ebright, R. H. Structural basis of ECF-sigma-factor-dependent transcription initiation. *Nat Commun* **10**, 710, doi:10.1038/s41467-019-08443-3 (2019).

Li, L., Fang, C., Zhuang, N., Wang, T. & Zhang, Y. Structural basis for transcription initiation by bacterial ECF sigma factors. *Nat Commun* **10**, 1153, doi:10.1038/s41467-019-09096-y (2019).

Bae, B., Feklistov, A., Lass-Napiorkowska, A., Landick, R. & Darst, S. A. Structure of a bacterial RNA polymerase holoenzyme open promoter complex. *Elife* **4**, doi:10.7554/eLife.08504 (2015).

Basu, R. S., Warner, B. A., Molodtsov, V., Pupov, D., Esyunina, D., Fernandez-Tornero, C., Kulbachinskiy, A. & Murakami, K. S. Structural basis of transcription initiation by bacterial RNA polymerase holoenzyme. *J Biol Chem* **289**, 24549-24559, doi:10.1074/jbc.M114.584037 (2014).

Fredrick, K. & Helmann, J. D. RNA polymerase sigma factor determines start-site selection but is not required for upstream promoter element activation on heteroduplex (bubble) templates. *Proc Natl Acad Sci U S A* **94**, 4982-4987, doi:10.1073/pnas.94.10.4982 (1997).

Marin-Gonzalez, A., Vilhena, J. G., Perez, R. & Moreno-Herrero, F. A molecular view of DNA flexibility. *Q Rev Biophys* **54**, e8, doi:10.1017/S0033583521000068 (2021).

Vishwakarma, R. K., Cao, A. M., Morichaud, Z., Perumal, A. S., Margeat, E. & Brodolin, K. Single-molecule analysis reveals the mechanism of transcription activation in M. tuberculosis. *Sci Adv* **4**, eaao5498, doi:10.1126/sciadv.aao5498 (2018).

Mooney, R. A., Darst, S. A. & Landick, R. Sigma and RNA polymerase: an on-again, off-again relationship? *Mol Cell* **20**, 335-345, doi:10.1016/j.molcel.2005.10.015 (2005).

Marchetti, M., Malinowska, A., Heller, I. & Wuite, G. J. L. How to switch the motor on: RNA polymerase initiation steps at the single-molecule level. *Protein Science* **26**, 1303-1313, doi:10.1002/pro.3183 (2017).

Lin, W., Mandal, S., Degen, D., Liu, Y., Ebright, Y. W., Li, S., Feng, Y., Zhang, Y., Mandal, S., Jiang, Y., Liu, S., Gigliotti, M., Talaue, M., Connell, N., Das, K., Arnold, E. & Ebright, R. H. Structural Basis of Mycobacterium tuberculosis Transcription and Transcription Inhibition. *Mol Cell* **66**, 169-179 e168, doi:10.1016/j.molcel.2017.03.001 (2017).

Nudler, E. RNA polymerase active center: the molecular engine of transcription. *Annu Rev Biochem* **78**, 335-361, doi:10.1146/annurev.biochem.76.052705.164655 (2009).

Hirata, A., Klein, B. J. & Murakami, K. S. The X-ray crystal structure of RNA polymerase from Archaea. *Nature* **451**, 851-854, doi:10.1038/nature06530 (2008).

Cramer, P., Bushnell, D. A. & Kornberg, R. D. Structural basis of transcription: RNA polymerase II at 2.8 angstrom resolution. *Science* **292**, 1863-1876, doi:10.1126/science.1059493 (2001).

Geszvain, K., Gruber, T. M., Mooney, R. A., Gross, C. A. & Landick, R. A hydrophobic patch
        on the flap-tip helix of E.coli RNA polymerase mediates sigma(70) region 4 function. *J Mol Biol*
        343, 569-587, doi:10.1016/j.jmb.2004.08.063 (2004).

55   Mathew, R. & Chatterji, D. The evolving story of the omega subunit of bacterial RNA poly-
        merase. *Trends Microbiol* **14**, 450-455, doi:10.1016/j.tim.2006.08.002 (2006).

56   Ghosh, P., Ishihama, A. & Chatterji, D. Escherichia coli RNA polymerase subunit omega and its
        N-terminal domain bind full-length beta' to facilitate incorporation into the alpha2beta subassem-
        bly. *Eur J Biochem* **268**, 4621-4627, doi:10.1046/j.1432-1327.2001.02381.x (2001).

57   Ghosh, P., Ramakrishnan, C. & Chatterji, D. Inter-subunit recognition and manifestation of seg-
        mental mobility in Escherichia coli RNA polymerase: a case study with omega-beta' interaction.
        Biophys Chem **103**, 223-237, doi:10.1016/s0301-4622(02)00271-5 (2003).

58   Mukherjee, K. & Chatterji, D. Studies on the omega subunit of Escherichia coli RNA poly-
        merase--its role in the recovery of denatured enzyme activity. *Eur J Biochem* **247**, 884-889,
        doi:10.1111/j.1432-1033.1997.00884.x (1997).

59   Bhowmik, D., Bhardwaj, N. & Chatterji, D. Influence of Flexible "omega" on the Activity of
        E. coli RNA Polymerase: A Thermodynamic Analysis. *Biophys J* **112**, 901-910, doi:10.1016/j.
        bpj.2017.01.015 (2017).

60   Bu, F., Wang, X., Li, M., Ma, L., Wang, C., Hu, Y., Cao, Z. & Liu, B. Cryo-EM Structure
        of Porphyromonas gingivalis RNA Polymerase. *J Mol Biol* **436**, 168568, doi:10.1016/j.
        jmb.2024.168568 (2024).

61   Lane, W. J. & Darst, S. A. The structural basis for promoter -35 element recognition by the group
        IV sigma factors. *PLoS Biol* **4**, e269, doi:10.1371/journal.pbio.0040269 (2006).

62   Li, J., Zhang, H., Li, D., Liu, Y. J., Bayer, E. A., Cui, Q., Feng, Y. & Zhu, P. Structure of the tran-
        scription open complex of distinct sigma(I) factors. *Nat Commun* **14**, 6455, doi:10.1038/s41467-
        023-41796-4 (2023).

63   Rohs, R., West, S. M., Sosinsky, A., Liu, P., Mann, R. S. & Honig, B. The role of DNA shape in
        protein-DNA recognition. *Nature* **461**, 1248-1253, doi:10.1038/nature08473 (2009).

64   Zhang, Y., Feng, Y., Chatterjee, S., Tuske, S., Ho, M. X., Arnold, E. & Ebright, R. H. Structural
        basis of transcription initiation. *Science* **338**, 1076-1080, doi:10.1126/science.1227786 (2012).

65   Campagne, S., Marsh, M. E., Capitani, G., Vorholt, J. A. & Allain, F. H. Structural basis for -10
        promoter element melting by environmentally induced sigma factors. *Nat Struct Mol Biol* **21**,
        269-276, doi:10.1038/nsmb.2777 (2014).

66   Feklistov, A. & Darst, S. A. Structural basis for promoter-10 element recognition by the bacterial
        RNA polymerase sigma subunit. *Cell* **147**, 1257-1269, doi:10.1016/j.cell.2011.10.041 (2011).

67   Lim, H. M., Lee, H. J., Roy, S. & Adhya, S. A "master" in base unpairing during isomeriza-
        tion of a promoter upon RNA polymerase binding. *Proc Natl Acad Sci U S A* **98**, 14849-14852,
        doi:10.1073/pnas.261517398 (2001).

68   Feng, Y., Zhang, Y. & Ebright, R. H. Structural basis of transcription activation. *Science* **352**,
        1330-1333, doi:10.1126/science.aaf4417 (2016).

Zuo, Y. & Steitz, T. A. Crystal structures of the E. coli transcription initiation complexes with a complete bubble. *Mol Cell* **58**, 534-540, doi:10.1016/j.molcel.2015.03.010 (2015).

Boyaci, H., Chen, J., Jansen, R., Darst, S. A. & Campbell, E. A. Structures of an RNA polymerase promoter melting intermediate elucidate DNA unwinding. *Nature* **565**, 382-385, doi:10.1038/s41586-018-0840-5 (2019).

Feklistov, A. RNA polymerase: in search of promoters. *Ann N Y Acad Sci* **1293**, 25-32, doi:10.1111/nyas.12197 (2013).

Sakata-Sogawa, K. & Shimamoto, N. RNA polymerase can track a DNA groove during promoter search. *Proc Natl Acad Sci U S A* **101**, 14731-14735, doi:10.1073/pnas.0406441101 (2004).

Bustamante, C., Guthold, M., Zhu, X. & Yang, G. Facilitated target location on DNA by individual Escherichia coli RNA polymerase molecules observed with the scanning force microscope operating in liquid. *J Biol Chem* **274**, 16665-16668, doi:10.1074/jbc.274.24.16665 (1999).

Wang, F., Redding, S., Finkelstein, I. J., Gorman, J., Reichman, D. R. & Greene, E. C. The promoter-search mechanism of Escherichia coli RNA polymerase is dominated by three-dimensional diffusion. *Nat Struct Mol Biol* **20**, 174-181, doi:10.1038/nsmb.2472 (2013).

Punjani, A., Rubinstein, J. L., Fleet, D. J. & Brubaker, M. A. cryoSPARC: algorithms for rapid unsupervised cryo-EM structure determination. *Nat Methods* **14**, 290-296, doi:10.1038/nmeth.4169 (2017).

Bepler, T., Morin, A., Rapp, M., Brasch, J., Shapiro, L., Noble, A. J. & Berger, B. Positive-unlabeled convolutional neural networks for particle picking in cryo-electron micrographs. *Nat Methods* **16**, 1153-1160, doi:10.1038/s41592-019-0575-8 (2019).

Sanchez-Garcia, R., Gomez-Blanco, J., Cuervo, A., Carazo, J. M., Sorzano, C. O. S. & Vargas, J. DeepEMhancer: a deep learning solution for cryo-EM volume post-processing. *Commun Biol* **4**, 874, doi:10.1038/s42003-021-02399-1 (2021).

Pettersen, E. F., Goddard, T. D., Huang, C. C., Couch, G. S., Greenblatt, D. M., Meng, E. C. & Ferrin, T. E. UCSF Chimera--a visualization system for exploratory research and analysis. *J Comput Chem* **25**, 1605-1612, doi:10.1002/jcc.20084 (2004).

Abramson, J., Adler, J., Dunger, J., Evans, R., Green, T., Pritzel, A., Ronneberger, O., Willmore, L., Ballard, A. J., Bambrick, J., Bodenstein, S. W., Evans, D. A., Hung, C. C., O'Neill, M., Reiman, D., Tunyasuvunakool, K., Wu, Z., Zemgulyte, A., Arvaniti, E., Beattie, C., Bertolli, O., Bridgland, A., Cherepanov, A., Congreve, M., Cowen-Rivers, A. I., Cowie, A., Figurnov, M., Fuchs, F. B., Gladman, H., Jain, R., Khan, Y. A., Low, C. M. R., Perlin, K., Potapenko, A., Savy, P., Singh, S., Stecula, A., Thillaisundaram, A., Tong, C., Yakneen, S., Zhong, E. D., Zielinski, M., Zidek, A., Bapst, V., Kohli, P., Jaderberg, M., Hassabis, D. & Jumper, J. M. Accurate structure prediction of biomolecular interactions with AlphaFold 3. *Nature* **630**, 493-500, doi:10.1038/s41586-024-07487-w (2024).

Emsley, P., Lohkamp, B., Scott, W. G. & Cowtan, K. Features and development of Coot. *Acta Crystallogr D Biol Crystallogr* **66**, 486-501, doi:10.1107/S0907444910007493 (2010).

Afonine, P. V., Poon, B. K., Read, R. J., Sobolev, O. V., Terwilliger, T. C., Urzhumtsev, A. & Adams, P. D. Real-space refinement in PHENIX for cryo-EM and crystallography. *Acta Crystallogr D Struct Biol* **74**, 531-544, doi:10.1107/S2059798318006551 (2018).

McWilliam, H., Li, W., Uludag, M., Squizzato, S., Park, Y. M., Buso, N., Cowley, A. P. & Lopez, R. Analysis Tool Web Services from the EMBL-EBI. *Nucleic Acids Res* **41**, W597-600, doi:10.1093/nar/gkt376 (2013).

Robert, X. & Gouet, P. Deciphering key features in protein structures with the new ENDscript server. *Nucleic Acids Res* **42**, W320-324, doi:10.1093/nar/gku316 (2014).

Kerpedjiev, P., Hammer, S. & Hofacker, I. L. Forna (force-directed RNA): Simple and effective online RNA secondary structure diagrams. *Bioinformatics* **31**, 3377-3379, doi:10.1093/bioinformatics/btv372 (2015).

Pettersen, E. F., Goddard, T. D., Huang, C. C., Meng, E. C., Couch, G. S., Croll, T. I., Morris, J. H. & Ferrin, T. E. UCSF ChimeraX: Structure visualization for researchers, educators, and developers. *Protein Sci* **30**, 70-82, doi:10.1002/pro.3943 (2021).