1

Impaired envelope integrity in the absence of SanA is linked to increased lipid II availability and an imbalance of septal peptidoglycan synthesis

Joseph F. Carr[a], Carolina Basurto De Santiago[a,‡], Shivani A. Bhut[a,‡], Daniel J. Warzecha[a,‡], Sarah A. Vastani[a], Robert Wei[a], Carmen M. Herrera[b], M. Stephen Trent[b,c], Beiyan Nan[a], Angela M. Mitchell[a,#]

[a] Department of Biology, Texas A&M University, College Station, Texas, USA

[b] Department of Infectious Diseases, College of Veterinary Medicine, University of Georgia, Athens, Georgia, USA

[c] Department of Microbiology, College of Art and Sciences; University of Georgia, Athens, Georgia, USA

Running Title: Envelope permeability from altered septum synthesis

Address correspondence to Angela M. Mitchell, amitchell@bio.tamu.edu

‡ Authors contributed equally to this work. Order was determined alphabetically.

Abstract: 236 words

Main Text: 7096 words

ABSTRACT

In Gram-negative bacteria, the outer membrane (OM) acts in conjunction with the peptidoglycan (PG) cell wall as a barrier against physical, osmotic, and chemical environmental stressors including antibiotics. SanA, an inner membrane protein in *Escherichia coli* K-12, is required for vancomycin resistance at high temperatures (>42 °C) and impacts sodium dodecyl sulfate (SDS) resistance during stationary phase reached from carbon limitation. However, its function remains unknown. Here, we show that Δ*sanA* has a synthetic genetic interaction with Δ*wecA*, a mutation that increases the availability of the isoprenoid carrier for PG synthesis. Specifically, the Δ*sanA* Δ*wecA* strain demonstrated heightened SDS-EDTA sensitivity, activation of the Rcs stress response, and increased cell length. Further investigation tied the SDS-EDTA sensitivity to increased lipid II available for PG synthesis. Spontaneous suppressor mutants of this phenotype harbored point mutations in *prc*, which encodes tail specific protease, or *ftsI*, which encodes the cell division DD-transpeptidase, a target of Prc. We focused on the *ftsI* mutations and demonstrated that the *ftsI* mutations had increased cell length but nevertheless enhanced PG incorporation at the septum compared to the Δ*sanA* mutant, returning PG incorporation to wild-type levels. Moreover, other mutations affecting septal PG synthesis, but not divisome assembly, also suppressed the SDS-EDTA sensitivity. These findings suggest that, in the absence of SanA, increased lipid II availability perturbs the balance between septal PG synthesis, lateral PG elongation, and other envelope biogenesis pathways, which leads to increased OM permeability.

IMPORTANCE

The Gram-negative cell envelope is a barrier that protects the cell from environmental stress. Therefore, the synthesis of each layer of this envelope needs to be closely coordinated throughout growth and division. Here, we investigated SanA, a protein in *Escherichia coli* K-12 that affects envelope permeability under cellular stress, including nutrient limitation and high temperature. We found that SanA plays a key role in maintaining the permeability barrier when precursor levels for peptidoglycan (PG) synthesis are elevated, linking envelope integrity to balanced septal PG production during cell division. Our results suggest that SanA modulates substrate availability to preserve envelope function, and that in its absence, imbalanced substrate flux to septal PG synthesis disrupts septum formation and compromises barrier integrity.

INTRODUCTION

Gram-negative bacteria have a multi-layered envelope consisting of the inner or cytoplasmic membrane (IM), the aqueous periplasm containing the peptidoglycan (PG) cell wall, and the outer membrane (OM) (1). While the IM is a phospholipid bilayer, the lipid portion of the OM is composed of phospholipids in the inner leaflet and primarily lipopolysaccharide (LPS) in the outer leaflet. The cell envelope is tightly interconnected. For instance, the OM and PG are physically connected by an OM lipoprotein (Lpp) and/or outer membrane proteins (OMPs) that bind to PG (e.g., OmpA) (2, 3). Properly composed, the cell envelope prevents entry of toxic molecules, resists osmotic and physical stress, and facilitates cell growth, all while allowing nutrients to enter the cell and metabolic waste to be expelled (1, 4-8). The envelope prevents entry of molecules that are either large or hydrophobic, making Gram-negative bacteria intrinsically resistant to many clinically used antibiotics and greatly complicating discovery of new antimicrobials (4). Yet, the pathways responsible for envelope biogenesis and the regulatory interactions between the biogenesis pathways are promising targets for development of new antimicrobials (9-17).

While synthesis of each envelope component depends on a separate pathway (1, 18-21), complex regulatory interactions between these pathways are essential to coordinate growth of the layers (22). For instance, in *Escherichia coli*, halting fatty acid synthesis, required for phospholipid and LPS synthesis, also causes PG synthesis to cease (23). Conversely, excess phospholipid synthesis can trigger PG expansion. In this case, phospholipids in the outer leaflet of the OM act as a signal to activate MepS, an endopeptidase that breaks PG crosslinks to allow new PG insertion (24). Interestingly, cell growth can be separated from new PG synthesis and endopeptidase activity, which can continue in the absence of PG synthesis until the point of cell lysis (25). YhcB (a.k.a. ZapG, LapD) has been suggested to interact with both the elongasome (the machinery responsible for PG synthesis during cell growth) and with the complex responsible for regulating LPS synthesis and to play a role in balancing the production of LPS, PG, and phospholipids (26-28). However, recent results have demonstrated that YhcB interacts directly with AccA, a member of the complex that

initiates fatty acid synthesis, and the phenotypes of a *yhcB* mutant are directly tied to an

increase in phospholipid production due to excess fatty acid synthesis (29, 30). This

suggests the effects on PG and LPS production are a secondary consequence of the

complex regulatory interactions between cell envelope pathways. The OM's physical

strength also plays a role in orienting PG synthesis, and LPS mutations strengthening

the OM can restore rod shape in mutants deficient in activation of the elongasome (31).

In *Pseudomonas aeruginosa*, MurA, which catalyzes the first committed step in PG

biosynthesis, directly interacts with LpxC, which catalyzes the first committed step in

LPS biosynthesis, causing stimulation of LpxC activity (32).

Interactions in biosynthesis can also occur through shared synthesis substrates. For

instance, most extracytoplasmic glycans are synthesized on an isoprenoid carrier,

undecaprenyl-phosphate (Und-P), and mutations that affect biosynthesis of one glycan

can disrupt synthesis of others due to sequestration of Und-P. Enterobacterial common

antigen (ECA) is an extracytoplasmic glycan found in the Enterobacterales envelope

and consists of a repeating, invariant trisaccharide unit that is found surface exposed as

the headgroup of a phospholipid or attached to LPS in place of O-antigen, as well as

cyclized in the periplasm (33). Mutations in later steps of ECA synthesis cause cell

shape defects and envelope stress response activation, presumably through inhibition

of PG synthesis (34-39). Loss of WecA, which catalyzes the first step in ECA synthesis,

causes an approximately 2-fold increase in free Und-P levels (40). Loss of later steps in

ECA biosynthesis cause Und-P levels to decrease compared to Δ*wecA*, with later

mutations causing larger decreases.

The need for coordination in synthesis of the cell envelope layers becomes more acute

during conditions that perturb synthesis of envelope components such as nutritional

stress or cell division. PG synthesis during cell division is carried out by a highly

regulated complex known as the divisome (41, 42). The divisome is assembled on the

FtsZ ring which consists of FtsZ, FtsA, and ZipA in *E. coli*. FtsK is then recruited,

followed by FtsQ, FtsL, and FtsB. Once this complex is complete, FtsW, FtsI, and FtsN

then join the complex. FtsA plays a key role in recruiting the downstream components of

the complex, while FtsN recruitment acts as a signal to begin septal PG synthesis (41,

42). FtsW and FtsI (PBP3) are responsible for septal PG synthesis with FtsW acting as

the transglycosylase linking the glycan backbone of PG and FtsI acting as the DD-

transpeptidase to link the peptide stems of the PG.

During septal PG synthesis, the OM invaginates with the PG allowing coordinated

separation of the cell envelope (42). This coordination is accomplished through a

protein, CpoB, that interacts with LpoB and PBP1B at the septum and with the Tol-Pal

complex, which has IM and OM components and contacts PG (43-45). Beyond its

importance for OM invagination, Tol-Pal, as well as two OM lipoproteins (NlpD and

YraP), are involved in the activation of amidases that split the septal PG during cell

division, ensuring the coordination of OM invagination with cell division (46, 47).

Interestingly, mutations inactivating NlpD lead to increased sensitivity to the detergent

SDS (sodium dodecyl sulfate) in combination with EDTA, which disrupts interactions

between LPS molecules, suggesting that the OM permeability barrier is disrupted by

loss of cell division amidase function (46). To complete OM invagination, an increase in

OM synthesis is required. Folding of OMPs into the OM by the β-barrel assembly

machine (BAM) is activated by immature PG allowing insertion of new OMPs into the

OM at the cell septum (48), and so allowing balanced OM synthesis during cell division.

We previously investigated changes in OM permeability during stationary phase initiated

by carbon limitation (49). We found that resistance to SDS during carbon limitation did

not rely on efflux pumps as it did in exponentially growing cells. Instead, a RpoS (the

master regulator of stationary phase)-dependent change in OM permeability occurs,

which requires the presence of three genes, *sanA*, *dacA* (PBP5)*,* and *yhdP* (49). YhdP

is a presumptive inter-membrane phospholipid transporter (50-53), the loss of which

causes increased OM permeability in the presence of ECA (54). PBP5 is a

carboxypeptidase that cleaves D-alanine from the PG pentapeptide, regulating the

amount of substrate available for transpeptidases that crosslink PG (55). Loss of PBP5

can prevent the maturation of PG that results in mid-cell insertion of new OMP proteins

(48). Together, these findings suggest that envelope coordination is important for the

stationary phase membrane permeability. However, the function of *sanA* has remained

unknown.

SanA is a bitopic IM protein with one N-terminal transmembrane helix and a periplasmic

DUF218 domain (56, 57). In *E. coli* K-12, *sanA* was first identified as a multicopy

suppressor of vancomycin sensitivity for a strain that had a large gene deletion including

sanA (58). It was also found to be involved in vancomycin resistance at high

temperature (i.e., 43 °C) (58). In *Salmonella enterica* serovar Typhimurium, SfiX, a

SanA homolog, is necessary for filamentation of a mutant overexpressing *hisFH* (His[C])

and also affects vancomycin resistance at high temperature (59). Recent work

demonstrated a decrease in hydrophobicity and charge of the *Salmonella* cell surface

and a corresponding change in envelope permeability in a *sfix* (*sanA*) mutant (60).

Here, we investigated the envelope permeability phenotypes caused by SanA's loss and

found a genetic interaction with ECA synthesis. Mutations expected to increase

availability of Und-P caused the *sanA* mutant to become sensitive to SDS-EDTA and

activated the Rcs (Regulator of Capsule Synthesis) envelope stress response. We then

selected suppressors of SDS-EDTA sensitivity and isolated partial loss-of-function

mutations in *ftsI* (PBP3) and in *prc*, a protease involved in the maturation of FtsI (61,

62). This finding suggests a connection between cell division and OM permeability.

Testing of additional cell division mutations demonstrated that mutations in septal PG

synthesis but not divisome assembly suppressed the OM permeability defect. Our

results tie the envelope permeability phenotype to lipid II availability and support a

model in which SanA modulates the availability of lipid II for septal PG synthesis. We

propose that, when lipid II levels are high such as during certain cellular stresses, lack

of balance between septal PG synthesis and other envelope biogenesis pathways alters

OM composition and envelope permeability.

RESULTS

Loss of SanA increases membrane permeability under cellular stress conditions.

We previously reported that, when treated during stationary phase reached from carbon

limitation, a Δ*sanA* strain is sensitive to the detergent SDS, as assayed by colony

forming units (CFU) 24 hours after treatment (49). When we examined the effect of

Δ*sanA* on SDS sensitivity in carbon-limited stationary phase at shorter time scales, we

found that the decrease in CFU began within the first hour of treatment and continued
through 7 hours (**Figure 1A**). In contrast, SDS-treated wild-type cells maintained
viability similar to that of untreated cells throughout the 7-hour treatment. An insertion
mutation in *sanA* was reported to increase vancomycin sensitivity at high temperatures
(43 °C) (58). To confirm this phenotype in a Δ*sanA* mutant, we performed efficiency-of-
plating assays (EOPs) with wild-type and Δ*sanA* cells carrying either an empty vector or
a low copy number plasmid expressing *sanA*. When treated with vancomycin at 43 °C,
we found the Δ*sanA* strain was more sensitive to vancomycin and this phenotype could
be complemented (**Figure 1B**). To confirm if we observed a true change in resistance
(i.e., the concentration of antibiotic that can kill cells), we performed minimum inhibitory
concentration (MIC) assays and found that the MIC of vancomycin for the Δ*sanA* strain
was significantly lower than that of wild type (**Figure 1C**).

Vancomycin is a large glycopeptide antibiotic for which entry into the cell is greatly
impeded by LPS in the OM (4, 63, 64) and anionic detergents such as SDS are
excluded from the cell by the strong hydrophilic network of lateral interactions between
LPS molecules in the OM (1, 4). Given the importance of LPS for exclusion of these
molecules, we examined the Δ*sanA* strain under various growth conditions for possible
changes in the lipid A component of LPS. Lipid A anchors LPS in the OM and is critical
for maintaining the OM permeability barrier. We observed a large increase in hepta-
acylated lipid A in the Δ*sanA* mutant in cells grown in defined MOPS media with glucose
but not in cells grown in LB media at 37 or 43 °C (**Figure 1D, S1**). The hepta-acylated
lipid A levels were returned to normal by plasmid-based complementation of *sanA*.
Hepta-acylated LPS is formed when PagP, an OMP, transfers a saturated acyl chain to
LPS from a phospholipid mislocalized to the outer leaflet of the OM (65). Thus,
increased hepta-acylated LPS indicates a loss of OM asymmetry causing increased
phospholipids on the cell surface (66). In support of this loss of asymmetry,
overexpression of *pldA*, an OM phospholipase that degrades phospholipids mislocalized
to the OM (67, 68), decreased the amount of hepta-acylated lipid A in the Δ*sanA* strain
(**Figure 1D, S1**). However, *pldA* overexpression did not restore the Δ*sanA* mutant's
vancomycin resistance at 43 °C or SDS resistance during carbon limitation (**Figure S2**),

suggesting the loss of OM asymmetry was secondary to another defect that caused OM
permeability.

SanA only affects vancomycin resistance at high temperatures (58) and *sanA* is
predicted to have a σ^H promoter (56, 57). Therefore, we investigated whether the loss of
SanA affected heat shock resistance in carbon-limiting conditions. Compared to the
wild-type strain, the Δ*sanA* strain showed a small but significant decrease in survival at
57 °C (**Figure 1E**). This increased sensitivity was not shared with deletion of the other
gene in *sanA*'s operon, *yeiS* (**Figure S3**). Overall, these data demonstrate that SanA
plays an important role in maintaining the OM permeability barrier specifically in
conditions of cellular stress such as carbon limitation and high temperatures.

***sanA* has genetic interactions with ECA mutations that induce PG stress.** To
investigate how SanA affects membrane permeability, we made double mutants of *sanA*
and non-essential members of various envelope biosynthesis pathways. A paralog of
sanA, *elyC*, has been shown to have genetic interactions with ECA mutations (69) and
is involved in regulating the synthesis of one form of ECA (70). In addition, in the same
screen as *sanA*, we identified another gene, *yhdP*, which encodes a putative
intermembrane phospholipid transporter (50-53), and *yhdP*'s membrane permeability
phenotypes are dependent on the presence of ECA (49). Therefore, we decided to
assay genetic interactions between ECA biosynthesis genes and *sanA*.

We found that Δ*sanA* had a synthetic SDS-EDTA sensitivity, even at 37 °C in LB media,
when paired with a deletion of *wecA*, the first gene in the ECA synthesis pathway
(**Figure 2A**). EDTA breaks the network of interactions between LPS molecules,
increasing cells susceptibility to SDS (71). Intriguingly, when we tested for genetic
interactions between *sanA* and *wecE*, which encodes a protein involved in synthesis of
the third sugar in the ECA trisaccharide, we found that Δ*wecE* causes SDS-EDTA
sensitivity and this sensitivity is suppressed by Δ*sanA* (**Figure 2A**). These genetic
interactions were apparent at 43 °C as well as 37 °C (**Figure S4A**) and the synthetic
SDS-EDTA phenotype of the Δ*sanA* Δ*wecA* mutant could be complemented with
plasmid-based expression of *sanA* (**Figure S4B**). The difference in phenotype between
Δ*wecA* and Δ*wecE* suggested that interactions between ECA and other biosynthetic

pathways are important for this phenotype, not the loss of ECA itself. Specifically, ECA
synthesis uses the same isoprenoid carrier (Und-P) as PG synthesis (33). Therefore, we
wondered if PG synthesis stress caused by sequestration of the isoprenoid carrier might
be important for the genetic interaction with *sanA*.

We proceeded to compare the phenotypes of Δ*sanA* with deletion of the three
transglycosylases in the ECA synthesis pathway. The PG stress caused by deletion of
each subsequent transglycosylase is more pronounced than the previous (37, 40, 69,
72). Δ*wecA* does not cause apparent PG stress, Δ*wecG* causes minimal PG stress, and
Δ*wecF* causes substantial PG stress as evidenced by envelope stress response
activation, permeability defects, and cell shape defects, although these defects are not
as strong as those of Δ*wecE* (37, 40, 69, 72). When we tested these mutants, we
indeed found the Δ*sanA* Δ*wecA* strain to have the highest SDS-EDTA sensitivity, while
the Δ*sanA* Δ*wecG* strain had slight sensitivity and the Δ*sanA* Δ*wecF* strain showed
similar SDS-EDTA sensitivity to that of the Δ*sanA* strain (**Figure 2B**). These data
indicated that, counterintuitively, the envelope permeability of a Δ*sanA* mutant is
increased by mutations that increase isoprenoid carrier availability.

ECA mutants causing PG stress have been shown to activate the Rcs stress response,
which directly responds to LPS defects but can also be activated by PG stress (73-75).
Therefore, we tested Rcs activation using luciferase genes driven by the *rprA* promoter
(76, 77) in single and double deletion mutants of *sanA*, *wecA* and *wecF*. Neither Δ*sanA*
or Δ*wecA* alone caused Rcs activation (**Figure 2C**); however, a Δ*sanA* Δ*wecA* mutant
showed Rcs activation greater than that of the Δ*wecF* or Δ*sanA* Δ*wecF* mutant. The
peak reporter activity in the Δ*sanA* Δ*wecA* strain corresponded to exponential growth.
As these data and the SDS-EDTA data point to interactions with isoprenoid carrier
availability and PG synthesis, we then assayed the cell shape of the Δ*sanA* and Δ*wecA*
single and double mutants. We grew cells for 7-10 generations in exponential phase
before the assay to ensure that cell shape was stable after exiting stationary phase (78,
79). The aspect ratio (length/width) of all mutants was larger than wild type with the
Δ*sanA* Δ*wecA* mutant showing a significant shape change from that of the Δ*sanA* single
mutant (**Figure 2DE**). Overall, these data demonstrate *sanA* has genetic interactions

causing increased permeability and envelope stress with increased isoprenoid carrier
availability.

Spontaneous mutation to *ftsI* and *prc* suppress the Δ*sanA* Δ*wecA* SDS-EDTA
sensitivity. When assaying the SDS-EDTA sensitivity of the Δ*sanA* Δ*wecA* double
mutant, we occasionally noticed the appearance of spontaneous suppressor mutants.
Therefore, we decided to select for these mutants to determine what types of mutations
could suppress this strain's envelope permeability. We plated Δ*sanA* Δ*wecA* strain on a
concentration of SDS-EDTA where it did not grow but the wild-type strain did and
collected 11 suppressor mutants from a total of 7 separate cultures. We grouped these
mutants by their antibiotic resistance patterns to find strains with similar mutations
(**Figure S5**) and then performed whole genome sequencing on representatives of each
group, following up to confirm the other mutants within the group had the same
mutation(s). These suppressor mutants each contained one of four mutations, *ftsI*$_{V98E}$,
ftsI$_{S165P}$, *prc*$_{F365L}$, or *prc*$_{R578C}$ (**Figure 3A**). FtsI is the DD-transpeptidase involved in
septal PG synthesis during cell division (41, 80), while Prc (a.k.a. Tsp) is a protease
known to cleave FtsI's C-terminal 11 amino acids to produce the fully mature protein
(61, 62). As Prc is necessary for FtsI maturation (61), we focused on the *ftsI* mutations
as the more direct suppressors. However, it is possible that the *prc* mutations may act
through an unrelated pathway (i.e., by increasing endopeptidase levels, which has been
recently shown to suppress another Δ*sanA* phenotype) (24, 81). We linked the *ftsI*$_{V98E}$
mutation to a nearby Tn*10* and used this marker to transfer the mutation to the wild-
type, Δ*sanA*, Δ*wecA*, and Δ*sanA* Δ*wecA* backgrounds. After being moved to a clean
background, the *ftsI*$_{V98E}$ mutation suppressed SDS-EDTA sensitivity in the Δ*sanA*
Δ*wecA* strain showing it was sufficient for suppression (**Figure 3B**). We were not
successful in moving the *ftsI*$_{S165P}$ mutation to other strain backgrounds; however, we
used P1*vir* transduction to remove the mutation from the suppressor strain and this
prevented the SDS-EDTA resistance of the strain (**Figure 3C**). Although we assayed the
effect of the *ftsI*$_{V98E}$ mutation on the vancomycin sensitivity of a Δ*sanA* strain at 43 °C,
the *ftsI*$_{V98E}$ mutation also caused vancomycin resistance in a wild-type background
making it difficult to assess whether its effect on the Δ*sanA* vancomycin sensitivity was
specific (**Figure S6**).

We then asked whether the *ftsI* mutations could suppress Rcs activation in the Δ*sanA*
Δ*wecA* strain. We found that the *ftsI*$_{S165P}$ mutation caused a growth delay and
maintained Rcs activation, while the *ftsI*$_{V98E}$ mutation brought Rcs activation in the
Δ*sanA* Δ*wecA* mutant back to nearly wild-type levels without a significant effect on
growth (**Figure 3D**). We made an AlphaFold model (82) of a partial divisome containing
FtsI, FtsW, FtsL, FtsB, and FtsQ, which corresponds well to the crystal structure of this
complex in *Pseudomonas aeruginosa* (83), and mapped locations of the suppressor
mutations onto this predicted structure (**Figure 3E**). FtsI has two periplasmic domains, a
pedestal domain thought to control activation of FtsW and mediate interactions with FtsL
and FtsB and the catalytic transpeptidase domain (80, 84). The Prc cleavage site in FtsI
is in an unstructured region adjacent to the transpeptidase domain, while the V98E and
S165P mutations are in the pedestal domain (**Figure 3E**). Neither of these mutations
has been previously reported and their effect on FtsI activity is yet to be determined.
Nevertheless, the consistent and repeated isolation of mutations in *ftsI* and *prc*
demonstrated that the envelope defect leading to SDS-EDTA sensitivity in the Δ*sanA*
Δ*wecA* strain is related to cell division.

Suppressor mutations in *ftsI* cause partial loss of FtsI function. As the Δ*sanA* and
Δ*sanA* Δ*wecA* strains showed a cell shape with a higher aspect ratio than wild-type cells
(**Figure 2D**) and the suppressor mutations we found are in cell division genes, we then
assayed the cell shape of the suppressor strains. Cells from both *ftsI* mutant strains
were significantly longer than the Δ*sanA* Δ*wecA* parent and had a greater aspect ratio
(**Figure 4AB**). In contrast, the aspect ratio of the *prc* mutants was indistinguishable from
that of the Δ*sanA* Δ*wecA* parent. The longer length of the *ftsI* mutants suggests that
they have a partial loss of *ftsI* function. Therefore, we tested the transpeptidase activity
of the mutants through resistance to aztreonam, a β-lactam antibiotic that is highly
specific to FtsI (85).

To show the correlation between FtsI activity and aztreonam resistance, we tested
ftsI23 (Y281D), a temperature sensitive *ftsI* allele that has reduced activity even at
permissive temperatures and can slow the rate of closure of the septum (86-88), and
ftsN overexpression which has been shown to result in increased divisome activity and

to suppress partial loss of function mutations in both divisome assembly and septal PG

synthesis (89-91). Given the temperature sensitive phenotype of *ftsI23,* we assayed

aztreonam MICs at both 30 and 37 °C. At 30 °C, *ftsN* overexpression caused a

significant increase in aztreonam MIC in the Δ*sanA* Δ*wecA* background (**Figure 4C**),

while none of the *ftsI* mutations caused a significant change in MIC. However, at 37 °C,

ftsI23 and *ftsN* overexpression caused a significant decrease and increase in MIC,

respectively (**Figure 4C**). Thus, FtsI activity positively correlates to aztreonam

resistance. At 37 °C, *ftsI*$_{V98E}$ decreased aztreonam's MIC compared to its Δ*sanA* Δ*wecA*

parent, while *ftsI*$_{S165P}$ did not cause a significant change. These data suggest that, while

ftsI$_{V98E}$ may have a direct or indirect decreased transpeptidase activity contributing to

increased cell length, *ftsI*$_{S165P}$ does not have a significant effect on transpeptidase

activity. Therefore, the effect of these mutations on protein-protein interactions, rather

than transpeptidase activity, may be more important for suppression.

To test if suppression of Δ*sanA* Δ*wecA* membrane permeability was unique to *ftsI*

mutations, we tested the effect of several gain- and loss-of-function mutations in

divisome activity on the SDS-EDTA sensitivity of the Δ*sanA* Δ*wecA* strain. In contrast to

ftsI$_{V98E}$ that suppressed the SDS-EDTA sensitivity of the Δ*sanA* Δ*wecA* strain at both 30

and 37 °C (**Figure 4D**), *ftsI23* suppressed only at 30 °C, suggesting there may be a

range of function loss for *ftsI* that can suppress envelope permeability. Neither *ftsA**

(R286W) (92, 93) nor *ftsN* overexpression, both of which are activating mutations,

suppressed the SDS-EDTA sensitivity of the Δ*sanA* Δ*wecA* mutant (**Figure 4D**). In fact,

the Δ*sanA* Δ*wecA* strain carrying the *ftsA** allele was more SDS-EDTA sensitive than

Δ*sanA* Δ*wecA* parent. Overall, these data demonstrate that other loss of function

mutations in *ftsI* can also suppress the Δ*sanA* Δ*wecA* envelope permeability.

Δ*sanA* Δ*wecA* envelope permeability is linked to septal PG synthesis. To test

whether SanA might influence the timing of Z ring assembly, the first step in cell division,

we expressed GFP-labeled FtsZ (94) and failed to observe significant differences in the

localization of FtsZ in the wild-type, Δ*sanA*, and Δ*sanA ftsI*$_{V98E}$ strains (**Figure S7AB**).

Moreover, the *ftsI*$_{V98E}$ mutation did not change FtsN localization in a Δ*sanA* Δ*wecA*

background (**Figure S7CD**). Therefore, we examined other cell division mutations to

determine which components of the divisome, its assembly and/or septal PG synthesis,
were important for suppression of SDS-EDTA sensitivity in the Δ*sanA* Δ*wecA* strain. We
examined the effect of three temperature sensitive loss-of-function mutations in
divisome assembly *ftsZ84* (G105S) (95), *ftsA12* (A188V) (96, 97), and *ftsA27* (S195P)
(90, 98), as well as a gain-of-function mutation, *ftsL** (E88K) (99, 100), and a
temperature sensitive loss-of-function mutation in septal PG synthesis, *ftsQ1* (E125K)
(96, 101). Given the temperature sensitivity caused by these mutations we performed
the experiment at 30 °C. None of the mutations affected SDS-EDTA sensitivity on their
own except *ftsQ1*, which showed sensitivity similar to that of the Δ*sanA* Δ*wecA* strain
(**Figure 5A**). Interestingly, only *ftsQ1* suppressed the SDS-EDTA sensitivity of the
Δ*sanA* Δ*wecA* strain (**Figure 5B**), demonstrating cross suppression between these
mutations. In addition, we found that the *ftsZ84*, *ftsA12*, and *ftsL** mutations cause SDS-
EDTA sensitivity in the Δ*sanA* background (**Figure 5C**). Overall, these data demonstrate
that the increased envelope permeability of the Δ*sanA* Δ*wecA* strain is suppressed by
septal PG synthesis mutants but not by mutations in divisome assembly. Instead,
divisome assembly mutants can increase the envelope permeability of a Δ*sanA* strain.

To directly evaluate the effect of *sanA* and *ftsI*$_{V98E}$ on septal PG synthesis, we incubated
cells with BODIPY-FL-3-amino-D-alanine (BADA) for 10 minutes to label newly
synthesized PG and compared PG incorporation at center-cell regions (center line ±5%
of cell length) vs. the whole cell in dividing and non-dividing cells (**Figure 6**). In both
dividing and non-dividing cells, Δ*sanA* mutants exhibited a lower ratio of center-
localized to total PG incorporation than the wild type, suggesting a spatial redistribution
of PG synthesis. In non-dividing cells, there was no difference between the Δ*sanA* and
Δ*sanA ftsI*$_{V98E}$ strain. However, in dividing Δ*sanA ftsI*$_{V98E}$ cells, center cell PG synthesis
was similar to that of wild type and significantly greater than that of Δ*sanA*. These data
suggest that septal PG synthesis is increased with the *ftsI* mutation, which is an
interesting contrast to its longer cell length.

Membrane permeability with loss of SanA is tied to lipid II availability and FtsW
activity. As the *ftsI* suppressor mutations reside in the FtsI pedestal domain that
interacts with FtsW and the *ftsI*$_{V98E}$ mutation increases septal PG synthesis, we

investigated whether SanA's envelope permeability phenotypes relate to FtsW activity.
We transferred a previously described (102) FtsW depletion system that contains an
arabinose-inducible *ftsW* expressing plasmid and chromosomal Δ*ftsW* to our strains.
We found that *ftsI*$_{V98E}$ could suppress the Δ*sanA* Δ*wecA* strain regardless of arabinose
concentration (**Figure 7A**). However, *ftsI*$_{V98E}$ in a wild-type background conferred
resistance to arabinose depletion, allowing cells to grow at a lower concentration of
arabinose than is tolerated by the parent strain, suggesting increased FtsW activity
(**Figure 7A**). We did not observe the same increase when *ftsI*$_{V98E}$ is expressed in either
the Δ*sanA* or Δ*sanA* Δ*wecA* background, suggesting that loss of SanA may increase the
activity of FtsW necessary for viability.

Given *sanA*'s genetic interaction with mutations altering isoprenoid carrier availability,
we further investigated whether the availability of lipid II, the substrate for PG synthesis,
was involved in SanA's envelope permeability phenotype. A chromosomal depletion
strain for MurJ, the lipid II flippase (102, 103), where the promoter for *murJ* was
replaced with the arabinose-inducible P$_{BAD}$ promoter has been described (103). We
transduced this promoter into our strains and found MurJ depletion did not prevent
suppression of Δ*sanA* Δ*wecA* SDS-EDTA sensitivity by *ftsI*$_{V98E}$ (**Figure 7B**). To test if
increasing lipid II by overproducing MurA, which catalyzes the first committed step in
PG synthesis (104), would affect OM permeability, we overexpressed *murA* in a Δ*sanA*
strain. We observed SDS-EDTA sensitivity that phenocopied that observed in a Δ*sanA*
Δ*wecA* strain and this sensitivity could be suppressed by *ftsI*$_{V98E}$ (**Figure 7C**).
Furthermore, overexpression of *murA* decreased the suppression observed in a Δ*sanA*
Δ*wecA* *ftsI*$_{V98}$ strain. These data demonstrate that multiple mutations expected to
increase lipid II availability lead to SDS-EDTA sensitivity in the absence of SanA and
that this sensitivity can be suppressed through mutations in septal PG synthesis genes.

A predicted pocket is important for SanA's function in envelope permeability. To
further investigate how SanA's structure affects its function, we modeled SanA's
structure using AlphaFold 3 (82). Using the predicted structure as a guide, we
constructed point mutations to several conserved SanA residues (**Figure 8A**).
Interestingly, SanA's predicted structure has a pocket that faces the direction of the

transmembrane helix and an amphipathic helix perpendicular to the transmembrane helix that may sit in the IM. In an attempt to validate this prediction, we first made charge switch mutation pairs of residues interacting with this amphipathic helix. Whereas two of the individual mutations (R152E and E182R) resulted in loss of *sanA* function (**Figure 8B**) without affecting protein stability (**Figure 8C**), a double mutation (R152E E182R) did not restore function. Mutations of another charge pair D192 and R198 resulted in loss of protein stability, complicating analysis (**Figure 8BC**). Therefore, we could not validate the interactions of the amphipathic helix. However, several structural models (AlphaFold 2, AlphaFold 3, and I-TASSER (82, 105, 106)) all predicted a pocket that contains several conserved residues (**Figure 8A**). Mutating conserved hydrophilic residues in the pocket resulted in loss of SanA function (**Figure 8D**), while generally maintaining protein stability (**Figure 8C**). Previous work also found the H149A mutation to cause loss of function (81). While charged residues are predicted near the outer edge of the pocket, the inner surface of the pocket is composed of hydrophobic residues (**Figure 8A**). Mutations to residues on the surface of the pocket (L51, G52) resulted in loss of function with minimal or no effect on protein stability, while mutations of buried residues did not change SanA function (**Figure 8CE**). Finally, we made several mutations to surface residues of SanA including those on the side of SanA predicted to face the IM and the side predicted to face away from the IM (**Figure 8A**). Generally, these mutations caused less severe phenotypes than mutations to SanA's pocket residues (**Figure 8F**). Overall, these structure-function analyses suggest that SanA's pocket may be important for binding or enzymatically altering an IM molecule.

DISCUSSION

Here, we set out to understand how SanA affects the cell's envelope permeability barrier. We confirmed that a Δ*sanA* strain has envelope permeability phenotypes that occur under carbon limitation and high temperatures. Additionally, we discovered that a Δ*sanA* Δ*wecA* strain, which has heightened isoprenoid carrier levels due to a lack of ECA synthesis (40), had a pronounced, synthetic SDS-EDTA sensitivity and that this sensitivity could be suppressed by partial loss of function mutations to either *ftsI* or to

prc. When investigating the suppression phenotype, we found that mutations to septal
PG synthesis machinery, but not early divisome assembly, could suppress the Δ*sanA*
Δ*wecA* SDS-EDTA sensitivity. Loss of SanA resulted in decreased PG incorporation in
the region of the septum and this was suppressed in dividing cells by the *ftsI* suppressor
mutation. Moreover, we observed that overexpressing *murA* in a Δ*sanA* strain
phenocopied Δ*sanA* Δ*wecA*, indicating increased lipid II is important for the SDS-EDTA
phenotype. Finally, at least one of the *ftsI* suppressor mutations we isolated, *ftsI*$_{V98E}$,
resulted in increased FtsW activity, which likely results in the increased septal PG
synthesis.

Based on these findings, we propose a model which provides a parsimonious
explanation for our results (**Figure 9**). During normal growing conditions, precursors are
carefully allocated to allow for balanced synthesis of lipid II and other envelope
constituents (**Figure 9A**). In our model, we reason that, under certain cellular stress
conditions, the amount of lipid II available for PG synthesis increases. For example,
enzymes involved in lipid II synthesis might have higher activity rates at higher
temperature leading to increased lipid II synthesis. Alternatively, under carbon-limited
conditions, decreased growth and the lower surface area of stationary phase cells could
lead to higher lipid II levels. The increase in substrate availability creates a potential
imbalance of septal and elongation PG synthesis, which is prevented by SanA (**Figure
9B**). When SanA is absent, lipid II use for lateral PG elongation and septal PG synthesis
is imbalanced and this imbalance ultimately leads to envelope permeability (**Figure 9C**).
This model would also suggest loss-of-function mutations in early divisome assembly
increase the membrane permeability of a Δ*sanA* strain by decreasing the number of
active septal PG synthesis complexes and so further decreasing septal PG synthesis
rates. In SanA's absence, mutations that restore septal PG synthesis rates restore
balance in septal PG synthesis and so the envelope permeability barrier.

In support of this model, a paper (81) was published while this manuscript was in
preparation that identified a suppressive phenotype of Δ*sanA* for the growth of several
cell division mutants including *ftsI23* and found that PG incorporation was increased in a
Δ*sanA* strain (81). Although the paper examined different conditions from our study, it

also found synthetic interactions between early genes in ECA synthesis and Δ*sanA*,
specifically for growth in low osmolarity (81). Furthermore, the paper demonstrated that
Δ*sanA* phenotypes could be suppressed by endopeptidase overexpression, which
provides a possible explanation for our *prc* suppressors, which could increase
endopeptidase levels as well as prevent the cleavage of FtsI (24, 81). From these
findings it was unclear whether SanA's phenotypes related to septal PG synthesis or PG
synthesis more broadly. However, our work clearly demonstrates that the membrane
permeability phenotype, which occurs when increased lipid II levels are coupled with the
absence of SanA, specifically relates to septal PG synthesis and that changes to the
elongasome (involved in cell lengthening) are not necessary to prevent permeability.

How SanA maintains septal PG synthesis with different lipid II levels remains an
unanswered question. The recent paper speculates that SanA may have enzymatic
activity, based on phenotypes of several *sanA* point mutations (81). We agree that
predicted structures of SanA suggest it has a potential binding pocket and possibly
enzymatic activity. Here, we present a large panel of *sanA* point mutations that
demonstrate conserved residues within this pocket are important for SanA function. If
SanA does have enzymatic activity, it is possible it alters lipid II in some way.
Alternatively, SanA may bind to and sequester lipid II from PG elongation. However,
SanA molecules in the cell are 10 to 50-fold fewer than lipid II under normal growth
conditions (107, 108). Thus, a significant increase in SanA levels under activating
conditions would be necessary for this to occur. Lipid II synthesis is regulated through
dimerization of MraY which binds to excess lipid II, inhibiting its activity and lipid II
synthesis (109). Given this, it is possible SanA is involved in this regulatory pathway.
However, it's perhaps more likely that SanA directly controls the availability of lipid II for
septal PG synthesis rather than overall lipid II levels (see **Figure 9**). For instance, SanA
could feed lipid II to the septal PG synthesis complex and ensure correct substrate
availability to balance septal and lateral PG synthesis. This hypothesis could explain the
slight increase in cell length observed in the Δ*sanA* strain, where cells elongate further
before division. Finally, a paper was published during revision of this manuscript that
suggests direct or indirect physical interaction between SanA and PBP1b (110). Thus,

SanA could act at least in part by modulating PBP1b activity. However, differentiating
these possibilities will require future investigation.

FtsI's cytoplasmic region and transmembrane helix are important for its recruitment to
the Z-ring through interaction with FtsW (80). The periplasmic portion of FtsI contains
the pedestal domain, which is important for interacting with other members of the septal
PG synthesis complex, and the catalytic domain that acts as the transpeptidase during
septal PG synthesis (80). Both FtsI variants that we identified, FtsI^{V98E} and FtsI^{S165P}, are
in the pedestal domain of FtsI. Notably, this domain is proposed to recruit FtsN (the final
activating protein for the complex), to interact with FtsW (the transglycosylase) and to
control its activation, and to control FtsI transpeptidase activity (80, 84, 111). It is
interesting then that although both FtsI^{V98E} and FtsI^{S165P} resulted in longer cells,
suggesting a partial loss-of-function, FtsI^{V98E} also increased PG incorporation in the
area of the septum and made wild type cells resistant to FtsW depletion, suggesting
increased septal PG synthesis. Thus, these mutations may affect both FtsI and FtsW
activity to alter septal PG synthesis. Mutations to residues near S165 (R166 and R167)
have been shown to either cause constitutive activation of FtsW or loss-of-function in
FtsI transpeptidase activity (80, 112, 113) so it may be that S165P causes functional
changes to both FtsI and FtsW as well. Further investigation of these mutants may lead
to greater insight into the complex regulatory interactions carried out by the pedestal
domain of FtsI.

Finally, it is worth considering how an imbalance in septal versus lateral PG synthesis
could lead to increased envelope permeability. The synthesis of PG and OM
components share many precursors. Therefore, it is possible that an increase in
elongation PG synthesis could limit some OM components, hindering proper OM
synthesis. However, increasing septal PG synthesis should not clear this defect. More
likely, the envelope biosynthesis phenotype is due to a lack of coordination between
different envelope biosynthetic pathways. For instance, OMP biosynthesis has been
shown to increase in the presence of immature PG (48). Therefore, it is possible that
decreased septal PG synthesis leads to an decrease in immature PG, causing reduced
OMP insertion near the septum. It is also possible that there is a defect in OM

invagination caused by the alteration in septal PG synthesis. These questions will be
interesting avenues for future investigations.

MATERIALS AND METHODS

Strains and growth conditions. All strains used in this study are listed in **Table S1**.
Cultures were inoculated, grown at 37°C unless otherwise noted in LB, M63 or MOPS
media, and supplemented with 20mg/liter chloramphenicol, 25mg/liter kanamycin,
10mg/liter tetracycline, or a combination thereof as necessary. M63 was supplemented
with 1mM $MgSO_4$, 50mg/ml thiamine, and 0.2% glucose or 0.05% glucose for carbon-
limitation. Mutations were moved using P1*vir* transduction (114) and resistance
cassettes were removed with the Flp recombinase-FRT system (115). To clone a
plasmid encoding *sanA*, genomic DNA was amplified using the primers pZS21-RBS-
sanA F and *sanA*-pZS21 R (**Table S2**). pZS21 (116) was amplified using pZS21 F and
pZS21-RBS R (**Table S2**). The fragments were assembled using HiFi Assembly Master
Mix (New England Biolabs [NEB]) as per the manufacturer's instructions. Site-directed
mutagenesis was performed by PCR with Q5 polymerase (NEB) and the indicated
primers (**Table S2**). PCR products were digested with DpnI (NEB), purified,
phosphorylated with T4 polynucleotide kinase (NEB), and then self-ligated and
transformed. Mutations were confirmed by sequencing.

Sensitivity assays. Resistance to SDS in stationary phase was assay as previously
described (49). For EOPs, overnight cultures were serially diluted, plated on the
indicated plates, and plates were incubated overnight. For Kirby-Bauer tests, 100 µL of
overnight cultures were added to 3 mL liquified top LB agar and spread on LB agar
plates. Preconditioned antibiotic disks were placed on the agar surface and plates were
incubated overnight. The diameter of the zone of clearance was measured in at least
two directions. For MICs, overnight cultures were normalized to an OD_{600} of 0.1 then
further diluted 1:1000. Diluted cultures were then added to 96-well plate followed by
two-fold dilutions of the indicated antibiotic. Plates were incubated at 37 °C overnight.
Then, OD_{600} was read using a BioTek Synergy H1 plate reader. The lowest
concentration where growth is inhibited was determined to be the MIC.

Lipid A modification assay. Cultures were grown overnight in the indicated media. Cells were then diluted 1:100 in fresh medium containing 2.5 µCi/mL of ^{32}P and grown to OD_{600} 0.6-0.8. LPS was purified and mid-acid hydrolysis was used to liberate lipid A which was visualized using thin-layer chromatography (TLC) as previously described (117).

Suppressor isolation. To isolate suppressors, cultures were serial diluted and plated with 0.5% SDS with 1.0 or 1.1 mM EDTA. Colonies were isolated and suppression was confirmed by EOP. Verified suppressors were categorized based on antibiotic resistance profiles. For a representative of each category and the parent strain, genomic DNA was purified using the DNeasy Blood and Tissue Kit (Qiagen) and provided to SeqCoast Genomics for library preparation (using a DNA Prep Tagmentation Kit (Illumina) and unique dual indexes) and sequencing with an Illumina NextSeq2000 using a 300-cycle flow cell producing 2x150bp reads. A PhiX control was used to optimize base calling. Demultiplexing, read trimming, and run analytics were performed using DRAGEN software version. SeqCoast quality trimmed the reads, mapped them to the MG1655 reference genome (Genebank ID U00096.3) and called variants using Trimmomatic and Breseq software (118-120). Mutations were confirmed in all suppressors using Sanger sequencing. *ftsI* mutations were linked to *zab*-3051:Tn*10* using P1*vir* transduction (121).

P*rprA* reporter activity assay. Overnight cultures carrying the pJW15-P*rprA* reporter plasmid (76, 77) were assayed for luciferase activity as has been previously described (70).

Cell imaging and analysis. Overnight cultures were diluted 1:10,000 and grown to OD_{600} ~0.5 before imaging. Imaging and cell shape analysis were performed as previously described (122, 123). Differential interference contrast (DIC) images of cells on an agarose pad were captured using an Andor iXon Ultra 897 EMCCD camera (effective pixel size 160 nm) on an inverted Nikon Eclipse-Ti™ microscope with a 100✕ 1.49 NA TIRF objective. Cell length and width was determined using a MATLAB (MathWorks, Inc) script (122). Raincloud plots were made and statistics calculated using Raincloud-shiny (124). For protein localization, overnight cultures were diluted to OD_{600} 0.1 and grown to OD_{600} 0.4. GFP-tagged proteins were excited by a 488 nm laser. GFP-

tagged *ftsZ* was expressed from pXY027 (94) (Addgene #98915). Demographs were
constructed using ImageJ plug-in MicrobeJ (125). Cells were aligned at mid-cell and
sorted by their length, and fluorescence intensity was normalized.

For assaying PG synthesis, cells were grown to exponential phase and then incubated
in LB supplemented with 100 µM BODIPY-FL-3-amino-D-alanine (BADA) in LB for 10
minutes at 37 °C. The cells were collected by centrifuging 1 minute at 9,000xg and
washed three times with LB before imaging as above. Images were analyzed using
code available on Zenodo (https://doi.org/10.5281/zenodo.18839357). Briefly,
fluorescence intensity across cells was normalized to their surface area assuming a
cylindrical shape with a width of 1 µm. Normalized single cell fluorescence intensity
profiles were integrated along the cell axis using trapezoidal numerical integration.
Center cell intensity was calculated as the intensity within ±5% of the midpoint of the
cell. Cells were classified as dividing or non-dividing based on the z-score of
fluorescence enrichment at mid cell (±10% of cell length) with a threshold of z>1.0.

Immunoblot Analysis. Immunoblots were performed as previously described (126,
127). The primary antibodies used were M2 αFLAG (Millipore Sigma, 1:50,000 dilution)
and αGroEL (Millipore Sigma, 1:60,000). α-mouse and α-rabbit secondary antibodies
(Prometheus) were used at a 1:100,000 dilution.

ACKNOWLEDGEMENTS

Thank you to the current and former members of the Mitchell Lab as well as Drs. Jolene
Ramsey, Jennifer Herman, Deborah Siegele, Ryland Young (Texas A&M University),
Anna Konovalova (University of Texas Health Science Center Houston), Petra Levin
(Washington University in St. Louis), and Natividad Ruiz (The Ohio State University) for
insights and productive discussions. Thank you to Dr. Srutha Venkatesan for helping
with the initial stages of the cell-length experiments. We would also like to thank Dr.
Thomas Silhavy (Princeton University) for all his assistance with the early stages of this
project. Thank you to Drs. Petra Levin (Washington University in St. Louis), Natividad
Ruiz (The Ohio State University), and Thomas Bernhardt (Harvard University Medical

School) for sharing strains with us to facilitate these experiments and to Dr. Tracy Raivio

(University of Alberta) for providing the pJW15-P*rpra* reporter plasmid and Dr. Jie Xiao

(Johns Hopkins University School of Medicine) for providing pXY027 (Addgene

#98915).

This work was supported by the National Institute of Allergy and Infectious Diseases

under award numbers R01-AI155915 (to A.M.M.) and R01-AI176766 (to M.S.T.), and

the National Institute of General Medical Sciences under award number R01-

GM129000 (to B.N.).

REFERENCES

1. Silhavy TJ, Kahne D, Walker S. 2010. The bacterial cell envelope. Cold Spring

Harbor Perspectives in Biology 2:a000414.

2. Braun V. 1975. Covalent lipoprotein from the outer membrane of *Escherichia coli*.

Biochimica et Biophysica Acta (BBA) - Reviews on Biomembranes 415:335–377.

3. Park JS, Lee WC, Yeo KJ, Ryu K-S, Kumarasiri M, Hesek D, Lee M, Mobashery

S, Song JH, Kim SI, Lee JC, Cheong C, Jeon YH, Kim H-Y. 2012. Mechanism of

anchoring of OmpA protein to the cell wall peptidoglycan of the gram-negative

bacterial outer membrane. The FASEB Journal 26:219–228.

4. Nikaido H. 2003. Molecular basis of bacterial outer membrane permeability

revisited. Microbiology and Molecular Biology Reviews 67:593–656.

5. Henderson JC, Zimmerman SM, Crofts AA, Boll JM, Kuhns LG, Herrera CM,

Trent MS. 2016. The power of asymmetry: Architecture and assembly of the

Gram-negative outer membrane lipid bilayer. Annual Review of Microbiology

70:255–278.

6. Berry J, Rajaure M, Pang T, Young R. 2012. The spanin complex is essential for

lambda lysis. J Bacteriol 194:5667–74.

7. Rojas ER, Billings G, Odermatt PD, Auer GK, Zhu L, Miguel A, Chang F, Weibel

DB, Theriot JA, Huang KC. 2018. The outer membrane is an essential load-

bearing element in Gram-negative bacteria. Nature 559:617–621.

8. Navarro Llorens JM, Tormo A, Martinez-Garcia E. 2010. Stationary phase in
gram-negative bacteria. FEMS Microbiol Rev 34:476–95.
9. MacNair CR, Brown ED. 2020. Outer membrane disruption overcomes intrinsic,
acquired, and spontaneous antibiotic resistance. mBio 11:e01615–20.
10. Lehman KM, Grabowicz M. 2019. Countering gram-negative antibiotic
resistance: Recent progress in disrupting the outer membrane with novel
therapeutics. Antibiotics 8:163.
11. Vetterli SU, Zerbe K, Müller M, Urfer M, Mondal M, Wang S-Y, Moehle K, Zerbe
O, Vitale A, Pessi G, Eberl L, Wollscheid B, Robinson JA. 2018. Thanatin targets
the intermembrane protein complex required for lipopolysaccharide transport in
Escherichia coli. Science Advances 4:eaau2634.
12. Srinivas N, Jetter P, Ueberbacher BJ, Werneburg M, Zerbe K, Steinmann J, Van
der Meijden B, Bernardini F, Lederer A, Dias RLA, Misson PE, Henze H,
Zumbrunn J, Gombert FO, Obrecht D, Hunziker P, Schauer S, Ziegler U, Käch A,
Eberl L, Riedel K, DeMarco SJ, Robinson JA. 2010. Peptidomimetic Antibiotics
Target Outer-Membrane Biogenesis in *Pseudomonas aeruginosa*. Science
327:1010–1013.
13. Steenhuis M, Abdallah AM, de Munnik SM, Kuhne S, Sterk G-J, van den Berg
van Saparoea B, Westerhausen S, Wagner S, van der Wel NN, Wijtmans M, van
Ulsen P, Jong WSP, Luirink J. Inhibition of autotransporter biogenesis by small
molecules. Molecular Microbiology 0.
14. Storek KM, Auerbach MR, Shi H, Garcia NK, Sun D, Nickerson NN, Vij R, Lin Z,
Chiang N, Schneider K, Wecksler AT, Skippington E, Nakamura G, Seshasayee
D, Koerber JT, Payandeh J, Smith PA, Rutherford ST. 2018. Monoclonal antibody
targeting the β-barrel assembly machine of *Escherichia coli* is bactericidal.
Proceedings of the National Academy of Sciences 115:3692–3697.
15. Hart EM, Mitchell AM, Konovalova A, Grabowicz M, Sheng J, Han X, Rodriguez-
Rivera FP, Schwaid AG, Malinverni JC, Balibar CJ, Bodea S, Si Q, Wang H,
Homsher MF, Painter RE, Ogawa AK, Sutterlin H, Roemer T, Black TA, Rothman
DM, Walker SS, Silhavy TJ. 2019. A small-molecule inhibitor of BamA impervious

to efflux and the outer membrane permeability barrier. Proc Natl Acad Sci U S A

116:21748–21757.

16.   Urfer M, Bogdanovic J, Lo Monte F, Moehle K, Zerbe K, Omasits U, Ahrens CH,

Pessi G, Eberl L, Robinson JA. 2016. A peptidomimetic antibiotic targets outer

membrane proteins and disrupts selectively the outer membrane in *Escherichia*

coli. Journal of Biological Chemistry 291:1921–1932.

17.   Steenhuis M, Corona F, ten Hagen-Jongman CM, Vollmer W, Lambin D, Selhorst

P, Klaassen H, Versele M, Chaltin P, Luirink J. 2021. Combining cell envelope

stress reporter assays in a screening approach to identify BAM complex

inhibitors. ACS Infectious Diseases 7:2250–2263.

18.   Konovalova A, Kahne DE, Silhavy TJ. 2017. Outer Membrane Biogenesis. Annu

Rev Microbiol 71:539–556.

19.   Garde S, Chodisetti Pavan K, Reddy M. 2021. Peptidoglycan: Structure,

Synthesis, and Regulation. EcoSal Plus 9:eESP–0010–2020.

20.   May KL, Silhavy TJ. 2017. Making a membrane on the other side of the wall.

Biochimica et Biophysica Acta (BBA) - Molecular and Cell Biology of Lipids

1862:1386–1393.

21.   John E. Cronan J, Rock CO. 2008. Biosynthesis of membrane lipids. EcoSal Plus

3:10.1128/ecosalplus.3.6.4.

22.   Fivenson EM, Dubois L, Bernhardt TG. 2024. Co-ordinated assembly of the

multilayered cell envelope of Gram-negative bacteria. Current Opinion in

Microbiology 79:102479.

23.   Rodionov DG, Ishiguro EE. 1996. Dependence of peptidoglycan metabolism on

phospholipid synthesis during growth of *Escherichia coli*. Microbiology (Reading)

142 (Pt 10):2871–7.

24.   Som N, Reddy M. 2023. Cross-talk between phospholipid synthesis and

peptidoglycan expansion by a cell wall hydrolase. Proceedings of the National

Academy of Sciences 120:e2300784120.

25.   Oldewurtel ER, Kitahara Y, Cordier B, Wheeler R, Özbaykal G, Brambilla E,

Boneca IG, Renner LD, van Teeffelen S. 2023. Cell envelope growth of Gram-

negative bacteria proceeds independently of cell wall synthesis. Embo j

42:e112168.

26.   Goodall ECA, Isom GL, Rooke JL, Pullela K, Icke C, Yang Z, Boelter G, Jones A,

Warner I, Da Costa R, Zhang B, Rae J, Tan WB, Winkle M, Delhaye A, Heinz E,

Collet J-F, Cunningham AF, Blaskovich MA, Parton RG, Cole JA, Banzhaf M,

Chng S-S, Vollmer W, Bryant JA, Henderson IR. 2021. Loss of YhcB results in

dysregulation of coordinated peptidoglycan, LPS and phospholipid synthesis

during *Escherichia coli* cell growth. PLOS Genetics 17:e1009586.

27.   Li G, Hamamoto K, Kitakawa M. 2012. Inner Membrane Protein YhcB Interacts

with RodZ Involved in Cell Shape Maintenance in *Escherichia coli*. ISRN Mol Biol

2012:304021.

28.   Wieczorek A, Sendobra A, Maniyeri A, Sugalska M, Klein G, Raina S. 2022. A

New Factor LapD Is Required for the Regulation of LpxC Amounts and

Lipopolysaccharide Trafficking. Int J Mol Sci 23.

29.   Stanley HM, Trent MS. 2024. Loss of YhcB results in overactive fatty acid

biosynthesis. mBio 15:e00790–24.

30.   Stanley HM, Trent MS. 2026. YhcB mediates growth-phase control of fatty acid

biosynthesis through regulation of acetyl-CoA carboxylase. mBio

doi:10.1128/mbio.03681-25:e0368125.

31.   Fivenson EM, Rohs PDA, Vettiger A, Sardis MF, Torres G, Forchoh A, Bernhardt

TG. 2023. A role for the Gram-negative outer membrane in bacterial shape

determination. Proceedings of the National Academy of Sciences

120:e2301987120.

32.   Hummels KR, Berry SP, Li Z, Taguchi A, Min JK, Walker S, Marks DS, Bernhardt

TG. 2023. Coordination of bacterial cell wall and outer membrane biosynthesis.

Nature 615:300–304.

33.   Rai AK, Mitchell AM. 2020. Enterobacterial common antigen: Synthesis and

function of an enigmatic molecule. mBio 11:e01914–20.

34.   Rick PD, Hubbard GL, Barr K. 1994. Role of the *rfe* gene in the synthesis of the

O8 antigen in *Escherichia coli* K-12. Journal of Bacteriology 176:2877–2884.

35.   Klena JD, Schnaitman CA. 1993. Function of the *rfb* gene cluster and the *rfe*
gene in the synthesis of O antigen by *Shigella dysenteriae* 1. Molecular
Microbiology 9:393–402.
36.   Alexander DC, Valvano MA. 1994. Role of the rfe gene in the biosynthesis of the
Escherichia coli O7-specific lipopolysaccharide and other O-specific
polysaccharides containing N-acetylglucosamine. Journal of Bacteriology
176:7079–7084.
37.   Danese PN, Oliver GR, Barr K, Bowman GD, Rick PD, Silhavy TJ. 1998.
Accumulation of the enterobacterial common antigen lipid II biosynthetic
intermediate stimulates *degP* transcription in *Escherichia coli*. Journal of
Bacteriology 180:5875–5884.
38.   Jorgenson MA, Kannan S, Laubacher ME, Young KD. 2016. Dead-end
intermediates in the enterobacterial common antigen pathway induce
morphological defects in *Escherichia coli* by competing for undecaprenyl
phosphate. Molecular Microbiology 100:1–14.
39.   Castelli ME, Véscovi EG. 2011. The Rcs signal transduction pathway is triggered
by enterobacterial common antigen structure alterations in *Serratia marcescens*.
Journal of Bacteriology 193:63–74.
40.   Eade CR, Wallen TW, Gates CE, Oliverio CL, Scarbrough BA, Reid AJ,
Jorgenson MA, Young KD, Troutman JM. 2021. Making the enterobacterial
common antigen glycan and measuring its substrate sequestration. ACS Chem
Biol 16:691–700.
41.   Cameron TA, Margolin W. 2024. Insights into the assembly and regulation of the
bacterial divisome. Nature Reviews Microbiology 22:33–45.
42.   Levin Petra A, Janakiraman A. 2021. Localization, Assembly, and Activation of
the *Escherichia coli* Cell Division Machinery. EcoSal Plus 9:eESP–0022–2021.
43.   Gray AN, Egan AJF, van't Veer IL, Verheul J, Colavin A, Koumoutsi A, Biboy J,
Altelaar AFM, Damen MJ, Huang KC, Simorre J-P, Breukink E, den Blaauwen T,
Typas A, Gross CA, Vollmer W. 2015. Coordination of peptidoglycan synthesis
and outer membrane constriction during *Escherichia coli* cell division. eLife
4:e07118.

44. Szczepaniak J, Holmes P, Rajasekar K, Kaminska R, Samsudin F, Inns PG,
Rassam P, Khalid S, Murray SM, Redfield C, Kleanthous C. 2020. The lipoprotein
Pal stabilises the bacterial outer membrane during constriction by a mobilisation-
and-capture mechanism. Nature Communications 11:1305.

45. Gerding MA, Ogata Y, Pecora ND, Niki H, De Boer PAJ. 2007. The trans-
envelope Tol–Pal complex is part of the cell division machinery and required for
proper outer-membrane invagination during cell constriction in *E. coli*. Molecular
Microbiology 63:1008–1025.

46. Tsang M-J, Yakhnina AA, Bernhardt TG. 2017. NlpD links cell wall remodeling
and outer membrane invagination during cytokinesis in *Escherichia coli*. PLOS
Genetics 13:e1006888.

47. Yakhnina AA, Bernhardt TG. 2020. The Tol-Pal system is required for
peptidoglycan-cleaving enzymes to complete bacterial cell division. Proceedings
of the National Academy of Sciences 117:6777–6783.

48. Mamou G, Corona F, Cohen-Khait R, Housden NG, Yeung V, Sun D, Sridhar P,
Pazos M, Knowles TJ, Kleanthous C, Vollmer W. 2022. Peptidoglycan maturation
controls outer membrane protein assembly. Nature 606:953–959.

49. Mitchell AM, Wang W, Silhavy TJ. 2017. Novel Rpos-dependent mechanisms
strengthen the envelope permeability barrier during stationary phase. J Bacteriol
199:e00708–16.

50. Grimm J, Shi H, Wang W, Mitchell AM, Wingreen NS, Huang KC, Silhavy TJ.
2020. The inner membrane protein YhdP modulates the rate of anterograde
phospholipid flow in *Escherichia coli*. Proc Natl Acad Sci U S A 117:26907–
26914.

51. Ruiz N, Davis RM, Kumar S. 2021. YhdP, TamB, and YdbH are redundant but
essential for growth and lipid homeostasis of the gram-negative outer membrane.
mBio 12:e02714–21.

52. Douglass MV, McLean AB, Trent MS. 2022. Absence of YhdP, TamB, and YdbH
leads to defects in glycerophospholipid transport and cell morphology in Gram-
negative bacteria. PLOS Genetics 18:e1010096.

53.   Rai AK, Sawasato K, Hill HC, Kozlova A, Sparagna GC, Bogdanov M, Mitchell
AM. 2024. Genetic evidence for functional diversification of gram-negative
intermembrane phospholipid transporters. PLOS Genetics 20:e1011335.

54.   Mitchell AM, Srikumar T, Silhavy TJ. 2018. Cyclic enterobacterial common
antigen maintains the outer membrane permeability barrier of *Escherichia coli* in
a manner controlled by YhdP. mBio 9:e01321–18.

55.   van Heijenoort J. 2011. Peptidoglycan hydrolases of Escherichia coli. Microbiol
Mol Biol Rev 75:636–63.

56.   Karp PD, Ong WK, Paley S, Billington R, Caspi R, Fulcher C, Kothari A,
Krummenacker M, Latendresse M, Midford PE, Subhraveti P, Gama-Castro S,
Muñiz-Rascado L, Bonavides-Martinez C, Santos-Zavaleta A, Mackie A, Collado-
Vides J, Keseler IM, Paulsen I. 2018. The EcoCyc database. EcoSal Plus
8:10.1128/ecosalplus.ESP–0006–2018.

57.   Keseler IM, Mackie A, Santos-Zavaleta A, Billington R, Bonavides-Martínez C,
Caspi R, Fulcher C, Gama-Castro S, Kothari A, Krummenacker M, Latendresse
M, Muñiz-Rascado L, Ong Q, Paley S, Peralta-Gil M, Subhraveti P, Velázquez-
Ramírez DA, Weaver D, Collado-Vides J, Paulsen I, Karp PD. 2016. The EcoCyc
database: reflecting new knowledge about *Escherichia coli* K-12. Nucleic Acids
Research 45:D543–D550.

58.   Rida S, Caillet J, Alix JH. 1996. Amplification of a novel gene, sanA, abolishes a
vancomycin-sensitive defect in *Escherichia coli*. Journal of Bacteriology 178:94–
102.

59.   Mouslim C, Cano AD, Casadesús J. 1998. The sfiX, rfe and metN genes of
Salmonella typhimurium and their involvement in the Hisc pleiotropic response.
Molecular and General Genetics MGG 259:46–53.

60.   Aleksandrowicz A, Kolenda R, Baraniewicz K, Thurston TLM, Suchański J,
Grzymajlo K. 2024. Membrane properties modulation by SanA: implications for
xenobiotic resistance in *Salmonella* Typhimurium. Frontiers in Microbiology 14.

61.   Hara H, Nishimura Y, Kato J, Suzuki H, Nagasawa H, Suzuki A, Hirota Y. 1989.
Genetic analyses of processing involving C-terminal cleavage in penicillin-binding
protein 3 of *Escherichia coli*. J Bacteriol 171:5882–9.

62. Chueh CK, Som N, Ke LC, Ho MR, Reddy M, Chang CI. 2019. Structural Basis
for the Differential Regulatory Roles of the PDZ Domain in C-Terminal Processing
Proteases. mBio 10.

63. Ruiz N, Falcone B, Kahne D, Silhavy TJ. 2005. Chemical conditionality: a genetic
strategy to probe organelle assembly. Cell 121:307–17.

64. Wu T, Malinverni J, Ruiz N, Kim S, Silhavy TJ, Kahne D. 2005. Identification of a
multicomponent complex required for outer membrane biogenesis in *Escherichia*
coli. Cell 121:235–45.

65. Bishop RE, Gibbons HS, Guina T, Trent MS, Miller SI, Raetz CR. 2000. Transfer
of palmitate from phospholipids to lipid A in outer membranes of gram-negative
bacteria. Embo j 19:5071–80.

66. Jia W, Zoeiby AE, Petruzziello TN, Jayabalasingham B, Seyedirashti S, Bishop
RE. 2004. Lipid trafficking controls endotoxin acylation in outer membranes of
Escherichia coli. Journal of Biological Chemistry 279:44966–44975.

67. Niek D. 2000. Outer-membrane phospholipase A: known structure, unknown
biological function. Molecular Microbiology 35:711–717.

68. Malinverni JC, Silhavy TJ. 2009. An ABC transport system that maintains lipid
asymmetry in the Gram-negative outer membrane. Proceedings of the National
Academy of Sciences 106:8009–8014.

69. Paradis-Bleau C, Kritikos G, Orlova K, Typas A, Bernhardt TG. 2014. A genome-
wide screen for bacterial envelope biogenesis mutants identifies a novel factor
involved in cell wall precursor metabolism. PLoS Genet 10:e1004056.

70. Rai AK, Carr JF, Bautista DE, Wang W, Mitchell AM. 2021. ElyC and cyclic
enterobacterial common antigen regulate synthesis of phosphoglyceride-linked
enterobacterial common antigen. mBio 12:e02846–21.

71. Marvin HJ, ter Beest MB, Witholt B. 1989. Release of outer membrane fragments
from wild-type *Escherichia coli* and from several *E. coli* lipopolysaccharide
mutants by EDTA and heat shock treatments. Journal of bacteriology 171:5262–
5267.

72. Jorgenson MA, Kannan S, Laubacher ME, Young KD. 2016. Dead-end
intermediates in the enterobacterial common antigen pathway induce

morphological defects in *Escherichia coli* by competing for undecaprenyl

phosphate. Mol Microbiol 100:1–14.

73.   Konovalova A, Mitchell AM, Silhavy TJ. 2016. A lipoprotein/beta-barrel complex

monitors lipopolysaccharide integrity transducing information across the outer

membrane. Elife 5:e15276.

74.   Wall E, Majdalani N, Gottesman S. 2018. The Complex Rcs Regulatory Cascade.

Annual Review of Microbiology 72:111–139.

75.   Tata M, Kumar S, Lach SR, Saha S, Hart EM, Konovalova A. 2021. High-

throughput suppressor screen demonstrates that RcsF monitors outer membrane

integrity and not Bam complex function. Proceedings of the National Academy of

Sciences 118:e2100369118.

76.   Macritchie DM, Ward JD, Nevesinjac AZ, Raivio TL. 2008. Activation of the Cpx

envelope stress response down-regulates expression of several locus of

enterocyte effacement-encoded genes in enteropathogenic Escherichia coli.

Infect Immun 76:1465–75.

77.   Wong JL, Vogt SL, Raivio TL. 2013. Using reporter genes and the *Escherichia*

coli ASKA overexpression library in screens for regulators of the Gram negative

envelope stress response. Methods Mol Biol 966:337–57.

78.   Santos JM, Freire P, Vicente M, Arraiano CM. 1999. The stationary-phase

morphogene bolA from *Escherichia coli* is induced by stress during early stages

of growth. Molecular Microbiology 32:789–798.

79.   Santos JM, Lobo M, Matos APA, De Pedro MA, Arraiano CM. 2002. The gene

bolA regulates dacA (PBP5), dacC (PBP6) and ampC (AmpC), promoting normal

morphology in *Escherichia coli*. Molecular Microbiology 45:1729–1740.

80.   Attaibi M, den Blaauwen T. 2022. An Updated Model of the Divisome: Regulation

of the Septal Peptidoglycan Synthesis Machinery by the Divisome. Int J Mol Sci

23.

81.   Gundavarapu B, Nallamotu KC, Murapaka VV, Venkataraman B, Saisree L,

Reddy M. 2025. Identification of SanA as a novel regulator of peptidoglycan

biogenesis in *Escherichia coli*. PLoS Genet 21:e1011712.

82. Abramson J, Adler J, Dunger J, Evans R, Green T, Pritzel A, Ronneberger O,
Willmore L, Ballard AJ, Bambrick J, Bodenstein SW, Evans DA, Hung C-C,
O'Neill M, Reiman D, Tunyasuvunakool K, Wu Z, Žemgulytė A, Arvaniti E, Beattie
C, Bertolli O, Bridgland A, Cherepanov A, Congreve M, Cowen-Rivers AI, Cowie
A, Figurnov M, Fuchs FB, Gladman H, Jain R, Khan YA, Low CMR, Perlin K,
Potapenko A, Savy P, Singh S, Stecula A, Thillaisundaram A, Tong C, Yakneen S,
Zhong ED, Zielinski M, Žídek A, Bapst V, Kohli P, Jaderberg M, Hassabis D,
Jumper JM. 2024. Accurate structure prediction of biomolecular interactions with
AlphaFold 3. Nature 630:493–500.

83. Käshammer L, van den Ent F, Jeffery M, Jean NL, Hale VL, Löwe J. 2023. Cryo-
EM structure of the bacterial divisome core complex and antibiotic target
FtsWIQBL. Nat Microbiol 8:1149–1159.

84. Li Y, Gong H, Zhan R, Ouyang S, Park K-T, Lutkenhaus J, Du S. 2021. Genetic
analysis of the septal peptidoglycan synthase FtsWI complex supports a
conserved activation mechanism for SEDS-bPBP complexes. PLOS Genetics
17:e1009366.

85. Kocaoglu O, Carlson EE. 2015. Profiling of β-lactam selectivity for penicillin-
binding proteins in *Escherichia coli* strain DC2. Antimicrob Agents Chemother
59:2785–90.

86. Begg KJ, Donachie WD. 1985. Cell shape and division in *Escherichia coli:*
experiments with shape and division mutants. J Bacteriol 163:615–22.

87. Coltharp C, Buss J, Plumer TM, Xiao J. 2016. Defining the rate-limiting
processes of bacterial cytokinesis. Proc Natl Acad Sci U S A 113:E1044–53.

88. Schoenemann KM, Krupka M, Rowlett VW, Distelhorst SL, Hu B, Margolin W.
2018. Gain-of-function variants of FtsA form diverse oligomeric structures on
lipids and enhance FtsZ protofilament bundling. Mol Microbiol 109:676–693.

89. Mueller EA, Westfall CS, Levin PA. 2020. pH-dependent activation of cytokinesis
modulates *Escherichia coli* cell size. PLOS Genetics 16:e1008685.

90. Dai K, Xu Y, Lutkenhaus J. 1993. Cloning and characterization of *ftsN*, an
essential cell division gene in *Escherichia coli* isolated as a multicopy suppressor
of *ftsA12*(Ts). J Bacteriol 175:3790–7.

91. Gerding MA, Liu B, Bendezu FO, Hale CA, Bernhardt TG, de Boer PA. 2009.
Self-enhanced accumulation of FtsN at Division Sites and Roles for Other
Proteins with a SPOR domain (DamX, DedD, and RlpA) in *Escherichia coli* cell
constriction. J Bacteriol 191:7383–401.

92. Geissler B, Elraheb D, Margolin W. 2003. A gain-of-function mutation in *ftsA*
bypasses the requirement for the essential cell division gene *zipA* in *Escherichia*
coli. Proceedings of the National Academy of Sciences 100:4197–4202.

93. Hill NS, Kadoya R, Chattoraj DK, Levin PA. 2012. Cell Size and the Initiation of
DNA Replication in Bacteria. PLOS Genetics 8:e1002549.

94. Buss J, Coltharp C, Shtengel G, Yang X, Hess H, Xiao J. 2015. A multi-layered
protein network stabilizes the *Escherichia coli* FtsZ-ring and modulates
constriction dynamics. PLoS Genet 11:e1005128.

95. Hirota Y, Ryter A, Jacob F. 1968. Thermosensitive mutants of E. coli affected in
the processes of DNA synthesis and cellular division. Cold Spring Harb Symp
Quant Biol 33:677–93.

96. Chen JC, Weiss DS, Ghigo JM, Beckwith J. 1999. Septal localization of FtsQ, an
essential cell division protein in Escherichia coli. J Bacteriol 181:521–30.

97. Wijsman HJ, Koopman CR. 1976. The relation of the genes envA and ftsA in
Escherichia coli. Mol Gen Genet 147:99–102.

98. Herricks JR, Nguyen D, Margolin W. 2014. A thermosensitive defect in the ATP
binding pocket of FtsA can be suppressed by allosteric changes in the dimer
interface. Mol Microbiol 94:713–27.

99. Ishino F, Jung HK, Ikeda M, Doi M, Wachi M, Matsuhashi M. 1989. New
mutations fts-36, lts-33, and ftsW clustered in the mra region of the Escherichia
coli chromosome induce thermosensitive cell growth and division. J Bacteriol
171:5523–30.

100. Tsang MJ, Bernhardt TG. 2015. A role for the FtsQLB complex in cytokinetic ring
activation revealed by an ftsL allele that accelerates division. Mol Microbiol
95:925–44.

101.   Begg KJ, Hatfull GF, Donachie WD. 1980. Identification of new genes in a cell
961   envelope-cell division gene cluster of Escherichia coli: cell division gene ftsQ. J
962   Bacteriol 144:435–7.

963   102.   Sham L-T, Butler EK, Lebar MD, Kahne D, Bernhardt TG, Ruiz N. 2014. MurJ is
964   the flippase of lipid-linked precursors for peptidoglycan biogenesis. Science
965   345:220–222.

966   103.   Ruiz N. 2008. Bioinformatics identification of MurJ (MviN) as the peptidoglycan
967   lipid II flippase in *Escherichia coli*. Proceedings of the National Academy of
968   Sciences 105:15553–15557.

969   104.   Marquardt JL, Siegele DA, Kolter R, Walsh CT. 1992. Cloning and sequencing of
970   Escherichia coli murZ and purification of its product, a UDP-N-acetylglucosamine
971   enolpyruvyl transferase. J Bacteriol 174:5748–52.

972   105.   Yang J, Zhang Y. 2015. I-TASSER server: new development for protein structure
973   and function predictions. Nucleic acids research 43:W174–W181.

974   106.   Varadi M, Anyango S, Deshpande M, Nair S, Natassia C, Yordanova G, Yuan D,
975   Stroe O, Wood G, Laydon A, Žídek A, Green T, Tunyasuvunakool K, Petersen S,
976   Jumper J, Clancy E, Green R, Vora A, Lutfi M, Figurnov M, Cowie A, Hobbs N,
977   Kohli P, Kleywegt G, Birney E, Hassabis D, Velankar S. 2021. AlphaFold Protein
978   Structure Database: massively expanding the structural coverage of protein-
979   sequence space with high-accuracy models. Nucleic Acids Research 50:D439–
980   D444.

981   107.   Li GW, Burkhardt D, Gross C, Weissman JS. 2014. Quantifying absolute protein
982   synthesis rates reveals principles underlying allocation of cellular resources. Cell
983   157:624–35.

984   108.   Heijenoort Yv, Gómez M, Derrien M, Ayala J, Heijenoort Jv. 1992. Membrane
985   intermediates in the peptidoglycan metabolism of *Escherichia coli:* possible roles
986   of PBP 1b and PBP 3. Journal of Bacteriology 174:3549–3557.

987   109.   Marmont LS, Orta AK, Baileeves BWA, Sychantha D, Fernández-Galliano A, Li
988   YE, Greene NG, Corey RA, Stansfeld PJ, Clemons WM, Jr., Bernhardt TG. 2024.
989   Synthesis of lipid-linked precursors of the bacterial cell wall is governed by a

feedback control mechanism in *Pseudomonas aeruginosa*. Nat Microbiol 9:763–
775.

110.   Yamaguchi H, Ago R, Tahara YO, Inoue M, Niki H, Miyata M, Shiomi D. 2026.
SanA Plays a Role in Peptidoglycan Integrity in Escherichia coli. Molecular
Microbiology doi:https://doi.org/10.1111/mmi.70058.

111.   Wissel MC, Weiss DS. 2004. Genetic analysis of the cell division protein FtsI
(PBP3): amino acid substitutions that impair septal localization of FtsI and
recruitment of FtsN. J Bacteriol 186:490–502.

112.   Yang X, McQuillen R, Lyu Z, Phillips-Mason P, De La Cruz A, McCausland JW,
Liang H, DeMeester KE, Santiago CC, Grimes CL, de Boer P, Xiao J. 2021. A
two-track model for the spatiotemporal coordination of bacterial septal cell wall
synthesis revealed by single-molecule imaging of FtsW. Nature Microbiology
6:584–593.

113.   Marrec-Fairley M, Piette A, Gallet X, Brasseur R, Hara H, Fraipont C, Ghuysen
JM, Nguyen-Distèche M. 2000. Differential functionalities of amphiphilic peptide
segments of the cell-septation penicillin-binding protein 3 of *Escherichia coli*.
Molecular Microbiology 37:1019–1031.

114.   Silhavy TJ, Berman ML, Enquist LW. 1984. Experiments with gene fusions. Cold
Spring Harbor Laboratory.

115.   Datsenko KA, Wanner BL. 2000. One-step inactivation of chromosomal genes in
Escherichia coli K-12 using PCR products. Proceedings of the National Academy
of Sciences 97:6640–6645.

116.   Lutz R, Bujard H. 1997. Independent and tight regulation of transcriptional units
in *Escherichia coli* via the LacR/O, the TetR/O and AraC/I1-I2 regulatory
elements. Nucleic Acids Res 25:1203–10.

117.   Hankins JV, Madsen JA, Needham BD, Brodbelt JS, Trent MS. 2013. The outer
membrane of Gram-negative bacteria: lipid A isolation and characterization.
Methods Mol Biol 966:239–258.

118.   Deatherage DE, Barrick JE. 2014. Identification of mutations in laboratory-
evolved microbes from next-generation sequencing data using breseq. Methods
Mol Biol 1151:165–88.

119. Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for
Illumina sequence data. Bioinformatics 30:2114–20.

120. Langmead B, Salzberg SL. 2012. Fast gapped-read alignment with Bowtie 2. Nat
Methods 9:357–9.

121. Singer M, Baker TA, Schnitzler G, Deischel SM, Goel M, Dove W, Jaacks KJ,
Grossman AD, Erickson JW, Gross CA. 1989. A collection of strains containing
genetically linked alternating antibiotic resistance elements for genetic mapping
of Escherichia coli. Microbiol Rev 53:1–24.

122. Fu G, Bandaria JN, Le Gall AV, Fan X, Yildiz A, Mignot T, Zusman DR, Nan B.
2018. MotAB-like machinery drives the movement of MreB filaments during
bacterial gliding motility. Proceedings of the National Academy of Sciences
115:2484–2489.

123. Zhang H, Venkatesan S, Ng E, Nan B. 2023. Coordinated peptidoglycan
synthases and hydrolases stabilize the bacterial cell wall. Nature
Communications 14:5357.

124. Allen M, Poggiali D, Whitaker K, Marshall T, van Langen J, Kievit R. 2021.
Raincloud plots: a multi-platform tool for robust data visualization [version 2; peer
review: 2 approved]. Wellcome Open Research 4.

125. Ducret A, Quardokus EM, Brun YV. 2016. MicrobeJ, a tool for high throughput
bacterial cell detection and quantitative analysis. Nat Microbiol 1:16077.

126. Morris KN, Mitchell AM. 2023. Phosphatidylglycerol Is the Lipid Donor for
Synthesis of Phospholipid-Linked Enterobacterial Common Antigen. Journal of
Bacteriology 205:e00403–22.

127. Carr JF, Rudolf JS, Warzecha DJ, Rezenom YH, Mitchell A. 2025. Chain-length
regulation by WzzE is necessary for, but genetically separable from, cyclic
enterobacterial common antigen synthesis. bioRxiv
doi:10.1101/2025.06.25.661564:2025.06.25.661564.

1048

1049

 FIGURES AND LEGENDS

1052

Figure 1: Loss of SanA causes increased envelope permeability and low levels of
temperature sensitivity. (A) The indicated strains were grown to carbon-limited stationary
phase and then treated with SDS-EDTA or a vehicle control and viability was assayed by CFU.
The Δ*sanA* strain is significantly more sensitive to SDS than the wild-type strain from early time
points post treatment. **(B)** The sensitivity of the indicated strains to vancomycin when grown at
43 °C was assayed by EOP. Plates were supplemented with MgSO$_4$ to support growth. The
Δ*sanA* strain was more sensitive to vancomycin than the wild-type strain and this sensitivity
could be complemented with plasmid-based expression of *sanA*. **(C)** To confirm a true change in
vancomycin resistance, the MIC for vancomycin was assayed for the wild-type and Δ*sanA*
strain. The Δ*sanA* strain was significantly more sensitive to vancomycin than the wild-type
strain. **(D)** Modifications to LPS were assayed by thin layer chromatography (TLC) for strains
grown in the indicated conditions. The bands for hexa- and hepta-acylated lipid A are shown.
The Δ*sanA* strain shows an increase in hepta-acylated lipid A which can be mitigated by
complementation of *sanA* or overexpression of *pldA*. The *mlaC* strain is known to have an
increase in phospholipids in the outer leaflet of the OM (68) leading to increased helpta-acylated

lipid A and serves as a control. **(E)** Cells were grown to carbon-limited stationary phase at 37 °C

and then their ability to survive at the indicated temperatures was assayed. The Δ*sanA* strain

shows a significant decrease in survival at 57 °C. Quantitative data are the average of three

biological replicates ± the SEM. For MICs, data are shown as the geometric average with

individual data points. Images are representative of three independent experiments. * $p < 0.05$ by

the Mann Whitney Test.

1074

Figure 2: SanA shows genetic interactions with ECA synthesis mutations differentially affecting PG stress. (A-B) The indicated strains were assayed for SDS-EDTA resistance by EOP. Images are representative of three independent experiments. (A) A double mutant of

Δ*sanA* Δ*wecA* showed synthetic SDS-EDTA sensitivity, while Δ*sanA* suppressed the SDS-EDTA

sensitivity of the Δ*wecE* strain. (B) Double mutants of each ECA transglycosylase showed less

SDS-EDTA sensitivity than the previous strain. **(C)** Activation of the Rcs stress response in the

indicated strains was assayed using a *luxCDABE* reporter. Data are shown as luminescence

relative to OD_{600}. The growth of the cultures (OD_{600}) over the course of the experiment is also

shown. Data are the average of three biological replicates ± the SEM. **(D)** Cell shape of the

indicated strains was assayed, and the length and aspect ratio (length/width) were calculated.

Raincloud plots are shown with individual data points, the population distribution, and a box plot.

**** $p < 0.00005$ vs. wild type MG1655 by Wilcoxon Test; ‡ $p < 0.05$ vs. the Δ*sanA* strain by

Wilcoxon Test. **(E)** Representative images demonstrating the cell shape of the wild type Δ*sanA*,

Δ*wecA*, and Δ*sanA* Δ*wecA* strains. Insets display the indicated regions.

1090

1091

Figure 3: The SDS-EDTA of a Δ*sanA* Δ*wecA* strain can be suppressed through mutations

to *ftsI* and *prc*. (A-B) The SDS-EDTA sensitivity of the indicated strains was assayed by EOP.

Images are representative of three independent experiments. (A) Suppressor mutations show

increased SDS-EDTA resistance compared to the parent Δ*sanA* Δ*wecA* strain. (B) A linked Tn*10*

was used to transfer the *ftsI*$_{V98E}$ mutation the indicated strains. The *ftsI*$_{V98E}$ mutation was

sufficient for suppression. (C) The *ftsI*$_{S165P}$ mutation was removed from the indicated suppressor

and the loss of this mutation prevented suppression demonstrating that it is necessary for

suppression. **(D)** Reporter assays for Rcs activation were performed as in Figure 2C. The P$_{rprA}$

activity (luminescence/OD$_{600}$) as well as the growth of the strains over time are shown. The

ftsI$_{V98E}$ mutation suppressed the Rcs activation phenotype of the Δ*sanA* Δ*wecA* strain but the

ftsI$_{S165P}$ mutation did not. The Δ*sanA* Δ*wecA* strain with the *ftsI$_{S165P}$* mutation also had a

noticeable growth defect. Data are the average of three biological replicates ± the SEM. **(E)** A

cartoon of the FtsQLBIW complex modeled with AlphaFold 3 is shown (82). V98 and S165, as

well as the localization of other selected variants, are shown in stick format. The locations of the

FtsI periplasmic domains, the pedestal domain and the transpeptidase (TPase) domain are

indicated.

1108

Figure 4: Suppressors of Δ*sanA* Δ*wecA* SDS-EDTA sensitivity cause partial loss of FtsI

function. (A) Cell shape of the indicated strains was assayed, and the aspect ratio

(length/width) and cell length are shown. Raincloud plots are shown with individual data points,

the population distribution, and a box plot. * $p<0.05$, **$p<0.005$, ***$p<0.0005$, **** $p<0.00005$ vs.

wild type MG1655 by Wilcoxon Test; ‡ $p<0.05$, ‡‡ $p<0.005$, ‡‡‡ $p<0.0005$, ‡‡‡‡ $p<0.00005$ vs.

the Δ*sanA* Δ*wecA* strain by Wilcoxon Test. Mutations in *ftsI* caused the Δ*sanA* Δ*wecA* cells to

become significantly longer, while *prc* mutations had less of an effect on length. **(B)**

Representative images demonstrating cell shape are shown for the indicated strains. Insets

display the indicated regions. **(C)** The sensitivity of the indicated strains to aztreonam, a β-

lactam targeting FtsI, was assayed by MIC. Data are shown as the geometric average of six

biological replicates ± the SEM with individual data points. * $p<0.05$, ** $p<0.005$ by Mann-

Whitney Test. *ftsI23* and overexpression of *ftsN* serve as controls for decreased and increased

activity of FtsI, respectively. The *ftsI*$_{V98E}$ mutation caused a significant increase in aztreonam

sensitivity at 37 °C, indicating decreased FtsI transpeptidase activity. **(D)** The SDS-EDTA

sensitivity of the indicated strains was assayed by EOP at both 30 and 37 °C. Of the cell division

mutants, only *ftsI23* was able to suppress the SDS-EDTA sensitivity of the Δ*sanA* Δ*wecA* strain.

Images are representative of three independent experiments.

1127

1128

Figure 5: Suppression of Δ*sanA* Δ*wecA* phenotypes is linked to septal PG synthesis and not divisome assembly. EOPs were performed at 30 °C to assay the SDS-EDTA sensitivity of the indicated strains. The effect of cell division mutations in a wild-type background (A), a Δ*sanA* Δ*wecA* background (B), and a Δ*sanA* background (C) are shown. Images are representative of three independent experiments. **(A)** Only the *ftsQ1* mutation causes SDS-EDTA sensitivity in a wild-type background. **(B)** Of the assayed mutations, only *ftsQ1* suppressed the SDS-EDTA sensitivity of the Δ*sanA* Δ*wecA* strain, while earlier loss-of-function mutations in divisome assembly or gain-of-function mutations in septal PG synthesis did not. **(C)** The *ftsZ84, ftsA12,* and *ftsL** mutations make the Δ*sanA* strain more sensitive to SDS-EDTA.

1139

Figure 6: FtsI suppressor restores balance between septal and elongation PG synthesis

in a Δ*sanA* strain. (A) To compare septal PG and elongation PG synthesis, newly synthesized

PG was labeled with fluorescent D-ala (BADA) and cells were imaged. The ratio of new PG

incorporation at the center 10% of the cell versus the total cell was calculated. In both dividing

and non-dividing cells, this ratio was lower in Δ*sanA* than wild-type cells. However, the *ftsI*$_{V98E}$

suppressor in the Δ*sanA* background restored the ratio to wild type levels only in dividing cells.

(B) Representative DIC and fluorescence images showing PG incorporation are shown for the

indicated strains. Expanded regions of fluorescence images are taken from the indicated

regions. Scale bars are 3 µm.

1149

1150

Figure 7: SanA membrane permeability phenotypes are tied to lipid II availability and
FtsW activity. EOPs were performed to assay SDS-EDTA resistance in the following strains in
the indicated conditions. Images are representative of three independent experiments. **(A)** FtsW
depletion strains were constructed by deleting *ftsW* in strains carrying a plasmid-based copy of
ftsW under an arabinose-inducible promoter. Deletion of *araH*, an inducible arabinose
permease, ensures linear induction. Wild-type cells carrying the *ftsI$_{V98E}$* mutation could grow at a
lower arabinose concentration, indicating increased activity of FtsW. Interestingly, this increased
activity is not observed with the *ftsI$_{V98E}$* mutation in the Δ*sanA* or Δ*sanA* Δ*wecA* strains. **(B)** MurJ
depletion constructs were made by transduction of an arabinose inducible promoter to replace
the native *murJ* promoter. Depletion of MurJ did not prevent suppression of Δ*sanA* Δ*wecA* SDS-
EDTA sensitivity by the *ftsI$_{V98E}$* mutation. **(C)** An IPTG inducible *murA* overexpression plasmid or
a vector control were introduced into the indicated strains. Overexpression of *murA* in a Δ*sanA*
strain phenocopies the SDS-EDTA sensitivity of the Δ*sanA* Δ*wecA* strain and this sensitivity can
be suppressed by the *ftsI$_{V98E}$* mutation.

1165

1166

Figure 8: Predicted SanA pocket is important for SanA function. (A) EOPs were performed

to assay SDS-EDTA resistance of the indicated strains. The effect of mutations of *sanA* charged

residues predicted by AlphaFold 3 (82) to interact is shown. The mutations decrease SanA

function and this function is not restored by double mutants expected to restore interaction. **(B)**

Cartoon and surface predicted structures of SanA are shown with mutated residues labeled.

Charge pair residues are shown in green, hydrophilic pocket residues in magenta, hydrophobic

pocket residues in blue, and surface residues in orange. N: Cytoplasmic N-terminus; TMH:

Transmembrane helix; C: C-terminus. **(C)** The stability of proteins produced by the C-terminally

FLAG-tagged SanA mutants was assayed by immunoblot. Constructs with reduced protein

accumulation are marked in orange and with undetectable protein accumulation in red. Images

are representative of three independent experiments. GroEL serves as a loading control. **(D)**

Mutations were made to conserved polar and charged residues in the predicted pocket of SanA

and SanA function was assayed by EOP. These mutations impede SanA function. **(E)** Mutations

were made to conserved non-polar residues in the predicted pocket of SanA and SanA function

was assayed by EOP. Mutations to the residues on the surface of the pocket affected SanA

function, while mutations to residues buried inside SanA did not. **(F)** Mutations were made to

residues predicted to be on or near the surface of SanA and SanA function was assayed by

EOP. Several mutations exhibited partial loss of function. R69E serves as a control as a pocket

mutation with apparent full loss of function. Images are representative of at least three

independent experiments.

1187

1188

Figure 9: Hypothesized model for the role of SanA in modulating lipid II availability for septal PG synthesis. (A) The synthesis of lipid II begins with MurA which catalyzes the first committed step of PG synthesis. The isoprenoid carrier (undecaprenyl-phosphate) is used for the synthesis of extracytoplasmic glycans including ECA and PG. This carrier is allocated between these synthesis pathways to maintain balanced synthesis. Once lipid II is synthesized, it is flipped across the IM by MurJ. Then, the available lipid II is used by the Rod complex for elongating the cell wall and by the septal PG complex for building the septum during cell division. We hypothesize that SanA controls the availability of lipid II to FtsW and FtsI in the septal PG complex. **(B)** Removing WecA prevents use of the isoprenoid carrier for ECA synthesis increasing its availability for the synthesis of other glycans including peptidoglycan (1). Alternatively, overexpressing *murA* increases flux through the lipid II synthesis pathway, increasing lipid II levels. We hypothesize that SanA balances the increased lipid II feeding into septal and lateral PG synthesis, maintaining balanced synthesis of the envelope components. **(C)** When SanA is absent (3) and lipid II levels are increased, the increased lipid II levels cause increased lateral PG synthesis compared to septal PG synthesis, and this causes increased envelope permeability.