Systematic characterization of the ovarian landscape across mouse menopause models

Systematic comparison of mouse menopause models

Minhoo Kim[1], Rajyk Bhala[1], Justin Wang[1,2,3], Xucheng Liang[1,2], Mingfei Chen[1], Ryan J. Lu[1], Evelyn H.

Lee[1], Julio L. Alvarenga[1], Rapheal G. Williams[1], and Bérénice A. Benayoun[1,4,5,6] *

1. Leonard Davis School of Gerontology, University of Southern California, Los Angeles, CA 90089, USA.

2. Department of Quantitative and Computational Biology, University of Southern California, Los Angeles,

CA 90089, USA.

3. Department of Chemistry, University of Southern California, Los Angeles, CA 90089, USA.

4. Cancer Biology Department, USC Keck School of Medicine, Los Angeles, CA 90033, USA.,

5. Pharmacology and Pharmaceutical Sciences Department, USC Mann School of Pharmacy, Los

Angeles, CA 90033, USA.

6. USC Norris Comprehensive Cancer Center, Epigenetics and Gene Regulation, Los Angeles, CA 90033,

USA.

*Correspondence: berenice.benayoun@usc.edu

Abstract

Menopause affects not only fertility but also systemic health. Yet mechanisms underlying this coupling are still poorly understood, partly due to the absence of robust, age-relevant preclinical models with comprehensive molecular and phenotypic characterization. To address this, we systematically compared three candidate mouse models of menopause: (1) intact aging, (2) chemical ovarian follicle depletion using 4-vinylcyclohexene diepoxide (VCD) administered at multiple ages, and (3) *Foxl2* haploinsufficiency, a genetic model based on a gene linked to human premature ovarian failure. Through histology, serum hormone profiling, single-cell transcriptomics and machine-learning approaches, we identify both shared and model-specific features of follicle loss, endocrine disruption, and transcriptional remodeling. Our comparative framework enables informed selection of context-appropriate preclinical rodent models to study menopause and the broader physiological consequences of ovarian aging.

Teaser

This systematic characterization of mouse menopause models lays the groundwork for future preclinical work on the systemic impacts of menopause.

MAIN TEXT

Introduction

Women are born with a finite oocyte reserve, and its progressive depletion with age leads to the

irreversible loss of ovarian function, known as menopause (*1-3*). Epidemiological studies link later age-at-

menopause with increased lifespan and reduced risk of age-related diseases (*e.g.* cardiovascular disease,

osteoporosis, and neurodegeneration) (*4-6*). Conversely, menopause is accompanied by a sharp rise in

age-associated pathologies, underscoring widespread consequences of ovarian aging (*7*). Despite its

public health relevance, how ovarian decline contributes to systemic aging is still poorly understood, in

part due to the dearth of well-characterized, aging-relevant preclinical models.

Menopause is characterized by endocrine reprogramming that extends beyond the reproductive

system (*8*). As follicular reserves decline, reduced steroidogenic activity disrupts feedback regulation of

the hypothalamic-pituitary-gonadal (HPG) axis, leading to sustained hormonal shifts (*8*). Circulating

estrogen and anti-Müllerian hormone (AMH) fall to near-undetectable levels, while follicle-stimulating

hormone (FSH) and inhibin A (INHBA) rise (*9-11*). These changes influence multiple somatic systems,

including metabolic, skeletal, and central nervous systems (*12, 13*). However, dissecting the mechanisms

linking ovarian dysfunction to systemic physiology requires robust preclinical models recapitulating key

biological features of menopause.

The most frequently used rodent models capture menopausal physiology only imperfectly (*14*).

Although intact aging mice are widely used, they do not undergo true menopause; instead, they enter a

state of "estropause," retaining low but detectable estrogen levels and lacking the postmenopausal

endocrine milieu observed in postmenopausal women (*15, 16*). Although ovariectomy (OVX) achieves

rapid estrogen depletion, it lack a gradual endocrine transition, and it depletes residual post-reproductive

ovarian tissue, which is known to exert androgenic and paracrine effects (*17*). Thus, while OVX models

acute hormone loss, it incompletely models the progressive ovarian decline characteristic of menopause.

Repeated injections with 4-vinylcyclohexene diepoxide (VCD) leads to dramatic follicle depletion through

selective targeting of small, pre-antral follicles, inducing progressive ovarian failure while preserving a

post-reproductive ovarian tissue (*15*). This approach produces estrous acyclicity, estrogen deficiency, and

compensatory FSH elevation, resembling several endocrine features of human menopause (*18, 19*).

However, prior studies have primarily applied VCD to young adult mice (~2-3 months), limiting insight into

how the timing of ovarian failure intersects with organismal aging (*20, 21*). Because aging alters

metabolic, immune, and endocrine states in profound ways (*22-25*), understanding menopausal biology

requires evaluating ovarian decline in the context of middle-aged biology.

Genetic models informed by human physiology can also provide useful tools to understand

ovarian decline and its systemic impacts. Haploinsufficiency of *Foxl2*, a gene encoding a transcription

factor essential for granulosa cell identity and ovarian maintenance (*26-29*), is causally linked to

premature ovarian insufficiency in humans (*30-32*). In adult mice, *Foxl2* expression is primarily restricted

to ovarian granulosa cells and pituitary gonadotroph-lineage cells (*33*). While complete *Foxl2* knockout

leads to severe developmental defects and perinatal lethality, heterozygous carriers have been

incidentally reported to be subfertile, suggestive of milder ovarian dysfunction that may better recapitulate

menopausal transition (*28, 29*). Unlike surgical or chemical models, *Foxl2* haploinsufficiency represents a

physiologically relevant genetic perturbation that enables investigation of early and potentially subclinical

features of ovarian decline without exogenous insult. Rather than modeling overt menopausal hormone

depletion alone, this model may provide insight into early trajectories of ovarian dysfunction that precede

systemic endocrine collapse.

Collectively, these models could help recapitulate distinct mechanisms of ovarian decline in a

preclinical setting. However, differential impact of VCD with age-at-exposure and *Foxl2* haploinsufficiency

have not been systematically characterized and benchmarked within a unified framework. Such

comparative characterization is essential for interpreting prior literature and for selecting appropriate

models tailored to understanding systemic impacts of menopause-like ovarian failure.

Here, we present a comprehensive resource evaluating three complementary mouse models of

ovarian aging: (1) intact aging, (2) VCD-induced follicle depletion initiated at multiple adult ages, and (3)

Foxl2 haploinsufficiency. Our objective is to systematically benchmark shared and divergent features

across multiple biological scales. We integrate histological profiling, endocrine measurements, and single-

cell transcriptomics to characterize ovarian structural, functional, and molecular trajectories. Additionally,

we develop hormone-based and transcriptome-based ovarian aging clocks to provide quantitative

measures of reproductive aging across models. This comparative framework establishes a foundational

resource to guide selection and interpretation of menopause models and to support future studies

investigating how ovarian dysfunction influences systemic aging trajectories.

Results

A systematic framework for modeling menopause in mice

To systematically assess and compare physiologically relevant menopause models, we

characterized three rodent models that represent distinct mechanisms of ovarian dysfunction: (1) the

intact aging model ("Aging model"), (2) a chemically-induced model using repeated intraperitoneal

injections of 4-vinylcyclohexene diepoxide (VCD, "VCD model") at various age-of-exposure (*15, 34*), and

(3) a genetic model of *Foxl2* haploinsufficiency ("*Foxl2* +/- model") (*26*). Our aim was to establish a

comparative framework to evaluate these models in terms of histological, hormonal, and integrated

ovarian health outcomes to evaluate their suitability for modeling key aspects of ovarian aging (Fig. 1A).

The Aging model involved comparison of young adult (4 months) and old (20 months) female

C57BL/6JNia mice. We selected these timepoints to capture the transition from reproductive to firmly

post-estropausal states, as C57BL/6 females typically enter estropause ~12 months of age and show

pronounced ovarian aging phenotypes by 20 months (*35, 36*). The VCD model included C57BL/6J female

mice injected with vehicle control (safflower oil, hereafter referred to as "CTL") or VCD starting at 3, 6, 8,

or 10 months of age. This scheme allowed us to examine how the timing of ovarian insult (potentially

modeling varying "age-at-menopause") could impact ovarian physiology, particularly when VCD exposure

occurs closer to natural estropause (*35*). Importantly, incorporating multiple age-at-injection paradigms

should help disentangle ovarian-intrinsic failure mechanisms and how it interacts with the aging milieu.

The *Foxl2* haploinsufficiency model compared *Foxl2*$^{+/+}$ and *Foxl2*$^{+/-}$ female littermate mice from our

transgenic colony at young (3-4 months), early middle-age (8-10 months) and later middle-age (12-13

months) to test whether partial loss of *Foxl2* leads to anticipated occurrence of age-associated ovarian

phenotypes compared to wild-type control mice. We selected the heterozygous model because complete

Foxl2 loss is associated with severe developmental defects and high perinatal lethality *(28)*, and

heterozygous mutations of FOXL2 are sufficient to drive both syndromic and isolate premature ovarian

failure in humans patients.

To evaluate the impact of *Foxl2* haploinsufficiency on ovarian aging phenotypes, we generated a

knock-out *Foxl2* allele using mice carrying a regenerated floxed *Foxl2* locus (*26*), which allowed us to

generate mice carrying a constitutive deletion of *Foxl2*'s single exon (fig. S1A; see Materials and

Methods). Importantly, RT-qPCR analysis confirmed that *Foxl2* expression was significantly reduced in

Foxl2$^{+/-}$ ovaries compared to *Foxl2*$^{+/+}$ controls (genotype effect, p-value ~0.0206; fig. S1B). Additionally,

fertility outcomes for *Foxl2*$^{+/+}$ and *Foxl2*$^{+/-}$ females from our breeding colony confirmed previous reports of

decreased fertility of *Foxl2* haploinsufficient mice (figs. S1C,D and table S1; see Materials and Methods).

Specifically, adult *Foxl2*$^{+/-}$ females showed significantly smaller first litters (p-value ~0.0161; fig. S1C), and

increased latency to first time to delivery after male pairing compared to matched *Foxl2*$^{+/+}$ female controls

(p-value ~0.0260; fig. S1D), suggesting impaired reproductive potential consistent with ovarian

dysfunction or premature ovarian aging.

Histological and tissue-level remodeling across mouse menopause models

A decline in ovarian follicle numbers is a well-established hallmark of ovarian aging and

menopause (*37*). Thus, we performed histological analysis of ovarian follicle populations using

hematoxylin and eosin (H&E) staining (Figs. 1B-G and fig. S2A). As expected, the Aging model showed a

significant reduction in follicles at all developmental stages in old females, including primordial, primary,

secondary, and antral follicles, as well as corpora lutea (Figs. 1B,E and fig. S2B), consistent with the

established progressive follicular depletion that accompanies natural ovarian aging (*38, 39*). At 5 months

post-injections, VCD-treated animals also showed significant depletion of follicle numbers, regardless of

age-at-injection (Figs. 1C,F and fig. S2C). To note, due to the terminal nature of histological analysis,

ovaries were collected for H&E staining at a single 5 months post-injection time point. This endpoint was

selected to align with the conclusion of longitudinal serum hormone profiling (see below). In contrast, we

did not observe any significant consistent differences in combined follicle counts between *Foxl2*$^{+/+}$ and

Foxl2$^{+/-}$ ovaries (Figs. 1D,G and fig. S2D).

Next, we leveraged Picrosirius Red and Sudan Black B staining of ovarian tissue to quantify

fibrosis and lipofuscin accumulation, respectively (Fig. 2A). Previous studies have demonstrated

increased fibrosis and lipofuscin deposition in aged ovaries, reflecting progressive remodeling with

ovarian aging (*40-42*). Consistently, ovaries from the Aging model exhibited significant increases in both

fibrosis and lipofuscin accumulation in old females compared to young controls (p ~0.0190 and p

~0.00433, respectively; Figs. 2B,E,H,K). VCD-treated animals similarly showed significant increases in

both fibrosis and lipofuscin staining relative to vehicle-treated controls, indicating that chemical follicle

depletion is accompanied by stromal remodeling and accumulation in a way consistent with accelerated

ovarian aging (treatment effect, p ~0.0242 and p ~0.00776, respectively; Figs. 2C,F,I,L). In the *Foxl2*

haploinsufficiency model, despite the absence of overt follicle depletion, genotype was significantly

associated with increased fibrosis and lipofuscin accumulation (genotype effect, p ~0.00279 and p

~0.0213, respectively; Figs. 2D,G,J,M), consistent with premature aging of the ovarian stroma. These

findings indicate that structural remodeling features can emerge independently of substantial reductions

in follicle number, highlighting both shared and model-specific aspects of ovarian aging phenotypes.

Comparative endocrine alterations across mouse menopause models

To evaluate how each model perturbs endocrine regulation, we measured serum levels of AMH,

FSH and INHBA, key markers of ovarian reserve and HPG axis regulation (*43, 44*) (Fig. 3A and table S2).

Estradiol was not quantified due to known technical limitations affecting assay reliability in rodent samples

(see Discussion). To note, due to a change of FSH quantitation assays offered at the UVA ligand core

over the course of our study, FSH measurements were harmonized across methods using a polynomial

correction model (fig. S3A; see Materials and Methods).

In the Aging model, old females showed expected decreases in circulating AMH and INHBA and

increased circulating FSH compared to young females, consistent with diminished ovarian reserve (*44, 45*)

(Figs. 3B,E,H). A similar pattern was observed in the VCD model, whereby AMH and INHBA were

decreased and FSH was increased compared to the vehicle control groups at 5 months post-injection

timepoint, regardless of age-at-injection (Figs. 3C,F,I). To enable a comprehensive analysis of endocrine

responses across reproductive life stages, we also performed longitudinal hormone profiling over a five-

month period following VCD exposure at various ages (3, 6, 8 and 10 months; figs. S3B-D). Interestingly,

VCD injections at older ages resulted in attenuated hormonal shifts compared to younger counterparts,

suggesting that ovarian or systemic factors at midlife modulate sensitivity to VCD-induced insult to

ovarian reserve. In the *Foxl2*$^{+/-}$ mice, we observed conflicting hormonal trends: AMH levels were

increased (consistent with younger follicle pools), whereas FSH levels were elevated and INHBA levels

were reduced (consistent with older follicle pools) compared to wild-type controls (Figs. 3D,G,J and fig.

S3A). To note, *FOXL2* has been shown to interact with *AMH* to modulate follicle recruitment in humans

(*46*). Thus, the elevated AMH levels in the *Foxl2*$^{+/-}$ animals likely reflect altered gene regulatory networks

driven by *Foxl2* deficiency, rather than preservation or rejuvenation of ovarian reserve (see Discussion).

Next, we applied our previously described composite ovarian health index, which integrates

information from follicle counts and serum hormone levels to yield a holistic measure of ovarian health (*36*)

(Figs. 3K-M). As expected, the index was significantly lower in old *vs.* young animals and VCD-treated

animals vs. CTL (regardless of age-at-injection), reflecting impaired ovarian health (Figs. 3K,L). In

contrast, *Foxl2*$^{+/-}$ mice did not differ significantly from wild-type animals (Fig. 3M), likely due to elevated

AMH levels.

Together, these results provide a comprehensive characterization of histological and endocrine

features of three mechanistically distinct mouse menopause models. Both the Aging and VCD models

exhibit robust phenotypic hallmarks of ovarian decline, including depletion of follicular reserves, tissue

remodeling, and disrupted endocrine profiles. In contrast, the *Foxl2* haploinsufficiency model displays a

distinct phenotype characterized by preserved follicle counts and elevated AMH levels, but with

anticipated aged-like stromal remodeling and reproductive impairment. These findings indicate that the

three models capture overlapping yet non-identical dimensions of ovarian aging biology, underscoring the

importance of selecting models based on specific structural, endocrine, or regulatory processes under

investigation.

A hormone-based clock, "OvAge", for non-terminal prediction of ovarian aging

Although the ovarian health index is a useful metric of ovarian health, its partial reliance on post-

mortem histological measurements limits its applicability in longitudinal studies. Thus, we aimed at

developing a complementary, non-terminal metric of ovarian aging, using a hormone-based predictive

machine-learning model, which we termed "OvAge" (Fig. 4A). By integrating information on AMH, FSH,

and INHBA levels, we sought to develop a model that provides a more comprehensive assessment of

endocrine function rather than evaluating each hormone individually. To increase the generalizability of

the clock, we trained a random forest regression model using serum hormone data (AMH, FSH, and

INHBA) from animals with intact ovarian function and no experimental perturbation, including animals

from the Aging model, vehicle control (CTL) groups from the VCD model, $Foxl2^{+/+}$ animals from the $Foxl2$

haploinsufficiency model. In addition, to further increase statistical power, we also took advantage of

similar data from $Fshr^{+/+}$ animals derived from our previously published study of $Fshr$ haploinsufficient

mice (*47*). This hormone level data was then randomly partitioned into distinct training (n = 212) and

testing (n = 68) sets to enable robust evaluation of model performance.

Model performance evaluation showed strong agreement between predicted OvAge and

chronological age (Spearman's Rho ~0.795; p-value ~5.5×10^{-16}), confirming successful model

construction (Fig. 4B). Additionally, feature importance analysis revealed that AMH contributed most

strongly to age prediction, followed by INHBA and FSH (Fig. 4C). We then calculated OvAge using

hormonal measurements from VCD-injected and $Foxl2^{+/-}$ animals (Figs. 4D,E). In the VCD model,

predicted ovarian age was consistently higher than chronological age across all age-at-injection groups

(Fig. 4D), suggestive of accelerated endocrine aging. In contrast, predicted ages of $Foxl2^{+/-}$ animals did

not substantially differ from their wild-type counterparts (Fig. 4E), likely due to the importance of AMH

levels in OvAge calculation.

To quantify the degree of divergence between predicted and chronological age, we computed

hormonal age acceleration, defined as the difference between OvAge-predicted and chronological age

(Figs. 4F-I). For the VCD model, we analyzed animals at 30-, 90-, and 150-days post-injection. In the

VCD model, predicted ovarian age exceeded chronological age following VCD exposure (Figs. 4F-H).

Analysis revealed significant treatment effects on age acceleration at 30, 90, and 150 days post-injection.

Post hoc analyses confirmed significant increases in age acceleration at most timepoints; however,

animals injected at 10 months of age did not exhibit a significant difference at 150 days post-injection

(adjusted p-value ~0.142; table S3). This pattern may reflect convergence of ovarian aging trajectories

between control and VCD-treated animals as they approach estropausal onset (~12-15 months in

C57BL/6 mice (*35*)). In contrast, $Foxl2$ haploinsufficient animals did not show a consistent increase in

OvAge or age acceleration (genotype effect, p-value ~0.401; Fig. 4I). This lack of hormonal age

acceleration, despite evidence of stromal remodeling and reproductive impairment, suggests that *Foxl2*

haploinsufficiency does not recapitulate the endocrine aging trajectory captured by OvAge, but instead

may reflect a distinct regulatory perturbation of ovarian physiology.

Together, our OvAge analysis captures divergent ovarian aging trajectories in response to distinct

biological perturbations. The clock recapitulates expected aging patterns in naturally aged animals,

detects accelerated ovarian aging following VCD exposure, and reveals a distinct endocrine trajectory in

the *Foxl2* haploinsufficiency model. These findings support the utility of OvAge as a non-terminal

framework for assessing ovarian aging and highlight model-specific differences that may inform the study

of reproductive senescence. To facilitate broader use of this framework, we developed an interactive

online R Shiny application that enables researchers to estimate ovarian age using serum hormone

measurements (https://minhooki.shinyapps.io/OvAge_Predictor/).

Importantly, although the *Foxl2* haploinsufficiency model did not exhibit overt ovarian dysfunction

based on ovarian health index or OvAge (Figs. 3M, 4I), the absence of endocrine age acceleration does

not exclude underlying molecular perturbations. Indeed, *Foxl2*$^{+/-}$ animals displayed significant

accumulation of ovarian fibrosis and lipofuscin (Figs. 2D,G,J,M), histological hallmarks associated with

tissue aging and cell stress. Together with the established role of *FOXL2* in granulosa cell identity and

ovarian maintenance, its association with premature menopause in humans (*30, 31, 48*), and the

reproductive impairment phenotypes observed in our cohort (Fig. S1C, D), our findings suggest that *Foxl2*

haploinsufficiency likely disrupts ovarian tissue homeostasis prior to overt depletion of follicular pools,

deserving further investigation of the accompanying transcriptional landscape across ovarian cell types.

Single-cell transcriptomic profiling of ovarian cell types across menopause models

To characterize the ovarian transcriptional landscapes across candidate mouse menopause

models, we performed single-cell RNA-sequencing (scRNA-seq) on ovaries collected from the Aging,

VCD, and *Foxl2* haploinsufficiency models (Fig. 5A). For the Aging model, samples were collected from

young (4 months) and old, estropausal (20 months) animals to capture transcriptional changes associated

with intact aging (Fig. 5B). For the VCD model, we focused on two age-at-injection groups, 3 and 10

months, to capture the extremes of VCD responsiveness observed in prior analyses, as well as two post-

injection time points, 30 and 90 days, based on differential effects on age acceleration at these stages

(Fig. 5B). For the *Foxl2* haploinsufficiency model, ovaries were collected from young adult (4 months),

middle-age (9 months) and old (17 months) *Foxl2$^{+/+}$* and *Foxl2$^{+/-}$* animals, enabling the assessment of

gene expression changes driven by partial loss of *Foxl2* function across adulthood (Fig. 5B). Indeed, we

reasoned that inclusion of a later time point for the *Foxl2* haploinsufficiency model should help amplify

regulatory impact and thus help reveal clearer trends in transcriptional remodeling. The resulting datasets

yielded 15,800, 47,386, and 39,110 high-quality single cells from the Aging, VCD, and *Foxl2*

haploinsufficiency models, respectively (table S4). Ovarian cell populations were defined through

dimensionality reduction, unsupervised clustering, and annotation using both established marker genes

and automated reference mapping to publicly available single-cell datasets (Figs. 5C-H and table S5; see

Materials and Methods).

Importantly, we identified all major ovarian cell types in our datasets across all models (Figs. 5C-

H). Among the abundant non-immune populations, granulosa cells (*Amh, Fshr, Cyp19a1*), theca cells

(*Cyp17a1, Cyp11a1*), stromal cells (*Pdgfra, Col1a1*), smooth muscle cells (SMCs; *Acta2*), pericytes

(*Notch3*, *Rgs5*), blood endothelial cells (BECs; *Flt1*), lymphatic endothelial cells (LECs; *Prox1*), and

epithelial cells (*Upk1b*) were identified. The immune compartment included neutrophils (*S100a8*),

monocytes (*Adgre1*), macrophages (*C1qa*), dendritic cells (*Flt3*), innate lymphoid cells (ILCs; *Il2ra*),

natural killer (NK) cells (*Klrb1c, Gzma, Nkg7*), NKT cells (*Cd3e, Klrb1c, Gzma, Nkg7*), CD8^{+} NKT cells

(*Cd3e, Klrb1c, Gzma, Nkg7, Cd8b1*), CD8^{+} T cells (*Cd3e , Cd8b1*), CD4^{+} T cells (*Cd3e, Cd4*), double-

negative T (DNT) cells (*Cd3e*), double-positive T (DPT) cells (*Cd3e, Cd8b1, Cd4*), and B cells (*Cd19*)

(Figs. 5F-H).

As an important control, given known ovotoxic effects of VCD and the absence of follicles 150

days post treatment (Figs. 1C,F and fig. S2C), we performed additional histological analyses on

condition-matched ovaries (matched for age-at-injection and time post-injection) to confirm the presence

of follicular structures in animals processed for scRNA-seq (fig. S4). H&E staining of condition-matched

samples revealed that, despite extensive follicle depletion, residual follicles were still present in most

VCD-treated animals at these time points, consistent with the presence of follicle-associated cell types in

our scRNA-seq data (Figs. 5D,G). We also performed Picrosirius Red and Sudan Black B staining to

quantify fibrosis and lipofuscin accumulation, respectively (fig. S5). Broadly, VCD-treated ovaries

exhibited increased fibrosis and lipofuscin deposition relative to matched control samples at the 30- and

90-day time points (fig. S5), consistent with our previous observations at 150 days (Figs. 2C,F,I,L).

RNAscope validation of ovarian cell type identities

To validate cell type assignments, we performed RNAscope, a high-sensitivity *in situ* hybridization

assay for both well-characterized and less frequently profiled ovarian cell populations (Fig. 5I and figs.

S6A-C). Granulosa cells (*Fshr*), theca cells (*Cyp11a1*), stromal cells (*Pdgfra*) and SMCs (*Acta2*) were

validated using established marker probes, confirming their presence in the dataset (fig. S6A). Epithelial

cells (*Upk1b*), BECs (*Flt1*) and LECs (*Prox1*) were similarly detected, supporting the robustness of

annotations (fig. S6B). Importantly, we validated the presence of various immune cell subsets, including

NK (*Klrb1c*), NKT (*Klrb1c*, *Cd3e*), CD8$^+$ T (*Cd3e*, *Cd8b1*), CD4$^+$ T cells (*Cd3e, Cd4*), DNT (*Cd3e,* with no

Cd8b1/Cd4 signal) and B (*Cd19*) cells (fig. S6C). We also confirmed presence of less commonly reported

ovarian cell types, including CD8+ NKT (*Klrb1c, Cd3e, Cd8b1*) and DPT (*Cd3e, Cd8b1, Cd4*) cells (Fig.

5I). These results further support the accuracy of the single-cell transcriptomic annotations and highlight

the diverse somatic and immune microenvironments present across the aging models.

Together, a total of 19 cell types, 8 non-immune and 11 immune types, were shared across all

three models (fig. S6D). Some cell types were not shared across models, potentially due to technical

factors (e.g., capture efficiency and sequencing depth), or underlying biological variation inherent to each

mouse model. To assist researchers in navigating the data efficiently, we developed an interactive R shiny

app that makes the annotated datasets directly accessible and explorable through a graphical interface:

https://minhooki.shinyapps.io/shinyappmulti/.

scRNA-seq analysis reveals shifts in ovarian cell composition in menopause models

Next, we examined changes in cell type proportions across our datasets (Fig. 6A and table S6).

We first assessed whether the proportion of immune cells shifted with ovarian aging by analyzing the

expression of *Ptprc*, which encodes Cd45, a pan-immune marker (Figs. 6B-D). In the Aging model, we

observed a significant increase in *Ptprc*$^+$ cells, indicating an expansion of the immune cell population with

age (Fig. 6B). This is consistent with previously published scRNA-seq studies in intactly aged mouse

ovaries, which reported an increase in immune cell abundance at even younger ages (9-15.5 months) (*41,*

49).

VCD-treated mice exhibited a drastic reduction in *Ptprc*$^+$ cells across all conditions, indicative of a

depletion of immune cells in ovarian tissue (Fig. 6C). This reduction may result from the loss of follicular

structures, alterations in immune recruitment, or uncharacterized systemic effects of VCD on immune

homeostasis. Additionally, broader tissue remodeling, including fibrosis, may contribute to the altered

immune cell landscape observed in VCD-treated ovaries. Thus, the VCD model does not recapitulate

immune features of intact aging ovaries in mice, regardless of age-at-injection. In contrast, we detected a

significant increase in *Ptprc*$^+$ cells in *Foxl2*$^{+/-}$ animals, regardless of age (Fig. 6D). These findings were

further validated by flow cytometric quantification of CD45$^+$ cells (Figs. 6E-G and fig. S7A). Intriguingly,

this suggests that the *Foxl2* haploinsufficiency model better recapitulates age-related shifts in ovarian

immune cells compared to the VCD model.

Then, we conducted a comprehensive analysis of all ovarian cell populations detected in the

datasets to identify shifts in cellular composition in response to intact aging, VCD exposure, and *Foxl2*

haploinsufficiency (Figs. 6H-J and figs. S7B-D). In the Aging model, nearly all non-immune cell

populations showed a decline in proportion with age, except for stromal cells and SMCs, which remained

stable (Fig. 6H). In contrast, several immune cell populations increased with age, including neutrophils,

ILCs, CD8$^+$ NKT, CD4$^+$ T, DNT, and DPT cells (fig. S7B). Notably, DNT cells have been reported to

increase in aging mouse ovaries, and a potential role for these cells in regulating ovarian aging has been

proposed (*41, 50*). Conversely, monocytes, NK, NKT, CD8$^+$ T, and B cells exhibited a significant decrease

with age (fig. S7B).

In the VCD model, cell proportion shifts were largely consistent within the treatment group,

regardless of age-at-injection or time post-injection (Fig. 6I and fig. S7C). The VCD-treated group showed

an increasing trend in stromal cells, BECs, LECs, epithelial cells, and CD4$^+$ T cells. In contrast, granulosa

cells, theca cells, SMC, and pericytes were significantly reduced in proportion (Fig. 6I). Among immune

cells, neutrophils, monocytes, macrophages, DCs, NKT, DPT, and B cells were all decreased following

VCD treatment (fig. S7C).

In the *Foxl2* haploinsufficiency model, granulosa cell proportions were consistently reduced in

Foxl2[+/-] animals across all ages (Fig. 6J). In the *Foxl2* haploinsufficiency model, granulosa cell proportions

were consistently reduced in *Foxl2*[+/-] animals across all ages (Fig. 6J). Notably, this reduction in

granulosa cell representation was not accompanied by a significant decrease in total follicle number at

the histological level (Fig. 1G and fig. S2D), suggesting that *Foxl2* haploinsufficiency may alter follicular

composition (*e.g.* promoting more immature states, that hold lower granulosa cell number) or granulosa

cell state (*e.g.* altered differentiation status, impaired maturation, or partial loss of canonical granulosa cell

identity) rather than simply causing overt follicle loss. Given the established role of *FOXL2* in granulosa

cell differentiation, maintenance (*51*), and AMH-associated regulation of follicle recruitment (*46*), one

possible explanation is that partial *Foxl2* loss shifts follicles toward altered maturation states or changes

granulosa cell transcriptional identity, thereby reducing their representation in scRNA-seq without

necessarily reducing total follicle counts. Theca and SMCs also showed an overall decrease, whereas

other non-immune populations showed relatively mild shifts (Fig. 6J). In contrast, immune cell populations

generally increased in *Foxl2*[+/-] ovaries, although macrophages, dendritic cells, and DNT cells showed less

consistent trends across ages (fig. S7D). Together, these results indicate reduced representation of key

somatic ovarian cell types alongside a relative enrichment of immune populations.

To validate the cell abundance patterns identified by scRNA-seq, we performed RNAscope *in situ*

hybridization for representative non-immune and immune populations (Figs. 6K-P and figs. S8, S9). We

examined granulosa cells, theca cells, stromal cells, SMCs, BECs, LECs, and epithelial cells, along with

NK, NKT, CD8[+] NKT, CD8[+] T, CD4[+] T, DNT, DPT, and B cells. In the Aging model, RNAscope confirmed

decreases in granulosa and theca cells with age (Figs. 6K,N). In old animals, residual granulosa signals

were largely non-specific and did not colocalize with DAPI (Fig. 6K, right panel). BECs also showed a

significant decline (fig. S8A). In contrast, stromal cells, SMCs, LECs, and epithelial cells did not show

significant changes in abundance (fig. S8A). Thus, RNAscope validated several of the major somatic

shifts detected by scRNA-seq while showing more limited changes for some populations. For immune

populations, RNAscope largely recapitulated the scRNA-seq trends, showing decreases in NK, NKT, and

B cells and increases in CD8[+] NKT, CD4[+] T, DNT, and DPT cells (figs. S7B, S9A). CD8[+] T cells did not

show a significant change by RNAscope despite a decrease with age in scRNA-seq, potentially reflecting

differences in detection sensitivity or spatial distribution.

In the VCD model, we focused RNAscope analysis on the 3-month age-at-injection and 30-day

post-injection condition, given consistent trends across groups (Figs. 6I,L,O and figs. S7C, S8B, S9B).

RNAscope confirmed a reduction in granulosa cells and an increase in stromal cells following VCD

exposure. Theca cells showed a decreasing trend but did not reach significance. Other non-immune

populations did not show significant differences, in contrast to scRNA-seq findings, likely reflecting probe

sensitivity or marker expression limitations. Among immune populations, RNAscope detected decreases

in NK, NKT, CD8$^+$ NKT, CD8$^+$ T, and DNT cells, broadly consistent with the scRNA-seq results (fig. S9B).

CD4$^+$ T, DPT and B cells did not reach statistical significance.

For the *Foxl2* haploinsufficiency model, we focused RNAscope analysis on young and middle-age

groups (Fig. 6J,M,P). Our analysis confirmed reduced granulosa cell proportions in *Foxl2*$^{+/-}$ animals

across both age groups (Fig. 6J,M,P). Other somatic populations showed modest or variable changes.

Immune populations showed patterns broadly consistent with the scRNA-seq analysis, including

increased NK, NKT cells and CD8$^+$ NKT cells (fig. S9C). Although some populations did not reach

statistical significance, likely due to lower detection sensitivity, the overall trends supported immune

expansion in *Foxl2*$^{+/-}$ ovaries. Together, these results validate the major cell composition shifts detected

by scRNA-seq and support model-specific remodeling of ovarian somatic and immune populations.

Global transcriptional perturbations of ovarian cell types across menopause models

To systematically evaluate how different menopause models influence global transcriptional

responses in the ovary, we applied "Augur" (*52*), a computational framework that quantifies cell type-

specific separability across experimental conditions based on single-cell level gene expression profiles

(fig. S10A). Augur scores (derived from underlying machine-learning algorithm performance, reported as

Area Under the Curve [AUC] values) represent the ability within each cell type to distinguish between

biological groups within a given model based on overall transcriptional landscapes (*52*). In the Aging

model, granulosa cells exhibited the highest AUC score (AUC ~0.721) among non-immune cells,

reinforcing their role as key transcriptional responders to chronological aging (figs. S10B,C). In the

immune compartment, NKT and ILCs ranked highest (fig. S10C). Interestingly, DNT cells, despite their

expansion in the aging ovary, showed the lowest AUC score (AUC ~0.519), indicating limited

transcriptional remodeling with age (fig. S10C).

In the VCD model, we observed marked transcriptional perturbations in theca, stromal, BEC, and

LEC cells, especially at 90 days post-injection, with these effects largely consistent across both 3-month

and 10-month age-at-injection groups (figs. S10D-G). However, epithelial and granulosa cells displayed

more pronounced age-at-injection-dependent differences: epithelial cells were more responsive in

younger animals, while granulosa cells were notably absent in the 10-month cohort, likely reflecting more

severe follicular depletion (fig. S10G, left panel). Among immune cell types, NK, DNT, and CD8$^+$ NKT

cells showed higher transcriptional sensitivity in younger animals (fig. S10G, right panel). At 30 days post-

injection, transcriptional divergence was generally more divergent in the VCD model compared to its 90-

day counterpart (figs. S10D,E). For example, LECs in the 10-month group displayed strong divergence

(AUC ~0.81), suggesting age-related vascular remodeling may emerge early after VCD exposure.

In the *Foxl2* haploinsufficiency model, transcriptional divergence was more pronounced in old

animals, suggesting that the effects of *Foxl2* loss may not fully manifest in the younger ovaries (fig. S11).

Among non-immune populations, granulosa, theca and epithelial cells showed the highest AUC scores in

older age groups (figs. S11C,E, left panels). In immune populations ILC, NK and monocytes exhibited the

strongest divergence, with all showing elevated AUC scores in the older age groups, indicating potential

immune activation or heightened sensitivity of specific immune compartments to *Foxl2* insufficiency (figs.

S11C,E, right panels).

Together, these results reveal both common and model-specific overall transcriptional trajectories

across ovarian aging paradigms. Granulosa cells consistently emerge as sensitive indicators of ovarian

dysfunction across all three models. Immune cells show particularly pronounced divergence in *Foxl2*$^{+/-}$

animals, suggesting that immune alterations may precede, accompany, or even drive ovarian changes in

this model. Notably, because *Foxl2* expression in adults is largely restricted to granulosa cells in the ovary

and gonadotroph-lineage cells in the pituitary and it is not expressed at meaningful levels in ovarian

immune cells (*33*), the immune-related phenotypes observed here are unlikely to be a cell autonomous

results of decreased *Foxl2* expression within immune cell types. These comparisons underscore the

importance of contextualizing cell type-specific transcriptional changes within each model's mechanistic

framework and highlight how distinct perturbations, such as chronological aging, chemical ablation, or

genetic loss, can produce divergent molecular responses across ovarian cell types.

Transcriptional and pathway-level signatures across menopause models

To investigate transcriptional remodeling across menopause models, we performed differential

expression and pathway enrichment analyses across cell types and models (Fig. 7A and table S7).

Importantly, we used a robust pseudobulking approach per annotated cell type in each individual scRNA-

seq library, via "muscat" (*53*), so as to limit false positive rate. Six cell types, granulosa, theca, stromal,

BEC, epithelial, and DNT cells, were consistently detected across all three models and thus included in

downstream analyses (fig. S12A). "DESeq2" (*54*) was used to identify differentially expressed genes

(DEGs) for each model: young vs. old in the Aging model; CTL vs. VCD in the VCD model (using age-at-

injection and time post-injection as modeling covariates); and *Foxl2*$^{+/+}$ vs. *Foxl2*$^{+/-}$ in the *Foxl2*

haploinsufficiency model (using age as a modeling covariate). As expected, we confirmed reduced *Foxl2*

expression in *Foxl2*$^{+/-}$ animals in our scRNAseq dataset (fig. S12B).

Using an FDR cutoff of 0.05, we observed substantial transcriptional remodeling across models

and cell types (Figs. 7B-D). Importantly, we asked such transcriptional changes may be unduly influenced

by underlying differences in estrous cyclicity of mice. For this purpose, we took advantage of a publicly

available scRNA-seq dataset profiling ovaries from young adult mice across the four stages of the estrous

cycle (fig. S13A) (*55*). After processing the data using our standardized pseudobulk workflow, we focused

on granulosa cells, which were among the most strongly perturbed populations across estrous cycle

stages (*55*). We first compared multidimensional scaling (MDS) structures across the estrous-cycle

dataset and our Aging, VCD, and *Foxl2* haploinsufficiency datasets (fig. S13B). Whereas our datasets

showed sample separation according to the primary biological variables of interest (*i.e.* age, treatment,

genotype), samples from the estrous-cycle dataset revealed substantial overlap in MDS space,

suggesting smaller or less consistent transcriptional remodeling. Consistently, differential expression

analysis identified only modest transcriptional differences between estrous stages, even in the proestrus-

versus-estrus comparison, which has previously been reported to show the strongest stage-dependent

effects (fig. S13C) (*55*). These findings indicate that the transcriptional changes observed across our

menopause models are substantially larger than those attributable to estrous cycle-effects and are

therefore unlikely to be driven primarily by stage differences.

Next, we performed gene set enrichment analysis (GSEA) to identify gene ontology (GO)

biological process terms recurrently enriched across models (Fig. 7E and table S8). Mitochondrial-related

gene sets were consistently downregulated across models and cell types, including "Respiratory electron

transport chain," "Oxidative phosphorylation" and "ATP synthesis coupled electron transport" (FDR < 0.1;

Fig. 7F). Immune-related processes, such as "Antigen processing and presentation of endogenous

peptide antigen," were also frequently identified as downregulated. Using independent gene sets from

Reactome (*56*) revealed consistent trends of mitochondria-related gene sets being downregulated (FDR <

0.1; Fig. 7F), suggesting convergent dysregulation of metabolic function during ovarian aging and in

menopause models. Notably, immune-related gene sets were more prominently altered in the Aging and

Foxl2 haploinsufficiency models compared to the VCD model, consistent with the immune cell shifts

observed in these datasets and suggesting that the *Foxl2* haploinsufficiency model could be particularly

informative for investigating immune dysregulation associated with ovarian aging.

Convergent and divergent aging-associated features are observed across menopause models

To identify transcriptional features shared or divergent across menopause models, we derived

age-associated gene signatures from the Aging model and used them as reference cell type-specific

"aging" gene sets (fig. S14A). The VCD model showed broad concordance, with consistent upregulation

of aging-associated genes across cell types. In contrast, the *Foxl2* haploinsufficiency model displayed

partial divergence. Granulosa and stromal cells showed reduced expression of genes typically

upregulated with age, while BECs exhibited inverse trends for both age-associated up- and

downregulated gene sets (fig. S14A). These cell type-specific deviations suggest that partial *Foxl2* loss

may engage distinct molecular programs relative to physiological aging. We also evaluated enrichment of

the published, validated SenMayo senescence-associated gene set (*57*) as a proxy for senescence

burden (fig. S14B). As expected, most cell types in the Aging model showed increased SenMayo

expression. The VCD model partially recapitulated this pattern, whereas the *Foxl2* haploinsufficiency

model showed mixed, cell type-dependent effects, with several non-immune cell populations exhibiting

inverse trends. These differences indicate that VCD and *Foxl2* haploinsufficiency models capture distinct

cell-specific aspects of accelerated ovarian aging.

Increased transcriptional variability is a characteristic of aging and has been observed across

multiple tissues, including the ovary (*49*) and hypothalamus (*58*) in mice, and is thought to reflect a loss of

transcriptional regulation/precision with aging. Thus, we next evaluated gene expression variability in

each cell type and condition using the coefficient of variation (CV) (fig. S14C). As expected, transcriptional

variability broadly increased with aging across cell types (fig. S14C). The VCD model showed elevated

variability primarily in theca cells, while other populations exhibited heterogeneous responses. In the

Foxl2 haploinsufficiency model, variability increased predominantly at middle-age, although inconsistently

across cell types. These mixed patterns suggest that intervention models may involve compensatory or

cell type-specific adaptations rather than uniform aging-like deregulation.

Finally, we examined transposable element (TE) transcription using "scTE" (*59*) (fig. S14D). TE

transcription is known to be derepressed with aging across tissues, driven in part by age-associated

chromatin remodeling, reduced heterochromatin integrity and impaired epigenetic repression (*60, 61*).

While aging was associated with increased TE expression in several somatic and immune populations,

intervention models showed more heterogeneous responses. VCD exposure led to increased TE

expression in stromal and vascular compartments but reductions in steroidogenic cells, whereas *Foxl2*

haploinsufficiency showed cell type-specific increases and decreases across ovarian lineages. These

findings indicate that TE regulation varies depending on the mechanism of ovarian perturbation rather

than uniformly mirroring chronological aging.

Age-associated gene expression trajectories reveal model-specific amplification patterns

To better characterize how VCD exposure and *Foxl2* haploinsufficiency modulate age-associated

transcriptional trajectories, we performed likelihood ratio test (LRT) analysis for the VCD and *Foxl2*

haploinsufficiency models (Fig. 8A). Significant genes were subsequently grouped by unsupervised

clustering to identify coherent gene expression patterns. Due to their relevance to ovarian function as key

steroidogenic cells, we focused our analyses on granulosa and theca cells.

To interpret cluster behavior relative to physiological aging, we first examined expression patterns

in corresponding control groups (CTL for the VCD model; *Foxl2*[+/+] for the *Foxl2* haploinsufficiency model)

to establish baseline age-associated trajectories. Clusters were then classified based on how the

perturbation modified these trajectories. When perturbations exaggerated age-associated increases or

decreases, we termed these "enhanced increase" (EI) or "enhanced decrease" (ED) patterns,

respectively (Fig. 8A).

In granulosa cells, both VCD exposure and *Foxl2* haploinsufficiency modified age-associated

transcriptional trajectories in distinct, but partially overlapping ways (Figs. 8B-E). In the VCD model, EI

genes were enriched for hormone metabolism pathways, whereas ED genes were associated with

mitochondrial respiration (Figs. 8B,C). In the *Foxl2* haploinsufficiency model, EI genes were enriched for

respiration-related pathways, while ED genes were linked to housekeeping processes, including cell cycle

regulation, DNA replication, and recombination (Figs. 8D,E). Notably, mitochondrial and respiration-

related pathways came up in both models, appearing in VCD ED clusters and *Foxl2* haploinsufficiency EI

clusters, indicating that mitochondrial programs are broadly perturbed across models, albeit in opposite

directions. Although additional significant gene regulatory patterns were also observed (fig. S15), these

did not show consistent aging-associated trajectories and we did not perform further analyses on them.

In theca cells, we also observed broadly similar amplification patterns (figs. S16, S17). VCD-

associated EI genes were enriched for metabolic pathways, whereas ED genes were enriched for

processes linked to actin organization and cell signaling (figs. S16A-C). In the *Foxl2* haploinsufficiency

model, ED genes were enriched for extracellular matrix and cell signaling pathways, suggesting altered

stromal and intercellular communication (fig. S16D). Similar to the granulosa analysis, we also detected

additional significant trajectory patterns in *Foxl2* haploinsufficient theca cells, but since these did not

demonstrate clear aging-related behavior or significant functional enrichment, suggestive of model-

specific effects (fig. S17), they were not explored further.

Together, these trajectory analyses reveal that both VCD exposure and *Foxl2* haploinsufficiency

modulate age-associated transcriptional programs but do so in distinct ways. VCD primarily amplifies

metabolic and mitochondrial changes in steroidogenic cells, whereas *Foxl2* haploinsufficiency

preferentially affects pathways related to respiration, cell cycle regulation, and extracellular matrix

organization. These results highlight how different menopause models reshape age-dependent

transcriptional trajectories across ovarian cell types.

Gene co-expression network and transcription factor activity across menopause models

To identify gene regulatory programs associated with ovarian aging, we performed weighted gene

co-expression network analysis (WGCNA) (*62*) and transcription factor (TF) activity inference using

"decoupleR" (*63*) (fig. S18A). As a reference framework, WGCNA was first applied to pseudobulked gene

expression data from the Aging model, which captures physiological ovarian decline. Gene modules

significantly correlated with age (FDR < 0.1) were retained for comparative analyses (figs. S18B, S19). As

expected, these modules showed strong enrichment within the Aging dataset, providing a benchmark for

assessing transcriptional convergence and divergence across intervention models (fig. S20A).

Module behavior in the VCD model largely paralleled that of the Aging model, indicating that

VCD-induced ovarian failure recapitulates major regulatory programs associated with chronological aging

(fig. S20A). In contrast, the *Foxl2* haploinsufficiency model displayed more divergent module dynamics.

Notably, stromal cell modules frequently showed opposite expression patterns relative to the Aging model,

suggesting that partial *Foxl2* loss alters regulatory programs governing stromal remodeling and tissue

organization (fig. S20A). These comparisons highlight both shared aging-related regulatory circuits and

model-specific deviations shaped by the underlying mechanism of ovarian dysfunction.

To investigate the functional basis of these modules, we performed Gene Ontology over-

representation analysis (ORA; figs. S20B-D). In granulosa cells, modules were enriched for pathways

related to mitochondrial metabolism and ribosomal function, consistent with high energetic and

translational demands in steroidogenic cells (fig. S20B) (*64, 65*). Stromal cell modules were enriched for

mitochondrial processes as well as mitotic machinery and chromosome segregation pathways,

suggesting coordinated changes in cellular proliferation and tissue remodeling during aging (fig. S20C)

(*41, 66*). In BECs, enriched terms included immune-related pathways, chromosome segregation, and

cytoskeletal organization, pointing to coupled regulation of vascular integrity, immune interactions, and

structural dynamics (fig. S20D) (*67*).

To complement co-expression network-based analyses, we also inferred transcription factor (TF)

activity across cell types using decoupleR (fig. S21A and table S10). Across datasets, we detected

shared changed TF activity signatures in multiple cell types (fig. S21B). Across datasets, shared TF

activity changes were observed in multiple cell types (fig. S21B). In granulosa cells, we consistently

detected reduced predicted activity of cell cycle-associated regulators (E2f1, E2f2, E2f3, E2f4, and Myc;

fig. S21C) (*68, 69*). In contrast, predicted activity increased for steroidogenic regulators (Nr5a1, Nr5a2)

(*70, 71*), metabolic regulators (Srebf1) (*72*), and stress- and hormone-responsive factors (Crem, Foxo1,

Jun, and Sp1) (*73-76*), suggesting coordinated remodeling of endocrine function, metabolic homeostasis,

and stress-adaptive programs in granulosa cells (fig. S21C).

Beyond granulosa cells, TF activity changes revealed distinct cell type-specific regulatory

patterns (fig. S21C). Theca cells showed reduced activity of Ar, Foxo3, Nr3c1, and Stat3, consistent with

attenuated steroidogenic signaling and stress-response pathways (*77, 78*). Stromal cells exhibited broad

decreases in TFs associated with developmental signaling, tissue remodeling, and cellular plasticity,

alongside increased immune function-linked regulatory activity, suggesting altered stromal homeostasis

with enhanced immune involvement (*79-81*). Among vascular endothelial populations, BECs displayed

reduced cell cycle-related TF activity (*82*), whereas LECs showed increased activity of immune and

inflammatory regulators (*83, 84*), indicating divergent adaptations between blood and lymphatic

vasculature. Epithelial cells exhibited reduced Sp1 activity, suggesting dampened transcriptional

programs linked to cellular maintenance and metabolic regulation (*85*).

Collectively, these findings reveal both conserved and model-specific regulatory programs

underlying ovarian aging, with coordinated transcription factor activity changes highlighting cell type-

specific shifts in endocrine function, metabolic regulation, tissue remodeling, and immune adaptation.

Transcriptome-based aging clocks capture evidence of accelerated ovarian aging across models

Finally, to obtain an integrative view of ovarian aging, we developed transcriptome-based clocks

that predict ovarian age from gene expression profiles (Fig. 9A and fig. S22A), similar to approaches used

in other tissues (*86, 87*). Given the central role of granulosa cells in ovarian function, we first trained a

lasso regression model using granulosa cell transcriptomes (Fig. 9A). In parallel, a comparable model

was constructed using theca cell transcriptomes to assess the fidelity of our menopause models in

another key steroidogenic population (fig. S22A).

Performance evaluation on held-out data demonstrated strong predictive accuracy for the

granulosa-based model (Spearman's Rho ~0.82, p-value ~1.3 × 10^{-585}; Fig. 9B). The theca-based model

also performed well (Spearman's Rho ~0.76, p-value ~3.6 × 10^{-447}; fig. S22B), although accuracy declined

more sharply at older ages. Notably, prediction accuracy decreased in samples derived from mice aged

≥12 months, coinciding with the reported onset of estropause in C57BL/6 mice (*35*) (fig. S22B; expected

age-at-estropause highlighted in gray).

We next applied the granulosa-based clock to VCD and *Foxl2* haploinsufficiency datasets (Figs.

9C,D). VCD-treated animals exhibited consistently elevated predicted transcriptomic ages relative to

chronological age, indicating molecular acceleration of ovarian aging. *Foxl2*$^{+/-}$ animals similarly showed

increased predicted ages compared to wild-type controls. To further quantify these differences, we next

calculated transcriptomic age acceleration (predicted minus chronological age; Figs. 9E,F). In the VCD

model, transcriptional age acceleration decreased with increasing age at exposure, mirroring trends

observed in the hormone-based OvAge model, consistent with reduced responsiveness of older ovaries

to chemical insult (Figs. 4F-H, 9E). In contrast, *Foxl2*$^{+/-}$ granulosa cells showed increased transcriptional

age acceleration relative to controls, consistent with acceleration of ovarian aging and contrasting with

our previous OvAge predictions (Figs. 4I, 9F). This discrepancy suggests that transcriptome-based clocks

may capture earlier, distinct physiological alterations preceding systemic hormonal changes.

In parallel, we also applied the theca cell-based clock to the VCD and *Foxl2* haploinsufficiency

models (figs. S22C,D). Despite its nominally lower accuracy, the theca clock recapitulated key aging

patterns observed in granulosa cells. VCD-treated animals again showed elevated predicted

transcriptomic ages. In the *Foxl2* haploinsufficiency model, genotype-dependent effects varied with age:

Foxl2$^{+/-}$ animals exhibited increased predicted ages at young and middle-age but reduced age

acceleration in the older cohort relative to wild-type animals. Age acceleration patterns were broadly

consistent with results from granulosa-based clock in the VCD model, while *Foxl2* haploinsufficiency

showed age-dependent genotype effects (figs. S22E,F). Analysis confirmed a significant overall genotype

effect on age acceleration, supporting the sensitivity of the theca-based clock in detecting molecular

ovarian aging across models. Together, these results demonstrate that transcriptome-based aging clocks

capture molecular features of accelerated ovarian aging and reveal both convergent and cell type-specific

responses to chemical and genetic perturbations.

Finally, to identify transcriptional features driving age prediction, genes with non-zero β

coefficients from the trained lasso model were extracted and their expression trajectories examined

across age in the pseudobulked granulosa cells (Fig. 9G). These genes displayed heterogeneous

temporal patterns, indicating involvement in diverse age-associated regulatory programs. To further

assess biological relevance, the top 500 genes based on absolute β coefficient were subjected to ORA

(Fig. 9H). Enriched terms were predominantly related to extracellular matrix organization, cell signaling,

and immune function, including "extracellular matrix," "signaling receptor regulator activity," and

"leukocyte chemotaxis." These results suggest that early alterations in tissue remodeling, intercellular

communication, and immune regulation contribute substantially to ovarian transcriptomic aging and

underscore the sensitivity of the transcriptome-based clock in detecting early molecular changes.

A similar approach was applied to the theca cell-based model (figs. S22G,H). Genes with non-

zero β coefficients also exhibited heterogeneous age-dependent expression patterns, suggesting

involvement in diverse regulatory programs during theca cell aging (fig. S22G). To evaluate their

biological relevance, the top 500 genes based on absolute β coefficient were subjected to ORA (fig.

S22H). Enriched terms were predominantly associated with reproductive and developmental processes,

including "sex differentiation" and "gonad development." Additional enrichment was observed for

structural and translational processes such as "actin cytoskeleton reorganization" and components of the

cytosolic ribosome, including "small ribosomal subunit." These findings suggest that transcriptional aging

in theca cells is characterized by coordinated remodeling of reproductive developmental programs

alongside with shifts in cytoskeletal organization and protein synthesis capacity.

Together, these data demonstrate the utility of transcriptome-based models for detecting

molecular aging trajectories in the mouse ovary. These predictive tools offer a valuable framework for

assessing molecular age across experimental models and for uncovering early transcriptomic alterations

that precede overt phenotypic decline.

Discussion

Understanding how ovarian decline contributes to systemic aging requires robust experimental models that accurately recapitulate the physiological features of menopause. In this study, we systematically evaluated three mechanistically distinct mouse models of menopause: an intact aging model, a chemical follicle depletion model using VCD exposure, and a genetic model of *Foxl2* haploinsufficiency. By integrating histological, hormonal, and transcriptomic profiling across these models, we provide a comparative framework for understanding shared and model-specific trajectories of ovarian aging. Findings from our study provide a critical comparative resource for identifying conserved and divergent features of ovarian aging and evaluating the suitability of each model, with the goal of advancing both mechanistic insights and translational strategies in menopause research.

The intact Aging model showed expected hallmarks of physiological ovarian decline, including depletion of follicles across developmental stages, endocrine disruption, increased fibrosis and lipofuscin accumulation, and broad transcriptomic remodeling. At the cellular level, aging was associated with expansion of the immune compartment, reduction of most non-immune populations, and strong transcriptional perturbation in granulosa cells. These findings are consistent with prior scRNA-seq studies of mouse ovarian aging and reinforce the view that granulosa cells are among the most transcriptionally responsive somatic populations during ovarian decline (*41, 49*). Aging was also accompanied by increased transcriptional variability, altered TE expression, and coordinated pathway-level changes involving mitochondrial dysfunction and immune remodeling, indicating loss of tissue homeostasis across multiple biological scales.

The VCD model led to the appearance of several key features of accelerated ovarian failure. VCD-treated animals exhibited marked follicle depletion, endocrine disruption, increased fibrosis and lipofuscin accumulation, and elevated ovarian age as measured by both OvAge and transcriptome-based clocks. Importantly, by varying the age at VCD exposure, we demonstrate that ovarian failure can be induced across different stages of the reproductive lifespan, including near the estropausal transition. This establishes the VCD paradigm as a flexible system for modeling age-at-menopause, enabling the generation of ovarian failure within an age-relevant systemic context. Additionally, we observed that the magnitude of endocrine and transcriptomic acceleration was greater when VCD exposure occurred at

younger ages and attenuated in animals treated closer to the estropausal transition. This finding indicates

that the physiological consequences of follicle depletion are shaped by the baseline ovarian and systemic

state at the time of insult. At the transcriptional level, VCD broadly recapitulated many features of the

Aging model, including module behavior, age-associated gene signature enrichment, and transcriptome-

based age acceleration. However, it also diverged from intact aging in important ways. Most notably,

VCD-treated ovaries showed reduced immune cell proportions, in contrast to the immune expansion

observed in naturally aged ovaries. This suggests that chemically induced follicle depletion captures

aspects of ovarian failure and accelerated endocrine aging, but does not fully model the immune

remodeling that accompanies physiological ovarian aging. Thus, the VCD model appears particularly

useful for studying acute follicular depletion and endocrine collapse, but less suitable for interrogating

immune-related features of intact ovarian aging.

The *Foxl2* haploinsufficiency model showed a more nuanced phenotype. Unlike the Aging and

VCD models, *Foxl2*$^{+/-}$ animals did not exhibit overt reductions in combined follicle counts or consistent

endocrine age acceleration by ovarian health index or OvAge. Nonetheless, this model displayed

significant fibrosis and lipofuscin accumulation, impaired fertility, increased immune cell abundance, and

marked transcriptional perturbation across several ovarian compartments. Granulosa cell proportions

were consistently reduced, and transcriptome-based clocks detected elevated molecular age in granulosa

cells despite the absence of a clear endocrine aging phenotype. Together, these findings indicate that

Foxl2 haploinsufficiency does not simply phenocopy overt menopause, but instead captures an

alternative trajectory of ovarian dysfunction in which structural and molecular perturbations emerge before

strong endocrine collapse. *FOXL2* interacts with *AMH* to regulate follicle recruitment in humans (*46*),

raising the possibility that the observed histological, endocrine, and transcriptional features reflect

enhanced AMH-mediated preservation of ovarian reserve in *Foxl2*$^{+/-}$ animals. Despite relatively preserved

endocrine output, the heightened transcriptional sensitivity of steroidogenic and immune compartments

positions this model as a valuable system for studying early or subclinical stages of ovarian aging.

This distinction is further supported by the transcriptomic analyses. Compared to the Aging and

VCD models, *Foxl2* haploinsufficiency showed more divergent behavior in aging-associated gene

signatures, senescence-associated programs, and co-expression modules, particularly in stromal cells. At

the same time, immune-related signals were more prominent in $Foxl2^{+/-}$ ovaries, consistent with the

increase in $Ptprc^+$ cells and the stronger transcriptional responsiveness of several immune populations.

These findings suggest that partial loss of $Foxl2$ may create an immune-primed ovarian environment and

reshape transcriptional programs in ways that only partially overlap with chronological aging. Accordingly,

the $Foxl2$ haploinsufficiency model may be particularly informative for studying early or subclinical stages

of ovarian dysfunction, especially those involving altered granulosa identity, tissue remodeling, and

immune dysregulation.

710   We want to acknowledge that our $Foxl2$ haploinsufficiency model uses a constitutive null allele

rather than an ovary/granulosa-specific one (see Materials and Methods). However, this is not really a

concern in the effort to model reproductive senescence due to the known adult expression pattern of

$Foxl2$. Indeed, adult expression of $Foxl2$ is largely restricted to ovarian granulosa cells and pituitary

gonadotroph-lineage cells, with no evidence of expression in (ovarian) immune cell populations or other

tissues (*33*). Thus, the immune phenotypes observed in $Foxl2^{+/-}$ ovaries are unlikely to arise from cell-

autonomous $Foxl2$ deficiency within immune cells, and instead more likely reflect secondary

consequences of altered ovarian or neuroendocrine homeostasis. At the same time, because decreased

$Foxl2$ pituitary expression is likely to contribute to aspects of the reproductive phenotype (as they likely do

in human $FOXL2$ mutation carriers with premature ovarian failure), future studies could also evaluate how

the hypothalamus/pituitary axis is impacted in the context of reproductive aging models, with relevance to

human health. Notably, $FOXL2$ has been shown to directly regulate expression of the common

glycoprotein hormone α-subunit (Cga), a shared component of both FSH and LH, in pituitary

gonadotrophs. Given that reproductive aging and menopause are characterized by elevated circulating

FSH levels, and that this feature is recapitulated across our three models, partial $Foxl2$ loss may

contribute to dysregulation of gonadotropin production, thereby influencing ovarian function in a non-cell-

autonomous manner (*88*).

727   Across models, several shared molecular themes emerged. Mitochondrial and respiration-related

728   pathways were consistently altered at the pseudobulk level, and trajectory analyses in granulosa cells

729   further highlighted mitochondrial programs as recurrent targets of remodeling in both VCD and $Foxl2$

730   haploinsufficiency, albeit in opposite transcriptional directions. Gene regulatory network analyses also

converged on processes related to mitochondrial metabolism, ribosomal function, cell cycle control, extracellular matrix organization, and immune adaptation. Transcriptome-based aging clocks reinforced these observations by identifying predictive features linked to extracellular matrix remodeling, signaling, immune regulation, and developmental programs. Together, these analyses point to common axes of ovarian aging biology while also showing that the direction, magnitude, and cellular context of these changes differ by model.

Our study also highlights the importance of cell type-specific readouts in evaluating ovarian aging. Granulosa cells consistently emerged as sensitive indicators across models, showing strong perturbation in Augur analyses, high DEG burden, and robust performance in transcriptome-based aging clocks. Theca cells similarly supported molecular age prediction and revealed complementary trajectory patterns, although the *Foxl2* model showed age-dependent divergence between granulosa- and theca-based clocks. These findings emphasize that ovarian aging is not uniform across cell types and that distinct compartments encode partially overlapping aspects of functional decline.

We would like to discuss several potential limitations of this work. First, this study was designed as a comparative resource and not to directly resolve how ovarian aging drives systemic aging phenotypes. Second, circulating estradiol was not quantified because technical constraints limit reliable measurement in mouse blood samples, including assay cross-reactivity with dietary phytoestrogens (including equol), extensive protein binding requiring complex extraction procedures, and limited sample volumes (*14*). Third, although the VCD age-at-injection paradigm enabled examination of age-dependent responsiveness to chemical follicle depletion, it does not directly model the aged systemic milieu. Finally, although cross-species parallels support the relevance of several pathways identified here, further work is needed to determine which model most faithfully captures specific aspects of human menopause biology.

In summary, our comparative analysis demonstrates that ovarian aging comprises both shared and model-specific structural, endocrine, and molecular features. The intact Aging model captures gradual physiological decline and immune remodeling; the VCD model induces robust follicle depletion and accelerated endocrine aging; and the *Foxl2* haploinsufficiency model reveals early molecular and tissue perturbations that may precede overt functional decline. By integrating histological, hormonal, and single-cell transcriptomic data, we provide a multidimensional framework for selecting and interpreting

759   preclinical models of ovarian aging. This resource should support future studies aimed at identifying

760   mechanisms, biomarkers, and interventions relevant to reproductive senescence and menopause-

761   associated health decline.

Materials and methods

Experimental Design

This study was designed as a comparative resource to systematically evaluate mouse models of menopause and ovarian aging. Our objective was to determine how distinct mechanisms of ovarian dysfunction (physiological aging, chemically induced follicle depletion, and genetic disruption of granulosa cell identity), shape structural, endocrine, and molecular features of ovarian aging. To this end, we performed integrated histological, hormonal, and single-cell transcriptomic profiling across three models: intact aging, VCD-induced ovarian failure, and *Foxl2* haploinsufficiency.

Prespecified components included standardized histological assessment of follicle abundance and tissue remodeling, endocrine profiling of ovarian reserve markers, single-cell RNA sequencing to characterize cellular composition and transcriptional states, and integrative computational analyses to quantify molecular aging trajectories and regulatory programs. Harmonized preprocessing and analytical pipelines were applied across models to enable direct comparisons.

Mouse husbandry

All animals were treated and housed in accordance with the Guide for Care and Use of Laboratory Animals. All experimental procedures were approved by the USC's Institutional Animal Care and Use Committee (IACUC) and are in accordance with institutional and national guidelines. Samples were derived from animals on approved IACUC protocol numbers 21155 and 21454. All animals were acclimated in the specific-pathogen-free animal facility at USC for two weeks prior to any experimental procedures. Mice were provided PicoLab Rodent Diet 20 (LabDiet, 5053) *ad libitum*. The facility was maintained on a 12-hour light/dark cycle, with housing rooms set to 72°F and 30% humidity.

For the Aging model, female C57BL/6JNia mice (4- and 20-month-old) were obtained from the National Institute on Aging (NIA) colony at Charles River Laboratories.

For the VCD model, female C57BL/6J mice (2.5-, 5.5-, 7.5-, and 9.5-month-old) were purchased from Jackson Laboratory. Animals (strain # 000664) received daily intraperitoneal injections of either vehicle (safflower oil; Sigma S8281) or VCD (160 mg/kg/day; Sigma 94956) for 15 consecutive days. All

injections were administered between 8:00 and 10:00 AM to minimize variability due to circadian
influences.

Foxl2 floxed (*Foxl2^lox/lox^*) mice were generated by Cyagen Biosciences Inc. using a previously
described targeting strategy (*28*). Briefly, a targeting construct was designed to replace a 2.177 kb
fragment containing the entire *Foxl2* coding region with a 1.157 kb *Neo* gene cassette (fig. S1A). The
construct was linearized and introduced into C57BL/6NTac embryonic stem (ES) cells via electroporation.
Candidate clones were assessed via PCR and Southern blot analysis. A targeted ES cell clone was
selected and injected into C57BL/6NTac albino embryos, which were then implanted into pseudo-
pregnant CD-1 females. Founder animals were identified based on coat color, and germline transmission
was confirmed through breeding with C57BL/6J females followed by genotyping of the offspring. Mice
carrying the desired floxed *Foxl2* allele were sent to USC for experiments and downstream phenotyping.

Foxl2 haploinsufficiency mice (*Foxl2^+/-^*) were generated by crossing homozygous *Foxl2^lox/lox^*
C57BL/6NTac mice with heterozygous B6.C-Tg(CMV-cre)1Cgn/J (JAX, stock #006054) mice. Offspring
were genotyped for both the *Foxl2* alleles and the CMV-Cre transgene. To eliminate the CMV-Cre
transgene, *Foxl2^+/-^* mice then were backcrossed to wild-type C57BL/6J mice. Progeny were selected to
carry the deleted *Foxl2* allele (*Foxl2^+/-^*), but not the CMV-Cre transgene. Afterwards, *Foxl2^+/-^* mice were
maintained as a stable heterozygous knock-out colony using allele transmission through parents of either
sex with *Foxl2^+/+^* by *Foxl2^+/-^* crossing to avoid generating full knockout animals. Routine genotyping was
performed using genomic DNA extracted from tail biopsies using specific PCR primers, as listed in table
S11A.

RT-qPCR analysis of *Foxl2* expression in *Foxl2* haploinsufficiency model mouse ovaries

Flash-frozen ovaries collected from mice were used for RNA extraction, cDNA synthesis and
quantitative RT-PCR. Ovaries were resuspended in 600 µL of TRIzol reagent (Thermo-Fisher, 15596018)
and lysed using the BeadBug Homogenizer (Benchmark Scientific, D1036). Homogenization was
performed at 3,500 rpm in 30-second intervals, repeated for a total of nine cycles. Total RNA was then
purified using the Direct-Zol RNA Miniprep kit (Zymo Research, R2052), following the manufacturer's
protocol. RNA was eluted in nuclease-free water and immediately quantified using the NanoDrop

spectrophotometer (Thermo Scientific). The purity of the RNA was assessed by measuring the 260/280nm and 260/230nm absorbance ratios. Samples with ratios close to ~2.0 were considered suitable for downstream applications.

Total RNA was reverse transcribed into cDNA, using the Thermo Scientific™ Maxima H Minus First Strand cDNA Synthesis Kit (Thermo Scientific, K1682), following the manufacturer's protocol. Quantitative PCR was performed using SensiFAST SYBR no-ROX kit (Bioline, BIO-98020) and the MIC Tubes and Caps kit (Bio Molecular Systems, 71e101) on the Magnetic Induction Cycler (MIC) machine (Bio Molecular Systems, MIC-2). micPCR v2.12.6 was used to capture and quantify Ct values. Gene expression levels were calculated using the ΔΔCt method, normalized to the geometric mean of two housekeeping genes: *Ubc* and *Hprt*. Primers and sequences used for RT-qPCR reactions can be found in table S11B.

Fertility assessment of *Foxl2* haploinsufficiency model mice

Fertility data were obtained from our ongoing breeding colony, where $Foxl2^{+/+}$ and $Foxl2^{+/-}$ females were paired with $Foxl2^{+/-}$ and $Foxl2^{+/+}$ males of our colony, respectively. Importantly, data was included in the fertility assessment only if (i) $Foxl2^{+/+}$ and $Foxl2^{+/-}$ female pairings were initiated on the same day, to avoid confounds from potential housing-related or handling-related stress events, and (ii) females were aged 2.5 and 3.5 months at the time of pairing, to avoid biases related to hyperfertility of younger females and to allow sufficient time for *Foxl2* haploinsufficiency–associated phenotypes to emerge.

Fertility was assessed based on the pup count of the first litter and the latency to the first litter (*i.e*. the time between pairing and birth of the first litter). Differences in litter size between $Foxl2^{+/+}$ and $Foxl2^{+/-}$ females were assessed using the aligned rank transform ANOVA to test effects. One outlier from the $Foxl2^{+/+}$ group was removed based on a significant Grubbs' test (p-value < 0.05), using the "outliers" package (v.0.15) (*89*) in R (v. 4.1.2). Latency to the first litter was analyzed using a log-rank test to compare survival curves, using the "survival" package (v. 3.6-4) (*90*) in R. Raw fertility data can be found in table S1A.

Hematoxylin-eosin staining of mouse ovarian tissues

Ovaries were fixed in Bouin's solution (Sigma, HT10132) for 24 hours at room temperature before

being transferred to 70% ethanol for storage. Paraffin embedding, tissue sectioning, and hematoxylin and

eosin (H&E) staining were carried out by the USC Norris Comprehensive Cancer Center Translational

Pathology Core Facility and Immunohistochemistry Laboratory at USC Labs. H&E-stained slides were

imaged using the Keyence BZ-X All-in-One Fluorescence Microscope platform using 20X objective and

automated stitching using default parameters.

Ovarian follicle counts

Follicle counts, including primordial, primary, secondary, and antral follicles, and corpus luteum,

were performed on three sections per ovary by three blinded observers (fig. S2A). Median values from

the three observers were used for data analysis. Statistical significance was assessed using the Wilcoxon

test for the Aging model and aligned rank transform ANOVA to test effects for the VCD and *Foxl2*

haploinsufficiency models.

Ovarian fibrosis and lipofuscin histological staining and analysis

Ovaries were fixed in Bouin's solution (Sigma, HT10132) for 24 h at room temperature and

transferred to 70% ethanol for storage. Paraffin embedding, tissue sectioning, and hematoxylin and eosin

(H&E) staining were performed by the USC Immunohistochemistry Laboratory. Histological staining was

conducted using the following kits according to the manufacturers' instructions: Picrosirius Red Stain Kit

(Abcam, ab150681) for fibrosis detection and Sudan Black B Histochemical Stain Kit (StatLab, KTSBB)

for lipofuscin detection. Stained slides were imaged using a Keyence BZ-X All-in-One Fluorescence

Microscope with a 20x objective and automated stitching under default settings. For each ovary, three 8-

μm sections were imaged and analyzed. Mean values across sections were used for downstream

analysis.

Image quantification was performed using "ImageJ2" (v. 2.14.0/1.54f). Non-ovarian regions were

manually traced and excluded from analysis. Color deconvolution was performed using the Colour

Deconvolution 2 plugin with the FastRed FastBlue DAB vector. For Picrosirius Red images, the Colour_1

channel was used to quantify fibrotic signal and the Colour_2 channel to define total ovarian area. For

Sudan Black B images, the Colour_2 channel was used to quantify lipofuscin signal and the Colour_1

channel to define total ovarian area. Deconvoluted grayscale images were thresholded to identify positive

signal. Background thresholds (entire ovary) were set between 210–230, and positive signal thresholds

were set between 150–170 depending on staining batch. Threshold parameters were determined

separately for each batch based on positive and negative control sections. Signal intensity was calculated

as the percentage of positive area relative to total ovarian area. All measurements were performed in a

blinded manner.

Statistical significance was assessed using the non-parametric Wilcoxon rank-sum test for the

Aging model. For the VCD and *Foxl2* haploinsufficiency models, factorial effects were evaluated using

aligned rank transform ANOVA.

Quantification of serum AMH, FSH and INHBA concentrations

Blood was collected either immediately following euthanasia via cardiac puncture or from live

animals via a facial vein blood draw. Blood was allowed to clot at room temperature for one hour. Serum

was then separated using the MiniCollect® Serum Tube (Greiner, 450472) and stored at -80°C until

further analysis. The University of Virginia Center for Research in Reproduction Ligand Assay and

Analysis Core quantified serum levels of AMH (Rat and Mouse Anti-Müllerian Hormone (AMH) ELISA kit,

Ansh Labs, AL-113), FSH (Millipore Pituitary Panel Multiplex kit, RPT86K or Ultra-Sensitive Mouse & Rat

FSH, UVA Ligand Core, in-house (*91*)), and INHBA (Inhibin A ELISA kit, Ansh Labs, AL-161), providing

standardized normalized values. Statistical significance was assessed using the Wilcoxon test for the

Aging model and aligned rank transform ANOVA to test effects for the VCD and *Foxl2* haploinsufficiency

models. Raw serum hormone quantification data can be found in table S2.

Two different FSH assay kits were used due to changes in offered FSH quantitation assays at the

UVA core during the time of our study. To ensure compatibility between measurements made using the

different kits, we applied a correction procedure as previously described (*36*). Briefly, matched serum

samples were analyzed using both kits, and a polynomial regression model was trained to capture the

relationship between the two outputs. This model was then applied to adjust FSH values obtained using

the ultra-sensitive assay, enabling direct computation of ovarian health index across datasets. The scripts

used for ovarian health index calculation are available on the Benayoun lab GitHub repository:

https://github.com/BenayounLaboratory/Ovarian_Aging_Microbiome/tree/main/1_Ovarian_health_index_c

alculation.

Ovarian health index calculation

The ovarian health index was calculated as previously described (*36*). The scoring framework

and reference thresholds were derived from the original cohort reported in that study. The index

integrates two components: ovarian hormone levels (AMH, FSH, and INHBA) and follicle counts, which

included the combined counts of primordial, primary, secondary, and antral follicles, as well as corpora

lutea. Each parameter was assigned a score based on a three-tier system. Values exceeding the median

of the young female group were assigned a score of 3, values positioned between the medians of the

young and old groups were assigned a score of 2, and values below the median of the old group were

assigned a score of 1. The hormone score was calculated as the mean of the individual scores for AMH,

FSH, and INHBA. This hormone score was then combined with the follicle score in a 1:1 ratio to generate

the overall ovarian health index, which was subsequently scaled to a 0-100 range for standardization.

Statistical significance was assessed using the Wilcoxon test for the Aging model and aligned rank

transform ANOVA to test effects for the VCD and *Foxl2* haploinsufficiency models.

OvAge clock model training

The OvAge clock was trained using serum hormone quantification data from multiple model and

cohorts, including animals from aging model, vehicle control animals from VCD model, wild-type animals

from *Foxl2* haploinsufficiency model (*Foxl2*[+/+]) and wild-type animals from a previously published *Fshr*

haploinsufficiency model (*Fshr*[+/+]) (*47*). To enhance the applicability of the model across different genetic

and experimental backgrounds, animals from multiple models were included in the training dataset.

Hormone data included serum levels of AMH, FSH, and INHBA, with corresponding age at blood

collection recorded in weeks.

The dataset was partitioned into training (75%) and testing (25%) subsets using stratified

sampling based on age, using the createDataPartition function from R package "caret" (v. 6.0-91) (*92*). A

random forest (RF) regression model was trained using the training set, with out-of-bag (OOB) error

correction applied for internal validation, using caret (v. 6.0-91) (*92*) and "randomForest" packages (v.

4.7-1) (*93*). The model was optimized by tuning the mtry parameter (best-performing value = 3), yielding

an OOB root mean square error (RMSE) of 13.13 weeks and an OOB R^2 of 0.65. A total of 500 trees

were used in the final model. Model performance was evaluated using Spearman correlation between

chronological and predicted ages in the test dataset. The RF model achieved a Spearman correlation

coefficient of 0.795 with a p-value of 5.5×10^{-16}. Age acceleration was calculated as difference between

predicted and chronological age. The final OvAge clock R object containing the trained RF model is

available   on   the   Benayoun   lab   GitHub   repository:

https://github.com/BenayounLaboratory/Mouse_Menopause_Models. We have also developed an

interactive R shiny app to enable users to input hormone values and estimate predicted ovarian age using

the OvAge clock: https://minhooki.shinyapps.io/OvAge_Predictor/.

Single-cell RNAseq sample and library preparation

Ovaries from two animals (four ovaries total) were used for most single-cell RNA sequencing

(scRNA-seq) sample and library preparations to optimize cellular viability and yield. In one instance

(*Foxl2$^{+/+}$ and Foxl$^{+/-}$* old group pair), a single animal (two ovaries) was used due to sample availability.

Single-cell suspensions were generated using Miltenyi's Multi Tissue Dissociation Kit (Miltenyi

Biotec, 130-110-201) with the gentleMACS Octo Dissociator with Heaters (Miltenyi Biotec, 130-096-427),

running the 37_m_LDK1 program. To maximize the recovery of viable single cells, the program was

deliberately aborted 10 sec after start of the highspeed phase, approximately 20 sec before program

completion. After dissociation, red blood cells were removed using Red Blood Cell Lysis Buffer (Miltenyi

Biotec, 130-094-183) according to the manufacturer's instructions. Dead cells were subsequently

eliminated using the EasySep Dead Cell Removal (Annexin V) Kit (STEMCELL Technologies, 17899).

Cell count and viability were assessed via Viobility™ 405/452 Fixable Dye (Miltenyi Biotec,

J66584-AB) staining and flow cytometry using a MACSQuant Analyzer 10 (Miltenyi Biotec, 130-130-420).

Flow cytometry data were analyzed with FlowLogic v8 (Inivai Technologies). Cells were gated based on

forward scatter area (FSC-A) vs. side scatter area (SSC-A), and singlets were identified by evaluating the

linear grouping of cells in FSC-A vs. FSC-H plots (fig. S8A).

Single-cell libraries were prepared using the Chromium Next GEM Single Cell 3☐ GEM, Library &

Gel Bead Kit v3.1 (10X Genomics, PN-1000121) according to the manufacturer's instructions. Based on

flow cytometry estimates, cell suspensions were loaded to achieve a targeted recovery of 6,000 cells per

sample. Samples were processed on a Chromium Next GEM Chip G (10X Genomics, 2000177) as per

the manufacturer's protocol.

Completed single-cell libraries were assessed for quality using the 4200 TapeStation system

(Agilent Technologies, G2991A) with High Sensitivity D1000 DNA ScreenTape (Agilent Technologies,

50675584). Libraries were sequenced on an Illumina NovaSeq 6000, generating 150 bp paired-end reads

at Novogene USA. Raw FASTQ reads have been deposited in the Sequence Read Archive under

accession PRJNA863443.

Single-cell RNAseq data analysis

<u>Data processing</u>

Raw sequencing reads were processed using "Cell Ranger" (v. 7.1.0) (*94*) from 10x Genomics

(table S4). Reads were aligned to the mm10 reference genome. To remove contamination from ambient

RNA, decontamination was performed using DecontX from the "celda" (v. 1.14.2) (*95*) package. Raw and

filtered feature-barcode matrices were used to estimate ambient RNA levels, and corrected expression

matrices were generated for each sample. Doublet removal was performed using a two-step approach.

First, "DoubletFinder" (v. 2.0.3) (*96*) was applied to each sample independently, estimating doublet rates

based on the expected multiplet rates from 10x Genomics. The optimal pK parameter was determined

using a parameter sweep, and predicted doublets were labeled. In parallel, a second doublet detection

method, "scds" (v. 1.10.0) (*97*), was applied using the cxds_bcds_hybrid function. Cells identified as

doublets by either method were excluded from further analysis.

The filtered single-cell expression matrices were loaded into "Seurat" (v. 4.3.0) (*98*), and initial

quality control (QC) steps were applied. Cells with fewer than 500 detected features, a total RNA count

outside the range of 500-100,000, mitochondrial RNA content exceeding 15%, or a DecontX

contamination score above 0.25 were removed. Following QC, datasets were normalized using

SCTransform, regressing out the number of detected features, mitochondrial content, DecontX

contamination score and batch effects.

<u>Data integration</u>

To correct for batch effects, integration was performed using "Harmony" (v. 1.0) (*99*). First,

datasets were processed separately by normalizing gene expression using SCTransform, regressing out

number of detected features (nFeature_RNA), mitochondrial percentage (percent.mito), ambient RNA

contamination (decontX_contamination), and sequencing library (Library). Highly variable genes were

selected using vst (nfeatures=2000 genes). The preprocessed datasets were merged using Seurat (v.

4.3.0) (*98*), ensuring shared variable features were maintained. Principal component analysis (PCA) was

performed on the merged dataset, and batch effects were assessed by visualizing PCA embeddings

colored by batch identity. Batch correction was applied using Harmony, where batch information was

explicitly modeled as a covariate. The corrected embeddings were used for downstream analysis,

including UMAP visualization, clustering, and differential expression analysis.

1000

<u>Cell type annotation</u>

Cell type annotation was conducted using a multi-step approach that combined automated

reference-based methods with manual marker-based annotation. First, *Ptprc* $^+$ and *Ptprc* $^-$ cell populations

were identified using "scGate" (v. 1.0.1) (*100*). Cells expressing *Ptprc* (Cd45) were classified as immune

cells, while those lacking *Ptprc* expression were designated as non-immune cells. The two subsets were

then processed separately for further annotation. Each subset underwent independent cell type

annotation using three complementary computational methods: "SingleR" (v. 1.8.1) (*101*), "scSorter" (v.

0.0.2) (*102*), and "scType" (v. 1.0) (*103*). The ImmGen dataset (*104*) was used as a reference for *Ptprc* $^+$

cells, while two publicly available ovarian single-cell RNA-seq datasets (Open Science Framework ID

924fz (*55*) and NCBI GEO accession no. GSE232309 (*41*)) were used for *Ptprc* $^-$ cells. To supplement

these automated methods, manual annotation was conducted based on the expression of canonical

marker genes specific to ovarian cell types (table S5). The final annotation was determined using a

majority voting approach - cell type labels from SingleR, scSorter, scType, and manual annotation were

compared, and the most frequently assigned label was selected as the final classification. All analyses

were performed using Seurat (v. 4.3.0) (*98*) within R.

1016

<u>Cell type proportion and Augur analysis</u>

Cell type proportions were quantified using "scProportionTest" (v. 0.0.0.9000) (*105*) within R. This

method performs a permutation-based test to compare cell-type abundances between two groups,

estimating the relative change in proportions along with a confidence interval for each cell type. The

analysis was conducted at two levels: the broader classification of *Ptprc*$^+$ vs. *Ptprc*$^-$ populations and the

more detailed sub-cell type level within each category. Relative proportion of detected cell types can be

found in table S6.

We used "Augur" (v. 1.0.3) (*52*) to identify cell types that exhibited the most transcriptional

changes associated with aging. Pairwise comparisons were conducted for each experimental condition,

assessing differences between (1) young and old females, (2) vehicle control (CTL) and VCD-treated

groups, and (3) *Foxl2*$^{+/+}$ and *Foxl2*$^{+/-}$ genotypes.

1028

<u>Pseudobulk analysis for differential gene expression</u>

Pseudobulk differential gene expression analysis was conducted separately for three datasets:

the Aging, VCD, and *Foxl2* haploinsufficiency models. Single-cell transcriptomic data were aggregated at

the cell type level using "muscat" (v. 1.18.0) (*53*) to generate pseudobulk expression profiles. To ensure

robust analysis, cell types were filtered based on dataset-specific criteria. In the Aging model, cell types

with at least 25 cells per sample across all samples were retained. For the VCD model, cell types were

included if they contained at least 25 cells in at least 12 samples, representing 75% of the dataset. In the

Foxl2 haploinsufficiency model, the threshold was set at a minimum of 25 cells in at least 8 samples, or

80% of the dataset.

Differential gene expression analysis was performed using "DESeq2" (v. 1.44.0) (*54*), with

comparisons made between young and old females in the Aging model, vehicle control (CTL) and VCD-

treated samples in the VCD model, and *Foxl2*$^{+/+}$ and *Foxl2*$^{+/-}$ samples in the *Foxl2* haploinsufficiency

model. Batch effects in the VCD and *Foxl2* haploinsufficiency models were assessed using "sva" (v.

3.52.0) (*106*) to determine the number of surrogate variables (SVs), and batch correction was applied

using the removeBatchEffect() function from "limma" (v. 3.60.6) (*107*). The adjusted counts were

subsequently used for DESeq2 analysis, incorporating age-at-injection and time post-injection as

covariates for the VCD model and age as a covariate for the *Foxl2* haploinsufficiency model. Variance-

stabilized transformed counts were computed using getVarianceStabilizedData(), and differentially

expressed genes (FDR < 0.05) were visualized using strip plots. The complete pseudobulk DESeq2

output, including log fold changes, adjusted p-values, and base mean expression values for each dataset

and cell type, can be found in table S7.

1050

<u>GO analysis for differential gene expression</u>

Gene set enrichment analysis (GSEA) was performed to identify Gene Ontology Biological

Process (GO BP) and Reactome terms enriched in differentially expressed genes across cell types in the

Aging, VCD, and *Foxl2* haploinsufficiency models. The analysis was conducted using the "clusterProfiler"

package (v. 4.2.2) (*108*) and the Molecular Signatures Database (MSigDB) gene sets retrieved via

"msigdbr" (v. 7.4.1) (*109*). Differential expression results from pseudobulk DESeq2 analysis were ranked

by t-statistic and used as input for GSEA. Enrichment analysis was performed separately for each cell

type in each dataset, with an FDR cutoff of 0.1 applied to identify significant terms. To identify recurrent

biological processes across datasets, GO-BP terms enriched in at least four cell types within each

dataset were extracted. The list of significantly enriched GO-BP terms (FDR < 0.1) for all datasets can be

found in table S8.

1062

<u>Estrous cycle transcriptomic variability analysis</u>

To evaluate potential confounding effects of estrous stage, we reanalyzed a publicly available

mouse ovary scRNA-seq dataset spanning the four estrous stages (*55*). Count matrices were obtained

from Open Science Framework (accession 924fz), and stage annotations were incorporated as metadata.

Granulosa cells were aggregated into pseudobulk profiles using muscat (v. 1.18.0) (*53*) and analyzed

using DESeq2 (v. 1.44.0) (*54*) following the same pipeline described above. Variance-stabilized counts

were used for principal component analysis (PCA; prcomp) and multidimensional scaling (MDS;

Spearman distance, cmdscale). Differential expression analysis was performed in DESeq2, focusing on

the proestrus versus estrus contrast.

1072

<u>GSEA of aging-associated transcriptional signatures across models</u>

GSEA was performed using differentially expressed genes (DEGs) from the Aging model dataset

as input gene sets with the clusterProfiler (v4.12.6) (*108*) package. Upregulated and downregulated gene

sets (FDR < 0.05) were extracted from the Aging model dataset and tested for enrichment in the Aging,

VCD and *Foxl2* haploinsufficiency model datasets. Genes were ranked based on the DESeq2-derived t-

statistic for analysis. Additionally, GSEA was conducted to assess the enrichment of the SenMayo (*57*)

gene set within the dataset. Differential expression results from the Aging, VCD, and *Foxl2*

haploinsufficiency datasets were used for GSEA, with genes ranked using the DESeq2-derived t-statistic.

1081

<u>Coefficient of variation analysis</u>

Gene expression variability across cell types was assessed by calculating the coefficient of

variation (CV) for each gene in the Aging, VCD, and *Foxl2* haploinsufficiency models. CV was computed

as the ratio of the standard deviation to the mean expression level per gene within each condition (young

vs. old for Aging, CTL vs. VCD for VCD, and $Foxl2^{+/+}$ vs. $Foxl2^{+/-}$ for *Foxl2* haploinsufficiency model) using

variance-stabilized transformed counts obtained from DESeq2 (v1.44.0) (*54*) pseudobulk analysis. For

the *Foxl2* haploinsufficiency model, the old-age group was excluded from CV analysis due to insufficient

biological replicates. Statistical differences in CV between groups were evaluated using the Wilcoxon

rank-sum test.

1091

<u>Transposable element (TE) analysis</u>

Transposable element (TE) expression was quantified using "scTE" (v1.0) (*59*) with default

parameters. BAM files generated from Cell Ranger (v. 7.1.0) (*94*) outputs were processed using the

mm10 genome index provided by scTE. Features detected by scTE but absent in the single-cell gene

expression data were extracted and merged, generating a Seurat object containing both gene and TE
expression profiles.

Cells were aggregated at the cell type level using muscat (v1.18.0) (*53*), and differential
expression analysis was performed using DESeq2 (v1.44.0) (*54*), following the same approach as
described above.

To assess functional enrichment of TE families, GSEA was conducted on ranked gene lists
derived from DESeq2 results. TE annotations were obtained from the UCSC database for mm10, the
same reference used by scTE. TE subfamilies were stratified to broader categories, including LINE, SINE,
LTR, and DNA transposons. TE family annotations were obtained from the *rmsk.txt* file downloaded from
the UCSC genome annotation database for the mm10 reference genome. For each cell type, genes were
ranked by t-statistic from differential expression analysis, and enrichment of TE families was assessed
using clusterProfiler (v4.12.6) (*108*). Significant TE families (FDR < 0.1) were reported for visualization.

1108

<u>Likelihood ratio test–based trajectory analysis</u>

Pseudobulk counts of granulosa and theca cells from each dataset were analyzed using
likelihood ratio tests (LRTs) implemented in DESeq2 (v1.44.0) (*54*) to identify genes exhibiting
perturbation-dependent age-associated transcriptional trajectories in the VCD and *Foxl2*
haploinsufficiency models. Resulting p-values were adjusted using the Benjamini-Hochberg method, and
genes with FDR < $\times 10^{-5}$ were retained. Variance-stabilized expression values of significant genes were
subjected to unsupervised clustering using the degPatterns function from "DEGreport" (v 1.40.1) (*110*).
Baseline age-associated trends were defined from control groups, and clusters were classified according
to whether perturbations amplified or attenuated age-related transcriptional changes.

1118

<u>WGCNA and ORA enrichment analysis</u>

Weighted gene co-expression network analysis (WGCNA) (*62*) was performed on the Aging
model dataset to identify gene co-expression modules. Variance-stabilized transformed counts from
DESeq2 (v1.44.0) (*54*) were used as input, with analyses conducted separately for each cell type. For
each cell type, an optimal soft-thresholding power was determined using pickSoftThreshold, followed by

the construction of an adjacency matrix and topological overlap matrix (TOM). Hierarchical clustering was

performed to identify gene modules, which were assigned colors using cutreeDynamic, and eigengenes

were computed to summarize module expression patterns. Gene sets from each module were classified

into upregulated and downregulated groups based on eigengene expression differences between young

and old samples. These gene sets were then subjected to GSEA via clusterProfiler (v. 4.2.2) (*108*), using

differential expression results from the VCD and *Foxl2* haploinsufficiency model datasets to assess

whether the identified modules were significantly enriched in these datasets. Modules significantly

associated with aging (FDR < 0.1) were further analyzed through over-representation analysis using

clusterProfiler (v. 4.2.2) (*108*) and the org.Mm.eg.db annotation package.

1133

<u>Transcription factor activity inference using decoupleR</u>

Transcription factor (TF) activity inference was performed using "decoupleR" (v2.10.0) (*63*) with

the CollecTRI transcriptional regulatory network for mouse. Differential expression results from the Aging,

VCD, and *Foxl2* haploinsufficiency models were used as input. For each dataset, fgsea scoring was

applied to rank TF activity based on DESeq2-derived t-statistics, and significant TFs were identified at a

threshold of FDR < 0.05. To compare TF activity across datasets, TFs were filtered to retain those with

consistent activity direction (positive or negative) in at least two datasets.

1141

1142 <u>R shiny application generation</u>

1143 An interactive R shiny application of Aging, VCD and *Foxl2* haploinsufficiency scRNA-seq

1144 datasets was generated using "ShinyCell" (v.2.1.0) (*111*), and made available at

1145 <u>https://minhooki.shinyapps.io/shinyappmulti/</u>.

1146

RNAscope sample preparation, imaging and data analysis

In situ <u>Hybridization Protocol</u>

Frozen ovaries were mounted in Tissue-Tek O.C.T Compound (Sakura, 4583) and sliced into 8

μm slices from the same ovarian region using PTFE-coated microtome blades (Duraedge, 7223) at

−20°C on a Cryostat CM1860 (Leica, 14-0491-46884). 3 ovarian sections from the same region were

mounted in succession on VWR Premium Superfrost Plus Microscope slides (VWR, 48311-703). All

batches of sample processing were prepared using the fresh-frozen sample preparation protocol along

with negative and positive controls (Advanced Cell Diagnostics, UM 323100-USM). 12 probes were

tested in 3 sets, with Set 1 containing markers for theca (*Cyp11a1*; C1), granulosa (*Fshr*; C2), smooth

muscle (*Acta2*; C3), and stromal (*Pdgfra*; C4) cells. Set 2 had markers for epithelial cells (*Upk1b*; C1),

lymphatic endothelial cells (*Prox1*; C2), blood endothelial cells (*Flt1*; C3), and B cells (*Cd19*; C4). Set 3

contained probes for T (*Cd3e*; C1), natural killer (*Klr1b1c*; C2), CD8+ (*Cd8b1*; C3), and CD4+ (*Cd4*; C4)

cells. Per the protocol, the RNAscope™ Multiplex Fluorescent Reagent Kit v2 (Advanced Cell Diagnostics,

323100) was used. The fluorescent signals TSA Vivid Fluorophore 520 (Advanced Cell Diagnostics,

323271), TSA Vivid Fluorophore 570 (Advanced Cell Diagnostics, 323272), TSA Vivid Fluorophore 650

(Advanced Cell Diagnostics, 323273), and Opal 690 (Akoya Biosciences, FP1497001KT) were used to

stain the probes in each set. The slides were preserved in ProLong Gold Antifade Mountant (Thermo

Fisher Scientific, P36930).

1165

1166   <u>Imaging</u>

1167   Imaging was performed using a Leica Stellaris 5 confocal microscope (Leica Microsystems)

1168   equipped with a HC PL APO 20x/0.75 IMM CORR CS2 objective (Leica, 11506343) and Leica

1169   Microsystems Type F Immersion liquid oil (Leica Microsystems, 11513859). Images were acquired using

1170   Leica Application Suite X software (LAS X v4.4.0).

1171

<u>Imaging analysis</u>

To quantify probe expression, the images were analyzed using the "QuPath" program (v0.5.1).

Cell boundaries based on DAPI signal were calculated using the following settings modified from the

default parameters – Requested pixel size: 0 μm, Nucleus background radius: 10 μm, Maximum area:

400 μm^2, Threshold: 0.5, and Cell expansion: 2. Sets 1 and 2 were analyzed using subcellular detection

to determine their expression levels relative to the number of cells. All thresholding values were uniform

across the date of processing of each ovarian tissue. For Set 3, single measurement classifiers were

created for each probe to generate colocalization data. Cells below the detection threshold were not

counted for analysis.

1181

Cell-type proportion analysis using flow cytometry

Single-cell suspensions of ovarian cells were prepared following the same protocol used for

single-cell RNA sequencing sample preparation, as described above. Cells were stained for APC-CD45

antibody (Miltenyi, 130-123-784) according to the manufacturer's recommendations. Flow cytometry data

were acquired using the MACSQuant10 flow cytometer (Miltenyi Biotec, 130-096-343) and analyzed

using FlowLogic v.8 (Inivai Technologies). The raw flow cytometry dataset was deposited to Figshare

(DOI: 10.6084/m9.figshare.28733222).

1189

Granulosa and theca transcriptional clock model training and feature assessment

Granulosa and theca cells were independently subsetted from scRNA-seq datasets

encompassing multiple experimental models, including natural aging, VCD, and *Foxl2* haploinsufficiency.

Additionally, an independent middle-aged female dataset (13, 76, and 86 weeks) was included to ensure

age coverage for intermediate ages not well represented in the datasets.

To develop the cell transcriptional clocks, cells from aging model, middle-aged females, vehicle

controls from the VCD model, and *Foxl2+/+* animals from the *Foxl2* haploinsufficiency model were

extracted from the dataset. These cells were used to train and evaluate the model. Data were partitioned

into training (75%) and testing (25%) sets using stratified sampling, using the createDataPartition function

from the "caret" package (v. 6.0-91). Training data were used for model optimization, while the test set

served as an independent validation dataset.

Predictive models were trained using L1-regularized lasso regression with the "glmnet" package

(v. 4.1-7) to estimate transcriptional age (in weeks) based on gene expression profiles. Five-fold nested

cross-validation was performed on the training dataset, and the optimal regularization parameter

(lambda.min) was determined based on cross-validation results. Spearman correlation analysis was

performed to assess the correlation between predicted and chronological age.

Transcriptional clocks were applied to experimental test groups, including VCD-treated and

Foxl2$^{+/-}$ animals. Model predictions for test groups were compared to those of control samples to assess

differences in predicted age patterns. Additionally, age acceleration was calculated as the difference

between predicted and chronological age. Statistical significance of group differences in predicted age

and age acceleration was assessed using the Wilcoxon rank-sum test.

To examine the age-associated expression dynamics of lasso features selected by the models,

we performed pseudobulk analysis using control samples. For each cell type, raw counts were

aggregated per sample using muscat (v. 1.18.0) (*53*), and DESeq2 (v. 1.44.0) (*54*) was used to compute

variance-stabilized transformed expression values. Batch effects were corrected using removeBatchEffect,

and expression heatmaps of lasso-selected features were generated to visualize age-dependent

expression trends.

The top 500 lasso features were selected based on the absolute values of their coefficients.

Over-representation analysis was performed using clusterProfiler (v. 4.2.2) and "org.Mm.eg.db" (v. 3.14.0),

analyzing "ALL" Gene Ontology terms. A p-value cutoff of 0.05 was applied to identify significantly

enriched pathways.

1221

Statistical Analysis

All statistical analysis was performed using the R software, version 4.1.2. For all boxplots, the

data is shown with the median, the 25th and 75th percentile of the data, and the whiskers represent 1.5 ∗

the inter-quartile range (IQR). Individual datapoints are overlayed, when possible, for transparency and

rigor. Specific statistical tests, number of biological replicates and animals used are indicated in the

corresponding figure legends and associated methods.

1228

References

1. S. R. Davis *et al.*, Menopause. *Nat Rev Dis Primers* **1**, 15004 (2015).
2. F. J. Broekmans, M. R. Soules, B. C. Fauser, Ovarian aging: mechanisms and clinical consequences. *Endocr Rev* **30**, 465-493 (2009).
3. M. J. Faddy, R. G. Gosden, A model conforming the decline in follicle numbers to the age of menopause in women. *Hum Reprod* **11**, 1484-1486 (1996).
4. M. E. Ossewaarde *et al.*, Age at menopause, cause-specific mortality and total life expectancy. *Epidemiology* **16**, 556-562 (2005).
5. J. S. Hong *et al.*, Age at menopause and cause-specific mortality in South Korean women: Kangwha Cohort Study. *Maturitas* **56**, 411-419 (2007).
6. A. H. Shadyab *et al.*, Association of Accelerometer-Measured Physical Activity With Leukocyte Telomere Length Among Older Women. *J Gerontol A Biol Sci Med Sci* **72**, 1532-1537 (2017).
7. M. R. Gubbels Bupp, Sex, the aging immune system, and chronic disease. *Cell Immunol* **294**, 102-110 (2015).
8. N. Santoro, C. Roeca, B. A. Peters, G. Neal-Perry, The Menopause Transition: Signs, Symptoms, and Management Options. *J Clin Endocrinol Metab* **106**, 1-15 (2021).
9. J. F. Randolph, Jr. *et al.*, Change in follicle-stimulating hormone and estradiol across the menopausal transition: effect of age at the final menstrual period. *J Clin Endocrinol Metab* **96**, 746-754 (2011).
10. T. W. Kelsey, P. Wright, S. M. Nelson, R. A. Anderson, W. H. Wallace, A validated model of serum anti-mullerian hormone from conception to menopause. *PLoS One* **6**, e22024 (2011).
11. G. Weiss, J. H. Skurnick, L. T. Goldsmith, N. F. Santoro, S. J. Park, Menopause and hypothalamic-pituitary sensitivity to estrogen. *JAMA* **292**, 2991-2996 (2004).
12. J. Xiong *et al.*, FSH blockade improves cognition in mice with Alzheimer's disease. *Nature* **603**, 470-476 (2022).
13. L. Mao, L. Wang, S. Bennett, J. Xu, J. Zou, Effects of follicle-stimulating hormone on fat metabolism and cognitive impairment in women during menopause. *Front Physiol* **13**, 1043237 (2022).
14. B. A. Benayoun, A. Kochersberger, J. L. Garrison, Studying ovarian aging and its health impacts: modern tools and approaches. *Genes Dev* **39**, 975-990 (2025).
15. T. A. Van Kempen, T. A. Milner, E. M. Waters, Accelerated ovarian failure: a novel, chemically induced animal model of menopause. *Brain Res* **1379**, 176-187 (2011).
16. C. L. Blenck, P. A. Harvey, J. F. Reckelhoff, L. A. Leinwand, The Importance of Biological Sex and Estrogen in Rodent Models of Cardiovascular Health and Disease. *Circ Res* **118**, 1294-1312 (2016).
17. M. Laszczynska, A. Brodowska, A. Starczewski, M. Masiuk, J. Brodowski, Human postmenopausal ovary--hormonally inactive fibrous connective tissue or more? *Histol Histopathol* **23**, 219-226 (2008).
18. J. Grub, H. Suss, J. Willi, U. Ehlert, Steroid Hormone Secretion Over the Course of the Perimenopause: Findings From the Swiss Perimenopause Study. *Front Glob Womens Health* **2**, 774308 (2021).
19. H. G. Burger, E. C. Dudley, D. M. Robertson, L. Dennerstein, Hormonal changes in the menopause transition. *Recent Prog Horm Res* **57**, 257-275 (2002).

20.   A. A. Mangoni, S. H. Jackson, Age-related changes in pharmacokinetics and
pharmacodynamics: basic principles and practical applications. *Br J Clin Pharmacol* **57**,
6-14 (2004).

21.   I. M. Conboy, T. A. Rando, Heterochronic parabiosis for the study of the effects of aging
on stem cells and their niches. *Cell Cycle* **11**, 2260-2267 (2012).

22.   H. S. Chahal, W. M. Drake, The endocrine system and ageing. *J Pathol* **211**, 173-180
(2007).

23.   A. W. van den Beld *et al.*, The physiology of endocrine systems with ageing. *Lancet*
Diabetes Endocrinol **6**, 647-658 (2018).

24.   T. A. Rando, D. L. Jones, Regeneration, Rejuvenation, and Replacement: Turning Back
the Clock on Tissue Aging. *Cold Spring Harb Perspect Biol* **13**, (2021).

25.   S. A. Villeda *et al.*, The ageing systemic milieu negatively regulates neurogenesis and
cognitive function. *Nature* **477**, 90-94 (2011).

26.   N. H. Uhlenhaut *et al.*, Somatic sex reprogramming of adult ovaries to testes by FOXL2
ablation. *Cell* **139**, 1130-1142 (2009).

27.   B. A. Benayoun, A. Dipietromaria, C. Bazin, R. A. Veitia, FOXL2: at the crossroads of
female sex determination and ovarian function. *Adv Exp Med Biol* **665**, 207-226 (2009).

28.   M. Uda *et al.*, Foxl2 disruption causes mouse ovarian failure by pervasive blockage of
follicle development. *Hum Mol Genet* **13**, 1171-1181 (2004).

29.   D. Schmidt *et al.*, The murine winged-helix transcription factor Foxl2 is required for
granulosa cell differentiation and ovary maintenance. *Development* **131**, 933-942 (2004).

30.   K. Gersak, S. E. Harris, W. J. Smale, A. N. Shelling, A novel 30 bp deletion in the
FOXL2 gene in a phenotypically normal woman with primary amenorrhoea: case report.
Hum Reprod **19**, 2767-2770 (2004).

31.   S. E. Harris *et al.*, Identification of novel mutations in FOXL2 associated with premature
ovarian failure. *Mol Hum Reprod* **8**, 729-733 (2002).

32.   P. Laissue *et al.*, Functional evidence implicating FOXL2 in non-syndromic premature
ovarian failure and in the regulation of the transcription factor OSR2. *J Med Genet* **46**,
455-457 (2009).

33.   L. M. Bilezikjian, W. W. Vale, The Local Control of the Pituitary by Activin Signaling
and Modulation. *Open Neuroendocrinol J* **4**, 90-101 (2011).

34.   S. M. Borman, P. J. Christian, I. G. Sipes, P. B. Hoyer, Ovotoxicity in female Fischer rats
and B6 mice induced by low-dose exposure to three polycyclic aromatic hydrocarbons:
comparison through calculation of an ovotoxic index. *Toxicol Appl Pharmacol* **167**, 191-
198 (2000).

35.   C. E. Finch, The menopause and aging, a comparative perspective. *J Steroid Biochem*
Mol Biol **142**, 132-141 (2014).

36.   M. Kim *et al.*, Estropausal gut microbiota transplant improves measures of ovarian
function in adult mice. *Nat Aging*, (2026).

37.   R. G. Gosden, M. J. Faddy, Ovarian aging, follicular depletion, and steroidogenesis. *Exp*
Gerontol **29**, 265-274 (1994).

38.   C. Lliberos *et al.*, Evaluation of inflammation and follicle depletion during ovarian
ageing in mice. *Sci Rep* **11**, 278 (2021).

39.   A. Sokalska, L. Valentin, Changes in ultrasound morphology of the uterus and ovaries
during the menopausal transition and early postmenopause: a 4-year longitudinal study.
Ultrasound Obstet Gynecol **31**, 210-217 (2008).

40. S. M. Briley *et al.*, Reproductive age-associated fibrosis in the stroma of the mammalian ovary. *Reproduction* **152**, 245-260 (2016).

41. J. V. V. Isola *et al.*, A single-cell atlas of the aging mouse ovary. *Nat Aging* **4**, 145-162 (2024).

42. A. Converse *et al.*, Multinucleated giant cells are hallmarks of ovarian aging with unique immune and degradation-associated molecular signatures. *PLoS Biol* **23**, e3003204 (2025).

43. X. Jiao *et al.*, Ovarian Reserve Markers in Premature Ovarian Insufficiency: Within Different Clinical Stages and Different Etiologies. *Front Endocrinol (Lausanne)* **12**, 601752 (2021).

44. D. R. Danforth *et al.*, Dimeric inhibin: a direct marker of ovarian aging. *Fertil Steril* **70**, 119-123 (1998).

45. M. E. Kevenaar *et al.*, Serum anti-mullerian hormone levels reflect the size of the primordial follicle pool in mice. *Endocrinology* **147**, 3228-3234 (2006).

46. M. Park, D. S. Suh, K. Lee, J. Bae, Positive cross talk between FOXL2 and antimullerian hormone regulates ovarian reserve. *Fertil Steril* **102**, 847-855 e841 (2014).

47. K. Mehalko *et al.*, Lack of accelerated ovarian aging in a follicle-stimulating hormone receptor haploinsufficiency model. *Transl Med Aging* **7**, 1-8 (2023).

48. L. Crisponi *et al.*, The putative forkhead transcription factor FOXL2 is mutated in blepharophimosis/ptosis/epicanthus inversus syndrome. *Nat Genet* **27**, 159-166 (2001).

49. X. Wang *et al.*, Transcriptomic signatures of mouse ovarian aging and estropausal transition at single cell resolution. *bioRxiv*, 2024.2008.2012.607592 (2024).

50. T. Ben Yaakov, T. Wasserman, E. Aknin, Y. Savir, Single-cell analysis of the aged ovarian immune system reveals a shift towards adaptive immunity and attenuated cell function. *Elife* **12**, (2023).

51. M. D. Pisarska, G. Barlow, F. T. Kuo, Minireview: roles of the forkhead transcription factor FOXL2 in granulosa cell biology and pathology. *Endocrinology* **152**, 1199-1208 (2011).

52. J. W. Squair, M. A. Skinnider, M. Gautier, L. J. Foster, G. Courtine, Prioritization of cell types responsive to biological perturbations in single-cell data with Augur. *Nat Protoc* **16**, 3836-3873 (2021).

53. H. L. Crowell *et al.*, muscat detects subpopulation-specific state transitions from multi-sample multi-condition single-cell transcriptomics data. *Nat Commun* **11**, 6077 (2020).

54. M. I. Love, W. Huber, S. Anders, Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol* **15**, 550 (2014).

55. M. E. Morris *et al.*, A single-cell atlas of the cycling murine ovary. *Elife* **11**, (2022).

56. M. Gillespie *et al.*, The reactome pathway knowledgebase 2022. *Nucleic Acids Res* **50**, D687-D692 (2022).

57. D. Saul *et al.*, A new gene set identifies senescent cells and predicts senescence-associated pathways across tissues. *Nat Commun* **13**, 4827 (2022).

58. K. H. Hajdarovic *et al.*, Single-cell analysis of the aging female mouse hypothalamus. *Nat Aging* **2**, 662-678 (2022).

59. J. He *et al.*, Identifying transposable element expression dynamics and heterogeneity during development at the single-cell level with a processing pipeline scTE. *Nat Commun* **12**, 1456 (2021).

1366   60.   J. I. Bravo, S. Nozownik, P. S. Danthi, B. A. Benayoun, Transposable elements, circular
1367   RNAs and mitochondrial transcription in age-related genomic regulation. *Development*
1368   **147**, (2020).
1369   61.   M. De Cecco *et al.*, Transposable elements become active and mobile in the genomes of
1370   aging mammalian somatic tissues. *Aging (Albany NY)* **5**, 867-883 (2013).
1371   62.   P. Langfelder, S. Horvath, WGCNA: an R package for weighted correlation network
1372   analysis. *BMC Bioinformatics* **9**, 559 (2008).
1373   63.   I. M. P. Badia *et al.*, decoupleR: ensemble of computational methods to infer biological
1374   activities from omics data. *Bioinform Adv* **2**, vbac016 (2022).
1375   64.   X. Zhang *et al.*, Enhanced glycolysis in granulosa cells promotes the activation of
1376   primordial follicles through mTOR signaling. *Cell Death Dis* **13**, 87 (2022).
1377   65.   Q. Zhang *et al.*, Mitochondrial and glucose metabolic dysfunctions in granulosa cells
1378   induce impaired oocytes of polycystic ovary syndrome through Sirtuin 3. *Free Radic Biol*
1379   *Med* **187**, 1-16 (2022).
1380   66.   J. N. Mara *et al.*, Ovulation and ovarian wound healing are impaired with advanced
1381   reproductive age. *Aging (Albany NY)* **12**, 9686-9713 (2020).
1382   67.   W. Tang *et al.*, Exploration of the mechanism and therapy of ovarian aging by targeting
1383   cellular senescence. *Life Med* **4**, lnaf004 (2025).
1384   68.   D. K. Dimova, N. J. Dyson, The E2F transcriptional network: old acquaintances with
1385   new faces. *Oncogene* **24**, 2810-2826 (2005).
1386   69.   J. Kim *et al.*, Knockdown of E2F4 suppresses the growth of ovarian cancer cells through
1387   the cell cycle pathway. *Int J Clin Exp Pathol* **14**, 866-874 (2021).
1388   70.   M. C. Meinsohn *et al.*, A role for orphan nuclear receptor liver receptor homolog-1
1389   (LRH-1, NR5A2) in primordial follicle activation. *Sci Rep* **11**, 1079 (2021).
1390   71.   C. H. K. Hughes *et al.*, Steroidogenic factor 1 (SF-1; Nr5a1) regulates the formation of
1391   the ovarian reserve. *Proc Natl Acad Sci U S A* **120**, e2220849120 (2023).
1392   72.   Y. X. Li, X. Guo, T. Gulappa, B. Menon, K. M. J. Menon, SREBP Plays a Regulatory
1393   Role in LH/hCG Receptor mRNA Expression in Human Granulosa-Lutein Cells. *J Clin*
1394   *Endocrinol Metab* **104**, 4783-4792 (2019).
1395   73.   T. Kameda, T. Mizutani, T. Minegishi, Y. Ibuki, K. Miyamoto, Regulation of cAMP
1396   responsive element binding modulator isoforms in cultured rat ovarian granulosa cells.
1397   *Biochim Biophys Acta* **1445**, 31-38 (1999).
1398   74.   M. Shen *et al.*, Involvement of FoxO1 in the effects of follicle-stimulating hormone on
1399   inhibition of apoptosis in mouse granulosa cells. *Cell Death Dis* **5**, e1475 (2014).
1400   75.   S. C. Sharma, J. S. Richards, Regulation of AP1 (Jun/Fos) factor expression and
1401   activation in ovarian granulosa cells. Relation of JunD and Fra2 to terminal
1402   differentiation. *J Biol Chem* **275**, 33718-33728 (2000).
1403   76.   J. Zhou *et al.*, SP1 impacts the primordial to primary follicle transition by regulating
1404   cholesterol metabolism in granulosa cells. *FASEB J* **37**, e22767 (2023).
1405   77.   Y. Ma *et al.*, Androgen Receptor in the Ovary Theca Cells Plays a Critical Role in
1406   Androgen-Induced Reproductive Dysfunction. *Endocrinology* **158**, 98-108 (2017).
1407   78.   E. Pelosi *et al.*, Constitutively active Foxo3 in oocytes preserves ovarian reserve in mice.
1408   *Nat Commun* **4**, 1843 (2013).
1409   79.   E. Havis, D. Duprez, EGR1 Transcription Factor is a Multifaceted Regulator of Matrix
1410   Production in Tendons and Other Connective Tissues. *Int J Mol Sci* **21**, (2020).

| 1411 | 80. | Z. Yang *et al.*, Reprogramming of stromal fibroblasts by SNAI2 contributes to tumor |
| 1412 | | desmoplasia and ovarian cancer progression. *Mol Cancer* **16**, 163 (2017). |

80. Z. Yang *et al.*, Reprogramming of stromal fibroblasts by SNAI2 contributes to tumor
desmoplasia and ovarian cancer progression. *Mol Cancer* **16**, 163 (2017).
81. S. Avraham *et al.*, ATF3 and JDP2 deficiency in cancer associated fibroblasts promotes
tumor growth via SDF-1 transcription. *Oncogene* **38**, 3812-3823 (2019).
82. J. Hsu, J. Sage, Novel functions for the transcription factor E2F4 in development and
disease. *Cell Cycle* **15**, 3183-3190 (2016).
83. M. Mussbacher *et al.*, Cell Type-Specific Roles of NF-kappaB Linking Inflammation and
Thrombosis. *Front Immunol* **10**, 85 (2019).
84. E. D. Lucas *et al.*, Type 1 IFN and PD-L1 Coordinate Lymphatic Endothelial Cell
Expansion and Contraction during an Inflammatory Immune Response. *J Immunol* **201**,
1735-1747 (2018).
85. X. Sun *et al.*, Role of post-translational modifications of Sp1 in cancer: state of the art.
Front Cell Dev Biol **12**, 1412461 (2024).
86. M. T. Buckley *et al.*, Cell-type-specific aging clocks to quantify aging and rejuvenation in
neurogenic regions of the brain. *Nat Aging* **3**, 121-137 (2023).
87. M. J. Peters *et al.*, The transcriptional landscape of age in human peripheral blood. *Nat*
Commun **6**, 8570 (2015).
88. B. S. Ellsworth *et al.*, FOXL2 in the pituitary: molecular, genetic, and developmental
analysis. *Mol Endocrinol* **20**, 2796-2805 (2006).
89. L. Komsta. (2022).
90. T. Therneau. (2022).
91. L. Ongaro *et al.*, Development of a Highly Sensitive ELISA for Measurement of FSH in
Serum, Plasma, and Whole Blood in Mice. *Endocrinology* **162**, (2021).
92. M. Kuhn, caret: Classification and Regression Training. (2022).
93. L. Breiman, A. Cutler, A. Liaw, M. Wiener.
94. G. X. Zheng *et al.*, Massively parallel digital transcriptional profiling of single cells. *Nat*
Commun **8**, 14049 (2017).
95. J. Campbell, S. Yang, Z. Wang, S. Corbett, Y. Koga. (2023).
96. . (2022).
97. A. Bais, D. Kostkam, scds: Computational Annotation of Doublets in Single Cell RNA
Sequencing Data. (2019).
98. Y. Hao *et al.*, Integrated analysis of multimodal single-cell data. *Cell* **184**, 3573-3587
e3529 (2021).
99. I. Korsunsky *et al.*, Fast, sensitive and accurate integration of single-cell data with
Harmony. *Nat Methods* **16**, 1289-1296 (2019).
100. M. Andreatta, A. J. Berenstein, S. J. Carmona, scGate: marker-based purification of cell
types from heterogeneous single-cell RNA-seq datasets. *Bioinformatics* **38**, 2642-2644
(2022).
101. D. Aran *et al.*, Reference-based analysis of lung single-cell sequencing reveals a
transitional profibrotic macrophage. *Nat Immunol* **20**, 163-172 (2019).
102. H. Guo, J. Li, scSorter: assigning cells to known cell types according to marker genes.
Genome Biol **22**, 69 (2021).
103. K. Nader *et al.*, ScType enables fast and accurate cell type identification from spatial
transcriptomics data. *Bioinformatics* **40**, (2024).

104. T. S. Heng, M. W. Painter, C. Immunological Genome Project, The Immunological
Genome Project: networks of gene expression in immune cells. *Nat Immunol* **9**, 1091-
1094 (2008).
105. S. A. Miller *et al.*, LSD1 and Aberrant DNA Methylation Mediate Persistence of
Enteroendocrine Progenitors That Support BRAF-Mutant Colorectal Cancer. *Cancer Res*
81, 3791-3805 (2021).
106. J. Leek *et al.*
107. M. E. Ritchie *et al.*, limma powers differential expression analyses for RNA-sequencing
and microarray studies. *Nucleic Acids Res* **43**, e47 (2015).
108. T. Wu *et al.*, clusterProfiler 4.0: A universal enrichment tool for interpreting omics data.
Innovation (Camb) **2**, 100141 (2021).
109. I. Dolgalev. (2021).
110. L. Pantano. (2024).
111. J. F. Ouyang, U. S. Kamaraj, E. Y. Cao, O. J. L. Rackham, ShinyCell: simple and sharable
visualization of single-cell gene expression data. *Bioinformatics* **37**, 3374-3376 (2021).
1470
1471
1472

Acknowledgements

We thank Dr. Gilbert Garcia for guidance and feedback on confocal imaging. We are grateful to Dr. Lynae M. Brayboy for providing the ovarian single-cell dissociation protocol and for valuable advice on protocol optimization. We thank Kristen Mehalko for maintaining the Foxl2 haploinsufficiency mouse colony and assisting with tissue collection. We acknowledge Younggyun Kim, Kelly Koh, and Sanjana Paye for assistance with follicle quantification from hematoxylin and eosin-stained ovarian sections. We thank Dr. Victor Ansere, Clayton Baker, and Aaron J.J. Lemus for constructive feedback on the manuscript.

Ovarian histological processing was performed by the Translational Pathology Core at the USC Norris Comprehensive Cancer Center and Immunohistochemistry Laboratory at USC Labs, and serum hormone measurements were conducted by the University of Virginia Center for Research in Reproduction Ligand Assay and Analysis Core. Some schematic panels were generated using the Generic Diagramming Platform.

Funding

Global Consortium for Reproductive Longevity and Equality Postdoctoral Fellowship GCRLE-2020 supported by the Bia-Echo Foundation (MK)

USC Provost's Undergraduate Research Fellowship (JW)

Global Consortium for Reproductive Longevity and Equality Junior Scholar Award GCRLE-0520 supported by the Bia-Echo Foundation (BAB)

Pew Biomedical Scholar Award #00034120 from The Pew Charitable Trusts (BAB)

Generous gifts from Kathleen Gilmore and Dr. Eric Hennigan (BAB)

National Cancer Institute Cancer Center Support Grant P30 CA014089 supporting the Translational Pathology Core at the USC Norris Comprehensive Cancer Center

Eunice Kennedy Shriver National Institute of Child Health and Human Development grant R24HD102061 supporting the University of Virginia Center for Research in Reproduction Ligand Assay and Analysis Core

Author contribution

Conceptualization: MK, BAB

Investigation: MK, RB, JW, XL, MC, RJL, EHL, JLA, RGW, BAB

Formal analysis: MK, RB, JW, EHL, BAB

Visualization: MK, JW, BAB

Supervision: BAB

Writing - original draft: MK, BAB

Writing - review & editing: MK, RB, JW, XL, RJL, EHL, JLA, RGW, BAB

1509

1510 **Competing interest**

1511 Authors declare that they have no competing interests.

1512

Data availability

The raw FASTQ files have been deposited in the Sequence Read Archive under accession

number PRJNA863443. Raw microscopy images of ovarian hematoxylin and eosin (H&E), Picrosirius

Red and Sudan Black B staining, compressed RNAscope images (due to file size constraints), and raw

flow cytometry datasets are available on Figshare (DOIs: 10.6084/m9.figshare.28745765,

10.6084/m9.figshare.28745792, 10.6084/m9.figshare.28745813, 10.6084/m9.figshare.29260391 and

10.6084/m9.figshare.28733222). Any additional information required to reproduce or reanalyze the data

presented in this study is available from the corresponding author upon reasonable request.

All new scripts used to analyze the datasets are available on the Benayoun lab GitHub at

https://github.com/BenayounLaboratory/Mouse_Menopause_Models.

1523

1524   **List of Supplementary Materials**

1525

1526   Figs. S1 to S22

1527   Tables S1 to S11

1528

1529 **Figures**

1530

Fig. 1. Ovarian follicle count quantification of animals from Aging, VCD and *Foxl2*

haploinsufficiency models. (A) Schematic of the experimental design. **(B-D)** Representative

hematoxylin-eosin staining images of ovarian tissues from Aging, VCD, and *Foxl2* haploinsufficiency

model mice. For the VCD model, ovarian tissues were analyzed 5 months post-injection. Scale bar: 250

µm. **(E-G)** Combined follicle counts (primordial, primary, secondary and antral follicles, and corpus luteum)

from Aging (n = 25 per group), VCD (n = 5-10 per group), and *Foxl2* haploinsufficiency (n = 11-24 per

group) model mice. Statistical significance was assessed using Wilcoxon test for the Aging model and

aligned rank transform ANOVA to test effects for the VCD and *Foxl2* haploinsufficiency models; p-values

are shown. Open data points indicate data used for ovarian health index calculation.

1540

Fig. 2. Quantification of ovarian fibrosis and lipofuscin accumulation across Aging, VCD, and

***Foxl2* haploinsufficiency models. (A)** Schematic of the experimental design. Picrosirius Red and Sudan

Black B staining were performed and quantified to assess fibrosis and lipofuscin accumulation in the

ovaries. **(B-D)** Representative Picrosirius Red staining images of ovarian tissues from Aging, VCD, and

Foxl2 haploinsufficiency model mice. Original and processed image are shown. **(E-G)** Quantification of

percent fibrotic area across Aging (n = 6 and 4 for Young and Old, respectively), VCD (n = 4-5 per group),

and *Foxl2* haploinsufficiency (n = 4-7 per group) model mice. **(H-J)** Representative Sudan Black B

staining images showing lipofuscin accumulation in ovarian tissues from Aging, VCD, and *Foxl2*

haploinsufficiency model mice. Original and processed image are shown. **(K-M)** Quantification of percent

lipofuscin-positive area across Aging (n = 5 and 6 for Young and Old, respectively), VCD (n=4-5 per

group), and *Foxl2* haploinsufficiency (n = 4-7 per group) model mice. For the VCD model, ovarian tissues

were analyzed 5 months post-injection. Scale bar: 200 µm. For (**E-G** and **K-M**), statistical significance

was assessed using the Wilcoxon test for the Aging model and aligned rank transform ANOVA to test

effects for the VCD and *Foxl2* haploinsufficiency models; p-values are shown.

1555

Fig. 3. Ovarian hormone marker quantification of animals from Aging, VCD and *Foxl2*
haploinsufficiency models. (A) Schematic of the experimental design. Serum levels of AMH, FSH and
INHBA were quantified via ELISA. Serum hormone levels and combined follicle counts were integrated to
calculate the ovarian health index. **(B-D)** Serum AMH level quantification from Aging (n = 10 per group),
VCD (n = 5 per group), and *Foxl2* haploinsufficiency (n = 10-21 per group) model mice. **(E-G)** Serum FSH
level quantification from Aging (n = 10 per group), VCD (n = 5 per group), and *Foxl2* haploinsufficiency (n
= 10-21 per group) model mice. **(H-J)** Serum INHBA level quantification from Aging (n = 10 per group),
VCD (n = 5 per group), and *Foxl2* haploinsufficiency (n = 10-21 per group) model mice. **(K-M)** Ovarian
health index from Aging (n = 5 per group), VCD (n = 4-5 per group), and *Foxl2* haploinsufficiency (n = 4-
10 per group) model mice. Statistical significance was assessed using Wilcoxon test for the Aging model
and aligned rank transform ANOVA to test effects for the VCD and *Foxl2* haploinsufficiency models; p-
values are shown. Open data points indicate data used for ovarian health index calculation.

1568

Fig. 4. Development and application of OvAge, a hormone-based ovarian aging clock. (A)
Schematic representation of the OvAge clock development pipeline. Data from all animals from the Aging
model, vehicle control (CTL) from the VCD model, *Foxl2*$^{+/+}$ from the *Foxl2* haploinsufficiency model, and
Fshr$^{+/+}$ from the *Fshr* haploinsufficiency model mice were used to train and test OvAge. **(B)** Scatter plot
comparing OvAge-predicted and chronological ages, reported in weeks, in the test dataset (n = 68). **(C)**
Bar plot of scaled feature importance for OvAge model. **(D,E)** Scatter plots comparing OvAge-predicted
and chronological ages for VCD (n = 147) and *Foxl2* haploinsufficiency (n = 62) model mice. **(F-H)** Age
acceleration analysis for VCD model animals at 30-, 90- and 150-days post-injection (n = 4-5 per group).
(I) Age acceleration analysis for *Foxl2* haploinsufficiency animals (n = 10-21 per group). Age acceleration
was measured as the difference between predicted OvAge and chronological age. Open data points
indicate data from out-of-bag predictions. For **(F-I)**, statistical significance was assessed using Wilcoxon
test for the Aging model and aligned rank transform ANOVA to test effects for the VCD and *Foxl2*
haploinsufficiency models; p-values are shown.

1582

Fig 5. scRNA-seq profiling of ovaries from Aging, VCD, and *Foxl2* haploinsufficiency model mice.

(A) Schematic of the experimental design. **(B)** Schematic of sample collection timepoints. **(C-E)** UMAP plots of scRNA-seq datasets from Aging, VCD and *Foxl2* haploinsufficiency models. **(F-H)** Dotplot representation of expression of marker genes of cell types detected in scRNA-seq datasets from Aging, VCD and *Foxl2* haploinsufficiency models. **(I)** Representative RNAscope images of DAPI, *Klrb1c*, *Cd3e*, *Cd8b1* and *Cd4* probes. Images were enhanced for visualization. The shown image is from a middle-aged *Foxl2*$^{+/+}$ animal.

Fig 6. Characterization of cell type proportions in Aging, VCD, and *Foxl2* haploinsufficiency model scRNA-seq datasets. (A) Schematic of the experimental design. **(B-D)** Proportion differences of *Ptprc*$^{-}$ and *Ptprc*$^{+}$ cells between young vs. old in Aging model, CTL (vehicle control) vs. VCD in VCD model, and *Foxl2*$^{+/+}$ vs. *Foxl2*$^{+/-}$ in the *Foxl2* haploinsufficiency model datasets. Colored points indicate statistically significant differences (p-value < 0.05) assessed using scProportionTest, whereas gray points denote non-significant comparisons. **(E-G)** Flow cytometry analysis of CD45^{+} cell proportions in ovaries of Aging, VCD, and *Foxl2* haploinsufficiency model mice. **(H-J)** Proportion differences of non-immune cell types detected in Aging, VCD, and *Foxl2* haploinsufficiency model scRNA-seq datasets. Color coding as in panels **(B-D)**. **(K-M)** Representative RNAscope images of *Fshr* (granulosa) and *Cyp11a1* (theca) probes from Aging, VCD, and *Foxl2* haploinsufficiency model mice. Scale bar: 100μm. **(N-P)** Cell abundance quantification data from RNAscope image analysis of *Fshr*$^{+}$ and *Cyp11a1*$^{+}$ cells from Aging, VCD, and *Foxl2* haploinsufficiency model mice. For panels **(E-G)** and **(N-P)**, statistical significance was assessed using Wilcoxon test for the Aging model and aligned rank transform ANOVA to test effects for the VCD and *Foxl2* haploinsufficiency models; p-values are shown.

Fig 7. Pseudobulk analysis of differential gene expression and gene ontology analysis across Aging, VCD, and *Foxl2* haploinsufficiency models. (A) Schematic of pseudobulk analysis pipeline. **(B-D)** Strip plots of differentially expressed genes (DEGs) from Aging, VCD and *Foxl2* haploinsufficiency models. DEGs were identified using DESeq2, comparing young vs. old (Aging model), CTL vs. VCD (VCD model), and *Foxl2*$^{+/+}$ vs. *Foxl2*$^{+/-}$ (*Foxl2* haploinsufficiency model). Genes that passed the FDR <

1611   0.05 threshold are colored and the numbers in parentheses following each cell type indicate the total

1612   number of DEGs that met the significance threshold. **(E-F)** Frequent Gene Set Enrichment Analysis

1613   (GSEA) term enrichment for GO Biological Process (GO BP, FDR < 0.1) and Reactome (FDR < 0.1)

1614   pathways across Aging, VCD, and *Foxl2* haploinsufficiency models.

1615

Fig. 8. Age-associated gene expression trajectory analysis reveals model-specific amplification

patterns in granulosa cells. (A) Schematic illustrating analytical workflow. Likelihood ratio tests (LRT)

and unsupervised clustering of genes were performed using DESeq2 and degPatterns, respectively, to

identify genes with significant expression changes across experimental groups in the VCD and *Foxl2*

haploinsufficiency models. Clusters were designated as enhanced increase (EI) or enhanced decrease

(ED) when perturbations amplified age-associated expression changes. Over-representation analysis

(ORA) was performed for each cluster via clusterProfiler. **(B,C)** Boxplots showing normalized gene

expression patterns of genes from granulosa EI and ED clusters in the VCD model. Bubble plots display

ORA results for enriched pathways for each cluster. **(D,E)** Boxplots showing normalized gene expression

patterns of representative genes from granulosa EI and ED clusters in the *Foxl2* haploinsufficiency model.

Bubble plots show ORA enrichment results.

1627

Fig 9. Development of a transcriptome-based aging clock for granulosa cells. **(A)** Schematic of

analysis and training workflow for the lasso regression-based clock. **(B)** Scatter plot of predicted age *vs*.

chronological age for the test set, with Spearman's correlation and p-value displayed. **(C,D)** Age

prediction for VCD and *Foxl2* haploinsufficiency datasets, comparing predicted age to chronological age

in weeks. Offsets were added to improve visualization. **(E,F)** Age acceleration analysis for VCD and *Foxl2*

haploinsufficiency datasets, calculated as the difference between predicted and chronological age.

Statistical significance was assessed using aligned rank transform ANOVA to test effects. **(G)** Heatmap of

expression lasso features from pseudobulked expression data. **(H)** Over-representation analysis of GO

ALL terms of the top 500 lasso features.

 Supplementary figures

 Fig. S1. Characterization of the *Foxl2* haploinsufficiency model. (A) Schematic representation of

 constructs for *Foxl2* haploinsufficiency mouse line generation. **(B)** *Foxl2* expression levels in *Foxl2*$^{+/+}$ and

 Foxl2$^{+/-}$ mice ovaries from young and mid-age groups, measured by RT-qPCR (n = 5 per group). P-values

 were calculated using aligned rank transform ANOVA to test effects of genotype and age. **(C,D)** Fertility

 assessment of *Foxl2* haploinsufficiency females, comparing first litter pup counts and latency to first

 pregnancy (n = 11 and 15 for *Foxl2*$^{+/+}$ and *Foxl2*$^{+/-}$, respectively). P-values were calculated using the

 Wilcoxon and log-rank tests, respectively.

 Fig. S2. Follicle counts across Aging, VCD and *Foxl2* haploinsufficiency models. (A) Follicle

 counting scheme. Pr: primordial follicle; P: primary follicle; S: secondary follicle; A: antral follicle; CL:

 corpus luteum. **(B-D)** Follicle counts for primordial, primary, secondary and antral follicles, and corpus

 luteum, from Aging (n = 25 per group), VCD (n = 5-10 per group), and *Foxl2* haploinsufficiency (n = 11-24

 per group) model mice. For panels **(B-D)**, statistical significance was assessed using Wilcoxon test for the

 Aging model and aligned rank transform ANOVA to test effects for the VCD and *Foxl2* haploinsufficiency

 models; p-values are shown.

 Fig. S3. Harmonization of FSH results for the *Foxl2* haploinsufficiency model and longitudinal

 analysis of serum AMH, FSH, and Inhibin A levels in the VCD model. (A) Polynomial regression-

 based correction model used to harmonize FSH measurements obtained from two different assay kits (n

 = 6 and 9 for *Foxl2*$^{+/+}$ and *Foxl2*$^{+/-}$, respectively). **(B-D)** Serum levels of AMH, FSH, and INHBA measured

 from 0 to 5 months post-injection in VCD model mice (n = 5 per group).

 Fig. S4. Histological characterization of the VCD model mice. (A) Representative hematoxylin and

 eosin staining images of ovarian tissues from condition-matched animals used in the VCD model scRNA-

 seq analysis. Scale bar: 250µm. **(B,C)** Follicle counts for the condition-matched animals used in the VCD

1664   model scRNA-seq analysis (n = 4-5 per group). Statistical significance was assessed using aligned rank

1665   transform ANOVA; p-values are shown.

1666

Fig. S5. Condition-matched analysis of ovarian fibrosis and lipofuscin accumulation following

VCD exposure. (A,B) Representative Picrosirius Red staining images of ovarian tissues collected at 30

days and 90 days post-VCD injection. **(C,D)** Quantification of percent fibrotic area at 30 and 90 days post-

injection (n = 5 per group). **(E,F)** Representative Sudan Black B staining images showing lipofuscin

accumulation in ovarian tissues at 30 and 90 days post-injection. **(G,H)** Quantification of percent

lipofuscin-positive area at 30 and 90 days post-injection (n = 4-5 per group). Scale bar: 200 μm. For

(C,D,G,H), statistical significance was assessed using aligned rank transform ANOVA; p-values are

shown.

1675

Fig. S6. Validation of detected ovarian cell types by *in situ* hybridization. (A) Representative

RNAscope images for *Fshr, Cyp11a1, Pdgfra* and *Acta2* probes. **(B)** Representative RNAscope images

for *Upk1b, Cd19, Flt1* and *Prox1* probes. **(C)** Representative RNAscope images for *Klrb1c, Cd3e, Cd8b2*

and *Cd4*. Images were enhanced for visualization. **(D)** Heatmap of detected cell types in the Aging, VCD

and *Foxl2* haploinsufficiency scRNA-seq datasets. For panels **(A-D)**, shown images are from a middle-

aged *Foxl2*$^{+/+}$ animal.

1682

Fig. S7. Cell type proportion analysis of ovarian cells via flow cytometry and scRNA-seq analysis.

(A) Gating strategy of CD45^{+} cells in flow cytometry analysis. **(B-D)** Proportion differences of *Ptprc*+ cell

types detected in Aging, VCD and *Foxl2* haploinsufficiency model scRNA-seq datasets. Colored points

indicate statistically significant differences (p-value < 0.05) assessed using scProportionTest, whereas

gray points denote non-significant comparisons. Asterisks (*) indicate comparisons in which the observed

log$_2$ fold change is infinite due to complete absence of the cell type in one of the groups.

1689

1690   **Fig. S8. Cell type proportion analysis of non-immune ovarian cells via *in situ* hybridization assays.**

1691   **(A-C)** Cell type abundance analysis of stromal, SMC, BEC, LEC and epithelial cells via RNAscope *in situ*

1692   hybridization assays from Aging, VCD and *Foxl2* haploinsufficiency model mice (n = 4-5 per group across

1693   models). Statistical significance was assessed using Wilcoxon test for the Aging and VCD models, and

1694   aligned rank transform ANOVA for the *Foxl2* haploinsufficiency model; p-values are shown.

1695

Fig. S9. Cell type proportion analysis of immune ovarian cells via *in situ* hybridization assays. (A-

C) Cell type abundance analysis of NK, NKT, CD8$^+$ NKT, CD8$^+$ T, CD4$^+$ T, DNT, DPT and B cells via

RNAscope *in situ* hybridization assays from Aging, VCD and *Foxl2* haploinsufficiency model mice (n = 4-5

per group across models). Statistical significance was assessed using Wilcoxon test for the Aging and

VCD models, and aligned rank transform ANOVA for the *Foxl2* haploinsufficiency model; p-values are

shown.

1702

Fig. S10. Comparative analysis of global gene expression in the Aging and VCD models via Augur.

(A) Schematic of global gene expression analysis comparisons using Augur. **(B,C)** UMAP visualization of

AUC scores and lollipop plot of AUC quantification from the Aging model. **(D-G)** UMAP visualization of

AUC scores and scatter plot of AUC quantification from the VCD model, comparing CTL vs. VCD at 3m

and 10m age-at-injection and 30 days vs. 90 days post-injection. For the scatter plot, Ptprc- and Ptprc+

cells were separately plotted to improve visualization. For **(E,G)**, data points with NA AUC scores (due to

low cell count or QC filtering) were assigned an AUC of 0.5 and colored gray to improve visualization.

1710

Fig. S11. Comparative analysis of global gene expression in the *Foxl2* haploinsufficiency model

via Augur. (A-E) UMAP visualization of AUC scores and scatter plot of AUC quantification from the *Foxl2*

haploinsufficiency model comparing *Foxl2*$^{+/+}$ vs. *Foxl2*$^{+/-}$ at young, mid-age and old. For the scatter plot,

Ptprc$^-$ and *Ptprc*$^+$ cells were separately plotted to improve visualization. For **(C,E)**, data points with NA

AUC scores (due to low cell count or QC filtering) were assigned an AUC of 0.5 and colored gray to

improve visualization.

1717

1718   **Fig. S12. Cell type representation across pseudobulked scRNA-seq datasets and validation of**

1719   **Foxl2 expression in Foxl2 haploinsufficiency mice. (A)** Heatmap of cell types detected in

1720   pseudobulked scRNA-seq datasets. **(B)** DESeq2 normalized $\log_2$ counts of *Foxl2* from young, mid-age

1721   and old *Foxl2*$^{+/+}$ and *Foxl2*$^{+/-}$ mice.

1722

Fig. S13. Assessment of estrous cycle effects on ovarian transcriptional profiles. (A) Overview of a

publicly available single-cell RNA-seq dataset profiling mouse ovaries across the four estrous cycle

stages. Granulosa cell transcriptomes were pseudobulked for downstream analyses. **(B)** Multidimensional

scaling plots of pseudobulked granulosa cell transcriptomes from Cycling, Aging, VCD and *Foxl2*

haploinsufficiency model data. **(C)** Differential expression analysis (FDR < 0.05) between estrous cycle

dataset (estrus vs. proestrus) and menopause model dataset.

1729

1730   **Fig. S14. Aging-associated signatures and transposable element enrichment analysis across**

1731   **Aging, VCD and *Foxl2* haploinsufficiency models. (A,B)** GSEA enrichment analysis of differentially

1732   expressed genes dentified in the Aging model (FDR < 0.05) and SenMayo gene list. **(C)** Dot plot showing

1733   the coefficient of variation analysis results. **(D)** GSEA enrichment analysis of transposable element (TE)

1734   families across Aging, VCD and *Foxl2* haploinsufficiency models.

1735

1736   **Fig. S15. Classification of age-associated transcriptional trajectory patterns in granulosa cells. (A)**

1737   Boxplot showing normalized gene expression patterns of genes from granulosa cluster 1 in the VCD

1738   model (VG Cluster 1). Bubble plot displays ORA results for enriched pathways for the cluster. **(C)** Boxplot

1739   showing normalized gene expression patterns of genes from granulosa cluster 2 in the *Foxl2*

1740   haploinsufficiency model (FG Cluster 2). Bubble plot shows ORA enrichment results.

1741

1742   **Fig. S16. Age-associated transcriptional trajectory patterns and pathway enrichment in theca cells.**

1743   **(A-C)** Boxplots showing normalized gene expression patterns of genes from theca EI, ED 1 and ED 2

1744   clusters in the VCD model. Bubble plots display ORA results for enriched pathways for each cluster. **(D)**

1745   Boxplot showing normalized gene expression patterns of genes from theca ED cluster in the *Foxl2*

1746   haploinsufficiency model. Bubble plot shows ORA enrichment results.

1747

Fig. S17. Age-associated transcriptional trajectories in *Foxl2* haploinsufficient theca cells. (A,B) Boxplots showing normalized gene expression patterns of genes from theca clusters 1 and 2 in the *Foxl2* haploinsufficiency model. No enrichment was identified via ORA.

Fig. S18. Module trees of WGCNA analysis. (A) Schematic of Weighted Gene Co-expression Network Analysis (WGCNA) and transcription factor inference analysis. **(B)** WGCNA module trees identified in the Aging model for granulosa, theca, stromal, BEC, LEC, epithelial, and DNT cells.

Fig. S19. Association analysis of WGCNA modules and ovarian aging traits from the Aging model. (A) Heatmap of trait significance for WGCNA modules in the Aging model. Modules that passed the FDR < 0.1 threshold are shown in bold.

Fig. S20. Preservation of co-expression modules in Aging, VCD, and *Foxl2* haploinsufficiency models. (A) GSEA enrichment of WGCNA module genes associated with aging traits across Aging, VCD and *Foxl2* haploinsufficiency models. **(B-D)** Bubble plots of over-representation analysis (ORA) results of GO ALL terms for WGCNA module genes (FDR < 0.1).

Fig. S21. Analysis of recurrent transcription factor activity inferred by decoupleR. (A) Heatmap of transcription factor (TF) activity for factors detected in at least three cell types within each model with an FDR < 0.05. (B) (C) **(B)** UpSet plot of decoupleR-inferred TF activity, showing the number of TF overlap across datasets and the corresponding cell types. **(C)** Heatmap of TF activity scores of common decoupleR-inferred TFs for each cell type.

Fig. S22. Development of a transcriptome-based aging clock for theca cells. **(A)** Schematic of analysis and training workflow for the lasso regression-based clock. **(B)** Scatter plot of predicted age *vs*. chronological age for the test set, with Spearman's correlation and p-value displayed. **(C,D)** Age prediction for VCD and *Foxl2* haploinsufficiency datasets, comparing predicted age to chronological age in weeks. Offsets were added for visualization purposes. **(E,F)** Age acceleration analysis for VCD and

1776   *Foxl2* haploinsufficiency datasets, calculated as the difference between predicted and chronological age.

1777   Statistical significance was assessed using aligned rank transform ANOVA to test effects. **(G)** Heatmap of

1778   expression lasso features from pseudobulked expression data. **(H)** Over-representation analysis of GO

1779   ALL terms of the top 500 lasso features.

1780   **Supplementary tables**

1781

Supplementary table 1. Raw fertility data from *Foxl2* haploinsufficiency model mice.

Supplementary table 2. Raw serum AMH, FSH and INHBA quantification data. Raw data from **(A)**

Aging, **(B)** VCD and **(C)** *Foxl2* haploinsufficiency models.

Supplementary table 3. BH-adjusted post hoc p-values for OvAge age acceleration in VCD and

Foxl2 models.

Supplementary table 4. 10xGenomics CellRanger QC metrix.

Supplementary table 5. List of canonical marker genes used to annotate cell types in scRNA-seq

datasets.

Supplementary table 6. Relative proportion of detected cell types from scRNA-seq datasets.

Proportion data from **(A)** Aging, **(B)** VCD and **(C)** *Foxl2* haploinsufficiency models.

Supplementary table 7. DESeq2 output from pseudobulk analysis. DESeq2 output from **(A)** Aging, **(B)**

VCD and **(C)** *Foxl2* haploinsufficiency model datasets.

Supplementary table 8. Significant GSEA enrichment terms (FDR < 10%) for GO Biological

Process and Reactome terms. GSEA enrichment terms for Biological Process from **(A)** Aging, **(B)** VCD

and **(C)** *Foxl2* haploinsufficiency model datasets. GSEA enrichment terms for Reactome from **(D)** Aging,

(E) VCD and **(F)** *Foxl2* haploinsufficiency model datasets.

Supplementary table 9. LRT DESeq2 output with cluster information. LRT DESeq2 output with

degPatterns cluster information from **(A)** VCD model granulosa cells, **(B)** *Foxl2* haploinsufficiency model

granulosa cells, **(C)** VCD model theca cells and **(D)** *Foxl2* haploinsufficiency model theca cells.

Supplementary table 10. List of decoupleR TF activity scores (FDR < 5%) for each cell type.

Supplementary table 11. Primers used in the study. a, Primers used for genotyping *Foxl2*

haploinsufficiency model animals. **b**, Primers used for *Foxl2* expression quantification in *Foxl2*

haploinsufficiency model animals via RT-qPCR.

Fig. 1

A Experimental scheme

Aging	VCD	*Foxl2 +/-*
Young *vs* Old (4 months) (20 months)	CTL *vs* VCD (Safflower oil)	*Foxl2⁺/⁺* *vs* *Foxl2⁺/⁻*

Ovarian health assessment
▪ Ovarian follicle count
▪ Ovarian fibrosis and lipofuscin quantification
▪ Serum AMH, FSH & INHBA concentrations
▪ Ovarian health index

Single-cell transcriptomic analysis

B Hematoxylin-eosin staining: Aging

Age
Young | Old

C Hematoxylin-eosin staining: VCD - 150d post-injection

Age-at-injection

	3m	6m	8m	10m
CTL				
VCD				

D Hematoxylin-eosin staining: *Foxl2 +/-*

Age

	3-4m	8-10m	12-13m
Foxl2⁺/⁺			
Foxl2⁺/⁻			

E Combined follicle count: Aging

Age: p ~2.73e-13

Follicle count (count / section)

Young | Old

F Combined follicle count: VCD - 150d post-injection

Treatment: p ~1.40e-13
Age: p ~2.25e-5

Follicle count (count / section)

CTL VCD | CTL VCD | CTL VCD | CTL VCD
3m | 6m | 8m | 10m
Age-at-injection

G Combined follicle count: *Foxl2 +/-*

Genotype: p ~2.24e-1
Age: p ~2.86e-10

Follicle count (count / section)

Foxl2⁺/⁺ *Foxl2⁺/⁻* | *Foxl2⁺/⁺* *Foxl2⁺/⁻* | *Foxl2⁺/⁺* *Foxl2⁺/⁻*
3-4m | 8-10m | 12-13m

Fig. 2

A Experimental scheme

Ovarian tissue collection & fixation → Picrosirius Red staining → Ovarian fibrosis quantification
→ Sudan Black B staining → Ovarian lipofuscin quantification

B Picrosirius Red: Aging

C Picrosirius Red: VCD

D Picrosirius Red: *Foxl2 +/-*

E Fibrosis quantification: Aging

F Fibrosis quantification: VCD

G Fibrosis quantification: *Foxl2 +/-*

H Sudan Black B: Aging

I Sudan Black B: VCD

J Sudan Black B: *Foxl2 +/-*

K Lipofuscin quantification: Aging

L Lipofuscin quantification: VCD

M Lipofuscin quantification: *Foxl2 +/-*

Fig. 3

A Experimental scheme

Serum collection → Quantification of AMH, FSH & INHBA via ELISA → Ovarian health index analysis

Combined follicle counts

B Serum AMH levels: Aging
Age: p ~1.63e-4

C Serum AMH levels: VCD
Treatment: p ~1.45e-6
Age: p ~1.99e-4

D Serum AMH levels: *Foxl2 +/-*
Genotype: p ~1.52e-2
Age: p ~3.41e-5

E Serum FSH levels: Aging
Age: p ~1.15e-2

F Serum FSH levels: VCD
Treatment: p ~3.42e-8
Age: p ~5.17e-1

G Serum FSH levels: *Foxl2 +/-*
Genotype: p ~3.11e-2
Age: p ~4.52e-2

H Serum INHBA levels: Aging
Age: p ~2.12e-3

I Serum INHBA levels: VCD
Treatment: p ~7.77e-12
Age: p ~1.90e-2

J Serum INHBA levels: *Foxl2 +/-*
Genotype: p ~5.21e-2
Age: p ~2.49e-1

K Ovarian health index: Aging
Age: p ~1.09e-2

L Ovarian health index: VCD
Treatment: p ~1.74e-10
Age: p ~2.36e-2

M Ovarian health index: *Foxl2 +/-*
Genotype: p ~3.65e-1
Age: p ~3.69e-3

Fig. 4

A OvAge clock training scheme

C57BL/6J, NTac
(3-20-months)

Aging model: All
VCD model: CTL
Foxl2 +/- model: *Foxl2*+/+
Fshr +/- model: *Fshr*+/+

AMH
FSH
INHBA

Serum ELISA

Training
(n = 212)

Testing
(n=68)

Random forest-based
OvAge clock

B OvAge performance

Rho ~0.795
p ~5.5e-16

Predicted age (weeks)
Chronological age (weeks)

● Aging model   (n=17)
▲ VCD model: CTL   (n=42)
◆ *Foxl2* +/- model: *Foxl2*+/+ (n=7)
■ *Fshr* +/- model: *Fshr*+/+   (n=2)

C Feature importance

Importance (scaled)

AMH   INHBA   FSH

D Age prediction: VCD

Predicted age (weeks)
Chronological age (weeks)

▲ CTL   (n=37)
▲ VCD   (n=110)

E Age prediction: *Foxl2* +/-

Predicted age (weeks)
Chronological age (weeks)

◆ *Foxl2*+/+ (n=8)
◆ *Foxl2*+/- (n=54)

F VCD - 30d post-injection

Treatment: p ~3.10e-11
Age: p ~8.79e-9

Age acceleration
(Predicted – Chronological)

CTL VCD CTL VCD CTL VCD CTL VCD
3m   6m   8m   10m
Age-at-injection

G VCD - 90d post-injection

Treatment: p ~1.56e-11
Age: p ~6.00e-8

CTL VCD CTL VCD CTL VCD CTL VCD
3m   6m   8m   10m
Age-at-injection

H VCD - 150d post-injection

Treatment: p ~2.33e-10
Age: p ~1.22e-3

CTL VCD CTL VCD CTL VCD CTL VCD
3m   6m   8m   10m
Age-at-injection

I *Foxl2* +/-

Genotype: p ~4.01e-1
Age: p ~2.75e-14

Age acceleration
(Predicted – Chronological)

Foxl2+/+ *Foxl2*+/- *Foxl2*+/+ *Foxl2*+/- *Foxl2*+/+ *Foxl2*+/-
3-4m   8-10m   12-13m

▲◆ Model prediction
△◇ Out-of-bag prediction

Fig. 5

Fig. 6

A Cell type proportion analysis scheme

Non-immune (*Ptprc⁻*) vs. Immune (*Ptprc⁺*) → Flow cytometry: CD45 expression

Cell subtypes → RNAscope: Marker gene expression

scRNA-seq datasets   Cell type proportion analysis via scProportion   Validation

Cell type proportions from scRNA-seq: *Ptprc⁻* vs. *Ptprc⁺*

B Aging

C VCD

D *Foxl2* +/−

Cell type proportions from flow cytometry: CD45⁺ cells

E Aging

F VCD

G *Foxl2* +/−

Cell type proportions from scRNA-seq: *Ptprc⁻* cell subtypes detected

H Aging

I VCD

J *Foxl2* +/−

Cell type proportions from RNAscope: Granulosa and theca cells

K Aging

L VCD

M *Foxl2* +/−

N Aging

O VCD

P *Foxl2* +/−

Fig. 7

A Pseudobulk data analysis pipeline scheme

scRNA-seq datasets → Subset by cell type → Pseudobulk data via muscat → DEG analysis via DESeq2 / GO analysis via clusterProfiler

B DEGs: Aging (FDR < 0.05)

C DEGs: VCD (FDR < 0.05)

D DEGs: *Foxl2 +/-* (FDR < 0.05)

E Frequent GSEA term enrichment: GO Biological Process (FDR < 0.1)

F Frequent GSEA term enrichment: Reactome (FDR < 0.1)

Fig. 8

A Model-specific age-gradient differential expression analysis scheme (DESeq2 LRT)

B Boxplots of ovarian age-regulated genes (FDR < 1e-5):
VCD model, Granulosa cells

Top functional enrichment of ovarian age-regulated genes (GO BP)

C

D Boxplots of ovarian age-regulated genes (FDR < 1e-5):
Foxl2 haploinsufficiency model, Granulosa cells

Top functional enrichment of ovarian age-regulated genes (GO BP)

E

Fig. 9

A Transcriptome-based clock training and analysis scheme - Granulosa cells

Gene 1 Gene 2 ... Gene X

Cell 1
Cell 2
...
Cell Y

23480 features
x 7768 cells

Predicted age

Chronological age

Train & Test
Lasso regression

Expression analysis by age

Over-representation analysis

scRNA-seq transcriptome data:
Granulosa cells

Lasso feature analysis

B Granulosa clock performance: Test set

Rho ~0.82
p ~1.3e-585

Predicted age (weeks)

Chronological age (weeks)

● Aging
▲ VCD: CTL
◆ Foxl2 +/-: Foxl2⁺/⁺

C Age prediction: VCD

Predicted age (weeks)

Chronological age (weeks, + offset)

▲ CTL
▲ VCD

D Age prediction: Foxl2 +/-

Predicted age (weeks)

Chronological age (weeks, + offset)

◆ Foxl2⁺/⁺
◆ Foxl2⁺/⁻

E Age acceleration: VCD

Treatment: p ~6.02e-121
Age   : p ~6.20e-16

Treatment: p ~6.38e-44
Age: p ~5.67e-8

Age acceleration
(Predicted - Chronological)

CTL VCD CTL VCD
3m   10m
30d

CTL VCD CTL VCD
3m   10m
90d

Groups:

F Age acceleration: Foxl2 +/-

Genotype: p ~2.35e-6
Age: p ~7.00e-29

Age acceleration
(Predicted - Chronological)

Foxl2⁺/⁺ Foxl2⁺/⁻   Foxl2⁺/⁺ Foxl2⁺/⁻   Foxl2⁺/⁺ Foxl2⁺/⁻
Young   Mid-age   Old

G Expression of lasso features by age (n=1063 genes)

Row z score

2
1
0
-1
-2

13 15 17 20 29 39 48 57 73 76 86 87
Chronological age (weeks)

H ORA enrichment of top 500 features: GO ALL

Extracellular matrix
External encapsulating structure
Collagen−containing extracellular matrix
Signaling receptor regulator activity
Negative regulation of locomotion
Positive regulation of response to external stimulus
Receptor ligand activity
Signaling receptor activator activity
Leukocyte chemotaxis
Negative regulation of cell motility
Negative regulation of cellular component movement
Maintenance of location
Cell chemotaxis
Maintenance of location in cell
Artery development

Count
● 10
● 20

FDR
0.10
0.05
0.00