Systematic characterization of the ovarian landscape across mouse menopause models

Minhoo Kim[1], Rajyk Bhala[1], Justin Wang[1,2,3], Ryan J. Lu[1], Evelyn H. Lee[1], Julio L. Alvarenga[1], Rapheal G. Williams[1], and Bérénice A. Benayoun[1,4,5,6,7*]

1. Leonard Davis School of Gerontology, University of Southern California, Los Angeles, CA 90089, USA.

2. Department of Quantitative and Computational Biology, University of Southern California, Los Angeles, CA 90089, USA.

3. Department of Chemistry, University of Southern California, Los Angeles, CA 90089, USA.

4. Molecular and Computational Biology Department, USC Dornsife College of Letters, Arts and Sciences, Los Angeles, CA 90089, USA.

5. Cancer Biology Department, USC Keck School of Medicine, Los Angeles, CA 90033, USA.

6. USC Norris Comprehensive Cancer Center, Epigenetics and Gene Regulation, Los Angeles, CA 90033, USA.

7. Pharmacology & Pharmaceutical Sciences Department, USC Alfred E. Mann School of Pharmacy, Los Angeles, CA 90033, USA.

*Correspondence: berenice.benayoun@usc.edu

Abstract

Menopause not only affects fertility but also has widespread impact on systemic health. Yet, the molecular mechanisms underlying this process are not fully understood, partly due to the absence of robust, age-relevant preclinical models with comprehensive molecular and phenotypic characterization. To address this, we systematically compared three candidate mouse models of menopause: (1) intact aging, (2) chemical ovarian follicle depletion using 4-vinylcyclohexene diepoxide (VCD) administered at multiple ages, and (3) *Foxl2* haploinsufficiency, a genetic model based on a transcription factor linked to human premature ovarian failure. Through histology, serum hormone profiling, single-cell transcriptomics and machine-learning approaches, we uncovered both shared and model-specific features of follicle loss, endocrine disruption, and transcriptional remodeling. The VCD and *Foxl2* haploinsufficiency models revealed distinct patterns of hormonal and immune alterations not captured by intact aging alone. This comparative framework enables informed selection of context-appropriate preclinical rodent models to study menopause and the broader physiological consequences of ovarian aging.

Keywords

Menopause, ovarian aging, preclinical models, single-cell RNAseq, ovarian age clock

Introduction

Women are born with a finite ovarian reserve, and its progressive depletion with age leads to the irreversible cessation of reproductive function, known as menopause[1-3]. Epidemiological studies have shown that later age-at-menopause is associated with increased lifespan and reduced risk of age-related diseases, including cardiovascular disease, osteoporosis, and neurodegeneration[4-6]. Conversely, the onset of menopause is accompanied by a sharp rise in the incidence of age-associated pathologies, underscoring the widespread physiological consequences of ovarian aging[7]. Despite its clinical relevance, the molecular mechanisms linking ovarian failure to systemic aging remain poorly understood.

Menopause marks the end of female fertility and is characterized by significant shifts in the hormonal milieu, which exert systemic effects[8]. As the ovarian follicular pool becomes depleted, declining steroidogenic activity disrupts feedback regulation of the hypothalamic–pituitary–gonadal (HPG) axis, leading to pronounced changes in reproductive hormones[8]. For example, circulating levels of estrogen and anti-Müllerian hormone (AMH) decline to near-undetectable levels, while follicle-stimulating hormone (FSH) and Inhibin A (INHBA) exhibit marked elevations[9-11]. This hormonal reprogramming impacts not only reproductive tissues but also multiple extragonadal systems, including the metabolic, skeletal, and central nervous systems[12,13]. Accurately modeling this complex endocrine transition in preclinical systems is essential for delineating the mechanisms through which ovarian dysfunction impacts systemic aging.

Currently available rodent models for menopause research each possess distinct limitations. Intact aging female mice remain the most widely used model due to their physiological relevance and ease of implementation. Moreover, decades of research across diverse systems have established a robust foundational understanding of age-related biological changes, providing a valuable reference point for comparative analyses. Yet, intact aging mice do not undergo true menopause; instead, they enter estropause, retaining low but detectable estrogen levels and lacking the postmenopausal hormonal milieu observed in humans[14,15]. Another popular preclinical model is the use of surgical removal of ovaries, known as ovariectomy (OVX). While OVX achieves complete estrogen depletion, the surgery bypasses the gradual endocrine transition associated with menopause, and eliminates post-reproductive ovarian tissue, which retains androgenic activity and contributes to aging phenotypes[16]. Thus, despite its popularity, OVX is a poor preclinical model of menopause.

An alternative approach involves chemical ovarian follicle depletion using 4-vinylcyclohexene

diepoxide (VCD), which selectively targets small pre-antral follicles (primordial and primary follicles), and

induces follicle atresia, and leads to progressive ovarian failure[14]. This strategy allows for temporal control

of ovarian depletion and results in estrus acyclicity, estrogen deficiency, and compensatory FSH elevation,

features that closely mirror human menopause[17,18]. However, prior applications of the VCD model have

been limited to young mice (~2–3 months old), restricting its relevance for studying menopause in the

context of organismal aging[19,20]. Given that age alters the metabolic, immune, and endocrine

landscapes[21-23], it is critical to evaluate how the timing of ovarian failure intersects with aging processes.

For example, heterochronic parabiosis experiments have demonstrated that the aging systemic milieu

can impair tissue function in young animals, with circulating factors from aged animals negatively

affecting neurogenesis and cognitive function[24]. Thus, evaluating VCD exposure in older animals may

yield important insights into how age-related changes in the systemic environment shape the

physiological consequences of menopause.

The genetics of premature menopause in humans may also help expand the toolkit to study

menopause in preclinical models[25,26]. Specifically, haploinsufficiency of *Foxl2*, a transcription factor

essential for granulosa cell identity and ovarian function[25-28], is a great candidate model for menopause in

mice. Indeed, heterozygous *FOXL2* mutations are causally linked to premature ovarian insufficiency in

humans[29-31]. Although most research efforts have focused on full *Foxl2* knockout mice[25,26], which show

high perinatal lethality and primary ovarian defects, previous studies did report anecdotal subfertility

phenotypes of heterozygous carriers. Despite these promising reports, the physiological consequences of

Foxl2 haploinsufficiency on ovarian health and function have not been characterized in mice *in vivo*.

Unlike surgical or chemically induced models, *Foxl2* haploinsufficiency enables the study of endogenous,

genetically encoded ovarian dysfunction and its downstream systemic effects. This model offers a unique

opportunity to interrogate gradual follicular depletion and endocrine dysregulation in the absence of

exogenous perturbation (surgical or chemical insult).

In this study, we characterized and systematically compared three complementary mouse models

that reflect distinct mechanisms of ovarian decline as candidate models of menopause: (1) the intact

aging model, (2) a VCD-induced follicle depletion initiated at different ages spanning young adulthood to

early middle-age, and (3) a genetic model of *Foxl2* haploinsufficiency. Using these models, we sought to

identify shared and model-specific features of ovarian functional decline across multiple biological scales

and phenotypic layers. We integrated histological analyses, serum hormone profiling, and single-cell

RNA-sequencing (scRNA-seq) to assess changes in ovarian structure, endocrine function, and

transcriptional landscape across menopause models. Additionally, we developed hormone-based and

transcriptome-based ovarian aging clocks to evaluate functional trajectories of reproductive aging.

Together, our data provides a comprehensive framework for systematically evaluating and comparing

physiologically relevant models of ovarian aging, thereby defining their distinct advantages and

applications for studying systemic and molecular effects of menopause.

Results

Selection of mouse models of menopause for the study

To systematically assess and compare physiologically relevant menopause models, we

characterized three rodent models that represent distinct mechanisms of ovarian dysfunction: (1) the

intact aging model ("Aging model"), (2) a chemically-induced model using repeated intraperitoneal

injections of 4-vinylcyclohexene diepoxide (VCD, "VCD model")[14,32], and (3) a genetic model of *Foxl2*

haploinsufficiency ("*Foxl2* +/- model")[27]. Our aim was to establish a comparative framework to evaluate

these models in terms of histological, hormonal, and integrated ovarian health outcomes to evaluate their

suitability for modeling key aspects of ovarian aging (**Fig. 1a**). The Aging model involved comparison of

young adult (4-month-old) and old (20-month-old) female C57BL/6JNia mice. We selected these

timepoints to capture the transition from adult reproductive with stable estrus cycling (4 months) to post-

estropausal age (20 months), as C57BL/6 females typically remain cyclic until ~12 months and show

pronounced ovarian aging phenotypes by 20 months[33]. The VCD model included C57BL/6J female mice

injected with vehicle control (safflower oil, hereafter referred to as "CTL") or VCD starting at 3, 6, 8, or 10

months of age. This scheme allowed us to examine how the timing of ovarian insult (potentially modeling

varying "age-at-menopause") can impact ovarian health outcomes, particularly when VCD exposure

occurs closer to natural estropause transition in mice (~12 months)[33]. The *Foxl2* haploinsufficiency model

compared *Foxl2*[+/+] and *Foxl2*[+/-] female littermate mice from our transgenic colony at young (3–4 months)

and early middle-age (8–10 months), prior to the onset of natural estropause, to test whether partial loss

of *Foxl2* can recapitulate age-associated ovarian changes at earlier ages than control mice.

For the *Foxl2* haploinsufficiency model, we generated *Foxl2* haploinsufficient mice using a

regenerated floxed targeting allele, as previously described [27], which introduces a null mutation in the

Foxl2 gene via deletion of the gene single exon, using services from Cyagen Biomodels LLC (**Extended**

Data Fig. 1a; see **Methods**). Unlike prior studies that employed this allele for temporally controlled,

conditional deletion in adulthood, our study used this strategy to achieve constitutive *Foxl2*

haploinsufficiency throughout life (see **Methods**). To confirm that reduced *Foxl2* expression was indeed

observed in ovarian tissue from *Foxl2*[+/-] animals, as required to observe phenotypic consequences of

haploinsufficiency, we performed RT-qPCR using ovaries from young and early middle-age matched

Foxl2^{+/+} and *Foxl2^{+/-}* female mice from our colony (**Extended Data Fig. 1b**). While the young group

showed only a non-significant trend towards decreased expression, the early middle-age *Foxl2^{+/-}* ovaries

showed a significant reduction in expression (p-value ~0.032; **Extended Data Fig. 1b**). To confirm

whether *Foxl2* haploinsufficiency had an impact on ovarian function in these mice, we examined fertility

outcomes for *Foxl2^{+/+}* and *Foxl2^{+/-}* females from our breeding colony (see **Methods**; **Extended Data Fig.**

1c,d; **Extended Data Table 1**). Interestingly, *Foxl2^{+/-}* females showed a trend towards a smaller first litter

size (p-value ~0.075; **Extended Data Fig. 1c**), as well as a significantly increased latency to first

pregnancy compared to matched *Foxl2^{+/+}* females (p-value ~0.035; **Extended Data Fig. 1d**), suggesting

impaired reproductive potential consistent with ovarian dysfunction or premature ovarian aging.

Histological and hormonal characterization of mouse menopause models

A decline in ovarian follicle numbers is a well-established hallmark of ovarian aging and

menopause[34]. Thus, we performed histological analysis of ovarian follicle populations using hematoxylin

and eosin (H&E) staining (**Fig. 1b-g**; **Extended Data Fig. 2**). As expected, the Aging model showed a

significant reduction in follicles at all developmental stages in old females, including primordial, primary,

secondary, and antral follicles, as well as corpora lutea (**Fig. 1b,e**; **Extended Data Fig. 2a**), consistent

with the established progressive follicular depletion that accompanies natural ovarian aging[35,36]. At five

months post-injections, VCD-treated animals also showed significant depletion of follicle numbers,

regardless of age-at-injection (**Fig. 1c,f**; **Extended Data Fig. 2b**). Due to the terminal nature of

histological analysis, ovaries were collected for H&E staining at the single 5 months post-injection time

point. This endpoint was selected to align with the conclusion of our longitudinal serum profiling, wherein

monthly hormone measurements enabled tracking of endocrine changes over time (see below). In

contrast, *Foxl2^{+/-}* mice did not show any reduction in combined follicle counts; instead, we observed a

trend toward increased follicle counts in the middle-aged *Foxl2^{+/-}* animals (p-value ~0.08; **Fig. 1d,g**;

Extended Data Fig. 2c).

To further assess ovarian function, we measured serum levels of AMH, FSH and INHBA, key

markers of ovarian reserve and hypothalamic–pituitary–gonadal (HPG) axis regulation[37,38] (**Fig. 1h-p**;

Extended Data Table 2). In the Aging model, old females had decreased AMH and INHBA and increased

FSH compared to young females, consistent with diminished ovarian reserve[38,39] (**Fig. 1h,k,n**). A similar

pattern was observed in the VCD model, whereby AMH and INHBA were decreased and FSH was

increased compared to the vehicle control groups at 5 months post-injection timepoint (**Fig. 1i,l,o**). To

enable a comprehensive analysis of endocrine responses across reproductive life stages, we also

performed longitudinal hormone profiling over a five-month period following VCD exposure at various

ages (3, 6, 8 and 10 months; **Extended Data Fig. 3a-c**). Interestingly, VCD injections at older ages

resulted in attenuated hormonal shifts compared to younger counterparts, suggesting that ovarian or

systemic factors at midlife may modulate sensitivity to VCD-induced insult to ovarian reserve. In the

Foxl2[+/-] mice, we observed distinct hormonal trends: AMH levels were increased, FSH levels were mildly

elevated, and INHBA levels were reduced compared to wild-type controls, although none of these

changes reached statistical significance (**Fig. 1j,m,p**). The increased AMH levels are consistent with our

histological findings of increased follicle counts in the *Foxl2*[+/-] animals (**Fig. 1d,g**). Importantly, *FOXL2* has

been shown to interact with *AMH* to modulate follicle recruitment in humans[40]. Thus, the increased follicle

counts and AMH levels in the *Foxl2*[+/-] animals likely reflect altered gene regulatory network driven by

Foxl2 deficiency, rather than true ovarian rejuvenation (see **Discussion**).

Next, we applied our previously described composite ovarian health index, which integrates

information from follicle counts and serum hormone levels to yield a holistic measure of ovarian health[41]

(**Fig. 1q-s**). As expected, this index was significantly lower in old *vs.* young animals and VCD-treated

animals vs. CTL (regardless of age-at-injection), reflecting impaired ovarian health (**Fig. 1q,r**). In contrast,

Foxl2[+/-] mice did not differ significantly from wild-type animals at either age group, consistent with their

preserved histological and hormonal profiles (**Fig. 1s**).

Together, these results provide a comprehensive characterization of three distinct candidate

mouse models of menopause. Both the Aging and VCD models exhibit robust phenotypic hallmarks of

ovarian decline, including depletion of follicular reserves and disrupted endocrine profiles, with the VCD

model offering unique advantages for temporally controlled induction of ovarian dysfunction. In contrast,

the *Foxl2* haploinsufficiency model presents an unexpected phenotype characterized by preserved or

even elevated follicle counts and increased AMH levels, opposite to typical features of ovarian aging[9-11],

despite clear evidence of reproductive dysfunction (**Extended Data Fig. 1c,d**). These findings raise the

possibility that *Foxl2* haploinsufficiency alters ovarian physiology through mechanisms distinct from

conventional aging, warranting deeper molecular investigation.

A hormone-based clock, "OvAge", for non-terminal prediction of ovarian aging

Although the ovarian health index is a useful metric of ovarian health, its partial reliance on post-

mortem histological measurements limits its applicability in longitudinal studies. Thus, we aimed at

developing a complementary non-terminal metric of ovarian aging, using a hormone-based predictive

machine-learning model, which we termed "OvAge", capable of estimating ovarian age from circulating

serum hormone levels (**Fig. 2a**). By incorporating AMH, FSH, and INHBA levels, we sought to develop a

model that provides a more comprehensive and integrative assessment of endocrine function than

evaluating each hormone individually. To increase the generalizability of the clock, we trained a random

forest regression model using serum hormone data (AMH, FSH, and INHBA) from animals with intact

ovarian function and no experimental perturbation, including animals from the Aging model, vehicle

control (CTL) groups from the VCD model, *Foxl2*[+/+] animals from the *Foxl2* haploinsufficiency model, and

data from *Fshr*[+/+] animals from our previously published *Fshr* haploinsufficiency study[42] (**Fig. 2a**). This

hormone level data was then randomly partitioned into training (n = 190) and testing (n = 62) sets to

enable robust evaluation of model performance (**Fig. 2a**).

Model performance evaluation showed strong agreement between predicted and chronological

age in the withheld test dataset (Spearman Rho ~0.621; p-value ~7.4 × 10[-8]), confirming the robustness

of our hormone-based OvAge clock (**Fig. 2b**). We then applied OvAge to assess ovarian age of VCD-

injected and *Foxl2*[+/-] animals (**Fig. 2c,d**). In the VCD model, predicted ovarian age was consistently

higher than chronological age across all age-at-injection groups, suggesting that VCD exposure

accelerates ovarian aging (**Fig. 2c**). Notably, the gap between predicted and chronological age narrowed

as the age-at-injection increased, a pattern consistent with both the hormone profiles and ovarian health

index (**Fig. 1l,o,r**). In contrast, no consistent pattern was observed in the *Foxl2* haploinsufficiency model;

predicted ages of *Foxl2*[+/-] animals varied and did not seem to differ substantially from their wild-type

counterparts (**Fig. 2d**). To quantify the degree of divergence between predicted and chronological age, we

computed hormonal age acceleration, defined as the difference between OvAge-predicted and true

chronological age (**Fig. 2e-h**). For the VCD model, we analyzed animals at 30-, 90-, and 150-days post-

injection (**Fig. 2e-g**). We observed significant increases in age acceleration in VCD-treated animals

compared to controls at both 30 and 90-days post-injection across all age-at-injection groups (**Fig. 2e,f**).

Interestingly, by 150 days post-injection, animals injected at 10 months of age (now aged ~15 months) no

longer showed significant age acceleration (**Fig. 2g**). This may reflect convergence in ovarian aging

trajectories in both vehicle control and VCD-treated animals as they approach or enter the estropausal

interval (12-15 months in the C57BL/6 strain[33]). In contrast, animals from the *Foxl2* haploinsufficiency

model did not show a consistent increase in OvAge or age acceleration (**Fig. 2h**). Instead, middle-age

Foxl2[+/-] animals exhibited a statistically significant reduction in age acceleration (p-value ~0.011),

consistent with other observed phenotypes in this model, including preserved follicle counts and higher

AMH levels (**Fig. 1d,g,j**). These findings may reflect a unique regulatory trajectory in this model, which

could offer a complementary perspective on ovarian aging as well as premature ovarian failure.

Together, our results demonstrate that OvAge effectively captures both accelerated and

attenuated ovarian aging trajectories in response to different biological perturbations. The clock

recapitulates expected aging patterns in intact naturally aged animals, reveals premature ovarian aging

following VCD exposure, and highlights a distinct endocrine aging profile in the *Foxl2* haploinsufficiency

model. These findings underscore the utility of OvAge as a non-terminal framework for assessing ovarian

aging and provide insight into model-specific differences that may be leveraged to explore various facets

of reproductive senescence.

Although the *Foxl2* haploinsufficiency model did not exhibit overt signs of ovarian dysfunction

based on the ovarian health index or OvAge (**Fig. 1s and 2d,h**), we reasoned that its preserved

endocrine profile does not preclude underlying molecular perturbations consistent with accelerated

ovarian aging. Given the established role of *FOXL2* in granulosa cell identity and ovarian maintenance, its

association with premature menopause in humans[29,30,43], and the fertility phenotypes that we observed

(**Extended Data Fig. 1c,d**), *Foxl2* haploinsufficiency may still disrupt transcriptional programs in the ovary

before functional decline becomes apparent.

Thus, we next examined the transcriptional landscape of ovaries from each model using single-

cell RNA-sequencing (scRNA-seq), enabling deeper investigation of molecular aging dynamics across

natural, chemical, and genetic models of menopause.

Single-cell transcriptomic profiling of ovarian cell types across menopause models

To characterize the ovarian transcriptional landscapes across candidate mouse menopause

models, we performed scRNA-seq on ovaries collected from the Aging, VCD, and *Foxl2*

haploinsufficiency models (**Fig. 3a**). For the Aging model, samples were collected from young (4-month-

old) and old (20-month-old) animals to capture transcriptional changes associated with intact aging (**Fig.**

3b). For the VCD model, we focused on two age-at-injection groups, 3 and 10 months, to capture the

extremes of VCD responsiveness observed in prior analyses, as well as two post-injection time points, 30

and 90 days, based on differential effects on age acceleration at these stages (**Fig. 3b**). For the *Foxl2*

haploinsufficiency model, ovaries were collected from *Foxl2$^{+/+}$* and *Foxl2$^{+/-}$* animals at young adult (~4

months) and early middle-age (~9 months), enabling the assessment of gene expression changes driven

by partial loss of *Foxl2* function across adulthood (**Fig. 3b**). The resulting datasets yielded 9,559, 47,386,

and 21,301 high-quality single cells from the Aging, VCD, and *Foxl2* haploinsufficiency models,

respectively (**Fig. 3a-e**; **Extended Data Table 3**). Ovarian cell populations were defined through

dimensionality reduction, unsupervised clustering, and annotation using both established marker genes

and automated reference mapping to publicly available single-cell datasets (**Fig. 3c-h**; **Extended Data**

Table 4; see **Methods**).

After annotation of our scRNA-seq datasets, we were able to identify all major ovarian cell types

(**Fig. 3c-h**). Among the abundant non-immune populations, granulosa cells (*Amh, Fshr, Cyp19a1*), theca

cells (*Cyp17a1, Cyp11a1*), stromal cells (*Pdgfra, Col1a1*), smooth muscle cells (SMCs; *Acta2*), pericytes

(*Notch3*, *Rgs5*), blood endothelial cells (BECs; *Flt1*), lymphatic endothelial cells (LECs; *Prox1*), and

epithelial cells (*Upk1b*) were identified. The immune compartment included neutrophils (*S100a8*),

monocytes (*Adgre1*), macrophages (*C1qa*), dendritic cells (*Flt3*), innate lymphoid cells (ILCs; *Il2ra*),

natural killer (NK) cells (*Klrb1c, Gzma, Nkg7*), NKT cells (*Cd3e, Klrb1c, Gzma, Nkg7*), CD8^{+} NKT cells

(*Cd3e, Klrb1c, Gzma, Nkg7, Cd8b1*), CD8^{+} T cells (*Cd3e , Cd8b1*), CD4^{+} T cells (*Cd3e, Cd4*), double-

negative T (DNT) cells (*Cd3e*), double-positive T (DPT) cells (*Cd3e, Cd8b1, Cd4*), and B cells (*Cd19*) (**Fig.**

3f-h).

Given the known ovotoxic effects of VCD, we performed additional histological analyses on

condition-matched ovaries (matched for age-at-injection and time post-injection) to confirm the presence

of follicular structures in animals processed for scRNA-seq (**Extended Data Fig. 4a**). H&E staining of

condition-matched samples revealed that despite extensive follicle depletion, residual follicles were

present in most VCD-treated animals (**Extended Data Fig. 4a**). These findings were consistent with the

detection of follicle-associated cell types in our scRNA-seq data (**Fig. 3d,g**). Quantification of total follicles

by blinded observers further supported the persistence of follicular structures in the VCD model animals

(**Extended Data Fig. 4b**). We also performed serum hormone quantification (AMH, FSH and INHBA) on

age- and condition-matched samples corresponding to those used for scRNA-seq (**Extended Data Fig.**

5a-c). Consistent with previous findings, AMH and INHBA levels were reduced in the VCD groups

(**Extended Data Fig. 5a,c**). To note, FSH levels were measured using an updated assay kit implemented

by the University of Virginia Ligand core facility, distinct from the kit used in prior analyses (see **Methods**).

Despite this change, we observed a trend toward increased FSH levels in the VCD-treated group relative

to matched controls, although statistical significance was not reached, likely due to limited sample size

(**Extended Data Fig. 5b**). Then, we evaluated the ovarian health index in these animals (**Extended Data**

Fig. 5e). Because FSH was measured using a different assay than the one used to establish the ovarian

health index, we implemented a previously reported correction procedure[41] to ensure consistency and

comparability of FSH values across cohorts (**Extended Data Fig. 5d**; see **Methods**). We observed a

trend for lower ovarian health index scores in the VCD group compared to matched controls, although it

did not reach statistical significance, again likely due to small sample size (**Extended Data Fig. 4e**).

To validate cell type assignments, we performed RNAscope, a high sensitivity *in situ* hybridization

assay for both well-characterized and less frequently profiled ovarian cell populations (**Fig. 3i**; **Extended**

Data Fig. 6). Granulosa cells (*Fshr*), theca cells (*Cyp11a1*), stromal cells (*Pdgfra*) and SMCs (*Acta2*)

were validated using established marker probes, confirming their presence in the dataset (**Extended**

Data Fig. 6a). Epithelial cells (*Upk1b*), BECs (*Flt1*) and LECs (*Prox1*) were similarly detected, supporting

the robustness of annotations (**Extended Data Fig. 6b**). Importantly, we also validated the presence of

various immune cell subsets, including NK (*Klrb1c*), NKT (*Klrb1c*, *Cd3e*), CD8$^+$ T (*Cd3e*, *Cd8b1*), CD4$^+$ T

cells (*Cd3e, Cd4*), DNT (*Cd3e,* with no Cd8b1/ Cd4 signal) and B (*Cd19*) cells (**Extended Data Fig. 6c**).

We also confirmed presence of less commonly reported ovarian cell types, including CD8+ NKT (*Klrb1c,*

Cd3e, Cd8b1) and DPT (*Cd3e, Cd8b1, Cd4*) cells (**Fig. 3i**). These results further support the accuracy of

the single-cell transcriptomic annotations and highlight the diverse somatic and immune

microenvironments present across the aging models.

To assess cell type consistency across models, we compared annotated cell populations across

datasets. A total of 19 cell types, 8 non-immune and 11 immune types, were shared across all three

models (**Extended Data Fig. 6d**). Some cell types were not shared across models, potentially due to

technical factors (e.g., capture efficiency and sequencing depth), or underlying biological variation

inherent to each mouse model. To assist researchers in navigating the data efficiently, we developed an

interactive R shiny app that makes the annotated datasets directly accessible and explorable through a

graphical interface: https://minhooki.shinyapps.io/shinyappmulti/.

scRNA-seq analysis reveals shifts in ovarian cell composition in menopause models

Next, we examined changes in cell type proportions across our datasets (**Fig. 4a**; **Extended Data**

Table 5). We first assessed whether the proportion of immune cells shifted with ovarian aging by

analyzing the expression of *Ptprc*, which encodes Cd45, a pan-immune marker (**Fig. 4b-d**). In the Aging

model, we observed a significant increase in *Ptprc*$^+$ cells, indicating an expansion of the immune cell

population with age (**Fig. 4b**). This is consistent with previous scRNA-seq studies in intactly aged mouse

ovaries, which reported an increase in immune cell abundance at even younger ages (3 vs. 9 months; 4.5

vs. 10.5 vs. 15.5 months)[44,45]. In contrast, single-nucleus RNA-sequencing data from human ovaries did

not identify a similar shift in immune cell proportions with age, consistent with potential species-specific

differences in immune remodeling for the aging ovary[46]. However, this discrepancy could also reflect

differences in age ranges examined or technical factors such as dissociation and capture methods

inherent to single-nucleus versus single-cell approaches.

VCD-treated mice exhibited a consistent reduction in *Ptprc*$^+$ cells across all conditions,

suggesting a depletion of immune cells following exposure to VCD (**Fig. 4c**). This reduction may result

from the loss of follicular structures, alterations in immune recruitment, or uncharacterized systemic

effects of VCD on immune homeostasis. Additionally, broader tissue remodeling processes, including

fibrosis, may contribute to the altered immune cell landscape observed in VCD-treated ovaries. Thus, the

VCD model does not seem to recapitulate immune features of intact aging ovaries in mice, regardless of

age-at-injection. In the *Foxl2* haploinsufficiency model, we detected a significant increase in *Ptprc*$^+$ cells in

the *Foxl2*$^{+/-}$ animals, regardless of age (**Fig. 4d**). These findings were further validated by flow cytometric

analysis of CD45$^+$ cells, using ovarian single-cell preparations generated with the same dissociation

protocol as the scRNA-seq experiments (**Fig. 4e-g**; **Extended Data Fig. 7a**). Intriguingly, this suggests

that the *Foxl2* haploinsufficiency model may better recapitulate immune shifts in the ovary seen with intact

aging compared to the VCD model.

Then, we conducted a comprehensive analysis of all ovarian cell populations detected in the

datasets to identify shifts in cellular composition in response to intact aging, VCD exposure, and *Foxl2*

haploinsufficiency (**Fig. 4h-j**; **Extended Data Fig. 7b-d**). In the Aging model, nearly all non-immune cell

populations showed a decline in proportion with age, except for stromal cells and LECs, which remained

unchanged (**Fig. 4h**). In contrast, several immune cell populations increased with age, including

neutrophils, DCs, ILCs, CD8$^+$ NKT, CD4$^+$ T, DNT, and DPT cells (**Extended Data Fig. 7b**). Notably, DNT

cells have been reported to increase in aging mouse ovaries, and a potential role for these cells in

regulating ovarian aging has been proposed[44,47]. Conversely, monocytes, NK, CD8$^+$ T, and B cells

exhibited a significant decrease with age (**Extended Data Fig. 7b**).

In the VCD model, cell proportion shifts were largely consistent within the treatment group,

regardless of age-at-injection or time post-injection (**Fig. 4i**; **Extended Data Fig. 7c**). The VCD-treated

group showed an increasing trend in stromal cells, BECs, LECs, epithelial cells, and CD4$^+$ T cells. In

contrast, granulosa cells, theca cells, SMC, and pericytes were significantly reduced in proportion (**Fig. 4i**).

Among immune cells, neutrophils, monocytes, macrophages, DCs, NKT, DPT, and B cells were all

decreased following VCD treatment (**Extended Data Fig. 7c**).

In the *Foxl2* haploinsufficiency model, granulosa and epithelial cell proportions were consistently

reduced in *Foxl2*$^{+/-}$ animals, regardless of age (**Fig. 4j**). Conversely, monocytes, NK, CD8$^+$ NKT, and B

cells were increased in *Foxl2*$^{+/-}$ animals (**Extended Data Fig. 7d**). Interestingly, we observed opposing

trends in stromal cells, macrophages, and DNT cells. Stromal cell proportions increased in *Foxl2*[+/+] young

mice but decreased in *Foxl2*[+/+] animals at early middle-age, whereas macrophages and DNT cells

showed the opposite trend. These findings highlight model-specific patterns of non-immune and immune

cell remodeling, underscoring distinct trajectories shaped by chronological, chemical, and genetic

perturbations of ovarian function.

To validate the cell abundance findings from our scRNA-seq data, we leveraged our RNAscope *in*

situ hybridization data for representative non-immune and immune populations (see above; **Fig. 4k-p**;

Extended Data Fig. 8,9). We examined granulosa cells, theca cells, stromal cells, SMCs, BECs, LECs,

and epithelial cells, along with NK, NKT, CD8[+] NKT, CD8[+] T, CD4[+] T, DNT, DPT, and B cells. In the Aging

model, RNAscope confirmed a decrease in granulosa and theca cell abundances with intact aging,

consistent with our scRNA-seq results (**Fig. 4k,n**). To note, there was non-specific staining for granulosa

and theca markers (which are histologically absent in the old), which did not colocalize with DAPI,

consistent with known technical limitations of the assay (**Fig. 4k**, right panel). Stromal cell proportions

remained unchanged, again in agreement with the scRNA-seq data (**Extended Data Fig. 8a**). While the

scRNA-seq data suggested a decrease in SMCs with age, we did not detect a significant difference by

RNAscope, likely due to low detection sensitivity of the *Acta2* probe. While epithelial cells showed a

decreasing trend with age in the RNAscope dataset without reaching statistical significance, BECs

exhibited a significant decline, consistent with scRNA-seq observations. We also examined LECs and did

not observe any significant age-associated changes, which aligns with findings from the scRNA-seq data.

For immune cells, RNAscope detected decreased abundances of NK, NKT, and B cells, and increased

abundances of CD8[+] NKT, CD4[+] T, DNT, and DPT cells, closely recapitulating the scRNA-seq trends

(**Extended Data Fig. 9a**). However, we did not observe a significant change in CD8[+] T cells by

RNAscope, despite a decrease in the scRNA-seq dataset (**Extended Data Fig. 7b,9a**). This discrepancy

may stem from differences in detection sensitivity or spatial distribution of CD8[+] T cells that may have

limited probe accessibility *in situ*.

In the VCD model, we focused our RNAscope analysis on samples from 3-month age-at-injection

and 30 days post-injection, given the consistent trends observed across conditions (**Fig. 4l,o**; **Extended**

Data Fig. 8b,9b). Among non-immune cells, we observed a significant decrease in granulosa and

increase in stromal with VCD exposure, consistent with scRNA-seq data (**Fig. 4l,o**; **Extended Data Fig.**

8b). We also observed some non-specific staining that did not colocalize with DAPI in the VCD-treated

ovaries, where granulosa is histologically absent (**Fig. 4l**, right panel). Theca cells showed a decreasing

trend in proportion with VCD exposure, but did not reach statistical significance (**Fig. 4l,o**). For SMCs,

BEC, LECs and epithelial cells, RNAscope revealed no significant change in proportions with age, which

contrasts with the scRNA-seq findings (**Extended Data Fig. 8b**). These inconsistencies may be

attributable to a numbers of technical factors: (i) reduced probe efficacy, (ii) limited marker expression or

(iii) under sampling of rare cell types in unique tissue slices compared to whole dissociated ovaries, all

potentially leading to under-detection of rare cell types by RNAscope. For immune populations,

RNAscope detected consistent decreases in NK, NKT, CD8$^+$ NKT, CD8$^+$ T and DNT cells, generally

aligning with findings from our scRNA-seq results (**Extended Data Fig. 7c,9b**). CD8$^+$ T and B cells did

not show any differences in proportion with VCD exposure (**Extended Data Fig. 9b**).

In the *Foxl2* haploinsufficiency model, RNAscope analysis revealed a reduction in granulosa cells

in *Foxl2$^{+/-}$* animals across both age groups, recapitulating the scRNA-seq findings (**Fig. 4j,m,p**). We also

detected a non-significant decrease in epithelial cell proportions in *Foxl2$^{+/-}$* animals in both age groups,

consistent with the trends observed in scRNA-seq data (**Fig. 4j**; **Extended Data Fig. 8c**). A significant

increase in SMCs and decrease in LECs in young and middle-age *Foxl2$^{+/-}$* animals were observed,

respectively, which aligns with the trends observed in our scRNA-seq datasets (**Extended Data Fig. 8c**).

Theca cells, stromal cells and BECs did not show significant changes in abundance, consistent with

scRNA-seq observations (**Extended Data Fig. 8c**). In the immune compartment, RNAscope identified

patterns that were generally consistent with our scRNA-seq results (**Extended Data Fig. 9c**). For

example, we observed increased proportions of NK and NKT cells in *Foxl2$^{+/-}$* animals. Additionally, we

observed increased DNT abundance in middle-age *Foxl2$^{+/-}$* animals, which agrees with our scRNA-seq

data (**Extended Data Fig. 9c**). As with other models, some cell types did not reach statistical significance

in RNAscope, often due to lower detection rates, but the overall trends were consistent. Together, these

findings further support model-specific shifts in immune and non-immune ovarian cell populations, as

captured by both scRNA-seq and RNAscope-based validation.

Global transcriptional perturbations of ovarian cell types across menopause models

To systematically evaluate how different models of menopause influence global transcriptional responses in the ovary, we applied Augur[48], a computational framework that quantifies cell type–specific separability across experimental conditions based on single-cell level gene expression profiles (**Fig. 5a**). Augur scores (derived from underlying machine-learning algorithm performance, reported as Area Under the Curve [AUC] values) represent the ability within each cell type to distinguish between biological groups within a given model based on overall transcriptional landscapes[48]. In the Aging model, granulosa cells exhibited the highest AUC score (AUC ~0.79), reinforcing their role as key transcriptional responders to chronological aging (**Fig. 5b,c**). Among other non-immune cell types, BECs and LECs also showed relatively high transcriptional divergence, suggesting that vascular and lymphatic compartments may undergo age-associated remodeling. Indeed, vascular decline has been closely linked to ovarian aging in multiple species, including humans, non-human primates, and mice[46,49,50]. In the immune compartment, CD4[+] T, NK, and B cells ranked highest (**Fig. 5b,c**). Interestingly, DNT cells, despite their expansion in the aging ovary, showed the lowest AUC score (AUC ~0.53), indicating limited transcriptional remodeling with age (**Fig. 5b,c , Extended Data Fig. 7b**).

In the VCD model, we observed marked transcriptional perturbations in theca, stromal, BEC, and LEC cells, especially at 90 days post-injection, with these effects largely consistent across both 3-month and 10-month age-at-injection groups (**Fig. 5d,e**). However, epithelial and granulosa cells displayed more pronounced age-at-injection-dependent differences: epithelial cells were more responsive in younger animals, while granulosa cells were notably absent in the 10-month cohort, likely reflecting more severe follicular depletion (**Fig. 5e**, left panel). Among immune cell types, NK, DNT, and CD8[+] NKT cells showed higher transcriptional sensitivity in younger animals (**Fig. 5e**, right panel). At 30 days post-injection, transcriptional divergence was generally more divergent in the VCD model compared to its 90-day counterpart (**Extended Data Fig. 10a,b**). For example, LECs in the 10-month group displayed strong divergence (AUC ~0.81), suggesting age-related vascular remodeling may emerge early after VCD exposure.

In the *Foxl2* haploinsufficiency model, transcriptional divergence was more pronounced at middle-age, suggesting that the effects of *Foxl2* loss may not fully manifest in the young ovary (**Fig. 5f,g**). Among

non-immune populations, granulosa and theca cells showed the highest AUC scores at mid-age, while

epithelial cells were more divergent in the young group (**Fig. 5g,** left panel). In immune populations,

neutrophils, NKT, CD8[+] NKT, and B cells exhibited the strongest divergence, with all showing elevated

AUC scores at middle-age, indicating potential immune activation or heightened sensitivity of specific

immune compartments to *Foxl2* insufficiency (**Fig. 5g**, right panel).

Together, these results reveal both common and model-specific overall transcriptional trajectories

across ovarian aging paradigms. Granulosa cells consistently emerge as sensitive indicators of ovarian

dysfunction across all three models. Immune cells show particularly pronounced divergence in *Foxl2*[+/-]

animals (**Fig. 5g**, right panel), suggesting that immune alterations may precede, accompany, or even

drive ovarian changes in this genetic model. These comparisons underscore the importance of

contextualizing cell type–specific transcriptional changes within each model's mechanistic framework and

highlight how distinct perturbations – such as chronological aging, chemical ablation, or genetic loss –

can produce divergent molecular responses across ovarian cell types.

Transcriptional and pathway-level signatures across menopause models

To investigate transcriptional remodeling across menopause models, we performed differential

expression and pathway enrichment analyses using data pseudobulked by cell type in each individual

scRNA-seq library (**Fig. 6a; Extended Data Table 6**). Six cell types, granulosa, theca, stromal, BEC,

epithelial, and DNT cells, were consistently detected across all three models and thus included in

downstream analyses (**Extended Data Fig. 11a**). DESeq2[51] was used to identify differentially expressed

genes (DEGs) for each model: young vs. old in the Aging model; CTL vs. VCD in the VCD model (using

age-at-injection and time post-injection as modeling covariates); and *Foxl2*[+/+] vs. *Foxl2*[+/-] in the *Foxl2*

haploinsufficiency model (using age as a modeling covariate). As expected, we confirmed reduced *Foxl2*

expression in *Foxl2*[+/-] animals at both ages based on our pseudobulked dataset (**Extended Data Fig.**

11b).

Using an FDR cutoff of 0.05, we observed substantial transcriptional changes that were both

model- and cell type-specific (**Fig. 6b-d**). Granulosa cells exhibited the highest number of DEGs in both

the Aging (n=1,194) and *Foxl2* haploinsufficiency (n=5,478) models, highlighting their sensitivity to

physiological and genetic perturbations. In the VCD model, theca cells showed the strongest

transcriptional response (n=5,385), followed by granulosa cells (n=3,664), suggesting robust

transcriptional remodeling in both steroidogenic cell types following VCD exposure (**Fig. 6b-d**).

We next performed gene set enrichment analysis (GSEA) to identify gene ontology (GO)

biological process terms recurrently enriched across models (**Fig. 6e; Extended Data Table 7**).

Mitochondrial-related gene sets were consistently downregulated across models and cell types, including

"Respiratory electron transport chain," "Oxidative phosphorylation" and "ATP synthesis coupled electron

transport" (**Fig. 6e**; FDR < 0.05). Immune-related processes, such as "Antigen processing and

presentation of endogenous peptide antigen," were also frequently identified as downregulated (**Fig. 6e**).

Using independent gene sets from Reactome[52] revealed consistent trends of mitochondria-related gene

sets being downregulated (**Fig. 6f**; FDR < 0.1), suggesting convergent dysregulation of metabolic function

during ovarian aging and in menopause models.

Together, these comparative analyses revealed both shared and model-specific transcriptional

features of ovarian aging. Granulosa cells consistently exhibited robust transcriptional changes across all

three models, underscoring their role as key responders to ovarian aging-related perturbations. Notably,

mitochondrial dysfunction and immune dysregulation emerged as common themes across models and

cell types, pointing to conserved biological pathways that may underlie the progression of ovarian aging.

Conserved and divergent aging-associated features are observed across menopause models

To identify features that may be broadly shared or uniquely diverge across menopause models,

we derived age-associated gene signatures from the Aging model, which reflects natural ovarian

physiological decline in mice (**Extended Data Fig. 11c**). These signatures were then used as a biological

reference to explore potential patterns of convergence and divergence across the VCD and *Foxl2*

haploinsufficiency models (**Extended Data Fig. 11c**). VCD dataset showed consistent upregulation of

aging-associated genes across cell types, whereas the *Foxl2* haploinsufficiency model exhibited partially

divergent expression patterns (**Extended Data Fig. 11c**). Notably, granulosa and stromal cells showed

decreased expression of genes typically upregulated with age, while BEC cells exhibited opposite trends

for both age-associated up- and down-regulated genes. These patterns in a subset of cell types may

reflect a distinct aging trajectory, or alternative molecular programs engaged in the setting of partial *Foxl2*

loss.

We also examined expression of the SenMayo[53] gene set, a curated gene set of senescence-

associated genes, which can be used as a proxy for senescence burden in transcriptional data

(**Extended Data Fig. 11d**). In the Aging model, all six cell types tested showed significant upregulation of

the SenMayo signature. In the VCD model, granulosa, stromal, BEC, and DNT cells mimicked this

enrichment, while theca and epithelial cells showed a negative correlation. In the *Foxl2* haploinsufficiency

model, theca and DNT cells showed consistent upregulation, whereas granulosa, stromal, BEC, and

epithelial cells showed an inverse pattern of downregulation (**Extended Data Fig. 11d**). These contrasting

results suggest that the VCD and *Foxl2* haploinsufficiency models may each capture distinct cell-specific

aspects of an accelerated ovarian aging process.

We next evaluated transcriptional noise by calculating the coefficient of variation (CV) for each

cell type (**Extended Data Fig. 11e**). Increased transcriptional variability is a characteristic of aging and

has been observed across multiple tissues, including the ovary[45] and hypothalamus[54] in mice, reflecting a

loss of transcriptional regulation with age. In the Aging model, CV increased across all detected cell types,

except in DNT cells. In the VCD model, theca cells consistently exhibited increased CV, while granulosa

and LECs showed more variable results. In the *Foxl2* haploinsufficiency model, theca cells again showed

increased CV regardless of age, although other cell types displayed inconsistent trends in young and

middle-aged animals. While increased CV is often linked to age-related loss of regulatory precision[54,55],

these mixed results in VCD and *Foxl2* haploinsufficiency models suggest a more nuanced transcriptional

landscape that may reflect compensatory or cell type-specific adaptations.

Lastly, we assessed transposable element (TE) expression using scTE[56] and DESeq2 (**Extended**

Data Fig. 11f). TE activity is known to increase with aging across tissues, driven in part by age-

associated chromatin remodeling, reduced heterochromatin integrity and impaired epigenetic

repression[57,58]. In the Aging model, granulosa, stromal, SMC, BEC, and LEC cells showed increased TE

expression with age, while theca, epithelial, NK, and B cells showed decreased TE expression. In the

VCD model, TE expression increased in stromal, BEC, LEC, and epithelial cells, and decreased in theca

cells. In the *Foxl2* haploinsufficiency model, TE expression increased in most cell types, except stromal

cells. Increase in TE expression is recognized as a hallmark of aging, contributing to genomic instability,

inflammation, and cellular stress[57,59]. Rather than reflecting uniform activation of TEs, these cell type–

specific shifts suggest that TE regulation may respond differently depending on the mechanism of ovarian

perturbation (**Extended Data Fig. 11f**).

Gene co-expression network and transcription factor activity across menopause models

To identify gene regulatory programs associated with ovarian aging and dysfunction, we

performed weighted gene co-expression network analysis (WGCNA)[60] and transcription factor (TF)

activity inference using decoupleR[61] (**Fig. 7a**). To obtain a comparative reference state, we first applied

WGCNA to pseudobulked gene expression data from the Aging model, reflecting naturally occurring

physiological changes. Modules with significant trait correlation to age (FDR < 0.1) were retained for

downstream analysis (**Extended Data Fig. 12,13**). GSEA of these gene modules revealed strong

enrichment in the Aging dataset (as expected by construction), providing a useful benchmark for

evaluating transcriptional convergence and divergence across intact aging and intervention models (**Fig.

7b**).

In the VCD model, most modules exhibited enrichment patterns consistent with the Aging model,

suggesting that VCD-induced ovarian failure recapitulates key transcriptional features observed during

normal chronological aging. In contrast, the *Foxl2* haploinsufficiency model revealed notable differences

in module behavior. For instance, in granulosa cells, genes from both reference modules were

downregulated in the *Foxl2* haploinsufficiency dataset, despite being upregulated in the Aging model (**Fig.

7b**). Conversely, the MEblue module, downregulated with age in the Aging model, was upregulated in

Foxl2[+/-] granulosa cells. Theca cells showed broadly consistent enrichment across models, whereas

stromal cells exhibited mixed and model-specific patterns (**Fig. 7b**). These findings illustrate both shared

and model-specific regulatory programs and highlight the utility of comparative approaches to dissect

diverse aspects of ovarian aging.

To further explore the biological relevance of these co-expression modules, we performed over-

representation analysis (ORA) on each module's genes using Gene Ontology. In granulosa cells,

enriched terms included components of the ribosome and translational machinery (e.g., "cytosolic

ribosome," "polysome," "cytoplasmic translation"), as well as actin binding and structural molecule activity

(e.g., "A band," "actin binding," and "structural molecule activity"), highlighting coordinated regulation of

protein synthesis and cytoskeletal organization. Notably, enrichment for NAD$^+$ ADP-ribosyltransferase

activity was also observed. This finding aligns with recent evidence linking the NAD$^+$ pathway to the

regulation of ovarian aging[62]. In theca cells, modules were enriched for mitochondrial and contractile

fiber-related terms, including "mitochondrial respiratory chain complex I," "respirasome," and "sarcomere,"

suggesting that transcriptional remodeling in aging theca cells may involve coordinated changes in

energy metabolism and cytoskeletal dynamics. The appearance of the "NADH dehydrogenase complex"

term further supports a role for the NAD$^+$ pathway in theca cell aging[62]. Stromal cell modules were

primarily enriched for extracellular matrix (ECM)-related pathways, such as "collagen-containing

extracellular matrix," "external encapsulating structure," and "basement membrane," consistent with

known stromal remodeling during ovarian aging[63,64]. In BECs, we observed enrichment of RNA-binding

activities, including "poly(U) RNA binding," while epithelial cell modules were enriched for pathways

related to pH regulation and organic acid metabolism, suggesting cell type–specific metabolic adaptations

during aging.

To complement co-expression network-based analyses, we also inferred TF activity across cell

types using decoupleR (**Fig. 7d,e; Extended Data Table 8**). Across datasets, we detected shared

changed TF activity signatures in multiple cell types (**Fig. 7d**). In granulosa cells, we consistently

observed decreased predicted activity of cell cycle–associated factors (E2f1, E2f2, E2f3, E2f4, E2f6, Myc)

and increased predicted activity of Pbx1 (**Fig. 7e**). Pbx1 has been implicated in anti-aging pathways, as

well as tissue and organ homeostasis and regeneration, in hair follicle-derived mesenchymal stem cells

and adrenocortical cells, respectively[65,66]; thus, increased activity in the context of ovarian aging may

reflect a compensatory response to maintain cellular function or structural integrity (**Fig. 7e**). Stromal cells

showed consistent decreased predicted activity of several signaling and remodeling-associated TFs,

including Runx2, Fosl1, Fli1, Klf4, Smad2/3/4, and Stat3. These factors are broadly involved in ECM

remodeling, vascular integrity, immune signaling, and cellular stress responses[46,67-69]; thus, their

coordinated decline may contribute to age-associated stromal dysfunction, fibrosis, and impaired tissue

homeostasis. In epithelial cells, Ppara activity was consistently predicted to decrease across menopause

models. Ppara has been shown to regulate lipid metabolism and oxidative stress responses[70]; its reduced

activity in aged/treated epithelial cells may contribute to metabolic dysregulation associated with

menopause. In DNT cells, Hivep2 and Nfatc1 showed increased activity (**Fig. 7e**). *HIVEP2* has been

implicated in regulatory T cell-mediated immunosuppression in humans[71], while *Nfatc1* is known to control

cytotoxic function in CD8+ T cells in mice[72,73]. Intriguingly, *Nfatc1* has been shown to be important for

regulation of CD8+ T cell metabolism[72]. However, the specific roles of these factors in the context of DNT

cells warrant further investigation.

These analyses reveal both conserved and model-specific gene regulatory programs underlying

ovarian aging. Network-based approaches identified modules and transcription factors linked to protein

synthesis, mitochondrial function, and ECM remodeling, with distinct patterns emerging across cell types

and models.

Transcriptome-based aging clocks capture evidence of accelerated ovarian aging across models

To gain a complementary and integrative view of ovarian aging, we developed transcriptome-

based clocks that predict ovarian age using gene expression data (**Fig. 8a; Extended Data Fig. 15a**),

similar to efforts in other tissues, including blood and brain[74,75]. Given the central role of granulosa cells in

ovarian function and their consistently high responsiveness in our models (see above), we first trained a

lasso regression model using single-cell transcriptomes from granulosa cells (**Fig. 8a**). In parallel, we also

constructed a comparable model using theca cell transcriptomes to evaluate performance in another key

steroidogenic ovarian cell type (**Extended Data Fig. 15a**).

Performance evaluation on the test dataset revealed strong predictive accuracy for the granulosa

cell-based model, with a Spearman correlation of ~0.76 and p-value of ~3.9×10^{-293} between predicted

and true chronological age (**Fig. 8b**). The theca cell-based model also performed well (Rho ~0.70, p-

value ~1.9×10^{-237}; **Extended Data Fig. 15b**), although it showed a dip in prediction accuracy at older

ages. Notably, predictive performance in the theca model starts to decline at ~12 months of age,

coinciding with the reported onset of estropause in mice of the C57BL/6 strain[33] (**Extended Data Fig. 15b;**

age of expected estropause onset highlighted in gray).

Having validated the predictive capacity of the model, we next applied the model to granulosa cell

data from the VCD-treated and *Foxl2*$^{+/-}$ animals (**Fig. 8c,d**). In the VCD model, transcriptome-predicted

ages were consistently higher than chronological age, supporting the notion that VCD exposure

accelerates ovarian aging at the molecular level (**Fig. 8c**). In the *Foxl2* haploinsufficiency model, *Foxl2*$^{+/-}$

animals also showed older predicted transcriptomic ages compared to wild-type controls (**Fig. 8d**). For

granulosa cells, the median predicted age was 27.3 weeks in young *Foxl2*$^{+/+}$, 43.0 weeks in young *Foxl2*$^{+/-}$,

36.3 weeks in middle-age *Foxl2*$^{+/+}$, and 38.5 weeks in middle-age *Foxl2*$^{+/-}$ (**Fig. 8d**).

To further quantify these differences, we calculated transcriptome-based age acceleration by

subtracting true chronological age from predicted age (**Fig. 8e,f**). In the VCD dataset, age acceleration

decreased as age at VCD exposure increased, a trend also observed in the hormone-based OvAge

model, suggesting a reduced magnitude of response to insult in older ovaries (**Fig. 2e-g, 8e**). Granulosa

cells from *Foxl2*$^{+/-}$ animals were predicted to be older than their chronological age, diverging from the

results of OvAge, where condition-matched *Foxl2*$^{+/-}$ animals were predicted to be younger (**Fig. 2h, 8f**).

This discrepancy suggests that the transcriptome-based clock likely detects early molecular alterations in

the ovary that have not yet manifested as systemic hormonal shifts.

We also applied the theca cell-based aging clock to the VCD and *Foxl2* haploinsufficiency models

to determine its predictions on ovarian transcriptional age (**Extended Data Fig. 15c,d**). Despite slightly

lower testing accuracy, the theca cell clock produced trends highly consistent with the granulosa-based

model. Specifically, VCD-treated animals showed elevated transcriptomic age estimates compared to

controls, and *Foxl2*$^{+/-}$ animals similarly exhibited increased predicted ages relative to wild-type

counterparts (**Extended Data Fig. 15c,d**). The median predicted age was 26.7 weeks in young *Foxl2*$^{+/+}$,

34.4 weeks in young *Foxl2*$^{+/-}$, 34.9 weeks in mid-age *Foxl2*$^{+/+}$, and 36.3 weeks in mid-age *Foxl2*$^{+/-}$. We

also calculated age acceleration for the theca cells and observed similar patterns to those seen in

granulosa cells (**Extended Data Fig. 15e,f**). Notably, the difference between age acceleration estimates

was statistically significant for all comparisons, including the middle-age group from the *Foxl2*

haploinsufficiency model, further supporting its sensitivity in detecting molecular ovarian aging

phenotypes across both VCD and *Foxl2* haploinsufficiency models. These findings reinforce the capacity

of the theca cell-based clock to detect molecular signatures of accelerated aging and suggest that both

granulosa and theca cells encode convergent transcriptional readouts of ovarian aging across models.

To explore the transcriptional features contributing to age prediction, features with non-zero β

coefficients from the trained lasso model were extracted and their expression trajectories assessed

across age in pseudobulked granulosa cell dataset (**Fig. 8g**). These features exhibited diverse patterns of

expression over time, suggesting involvement in distinct age-related regulatory programs (**Fig. 8g**). To

further investigate their biological relevance, the top 500 features, ranked by the absolute value of β

coefficients in the final lasso model, were subjected to ORA enrichment (**Fig. 8h**). Enriched terms were

primarily related to immune processes and vascular function, including "leukocyte migration," "interferon-

gamma production," "positive regulation of antigen processing and presentation," and "blood circulation"

and "vascular remodeling" (**Fig. 8h**). These findings suggest that early shifts in immune signaling and

vascular architecture may play a central role in shaping the transcriptomic landscape of ovarian aging and

highlight the sensitivity of the transcriptome-based clock in detecting early molecular changes.

A similar approach was applied to the theca cell-based model, where features with non-zero β

coefficients were extracted and their expression trajectories examined across age in the pseudobulked

theca cell dataset (**Extended Data Fig. 15g**). These features also showed heterogeneous age-dependent

expression patterns, indicating involvement of diverse regulatory programs during theca cell aging. To

evaluate their biological relevance, the top 500 features, ranked by the absolute value of their β

coefficients as above, were also subjected to ORA enrichment (**Extended Data Fig. 15h**). Enriched terms

included immune-related processes such as "regulation of myeloid cell apoptotic process," "regulation of

cytokine production involved in immune response," and "positive regulation of response to external

stimulus," as well as metabolic and structural pathways including "cellular response to lipid," "collagen-

containing extracellular matrix," and components of the "mitochondrial outer membrane" and "membrane

microdomains." These results suggest that transcriptional aging in theca cells may be shaped by

coordinated shifts in immune regulation, lipid metabolism, and mitochondrial and membrane-associated

processes.

Together, these data demonstrate the utility of transcriptome-based models for detecting

molecular aging trajectories in the mouse ovary. These predictive tools offer a valuable framework for

assessing molecular age across experimental models and for uncovering early transcriptomic alterations

that precede overt phenotypic decline.

Discussion

Understanding how ovarian decline contributes to systemic aging requires robust experimental

models that accurately recapitulate the physiological features of menopause. In this study, we

systematically evaluated three mechanistically distinct mouse models of menopause: an intact aging

model, a chemical follicle depletion model using VCD exposure, and a genetic model of *Foxl2*

haploinsufficiency. By integrating histological, hormonal, and transcriptomic profiling across these models,

we provide a comparative framework for understanding shared and model-specific trajectories of ovarian

aging. Findings from our study provide a critical comparative resource for identifying conserved and

divergent features of ovarian aging and evaluating the suitability of each model, with the goal of

advancing both mechanistic insights and translational strategies in menopause research.

In the Aging model, we observed hallmark features of physiological ovarian decline, including

depletion of all follicle stages and disrupted endocrine profiles. These changes were accompanied by

immune cell expansion, increased gene expression variability, elevated TE expression, and significant

transcriptional remodeling in somatic and immune cell types. Granulosa cells consistently exhibited the

highest transcriptional responsiveness, reinforcing their central role in orchestrating ovarian function and

aging. Among immune populations, DNT cells displayed a marked increase in abundance with age. This

expansion was also reported in two independent studies using mouse scRNA-seq datasets[44,47],

reinforcing the emerging role of DNT cells as conserved modulators of ovarian immune remodeling and

aging. Mitochondrial dysfunction, immune activation, and vascular remodeling emerged as common

transcriptional themes in the aging ovary. Gene regulatory network analyses further implicated

downregulation of key transcription factors involved in cell cycle regulation, stress responses, and tissue

homeostasis.

In the VCD model, animals displayed features reminiscent of premature ovarian failure, including

acute follicular depletion, endocrine disruption, and elevated predicted ovarian age based on both OvAge

and transcriptomic clocks. The extent of age acceleration was most pronounced in animals exposed to

VCD at younger ages. This is particularly relevant given that early menopause is a strong predictor of

systemic aging and reduced lifespan in humans[4]. By varying the timing of VCD exposure, this model

could be leveraged to probe the systemic impact of menopausal timing and explore interactions between

reproductive and systemic aging trajectories.

The *Foxl2* haploinsufficiency model revealed a more nuanced aging trajectory. Despite normal or

elevated follicle counts and serum AMH levels in young and mid-age $Foxl2^{+/-}$ animals, transcriptome-

based clocks detected increased molecular age in granulosa and theca cells. Additionally, immune

populations, including neutrophil, NKT and B cells showed enhanced transcriptional responsiveness,

consistent with an immune-primed ovarian environment[35,47]. These results suggest that partial loss of

Foxl2 function may induce early molecular reprogramming prior to overt histological or endocrine decline.

Interestingly, granulosa and stromal cells showed inverse expression of classical aging signatures, and

the model exhibited downregulation of typical senescence pathways and aging modules identified in the

Aging model. *FOXL2* has been shown to interact with *AMH* to modulate follicle recruitment in humans[40].

Thus, the observed histological, endocrine, and transcriptional features may reflect enhanced *Amh*-

mediated reserve preservation in the $Foxl2^{+/-}$ animals. Nonetheless, the preserved endocrine output,

coupled with increased transcriptional sensitivity in steroidogenic and immune compartments, positions

the *Foxl2* haploinsufficiency model as a unique tool for interrogating early or subclinical features of

ovarian aging. Based on the increased transcriptional responsiveness of immune cells and the nuanced

acceleration in transcriptomic aging of granulosa and theca cells, it may be valuable to assess these

animals at older age ranges. The extended timeline could reveal whether early transcriptomic shifts

translate into functional decline and help uncover regulatory mechanisms that are obscured in more overt

aging models.

Cross-species single-cell analyses of ovarian aging further support the translational relevance of

our findings. Studies in mice, non-human primates, and humans have consistently identified conserved

molecular signatures associated with ovarian aging, including reduced oxidative phosphorylation,

diminished antioxidant gene expression, and elevated immune activation across multiple ovarian cell

types[44-46,76]. These transcriptional features were similarly prominent across all three models examined in

our study. Additionally, regulatory network analyses from human ovarian datasets have implicated

transcription factors such as *FOXL1* and *RUNX2* as age-associated regulators[46]. Although these factors

were detected in different cell types in our dataset, their recurrent identification across species suggests

that they may represent broader regulators of ovarian homeostasis. These cross-species parallels

underscore the robustness of the aging signatures captured by our models and highlight the value of this

comparative framework for investigating conserved mechanisms of ovarian aging.

Several mouse genetic models have provided additional insights into molecular features

underlying ovarian decline. In a model with impaired interferon-gamma regulation ($ARE^{+/-}$ and $ARE^{/-}$),

elevated systemic interferon-gamma led to $CD8^+$ T cell infiltration in reproductive tissues and resulted in

infertility[77]. In contrast, we observed a decrease in $CD8^+$ T cell abundance in aged ovaries from our Aging

model, raising the possibility that infiltration may occur earlier in the aging process, followed by depletion

or exhaustion at advanced stages. $CD8^+$ T cells were also sparse in the VCD and Foxl2 models,

suggesting that interferon-mediated immune activation may represent a distinct and context-dependent

mechanism.

A separate study investigating *Cd38* deletion identified early ovarian inflammation and NAD^+

depletion as key features of ovarian aging, with loss of *Cd38* rescuing both transcriptional profiles and

follicular health[62]. In the study, immune cell abundance decreased following *Cd38* deletion[62]. In contrast,

we observed increased immune cell abundance in the Aging and *Foxl2* haploinsufficiency models, and

decreased abundance in the VCD model. Despite these differences, our results converge with the *Cd38*

study on a set of transcription factors potentially involved in regulating ovarian aging. Notably,

transcription factors such as *Myc*, *Egr1*, *Fli1*, and *Klf4*, reported to be restored upon *Cd38* deletion, were

also identified across all three models in our analysis[62]. These shared regulatory nodes point to

conserved, age-sensitive networks that may facilitate physiological decline.

In summary, our comparative analysis reveals that ovarian aging is shaped by a combination of

cell type–specific, model-specific, and shared molecular processes. Intact aging reflects gradual,

widespread physiological decline; VCD exposure triggers acute and accelerated failure; and *Foxl2*

haploinsufficiency initiates early molecular perturbations that may precede functional impairment. By

integrating histological, hormonal, and transcriptomic datasets, we provide a multidimensional framework

for dissecting the mechanistic landscape of ovarian aging. This platform will be valuable for both basic

and translational studies, offering new opportunities to define early biomarkers, and design interventions

to delay or reverse reproductive senescence.

Acknowledgements

We thank Dr. Gilbert Garcia for his feedback on confocal imaging. We also thank Dr. Lynae M. Brayboy for providing the ovarian single-cell dissociation protocol and for her valuable feedback on optimization. We thank Kristen Mehalko for maintaining the *Foxl2* haploinsufficiency colony and contributing to mouse tissue collection. We thank Younggyun Kim, Kelly Koh, and Sanjana Paye for assisting with follicle counting from hematoxylin and eosin-stained ovarian sections. We thank Dr. Victor Ansere, Clayton Baker and Aaron J.J. Lemus for feedback on our manuscript.

This work was supported by the GCRLE-2020 post-doctoral fellowship from the Global Consortium for Reproductive Longevity and Equality at the Buck Institute, made possible by the Bia-Echo Foundation to M.K.; USC Provost's Undergraduate Research Fellowship to J.W.; a Diana Jacobs Kalman/AFAR Scholarships for Research in the Biology of Aging, NIA T32 AG052374, and AthenaDAO student leader award to R.J.L.; GCRLE-0520 Junior Scholar Award from the Global Consortium for Reproductive Longevity and Equality at the Buck Institute, made possible by the Bia-Echo Foundation, Pew Biomedical Scholar award #00034120 from the Pew Charitable Trust, and generous gifts from Kathleen Gilmore and Dr. Eric Hennigan to B.A.B.

Ovarian histological analysis was performed by the Translational Pathology Core at the USC Norris Comprehensive Cancer Center (supported by NCI P30 CA014089). Serum hormone quantification was performed by the University of Virginia Center for Research in Reproduction Ligand Assay and Analysis Core (supported by the Eunice Kennedy Shriver NICHD/NIH Grant R24HD102061).

Some panels were generated using the Generic Diagramming Platform[78].

Author contributions

M.K. and B.A.B. designed the study. R.J.L set-up the original *Foxl2* haploinsufficiency colony. M.K. and R.J.L dissected and collected mouse tissues. E.H.L performed quantitative RT-PCR experiments. M.K. and B.A.B. processed and generated scRNAseq libraries. M.K., R.B., and J.W. analyzed ovarian histology data. R.B. performed RNAscope experiments. M.K. and J.W. performed computational analyses. M.K. and R.W. analyzed and quantified RNAscope images. M.K. and B.A.B. wrote the manuscript with input from all authors. All authors edited and commented on the manuscript.

Declaration of interests

The authors declare no competing interests.

Methods and materials

Mouse husbandry

All animals were treated and housed in accordance with the Guide for Care and Use of

Laboratory Animals. All experimental procedures were approved by the USC's Institutional Animal Care

and Use Committee (IACUC) and are in accordance with institutional and national guidelines. Samples

were derived from animals on approved IACUC protocol numbers 21155 and 21454. All animals were

acclimated in the specific-pathogen-free animal facility at USC for two weeks prior to any experimental

procedures. Mice were provided PicoLab Rodent Diet 20 (LabDiet, 5053) *ad libitum*. The facility was

maintained on a 12-hour light/dark cycle, with housing rooms set to 72°F and 30% humidity.

For the Aging model, female C57BL/6JNia mice (4- and 20-month-old) were obtained from the

National Institute on Aging (NIA) colony at Charles River Laboratories.

For the VCD model, female C57BL/6J mice (2.5-, 5.5-, 7.5-, and 9.5-month-old) were purchased

from Jackson Laboratory. Animals received daily intraperitoneal injections of either vehicle (safflower oil;

Sigma S8281) or VCD (160 mg/kg/day; Sigma 94956) for 15 consecutive days. All injections were

administered between 8:00 and 10:00 AM to minimize variability due to circadian influences.

Foxl2 floxed (*Foxl2^lox/lox^*) mice were generated by Cyagen Biosciences Inc. using a previously

described targeting strategy[25]. Briefly, a targeting construct was designed to replace a 2.177 kb fragment

containing the entire *Foxl2* coding region with a 1.157 kb *Neo* gene cassette (**Extended Data Fig. 1a**).

The construct was linearized and introduced into C57BL/6NTac embryonic stem (ES) cells via

electroporation. Candidate clones were assessed via PCR and Southern blot analysis. A targeted ES cell

clone was selected and injected into C57BL/6 NTac albino embryos, which were then implanted into

pseudo-pregnant CD-1 females. Founder animals were identified based on coat color, and germline

transmission was confirmed through breeding with C57BL/6 females followed by genotyping of the

offspring. Mice carrying the desired floxed *Foxl2* allele were sent to USC for experiments and downstream

phenotyping.

Foxl2 haploinsufficiency mice (*Foxl2^+/-^*) were generated by crossing homozygous *Foxl2^lox/lox^*

C57BL/6NTac mice with heterozygous B6.C-Tg(CMV-cre)1Cgn/J (JAX, stock #006054) mice. Offspring

were genotyped for both the *Foxl2* alleles and the CMV-Cre transgene. To eliminate the CMV-Cre

transgene, *Foxl2*[+/-] mice were crossed with wild-type C57BL/6J mice. Progeny were selected to carry the

deleted Foxl2 allele (*Foxl2*[+/-]), but not the CMV-Cre transgene. Afterwards, *Foxl2*[+/-] mice were maintained

as a stable heterozygous knockout colony using allele transmission through both parent sexes. Routine

genotyping was performed using genomic DNA extracted from tail biopsies using specific PCR primers,

as listed in **Extended Data Table 9a**.

RT-qPCR analysis of *Foxl2* expression in *Foxl2* haploinsufficiency model mouse ovaries

Flash-frozen ovaries collected from mice were used for RNA extraction, cDNA synthesis and

quantitative RT-PCR. Ovaries were resuspended in 600 μL of TRIzol reagent (Thermo-Fisher, 15596018)

and lysed using the BeadBug Homogenizer (Benchmark Scientific, D1036). Homogenization was

performed at 3,500 rpm in 30-second intervals, repeated for a total of nine cycles. Total RNA was then

purified using the Direct-Zol RNA Miniprep kit (Zymo Research, R2052), following the manufacturer's

protocol. RNA was eluted in nuclease-free water sand immediately quantified using the NanoDrop

spectrophotometer (Thermo Scientific). The purity of the RNA was assessed by measuring the

260/280nm and 260/230nm absorbance ratios. Samples with ratios close to ~2.0 were considered

suitable for downstream applications.

Total RNA was reverse transcribed into cDNA, using the Thermo Scientific™ Maxima H Minus

First Strand cDNA Synthesis Kit (Thermo Scientific, K1682), following the manufacturer's protocol.

Quantitative PCR was performed using SensiFAST SYBR no-ROX kit (Bioline, BIO-98020) and the MIC

Tubes and Caps kit (Bio Molecular Systems, 71e101) on the Magnetic Induction Cycler (MIC) machine

(Bio Molecular Systems, MIC-2). micPCR v2.12.6 was used to capture and quantify Ct values. Gene

expression levels were calculated using the ΔΔCt method, normalized to the geometric mean of two

housekeeping genes: *Ubc* and *Hprt*. Primers and sequences used for RT-qPCR reactions can be found in

Extended Data Table 9b.

Fertility assessment of *Foxl2* haploinsufficiency model mice

Fertility data were obtained from our breeding colony, where $Foxl2^{+/+}$ and $Foxl2^{+/-}$ females were

paired with $Foxl2^{+/-}$ and $Foxl2^{+/+}$ males, respectively. Data were included in the fertility assessment only if

$Foxl2^{+/+}$ and $Foxl2^{+/-}$ female pairings were initiated on the same day. Pairings conducted on days when

only $Foxl2^{+/+}$ or only $Foxl2^{+/-}$ females were enrolled were excluded from analysis to avoid any batch-

specific effects on fertility assessments due to uncontrolled routine handling fluctuations.

Female mice between 2.5 and 3 months of age were paired one-to-one with age-matched males.

Fertility was assessed based on the pup count of the first litter and the latency to the first litter (i.e., the

time between pairing and birth of the first litter). Differences in litter size between $Foxl2^{+/+}$ and $Foxl2^{+/-}$

females were assessed using the non-parametric Wilcoxon rank-sum test. One outlier from the $Foxl2^{+/+}$

group was removed based on a significant Grubbs' test (p-value < 0.05), using the outliers package

(v.0.15)[79] in R (v. 4.1.2). Latency to the first litter was analyzed using a log-rank test to compare "survival"

curves, using the survival package (v. 3.6-4)[80] in R. Raw fertility data can be found in **Extended Data**

Table 1a.

Hematoxylin-eosin staining of mouse ovarian tissues

Ovaries were fixed in Bouin's solution (Sigma, HT10132) for 24 hours at room temperature before

being transferred to 70% ethanol for storage. Paraffin embedding, tissue sectioning, and hematoxylin and

eosin (H&E) staining were carried out by the USC Norris Comprehensive Cancer Center Translational

Pathology Core Facility. H&E-stained slides were imaged using the Keyence BZ-X All-in-One

Fluorescence Microscope platform using 20X objective and automated stitching using default parameters.

Ovarian follicle counts

Follicle counts, including primordial, primary, secondary, and antral follicles, and corpus luteum,

were performed on three sections per ovary by three blinded observers. Median values from the three

observers were used for data analysis. Statistical differences between groups were determined using the

non-parametric Wilcoxon rank-sum test.

Quantification of serum AMH, FSH and Inhibin A concentrations

Blood was collected either immediately following euthanasia via cardiac puncture or from live

animals via a facial vein blood draw. Blood was allowed to clot at room temperature for one hour. Serum

was then separated using the MiniCollect® Serum Tube (Greiner, 450472) and stored at -80°C until

further analysis. The University of Virginia Center for Research in Reproduction Ligand Assay and

Analysis Core quantified serum levels of AMH (Rat and Mouse Anti-Müllerian Hormone (AMH) ELISA kit,

Ansh Labs, AL-113), FSH (Millipore Pituitary Panel Multiplex kit, RPT86K or Ultra-Sensitive Mouse & Rat

FSH, UVA Ligand Core, in-house[81]), and Inhibin A (Inhibin A ELISA kit, Ansh Labs, AL-161), providing

standardized normalized values. Statistical differences between groups were evaluated using the non-

parametric Wilcoxon rank-sum test. Raw serum hormone quantification data can be found in **Extended**

Data Table 2.

Ovarian health index calculation

The ovarian health index was calculated as previously described[41]. The index was determined by

integrating two components: ovarian hormone levels (AMH, FSH, and Inhibin A) and follicle counts, which

included the combined counts of primordial, primary, secondary, and antral follicles, as well as corpus

luteum. Each parameter was assigned a score based on a three-tier system. Values exceeding the

median of the young female group were assigned a score of 3, values positioned between the medians of

the young and old groups were assigned a score of 2, and values below the median of the old group were

assigned a score of 1. The hormone score was calculated as the mean of the individual scores for AMH,

FSH, and Inhibin A. This hormone score was then combined with the follicle score in a 1:1 ratio to

generate the overall ovarian health index, which was subsequently scaled to a 0–100 range for

standardization. Statistical differences between groups were evaluated using the Wilcoxon rank-sum test.

Two different FSH assay kits were used due to changes in offered FSH quantitation assays at the

UVA core during the time of our study. To ensure compatibility between measurements made using the

different kits, we applied a correction procedure as previously described[41]. Briefly, matched serum

samples were analyzed using both kits, and a polynomial regression model was trained to capture the

relationship between the two outputs. This model was then applied to adjust FSH values obtained using

the ultra-sensitive assay, enabling direct computation of ovarian health index across datasets.

The scripts used for ovarian health index calculation are available on the Benayoun lab GitHub

repository:

https://github.com/BenayounLaboratory/Ovarian_Aging_Microbiome/tree/main/1_Ovarian_health_index_c

alculation.

OvAge clock model training

The OvAge clock was trained using serum hormone quantification data from multiple model and

cohorts, including animals from aging model, vehicle control animals from VCD model, wild-type animals

from *Foxl2* haploinsufficiency model (*Foxl2$^{+/+}$*) and wild-type animals from a previously published *Fshr*

haploinsufficiency model (*Fshr$^{+/+}$*)[42]. To enhance the applicability of the model across different genetic and

experimental backgrounds, animals from multiple models were included in the training dataset. Hormone

data included serum levels of AMH, FSH, and INHBA, with corresponding age at blood collection

recorded in weeks.

The dataset was partitioned into training (75%) and testing (25%) subsets using stratified

sampling based on age, using the createDataPartition function from R package caret (v. 6.0-91)[82]. A

random forest (RF) regression model was trained using the training set, with out-of-bag (OOB) error

correction applied for internal validation, using caret (v. 6.0-91)[82] and randomForest packages (v. 4.7-1)[83].

The model was optimized by tuning the mtry parameter (best-performing value = 2), yielding an OOB root

mean square error (RMSE) of 12.31 weeks and an OOB R² of 0.67. A total of 500 trees were used in the

final model. Model performance was evaluated using Spearman correlation between chronological and

predicted ages in the test dataset. The RF model achieved a Spearman correlation coefficient of 0.62 with

a p-value of 7.35 × 10^{-8}. Age acceleration was calculated as difference between predicted and

chronological age. The final OvAge clock R object containing the trained RF model is available on the

Benayoun lab GitHub repository: https://github.com/BenayounLaboratory/Mouse_Menopause_Models.

We have also developed an interactive R shiny app to enable users to input hormone values and

estimate predicted ovarian age using the OvAge clock: https://minhooki.shinyapps.io/OvAge_Predictor/.

Single-cell RNAseq sample and library preparation

Ovaries from two animals, totaling four ovaries, were used for each single-cell RNA sequencing

(scRNA-seq) sample and library preparation to enable optimal cellular viability and yield. Single-cell

suspensions were generated using Miltenyi's Multi Tissue Dissociation Kit (Miltenyi Biotec, 130-110-201)

with the gentleMACS Octo Dissociator with Heaters (Miltenyi Biotec, 130-096-427), running the

37_m_LDK1 program. To maximize the recovery of viable single cells, the program was deliberately

aborted 10 sec after start of the highspeed phase, approximately 20 sec before program completion. After

dissociation, red blood cells were removed using Red Blood Cell Lysis Buffer (Miltenyi Biotec, 130-094-

183) according to the manufacturer's instructions. Dead cells were subsequently eliminated using the

EasySep Dead Cell Removal (Annexin V) Kit (STEMCELL Technologies, 17899).

Cell count and viability were assessed via Viobility™ 405/452 Fixable Dye (Miltenyi Biotec,

J66584-AB) staining and flow cytometry using a MACSQuant Analyzer 10 (Miltenyi Biotec, 130-130-420).

Flow cytometry data were analyzed with FlowLogic v8 (Inivai Technologies). Cells were gated based on

forward scatter area (FSC-A) vs. side scatter area (SSC-A), and singlets were identified by evaluating the

linear grouping of cells in FSC-A vs. FSC-H plots (**Extended Data Fig. 7a**).

Single-cell libraries were prepared using the Chromium Next GEM Single Cell 3□ GEM, Library &

Gel Bead Kit v3.1 (10X Genomics, PN-1000121) according to the manufacturer's instructions. Based on

flow cytometry estimates, cell suspensions were loaded to achieve a targeted recovery of 6,000 cells per

sample. Samples were processed on a Chromium Next GEM Chip G (10X Genomics, 2000177) as per

the manufacturer's protocol.

Completed single-cell libraries were assessed for quality using the 4200 TapeStation system

(Agilent Technologies, G2991A) with High Sensitivity D1000 DNA ScreenTape (Agilent Technologies,

50675584). Libraries were sequenced on an Illumina NovaSeq 6000, generating 150 bp paired-end reads

at Novogene USA. Raw FASTQ reads have been deposited in the Sequence Read Archive under

accession PRJNA863443.

Single-cell RNAseq data analysis

Data processing

Raw sequencing reads were processed using Cell Ranger (v. 7.1.0)[84] from 10x Genomics

(**Extended Data Table 3**). Reads were aligned to the mm10 reference genome. To remove contamination

from ambient RNA, decontamination was performed using DecontX from the celda (v. 1.14.2)[85] package.

Raw and filtered feature-barcode matrices were used to estimate ambient RNA levels, and corrected

expression matrices were generated for each sample. Doublet removal was performed using a two-step

approach. First, DoubletFinder (v. 2.0.3)[86] was applied to each sample independently, estimating doublet

rates based on the expected multiplet rates from 10x Genomics. The optimal pK parameter was

determined using a parameter sweep, and predicted doublets were labeled. In parallel, a second doublet

detection method, scds (v. 1.10.0)[87], was applied using the cxds_bcds_hybrid function. Cells identified as

doublets by either method were excluded from further analysis.

The filtered single-cell expression matrices were loaded into Seurat (v. 4.3.0)[88], and initial quality

control (QC) steps were applied. Cells with fewer than 500 detected features, a total RNA count outside

the range of 500–100,000, mitochondrial RNA content exceeding 15%, or a DecontX contamination score

above 0.25 were removed. Following QC, datasets were normalized using SCTransform, regressing out

the number of detected features, mitochondrial content, DecontX contamination score and batch effects.

Data integration

To correct for batch effects in the VCD and *Foxl2* haploinsufficiency datasets, integration was

performed using Harmony (v. 1.0)[89]. First, datasets were processed separately by normalizing gene

expression using SCTransform, regressing out number of detected features (nFeature_RNA),

mitochondrial percentage (percent.mito), ambient RNA contamination (decontX_contamination), and

sequencing library (Library). Highly variable genes were selected using vst (nfeatures=2000 genes). The

preprocessed datasets were merged using Seurat (v. 4.3.0)[88], ensuring shared variable features were

maintained. Principal component analysis (PCA) was performed on the merged dataset, and batch effects

were assessed by visualizing PCA embeddings colored by batch identity. Batch correction was applied

using Harmony, where batch information was explicitly modeled as a covariate. The corrected

embeddings were used for downstream analysis, including UMAP visualization, clustering, and differential

expression analysis.

<u>Cell type annotation</u>

Cell type annotation was conducted using a multi-step approach that combined automated

reference-based methods with manual marker-based annotation. First, *Ptprc⁺* and *Ptprc⁻* cell populations

were identified using scGate (v. 1.0.1)[90]. Cells expressing *Ptprc* (Cd45) were classified as immune cells,

while those lacking *Ptprc* expression were designated as non-immune cells. The two subsets were then

processed separately for further annotation. Each subset underwent independent cell type annotation

using three complementary computational methods: SingleR (v. 1.8.1)[91], scSorter (v. 0.0.2)[92], and scType

(v. 1.0)[93]. The ImmGen dataset[94] was used as a reference for *Ptprc⁺* cells, while two publicly available

ovarian single-cell RNA-seq datasets (Open Science Framework ID 924fz[95] and NCBI GEO accession no.

GSE232309[44]) were used for *Ptprc⁻* cells. To supplement these automated methods, manual annotation

was conducted based on the expression of canonical marker genes specific to ovarian cell types

(**Extended Data Table 4**). The final annotation was determined using a majority voting approach – cell

type labels from SingleR, scSorter, scType, and manual annotation were compared, and the most

frequently assigned label was selected as the final classification. All analyses were performed using

Seurat (v. 4.3.0)[88] within R.

<u>Cell type proportion and Augur analysis</u>

Cell type proportions were quantified using scProportionTest (v. 0.0.0.9000)[96] within R. This

method performs a permutation-based test to compare cell-type abundances between two groups,

estimating the relative change in proportions along with a confidence interval for each cell type. The

analysis was conducted at two levels: the broader classification of *Ptprc⁺* vs. *Ptprc⁻* populations and the

more detailed sub-cell type level within each category.

We used Augur (v. 1.0.3)[48] to identify cell types that exhibited the most transcriptional changes

associated with aging. Pairwise comparisons were conducted for each experimental condition, assessing

differences between (1) young and old females, (2) vehicle control (CTL) and VCD-treated groups, and (3)

Foxl2⁺/⁺ and *Foxl2⁺/⁻* genotypes.

Pseudobulk analysis for differential gene expression

Pseudobulk differential gene expression analysis was conducted separately for three datasets:

the Aging, VCD, and *Foxl2* haploinsufficiency models. Single-cell transcriptomic data were aggregated at

the cell type level using muscat (v. 1.18.0)[97] to generate pseudobulk expression profiles. To ensure

robust analysis, cell types were filtered based on dataset-specific criteria. In the Aging model, cell types

with at least 25 cells per sample across all samples were retained. For the VCD model, cell types were

included if they contained at least 25 cells in at least 12 samples, representing 75% of the dataset. In the

Foxl2 haploinsufficiency model, the threshold was set at a minimum of 25 cells in at least 8 samples, or

80% of the dataset.

Differential gene expression analysis was performed using DESeq2 (v. 1.44.0)[51], with

comparisons made between young and old females in the Aging model, vehicle control (CTL) and VCD-

treated samples in the VCD model, and *Foxl2*$^{+/+}$ and *Foxl2*$^{+/-}$ samples in the *Foxl2* haploinsufficiency

model. Batch effects in the VCD and *Foxl2* haploinsufficiency models were assessed using sva (v.

3.52.0)[98] to determine the number of surrogate variables (SVs), and batch correction was applied using

the removeBatchEffect() function from limma (v. 3.60.6)[99]. The adjusted counts were subsequently used

for DESeq2 analysis, incorporating age-at-injection and time post-injection as covariates for the VCD

model and age as a covariate for the *Foxl2* haploinsufficiency model. Variance-stabilized transformed

counts were computed using getVarianceStabilizedData(), and differentially expressed genes (FDR <

0.05) were visualized using strip plots. The complete pseudobulk DESeq2 output, including log fold

changes, adjusted p-values, and base mean expression values for each dataset and cell type, can be

found in **Extended Data Table 6**.

1013

GO analysis for differential gene expression

Gene set enrichment analysis (GSEA) was performed to identify Gene Ontology Biological

Process (GO BP) and Reactome terms enriched in differentially expressed genes across cell types in the

Aging, VCD, and *Foxl2* haploinsufficiency models. The analysis was conducted using the clusterProfiler

package (v. 4.2.2)[100] and the Molecular Signatures Database (MSigDB) gene sets retrieved via msigdbr

(v. 7.4.1)[101]. Differential expression results from pseudobulk DESeq2 analysis were ranked by t-statistic

and used as input for GSEA. Enrichment analysis was performed separately for each cell type in each

dataset, with an FDR cutoff of 0.1 applied to identify significant terms. To identify recurrent biological

processes across datasets, GO-BP terms enriched in at least four cell types within each dataset were

extracted. The list of significantly enriched GO-BP terms (FDR < 0.1) for all datasets can be found in

Extended Data Table 7.

1025

GSEA of aging-associated transcriptional signatures across models

GSEA was performed using differentially expressed genes (DEGs) from the Aging model dataset

as input gene sets with the clusterProfiler (v4.12.6)[100] package. Upregulated and downregulated gene

sets (FDR < 0.05) were extracted from the Aging model dataset and tested for enrichment in the Aging,

VCD and Foxl2 haploinsufficiency model datasets. Genes were ranked based on the DESeq2-derived t-

statistic for analysis. Additionally, GSEA was conducted to assess the enrichment of the SenMayo[53] gene

set within the dataset. Differential expression results from the Aging, VCD, and Foxl2 datasets were used

for GSEA, with genes ranked using the DESeq2-derived t-statistic.

1034

Coefficient of variation analysis

Gene expression variability across cell types was assessed by calculating the coefficient of

variation (CV) for each gene in the Aging, VCD, and *Foxl2* haploinsufficiency models. CV was computed

as the ratio of the standard deviation to the mean expression level per gene within each condition (young

vs. old for Aging, CTL vs. VCD for VCD, and *Foxl2$^{+/+}$* vs. *Foxl2$^{+/-}$* for *Foxl2* haploinsufficiency model) using

variance-stabilized transformed counts obtained from DESeq2 (v1.44.0)[51] pseudobulk analysis. Statistical

differences in CV between groups were evaluated using the Wilcoxon rank-sum test.

1042

Transposable element (TE) analysis

Transposable element (TE) expression was quantified using scTE (v1.0)[56] with default

parameters. BAM files generated from Cell Ranger (v. 7.1.0)[84] outputs were processed using the mm10

genome index provided by scTE. Features detected by scTE but absent in the single-cell gene

expression data were extracted and merged, generating a Seurat object containing both gene and TE

expression profiles.

Cells were aggregated at the cell type level using muscat (v1.18.0)[97], and differential expression

analysis was performed using DESeq2 (v1.44.0)[51], following the same approach as described above. The

VCD and *Foxl2* haploinsufficiency datasets underwent batch correction as before.

To assess functional enrichment of TE families, GSEA was conducted on ranked gene lists

derived from DESeq2 results. TE annotations were obtained from the UCSC database for mm10, the

same reference used by scTE. TE subfamilies were stratified to broader categories, including LINE, SINE,

LTR, and DNA transposons. TE family annotations were obtained from the *rmsk.txt* file downloaded from

the UCSC genome annotation database for the mm10 reference genome. For each cell type, genes were

ranked by t-statistic from differential expression analysis, and enrichment of TE families was assessed

using clusterProfiler (v4.12.6)[100]. Significant TE families (FDR < 0.1) were reported for visualization.

1059

WGCNA and ORA enrichment analysis

Weighted gene co-expression network analysis (WGCNA)[60] was performed on the Aging model

dataset to identify gene co-expression modules. Variance-stabilized transformed counts from DESeq2

(v1.44.0)[51] were used as input, with analyses conducted separately for each cell type. For each cell type,

an optimal soft-thresholding power was determined using pickSoftThreshold, followed by the construction

of an adjacency matrix and topological overlap matrix (TOM). Hierarchical clustering was performed to

identify gene modules, which were assigned colors using cutreeDynamic, and eigengenes were

computed to summarize module expression patterns. Gene sets from each module were classified into

upregulated and downregulated groups based on eigengene expression differences between young and

old samples. These gene sets were then subjected to GSEA via clusterProfiler (v. 4.2.2)[100], using

differential expression results from the VCD and *Foxl2* haploinsufficiency model datasets to assess

whether the identified modules were significantly enriched in these datasets. Modules significantly

associated with aging (FDR < 0.1) were further analyzed through over-representation analysis using

clusterProfiler (v. 4.2.2) [100] and the org.Mm.eg.db annotation package.

1074

 Transcription factor activity inference using decoupleR

 Transcription factor (TF) activity inference was performed using decoupleR (v2.10.0)[61] with the

 CollecTRI transcriptional regulatory network for mouse. Differential expression results from the Aging,

 VCD, and *Foxl2* haploinsufficiency models were used as input. For each dataset, fgsea scoring was

 applied to rank TF activity based on DESeq2-derived t-statistics, and significant TFs were identified at a

 threshold of FDR < 0.05. To compare TF activity across datasets, TFs were filtered to retain those with

 consistent activity direction (positive or negative) in at least two datasets.

 R shiny application generation

 An interactive R shiny application of Aging, VCD and Foxl2 haploinsufficiency scRNA-seq

 datasets was generated using ShinyCell (v. 2.1.0)[102], and made available at

 https://minhooki.shinyapps.io/shinyappmulti/.

 RNAscope sample preparation, imaging and data analysis

 In situ Hybridization Protocol

 Frozen ovaries were mounted in Tissue-Tek O.C.T Compound (Sakura, 4583) and sliced into 8

 µm slices from the same ovarian region using PTFE-coated microtome blades (Duraedge, 7223) at

 −20°C on a Cryostat CM1860 (Leica, 14-0491-46884). 3 ovarian sections from the same region were

 mounted in succession on VWR Premium Superfrost Plus Microscope slides (VWR, 48311-703). All

 batches of sample processing were prepared using the fresh-frozen sample preparation protocol along

 with negative and positive controls (Advanced Cell Diagnostics, UM 323100-USM). 12 probes were

 tested in 3 sets, with Set 1 containing markers for theca (*Cyp11a1*; C1), granulosa (*Fshr*; C2), smooth

 muscle (*Acta2*; C3), and stromal (*Pdgfra*; C4) cells. Set 2 had markers for epithelial cells (*Upk1b*; C1),

 lymphatic endothelial cells (*Prox1*; C2), blood endothelial cells (*Flt1*; C3), and B cells (*Cd19*; C4). Set 3

 contained probes for T (*Cd3e*; C1), natural killer (*Klr1b1c*; C2), CD8+ (*Cd8b1*; C3), and CD4+ (*Cd4*; C4)

 cells. Per the protocol, the RNAscope[TM] Multiplex Fluorescent Reagent Kit v2 (Advanced Cell Diagnostics,

 323100) was used. The fluorescent signals TSA Vivid Fluorophore 520 (Advanced Cell Diagnostics,

 323271), TSA Vivid Fluorophore 570 (Advanced Cell Diagnostics, 323272), TSA Vivid Fluorophore 650

1103   (Advanced Cell Diagnostics, 323273), and Opal 690 (Akoya Biosciences, FP1497001KT) were used to

1104   stain the probes in each set. The slides were preserved in ProLong Gold Antifade Mountant (Thermo

1105   Fisher Scientific, P36930).

1106

1107   <u>Imaging</u>

1108   Imaging was performed using a Leica Stellaris 5 confocal microscope (Leica Microsystems)

1109   equipped with a HC PL APO 20x/0.75 IMM CORR CS2 objective (Leica, 11506343) and Leica

1110   Microsystems Type F Immersion liquid oil (Leica Microsystems, 11513859). Images were acquired using

1111   Leica Application Suite X software (LAS X v4.4.0).

1112

<u>Imaging analysis</u>

To quantify probe expression, the images were analyzed using the QuPath program v0.5.1. Cell

boundaries based on DAPI signal were calculated using the following settings modified from the default

parameters – Requested pixel size: 0 μm, Nucleus background radius: 10 μm, Maximum area: 400 $μm^2$,

Threshold: 0.5, and Cell expansion: 2. Sets 1 and 2 were analyzed using subcellular detection to

determine their expression levels relative to the number of cells. All thresholding values were uniform

across the date of processing of each ovarian tissue. For Set 3, single measurement classifiers were

created for each probe to generate colocalization data. Cells below the detection threshold were not

counted for analysis.

1122

Cell-type proportion analysis using flow cytometry

Single-cell suspensions of ovarian cells were prepared following the same protocol used for

single-cell RNA sequencing sample preparation, as described above. Cells were stained for APC-CD45

antibody (Miltenyi, 130-123-784) according to the manufacturer's recommendations. Flow cytometry data

were acquired using the MACSQuant10 flow cytometer (Miltenyi Biotec, 130-096-343) and analyzed

using FlowLogic v.8 (Inivai Technologies). The raw flow cytometry dataset was deposited to Figshare

(DOI: 10.6084/m9.figshare.28733222).

1130

Granulosa and theca transcriptional clock model training and feature assessment

Granulosa and theca cells were independently subsetted from scRNA-seq datasets encompassing multiple experimental models, including natural aging, VCD, and *Foxl2* haploinsufficiency. Additionally, an independent middle-aged female dataset (13, 76, and 86 weeks) was included to ensure age coverage for intermediate ages not well represented in the datasets.

To develop the cell transcriptional clocks, cells from aging model, middle-aged females, vehicle controls from the VCD model, and *Foxl2+/+* animals from the *Foxl2* haploinsufficiency model were extracted from the dataset. These cells were used to train and evaluate the model. Data were partitioned into training (75%) and testing (25%) sets using stratified sampling, using the createDataPartition function from the caret package (v. 6.0-91). Training data were used for model optimization, while the test set served as an independent validation dataset.

Predictive models were trained using L1-regularized lasso regression with the glmnet package (v. 4.1-7) to estimate transcriptional age (in weeks) based on gene expression profiles. Five-fold nested cross-validation was performed on the training dataset, and the optimal regularization parameter (lambda.min) was determined based on cross-validation results. Spearman correlation analysis was performed to assess the correlation between predicted and chronological age.

Transcriptional clocks were applied to experimental test groups, including VCD-treated and *Foxl2*[+/-] animals. Model predictions for test groups were compared to those of control samples to assess differences in predicted age patterns. Additionally, age acceleration was calculated as the difference between predicted and chronological age. Statistical significance of group differences in predicted age and age acceleration was assessed using the Wilcoxon rank-sum test.

To examine the age-associated expression dynamics of lasso features selected by the models, we performed pseudobulk analysis using control samples. For each cell type, raw counts were aggregated per sample using muscat (v. 1.18.0)[97], and DESeq2 (v. 1.44.0)[51] was used to compute variance-stabilized transformed expression values. Batch effects were corrected using removeBatchEffect, and expression heatmaps of lasso-selected features were generated to visualize age-dependent expression trends.

1158   The top 500 lasso features were selected based on the absolute values of their coefficients.

1159   Over-representation analysis was performed using clusterProfiler (v. 4.2.2) and org.Mm.eg.db (v. 3.14.0),

1160   analyzing "ALL" Gene Ontology terms. A p-value cutoff of 0.05 was applied to identify significantly

1161   enriched pathways.

1162

Quantification and statistical analysis

All statistical analysis was performed using the R software, version 4.1.2. For all boxplots, the

data is shown with the median, the 25th and 75th percentile of the data, and the whiskers represent 1.5 $*$

the inter-quartile range (IQR). Individual datapoints are overlayed, when possible, for transparency and

rigor. Specific statistical tests, number of biological replicates and animals used are indicated in the

corresponding figure legends and associated methods.

1169

Data availability

The raw FASTQ files have been deposited in the Sequence Read Archive under accession

number PRJNA863443. Raw microscopy images of ovarian hematoxylin and eosin (H&E) staining,

compressed RNAscope images (due to file size constraints), and raw flow cytometry datasets are

available on Figshare (DOIs: 10.6084/m9.figshare.28745765, 10.6084/m9.figshare.28745792,

10.6084/m9.figshare.28745813, 10.6084/m9.figshare.29260391 and 10.6084/m9.figshare.28733222).

The final annotated Seurat objects have been deposited to Figshare (10.6084/m9.figshare.28934126)

and can be explored via an interactive R shiny application: https://minhooki.shinyapps.io/shinyappmulti/.

All analyses were performed using R version 4.1.2. Any additional information required to reproduce or

reanalyze the data presented in this study is available from the corresponding author upon reasonable

request.

1181

1182   **Code availability**

1183   All new scripts used to analyze the datasets are available on the Benayoun lab GitHub at

1184   https://github.com/BenayounLaboratory/Mouse_Menopause_Models.

    Figure legends

    Fig 1. Characterization of ovarian health markers of animals from Aging, VCD and *Foxl2*

    haploinsufficiency models. a, Schematic of the experimental design. **b-d,** Representative hematoxylin-

    eosin staining images of ovarian tissues from Aging, VCD, and *Foxl2* haploinsufficiency model mice. For

    the VCD model, ovarian tissues were analyzed 5 months post-injection. **e-g,** Combined follicle counts

    (primordial, primary, secondary and antral follicles, and corpus luteum) from Aging, VCD, and *Foxl2*

    haploinsufficiency model mice. **h-j,** Serum AMH level quantification from Aging, VCD, and *Foxl2*

    haploinsufficiency model mice. **k-m,** Serum FSH level quantification from Aging, VCD, and *Foxl2*

    haploinsufficiency model mice. **n-p,** Serum INHBA level quantification from Aging, VCD, and *Foxl2*

    haploinsufficiency model mice. **q-s,** Ovarian health index from Aging, VCD, and *Foxl2* haploinsufficiency

    model mice. For panels **e-s**, statistical significance was assessed using the Wilcoxon test, and p-values

    are reported. Open data points indicate data used for ovarian health index calculation. Scale bar: 250 µm.

    Fig 2. Development and application of OvAge, a hormone-based ovarian aging clock. a, Schematic

    representation of the OvAge clock development pipeline. Data from all animals from the Aging model,

    vehicle control (CTL) from the VCD model, *Foxl2*[+/+] from the *Foxl2* haploinsufficiency model, and *Fshr*[+/+]

    from the *Fshr* haploinsufficiency model mice were used to train and test OvAge. **b,** Scatter plot comparing

    predicted and chronological age, reported in weeks, in the test dataset. **c-d,** Scatter plots comparing

    predicted and chronological age for VCD and *Foxl2* haploinsufficiency model mice. **e-g,** Age acceleration

    analysis for VCD model animals at 30-, 90- and 150-days post-injection. **h,** Age acceleration analysis for

    Foxl2 haploinsufficiency animals. Age acceleration was measured as the difference between predicted

    OvAge and chronological age. Open data points indicate data from out-of-bag predictions. For panels **e-h**,

    statistical significance was assessed using the Wilcoxon test, and p-values are reported.

    Fig 3. scRNA-seq profiling of ovaries from Aging, VCD, and *Foxl2* haploinsufficiency model mice.

    a, Schematic of the experimental design. **b,** Schematic of sample collection timepoints. **c-e**, UMAP plots

    of scRNA-seq datasets from Aging, VCD and *Foxl2* haploinsufficiency models. **f-h**, Dotplot representation

1213   of expression of marker genes of cell types detected in scRNA-seq datasets from Aging, VCD and *Foxl2*

1214   haploinsufficiency models. **i**, Representative RNAscope images of DAPI, *Klrb1c*, *Cd3e*, *Cd8b1* and *Cd4*

1215   probes. Images were enhanced for visualization. The shown image is from a middle-aged *Foxl2*$^{+/+}$ animal.

1216

Fig 4. Characterization of cell type proportions in Aging, VCD, and *Foxl2* haploinsufficiency model

scRNA-seq datasets. a, Schematic of the experimental design. **b-d**, Proportion differences of *Ptprc*$^{-}$ and

Ptprc^{+} cells between young *vs.* old in Aging model, CTL (vehicle control) *vs.* VCD in VCD model, and

Foxl2$^{+/+}$ *vs. Foxl2*$^{+/-}$ in the *Foxl2* haploinsufficiency model datasets. **e-g,** Flow cytometry analysis of CD45^{+}

cell proportions in ovaries of Aging, VCD, and *Foxl2* haploinsufficiency model mice. **h-j,** Proportion

differences of non-immune cell types detected in Aging, VCD, and *Foxl2* haploinsufficiency model

scRNA-seq datasets. **k-m,** Representative RNAscope images of *Fshr* (granulosa) and *Cyp11a1* (theca)

probes from Aging, VCD, and Foxl2 haploinsufficiency model mice. Scale bar: 100µm.   **n-o,** Cell

abundance quantification data from RNAscope image analysis of *Fshr*$^{+}$ and *Cyp11a1*$^{+}$ cells from Aging,

VCD, and *Foxl2* haploinsufficiency model mice. For panels **e-g** and **n-p**, statistical significance was

assessed using the Wilcoxon test, and p-values are reported.

1228

Fig 5. Comparative analysis of global gene expression in Aging, VCD and *Foxl2* haploinsufficiency

models via Augur. a, Schematic of global gene expression analysis comparisons using Augur. **b-c,**

UMAP visualization of AUC scores and lollipop plot of AUC quantification from the Aging model. **d-e**,

UMAP visualization of AUC scores and scatter plot of AUC quantification from the VCD model, comparing

CTL *vs.* VCD at 3m and 10m age-at-injection. For the scatter plot, *Ptprc*$^{-}$ and *Ptprc*$^{+}$ cells were separately

plotted to improve visualization. All data are from the 90d post-injection timepoint. **f-g,** UMAP visualization

of AUC scores and scatter plot of AUC quantification from the *Foxl2* haploinsufficiency model comparing

Foxl2$^{+/+}$ *vs. Foxl2*$^{+/-}$ at young and mid-age. For the scatter plot, *Ptprc*$^{-}$ and *Ptprc*$^{+}$ cells were separately

plotted to improve visualization. For **e** and **g**, data points with NA AUC scores (due to low cell count or QC

filtering) were assigned an AUC of 0.5 and colored gray to improve visualization.

1239

Fig 6. Pseudobulk analysis of differential gene expression and gene ontology analysis across

Aging, VCD, and *Foxl2* haploinsufficiency models. a, Schematic of pseudobulk analysis pipeline. **b-d,**

Strip plots of differentially expressed genes (DEGs) from Aging, VCD and *Foxl2* haploinsufficiency models.

DEGs were identified using DESeq2, comparing young *vs.* old (Aging model), CTL *vs.* VCD (VCD model),

and *Foxl2*$^{+/+}$ *vs. Foxl2*$^{+/-}$ (*Foxl2* haploinsufficiency model). Genes that passed the FDR < 0.05 threshold

are colored and the numbers in parentheses following each cell type indicate the total number of DEGs

that met the significance threshold. **e-f**, Frequent Gene Set Enrichment Analysis (GSEA) term enrichment

for GO Biological Process (GO BP, FDR < 0.05) and Reactome (FDR < 0.1) pathways across Aging, VCD,

and *Foxl2* haploinsufficiency models. Terms enriched in at least four and three cell types within each

dataset were extracted for GO BP and Reactome, respectively.

1250

Fig 7. Preservation of co-expression modules and transcription factor activity in Aging, VCD, and

***Foxl2* haploinsufficiency models. a,** Schematic of WGCNA and transcription factor (TF) activity

inference analysis. **b,** GSEA enrichment of WGCNA module genes associated with aging traits across

Aging, VCD and *Foxl2* haploinsufficiency models. WGCNA modules significantly associated with aging

traits (FDR < 0.1) were identified for each cell type from the Aging model. **c,** Over-representation analysis

of GO ALL terms for WGCNA module genes (FDR < 0.1). **d,** UpSet plot of decoupleR-inferred TF activity,

showing the number of TF overlap across datasets and the corresponding cell types. **e**, Heatmap of TF

activity scores of common decoupleR-inferred TFs for each cell type.

1259

Fig 8. Development of a transcriptome-based aging clock for granulosa cells. **a,** Schematic of

analysis and training workflow for the lasso regression-based clock. **b,** Scatter plot of predicted age *vs.*

chronological age for the test set, with Spearman's correlation and p-value displayed. **c-d,** Age prediction

for VCD and *Foxl2* haploinsufficiency datasets, comparing predicted age to chronological age in weeks.

Offsets were added to improve visualization. **e-f,** Age acceleration analysis for VCD and *Foxl2*

haploinsufficiency datasets, calculated as the difference between predicted and chronological age.

Statistical significance was assessed using the Wilcoxon test. **g,** Heatmap of expression lasso features

from pseudobulked expression data. **h,** Over-representation analysis of GO ALL terms of the top 500

lasso features.

Extended Data Figure legends

1270

Extended Data Fig. 1. Characterization of the *Foxl2* haploinsufficiency model. a, Schematic

representation of constructs for *Foxl2* haploinsufficiency mouse line generation. **b,** *Foxl2* expression

levels in *Foxl2*[+/-] and *Foxl2*[+/-] mice ovaries from young and mid-age groups, measured by RT-qPCR. P-

values were calculated using the Wilcoxon test. **c-d,** Fertility assessment of *Foxl2* haploinsufficiency

females, comparing first litter pup counts and latency to first pregnancy. P-values were calculated using

the Wilcoxon and log-rank tests, respectively.

1277

1278 **Extended Data Fig. 2. Follicle counts across Aging, VCD and *Foxl2* haploinsufficiency models. a-c,**

1279 Follicle counts for primordial, primary, secondary and antral follicles, and corpus luteum, from Aging, VCD

1280 and *Foxl2* haploinsufficiency model mice. P-values were calculated using the Wilcoxon test. Illustrations

1281 of ovarian follicles were obtained from Generic Diagramming Platform[78].

1282

1283 **Extended Data Fig. 3. Longitudinal analysis of serum AMH, FSH, and Inhibin A levels in the VCD**

1284 **model. a-c,** Serum levels of AMH, FSH, and Inhibin A (INHBA) measured from 0 to 5 months post-

1285 injection in VCD model mice.

1286

1287 **Extended Data Fig. 4. Histological characterization of the VCD model mice. a,** Representative

1288 hematoxylin and eosin staining images of ovarian tissues from condition-matched animals used in the

1289 VCD model scRNA-seq analysis. **b,** Follicle counts for the condition-matched animals used in the VCD

1290 model scRNA-seq analysis. Satistical significance was assessed using the Wilcoxon test, and p-values

1291 are reported.

1292

Extended Data Fig. 4. Serum hormone quantification and ovarian health index calculation of the

VCD model mice. a-c, Serum levels of AMH, FSH, and Inhibin A (INHBA) measured from condition-

matched animals used in the VCD model scRNA-seq analysis. **d**, Adjusted FSH values to account for

inter-assay differences between measurement assays. **e,** Ovarian health index from condition-matched

animals used in the VCD model scRNA-seq analysis. Statistical significance was assessed using the
Wilcoxon test, and p-values are reported.

1299

Extended Data Fig. 6. Validation of detected ovarian cell types by *in situ* hybridization. a,
Representative RNAscope images for *Fshr, Cyp11a1, Pdgfra* and *Acta2* probes. **b,** Representative
RNAscope images for *Upk1b, Cd19, Flt1* and *Prox1* probes. **c,** Representative RNAscope images for
Klrb1c, Cd3e, Cd8b2 and *Cd4*. Images were enhanced for visualization. **d,** Heatmap of detected cell
types in the Aging, VCD and *Foxl2* haploinsufficiency scRNA-seq datasets. For panels **a-c**, shown
images are from a middle-aged *Foxl2*$^{+/+}$ animal.

1306

1307 **Extended Data Fig. 7. Cell type proportion analysis of ovarian cells via flow cytometry and scRNA-**
1308 **seq analysis. a,** Gating strategy of CD45^{+} cells in flow cytometry analysis. **b-d**, Proportion differences of
1309 *Ptprc*+ cell types detected in Aging, VCD and *Foxl2* haploinsufficiency model scRNA-seq datasets.
1310 Asterisks (*) indicate comparisons in which the observed log$_2$ fold change is infinite due to complete
1311 absence of the cell type in one of the groups.

1312

1313 **Extended Data Fig. 8. Cell type proportion analysis of non-immune ovarian cells via *in situ***
1314 **hybridization assays. a-c,** Cell type abundance analysis of stromal, SMC, BEC, LEC and epithelial cells
1315 via RNAscope *in situ* hybridization assays from Aging, VCD and *Foxl2* haploinsufficiency model mice.

1316

1317 **Extended Data Fig. 9. Cell type proportion analysis of immune ovarian cells via *in situ***
1318 **hybridization assays. a-c,** Cell type abundance analysis of NK, NKT, CD8^{+} NKT, CD8^{+} T, CD4^{+} T, DNT,
1319 DPT and B cells via RNAscope *in situ* hybridization assays from Aging, VCD and *Foxl2* haploinsufficiency
1320 model mice.

1321

Extended Data Fig. 10. Comparative analysis of global gene expression in VCD model via Augur.
a-b, UMAP visualization of AUC scores and scatter plot of AUC quantification in the VCD model,
comparing CTL vs. VCD at 3m and 10m age-at-injection. For the scatter plot, *Ptprc*$^{-}$ and *Ptprc*$^{+}$ cells were

separately plotted to improve visualization. All data are from the 30d post-injection timepoint. For **b**, data

points with NA AUC scores (due to low cell count or QC filtering) were assigned an AUC of 0.5 and

colored gray to improve visualization.

1328

Extended Data Fig 11. Aging-associated signatures and transposable element enrichment

analysis across Aging, VCD and *Foxl2* haploinsufficiency models. a, Heatmap of cell types detected

in pseudobulked scRNA-seq datasets. **b,** DESeq2 normalized $\log_2$ counts of *Foxl2* from young and mid-

age *Foxl2*$^{+/+}$ and *Foxl2*$^{+/-}$ mice. **c-d,** GSEA enrichment analysis of differentially expressed genes (DEGs)

identified in the Aging model (FDR < 0.05) and SenMayo gene list. **e,** Dot plot showing the coefficient of

variation analysis results. **f,** GSEA enrichment analysis of transposable element (TE) families across

Aging, VCD and *Foxl2* haploinsufficiency models.

1336

1337 **Extended Data Fig 12. Module trees of WGCNA analysis. a,** WGCNA module trees identified in the

1338 Aging model for granulosa, theca, stromal, BEC, epithelial, and DNT cells.

1339

1340 **Extended Data Fig 13. Association analysis of WGCNA modules and ovarian aging traits from the**

1341 **Aging model. a,** Heatmap of trait significance for WGCNA modules in the Aging model. Modules that

1342 passed the FDR < 0.1 threshold are shown in bold.

1343

1344 **Extended Data Fig 14. Analysis of recurrent transcription factor activity inferred by decoupleR. a-c,**

1345 Heatmap of transcription factor (TF) activity for factors detected in at least three cell types within each

1346 model with an FDR < 0.05.

1347

Extended Data Fig 15. Development of a transcriptome-based aging clock for theca cells. **a,**

Schematic of analysis and training workflow for the lasso regression-based clock. **b,** Scatter plot of

predicted age *vs.* chronological age for the test set, with Spearman's correlation and p-value displayed. **c-**

d, Age prediction for VCD and *Foxl2* haploinsufficiency datasets, comparing predicted age to

chronological age in weeks. Offsets were added for visualization purposes. **e-f,** Age acceleration analysis

for VCD and *Foxl2* haploinsufficiency datasets, calculated as the difference between predicted and

chronological age. Statistical significance was assessed using the Wilcoxon test. **f-g,** ORA enrichment

analysis of GO ALL and Reactome terms of the top 500 lasso features.

Extended Data Tables

1357

Extended Data Table 1. Raw fertility data from *Foxl2* haploinsufficiency model mice.

Extended Data Table 2. Raw serum AMH, FSH and INHBA quantification data. Raw data from **a**,

Aging, **b**, VCD and **c**, *Foxl2* haploinsufficiency models.

Extended Data Table 3. 10xGenomics CellRanger QC metrix.

Extended Data Table 4. List of canonical marker genes used to annotate cell types in scRNA-seq

datasets.

Extended Data Table 5. Relative proportion of detected cell types from scRNA-seq datasets.

Proportion data from **a**, Aging, **b**, VCD and **c**, *Foxl2* haploinsufficiency models.

Extended Data Table 6. DESeq2 output from pseudobulk analysis. DESeq2 output from **a**, Aging, **b**,

VCD and **c**, *Foxl2* haploinsufficiency model datasets.

Extended Data Table 7. Significant GSEA enrichment terms (FDR < 10%) for GO Biological

Process and Reactome terms. GSEA enrichment terms for Biological Process from **a**, Aging, **b**, VCD

and **c**, *Foxl2* haploinsufficiency model datasets. GSEA enrichment terms for Reactome from **d**, Aging, **e**,

VCD and **f**, *Foxl2* haploinsufficiency model datasets.

Extended Data Table 8. List of decoupleR TF activity scores (FDR < 5%) for each cell type.

Extended Data Table 9. Primers used in the study. a, Primers used for genotyping *Foxl2*

haploinsufficiency model animals. **b**, Primers used for *Foxl2* expression quantification in *Foxl2*

haploinsufficiency model animals via RT-qPCR.

References

Davis, S. R. *et al.* Menopause. *Nat Rev Dis Primers* **1**, 15004, doi:10.1038/nrdp.2015.4 (2015).

Broekmans, F. J., Soules, M. R. & Fauser, B. C. Ovarian aging: mechanisms and clinical consequences. *Endocr Rev* **30**, 465-493, doi:10.1210/er.2009-0006 (2009).

Faddy, M. J. & Gosden, R. G. A model conforming the decline in follicle numbers to the age of menopause in women. *Hum Reprod* **11**, 1484-1486, doi:10.1093/oxfordjournals.humrep.a019422 (1996).

Ossewaarde, M. E. *et al.* Age at menopause, cause-specific mortality and total life expectancy. *Epidemiology* **16**, 556-562, doi:10.1097/01.ede.0000165392.35273.d4 (2005).

Hong, J. S. *et al.* Age at menopause and cause-specific mortality in South Korean women: Kangwha Cohort Study. *Maturitas* **56**, 411-419, doi:10.1016/j.maturitas.2006.11.004 (2007).

Shadyab, A. H. *et al.* Ages at menarche and menopause and reproductive lifespan as predictors of exceptional longevity in women: the Women's Health Initiative. *Menopause* **24**, 35-44, doi:10.1097/GME.0000000000000710 (2017).

Gubbels Bupp, M. R. Sex, the aging immune system, and chronic disease. *Cell Immunol* **294**, 102-110, doi:10.1016/j.cellimm.2015.02.002 (2015).

Santoro, N., Roeca, C., Peters, B. A. & Neal-Perry, G. The Menopause Transition: Signs, Symptoms, and Management Options. *J Clin Endocrinol Metab* **106**, 1-15, doi:10.1210/clinem/dgaa764 (2021).

Randolph, J. F., Jr. *et al.* Change in follicle-stimulating hormone and estradiol across the menopausal transition: effect of age at the final menstrual period. *J Clin Endocrinol Metab* **96**, 746-754, doi:10.1210/jc.2010-1746 (2011).

Kelsey, T. W., Wright, P., Nelson, S. M., Anderson, R. A. & Wallace, W. H. A validated model of serum anti-mullerian hormone from conception to menopause. *PLoS One* **6**, e22024, doi:10.1371/journal.pone.0022024 (2011).

Weiss, G., Skurnick, J. H., Goldsmith, L. T., Santoro, N. F. & Park, S. J. Menopause and hypothalamic-pituitary sensitivity to estrogen. *JAMA* **292**, 2991-2996, doi:10.1001/jama.292.24.2991 (2004).

Xiong, J. *et al.* FSH blockade improves cognition in mice with Alzheimer's disease. *Nature* **603**, 470-476, doi:10.1038/s41586-022-04463-0 (2022).

Mao, L., Wang, L., Bennett, S., Xu, J. & Zou, J. Effects of follicle-stimulating hormone on fat metabolism and cognitive impairment in women during menopause. *Front Physiol* **13**, 1043237, doi:10.3389/fphys.2022.1043237 (2022).

Van Kempen, T. A., Milner, T. A. & Waters, E. M. Accelerated ovarian failure: a novel, chemically induced animal model of menopause. *Brain Res* **1379**, 176-187, doi:10.1016/j.brainres.2010.12.064 (2011).

Blenck, C. L., Harvey, P. A., Reckelhoff, J. F. & Leinwand, L. A. The Importance of Biological Sex and Estrogen in Rodent Models of Cardiovascular Health and Disease. *Circ Res* **118**, 1294-1312, doi:10.1161/CIRCRESAHA.116.307509 (2016).

16 Laszczynska, M., Brodowska, A., Starczewski, A., Masiuk, M. & Brodowski, J.
Human postmenopausal ovary--hormonally inactive fibrous connective tissue or
more? *Histol Histopathol* **23**, 219-226, doi:10.14670/HH-23.219 (2008).
17 Grub, J., Suss, H., Willi, J. & Ehlert, U. Steroid Hormone Secretion Over the
Course of the Perimenopause: Findings From the Swiss Perimenopause Study.
Front Glob Womens Health **2**, 774308, doi:10.3389/fgwh.2021.774308 (2021).
18 Burger, H. G., Dudley, E. C., Robertson, D. M. & Dennerstein, L. Hormonal
changes in the menopause transition. *Recent Prog Horm Res* **57**, 257-275,
doi:10.1210/rp.57.1.257 (2002).
19 Mangoni, A. A. & Jackson, S. H. Age-related changes in pharmacokinetics and
pharmacodynamics: basic principles and practical applications. *Br J Clin
Pharmacol* **57**, 6-14, doi:10.1046/j.1365-2125.2003.02007.x (2004).
20 Conboy, I. M. & Rando, T. A. Heterochronic parabiosis for the study of the effects
of aging on stem cells and their niches. *Cell Cycle* **11**, 2260-2267,
doi:10.4161/cc.20437 (2012).
21 Chahal, H. S. & Drake, W. M. The endocrine system and ageing. *J Pathol* **211**,
173-180, doi:10.1002/path.2110 (2007).
22 van den Beld, A. W. *et al.* The physiology of endocrine systems with ageing.
Lancet Diabetes Endocrinol **6**, 647-658, doi:10.1016/S2213-8587(18)30026-3
(2018).
23 Rando, T. A. & Jones, D. L. Regeneration, Rejuvenation, and Replacement:
Turning Back the Clock on Tissue Aging. *Cold Spring Harb Perspect Biol* **13**,
doi:10.1101/cshperspect.a040907 (2021).
24 Villeda, S. A. *et al.* The ageing systemic milieu negatively regulates neurogenesis
and cognitive function. *Nature* **477**, 90-94, doi:10.1038/nature10357 (2011).
25 Uda, M. *et al.* Foxl2 disruption causes mouse ovarian failure by pervasive
blockage of follicle development. *Hum Mol Genet* **13**, 1171-1181,
doi:10.1093/hmg/ddh124 (2004).
26 Schmidt, D. *et al.* The murine winged-helix transcription factor Foxl2 is required
for granulosa cell differentiation and ovary maintenance. *Development* **131**, 933-
942, doi:10.1242/dev.00969 (2004).
27 Uhlenhaut, N. H. *et al.* Somatic sex reprogramming of adult ovaries to testes by
FOXL2 ablation. *Cell* **139**, 1130-1142, doi:10.1016/j.cell.2009.11.021 (2009).
28 Benayoun, B. A., Dipietromaria, A., Bazin, C. & Veitia, R. A. FOXL2: at the
crossroads of female sex determination and ovarian function. *Adv Exp Med Biol*
665, 207-226, doi:10.1007/978-1-4419-1599-3_16 (2009).
29 Gersak, K., Harris, S. E., Smale, W. J. & Shelling, A. N. A novel 30 bp deletion in
the FOXL2 gene in a phenotypically normal woman with primary amenorrhoea:
case report. *Hum Reprod* **19**, 2767-2770, doi:10.1093/humrep/deh496 (2004).
30 Harris, S. E. *et al.* Identification of novel mutations in FOXL2 associated with
premature ovarian failure. *Mol Hum Reprod* **8**, 729-733,
doi:10.1093/molehr/8.8.729 (2002).
31 Laissue, P. *et al.* Functional evidence implicating FOXL2 in non-syndromic
premature ovarian failure and in the regulation of the transcription factor OSR2. *J
Med Genet* **46**, 455-457, doi:10.1136/jmg.2008.065086 (2009).

32 Borman, S. M., Christian, P. J., Sipes, I. G. & Hoyer, P. B. Ovotoxicity in female
Fischer rats and B6 mice induced by low-dose exposure to three polycyclic
aromatic hydrocarbons: comparison through calculation of an ovotoxic index.
Toxicol Appl Pharmacol **167**, 191-198, doi:10.1006/taap.2000.9006 (2000).
33 Finch, C. E. The menopause and aging, a comparative perspective. *J Steroid*
Biochem Mol Biol **142**, 132-141, doi:10.1016/j.jsbmb.2013.03.010 (2014).
34 Gosden, R. G. & Faddy, M. J. Ovarian aging, follicular depletion, and
steroidogenesis. *Exp Gerontol* **29**, 265-274, doi:10.1016/0531-5565(94)90006-x
(1994).
35 Lliberos, C. *et al.* Evaluation of inflammation and follicle depletion during ovarian
ageing in mice. *Sci Rep* **11**, 278, doi:10.1038/s41598-020-79488-4 (2021).
36 Sokalska, A. & Valentin, L. Changes in ultrasound morphology of the uterus and
ovaries during the menopausal transition and early postmenopause: a 4-year
longitudinal study. *Ultrasound Obstet Gynecol* **31**, 210-217,
doi:10.1002/uog.5241 (2008).
37 Jiao, X. *et al.* Ovarian Reserve Markers in Premature Ovarian Insufficiency:
Within Different Clinical Stages and Different Etiologies. *Front Endocrinol*
(Lausanne) **12**, 601752, doi:10.3389/fendo.2021.601752 (2021).
38 Danforth, D. R. *et al.* Dimeric inhibin: a direct marker of ovarian aging. *Fertil Steril*
70, 119-123, doi:10.1016/s0015-0282(98)00127-7 (1998).
39 Kevenaar, M. E. *et al.* Serum anti-mullerian hormone levels reflect the size of the
primordial follicle pool in mice. *Endocrinology* **147**, 3228-3234,
doi:10.1210/en.2005-1588 (2006).
40 Park, M., Suh, D. S., Lee, K. & Bae, J. Positive cross talk between FOXL2 and
antimullerian hormone regulates ovarian reserve. *Fertil Steril* **102**, 847-855 e841,
doi:10.1016/j.fertnstert.2014.05.031 (2014).
41 Kim, M. *et al.* Estropausal gut microbiota transplant improves measures of
ovarian function in adult mice. *bioRxiv*, doi:10.1101/2024.05.03.592475 (2025).
42 Mehalko, K. *et al.* Lack of accelerated ovarian aging in a follicle-stimulating
hormone receptor haploinsufficiency model. *Transl Med Aging* **7**, 1-8,
doi:10.1016/j.tma.2023.01.001 (2023).
43 Crisponi, L. *et al.* The putative forkhead transcription factor FOXL2 is mutated in
blepharophimosis/ptosis/epicanthus inversus syndrome. *Nat Genet* **27**, 159-166,
doi:10.1038/84781 (2001).
44 Isola, J. V. V. *et al.* A single-cell atlas of the aging mouse ovary. *Nat Aging* **4**, 145-
162, doi:10.1038/s43587-023-00552-5 (2024).
45 Wang, X. *et al.* Transcriptomic signatures of mouse ovarian aging and
estropausal transition at single cell resolution. *bioRxiv*, 2024.2008.2012.607592,
doi:10.1101/2024.08.12.607592 (2024).
46 Jin, C. *et al.* Molecular and genetic insights into human ovarian aging from
single-nuclei multi-omics analyses. *Nat Aging* **5**, 275-290, doi:10.1038/s43587-
024-00762-5 (2025).
47 Ben Yaakov, T., Wasserman, T., Aknin, E. & Savir, Y. Single-cell analysis of the
aged ovarian immune system reveals a shift towards adaptive immunity and
attenuated cell function. *Elife* **12**, doi:10.7554/eLife.74915 (2023).

48   Squair, J. W., Skinnider, M. A., Gautier, M., Foster, L. J. & Courtine, G.
Prioritization of cell types responsive to biological perturbations in single-cell data
with Augur. *Nat Protoc* **16**, 3836-3873, doi:10.1038/s41596-021-00561-x (2021).
49   Xu, X. *et al.* Imaging and tracing the pattern of adult ovarian angiogenesis implies
a strategy against female reproductive aging. *Sci Adv* **8**, eabi8683,
doi:10.1126/sciadv.abi8683 (2022).
50   Lu, H. *et al.* Aging hallmarks of the primate ovary revealed by spatiotemporal
transcriptomics. *Protein Cell* **15**, 364-384, doi:10.1093/procel/pwad063 (2024).
51   Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. *Genome Biol* **15**, 550,
doi:10.1186/s13059-014-0550-8 (2014).
52   Gillespie, M. *et al.* The reactome pathway knowledgebase 2022. *Nucleic Acids
Res* **50**, D687-D692, doi:10.1093/nar/gkab1028 (2022).
53   Saul, D. *et al.* A new gene set identifies senescent cells and predicts
senescence-associated pathways across tissues. *Nat Commun* **13**, 4827,
doi:10.1038/s41467-022-32552-1 (2022).
54   Hajdarovic, K. H. *et al.* Single-cell analysis of the aging female mouse
hypothalamus. *Nat Aging* **2**, 662-678, doi:10.1038/s43587-022-00246-4 (2022).
55   Harris, S. E. *et al.* Age-related gene expression changes, and transcriptome wide
association study of physical and cognitive aging traits, in the Lothian Birth
Cohort 1936. *Aging (Albany NY)* **9**, 2489-2503, doi:10.18632/aging.101333
(2017).
56   He, J. *et al.* Identifying transposable element expression dynamics and
heterogeneity during development at the single-cell level with a processing
pipeline scTE. *Nat Commun* **12**, 1456, doi:10.1038/s41467-021-21808-x (2021).
57   Bravo, J. I., Nozownik, S., Danthi, P. S. & Benayoun, B. A. Transposable
elements, circular RNAs and mitochondrial transcription in age-related genomic
regulation. *Development* **147**, doi:10.1242/dev.175786 (2020).
58   De Cecco, M. *et al.* Transposable elements become active and mobile in the
genomes of aging mammalian somatic tissues. *Aging (Albany NY)* **5**, 867-883,
doi:10.18632/aging.100621 (2013).
59   Herr, L. M., Schaffer, E. D., Fuchs, K. F., Datta, A. & Brosh, R. M., Jr. Replication
stress as a driver of cellular senescence and aging. *Commun Biol* **7**, 616,
doi:10.1038/s42003-024-06263-w (2024).
60   Langfelder, P. & Horvath, S. WGCNA: an R package for weighted correlation
network analysis. *BMC Bioinformatics* **9**, 559, doi:10.1186/1471-2105-9-559
(2008).
61   Badia, I. M. P. *et al.* decoupleR: ensemble of computational methods to infer
biological activities from omics data. *Bioinform Adv* **2**, vbac016,
doi:10.1093/bioadv/vbac016 (2022).
62   Yang, Q. *et al.* NADase CD38 is a key determinant of ovarian aging. *Nat Aging* **4**,
110-128, doi:10.1038/s43587-023-00532-9 (2024).
63   Dipali, S. S. *et al.* Proteomic quantification of native and ECM-enriched mouse
ovaries reveals an age-dependent fibro-inflammatory signature. *Aging (Albany
NY)* **15**, 10821-10855, doi:10.18632/aging.205190 (2023).

64 Ouni, E. *et al.* A blueprint of the topology and mechanics of the human ovary for next-generation bioengineering and diagnosis. *Nat Commun* **12**, 5603, doi:10.1038/s41467-021-25934-4 (2021).

65 Wang, Y. *et al.* PBX1 Attenuates Hair Follicle-Derived Mesenchymal Stem Cell Senescence and Apoptosis by Alleviating Reactive Oxygen Species-Mediated DNA Damage Instead of Enhancing DNA Damage Repair. *Front Cell Dev Biol* **9**, 739868, doi:10.3389/fcell.2021.739868 (2021).

66 Lichtenauer, U. D. *et al.* Pre-B-cell transcription factor 1 and steroidogenic factor 1 synergistically regulate adrenocortical growth and steroidogenesis. *Endocrinology* **148**, 693-704, doi:10.1210/en.2006-0681 (2007).

67 Shen, L., Liu, J., Luo, A. & Wang, S. The stromal microenvironment and ovarian aging: mechanisms and therapeutic opportunities. *J Ovarian Res* **16**, 237, doi:10.1186/s13048-023-01300-4 (2023).

68 Hao, X., Reyes Palomares, A., Anastacio, A., Liu, K. & Rodriguez-Wallberg, K. A. Evidence of apoptosis as an early event leading to cyclophosphamide-induced primordial follicle depletion in a prepubertal mouse model. *Front Endocrinol (Lausanne)* **15**, 1322592, doi:10.3389/fendo.2024.1322592 (2024).

69 Ravelojaona, M. *et al.* Oncostatin M and STAT3 Signaling Pathways Support Human Trophoblast Differentiation by Inhibiting Inflammatory Stress in Response to IFNgamma and GM-CSF. *Cells* **13**, doi:10.3390/cells13030229 (2024).

70 Devchand, P. R., Ziouzenkova, O. & Plutzky, J. Oxidative stress and peroxisome proliferator-activated receptors: reversing the curse? *Circ Res* **95**, 1137-1139, doi:10.1161/01.RES.0000151331.69399.b2 (2004).

71 Schumann, K. *et al.* Functional CRISPR dissection of gene networks controlling human regulatory T cell identity. *Nat Immunol* **21**, 1456-1466, doi:10.1038/s41590-020-0784-4 (2020).

72 Klein-Hessling, S. *et al.* NFATc1 controls the cytotoxicity of CD8(+) T cells. *Nat Commun* **8**, 511, doi:10.1038/s41467-017-00612-6 (2017).

73 Kostel Bal, S. *et al.* Biallelic NFATC1 mutations cause an inborn error of immunity with impaired CD8+ T-cell function and perturbed glycolysis. *Blood* **142**, 827-845, doi:10.1182/blood.2022018303 (2023).

74 Buckley, M. T. *et al.* Cell-type-specific aging clocks to quantify aging and rejuvenation in neurogenic regions of the brain. *Nat Aging* **3**, 121-137, doi:10.1038/s43587-022-00335-4 (2023).

75 Peters, M. J. *et al.* The transcriptional landscape of age in human peripheral blood. *Nat Commun* **6**, 8570, doi:10.1038/ncomms9570 (2015).

76 Wang, S. *et al.* Single-Cell Transcriptomic Atlas of Primate Ovarian Aging. *Cell* **180**, 585-600 e519, doi:10.1016/j.cell.2020.01.009 (2020).

77 Bafor, E. E. *et al.* Aberrant CD8(+)T cells drive reproductive dysfunction in female mice with elevated IFN-gamma levels. *Front Immunol* **15**, 1368572, doi:10.3389/fimmu.2024.1368572 (2024).

78 Jiang, S. *et al.* Generic Diagramming Platform (GDP): a comprehensive database of high-quality biomedical graphics. *Nucleic Acids Res* **53**, D1670-D1676, doi:10.1093/nar/gkae973 (2025).

79 Komsta, L. *outliers: Tests for Outliers.*, <https://CRAN.R-project.org/package=outliers> (2022).

80 Therneau, T. *A Package for Survival Analysis in R*, <https://CRAN.R-
project.org/package=survival> (2022).
81 Ongaro, L. *et al.* Development of a Highly Sensitive ELISA for Measurement of
FSH in Serum, Plasma, and Whole Blood in Mice. *Endocrinology* **162**,
doi:10.1210/endocr/bqab014 (2021).
82 Kuhn, M. caret: Classification and Regression Training. (2022).
83 Breiman, L., Cutler, A., Liaw, A. & Wiener, M. *Breiman and Cutlers Random
Forests for Classification and Regression*,
<https://www.stat.berkeley.edu/~breiman/RandomForests/> (
84 Zheng, G. X. *et al.* Massively parallel digital transcriptional profiling of single cells.
Nat Commun **8**, 14049, doi:10.1038/ncomms14049 (2017).
85 Campbell, J., Yang, S., Wang, Z., Corbett, S. & Koga, Y. *celda: CEllular Latent
Dirichlet Allocation*, 2023).
86 *DoubletFinder: DoubletFinder is a suite of tools for identifying doublets in single-
cell RNA sequencing data*, 2022).
87 Bais, A. & Kostkam, D. scds: Computational Annotation of Doublets in Single Cell
RNA Sequencing Data. (2019).
88 Hao, Y. *et al.* Integrated analysis of multimodal single-cell data. *Cell* **184**, 3573-
3587 e3529, doi:10.1016/j.cell.2021.04.048 (2021).
89 Korsunsky, I. *et al.* Fast, sensitive and accurate integration of single-cell data with
Harmony. *Nat Methods* **16**, 1289-1296, doi:10.1038/s41592-019-0619-0 (2019).
90 Andreatta, M., Berenstein, A. J. & Carmona, S. J. scGate: marker-based
purification of cell types from heterogeneous single-cell RNA-seq datasets.
Bioinformatics **38**, 2642-2644, doi:10.1093/bioinformatics/btac141 (2022).
91 Aran, D. *et al.* Reference-based analysis of lung single-cell sequencing reveals a
transitional profibrotic macrophage. *Nat Immunol* **20**, 163-172,
doi:10.1038/s41590-018-0276-y (2019).
92 Guo, H. & Li, J. scSorter: assigning cells to known cell types according to marker
genes. *Genome Biol* **22**, 69, doi:10.1186/s13059-021-02281-7 (2021).
93 Nader, K. *et al.* ScType enables fast and accurate cell type identification from
spatial transcriptomics data. *Bioinformatics* **40**,
doi:10.1093/bioinformatics/btae426 (2024).
94 Heng, T. S., Painter, M. W. & Immunological Genome Project, C. The
Immunological Genome Project: networks of gene expression in immune cells.
Nat Immunol **9**, 1091-1094, doi:10.1038/ni1008-1091 (2008).
95 Morris, M. E. *et al.* A single-cell atlas of the cycling murine ovary. *Elife* **11**,
doi:10.7554/eLife.77239 (2022).
96 Miller, S. A. *et al.* LSD1 and Aberrant DNA Methylation Mediate Persistence of
Enteroendocrine Progenitors That Support BRAF-Mutant Colorectal Cancer.
Cancer Res **81**, 3791-3805, doi:10.1158/0008-5472.CAN-20-3562 (2021).
97 Crowell, H. L. *et al.* muscat detects subpopulation-specific state transitions from
multi-sample multi-condition single-cell transcriptomics data. *Nat Commun* **11**,
6077, doi:10.1038/s41467-020-19894-4 (2020).
98 Leek, J. *et al. sva: Surrogate Variable Analysis*.

99 Ritchie, M. E. *et al.* limma powers differential expression analyses for RNA-
sequencing and microarray studies. *Nucleic Acids Res* **43**, e47,
doi:10.1093/nar/gkv007 (2015).
100 Wu, T. *et al.* clusterProfiler 4.0: A universal enrichment tool for interpreting omics
data. *Innovation (Camb)* **2**, 100141, doi:10.1016/j.xinn.2021.100141 (2021).
101 Dolgalev, I. *msigdbr: MSigDB Gene Sets for Multiple Organisms in a Tidy Data*
Format, <https://CRAN.R-project.org/package=msigdbr> (2021).
102 Ouyang, J. F., Kamaraj, U. S., Cao, E. Y. & Rackham, O. J. L. ShinyCell: simple
and sharable visualization of single-cell gene expression data. *Bioinformatics* **37**,
3374-3376, doi:10.1093/bioinformatics/btab209 (2021).
1656

Fig. 1

a Experimental scheme

Aging
Young (4 months) *vs* Old (20 months)

VCD
CTL (Safflower oil) *vs* VCD

Foxl2 +/-
Foxl2+/+ *vs* *Foxl2*+/-

Ovarian health assessment
▪ Ovarian follicle count
▪ Serum AMH, FSH & INHBA concentrations
▪ Ovarian health index

Single-cell transcriptomic analysis

b Hematoxylin-eosin staining: Aging
c Hematoxylin-eosin staining: VCD
d Hematoxylin-eosin staining: *Foxl2* +/-

e Combined follicle count: Aging
f Combined follicle count: VCD
g Combined follicle count: *Foxl2* +/-

h Serum AMH levels: Aging
i Serum AMH levels: VCD
j Serum AMH levels: *Foxl2* +/-

k Serum FSH levels: Aging
l Serum FSH levels: VCD
m Serum FSH levels: *Foxl2* +/-

n Serum INHBA levels: Aging
o Serum INHBA levels: VCD
p Serum INHBA levels: *Foxl2* +/-

q Ovarian health index: Aging
r Ovarian health index: VCD
s Ovarian health index: *Foxl2* +/-

Fig. 2

a OvAge clock training scheme

C57BL/6J, NTac (3-20-months)
— Aging model: All
— VCD model: CTL
— *Foxl2 +/-* model: *Foxl2⁺ᐟ⁺*
— *Fshr +/-* model: *Fshr⁺ᐟ⁺*

→ AMH FSH INHBA
Serum ELISA

→ Training (n = 190)
Testing (n=62)

→ Random forest-based OvAge clock

b OvAge performance: Test dataset
Rho ~0.621
P-value ~7.4e-8
Predicted age (weeks) vs Chronological age (weeks)

- Aging (n=11) ◆ *Foxl2 +/-* (n=7)
- ▲ VCD (n=43) ■ *Fshr +/-* (n=1)

c Age prediction: VCD
Predicted age (weeks) vs Chronological age (weeks)

- ▲ CTL (n=43)
- ▲ VCD (n=110)

d Age prediction: *Foxl2 +/-*
Predicted age (weeks) vs Chronological age (weeks)

- ◆ *Foxl2⁺ᐟ⁺* (n=7)
- ◆ *Foxl2⁺ᐟ⁻* (n=46)

e VCD - 30d post-injection
Age acceleration (Predicted − Chronological)
~7.9e-3 ~7.9e-3 ~7.9e-3 ~7.9e-3
CTL/VCD 3m 6m 8m 10m
Age-at-injection

f VCD - 90d post-injection
~7.9e-3 ~7.9e-3 ~7.9e-3 ~7.9e-3
CTL/VCD 3m 6m 8m 10m
Age-at-injection

g VCD - 150d post-injection
~7.9e-3 ~7.9e-3 ~7.9e-3 ~0.151
CTL/VCD 3m 6m 8m 10m
Age-at-injection

h *Foxl2 +/-*
Age acceleration (Predicted − Chronological)
~0.846 ~0.011
Foxl2⁺ᐟ⁺ *Foxl2⁺ᐟ⁻* *Foxl2⁺ᐟ⁺* *Foxl2⁺ᐟ⁻*
Young Mid-age

▲◆ Model prediction
△◇ Out-of-bag prediction

Fig. 3

a scRNA-seq experimental scheme

4 ovaries per library → Single-cell suspension → scRNA-seq

b Sample collection scheme

Aging: 4m — 20m
VCD: 3m, 30d, 90d; 10m, 30d, 90d
Foxl2 +/-: 4m — 9m

CTL/VCD injection
Sample collection

c Cell type UMAP: Aging

Epithelial, SMC, Pericyte, Stroma, Theca, Granulosa, BEC, LEC

d Cell type UMAP: VCD

BEC, Granulosa, SMC, Theca, Pericyte, Stroma, LEC, Epithelial

e Cell type UMAP: *Foxl2 +/-*

LEC, BEC, Epithelial, Stroma, Pericyte, SMC, Granulosa, Theca

Non-immune cell types:
- Granulosa
- Theca
- Stroma
- SMC
- Pericyte
- BEC
- LEC
- Epithelial

Immune cell types:
- Neutrophil
- Monocyte
- Macrophage
- DC
- ILC
- NK
- NKT
- CD8+ NKT
- CD8+ T
- CD4+ T
- DNT
- DPT
- B

f Marker gene expression: Aging

g Marker gene expression: VCD

h Marker gene expression: *Foxl2 +/-*

Features: Cd19, Cd4, Cd8b1, Cd3e, Nkg7, Gzma, Klrb1c, Il2ra, Flt3, C1qa, Adgre1, S100a8, Upk1b, Prox1, Flt1, Rgs5, Notch3, Acta2, Col1a1, Pdgfra, Cyp11a1, Cyp17a1, Cyp19a1, Fshr, Amh

Cell types (x-axis): Granulosa, Theca, Stroma, SMC, Pericyte, BEC, LEC, Epithelial, Neutrophil, Monocyte, Macrophage, DC, ILC, NK, CD8+ NKT, CD8+ T, CD4+ T, DNT, DPT, B

% Expressed: 0, 25, 50, 75, 100
Ave. Expression: 2, 1, 0

i Representative RNAscope images: *Klrb1c*, *Cd3e*, *Cd8b1*, *Cd4*

DAPI *Klrb1c* *Cd3e* *Cd8b1* *Cd4*

Merged | DAPI | *Klrb1c* | *Cd3e* | *Cd8b1* | *Cd4*

CD8+ NKT

DPT

10μm

Fig. 4

a Cell type proportion analysis scheme

scRNA-seq datasets → Non-immune (*Ptprc⁻*) vs. Immune (*Ptprc⁺*) → Flow cytometry: CD45 expression

Cell subtypes → RNAscope: Marker gene expression

Cell type proportion analysis via scProportion — Validation

Cell type proportions from scRNA-seq: *Ptprc⁻* vs. *Ptprc⁺*

b Aging

Ptprc⁻
Ptprc⁺

Log$_2$FD

More abundant in:
● Young
◆ Old

c VCD

Ptprc⁻
Ptprc⁺

Log$_2$FD

More abundant in:
■ CTL   ● 3m / 30d   ■ 10m / 30d
■ VCD   ▲ 3m / 90d   ◆ 10m / 90d

d *Foxl2 +/-*

Ptprc⁻
Ptprc⁺

Log$_2$FD

More abundant in:
■ *Foxl2⁺/⁺*   ● Young
■ *Foxl2⁺/-*   ▲ Mid-age

Cell type proportions from flow cytometry: CD45⁺ cells

e Aging

Young
Old

~1.1e-5

CD45⁺ proportion (%)

f VCD

Age-at-injection: 3m
CTL
VCD

~3.2e-2

CD45⁺ proportion (%)

g *Foxl2 +/-*

Young: *Foxl2⁺/⁺*, *Foxl2⁺/-*   ~1.5e-2
Mid-age: *Foxl2⁺/⁺*, *Foxl2⁺/-*   ~4.6e-2

CD45⁺ proportion (%)

Cell type proportions from scRNA-seq: *Ptprc⁻* cell subtypes detected

h Aging

Granulosa
Theca
Stroma
SMC
Pericyte
BEC
LEC
Epithelial

Log$_2$FD

i VCD

Granulosa
Theca
Stroma
SMC
Pericyte
BEC
LEC
Epithelial

Log$_2$FD

j *Foxl2 +/-*

Granulosa
Theca
Stroma
SMC
Pericyte
BEC
LEC
Epithelial

Log$_2$FD

Cell type proportions from RNAscope: Granulosa and theca cells

k Aging

Young | Old

DAPI *Fshr* *Cyp11a1*

l VCD

Age-at-injection: 3m
CTL | VCD

DAPI *Fshr* *Cyp11a1*

m *Foxl2 +/-*

Young: *Foxl2⁺/⁺* | *Foxl2⁺/-*   Mid-age: *Foxl2⁺/⁺* | *Foxl2⁺/-*

DAPI *Fshr* *Cyp11a1*

n Aging

Granulosa
Young
Old
~1.6e-2
Fshr⁺ proportion (%)

Theca
Young
Old
~3.6e-2
Cyp11a1⁺ proportion (%)

o VCD

Granulosa
Age-at-injection: 3m
CTL
VCD
~1.6e-2
Fshr⁺ proportion (%)

Theca
Age-at-injection: 3m
CTL
VCD
~0.29
Cyp11a1⁺ proportion (%)

p *Foxl2 +/-*

Granulosa
Young: *Foxl2⁺/⁺*, *Foxl2⁺/-*   ~2.9e-2
Mid-age: *Foxl2⁺/⁺*, *Foxl2⁺/-*   ~2.9e-2
Fshr⁺ proportion (%)

Theca
Young: *Foxl2⁺/⁺*, *Foxl2⁺/-*   1
Mid-age: *Foxl2⁺/⁺*, *Foxl2⁺/-*   1
Cyp11a1⁺ proportion (%)

Fig. 5

a Global gene expression analysis comparisons: Augur

Aging model
Young *vs* Old

VCD model
CTL *vs* VCD
- Age-at-injection: 3m ⌐30d ⌐90d
- Age-at-injection: 10m ⌐30d ⌐90d

Foxl2 +/- model
Foxl2+/+ *vs* *Foxl2+/-*
- Young
- Mid-age

b UMAP of Augur AUC scores: Aging

c Augur AUC quantified by cell type: Aging

Cell type	AUC
Granulosa	0.790
CD4+ T	0.724
NK	0.722
B	0.710
Macrophage	0.706
DC	0.666
CD8+ NKT	0.634
BEC	0.629
LEC	0.621
Stroma	0.617
Theca	0.607
NKT	0.601
SMC	0.599
Epithelial	0.586
DNT	0.530

d UMAP of Augur AUC scores: VCD - 90d post-injection
Age-at-injection: 3m   Age-at-injection: 10m

e Augur AUC quantified by cell type: VCD - 90d post-injection
Ptprc- cells   *Ptprc+* cells

f UMAP of Augur AUC scores: *Foxl2 +/-*
Young   Mid-age

g Augur AUC quantified by cell type: *Foxl2 +/-*
Ptprc- cells   *Ptprc+* cells

Fig. 6

a Pseudobulk data analysis pipeline scheme

scRNA-seq datasets → Subset by cell type → Pseudobulk data via muscat → DEG analysis via DESeq2 / GO analysis via clusterProfiler

b DEGs: Aging (FDR < 0.05)

Log_2FC

Granulosa (1194), Theca (31), Stroma (458), SMC (22), BEC (182), LEC (49), Epithelial (50), NK (23), DNT (13), B (8)

c DEGs: VCD (FDR < 0.05)

Log_2FC

Granulosa (3664), Theca (5385), Stroma (3418), SMC (NA), BEC (1727), LEC (173), Epithelial (2975), NK (NA), DNT (7), B (NA)

d DEGs: *Foxl2* +/- (FDR < 0.05)

Log_2FC

Granulosa (5478), Theca (134), Stroma (1675), SMC (NA), BEC (432), LEC (NA), Epithelial (6), NK (33), DNT (84), B (4)

e Frequent GSEA term enrichment: GO Biological Process (FDR < 0.05)

Aging | VCD | *Foxl2* +/-

Rows:
- Respiratory electron transport chain
- Proton transmembrane transport
- Oxidative phosphorylation
- Energy derivation by oxidation of organic compounds
- Electron transport chain
- Cellular respiration
- ATP synthesis coupled electron transport
- ATP metabolic process
- Antigen processing and presentation of endogenous peptide antigen
- Antigen processing and presentation of endogenous antigen
- Aerobic respiration

Columns: Granulosa, Theca, Stroma, SMC, BEC, LEC, Epithelial, NK, DNT, B

NES: 3, 0, -3
$-Log_{10}(FDR)$: 2, 3, 4, 5

f Frequent GSEA term enrichment: Reactome (FDR < 0.1)

Aging | VCD | *Foxl2* +/-

Rows:
- The citric acid tca cycle and respiratory electron transport
- Respiratory electron transport ATP synthesis by chemiosmotic coupling and heat production by uncoupling proteins
- Respiratory electron transport

Columns: Granulosa, Theca, Stroma, SMC, BEC, LEC, Epithelial, NK, DNT, B

NES: 3, 0, -3
$-Log_{10}(FDR)$: 1.5, 2.0, 2.5

Fig. 7

a WGCNA and TF inference analysis scheme

b GSEA enrichment: WGCNA modules

c ORA enrichment: WGCNA modules per cell type

d Intersection size of decoupleR-inferred TF activity

e Common decoupleR-inferred TF activity for each cell type

Fig. 8

a Gene 1 Gene 2 ... Gene X

Cell 1
Cell 2
...
Cell Y

24503 features
x 7135 cells

→

Predicted age

Chronological age

Train & Test
Lasso regression

→

Expression analysis by age

Over-representation analysis

scRNA-seq transcriptome data:
Granulosa cells

Lasso feature analysis

b Granulosa clock performance: Test set

Rho ~0.76
P-value ~3.9e-293

Predicted age (weeks)

Chronological age (weeks)

• Aging
▲ VCD: CTL
◆ Foxl2 +/-: Foxl2⁺/⁺

c Age prediction: VCD

Predicted age (weeks)

Chronological age (weeks, + offset)

▲ CTL
▲ VCD

d Age prediction: Foxl2 +/-

Predicted age (weeks)

Chronological age (weeks, + offset)

◆ Foxl2⁺/⁺
◆ Foxl2⁺/⁻

e Age acceleration: VCD

~2.6e-51   ~8.7e-42   ~3.4e-23   ~5.4e-9

Age acceleration

CTL VCD   CTL VCD   CTL VCD   CTL VCD
30d   90d   30d   90d
3m   10m

Groups:

f Age acceleration: Foxl2 +/-

~1.7e-36   ~0.11

Age acceleration

Foxl2⁺/⁺ Foxl2⁺/⁻   Foxl2⁺/⁺ Foxl2⁺/⁻
Young   Mid-age

g Expression of lasso features by age (n=907 genes)

Row z score
2
1
0
-1
-2

13 15 17 20 29 39 48 57 76 87
Chronological age (weeks)

h ORA enrichment of top 500 features: GO ALL

Blood circulation
Leukocyte migration
Cell chemotaxis
Vascular process in circulatory system
Regulation of body fluid levels
Regulation of tube diameter
Blood vessel diameter maintenance
Regulation of tube size
Myeloid leukocyte migration
Leukocyte chemotaxis
Blood coagulation
Hemostasis
Coagulation
Interferon−gamma production
Regulation of interferon−gamma production
Interleukin−2 production
Smooth muscle cell differentiation
Positive regulation of antigen processing and presentation
Glutamate receptor clustering
CD4−positive, alpha−beta T cell proliferation

Count
● 10
● 20

FDR
0.10
0.05
0.00