Divergent effects of a Treg-selective IL-2 mutein on Influenza specific T cell responses

Joseph R Albe[*,†], Anita Chaudhary[#], Asheema Khanna[#], Kristen N Weinstein[*,†], Steven F. Ziegler[*,†],Vandana Kalia[§] ,Surojit Sarkar[#,¶], Daniel J Campbell[*,†].

[*]Center for Fundamental Immunology, Benaroya Research Institute, Seattle, WA 98101

[†]Department of Immunology, University of Washington School of Medicine, Seattle, WA 98195

[#ᵛ]Ben Towne Center for Childhood Cancer Research, Seattle Children's Research Institute, Seattle, WA, 98101, USA;

[§]Department of Pathology, University of Washington School of Medicine, Seattle, WA, 98195, USA; Department of Pediatrics, Division of Hematology and Oncology, University of Washington, Seattle, WA, 98195, USA.

[¶]Ben Towne Center for Childhood Cancer Research, Seattle Children's Research Institute, Seattle, WA, 98101, USA; Department of Pediatrics, Division of Hematology and Oncology, University of Washington, Seattle, WA, 98195, USA.

[*]Corresponding Author: Daniel J. Campbell, Benaroya Research Institute, 1201 Ninth Avenue, Seattle, WA 98101-2795. Phone: (206) 287-10565, email address: campbell@benaroyaresearch.org

Running Title: A Treg-selective IL-2 mutein alters flu-specific T cell responses

Abstract

Enhancing regulatory T cell (Treg) function offers a compelling therapeutic strategy for autoimmune disease. Engineered IL-2 muteins selectively expand functional Tregs with minimal impact on other immune cells, but their potential to compromise antiviral immunity remains largely unexplored. Here, we used a murine model of Influenza A virus (Flu) infection to determine how IL-2 mutein shapes T cell responses to respiratory virus infection. IL-2 mutein administration prior to infection suppressed Flu-specific (Flu-sp) CD8 T cell responses and altered their localization and phenotype within the lungs, without affecting bystander CD8 T cells. This suppression correlated with reduced antigen presentation molecule expression on conventional dendritic cells (cDCs) early after infection but did not impact Flu-sp CD8 T cell priming. In contrast, administering IL-2 mutein during infection exacerbated disease and drove CD25-dependent expansion of Flu-sp CD8 T cells. Despite these opposing effects on effector responses, Fc.Mut24-treated mice generated robust antibody responses and protective T cell memory which were maintained for at least 170 days. These findings reveal that Fc.Mut24 has temporally distinct effects on antiviral immunity, dampening early effector responses when given before infection, but enhancing effector expansion and disease severity when delivered during infection. Our results provide critical context for the therapeutic application of IL-2 muteins and highlight the importance of treatment timing in balancing immune modulation with protective immunity.

Keywords: IL-2 Mutein, Regulatory T cells, Influenza-specific CD8 T cells, Respiratory Infection

Introduction

Increasing regulatory T cell (Treg) abundance and function is a promising strategy for treating autoimmune diseases, as Tregs suppress inflammatory responses that drive disease pathology (1, 2). This can be accomplished by either expanding and re-infusing Tregs in various cell therapy approaches, or by boosting the abundance and/or function of endogenous Tregs. In the second category, one approach involves targeting the interleukin-2 receptor (IL-2R), which is critical for Treg development and function (3). The IL-2R exists in two distinct forms - a high-affinity form (composed of CD25, CD122, and CD132) is constitutively expressed by Tregs and activated CD4 and CD8 conventional T cells (Tconv), whereas the low-affinity IL-2R (composed of CD122 and CD132), is primarily found on resting naïve and memory Tconv and is highly expressed by NK cells. This differential expression of IL-2R forms helps support Treg homeostasis and function by allowing them to effectively compete for limiting amounts of IL-2 thereby preventing excessive immune activation (4). However, when IL-2 is produced in excess it can act on Tconv to support their differentiation into effector and memory populations (5). This receptor expression pattern also provides the rationale for low-dose IL-2 therapy, which aims to preferentially expand Tregs in autoimmune disease (6).

An emerging approach to more effectively target Treg cells for treatment of autoimmunity is to use engineered IL-2 mutant proteins ('muteins'), which are IL-2 variants designed to enhance Treg selectivity (2, 7). Several Treg-biased IL-2 muteins have been developed, including the Fc-fused IL-2 mutein from Merck (MK-6194) and

pegylated IL-2 from Nektar (Rezpegaldesleukin), each of which show enhanced Treg-
selectivity and potent Treg expansion in pre-clinical models or early clinical trials (8, 9).
Data from Rezpegaldesleukin trials have demonstrated efficacy in improving atopic
dermatitis in Phase 1b (9), showing the potential of this therapeutic approach. We
developed an Fc-fused murine IL-2 mutein (known as Fc.Mut24) with two amino acid
substitutions that decreased CD122 binding and thus increased dependence on CD25
to induce functional signaling (10, 11). Although Fc.Mut24 is a weaker IL-2R agonist *in*
vitro than an Fc-fused version of wild-type IL-2 (Fc.WT), it is highly Treg-selective and
more effectively expands Tregs *in vivo*. In addition to increasing Treg abundance,
Fc.Mut24 also enhances suppressive functions of Tregs, including CTLA-4-mediated
transendocytosis of the co-stimulatory ligands CD80 and CD86 (12). Importantly, unlike
Fc.WT, Fc.Mut24 can be administered at high doses while maintaining Treg selectivity
and effectively prevented disease progression in the non-obese diabetic (NOD) mouse
model of Type 1 Diabetes (10). Thus, Fc.Mut24 provides a novel tool for pre-clinical
mechanistic studies of IL-2 mutein therapy.

Given their ability to potently expand Tregs and broadly suppress the immune system,
IL-2 muteins represent a promising new class of autoimmune therapies. However,
immunosuppressive treatments—including corticosteroids, IL-6 inhibitors, co-stimulation
inhibitors, tumor necrosis factor (TNF)-α inhibitors—also impair immune responses to
infections and reduce vaccine efficacy (13-17). While these therapies target specific
inflammatory pathways, Tregs use multiple immunosuppressive mechanisms to
dampen immune responses (18), and therefore therapies that augment Treg abundance
and function may be even more detrimental for responses to infection. Respiratory virus
infections, such as RSV, Influenza and SARS-CoV-2, are particularly relevant in this
context, as several immunosuppressive therapies increase the risk of severe infection
(19, 20), and Tregs play a role in dampening immune responses and promoting tissue
repair during respiratory virus infections (21, 22), raising the possibility that IL-2 mutein
therapy could increase the risk of poor outcomes. Influenza, in particular, remains a
major global health burden causing significant morbidity and mortality each year. Given
these risks, it is important to determine whether IL-2 muteins compromise antiviral
immunity. In addition, IL-2-based therapies could also promote inflammation by
activating T cells that upregulate CD25 and express the high-affinity IL-2R. Therefore,
administering IL-2 mutein during infection, when T cells are already activated and have
upregulated the high-affinity IL-2 receptor, could enhance signaling and drive excessive
T cell activation, proliferation, and immunopathology. Thus how the various activities of
IL-2 mutein combine to impact the outcome of respiratory virus infection is complex, and
may depend on the timing of IL-2 mutein administration.

We used a murine model of Influenza A virus (Flu) infection to investigate how Fc.Mut24
influences antiviral T cell responses during respiratory infection. Administering Fc.Mut24
before infection suppressed Flu-specific (Flu-sp) CD8 T cells at 9 days post-infection
(dpi) and altered their localization and phenotype within the lung. This suppression was
specific to Flu-sp CD8 T cells, as bystander cells remained unaffected. Although
Fc.Mut24 reduced antigen presentation molecule expression on conventional dendritic
cells (cDCs) at 3dpi, the number of Flu-sp CD8 T cells was unchanged at 4dpi. In

contrast, administering Fc.Mut24 during infection worsened disease severity and drove
an expansion of Flu-sp CD8 T cells that was dependent on cell-intrinsic CD25
expression. Despite the divergent effects of Fc.Mut24 treatment before or during
infection, Fc.Mut24-treated mice generated protective memory that persisted through at
least 170dpi. These findings demonstrate that the immunologic context in which IL-2
mutein therapy is administered critically shapes antiviral T cell responses, highlighting
the need to evaluate how immune-modulating treatments for autoimmunity may
influence host defense.

Results

Fc.Mut24 treatment before infection suppresses Flu-specific CD8 T cell responses
In patients receiving IL-2 muteins for autoimmunity, administration of mutein and
expansion of Tregs just prior to respiratory virus infection could dampen the immune
response, worsen outcomes, and compromise development of durable memory against
re-infection. To model this possibility, we gave 10□µg of Fc.Mut24 to female C57BL/6
(B6) mice one day before intranasal infection with 2000 plaque-forming units (PFU) of
Influenza A virus (strain A/HK-x31(H3N2)) (X31). The relative timing of Fc.Mut24
treatment and infection aimed to maximize Treg expansion during the T cell priming
phase to determine how this impacted the Flu-sp immune response. We assessed T
cell responses in the lungs and spleen at 9dpi, corresponding to the peak of the effector
T response in our model.

Consistent with its ability to potently expand Tregs, Fc.Mut24-treated mice exhibited a
higher frequency and total number of Tregs in both lungs and spleen compared to
vehicle controls (Figure 1A and Supplemental Fig. 1A). Flu-sp CD8$^+$ T cells play a
critical role in viral clearance, memory formation, and cross-protection against
heterosubtypic influenza strains, and we therefore focused our analyses on the impact
of Fc.Mut24 treatment on this population, using an MHC class I tetramer recognizing the
immunodominant nucleoprotein antigen (NP)$_{366-374}$ to identify Flu-sp cells. At 9 dpi
Fc.Mut24-treated mice had a significant reduction in both the frequency and total
number of Flu-sp CD8$^+$ T cells in the lungs (Figure 1B) and spleen (Supplemental
Figure 1B) compared to vehicle-treated controls. The lung is a complex organ that is
highly vascularized, and T cells specific for respiratory pathogens can be found in both
the lung parenchyma and patrolling the lung vasculature (23). Therefore, we used
intravenous (IV) CD45 antibody labeling to distinguish Flu-sp T cells found in the lung
vasculature (IV$^+$) and lung parenchymal (IV$^-$) Flu-sp CD8$^+$ T cells. Fc.Mut24-treated
mice had a significant enrichment of Flu-sp CD8$^+$ T cells within the lung parenchyma
compared to vehicle-treated controls (Figure 1C). However, despite this skewed
distribution in the lung parenchyma, the total number of Flu-sp CD8$^+$ T cells remained
lower in both the lung vasculature (~7-fold reduction) and parenchyma (~3-fold
reduction) compared to control animals (Figure 1C).

To further investigate the altered differentiation and migration of Flu-sp CD8$^+$ T cells, we
analyzed markers associated with tissue residency and vascular retention. Consistent
with IV labeling results, the frequency of Flu-sp CD8$^+$ T cells expressing KLRG1 and
CX3CR1 (molecules linked to retention in the lung vasculature (24) was reduced in
Fc.Mut24-treated mice, whereas expression of the tissue-residency markers CD103 and

CD69 was increased (Figure 1D). This did not simply reflect their altered distribution, as
Fc.Mut24-treated mice had higher expression of CD69 and CD103 compared with
control mice even within the lung parenchyma fraction (Supplemental Figure 1C), and
similarly showed significantly reduced expression of CX3CR1 and KLRG1 in the blood-
exposed fraction (Supplemental Figure 1D) compared to the parenchymal compartment.
To further determine how Fc.Mut24 treatment altered the functional differentiation of
Flu-sp CD8$^+$ T cells, we examined expression of transcription factors and cytokine
receptors that promote effector and memory cell differentiation (Supplemental Figure
2A). Flu-sp CD8$^+$ T cells from Fc.Mut24-treated mice showed a slight reduction in
expression of the transcription factor EOMES, but no difference in expression of TCF1
or T-bet. However, consistent with increased memory potential, Fc.Mut24 treatment
increased expression of the IL-7R component CD127 (25).

Despite the reduced magnitude and altered distribution and phenotype of the Flu-sp
CD8$^+$ T cell response, Fc.Mut24 treatment did not affect disease severity, as measured
by weight loss (Figure 1E), viral RNA load (Figure 1F), or NP-specific IgG antibody titers
(Supplemental Figure 1E) compared to vehicle controls. Notably, while Fc.Mut24
treatment did not alter the magnitude of the Flu-sp CD4$^+$ conventional T cell response
(Supplemental Figure 1F), there was a decrease in FoxP3$^-$ CD4$^+$ T cells in the lungs,
suggesting a broader immunomodulatory effect (Supplemental Figure 1G). Further we
observed no changes in the interferon gamma or IL-2 production by Flu-sp CD8 T cells
that were stimulated *ex vivo* between vehicle treated and Fc.Mut24, suggesting there
was no direct impact on functionality of the resulting response (Supplemental Figure 1H
and I). Thus, Fc.Mut24 treatment resulted in sustained Treg expansion, and decreased
Flu-sp CD8$^+$ T cell abundance associated with altered localization and phenotype, with
a larger fraction of cells localized to the tissue parenchyma, potentially positioning them
for effective viral clearance. This is supported by the lack of change in disease severity
or viral loads despite the overall reduction in Flu-sp CD8$^+$ T cells.

Fc.Mut24 Treatment Selectively Limits Flu-Specific Responses Without Suppressing
Bystander CD8 T Cells
In addition to reducing the abundance and localization of NP-specific CD8$^+$ T cells,
Fc.Mut24 treatment also reduced the number of tetramer-negative CD44^{hi} (CD44^{hi})
CD8$^+$ T cells in the lungs at 9 dpi (Figure 2A), and these tetramer-negative cells also
tended to localize more toward the lung parenchyma, although this trend did not reach
statistical significance (Supplemental Figure 3A). These cells also showed a similar shift
in expression of markers of T cell migration and functional differentiation as the NP-
specific CD8$^+$ T cell population (Supplemental Figure 3B). Ki-67 expression was also
reduced in CD44^{hi} cells from Fc.Mut24-treated mice, suggesting decreased proliferation
of these antigen-experienced cells (Supplemental Figure 3B). These tetramer-negative
cells could represent Flu-sp cells recognizing epitopes other than the (NP)$_{366-374}$
epitope, or alternatively could be bystander activated CD8$^+$ T cells that also infiltrate the
lungs during influenza infection (26, 27). To determine if Fc.Mut24 altered this
bystander response, we infected mice intraperitoneally with 2000 PFU of Vesicular
Stomatitis virus that expresses the SIINFEKL epitope from ovalbumin (VSV-OVA) and
allowed 28 days for resolution. This generated SIINFEKL-specific CD8$^+$ T cells,

providing a bystander population that could be identified and tracked via staining with a
K^b/SIINFEKL MHCI tetramer. We then treated the mice with Fc.Mut24 or vehicle and,
one day later, challenged half the mice intranasally with X31 or mock infection, creating
four experimental groups for tracking the SIINFEKL-sp CD8$^+$ T cell response. In
contrast with the response of Flu-sp CD8$^+$ T cells, we observed no significant
differences in the number of SIINFEKL-sp CD8$^+$ T cells in the lungs across all groups
(Figure 2B). Although the frequency of SIINFEKL-sp was reduced in the VSV + Flu +
vehicle group, this likely reflects the influx of Flu-sp CD8$^+$ T cells (Figure 2B). Moreover,
SIINFEKL-sp CD8$^+$ T cells showed no differences in localization between the
parenchyma and blood-exposed compartments across all groups (Supplemental Figure
3C). We also found no changes in the expression of selected markers that were altered
on CD44^{hi} CD8$^+$ T cells (TCF, CD127, CD69, Ki-67, KLRG1, or CX3CR1) on SIINFEKL -
sp CD8$^+$ T cells (Supplemental Figure 3D). Together, these results indicate that
Fc.Mut24 does not suppress the SIINFEKL-sp response, suggesting that the observed
suppression in the CD44^{hi} population likely reflects effects on Flu-sp cells that were not
directly tracked, rather than non-specific bystander responses.

Fc.Mut24 treatment prior to infection reduces cDC antigen presentation capacity with
minimal impact on CD8 T cell proliferation at 4 days post-Influenza infection
Given the significant reduction in Flu-sp CD8$^+$ T cells during the effector phase of the
response, we wanted to determine if this was associated with changes in antigen
presentation capacity of conventional dendritic cells (cDCs), which prime Flu-sp CD8$^+$ T
cell responses in the draining mediastinal lymph nodes (medLN) and spleen following
Flu infection in mice (28). Indeed, by 3dpi, both cDC1 and cDC2 subsets strongly
upregulated key antigen presentation markers, including CD80, CD86, and MHC I
(Supplemental Figure 4A and B). We have previously shown that Tregs expanded
through Fc.Mut24 upregulate CTLA-4 which targets co-stimulatory molecules CD80 and
CD86 on cDCs, and that IL-10 production from expanded Tregs can modulate surface
MHC expression (12). Therefore, to further assess how Fc.Mut24 affects Treg
expansion and cDC activation in the context of Flu infection, we analyzed lung, medLN
and spleen at 3 dpi. Fc.Mut24-treated mice showed a marked increase in FoxP3$^+$ Tregs
in both tissue sites (Figure 3A), which exhibited an effector phenotype defined by high
expression of CTLA-4 and CD25 (Supplemental Figure 4C). Consistent with our
previous results in LPS-treated mice and in NOD mice, the fraction of activated
CD80^{hi}CD86^{hi} cDC1 and cDC2 subsets in the lungs and spleen were significantly
reduced in Fc.Mut24-treated mice at 3 dpi, whereas only cDC2 cells were significantly
impacted in the medLN, (Figure 3B).

Decreased antigen presentation capacity by cDCs could impair Flu-sp CD8$^+$ T cell
priming, resulting in the decreased expansion and infiltration of Flu-sp CD8$^+$ T cells in
the lungs we observed in Fc.Mut24-treated mice. To test this, we adoptively transferred
congenically marked OT-I CD8$^+$ T cells into Fc.Mut24- or vehicle-treated mice, then
infected them with a recombinant X31 expressing the SIINFEKL peptide (X31-
SIINFEKL). At 4 dpi, we measured OT-I expansion based on cell counts in the lung,
medLN, and spleen. Surprisingly, although the number of OT-I cells trended lower in
both the lungs (Figure 3C), spleen (Supplemental Figure 4D), Fc.Mut24-treated mice

showed no significant reduction in OT-I expansion in any tissue, including the medLN compared to controls at 4dpi, and there were no significant differences in expression of the activation markers CD69, CD25, and CD44 in the lung or spleen (Supplemental Figure 4E). Thus, despite reducing co-stimulatory ligand expression by cDC2, pre-treatment with Fc.Mut24 had only minimal effects on the initial activation and tissue localization of virus-specific CD8$^+$ T cells.

In addition to potentially impacting their initial priming, Fc.Mut24-mediated Treg expansion may suppress the migration and activation of Flu-sp CD8$^+$ T cells within the lungs themselves. To test this, we activated OT-I CD8$^+$ T cells *in vitro* with anti-CD3/CD28 Dynabeads for 48 hours to mimic T cell priming and then transferred them into recipient mice that had been pre-treated with vehicle or Fc.Mut24 and infected with X31-SIINFEKL three days earlier and sacrificed the mice for analysis 6 days later (9dpi). Although the frequency and number of transferred cells were reduced in the spleens of Fc.Mut24-treated mice, we observed no reduction in the number of OT-I cells in the lungs (Figure 3D, Supplemental Figure 4F). Additionally, rather than being suppressed the transferred OT-I cells showed increased expression of activation markers in the lungs but not the spleen (Supplemental Figure 4G). Together, these results indicate that while Fc.Mut24-driven Treg expansion dampens cDC activation and reduces the expansion and differentiation of Flu-sp cells, it has limited impact on their initial priming, and may even potentiate reactivation of virus-specific CD8$^+$ T cells in the lungs.

Fc.Mut24 administration during infection enhances Flu-specific CD8 T cell responses
A second important clinical scenario to model for how IL-2 mutein therapy may impact anti-viral immune responses is a patient receiving IL-2 mutein after initiation of infection, which would likely impact the Flu-sp T cell response differently than pre-infection treatment. These effects could include 'off-target' stimulation of CD25$^+$ responding T cells leading to increased proliferation and effector cell differentiation that could enhance viral clearance but exacerbate detrimental immunopathology. However, Tregs also help coordinate tissue repair responses in the lungs during Flu infection, and this may help offset these inflammatory effects. Therefore, to model the effects of post-infection treatment, we administered Fc.Mut24 or vehicle to mice at 4dpi with X31 and analyzed immune responses at 9 dpi (five days after mutein treatment). Fc.Mut24 treatment at 4dpi robustly increased both the number and frequency of Tregs in the lung and spleen (Supplemental Figure 5A), confirming that it retained its ability to expand Treg even in highly inflammatory conditions as previously reported (10). However, rather than dampening the Flu-sp CD8$^+$ T cell response, Fc.Mut24 given at 4dpi did not significantly change the frequency of Flu-sp CD8$^+$ T cells, and increased their absolute number in the lung (Figure 4A) and spleen (Supplemental Figure 5B) at 9dpi. In contrast to results from mice pre-treated with Fc.Mut24, we observed a trend toward increased Flu-sp CD8 T cell frequency in the lung vasculature in mice given Fc.Mut24-after infection, and the absolute number of cells in this compartment was significantly increased whereas cell number in the lung parenchyma was not changed (Figure 4B). Fc.Mut24-treatment after infection did not alter expression of KLRG1, EOMES T-bet or TCF1, however similar to mice receiving Fc.Mut24 before infection, expression of CX3CR1 was reduced and CD127 expression was dramatically increased in Flu-sp

CD8 T cells from the lungs of mice treated with Fc.Mut24 at 4dpi (Supplemental Figure
6A).

Unlike mice given Fc.Mut24 before Flu infection, mice treated at 4dpi lost more weight
and recovered more slowly than vehicle-treated controls (Figure 4C). This was not
associated with increased viral RNA in the lungs (Figure 4D) and thus is likely due to
exacerbated immunopathology (Figure 4E). Importantly, this was not due to heightened
functionality of Flu-specific CD8$^+$ T cells, as they did not show increased IFN-γ or IL-2
production following ex vivo peptide stimulation (Supplemental Figure 5E and F). As we
observed with Flu-sp CD8+ T cells, Flu-sp CD4+ T cell responses were also increased
in mice given Fc.Mut24 at 4dpi (Supplemental Figure 5C). However, despite the ability
of IL-2 to inhibit the differentiation of Tfh cells and subsequent antibody responses
during infection (29, 30), NP-specific IgG titers were not altered by Fc.Mut24 treatment
(Supplemental Figure 5D).

Expansion of Flu-specific CD8 T cells by Fc.Mut24 requires CD25 expression
Fc.Mut24 requires CD25 to initiate IL-2R signaling, and activated STAT5 downstream of
this pathway can further amplify CD25 expression through a positive feedback loop
(31). We therefore hypothesized that in mice given Fc.Mut24 at 4dpi, CD25 expression
on primed Flu-sp CD8$^+$ T cells enables direct stimulation by Fc.Mut24, leading to
enhanced expansion and altered differentiation and localization. Indeed, Flu-sp CD8$^+$ T
cells expressed higher levels of CD25 in 4dpi Fc.Mut24-treated mice, and thus
Fc.Mut24 may directly stimulate responding CD8$^+$ T cells and promote their excessive
proliferation (Figure 5A). To directly test this hypothesis, we used CRISPR/CAS9 RNPs
to delete the *Il2ra* gene (which encodes CD25) in naïve OT-I T cells (32), and
successful targeting was confirmed by assessing CD25 upregulation in targeted cells
after *in vitro* stimulation with anti-CD3/anti-CD28 (Supplemental Figure 7A). We
transferred 5 × 10□ cells to recipient CD45.2+ animals prior to infection with X31-
SIINFEKL, and infected animals were treated with either vehicle or Fc.Mut24 at 4dpi.
Indeed, loss of CD25 significantly impaired the Fc.Mut24-mediated hyper-expansion of
OT-I cells following X31-SIINFEKL infection, whereas there was no significant
difference in expansion of control and *Il2ra*-targeted cells in vehicle-treated mice (Figure
5B). We also analyzed the response of endogenous host-derived SIINFEKL-sp T cells
using K^b/SIINFEKL tetramers. Similar to what we observed for CD8$^+$ T cells specific for
the NP$_{366-374}$ Flu epitope, Fc.Mut24 treatment led to an increased expansion of the
endogenous SIINFEKL-sp cells compared to mice that received vehicle (Figure 5C),
and thus the changes in expansion we observed are not epitope-specific, but rather
reflect direct stimulation of CD25-expressing CD8 T cells by Fc.Mut24 during infection,
which likely contributes to the increased presence Flu-sp CD8$^+$ T cells and the elevated
immune-pathology associated with Fc.Mut24 administration post-infection.

Protective Memory formation and durability are maintained with Fc.Mut24 treatment
Mice treated with Fc.Mut24 before Flu infection showed reduced CD8$^+$ T cell responses,
which could impair memory formation and durability. Conversely, mice treated at 4dpi
exhibited enhanced proliferation and effector phenotypes, which might also compromise
memory development. However, regardless of treatment time, Flu-sp CD8$^+$ T cells from

Fc.Mut24-treated mice at 9dpi expressed elevated levels of the memory precursor
marker CD127 which allows cells to respond to IL-7 and promotes memory cell
formation. Therefore, we evaluated protective memory at 30 dpi. For this, we challenged
previously infected mice with a lethal dose (5000 PFU) of the A/PR/8 (PR8, H1N1)
(PR8) influenza strain. Importantly, PR8 (H1N1) and X31 (H3N2) are serologically
distinct strains, and protection against this heterosubtypic infection is partially T cell
dependent (33). All mice treated with Fc.Mut24 (either before or after initial X31
infection) were fully protected from PR8 infection at this timepoint, showing weight loss
and recovery comparable to vehicle-treated X31 immune controls and surviving the
challenge, whereas naïve mice all succumbed to PR8 (Figure 6A).

To assess long-term durability of protective memory, we treated mice with Fc.Mut24
either one day before or 4 days after X31 infection, and evaluated memory T cell
abundance and protective responses at 170 dpi. Flu-sp CD8$^+$ T cells remained
detectable in the lungs of all animals, and both their frequency and number were
comparable between Fc.Mut24- and vehicle-treated groups (Figure 6B and C). We also
examined localization of memory cells in the lung parenchyma vs. vasculature, and
expression of compartment-specific residency markers (CD69, CX3CR1, KLRG1), but
found no significant differences at 170 dpi in any of our experimental groups
(Supplemental Figure 8A and B). Upon lethal PR8 challenge infection, both vehicle- and
Fc.Mut24-treated X31 immune were protected from lethal disease, and mice treated
with Fc.Mut24 at 4dpi during initial X31 infection trended toward milder disease and less
weight loss compared to vehicle controls, although this difference did not reach
statistical significance (Figure 6 D and E). Moreover, the Flu-sp CD8$^+$ T cell response
showed no differences in magnitude, phenotype, or tissue localization between any of
the X31 immune groups (Figure 6E; Supplemental Figure 8C and D). Thus, although
Fc.Mut24 alters the acute CD8 T cell response—either suppressing or enhancing
expansion of Flu-sp CD8$^+$ T cells depending on timing—it does not compromise the
generation or maintenance of protective T cell memory. Similarly, there were no
differences in the maintenance of IgG responses to NP at 170dpi (Supplemental Figure
8E), and thus our findings show that regardless of timing relative to initial infection,
Fc.Mut24 treatment does not impact the integrity and durability of memory Flu-sp CD8$^+$
T cell responses.

Discussion
370
Our study demonstrates that the timing of IL-2-based immunotherapy critically shapes
its impact on anti-viral CD8$^+$ T cell responses. Using Fc.Mut24, a Treg-selective IL-2
mutein, we show that treatment before respiratory virus infection suppressed the
expansion and function of Flu-specific CD8$^+$ T cells and alters their localization within
the lungs, whereas treatment after infection expands virus-specific CD8$^+$ T cells through
CD25-dependent signaling and worsens symptomatic disease, likely through enhanced
immunopathology. However, regardless of timing of treatment, mice recovered from
infection and developed durable protective immunity against lethal heterosubtypic
infections. These divergent and context-dependent effects appear to reflect a balance
between suppressive Treg activity, and direct IL-2 signaling that promotes differentiation
of CD25$^+$ effector CD8$^+$ T cells.

Administering Fc.Mut24 before infection significantly reduced the abundance of Flu-
specific CD8$^+$ T cells. Interestingly, despite this overall reduction, these cells showed a
skewed localization toward the lung parenchyma. Viral load and disease severity
remained unchanged despite the reduced number and altered distribution of CD8$^+$ T
cells, and this indicates that even in the face of massive Treg expansion, it is possible to
generate protective Flu-specific CD8$^+$ T cell responses to efficiently clear the infection.
Through expansion of Treg cells expressing high levels of CTLA-4 Fc.Mut24 treatment
before infection likely limits the availability of the CD80 and CD86 co-stimulatory
ligands. Similarly, the large reservoir of CD25^{hi} Treg cells may reduce IL-2 availability
for Flu-sp CD8$^+$ T cells during the expansion phase, and together these factors may
favor the priming and expansion of CD8$^+$ T cells that receive strong TCR signals and
effectively clear virus and form protective memory despite the immunosuppressive
environment present during priming. Indeed, their function as an IL-2 sink is a key
mechanism by which Treg can suppress CD8$^+$ T cell responses (34, 35), and this takes
place in a key priming window that occurs 2-3 days after initial activation (36). Since IL-
2 signaling directly inhibits CD127 expression by responding CD8$^+$ T cells (37), this
could also help explain why formation of durable heterosubtypic immunity was intact
despite a reduced initial effector response. IL-2 signaling during priming also drives
CX3CR1 expression (38), and decreased IL-2 availability in Fc.Mut24 pre-treated mice
may therefore decrease expression of this receptor and promote parenchymal rather
than vascular T cell localization in the lung. Despite limiting expansion of Flu-sp CD8$^+$ T
cells, Fc.Mut24 had minimal effects on bystander T cells during X31 infection. As the
recruitment and activation of bystander CD8$^+$ T cells is driven largely by chemokines
and cytokines produced by cells of the innate immune system (39), this supports a
mechanism whereby Treg cells interrupt key pathways central to elaboration of the
antigen-specific adaptive immune response.

Interestingly, we also observed no differences in total IgG specific for the viral NP
protein at any time point examined. This is surprising given that IL-2 is a key inhibitor of
Tfh differentiation and function (29, 30), and therefore restricting IL-2 availability by
expanding Treg prior to infection may be expected to promote better antibody
responses. Indeed, despite the reduced CD8$^+$ T cell response we observed in Fc.Mut24
pre-treated mice, these animals successfully cleared the virus and established durable
protective memory that was quantitatively indistinguishable from that observed in
vehicle-treated mice, indicating that long-term memory formation and maintenance was
not strictly dependent on the magnitude of the initial clonal expansion.

When administered 4 days after infection had begun, Fc.Mut24 drove hyper-expansion
of Flu-sp CD8$^+$ T cells. Under these conditions, we found that CD25-dependent
signaling in responding Flu-sp CD8$^+$ T cells promoted their increased expansion and
accumulation in the lungs. We hypothesize that this results in enhanced immune-
mediated pathology in the lungs (40), consistent with the increased weight loss
observed in these mice. Interestingly, as we observed with Fc.Mut24 pre-treatment, flu-
specific CD8$^+$ T cells in mice given Fc.Mut24 four days after infection also expressed
elevated CD127 and decreased CX3CR1, although this did not result in enhanced
parenchymal localization. The direct impact of IL-2 on CD8$^+$ T cell differentiation is

thought to take place in a distinct priming window that would occur before Fc.Mut24 treatment in our system (36), and therefore we hypothesize that the changes in receptor expression we observe are due to enhanced proliferation of CD127$^+$CX3CR1$^-$ memory precursor cells in response to Fc.Mut24 after this differentiation window has closed. In this way, Fc.Mut24 treatment during infection may actually boost lasting protective memory formation, as mice treated as such trended towards less weight loss than vehicle controls upon heterosubtypic PR/8 rechallenge. Consistent with this, others have also described IL-2 administration as boosting CD8$^+$ memory T cell generation and function in the context of active inflammation (41, 42). Sustained antibody responses may be explained by the enhanced proliferation of Tregs which have been shown to favor Tfh generation during influenza infection (43), thus enabling sufficient antibody responses and explain why we did not see reduced total IgG against the viral NP.

Together, our findings demonstrate that although treatment with Fc.Mut24 exerts immunosuppressive effects through Treg expansion, respiratory viral infection is still effectively controlled and durable memory responses are established, even when peak Treg expansion coincides with the initial priming phase. This observation has important implications for use of IL-2 muteins and other Treg-expanding immunotherapies, suggesting that the presence of Treg-mediated suppression does not necessarily preclude effective anti-viral T cell responses, viral clearance and development of protective immunity. However, these results also highlight a critical caveat in that IL-2-based therapies may exacerbate immunopathology if administered during active infection. This is especially relevant in the context of respiratory viral infections, where immune-driven tissue damage significantly contributes to morbidity and mortality (44). These results underscore the need for careful consideration of the timing and context of Treg-targeted interventions in infectious disease settings, particularly when balancing immune regulation with the necessity of effective pathogen clearance.

Materials and Methods

Mice
C57BL/6 (B6) (JAX Strain# 000664) mice were purchased from The Jackson Laboratory. B6.SJL-Ptprca Pepcb/BoyJ (CD45.1$^+$, JAX Strain# 002014) and C57BL/6-Tg(TcraTcrb)1100Mjb/J (OT-I, JAX Strain # 003831) mice were bred and maintained at Benaroya Research Institute (BRI). Female Mice used in experiments between the ages of 6-10 weeks at start of experiment. Female mice are used due to male B6 mice being more resistant to infection (45). For histological analyses, euthanasia was performed with tribromoethanol overdose. For all other experiments mice were euthanized by CO$_2$ inhalation. All experiments were performed in accordance with the guidelines and with the approval of the Institutional Animal Care and Use Committee of the BRI.

Viruses and infections
Influenza viruses A/HK-x31(x31, H3N2), A/PR/8 (PR8, H1N1) (46), and the X31-ova expressing the CD8 H2-K^b restricted SIINFEKL epitope (47), were all generously provided by Dr. Kimberly D. Klonowski (University of Georgia). Animals were infected with 2000 PFU of either X31 strain or 5000PFU of PR8 strain intranasally in 50uL of

PBS. All mice infected with Influenza were monitored at least daily for weight loss and
twice a day when mice reached 80% of baseline weight. If mice reached 70% of
baseline weight, mice were promptly euthanized using aforementioned methods. For
longer term studies, once mice recovered they were monitored twice weekly. VSV-OVA
(48) was generously provided from Dr. Pamela Fink (University of Washington). Mice
were infected with 2000PFU of VSV-OVA, intraperitoneally in 100 uL of PBS.

IL-2 mutein treatment
The development of the murine Fc.IL-2 mutein (Fc.Mut24) was previously described
(10). Fc.Mut24 was produced and purified by Olympic Protein Technologies (Seattle,
WA), and contained less than 15 endotoxin units (EU)/mL. For all experiments, 10 µg of
Fc.Mut24 or PBS vehicle was administered via intraperitoneal injection.

Cell isolation and flow cytometry
Where applicable, mice were anesthetized using 4% isoflurane and intravenously
injected with 3µg of anti-CD45 or anti-CD45.2 antibody in 100uL for intravascular
labeling. Mice were allowed to recover for two minutes before euthanasia, performed as
described above. For flow cytometry, lungs were minced and digested in RPMI
supplemented with Liberase (50 µg/mL) and DNase I (10 U/mL) for 20 minutes at
37 °C with agitation. Cell suspensions were filtered through 70 µm strainers into RPMI
containing 10% FBS (RPMI-10), followed by red blood cell lysis with ACK lysis buffer
and washing in RPMI-10. The same protocol was used for lymphoid tissues (mediastinal
lymph node and spleen) when analyzing conventional dendritic cells (cDCs). Otherwise,
spleens and medLNs were mechanically dissociated through a 70 µm strainer,
followed by red blood cell lysis and washing as above. For dendritic cell analysis,
CD11c[+] cells were enriched using anti-CD11c microbeads (Miltenyi Biotec) according to
the manufacturer's instructions. Enriched cells were stained with selected antibodies
targeting cDC surface markers (see reagent table).

For detection of antigen-specific CD4[+] T cells, MHC class II tetramer staining was
performed using the I-A^b/NP$_{311-325}$ (QVYSLIRPNENPAHK) tetramer (49), kindly provided
by Marion Pepper (University of Washington). Samples were incubated with tetramer at
room temperature for 1 hour, washed, and enriched using anti-APC microbeads
(Miltenyi Biotec). Enriched cells were then stained with the CD4 flow cytometry panel,
while the flow-through was used for additional staining and analysis. For CD8[+] T cell
analysis, H-2D^b/NP$_{366-374}$ (ASNENMETM) or H-2K^b/SIINFEKL tetramers (generated by
the NIH Tetramer Facility, Emory University) were used. Cells were stained with
tetramer for 1 hour at room temperature prior to surface staining.

Cell surface staining for flow cytometry was performed in FACS buffer (PBS-2% BCS)
using a selected antibody cocktail (See reagent table). Cells were incubated in the
antibody cocktail for 20 min at room temperature and then washed in FACS buffer
before collecting events on an BD Symphony flow cytometer. For intracellular staining,
surface antigens were stained before fixation and permeabilization with FixPerm buffer

(eBioscience) (See reagent table). Cells were washed and stained with selected
antibodies. Flow cytometry data was analyzed using FlowJo software.

Cell stimulations:

Single-cell suspensions were prepared from tissues (as described above) and cultured
in complete RPMI in the presence of influenza $NP_{366-374}$ peptide (ASNENMETM, H-2D□
restricted) at 10 microMolar in 200 µL of cRPMI. Cells were incubated with secretion
inhibitors (brefeldin A and monensin at 1x manufactures recommendation) throughout
the stimulation period. Following stimulation, cells were stained with a viability dye and
fluorochrome-conjugated antibodies against surface markers, then fixed, permeabilized,
with BD cytoperm/Fix (cat. 554722) and stained for intracellular cytokines IL-2 and IFN-
γ.

Quantification of viral RNA
Total RNA was extracted from lung tissue using the Qiagen RNeasy Plus Mini Kit (cat.
74134) following the manufacturer's protocol. Tissue homogenization was performed in
RLT buffer containing β-mercaptoethanol (BME), with buffer volume adjusted according
to tissue weight as specified by the kit. Approximately 15 mg of tissue was used per
sample to avoid overloading the RNA-binding column. Samples were lysed using 5 mm
stainless steel beads (Qiagen, cat. 69989) in 2.0 mL Safe-lock Eppendorf tubes on a
TissueLyser II (Qiagen) at 30 Hz for 2 minutes. RNA yield was quantified using a
NanoDrop spectrophotometer. Complementary DNA (cDNA) was synthesized from
purified RNA using SuperScript II Reverse Transcriptase (Invitrogen, cat. 18064-014)
and IDT ReadyMade Random Hexamer primers (cat. 51-01-18-26), including a no-RNA
negative control in each batch. Reactions were assembled using a two-step thermal
protocol. First, Master Mix 1 was added and samples were incubated at 65□°C for 5
minutes, then cooled to 4□°C. After addition of Master Mix 2, samples were incubated
at 30□°C for 10 minutes followed by 42□°C for 50 minutes. The resulting cDNA was
quantified using a NanoDrop spectrophotometer. Quantitative PCR (qPCR) was used to
quantify Influenza A virus matrix (M1/M2) gene expression from PR8 and X31 strains.
Reactions were prepared using IDT PrimeTime Std qPCR Assay containing a
fluorescent probe and primers specific to the M segment of IAV with the following primer
and probe design: Probe: (6-FAM/ZEN/IBFQ) 5′-CCTCTGCTGCTTGCTCACTCGATC-
3′; Forward Primer: 5′-CAGCACTACAGCTAAGGCTATG-3′; Reverse Primer: 5′-
CTCATCGCTTGCACCATTTG-3′. The probe/primer mix was resuspended at a 20x
concentration. Each qPCR reaction (20□µL total volume) contained: 10□µL 2x IDT
PrimeTime Gene Expression Master Mix (cat. 1055770),1□µL 20x PrimeTime qPCR
Assay Mix, 4□µL diluted cDNA (~75 ng), 5□µL nuclease-free water. An internal control
targeting hypoxanthine-guanine phosphoribosyltransferase (HPRT) was run in parallel
to normalize viral gene expression. The Viia7 Real-Time PCR System (Thermo Fisher
Scientific) was used for amplification and detection.

Adoptive transfers

Spleen and lymph nodes were harvested from OT-I mice, passed through a 70 μm
strainer and ACK lysed. CD8$^+$ cells were enriched by negative selection using a CD8 T
cell isolation kit according to the manufacturer's instructions (Miltenyi). Cells were then
enumerated, washed in PBS before transfer by retroorbital injection into recipient mice
(0.005-0.5×10^6 cells per mouse depending on the experiment). For transfer of activated
OT-I CD8 T cells, spleens were collected from CD45.1/.2 mice and processed as
described above. Once in single cell suspension, 1x10^6 cells were plated in a 12-well
plate (see reagent table) in 1mL of cRPMI supplemented with 50UI/mL of rIL-2 (see
reagent table). Cells were plated at a ratio 1:1 ratio with washed Dynabeads and
allowed to culture for 48hrs. Following 48hrs, Dynabeads (see reagent table) were
extracted using a magnet and cells washed before being transferred retroorbitally to
mice at 0.5×10^6 cells per mouse.)

CRISPR deletion of Il2ra
For deletion of the *Il2ra* gene in primary naive OT-I TCR T cells, we utilized the
CRISPR/CAS9 system as previously described (32). Purified CAS9 protein, *Il2ra* crRNA
(AGAUGAAGUGUGGGAAAACGGUUUUAGAGCUAUGCU), and trans-activator RNA
(tracrRNA) were purchased from Integrated DNA Technologies. CD8$^+$ T cells were
isolated from OT-I mouse spleen using CD8$^+$ T cell negative selection MojoSort Kit
(Biolegend). T cells were cultured in IL-7 (10ng/ml) for 24 hrs prior to transfection.
CRISPR/CAS9 reagents were prepared as per the manufacturer's instructions for IDT
Alt-R CRISPR-CAS9 system. The pre-cultured CD8$^+$ cells were were resuspended in T
buffer (Invitrogen MPK1096) and electroporated using 6 2200V, 10ms, 3 pulses, with
the Neon electroporation system. Following transfection, cells were transferred into
warm RPMI with 10% FCS and were incubated at 37^{o}C, 5% CO2 for 4- 5 hours. Prior to
adoptive transfer, the electroporation efficiency was evaluated by flowcytometry
measuring ATTO uptake. Control cells were transfected with ATTO and CAS9 alone
without *Il2ra* crRNA. To validate *Il2ra* deletion efficiency, the neon-transfected cells were
plated on αCD3/αCD28 coated plates for activation for 48 hours followed by flow
cytometry analysis of cell surface CD25 expression.

Enzyme-Linked Immunosorbent Assay (ELISA)
To quantify serum antibody responses against influenza PR8 nuclear protein, ELISA
plates were coated overnight at 4°C with anti-His antibody (1:1000 dilution in PBS,
50 μL/well). Plates were then washed three times with 100 μL PBS containing 0.1%
Tween-20 (PBS-T), followed by incubation with His-tagged PR8 nuclear protein
(SinoBiological, Cat# 11675-V08B) at 1:100 dilution in PBS-T (50 μL/well) for 1 hour
at 37°C. After three additional PBS-T washes, wells were blocked with 100 μL of
blocking buffer (PBS-T supplemented with 5% BSA) for 1 hour at 37°C. Diluted serum
samples were added to the plates (50 μL/well) and incubated for 1 hour at 37°C.
Plates were then washed three times with PBS-T and incubated with 50 μL of biotin-
conjugated anti-mouse IgG antibody (Jackson ImmunoResearch, Cat# 115-035-062)
diluted 1:1000 in blocking buffer for 1 hour at 37°C. After five washes with PBS-T,

50 µL of 1× TMB substrate was added to each well and color development was
monitored. The reaction was stopped with 50 µL of H_2SO_4, and absorbance was
measured at 450 nm using a microplate reader.

Histological analysis
Lungs were excised and immediately fixed in 10% formalin for at least 24hrs and then
paraffin embedded. H&E staining was performed on a 5µm tissue sections by Benaroya
Research Institute Histology Core. Resulting slides were imaged using a Molecular
Devices ImageXpress Confocal with the 10x objective by Benaroya Research Institute
Histology Core.

Statistics
All data are presented as mean values ± SD, and graphs were created and analyzed
using Prism Software (GraphPad). Comparisons between treatment groups were
analyzed using two-tailed unpaired student T tests, one-, or two-way ANOVA where
appropriate, adjusted for multiple comparisons using Tukey's post-test. To compare
weight loss between experimental groups over time, a two-way mixed-effects ANOVA
was performed. This mixed-effects approach allows for modeling the correlation
structure of repeated observations and provides more accurate estimates by adjusting
for inter-subject variability.

Writing disclosure
Portions of the manuscript text were edited for grammar and clarity using OpenAI's
ChatGPT (version GPT-4). The tool was used solely for language refinement; all
scientific content and interpretations are the original work of the authors.

Acknowledgements:

We would like to thank Pam Johnson and Riley Snodgrass for assistance with histology;
Adam Wojno and Adin Pierce for help with flow cytometry; Andrew Biurch and the BRI
veterinary staff for support with mouse work; and Marion Pepper, Mikel Ruterbusch, Kurt
Pruner, and Geoff Hutchinson for providing the CD4 T cell tetramer. This work was
supported by grants R01AI154773 and R21AI172140 from the NIH to D.J.C.

 Figure Legends

Figure 1: Fc.Mut24 administration before X31 infection limits Flu-sp CD8$^+$ T cell
responses. Mice were treated one day before infection with 2000PFU of X31 with
either vehicle (blue) or Fc.Mut24 (red). All data represent takedown at 9dpi. **A)**
Representative flow cytometry analysis of CD25 and Foxp3 expression by gated CD4$^+$ T
cells in the lungs of vehicle- and Fc.Mut24-treated mice (left), and quantification of the
number and frequency of Tregs in the lungs (right). **B)** Representative flow cytometry
analysis of CD44 expression and NP$_{366-374}$ tetramer staining by gated CD8$^+$ T cells in
the lungs of vehicle- and Fc.Mut24-tretted mice (left), and quantification of the number
and frequency of Flu-sp T cells in the lungs (right). **C)** Representative histogram of IV
label staining on gated Flu-sp T cells in lungs of vehicle- and Fc.Mut24-treted mice (left),
and quantification of the frequency and number Flu-sp CD8 T cells in parenchymal vs
blood exposed compartments (right). **D)** Representative flow cytometry staining and
quantitative analyses showing expression of the selected markers by gated CD44^{lo}
naïve CD8$^+$ T cells in the lungs of vehicle-treated mice, or by Flu-sp T cells in the lungs
of vehicle- or Fc.Mut24-treated mice as indicated. **E)** Weight loss as a percent of
baseline in vehicle- or Fc.Mut24-treated mice during the course of X31 infection. **F)**
Quantification of viral genes *M1/M2* relative to *HPRT* by PCR in vehicle- or Fc.Mut24-
treated mice as indicated. Each point represents analysis of an individual animal. All
data are representative of two separate experiments *$p < 0.05$, **$p < 0.01$,
****$p < 0.0001$, by unpaired T test for A-C and one way ANOVA with multiple
comparison for D and F and with mixed effects for E.

Figure 2: Fc.Mut24 treatment does not impact bystander CD8 T Cells. **A)** Mice
were treated one day before infection with 2000PFU of X31 with either vehicle (blue) or
Fc.Mut24 (red). Representative flow cytometry analysis of CD44 expression and NP$_{366-}$
$_{374}$ tetramer staining by gated CD8$^+$ T cells in the lungs of vehicle- and Fc.Mut24-tretted
mice (left), and quantification of the number and frequency of CD44^{hi}Tetramer$^-$ T cells in
the lungs (right). **B)** Mice were infected with VSV-OVA 30d before treatment with vehicle
or Fc.Mut24 and X31 infection. All data represent takedown at 9dpi with X31.
Representative flow cytometry analysis of CD44 expression and SIINFEKL tetramer
staining by gated CD8+ T cells in the lungs of mice treated as indicated (left), and
quantitative analysis of the number and frequency of SIINFEKL-sp CD8 T cells across
groups (right). N = 5 per group. Each point represents analysis of an individual animal.
*$p < 0.05$, **$p < 0.01$, ****$p < 0.0001$, by by unpaired T test for A and one way
ANOVA with multiple comparison for B.

Figure 3: Fc.Mut24 treatment before X31 infection reduces antigen presentation
capacity of DCs in the lungs and spleen. Mice were treated one day before infection
with 2000PFU of X31 with either vehicle (blue) or Fc.Mut24 (red). **A)** Representative
flow cytometry analysis of CD25 and Foxp3 expression by gated CD4$^+$ T cells in the
lungs of vehicle- and Fc.Mut24-treated mice (left), and quantification of the number and
frequency of Tregs in the lungs (right) at 3dpi. **B)** Representative flow cytometry

analysis of CD80 and CD86 expression by gated cDC1 (CD3/B220⁻Ly6G⁻F4/80⁻

MHCII^high CD11c^high XCR1⁺SIRPalpha⁻) and cDC2 (CD3/B220⁻Ly6G⁻F4/80⁻

MHCII^high CD11c^high XCR1⁻SIRPalpha⁺) cells from lungs at 3dpi with X31 (top).

Quantification of activated CD80⁺CD86⁺ cDC1 or cDC2 in vehicle- or Fc.Mut24-treated

mice as indicated. **C)** CD8⁺ T cells from CD45.1⁺CD45.2⁺ OT-I mice were enriched and

transferred to CD45.2⁺ C57BL/6 recipients concurrent with vehicle- or Fc.Mut24-

treatment one day prior to infection with X31-SIINFEKL. Representative flow cytometry

analysis of CD8 and CD45.1 expression by gated live lymphocytes (left), and

quantification of the frequency and number of transferred OT-I T cells (right) from the

lungs of recipient mice at 4dpi (right). **D)** CD8⁺ T cells from CD45.1⁺CD45.2⁺ OT-I mice

were activated for 48hrs and then transferred into transferred to CD45.2⁺ C57BL/6

recipients at 3dpi with X31-SIINFEKL. Representative flow cytometry analysis of CD8

and CD45.1 expression by gated live lymphocytes (left), and quantification of the

frequency and number of transferred OT-I T cells (right) from the lungs of recipient mice

(right) at 9dpi (6 days after cell transfer). Panels A, C and D are representative of one

experiment n = 5 mice per group. Panel B is representative of two separate experiments

n = 10 per group. Each point represents analysis of an individual animal. *p < 0.05,

**p < 0.01, ****p < 0.0001, by unpaired T test for all panels.

Figure 4: Fc.Mut24 treatment during infection enhances the Flu-specific CD8 T

cell response. Mice were treated at 4dpi with X31 with either vehicle (blue) or

Fc.Mut24 (green)**.** All data represent takedown at 9dpi. **A)** Representative flow

cytometry analysis of CD44 expression and NP$_{366-374}$ tetramer staining by gated CD8⁺ T

cells in the lungs of vehicle- and Fc.Mut24-tretted mice (left), and quantification of the

number and frequency of Flu-sp T cells in the lungs (right) **C)** Representative histogram

of IV label staining on gated Flu-sp T cells in lungs of vehicle- and Fc.Mut24-treted mice

(left), and quantification of the frequency and number Flu-sp CD8 T cells in parenchymal

vs blood exposed compartments (right). **C)** Weight loss as a percent of baseline in

vehicle- or Fc.Mut24-treated mice during the course of X31 infection. Dotted line

indicates time of vehicle or Fc.Mut24 treatment. **D)** Quantification of viral genes *M1/M2*

relative to *HPRT* by PCR in vehicle- or Fc.Mut24-treated mice as indicated. **E)**

Representative H&E images from mice at 12 days post infection. All data are

representative of two separate experiments with n = 10. Each point represents analysis

of an individual animal. *p < 0.05, **p < 0.01, ****p < 0.0001, unpaired T test for

panels A and B and one way ANOVA with multiple comparison for D and Mixed Effects

for panel C.

Figure 5: Fc.Mut24 directly expands IAV-specific CD8 T cells. 5x10⁵ congenic OT-I

CD8 T cells were transferred to mice, that were a day later infected with X31-SIINFEKL

and treated with 10ug of Fc.Mut24 at 4dpi. Data represents 9dpi takedown of lungs of

indicated mice. **A)** Mice were treated at 4dpi with X31 with either vehicle (blue) or

Fc.Mut24 (green)**.** Data represent takedown at 9dpi. Representative flow cytometry

staining and quantitative analyses showing CD25 expression by gated CD44^lo naïve

CD8⁺ T cells in the lungs of vehicle-treated mice, or by Flu-sp T cells in the lungs of

vehicle- or Fc.Mut24-treated mice as indicated. **B)** CD8$^+$ T cells from CD45.1$^+$ OT-I mice were enriched transfected with CAS9 RNPs containing a guide (g)RNA targeting the *Il2ra* locus or a control non-targeting gRNA. Cells were transferred to CD45.2$^+$ C57BL/6 recipients that were infected with X31-SIINFEKL and treated with vehicle of Fc.Mut24 4 days later as indicated. Representative flow cytometry analysis of CD45.1 and CD45.2 expression by gated CD8+ T cells from the lungs of mice at 9dpi showing gate for donor OT-I T cells (left), and quantitative analysis of the abundance and frequency of donor OT-I cells (right). **C)** Representative flow cytometry analysis of CD44 expression and SIINFEKL tetramer binding by CD45.2+CD45.1- recipient cells (left) and quantitative analysis of endogenous SIINFEKL-sp CD8$^+$ T cells from lungs of mice treated as in B. Data are from one experiment with n=5 mice/group, representative of two separate experiments. *p$<$0.05, **p$<$0.01, ****p$<$0.0001, by one way ANOVA with multiple comparisons.

Figure 6: Fc.Mut24 does not alter generation or durability of protective T cell memory to influenza. Mice were infected with X31 and treated with vehicle (blue), or with Fc.Mut24 either a day before (red) or 4 days after (green) infection.. **A)** Mice were lethally challenged with 5000 FU of PR8 at 30 days post X31 infection. Disease severity based on weight loss from start of lethal challenge in mice treated with Fc.Mut24 one day before (left) or 4 days after (right) initial X31 infection (left) compared with vehicle-treated and X31-naïve mice. **B), C)** Representative flow cytometry analysis of CD44 expression and NP$_{366-374}$ tetramer staining by gated CD8$^+$ T cells in the lungs of vehicle- and Fc.Mut24-tretted mice, and quantification of the number and frequency of CD44^{hi}Tetramer$^-$ T cells in the lungs at 170dpi in mice treated one day prior to **B)** or 4 days after **C)** initial X31infection. **D), E)** X31-immune mice treated with vehicle or Fc.Mut24 were lethally challenged with 5000 FU of PR8 at 170dpi. Disease severity based on weight loss from start of lethal challenge (left). Quantification of Flu-sp CD8 T cells in lung (right) in mice treated with Fc.Mut24 one day prior to **D)** or 4 days after **E)** initial X31 infection. *p$<$0.05, **p$<$0.01, ****p$<$0.0001, by unpaired T test and two way ANOVA with mixed effects for weight loss curves in panels A, C, D and E.

References

1. Raffin, C., L. T. Vo, and J. A. Bluestone. 2020. T(reg) cell-based therapies: challenges and perspectives. *Nat Rev Immunol* 20: 158-172.
2. Wright, R. C., D. J. Campbell, and M. K. Levings. 2025. Pharmacotherapeutic strategies to promote regulatory T cell function in autoimmunity. *Curr Opin Immunol* 94: 102554.
3. Shouse, A. N., K. M. LaPorte, and T. R. Malek. 2024. Interleukin-2 signaling in the regulation of T cell biology in autoimmunity and cancer. *Immunity* 57: 414-428.
4. Chinen, T., A. K. Kannan, A. G. Levine, X. Fan, U. Klein, Y. Zheng, G. Gasteiger, Y. Feng, J. D. Fontenot, and A. Y. Rudensky. 2016. An essential role for the IL-2 receptor in T(reg) cell function. *Nat Immunol* 17: 1322-1333.
5. Osum, K. C., and M. K. Jenkins. 2023. Toward a general model of CD4(+) T cell subset specification and memory cell formation. *Immunity* 56: 475-484.
6. Klatzmann, D., and A. K. Abbas. 2015. The promise of low-dose interleukin-2 therapy for autoimmune and inflammatory diseases. *Nat Rev Immunol* 15: 283-294.
7. Dashwood, A., A. A. Ghodsinia, J. Dooley, and A. Liston. 2025. Cytokine Couture: Designer IL2 Molecules for the Treatment of Disease. *Immunotargets Ther* 14: 403-431.
8. Scheid, J. F., K. Cunningham-Bussel, N. Kim, S. Agarwal, G. Nieddu, J. Cote, L. Lemoine, T. Decaesteker, L. Mendez, E. Paul, L. Love-Gregory, A. V. Contreras, X. Zhao, L. Franco-Dilone, L. Pang, G. A. Baltus, M. Beaumont, K. Shah, N. Higginson-Scott, K. Kis-Toth, K. L. Otipoby, J. L. Viney, E. Sicard, S. Rottey, J. S. Sundy, K. Van Dyck, T. Laethem, P. Larson, S. Sutradhar, R. Wnek, T. Bueters, E. Lai, S. A. Stoch, M. Iwamoto, and J. A. Robbins. 2025. Safety, pharmacokinetics, and pharmacodynamics of MK-6194, an IL-2 mutein designed to selectively activate regulatory T cells: single ascending dose and multiple ascending dose trial data. *Immunohorizons* 9.
9. Silverberg, J. I., D. Rosmarin, R. Chovatiya, T. Bieber, S. Schleicher, L. Beck, M. Gooderham, S. Chaudhry, C. Fanton, D. Yu, J. Levy, Y. Liu, T. Miyazaki, M. Tagliaferri, C. Schmitz, A. Nirula, B. Kotzin, and J. Zalevsky. 2024. The regulatory T cell-selective interleukin-2 receptor agonist rezpegaldesleukin in the treatment of inflammatory skin diseases: two randomized, double-blind, placebo-controlled phase 1b trials. *Nat Commun* 15: 9230.
10. Khoryati, L., M. N. Pham, M. Sherve, S. Kumari, K. Cook, J. Pearson, M. Bogdani, D. J. Campbell, and M. A. Gavin. 2020. An IL-2 mutein engineered to promote expansion of regulatory T cells arrests ongoing autoimmunity in mice. *Sci Immunol* 5.
11. Pham, M. N., L. Khoryati, B. L. Jamison, E. Hayes, J. M. Sullivan, D. J. Campbell, and M. A. Gavin. 2021. In Vivo Expansion of Antigen-Specific Regulatory T Cells through Staggered Fc.IL-2 Mutein Dosing and Antigen-Specific Immunotherapy. *Immunohorizons* 5: 782-791.
12. Jamison, B. L., M. Lawrance, C. J. Wang, H. A. DeBerg, L. J. Ziegler, D. M. Sansom, M. A. Gavin, L. S. K. Walker, and D. J. Campbell. 2024. An IL-2 mutein

increases regulatory T cell suppression of dendritic cells via IL-10 and CTLA-4 to
promote T cell anergy. *Cell Rep* 43: 114938.

13. Klebanoff, S. D., L. B. Rodda, C. Morishima, M. H. Wener, Y. Yuzefpolskiy, E.
Bettelli, J. H. Buckner, C. Speake, M. Pepper, and D. J. Campbell. 2023.
Diminished responses to mRNA-based SARS-CoV-2 vaccines in individuals with
rheumatoid arthritis on immune-modifying therapies. *JCI Insight* 8.

14. Dey, M., K. Bechman, S. Zhao, G. E. Fragoulis, C. Smith, A. Cope, E. Nikiphorou,
K. L. Hyrich, and J. Galloway. 2022. Infection profile of immune-modulatory drugs
used in autoimmune diseases: analysis of summary of product characteristic
data. *RMD Open* 8.

15. Yuzefpolskiy, Y., P. Morawski, M. Fahning, C. Speake, S. Lord, A. Chaudhary, C.
Morishima, M. H. Wener, M. Kita, L. McCarthy, J. H. Buckner, D. J. Campbell,
and E. Bettelli. 2022. Cutting Edge: Effect of Disease-Modifying Therapies on
SARS-CoV-2 Vaccine-Induced Immune Responses in Multiple Sclerosis Patients.
J Immunol 208: 1519-1524.

16. Garcillan, B., M. Salavert, J. R. Regueiro, and S. Diaz-Castroverde. 2022.
Response to Vaccines in Patients with Immune-Mediated Inflammatory Diseases:
A Narrative Review. *Vaccines (Basel)* 10.

17. Salliot, C., M. Dougados, and L. Gossec. 2009. Risk of serious infections during
rituximab, abatacept and anakinra treatments for rheumatoid arthritis: meta-
analyses of randomised placebo-controlled trials. *Ann Rheum Dis* 68: 25-32.

18. Goldmann, O., O. V. Nwofor, Q. Chen, and E. Medina. 2024. Mechanisms
underlying immunosuppression by regulatory cells. *Front Immunol* 15: 1328193.

19. Patterson, C. M., T. D. Shaw, V. Gerovasili, U. Khatana, and R. J. Jose. 2024.
Emerging therapies and respiratory infections: Focus on the impact of
immunosuppressants and immunotherapies. *Clin Med (Lond)* 24: 100015.

20. Furst, D. E. 2010. The risk of infections with biologic therapies for rheumatoid
arthritis. *Semin Arthritis Rheum* 39: 327-346.

21. Arpaia, N., J. A. Green, B. Moltedo, A. Arvey, S. Hemmers, S. Yuan, P. M.
Treuting, and A. Y. Rudensky. 2015. A Distinct Function of Regulatory T Cells in
Tissue Protection. *Cell* 162: 1078-1089.

22. Sjaastad, L. E., D. L. Owen, S. Joo, T. P. Knutson, C. H. O'Connor, B.
McCluskey, R. S. LaRue, R. A. Langlois, and M. A. Farrar. 2022. Influenza
infection recruits distinct waves of regulatory T cells to the lung that limit lung
resident IgA+ B cells. *bioRxiv*: 2022.2009.2019.508325.

23. Anderson, K. G., H. Sung, C. N. Skon, L. Lefrancois, A. Deisinger, V. Vezys, and
D. Masopust. 2012. Cutting edge: intravascular staining redefines lung CD8 T
cell responses. *J Immunol* 189: 2702-2706.

24. Hoft, S. G., M. A. Sallin, K. D. Kauffman, S. Sakai, V. V. Ganusov, and D. L.
Barber. 2019. The Rate of CD4 T Cell Entry into the Lungs during Mycobacterium
tuberculosis Infection Is Determined by Partial and Opposing Effects of Multiple
Chemokine Receptors. *Infect Immun* 87.

25. Hand, T. W., M. Morre, and S. M. Kaech. 2007. Expression of IL-7 receptor alpha
is necessary but not sufficient for the formation of memory CD8 T cells during
viral infection. *Proc Natl Acad Sci U S A* 104: 11730-11735.

26. Sckisel, G. D., J. K. Tietze, A. E. Zamora, H. H. Hsiao, S. O. Priest, D. E. Wilkins,
L. L. Lanier, B. R. Blazar, N. Baumgarth, and W. J. Murphy. 2014. Influenza
infection results in local expansion of memory CD8(+) T cells with antigen non-
specific phenotype and function. *Clin Exp Immunol* 175: 79-91.
27. Ely, K. H., L. S. Cauley, A. D. Roberts, J. W. Brennan, T. Cookenham, and D. L.
Woodland. 2003. Nonspecific recruitment of memory CD8+ T cells to the lung
airways during respiratory virus infections. *J Immunol* 170: 1423-1429.
28. Jenkins, M. M., H. Bachus, D. Botta, M. D. Schultz, A. F. Rosenberg, B. Leon,
and A. Ballesteros-Tato. 2021. Lung dendritic cells migrate to the spleen to prime
long-lived TCF1(hi) memory CD8(+) T cell precursors after influenza infection.
Sci Immunol 6: eabg6895.
29. Ballesteros-Tato, A., B. Leon, B. A. Graf, A. Moquin, P. S. Adams, F. E. Lund, and
T. D. Randall. 2012. Interleukin-2 inhibits germinal center formation by limiting T
follicular helper cell differentiation. *Immunity* 36: 847-856.
30. Pepper, M., A. J. Pagan, B. Z. Igyarto, J. J. Taylor, and M. K. Jenkins. 2011.
Opposing signals from the Bcl6 transcription factor and the interleukin-2 receptor
generate T helper 1 central and effector memory cells. *Immunity* 35: 583-595.
31. Li, P., S. Mitra, R. Spolski, J. Oh, W. Liao, Z. Tang, F. Mo, X. Li, E. E. West, D.
Gromer, J. X. Lin, C. Liu, Y. Ruan, and W. J. Leonard. 2017. STAT5-mediated
chromatin interactions in superenhancers activate IL-2 highly inducible genes:
Functional dissection of the Il2ra gene locus. *Proc Natl Acad Sci U S A* 114:
12111-12119.
32. Kalia, V., J. Reed, A. Khanna, R. Toumi, H. Xiao, T. Pulliam, R. Deo, P. Nghiem,
and S. Sarkar. 2023. Tempered IL-2 Signals Program PD-1 Checkpoint
Blockade-Responsive Stem-like Exhausted T Cells During Priming. *bioRxiv*:
2023.2005.2025.541936.
33. Liang, S., K. Mozdzanowska, G. Palladino, and W. Gerhard. 1994.
Heterosubtypic immunity to influenza type A virus in mice. Effector mechanisms
and their longevity. *J Immunol* 152: 1653-1661.
34. Kastenmuller, W., G. Gasteiger, N. Subramanian, T. Sparwasser, D. H. Busch, Y.
Belkaid, I. Drexler, and R. N. Germain. 2011. Regulatory T cells selectively
control CD8+ T cell effector pool size via IL-2 restriction. *J Immunol* 187: 3186-
3197.
35. McNally, A., G. R. Hill, T. Sparwasser, R. Thomas, and R. J. Steptoe. 2011.
CD4+CD25+ regulatory T cells control CD8+ T-cell effector differentiation by
modulating IL-2 homeostasis. *Proc Natl Acad Sci U S A* 108: 7529-7534.
36. Jobin, K., D. Seetharama, L. Ruttger, C. Fenton, E. Kharybina, A. Wirsching, A.
Huang, K. Knopper, T. Kaisho, D. H. Busch, M. Vaeth, A. E. Saliba, F. Graw, A.
Pulfer, S. F. Gonzalez, D. Zehn, Y. Liang, M. Ugur, G. Gasteiger, and W.
Kastenmuller. 2025. A distinct priming phase regulates CD8 T cell immunity by
orchestrating paracrine IL-2 signals. *Science* 388: eadq1405.
37. Kalia, V., S. Sarkar, S. Subramaniam, W. N. Haining, K. A. Smith, and R. Ahmed.
2010. Prolonged interleukin-2Ralpha expression on virus-specific CD8+ T cells
favors terminal-effector differentiation in vivo. *Immunity* 32: 91-103.
38. Barlic, J., D. H. McDermott, M. N. Merrell, J. Gonzales, L. E. Via, and P. M.
Murphy. 2004. Interleukin (IL)-15 and IL-2 reciprocally regulate expression of the

chemokine receptor CX3CR1 through selective NFAT1- and NFAT2-dependent
mechanisms. *J Biol Chem* 279: 48520-48534.
39.   Paprckova, D., E. Salyova, J. Michalik, and O. Stepanek. 2023. Bystander
activation in memory and antigen-inexperienced memory-like CD8 T cells. *Curr*
Opin Immunol 82: 102299.
40.   Duan, S., and P. G. Thomas. 2016. Balancing Immune Protection and Immune
Pathology by CD8(+) T-Cell Responses to Influenza Infection. *Front Immunol* 7:
25.
41.   Williams, M. A., A. J. Tyznik, and M. J. Bevan. 2006. Interleukin-2 signals during
priming are required for secondary expansion of CD8+ memory T cells. *Nature*
441: 890-893.
42.   Cheng, L. E., and P. D. Greenberg. 2002. Selective delivery of augmented IL-2
receptor signals to responding CD8+ T cells increases the size of the acute
antiviral response and of the resulting memory T cell pool. *J Immunol* 169: 4990-
4997.
43.   Leon, B., J. E. Bradley, F. E. Lund, T. D. Randall, and A. Ballesteros-Tato. 2014.
FoxP3+ regulatory T cells promote influenza-specific Tfh responses by controlling
IL-2 availability. *Nat Commun* 5: 3495.
44.   Newton, A. H., A. Cardani, and T. J. Braciale. 2016. The host immune response
in respiratory virus infection: balancing virus clearance and immunopathology.
Semin Immunopathol 38: 471-482.
45.   Lorenzo, M. E., A. Hodgson, D. P. Robinson, J. B. Kaplan, A. Pekosz, and S. L.
Klein. 2011. Antibody responses and cross protection against lethal influenza A
viruses differ between the sexes in C57BL/6 mice. *Vaccine* 29: 9246-9255.
46.   Verbist, K. C., C. J. Cole, M. B. Field, and K. D. Klonowski. 2011. A role for IL-15
in the migration of effector CD8 T cells to the lung airways following influenza
infection. *J Immunol* 186: 174-182.
47.   Jenkins, M. R., R. Webby, P. C. Doherty, and S. J. Turner. 2006. Addition of a
prominent epitope affects influenza A virus-specific CD8+ T cell
immunodominance hierarchies when antigen is limiting. *J Immunol* 177: 2917-
2925.
48.   Turner, M. J., E. R. Jellison, E. G. Lingenheld, L. Puddington, and L. Lefrancois.
2008. Avidity maturation of memory CD8 T cells is limited by self-antigen
expression. *J Exp Med* 205: 1859-1868.
49.   Ruterbusch, M. J., B. D. Hondowicz, K. K. Takehara, K. B. Pruner, T. S. Griffith,
and M. Pepper. 2023. Allergen exposure functionally alters influenza-specific
CD4+ Th1 memory cells in the lung. *J Exp Med* 220.

Figure 1

Figure 2