Rift Valley fever virus dynamics in a transhumant cattle system in The Gambia

Essa Jarra[1,2*], Divine Ekwem[2], Sarah Cleaveland[2], Daniel T. Haydon[2*]

[1] Department of Livestock Services, Ministry of Agriculture, Abuko, The Gambia,

[2] School of Biodiversity, One Health, and Veterinary Medicine, College of Medical, Veterinary and Life Sciences, University of Glasgow, Glasgow, United Kingdom,

* jarraessa@yahoo.com (EJ); Daniel.Haydon@glasgow.ac.uk (DTH)

Abstract

Rift Valley fever (RVF) is a zoonotic disease of global concern, driven by environmental conditions, vector activity, and livestock mobility. Although RVF has been reported in The Gambia, its epidemiology remains poorly understood. This study developed a compartmental model to study RVF dynamics in the cattle population of the country. The model incorporated seasonally dynamic transmission parameters reflecting transhumant movement and ecological differences between two distinct ecoclimatic regions: the Sahelian area and the Gambia river. Parameterised using serological data linked to household survey data, the model predicted endemic RVF virus (RVFV) circulation within The Gambia, and captured temporal infection trends that closely match empirical data. Weak decay rates of seropositivity were required to match predicted and observed age-seroprevalence. Results indicated sustained RVFV transmission during the dry season in the Gambia river eco-region, with a high risk of seasonal virus introductions to the Sahelian eco-region at the start of the wet season via the returning transhumant cattle. Our study highlighted the role of livestock mobility in RVFV epidemiology in The Gambia and the need for targeted control strategies that might include, for example, targeted cattle vaccination or application of topical insecticide treatments for transhumant herds.

Introduction

Livestock production is a cornerstone of rural livelihoods across sub-Saharan Africa, contributing to economic stability, socio-cultural cohesion, and food security (Herrero et al., 2013). However, the sector is increasingly threatened by infectious diseases and climate change impacts – ranging from conditions intensifying vector proliferation to those exacerbating water and pasture scarcity (Stanimirova et al., 2019; Thornton et al., 2009). In this context, transhumant and nomadic practices across diverse settings serve as critical resilience strategies (Aryal et al., 2018; Motta et al., 2018), allowing pastoral communities to respond to seasonal resource challenges. In the West African Sahel, these transhumant movements typically lead to resource-rich regions along major rivers where abundance of water and high-quality pasture support large populations of livestock (Bassett & Turner, 2007; Freudenberger & Freudenberger, 1993). Despite their importance, these movements are poorly understood due to their largely unregulated nature and limited documentation (Belkhiria et al., 2019; Erdaw, 2023). The association between livestock movements and the spread of infectious diseases has long been recognised (Fevre et al., 2006; Kim et al., 2021; Prentice et al., 2017), allowing introduction of pathogens to local susceptible populations. The spread of infectious diseases is further exacerbated by the very limited veterinary surveillance and disease prevention services available to nomadic and transhumant pastoralists (Belkhiria et al., 2019; Schelling et al., 2008), who often inhabit remote and hard-to-reach regions.

The epidemiology of Rift Valley fever (RVF), a WHO priority zoonotic disease caused by the Rift Valley fever virus (RVFV) [Family: *Phenuiviridae*], exemplifies the interplay between ecological conditions, vector activity, and livestock mobility (Durand et al., 2020; Walsh et al., 2017). RVFV is primarily transmitted by mosquitoes, with competent species identified across the *Aedes, Culex,* and *Anopheles* genera (Linthicum et al., 2016; Lumley et al., 2017). In addition to vector transmission, both human and ruminant infections can also occur through direct contact with infectious tissues, and fluids such as blood and milk during outbreaks (Gerken et al., 2022). Outbreaks in sheep and goats are characterised by devastating abortions and neonatal mortalities (Ksiazek et al., 1989; Munyua et al., 2010). Although cattle and camels are generally less susceptible to RVFV infection compared to sheep (Nair et al., 2023), the movements of asymptomatic but infectious cattle within and between districts (particularly along cattle corridors that connect communities to grazing and water resources) can transport the virus from areas of active transmission to where no outbreaks have been reported (Bird et al., 2009; Shoemaker et al., 2002; Tumusiime et al., 2023). Human cases range from asymptomatic or mild flu-like symptoms to a severe and sometimes fatal syndrome, with complications including haemorrhagic fever or encephalitis (Hoogstraal et al., 1979; Madani et al., 2003). Since its discovery in Kenya in 1931 (Daubney et al., 1931), RVF has caused repeated outbreaks across sub-Saharan Africa, with notable expansions into Egypt (Hoogstraal et al., 1979), the Arabian Peninsula (Balkhy & Memish, 2003) and Comoros Archipelago (Sissoko et al., 2009), underscoring its global threat potential.

In The Gambia, situated in the Sahelian zone of West Africa, the livestock sector contributes approximately 30% of the agricultural GDP and 10% of the national GDP (Rich et al., 2020). The majority of livestock management practices are traditional, with 98% of livestock-owning households engaged in either pastoral (entirely dependent on livestock production) or semi-pastoral systems (integrating crop husbandry and livestock rearing) (Department of Livestock Services, 2016, pers. comms.). The floodplains of the Gambia river serve as a perennial resource hub, providing water and pasture that attract transhumant herds migrating from the Sudano-Sahelian zone during the prolonged harsh dry season

(November – June). Additionally, riverine ecosystems offer ideal breeding sites for RVFV-competent mosquito species in the *Culex* and *Mansonia* genera (Diallo et al., 2000; Saluzzo et al., 1984; Snow, 1983), supporting year-round mosquito proliferation. The Gambia river floodplains thus present an environment conducive to the spread of both vector-borne and non-vector-borne livestock diseases. RVF cases affecting both humans and livestock were first reported in The Gambia in 2002 (FAO, 2003), however, surveillance remains limited, leaving critical gaps in understanding of its epidemiology. Our recent serological investigation in The Gambia revealed ongoing RVFV transmission within ruminant livestock, and highlighted the importance of transhumant movements to the Gambia river as risk factors for livestock seropositivity (Jarra et al., 2025). These findings underscore the need to explore how livestock movement and ecological variability interact to shape RVF dynamics in The Gambia.

In a disease system where transmission is influenced by the movement of hosts, the incorporation of movement processes between habitats is essential for accurately modelling its dynamics (Merkle et al., 2017). Recent studies have integrated livestock movement patterns into the classical compartmental model framework to enhance our understanding of RVF dynamics. For instance, simulations have explored how nomadic ruminants, which occasionally or seasonally migrate from pond to pond according to the availability of water, sustain endemic RVFV circulation in the Ferlo region of Senegal (Durand et al., 2020). Similarly, Tennant et al. (2021) demonstrated that localised environmental conditions and inter-island livestock movements could sustain RVFV circulation in the Comoros Archipelago, without reintroduction from mainland Africa. Notably, this study used vegetation index as a proxy parameter for mosquito-borne transmission in a data-constraint context and achieved predictions consistent with empirical observations (Tennant et al., 2021). While these studies provide valuable insights, their focus on specific ecological and epidemiological contexts limits their applicability to The Gambia, which presents distinct ecological and livestock movement characteristics.

This study aimed to develop and apply an SIR model to simulate RVF dynamics in the cattle population of The Gambia, with a focus on incorporating seasonal transhumant movement as a key mechanism in explaining observed RVF epidemiology. We used household survey data to reconstruct network connections between cattle herds at water and grazing locations, and estimated seasonally dynamic transmission matrices for distinct ecoclimatic regions based on livestock movements. These transmission matrices were integrated into the model to estimate basic reproduction numbers and simulate temporal trends and infection magnitudes that approximate observed RVF dynamics. Using estimates of the force of infection in specific ecoclimatic regions, we predicted age-dependent seroprevalence, and estimated seropositivity waning rates consistent with observed data. By exploring the behaviour of both deterministic and stochastic versions of the same model, this study sought to provide insight into the intersection of livestock movement and ecological conditions that supports RVF dynamics in The Gambia.

Methods

RVFV seroprevalence and household survey data

The data underpinning this modelling study were derived from a previously published cross-sectional serosurvey, which investigated population-level RVFV seroprevalence patterns and household-level risk factors associated with seropositivity in ruminant livestock in The Gambia (Jarra et al., 2025). Briefly, a cross-sectional sero-epidemiological survey was conducted in 2022, collecting 3,602 serum samples from ruminant livestock (cattle, sheep and goats), linked to questionnaire data from 202 livestock-owning households in 52

villages. Details of the study location and data collection procedures are provided in the
Supplementary Text. Due to the very limited number of small ruminants (sheep and goats)
involved in transhumant movement in our household survey data, we focus this study on the
cattle population. The estimated RVFV-specific IgG seroprevalence in cattle was 36.8% (n =
1416, 95% Confidence Interval [CI]: 31.6 – 40.0), with seroprevalence ranging from 0% to
69.8% (95% CI: 47.1 – 92.4) in cattle-owning villages (Jarra et al., 2025). In the present
work, we use this dataset to develop and apply a mechanistic modelling framework aimed at
exploring RVFV transmission dynamics across ecological regions.

Epidemiological setting and cattle population structure

Following Sumaye et al. (2019), we define two regions with distinct ecological and climatic
characteristics (hereafter referred to as ecoclimatic regions or eco-regions) in The Gambia:
the Gambia river ecoclimatic region with its year-round suitability for RVFV transmission
contrasts with the Sahelian ecoclimatic region, which experiences seasonal rainfall and hot
arid dry seasons. Our data revealed that 30% of the cattle-owning households participate in
transhumant movements from the Sahelian ecoclimatic region to the Gambia river during the
preceding dry season (Figure 1). Resident cattle herds are assumed to remain at their
homestead villages year-round. This observation led to the classification of cattle into three
structured subpopulations; M: resident cattle in Sahelian villages, present year-round in this
eco-region without access to the Gambia river. L: resident cattle in villages within the
Gambia river ecoclimatic region, with year-round access to its floodplains. T: transhumant
cattle in Sahelian villages which seasonally move to the Gambia river eco-region during the
dry season (November - June) and return at the start of the wet season (July - October).

144
Figure 1: *Spatial range of transhumant cattle movements in The Gambia identified from the*
household survey. Movements originated from study villages (represented as red squares)
and extended to destination villages, which are either other study villages or villages not
selected for this study (represented as black triangles). This study was conducted during the
dry season, and all transhumant herds were sampled at their destination villages. Distinct
directional movements between homestead villages and their respective destinations are
illustrated with blue arrows. The Gambia river is depicted as the meandering white line
running the length of the country. The map was generated using the igraph package in R.

The observed RVFV IgG seroprevalence in each of the three populations was estimated as
M: 30% (95% CI: 27 – 34), L: 36% (95% CI: 28 – 45) and T: 45% (95% CI: 41 – 48). A 2016
national livestock census of The Gambia estimated the total cattle population at 292,837
(Department of Livestock Services, 2016, pers. comms.). The livestock census includes
complementary data on livestock management practices and grazing patterns. These data
were used to disaggregate the total cattle population across the country into Sahelian

resident (M), transhumant (T), and river resident (L) subpopulations, providing the basis for
the subpopulation estimates in Table 1.

Movement network analysis

We conducted a descriptive movement network analysis to characterise herd mobility
patterns and connectivity within and across eco-climatic regions The household herd served
as the mapping unit to capture both daily local movements around homestead villages and
seasonal transhumant migrations. Herds within a village typically share grazing and water
resources and were assumed to mix homogeneously with those in neighbouring villages
through use of shared resources. Seasonal transhumant movements established
epidemiological links between otherwise distant herds (Figure 1). To represent these
movement processes, we constructed an undirected network in which nodes corresponded
to herds and edges linked herds that shared grazing or water locations. For each node, we
calculated normalised centrality measures (degree, betweenness, and eigenvector
centrality) to quantify relative influence within the network (see Supplementary Table S1).
Edge-level geographic distances were estimated using the mean Euclidean distance
between connected herds. We then used Spearman's rank correlations to examine
associations between network metrics, geographic distance, and herd-specific RVFV
seropositivity. This network analysis was used to contextualise movement structure but was
not incorporated directly into the transmission model, which was parameterised at the eco-
region level to capture broader-scale transmission dynamics.

General overview of the deterministic model

To better understand and simulate the dynamics of RVFV transmission among the cattle
population in The Gambia, we developed a deterministic susceptible-infectious-recovered
(SIR) model that incorporated data from the 2022 survey and accounted for transhumant
cattle movements across the two different ecoclimatic $\text{regions } i \ (i \in \{$Gambia river eco-
region, Sahelian eco-region$\})$ The model operated in continuous time with daily observations
and incorporated matrices reflecting cattle movements and region-specific transmission
parameters, under the assumption of random mixing among varyingly combined cattle
subpopulations within each eco-region. The model was required to track the three cattle
subpopulations (M, L, and T) as they fuse and fission over the wet and dry seasons in the
different eco-regions. Figure 2 represents the three distinct processes in the model:
seasonal cattle movements (arrival and departure of transhumant cattle herds), the
population dynamics (birth and death of cattle), and the RVFV dynamics (infection and
transmission) in each eco-region.

Entry and exit of cattle through trade or gifting from outside the system were not considered.
Each structured cattle subpopulation j ($j \in \{$M, L, T$\}$) was divided into number of susceptible
(S_j), infectious (I_j), and recovered (R_j) compartments. The processes driving population
dynamics were per-capita birth (b) and natural death (μ). Susceptible cattle became
exposed to infection at a transmission rate $\beta_{i,season}$ that depends on the eco-region (i) and
the season (Figure 2B). Infectious cattle recovered from RVF at rate δ or died of the disease
at mortality rate γ. Birth and natural death were modelled as constant rates to maintain a
stable population within each eco-region, reflecting observed herd size in The Gambia over
the past 20 years. This approach assumes that RVF does not cause substantial change in
cattle numbers under the transmission conditions in The Gambia. In addition, we did not
include RVFV-induced abortion in our model. However, the model accounted for temporal

variations in population sizes due to transhumant movements, and explicitly tracks seasonal
population changes in each region:

- **Sahelian eco-region** (s)**:** During the wet season, both resident (M) and transhumant (T) cattle are combined ($N_{s,wet}(t) = T + M$). Only resident cattle (M) remain during the dry season ($N_{s,dry}(t) = M$).
- **Gambia river eco-region** (r)**:** During the wet season, only the resident cattle (L) are present ($N_{r,wet}(t) = L$), while transhumant cattle (T) join during the dry season ($N_{r,dry}(t) = L + T$).

We modelled the RVFV transmission rate ($\beta_{i,season}$) to account for both mosquito-borne, and
direct transmission via contact with highly infectious abortion (calving) products. Cattle of all
age groups were assumed to participate in the seasonal movement between the two eco-
regions, a practice consistently observed among transhumant households over the years.
Only movement of the transhumant subpopulation (T), serving as the primary
epidemiological link between the resident cattle in the Sahelian (M) and those in the Gambia
river eco-region (L), was considered. This necessitated changing the RVFV transmission
matrices (Figure 2B) between wet and dry seasons to accurately reflect the effects of
seasonal movement on RVF transmission dynamics.

221
Figure 2: *Epidemiological model of RVFV transmission and infection dynamics among the*
cattle subpopulations in The Gambia. ***A.*** *Schematic representation of ecoclimatic region and*
seasonal combinations that influences RVFV transmission between the Sahelian areas and
the Gambia river eco-region. ***B.*** *The transmission matrices that determined possible RVFV*
transmissions between the subpopulations during each season. ***C.*** *Transmission framework*
of the within eco-region RVFV transmission, parameters are defined within the main text.

Estimation of model parameters

We estimated basic reproduction numbers ($R_{0,i}$) – defined as the number of secondary
infections caused by one infectious cattle in an entirely susceptible population – separately

for each ecoclimatic region, using a standard formulation adapted from Anderson and May
([1991](#)). This estimation was based on the proportion of the total cattle population in each
eco-region that is susceptible (s^*_i), assuming homogenous mixing of the cattle
subpopulations. Preliminary estimates of season-specific transmission parameters, $\beta_{r,wet}$
and $\beta_{r,dry}$, were then calculated as the remaining unknown that can be solved for from a
standard expression for $R_{0,i}$ derived from the SIR model (see Supplementary Text). RVFV
transmission in the West African Sahel is considered to be negligible during the prolonged
dry season ([Chevalier et al., 2004](#)), and therefore $\beta_{s,dry}$ was set to zero. To align the wet
season transmission in the Sahelian eco-region $(\beta_{s,wet})$ and the potential for year-round
transmission in the Gambia river eco-region, a scaling factor ψ was applied to relate the two
transmission regimes. Specifically, $\beta_{s,wet}$ was defined as $\frac{(\beta_{r,wet} + \beta_{r,dry})}{\psi}$. This formulation
assumes that the annual transmission in the Sahel is predominantly seasonal and confined
to the shorter wet season, whereas transmission in the river eco-region occurs across both
wet and dry seasons due to persistent suitability for mosquito breeding. The scaling factor
was initially set to 2, due to the absence of an established value in the literature.

Preliminary estimates of four key parameters (b, μ, δ and γ) were obtained from existing
literature (Table 1). A final re-estimation of all eight parameters was conducted by fitting the
SIR model to the observed seroprevalence using a simulation-based Approximate Bayesian
Computation (ABC) framework. Informative priors for each parameter were defined as
triangular distributions, with modes based on the preliminary estimates with upper and lower
bounds set to $\pm70\%$ of these estimates. The rejection method of the ABC algorithm was
implemented in the abc package in R [version 2.2.1] ([Csillery et al., 2012](#)). The priors were
used to generate 500,000 randomly parameterised models, and a snapshot RVFV
seroprevalence (summary statistics) was predicted for each model at the end of each
simulation. These predicted seroprevalence values were then compared to the observed
RVFV seroprevalence for each cattle subpopulation. Sets of proposed parameters were
accepted for the 0.2% best fitting models (as determined by the Euclidean distance between
predicted and observed seroprevalence values for each subpopulation). The final parameter
values were taken as the means of the posterior distributions and used for subsequent
model analysis and simulations.

The RVFV dynamics were mathematically formalised using ordinary differential equations
(ODEs). A full description of the equations is provided in the Supplementary Text. An
epidemic was initiated at t = 0 by introducing one infectious cattle into each previously
RVFV-free subpopulation. The simulation was run for 20 years (representing the
approximate period from the initial RVFV outbreak in The Gambia in 2002 to our serological
survey in 2022). The model incorporated distinct seasonal values of the transmission
parameters ($\beta_{i,season}$) (Figure 2B). The dry season spanned 245 days (days 1 – 245), while
the wet season lasted 120 days (days 246 – 365) of each year. The model iteratively
checked the time variable daily to apply the appropriate seasonal parameters. The ODEs are
implemented and solved in R statistical software [v4.2.2] ([R Core Team, 2022](#)) using the
package deSolve ([Soetaert et al., 2010](#)). All plots are generated using the ggplot2 ([Wickham,](#)
[2009](#)).

Estimating the per capita rate of decay of RVFV seropositivity

The force of infection (FOI; $\lambda_{i,\text{season}}$) represents the per capita rate at which susceptible cattle become infected with RVFV within each eco-climatic region during the wet and dry seasons. In this study, FOI was estimated mechanistically as a function of the infectious cattle population, rather than inferred directly from the serological data. A modified version of our transmission model, in which cattle age was substituted for time, was used to generate predicted age-seroprevalence profiles for each subpopulation within each eco-region under quasi-equilibrium conditions (see Supplementary Text). Given that cattle typically remain infectious for approximately one week, temporal variation in $\lambda_{i,\text{season}}$ was evaluated at weekly intervals over a 10-year period, corresponding to the assumed lifespan of cattle in the model. Disease-induced mortality influences FOI indirectly by reducing the size and duration of the infectious compartment and therefore does not enter explicitly into the FOI formulation.

Cattle that recover from RVFV infection were assigned to an immune seropositive compartment (P_j), and the age-seroprevalence curves predicted were compared with observed data. To adjust for inconsistencies between the predicted and observed curves, the predicted age-seroprevalence in each subpopulation was fitted to the observed seroprevalence by including a decay parameter (π) which described the rate of waning of RVFV seropositivity in recovered cattle over time (Figure 2C). The proportion of immune cattle in which seropositivity decayed over time transitioned to an immune seronegative compartment (D_j) (see Supplementary Text).

The age-seroprevalence model was initialised with a fully susceptible population of newly born calves, aging over a 10-year lifespan, and subjected to seasonal variations in the $\lambda_{i,season}$ observed at conditions of quasi-equilibrium. A preliminary decay rate of 0.005 per immune cattle per week was set, and a final value was estimated by fitting the predicted seroprevalence to the observed data using ABC. Triangular priors were assigned to π, with limits extending $\pm 70\%$ from the preliminary rate. The posterior mean of π was used to calculate the half-life of RVFV seropositivity decay in cattle in The Gambia.

Estimation of basic and seasonal reproduction numbers for the entire system

The basic reproduction number (R_0) for the fully susceptible cattle population in the entire system during the wet and dry season was calculated by numerically solving for the leading eigenvalues of a matrix **Q**, using the next-generation matrix method described by Diekmann et al. ([2010](#)) (Supplementary Text). Additionally, to account for dynamic susceptibility and transmissibility within the cattle population at close to quasi-equilibrium, a seasonal reproduction number (R_{st}) was computed at weekly intervals over the course of the 10-year simulation. At each weekly time step, the model output provided the number of new infected cattle in each partially susceptible subpopulation, which is used to determine R_{st}.

Stochastic modelling and RVFV extinction risk

To complement the deterministic solution, a parallel stochastic simulation was conducted to capture demographic stochasticity arising from the inherent probabilistic dynamics of RVFV transmission. The model employed the tau-leap method ([Gillespie, 2007](#)), in combination with the Gillespie algorithm to improve computational efficiency. Within our model, the expected rate for each event type (e.g., population dynamics, transmission dynamics, and infection dynamics) was calculated based on the current state of the system, and the number of events occurring within a time step ($\tau = 0.1$ simulated day) was then assumed to

be Poisson-distributed. The states of all subpopulations were iteratively updated according to the number of sampled events.

We approximated the extinction risk of RVFV in two contexts: (i) within the transhumant subpopulation (T) returning to the Sahelian eco-region at the start of each wet season (stochastic fade-out/local extinction), and (ii) across all three subpopulations (system-wide extinction). For this purpose, 1,000 independent stochastic realisations of the model were conducted, to ensure statistical robustness. RVFV extinction risks were quantified by recording the frequencies at which the number of infectious cattle declined to zero within the transhumant subpopulation (Supplementary Text) and across the entire system. It is important to note that the impact of environmental stochasticity was not considered in this analysis.

Table 1: Preliminary parameter values for the SIR model.

Parameter description	Notation	Value	Unit	Reference
Per capita birth rate	b	0.00215	day^{-1}	Gachohi et al., 2016
Per capita natural death rate	μ	0.0016	day^{-1}	Nicholas et al., 2014
RVF specific mortality rate	γ	0.075	day^{-1}	Gachohi et al., 2016
RVF Recovery rate	δ	1/8	day^{-1}	Durand et al., 2020
Scaling factor	ψ	2	–	assumed
Decay rate of RVFV seropositivity	π	0.005	week^{-1}	assumed
Total M population	N_M	177,004		2016 Livestock Census
Total L population	N_L	80,433		2016 Livestock Census
Total T population	N_T	35,400		2016 Livestock Census

Elasticity Analyses

We evaluated the sensitivity of the predicted RVFV seroprevalence in each of the three subpopulations to individual variations of the eight posterior parameter values: $b, \mu, \gamma, \delta, \psi, \beta_{s,wet}, \beta_{r,wet}, \beta_{r,dry}$. Each parameter was independently sampled 1,000 times from a uniform distribution, constrained within $\pm 20\%$ of its mean posterior value, while the remaining seven parameters are held at their respective mean posterior values. For each modified parameter set, a new predicted seroprevalence was estimated for each subpopulation. The percentage change in predicted seroprevalence was calculated as the difference between the predicted seroprevalence under the original posterior means and the predicted seroprevalence obtained with the modified parameter value. Likewise, the percentage change in parameter value was calculated as the deviation of the randomly sampled parameter value from its mean posterior value. The elasticity was defined as the percentage change in the seroprevalence (the response variable) per unit percentage change in the parameter value (the explanatory variable) and estimated using a general linear model (GLM). Additionally, a Loess smoothing function was applied to evaluate the adequacy of the linear approximation and to identify potential nonlinear trends.

Results

Household herd connectivity

The network analysis revealed variations in between-herd connectivity through shared water and grazing locations (Supplementary Figure S2). Centrality measures were disproportionately influenced by a small subset of transhumant herds with relatively high centrality values, highlighting their prominent role within the network (Figure 3). However, these connections show no significant correlation with herd-specific RVFV seropositivity (Supplementary Figure S3).

Figure 3: *Distribution of degree, normalised betweenness and eigenvector centrality values based on cross-sectional household survey data on livestock movement. Only a few of households showed high centrality values, highlighting their importance in the network.*

Transhumant movements during the dry season covered considerable distances, with herds directed towards the Gambia river from within the Sahelian eco-region. The pairwise geographic distances between connected herds from different study villages ranged from 3.9 to 81.4 km (median 13.5 km). A weak negative correlation (Spearman's R = -0.073, p < 0.05) was observed between the geographic distance of connected herds and their mean RVFV seroprevalence (Supplementary Figure S4), as might be expected from a transmission process influenced by both shared environmental factors and livestock management practices.

Estimated model parameters

The posterior distributions of the eight estimated parameters, derived through the Approximate Bayesian framework, are shown in Supplementary Figure S5. The mean posterior values and their 95% credible intervals (CrI) are summarised in Table 2. These posterior means are used in all subsequent simulations.

371 Table 2: Mean posterior values and 95% CrI of the estimated parameters.

372 * (infected·susceptible·day)$^{-1}$

Parameter description	Notation	Unit	Mean value	95% CrI
Per capita birth rate	b	day^{-1}	0.0022	0.0012 – 0.0033
Per capita natural death rate	μ	day^{-1}	0.0017	0.0010 – 0.0023
RVF specific mortality rate	γ	day^{-1}	0.0758	0.0358 – 0.1868
RVF Recovery rate	δ	day^{-1}	0.1161	0.0500 – 0.1917
Scaling factor	ψ	-	2.0508	0.9224 – 3.1282
Wet season transmission parameter in the Sahelian eco-region	$\beta_{s,wet}$	*	1.4×10^{-6}	6.0×10^{-7} – 2.2×10^{-6}
Wet season transmission parameter in the Gambia river eco-region	$\beta_{r,wet}$	*	3.7×10^{-6}	1.9×10^{-6} – 5.3×10^{-6}
Dry season transmission parameter in the Gambia river eco-region	$\beta_{r,dry}$	*	3.1×10^{-6}	1.7×10^{-6} – 4.4×10^{-6}

373 Seasonal and regional RVF dynamics

We simulated the timing and magnitude of RVFV infections across three structured cattle
subpopulations, which seasonally occupy two eco-regions with distinct ecological
characteristics. At the end of the 20-year simulations, the model remains in a transient phase
without fully attaining a quasi-equilibrium state (Figure 3). The results exhibited an oscillatory
pattern, with recurrent spikes in the proportion of infectious cattle aligning with the wet
season. In the Sahelian eco-region, infections peaked at approximately 5% of the cattle
population between the 64th and 111th days of the wet season (September – October)
(Figure 3 insert), followed by a secondary phase characterised by sharp decline of infection
during the dry season, accompanied by gradual decrease in herd immunity.

While both ecoclimatic regions showed similar temporal RVFV infection dynamics during the
wet season, simulations indicated sustained low-level transmission in the Gambia river eco-
region throughout the dry season (Figure 3 insert). This continued virus activity suggests a
potential "source-sink" effect, as a small fraction (approximately 1%) of the transhumant
subpopulation (T) returned to the Sahelian eco-region at the start of the wet season while
infectious.

At the end of the deterministic simulation, the maximum number of infectious cattle in the M,
L, and T subpopulations during the wet season were approximately 4,562, 2,467, and 443,
respectively. During the dry season, the minimum number of infectious cattle remaining in
the L and T subpopulations within the Gambia river eco-region were approximately 37 and 4,
respectively. By simulating seasonal transmission, and incorporating cattle movement
matrices, our deterministic model provided a close approximation to the RVFV infection
fluctuations demonstrated by the stochastic model (Figure 3).

Figure 3: *Time-series of the proportion of infectious cattle in the three subpopulations across the two eco-regions.* **Main:** *The simulated RVF infection dynamics from the deterministic model (solid lines) together with twenty realisations of the stochastic model (dashed lines).* **Insert:** *A magnified view highlighting the seasonal dynamics of RVFV transmission focused on the last two years of the 20-year deterministic simulation in the cattle subpopulations, highlighting finer seasonal variations (wet season = beige; dry season = cyan). The proportion of infectious cattle peaked at the latter part of the wet season, but infections quickly disappeared in the dry season in the Sahelian eco-region. Once transhumant herds begin to arrive in the Gambia river region, infections are predicted to rise. I = infectious cattle in each subpopulation. A square root transformation was applied to the y-axis for visualisation purposes using coord_trans(y = "sqrt"), while the data remained in its original scale.*

The stochastic simulation indicated that RVFV infection undergoes periodic demographic fade-outs (local extinctions) in the transhumant (T) subpopulation, occurring in 73.8% of realisations (738 of 1,000) during the dry season in the Gambia river eco-region. Each of these realisations predicted between one and five distinct local extinction events (Supplementary Figure S6). Despite these fade-outs, RVFV infection persisted within the broader river eco-region through continued transmission in the river resident (L) subpopulation, enabling recurrent re-infection of the T subpopulation in subsequent years. Across all 1,000 stochastic realisations, a total of 1,363 local extinction events were recorded in the T subpopulation, with a mean transmission duration of 13.9 years. This corresponded to an estimated 9.8% probability of local extinction per dry season within the river eco-region. Following these local extinctions, the annual re-introduction of RVFV into the Sahelian eco-region at the start of the wet season, mediated by the T subpopulation returning from the Gambia river eco-region, was predicted to occur at a high rate. Finally, complete extinction of RVFV across the entire system occurred in 40.2% of realisations (402 of 1,000) before the end of the 20-year simulation period.

Our results showed that, in most realisations, the transhumant subpopulation experienced local extinctions within the first 1,000 days, coinciding with the initial transient phase of the RVF dynamics – big outbreaks within a fully susceptible subpopulation. As the system gradually converged on a quasi-equilibrium state, the probability of RVFV persistence in the T subpopulation stabilised (Figure 4A). In contrast, RVFV exhibited a higher initial

429 persistence probability within the entire system, with most system-wide extinctions occurring
430 beyond the 5,000th day of the simulation as infection dynamics neared a quasi-equilibrium
431 state (Figure 4B).

432
Figure 4: *Extinction rate of RVFV over time (red line with 95% CI – grey ribbon), in the*
*transhumant subpopulation **(A)** and the entire cattle population within the system **(B)** based*
on 1,000 stochastic realisations, illustrating differences in the timing of local and system-
wide extinctions, respectively. Most of the local extinction occurred shortly after RVFV
*introduction into a fully susceptible population in the T subpopulation. **Note:** the timing of the*
local extinction in the T subpopulation depicted here was the first extinction event within this
subpopulation, although re-infection occurs when the subpopulation returned to the river.

Estimated R_{st} and $\lambda_{i,season}$

The R_0 computed for the entire system assuming a fully susceptible cattle population was
3.15 during the wet season and 1.33 during the dry season. The seasonality of transmission
was further illustrated by variations in the R_{st} value (Figure 5), which revealed peaks
($R_{st} > 1$) during the wet season across the entire system, aligning with the amplification of
infection as epidemiological conditions are assumed to be favourable for transmission. In the
latter part of the dry season, $R_{st} > 1$ values were also observed as the susceptible
population recovered, supporting the localised persistence of transmission in the Gambia
river eco-region (Figure 5). Table 3 summarises R_{st} and $\lambda_{i,season}$ values across both eco-
regions and seasons, providing insights into the RVFV dynamics across temporal and spatial
scales.

451
Figure 5: *The full 10-year simulation of the weekly R_{st} (green), and force of infection,*
$\lambda_{region,season}$, in each eco-region (blue = Sahelian, red = Gambia river eco-region). Shaded

areas correspond to the seasons (wet season = beige; dry season = cyan) in the last two
years of the simulation. A square root transformation was applied to the y-axis for
visualization purposes.

Table 3: Summary of estimated seasonal reproduction number (R_{st}) and regio-specific force
of infection (λ_i)

Seasons	Mean R_{st}	Maximum R_{st}	Minimum R_{st}	Avg. weeks $R_{st} > 1$
Wet	1.15	1.40	0.91	9 / 17
Dry	1.02	1.11	0.87	23 / 35

Seasons	Region	Mean λ_i	Maximum λ_i	Minimum λ_i
Wet	Sahelian	0.008	0.05	0.0001
	river	0.003	0.005	0.0002
Dry	Sahelian	–	–	–
	river	0.002	0.018	0.0001

Predicted age-seroprevalence curve

Here, we fitted a model using the weekly $\lambda_{i,season}$ values estimated for each ecoclimatic
region at conditions of near quasi-equilibrium to explore the observed trend in age-
seroprevalence. Age was used as a proxy for cumulative exposure duration, consistent with
prior methodologies (Salje et al., 2016). A cohort of newly born calves was subjected to the
specific $\lambda_{i,season}$ associated with the eco-region they occupy across their lifespan to estimate
the cumulative exposure to RVFV in the absence of waning seropositivity. In the absence of
any waning of seropositivity, the model substantially overestimated seroprevalence,
predicting values of approximately M: 63%, L: 64% and T: 84% after 10-years (Figure 6A).
These predicted seroprevalence values were then re-estimated accounting for potential
waning of seropositivity in recovered cattle, enabling reconstruction of the age-stratified
seroprevalence profile observed in The Gambia (Figure 6B). The posterior distribution of the
rate of decay of RVFV seropositivity (π) yielded a mean value of 0.00217 per week (95% CrI:
0.00216 – 0.00219). This decay rate corresponded to a seropositivity half-life of
approximately 319 weeks (~6 years). These results suggests that the observed
seroprevalence was influenced by gradual loss of seropositivity over time in immune cattle.

*Figure 6: Predicted RVFV seroprevalence in each population subject to FOI during the wet and dry seasons. **A:** Cumulative growth of RVFV immunity over a 10-year period, representing the hypothetical lifespan of cattle, with no consideration for decay of RVFV seropositivity ($\pi = 0$). **B:** Proportion of immune cattle after introducing a seropositivity decay parameter ($\pi = 0.0035$), aligning the predicted seroprevalence with observed data. Observed seroprevalence across different age classes in the three structured cattle populations in The Gambia is shown as dots, with 95% confidence intervals (CI) as error bars. P: Cattle exposed to RVFV infection and recovered, assumed to be seropositive.*

Elasticity Analysis

A 1% change in the mean posterior value of each of the eight parameters resulted in a corresponding percentage change in predicted seroprevalence of each subpopulation (Table 4). The analysis indicated that the per-capita birth (b) and natural death (μ) rates exerted the strongest influence on predicted seroprevalence (Table 4). In contrast, parameters related to transmission ($\beta_{s,wet}$, $\beta_{r,wet}$, $\beta_{r,dry}$) and infection dynamics (τ, γ and δ) exhibited relatively lower elasticities. Additionally, comparisons between GLM and Loess fits suggested that the relationship between the percentage change in the mean posterior parameter values and the corresponding percentage change in predicted seroprevalence was well-approximated as linear (Supplementary Figure S7).

496   Table 4: The elasticity for all eight parameters in the GLM. The parameters with the largest
497   coefficient are in shown in bold.

Parameter	Subpopulation M		Subpopulation L		Subpopulation T	
	Intercept	Coefficient	Intercept	Coefficient	Intercept	Coefficient
$\beta_{s,wet}$	-0.004	0.140	-0.005	-0.027	0.008	-0.032
$\beta_{r,wet}$	-0.014	0.018	0.064	-1.023	0.059	-1.485
$\beta_{r,dry}$	-0.028	0.180	0.074	1.010	0.056	1.604
μ	-0.104	-3.965	0.151	-4.731	0.091	-4.251
b	-0.154	3.603	0.108	4.784	0.059	4.230
γ	-0.001	-1.016	0.055	-1.735	0.057	-1.875
δ	-0.019	0.874	0.059	1.500	0.073	1.571
τ	-0.004	0.140	-0.005	-0.027	0.008	-0.032

Discussion

Previous modelling studies have significantly advanced our understanding of RVF epidemiology, particularly regarding the disease dynamics in ruminant hosts and mosquito vectors. These models have informed the development of predictive tools, surveillance frameworks, control interventions, as well as the evaluation of their impact (Anyamba et al., 2006; Gachohi et al., 2016; Gaff et al., 2007; Tennant et al., 2021). Despite these advancements, the persistence of RVFV in the Sahelian zone, where unfavourable dry season conditions make permanent RVFV circulation unlikely, remains poorly understood (Durand et al., 2020; Lumley et al., 2017). Increasing evidence suggests that livestock movements play a critical role in influencing RVF epidemiology in this zone (Belkhiria et al., 2019; Jarra et al., 2025). In this context, risk exposure to RVFV is primarily driven by movements into regions with high environmental suitability for RVFV-competent vectors in search of water and pasture, rather than transmission through direct livestock-to-livestock contact.

To investigate this hypothesis, we developed deterministic and stochastic versions of the same model, parameterised using seroprevalence data collected in 2022 in The Gambia, to stimulate RVF dynamics. Our study focuses on the transhumant movement of cattle from the Sahelian ecoclimatic region of the country to the floodplains of the Gambia river as a potential mechanism accounting for the observed RVF dynamics among three cattle subpopulations. Key findings from our study include: (i) seasonal infection patterns peaking at approximately 3% and 1% of cattle in the wet and dry seasons, respectively, in the Gambia river eco-region, whereas infections peaked at approximately 5% in the wet season and disappeared in the dry season in the Sahelian region; (ii) local extinction and re-infection dynamics in the transhumant (T) subpopulation predicted by our stochastic model, with a high risk (90.2%) of re-introducing RVFV into their homestead villages in the Sahelian eco-region; (iii) the R_0 for the entire system was estimated as 3.15 during the wet season and 1.33 during the dry season, while the mean seasonal reproduction number (R_{st}) in both seasons was >1; (iv) the age-seroprevalence data in combination with the predicted FOI in each eco-region implied a gradual decay in seropositivity in recovered cattle over time.

RVFV inter-epidemic transmission dynamics

Our modelling approach incorporates transhumant cattle movement, specific eco-regional transmission rates and ecological assumptions. Based on predictions that mosquito-borne transmission is responsible for 84% of RVFV infections in cattle (Nicolas et al., 2014), we pre-suppose the existence of an active mosquito population in the Gambia river eco-region (Snow, 1983) and show that RVFV infections can persist year-round therein. Despite the relatively low proportion of infectious cattle across the three structured subpopulations, RVFV transmission could be sustained in an endemic state over an extended period. Our simulations predicted there is a 90.2% chance the transhumant subpopulation returning to their homestead villages at the start of the wet season would include a small proportion of infectious cattle (approximately 1% of the subpopulation). At the start and end of transhumance, GPS-tracked cattle herds in Cameroon exhibit active movement behaviour at an average daylight speed of 3 – 4 km/hour (Motta et al., 2018), a pattern corroborated by survey data from transhumant households in The Gambia. Given that RVFV incubation and viraemia last approximately seven to eight days, virus re-introduction to the Sahelian eco-region through the returning transhumant herds is plausible, posing a recurrent transmission risk. However, herd immunity among the Sahelian resident cattle population in our simulation prevents annual large-scale outbreaks.

The possibility of RVFV re-introduction between the two eco-regions is consistent with explanations given for the dissemination of the virus during outbreaks within and across

countries through movement of potentially viraemic but asymptomatic ruminant livestock
(Abdo-Salem et al., 2011; Shoemaker et al., 2002). In addition, similar patterns have been
observed in northern Senegal (Cecilia et al., 2020; Durand et al., 2020) and Tanzania
(Sumaye et al., 2013), where movement of infectious livestock between regions with
favourable ecological conditions is thought to maintain RVFV endemicity. Our simulation
results of the timing and magnitude of infections are supported by evidence showing RVFV
amplification during the wet season, with peak circulation occurring at the latter part of the
season in Senegal (Durand et al., 2020). The R_{st} reflected this, with mean R_{st} values greater
than one in the first half of the wet season (9 of 17 weeks) in all eco-regions, and in over
two-thirds of the dry season (23 of 35 weeks) in the Gambia river eco-region. Earlier studies
conducted during an inter-epidemic period in Mayotte estimated seasonal reproduction
numbers broadly consistent with those estimated by our study, including Metras et al.,
(2017) (maximum R_{st} = 2.19) and Tennant et al., (2021) (maximum R_{st} = 2.77).

Traditionally, RVFV epidemiology has been characterised as having episodic outbreaks
separated by long inter-epidemic periods (Murithi et al., 2011; Sindato et al., 2014).
However, growing evidence across sub-Saharan Africa suggests that RVFV dynamics do not
conform to a single archetype but instead occur along a continuum (Rostal et al., 2025),
ranging from epidemic-prone systems where viral persistence is low (Sindato et al., 2014) to
hyperendemic systems with sustained, moderate transmission without necessarily resulting
in substantial outbreaks (Jarra et al., 2025; van den Bergh et al., 2019). Within this
conceptual framework, the stable seasonal dynamics predicted by our model likely position
The Gambia toward the hyperendemic end of this spectrum, where ecological conditions,
RVFV-competent vectors (potentially including transovarial transmission) and seasonal
rainfall maintain transmission without requiring repeated viral re-introduction. In addition,
livestock demography, movement patterns, and acquired immunity may further stabilise
these dynamics thereby supporting long-term circulation without substantial outbreaks.

While our model incorporates the possibility of stochastic extinction, its structure assumes
potential for continuous local transmission and therefore tends toward an endemic quasi-
equilibrium rather than higher amplitude epidemic cycles. This highlights the plausibility of
ecologically sustained RVFV circulation in cattle in The Gambia with high seroprevalence
despite the absence of substantive outbreaks. These findings are consistent with patterns
observed in other hyperendemic systems such as northeastern KwaZulu-Natal, South Africa
(van den Bergh et al., 2019). Nonetheless, longitudinal serological and entomological studies
are needed to determine whether the apparent stability reflects true endemic persistence.

Age-seroprevalence curve and seropositivity decay
Acquired immunity following natural infection with many communicable diseases is often
imperfect, with immunity potentially waning over time, offering partial protection or failing
entirely (Le et al., 2021). Although acquired immunity to RVFV is often considered long-
lasting (Swanepoel & Coetzer., 2004; Wilson, 1994), there is a lack of empirical data
regarding the nature and longevity of IgG seropositivity following natural infection in ruminant
livestock. In a long-term cohort study, Wright et al. (2020) observed that RVFV-specific IgG
and neutralising antibodies remained detectable in adult humans for up to eleven years post-
infection, although antibody titers progressively declined. This finding has important
implications for the interpretation of serological data: while antibodies may be present, the
ability to detect them (i.e. seropositivity) may decline over time. Serological diagnostics
reliant on detecting specific antibody thresholds may classify immune individuals as
seronegative, potentially underestimating the true extent of population immunity. This
implication has been observed in other viral infections. For instance, in a 13-year longitudinal
study of 34 healthcare workers infected with SARS-CoV, the 50th percentile exhibited anti-

nucleocapsid IgG titers below the assay detection limit by year nine, and 75th percentile fell
below the limit by year thirteen (Guo et al., 2020). These observations reinforce the
importance of considering seropositivity decay when interpreting long-term antibody
persistence.

Our sero-epidemiological study in The Gambia revealed an age-seroprevalence pattern in
cattle that is characterised by high IgG seropositivity during early calfhood, but that quickly
plateaued or declined with advancing age (Jarra et al., 2025). To reconcile the discrepancies
between this observed age-seroprevalence pattern and the predictions based on our FOI
estimates from each eco-region, we incorporated a seropositivity decay parameter into the
model. Specifically, RVFV-exposed cattle in which seropositivity decayed over time
transitioned into an immune but seronegative compartment. This adjustment is supported by
evidence that cellular immunity (T cell-mediated responses) can confer durable protection
even in the presence of substantially low RVFV-specific antibody levels following both
natural RVFV infection in humans (Wright et al., 2020) and wildtype RVFV challenge in
vaccinated mice (Doyle et al., 2022). Therefore, our interpretation of antibody decay reflects
waning antibody detectability rather than true loss of immunity.

Although earlier RVF seroprevalence-based modelling studies have not explicitly
incorporated or contextualized the kinetics of IgG decay, Le et al. (2021) provides a valuable
theoretical framework of imperfect infection-derived immunity by outlining seven distinct
models. These models span between the two classical extremes: the "perfect" immunity
assumed in SEIR frameworks (recovered individuals are permanently protected) and the
"none at all" immunity characteristic of SEIS models (individuals become susceptible again
after recovery). Each of these immune decay models produced unique implications for the
overall disease dynamics within a population (Le et al., 2021). Although a SEIS or SIRS
framework could theoretically have allowed recovered cattle to revert to susceptibility,
stronger empirical evidence on the duration and durability of natural RVFV antibody
responses in ruminant livestock and genuine loss of acquired immune is required to justify
this assumption. Future studies incorporating longitudinal serological data could refine
parameterisation of our model to explicitly quantify seropositivity decay and its implications
for RVFV transmission dynamics, particularly in systems where seropositivity may wane or
become undetectable over time despite persistent protection.

Implications of the model and future enhancements

While this study provides important insight into RVFV dynamics in The Gambia, several data
constraints should be considered. The most influential source of uncertainty arises from
limited resolution in the household survey data, particularly gaps in spatial data coverage
and the coarse temporal information on cattle movements. As a result, the extent of
transhumant movements the model relies on may not fully represent the diversity of
movements across the country. These limitations may have implications for our conclusions,
and may lead to under- or over-estimation of periods of heightened transmission risk.
Despite these uncertainties, the finding of low-level, spatially diffuse circulation is robust,
even though the overall network structure observed suggests a fragmented population with
modest but epidemiologically meaningful transhumance distances for The Gambia livestock
management systems. Nonetheless, future work incorporating high-resolution movement
data, herd-level demographic monitoring, and spatially explicit network models would better
capture heterogeneity in contact patterns and improve the accuracy of the predicted RVFV
dynamics.

Although RVF can cause substantial abortion and neonatal losses during large outbreaks,
our model predicts only low-level endemic circulation in The Gambia. The dampening effect

of managed herd demography – through routine sales, replacements, and movement –
supports our assumption of near-constant births, consistent with other models of endemic or
inter-epidemic RVF dynamics (Nicolas et al., 2014). However, this assumption should be
revisited in future analyses that incorporate higher-resolution demographic data.

Conclusion

This study contributes to our understanding RVFV epidemiology in The Gambia by
integrating transhumant cattle movement into a modelling framework that accounted for eco-
regional transmission dynamics. Our findings highlighted seasonal transmission patterns,
with year-round infections predicted in the Gambia river eco-region reinforcing the role of
ecological suitability in sustaining RVFV. While transhumance played a role in shaping RVFV
transmission through a high risk of RVFV re-introduction, herd immunity limited large-scale
outbreaks in the Sahelian eco-region. The predicted seropositivity decay underscored the
importance of considering waning of antibody titers in serological assessments. Future
refinements incorporating high-resolution livestock movement data and environmental
variability will enhance predictive accuracy. There is need for strategies informed by
transhumance practices in managing RVF, particularly in remote livestock production
communities in The Gambia. Understanding livestock owners' perceptions of seasonal
movements, and the determinants of these movements could inform the design of socially
acceptable measures such as targeted vaccination and the strategic or pre-emptive
application of topical acaricides for ruminants during peak vector activity in the wet season or
before transhumant movements into the river floodplain.

Supplementary Information

666 Supplementary Text

Acknowledgements

668 We thank the communities and livestock-owning households for participating in the study.
669 We are also deeply grateful to the livestock assistants of the Department of Livestock
670 Services in The Gambia for their contributions to the field work.

Ethical clearance

The study protocol received ethical approval from the National Agricultural Research Institute
of The Gambia [Approval Reference: NARI/DLS/01/(01)] and the University of Glasgow
School of Veterinary Medicine Research Ethics Committee (Reference: EA47/21). Approval
for interactions with human participant were obtained from The Gambia
Government/MRCG@LSHTM Joint Ethics Committee (Project ID/Review Ref: 26750) and
the University of Glasgow College of Medical, Veterinary and Life Sciences Ethics
Committee (Project No: 200210057).

We adhere to the STROBE Statement–Checklist of items that should be included in reports
of cross-sectional studies.

Consent

Community engagement meetings were held to introduce the study, explain sampling
procedures, and obtain informed consent. Participants provided written consent or a
thumbprint as a substitute in cases of illiteracy. Consent procedures emphasized voluntary
participation, anonymity, and participants' right to withdraw. Collaboration with the local
livestock assistants of the Department of Livestock Services (DLS), who are familiar with and
trusted by the communities, facilitated recruitment of participants. All DLS livestock
assistants received biosafety and ethics training prior to data collection. No human samples
were collected for this study.

Author Contributions

E.J. and D.T.H. conceptualized the study; E.J. acquired the funding; E.J. coordinated and administered the field work; E.J. and D.T.H developed the deterministic and stochastic models; D.E. conducted the network analysis; E.J. wrote the original draft with input from D.T.H. and D.E.; D.T.H. and S.C. supervised the project; All authors reviewed, edited and approved the manuscript.

Conflict of Interest Statement

The authors declare no conflicts of interest.

Data Availability Statement

The codes that support the findings and produce the figures of this manuscript are available on Github: https://github.com/enz-j/Rvfv_dynamics_gambia

Funding

This work was funded by the Foreign, Commonwealth & Development Office of the UK Government through the Commonwealth Scholarship Commission for a PhD studentship for Essa Jarra at University of Glasgow. The Commonwealth Scholarship Commission was not involved in study design, data collection and interpretation, or the decision to submit the work for publication. The opinions expressed in this publication are entirely those of the authors.

References

Abdo-Salem, S., Waret-Szkuta, A., Roger, F., Olive, M. M., Saeed, K., & Chevalier, V. (2011). Risk assessment of the introduction of Rift Valley fever from the Horn of Africa to Yemen via legal trade of small ruminants. *Tropical Animal Health and Production, 43*(2), 471-480. doi:https://doi.org/10.1007/s11250-010-9719-7

Anderson, R. M., & May, R. M. (1991). *Infectious Diseases of Humans: Dynamics and Control*. Oxford: Oxford Univesrity Press.

Anyamba, A., Chretien, J. P., Small, J., Tucker, C. J., & Linthicum, K. J. (2006). Developing global climate anomalies suggest potential disease risks for 2006-2007. *International Journal of Health Geographics, 5*, 60. doi:https://doi.org/10.1186/1476-072X-5-60

Aryal, S., Maraseni, T., Cockfield, G., & de Bruyn, L. L. (2018). *Transhumance, Livestock Mobility and Mutual Benefits Between Crop and Livestock Production* (Vol. 31): Springer.

Balkhy, H. H., & Memish, Z. A. (2003). Rift Valley fever: an uninvited zoonosis in the Arabian peninsula. *International Journal of Antimicrobial Agents, 21*(2), 153-157. doi:https://doi.org/10.1016/s0924-8579(02)00295-9

Bassett, T. J., & Turner, M. D. (2007). Sudden Shift or Migratory Drift? FulBe Herd Movements to the Sudano-Guinean Region of West Africa. *The Journal of Human Ecology, 35*(1), 33–49 doi:https://doi.org/10.1007/s10745-006-9067-4

Belkhiria, J., Lo, M. M., Sow, F., Martinez-Lopez, B., & Chevalier, V. (2019). Application of exponential random graph models to determine nomadic herders' movements in Senegal. *Transboundary and Emerging Diseases, 66*(4), 1642-1652. doi:https://doi.org/10.1111/tbed.13198

Bird, B. H., Ksiazek, T. G., Nichol, S. T., & Maclachlan, N. J. (2009). Rift Valley fever virus. *Journal of American Veterinary Medical Association, 234*(7), 883-893. doi:10.2460/javma.234.7.883

Cecilia, H., Metras, R., Fall, A. G., Lo, M. M., Lancelot, R., & Ezanno, P. (2020). It's risky to wander in September: Modelling the epidemic potential of Rift Valley fever in a Sahelian setting. *Epidemics, 33*, 100409. doi:https://doi.org/10.1016/j.epidem.2020.100409

Chevalier, V., de la Rocque, S., Baldet, T., Vial, L., & Roger, F. (2004). Epidemiological processes involved in the emergence of vector-borne diseases: West Nile fever, Rift Valley fever, Japanese encephalitis and Crimean-Congo haemorrhagic fever. *Scientific and Technical Review, 23*(2), 535-555. doi:https://doi.org/10.20506/rst.23.2.1505

Csillery, K., Francois, O., & Blum, M. G. B. (2012). abc: an R package for approximate Bayesian computation (ABC). *Methods in Ecology and Evolution, 3*(3), 475-479. doi:https://doi.org/10.1111/j.2041-210X.2011.00179.x

Daubney, R., Hudson, J. R., & Garnham, P. C. (1931). Enzootic hepatitis or rift valley fever. An undescribed virus disease of sheep cattle and man from east africa. *The Journal of Pathology and Bacteriology, 34*(4), 545-579. doi:https://doi.org/10.1002/path.1700340418

Department of Livestock Services. (2016). *National Livestock Census Report*. Department of Livestock Services. Ministry of Agriculture. The Gambia.

Diallo, M., Lochouarn, L., Ba, K., Sall, A. A., Mondo, M., Girault, L., & Mathiot, C. (2000). First isolation of the Rift Valley fever virus from Culex poicilipes (Diptera: Culicidae) in nature. *American Journal of Tropical Medicine and Hygiene, 62*(6), 702-704. doi:https://doi.org/10.4269/ajtmh.2000.62.702

Diekmann, O., Heesterbeek, J. A. P., & Roberts, M. G. (2010). The construction of next-generation matrices for compartmental epidemic models. *Journal of The Royal Society Interface, 7*(46), 873-885. doi:https://doi.org/10.1098/rsif.2009.0386

Doyle, J. D., Barbeau, D. J., Cartwright, H. N., & McElroy, A. K. (2022). Immune correlates of protection following Rift Valley fever virus vaccination. *NPJ Vaccines, 7*(1), 129. doi:https://doi.org/10.1038/s41541-022-00551-4

Durand, B., Lo Modou, M., Tran, A., Ba, A., Sow, F., Belkhiria, J., . . . Chevalier, V. (2020). Rift Valley fever in northern Senegal: A modelling approach to analyse the processes underlying virus circulation recurrence. *PLOS Neglected Tropical Diseases, 14*(6), e0008009. doi:https://doi.org/10.1371/journal.pntd.0008009

Erdaw, M. M. (2023). Contribution, prospects and trends of livestock production in sub-Saharan Africa: a review. *International Journal of Agricultural Sustainability, 21*(1), 1-15. doi:https://doi.org/10.1080/14735903.2023.2247776

Fevre, E. M., Bronsvoort, B. M., Hamilton, K. A., & Cleaveland, S. (2006). Animal movements and the spread of infectious diseases. *Trends in Microbiology, 14*(3), 125-131. doi:https://doi.org/10.1016/j.tim.2006.01.004

Food and Agriculture Organisation. (2003). Rift Valley fever monitoring in West Africa in 2002: Early warning message - Rift Valley fever in The Gambia. *EMPRES Transboundary Animal Diseases Bulletin, No 23*, 9-11. Retrieved from https://www.fao.org/4/y4809e/y4809e00.pdf

Freudenberger, M. S., & Freudenberger, K. S. (1993). Pastoralism in Peril: Pressures on Grazing land in Senegal. *IIED Drylands Programme, Pastoral Land Tenure Series No. 4.* Retrieved from https://www.iied.org/sites/default/files/pdfs/migrate/7351IIED.pdf

Gachohi, J. M., Njenga, M. K., Kitala, P., & Bett, B. (2016). Modelling Vaccination Strategies against Rift Valley Fever in Livestock in Kenya. *PLOS Neglected Tropical Diseases, 10*(12), e0005049. doi:https://doi.org/10.1371/journal.pntd.0005049

Gaff, H. D., Hartley, D. M., & Leahy, N. P. (2007). An epidemiological model of Rift Valley fever. *Electronic Journal of Differential Equations*(115), 1-12. Retrieved from https://digitalcommons.odu.edu/biology_fac_pubs/49

Gerken, K. N., LaBeaud, A. D., Mandi, H., L'Azou Jackson, M., Breugelmans, J. G., & King, C. H. (2022). Paving the way for human vaccination against Rift Valley fever virus: A systematic literature review of RVFV epidemiology from 1999 to 2021. *PLOS Neglected Tropical Diseases, 16*(1), e0009852. doi:https://doi.org/10.1371/journal.pntd.0009852

Gillespie, D. T. (2007). Stochastic simulation of chemical kinetics. *Annual Review of Physical Chemistry, 58*, 35-55. doi:https://doi.org/10.1146/annurev.physchem.58.032806.104637

Guo, X., Guo, Z., Duan, C., Chen, Z., Wang, G., Lu, Y., . . . Lu, J. (2020). Long-Term Persistence of IgG Antibodies in SARS-CoV Infected Healthcare Workers. *medRxiv*, 2020.2002.2012.20021386. doi:https://doi.org/10.1101/2020.02.12.20021386

Herrero, M., Grace, D., Njuki, J., Johnson, N., Enahoro, D., Silvestri, S., & Rufino, M. C. (2013). The roles of livestock in developing countries. *Animal, 7 Suppl 1*, 3-18. doi:https://doi.org/10.1017/S1751731112001954

Hoogstraal, H., Meegan, J. M., Khalil, G. M., & Adham, F. K. (1979). The Rift Valley fever epizootic in Egypt 1977-78. 2. Ecological and entomological studies. *Transactions of the Royal Society of Tropical Medicine and Hygiene, 73*(6), 624-629. doi:https://doi.org/10.1016/0035-9203(79)90005-1

Jarra, E., Beyai, Y., Mendy, A., Ceesay, O., Ceesay, A., Daffeh, K., . . . Cleaveland, S. (2025). Sero-epidemiology of Rift Valley fever virus in ruminant livestock in The Gambia. *bioRxiv*(2025.03.14.640388). doi:https://doi.org/10.1101/2025.03.14.640388

Kim, Y., Metras, R., Dommergues, L., Youssouffi, C., Combo, S., Le Godais, G., . . . Fournie, G. (2021). The role of livestock movements in the spread of Rift Valley fever virus in animals and humans in Mayotte, 2018-19. *PLOS Neglected Tropical Diseases, 15*(3), e0009202. doi:https://doi.org/10.1371/journal.pntd.0009202

Ksiazek, T. G., Jouan, A., Meegan, J. M., Le Guenno, B., Wilson, M. L., Peters, C. J., . . . Touray, E. M. (1989). Rift Valley fever among domestic animals in the recent West African outbreak. *Research in Virology, 140*(1), 67-77. doi:https://doi.org/10.1016/s0923-2516(89)80086-x

Le, A., King, A. A., Magpantay, F. M. G., Mesbahi, A., & Rohani, P. (2021). The impact of
infection-derived immunity on disease dynamics. *Journal of Mathematical Biology,*
83(6-7), 61. doi:https://doi.org/10.1007/s00285-021-01681-4
Linthicum, K. J., Britch, S. C., & Anyamba, A. (2016). Rift Valley Fever: An Emerging
Mosquito-Borne Disease. *Annual Review of Entomology, 61,* 395-415.
doi:https://doi.org/10.1146/annurev-ento-010715-023819
Lumley, S., Horton, D. L., Hernandez-Triana, L. L. M., Johnson, N., Fooks, A. R., & Hewson,
R. (2017). Rift Valley fever virus: strategies for maintenance, survival and vertical
transmission in mosquitoes. *Journal of General Virology, 98*(5), 875-887.
doi:https://doi.org/10.1099/jgv.0.000765
Madani, T. A., Al-Mazrou, Y. Y., Al-Jeffri, M. H., Mishkhas, A. A., Al-Rabeah, A. M., Turkistani,
A. M., . . . Shobokshi, O. (2003). Rift Valley fever epidemic in Saudi Arabia:
epidemiological, clinical, and laboratory characteristics. *Clinical infectious diseases,*
37(8), 1084-1092. doi:https://doi.org/10.1086/378747
Merkle, J. A., Cross, P. C., Scurlock, B. M., Cole, E. K., Courtemanch, A. B., Dewey, S. R., &
Kauffman, M. J. (2017). Linking spring phenology with mechanistic models of host
movement to predict disease transmission risk. *Journal of Applied Ecology, 55*(2),
810-819. doi:https://doi.org/10.1111/1365-2664.13022
Metras, R., Fournie, G., Dommergues, L., Camacho, A., Cavalerie, L., Merot, P., . . .
Edmunds, W. J. (2017). Drivers for Rift Valley fever emergence in Mayotte: A
Bayesian modelling approach. *PLOS Neglected Tropical Diseases, 11*(7), e0005767.
doi:https://doi.org/10.1371/journal.pntd.0005767
Motta, P., Porphyre, T., Hamman, S. M., Morgan, K. L., Ngwa, V. N., Tanya, V. N., . . .
Bronsvoort, B. M. (2018). Cattle transhumance and agropastoral nomadic herding
practices in Central Cameroon. *BMC Veterinary Research, 14*(1), 214.
doi:https://doi.org/10.1186/s12917-018-1515-z
Munyua, P., Murithi, R. M., Wainwright, S., Githinji, J., Hightower, A., Mutonga, D., . . .
Njenga, M. K. (2010). Rift Valley fever outbreak in livestock in Kenya, 2006-2007.
American Journal of Tropical Medicine and Hygiene, 83(2 Suppl), 58-64.
doi:https://doi.org/10.4269/ajtmh.2010.09-0292
Murithi, R. M., Munyua, P., Ithondeka, P. M., Macharia, J. M., Hightower, A., Luman, E. T., . .
. Njenga, M. K. (2011). Rift Valley fever in Kenya: history of epizootics and
identification of vulnerable districts. *Epidemiol Infect, 139*(3), 372-380.
doi:10.1017/S0950268810001020
Nair, N., Osterhaus, A., Rimmelzwaan, G. F., & Prajeeth, C. K. (2023). Rift Valley Fever
Virus-Infection, Pathogenesis and Host Immune Responses. *Pathogens, 12*(9).
doi:10.3390/pathogens12091174
Nicolas, G., Chevalier, V., Tantely, L. M., Fontenille, D., & Durand, B. (2014). A spatially
explicit metapopulation model and cattle trade analysis suggests key determinants
for the recurrent circulation of rift valley Fever virus in a pilot area of madagascar
highlands. *PLOS Neglected Tropical Diseases, 8*(12), e3346.
doi:https://doi.org/10.1371/journal.pntd.0003346
Prentice, J. C., Marion, G., Hutchings, M. R., McNeilly, T. N., & Matthews, L. (2017).
Complex responses to movement-based disease control: when livestock trading
helps. *Journal of The Royal Society Interface, 14*(126).
doi:https://doi.org/10.1098/rsif.2016.0531
R Core Team. (2022). R: A Language and Environment for Statistical Computing. *R
Foundation for Statistical Computing.* Retrieved from https://www.R-project.org
Rich, K. M., Bahta, S., Wane, A., Wanyoike, F., & Baltenweck, I. (2020). Initial assessment
report of the Gambian livestock sector: a report to the Small Ruminant Production
Enhancement Project (SRPEP) by the International Livestock Research Institute in
support of the Gambia's livestock master plan. *Nairobi, Kenya: ILRI.* Retrieved from
https://hdl.handle.net/10568/113977
Rostal, M. K., Thompson, P. N., Anyamba, A., Bett, B., Cetre-Sossah, C., Chevalier, V., . . .
Swanepoel, R. (2025). Rift Valley fever epidemiology: shifting the paradigm and

rethinking research priorities. *Lancet Planet Health, 9*(9), 101299. doi:10.1016/j.lanplh.2025.101299

Salje, H., Cauchemez, S., Alera, M. T., Rodriguez-Barraquer, I., Thaisomboonsuk, B., Srikiatkhachorn, A., . . . Yoon, I. K. (2016). Reconstruction of 60 Years of Chikungunya Epidemiology in the Philippines Demonstrates Episodic and Focal Transmission. *The Journal of Infectious Diseases, 213*(4), 604-610. doi:https://doi.org/10.1093/infdis/jiv470

Saluzzo, J. F., Digoutte, J. P., Cornet, M., Baudon, D., Roux, J., & Robert, V. (1984). Isolation of Crimean-Congo haemorrhagic fever and Rift Valley fever viruses in Upper Volta. *Lancet, 1*(8387), 1179. doi:https://doi.org/10.1016/s0140-6736(84)91419-3

Schelling, E., Wyss, K., Diguimbaye, C., Béchir, M., Taleb, M. O., Bonfoh, B., . . . Zinsstag, J. (2008). *Towards Integrated and Adapted Health Services for Nomadic Pastoralists and their Animals: A North–South Partnership*. Dordrecht: Springer.

Shoemaker, T., Boulianne, C., Vincent, M. J., Pezzanite, L., Al-Qahtani, M. M., Al-Mazrou, Y., . . . Nichol, S. T. (2002). Genetic analysis of viruses associated with emergence of Rift Valley fever in Saudi Arabia and Yemen, 2000-01. *Emerging Infectious Diseases, 8*(12), 1415-1420. doi:https://doi.org/10.3201/eid0812.020195

Sindato, C., Karimuribo, E. D., Pfeiffer, D. U., Mboera, L. E., Kivaria, F., Dautu, G., . . . Paweska, J. T. (2014). Spatial and temporal pattern of Rift Valley fever outbreaks in Tanzania; 1930 to 2007. *PLoS One, 9*(2), e88897. doi:10.1371/journal.pone.0088897

Sissoko, D., Giry, C., Gabrie, P., Tarantola, A., Pettinelli, F., Collet, L., . . . Pierre, V. (2009). Rift Valley fever, Mayotte, 2007-2008. *Emerging Infectious Diseases, 15*(4), 568-570. doi:https://doi.org/10.3201/eid1504.081045

Snow, W. F. (1983). Mosquito production and species succession from an area of irrigated rice fields in The Gambia, West Africa. *American Journal of Tropical Medicine and Hygiene, 86*(6), 237-245.

Soetaert, K., Petzoldt, T., & Setzer, R. (2010). Solving Differential Equations in R: Package deSolve. *Journal of Statistical Software, 33*(9), 1-25. doi:https://doi.org/

Stanimirova, R., Arévalo, P., Kaufmann, R. K., Wegner Maus, V., Lesiv, M., Havlík, P., & Friedl, M. A. (2019). Sensitivity of global pasturelands to climate variation. *Earths Future, 7*(12), 1353 -1366. doi:https://doi.org/10.1029/2019EF001316

Sumaye, R. D., Geubbels, E., Mbeyela, E., & Berkvens, D. (2013). Inter-epidemic transmission of Rift Valley fever in livestock in the Kilombero River Valley, Tanzania: a cross-sectional survey. *PLOS Neglected Tropical Diseases, 7*(8), e2356. doi:https://doi.org/10.1371/journal.pntd.0002356

Sumaye, R. D., Jansen, F., Berkvens, D., De Baets, B., Geubels, E., Thiry, E., & Krit, M. (2019). Rift Valley fever: An open-source transmission dynamics simulation model. *PLoS One, 14*(1), e0209929. doi:https://doi.org/10.1371/journal.pone.0209929

Swanepoel, R., & Coetzer., J. A. W. (2004). *Rift Valley fever*. Oxford, UK: Oxford University Press.

Tennant, W. S. D., Cardinale, E., Cetre-Sossah, C., Moutroifi, Y., Le Godais, G., Colombi, D., . . . Metras, R. (2021). Modelling the persistence and control of Rift Valley fever virus in a spatially heterogeneous landscape. *Nature Communications, 12*(1), 5593. doi:https://doi.org/10.1038/s41467-021-25833-8

Thornton, P., van de Steeg, J., Notenbaert, M. H., & Herrero, M. (2009). The impacts of climate change on livestock and livestock systems in developing countries: A review of what we know and what we need to know. *Agricultural Systems, 101*(3), 113-127. doi:https://doi.org/10.1016/j.agsy.2009.05.002

Tumusiime, D., Isingoma, E., Tashoroora, O. B., Ndumu, D. B., Bahati, M., Nantima, N., . . . Bett, B. (2023). Mapping the risk of Rift Valley fever in Uganda using national seroprevalence data from cattle, sheep and goats. *PLoS Negl Trop Dis, 17*(5), e0010482. doi:10.1371/journal.pntd.0010482

van den Bergh, C., Venter, E. H., Swanepoel, R., & Thompson, P. N. (2019). High seroconversion rate to Rift Valley fever virus in cattle and goats in far northern

925   KwaZulu-Natal, South Africa, in the absence of reported outbreaks. *PLoS Negl Trop*
926   *Dis, 13*(5), e0007296. doi:10.1371/journal.pntd.0007296
927 Walsh, M. G., Willem de Smalen, A., & Mor, S. M. (2017). Wetlands, wild Bovidae species
928   richness and sheep density delineate risk of Rift Valley fever outbreaks in the African
929   continent and Arabian Peninsula. *PLOS Neglected Tropical Diseases, 11*(7),
930   e0005756. doi:https://doi.org/10.1371/journal.pntd.0005756
931 Wickham, H. (2009). ggplot2: Elegant Graphics for Data Analysis. Retrieved from
932   https://CRAN.R-project.org/package=ggplot2
933 Wilson, M. L. (1994). Rift Valley fever virus ecology and the epidemiology of disease
934   emergence. *Annals of the New York Academy of Sciences, 740*, 169-180.
935   doi:https://doi.org/10.1111/j.1749-6632.1994.tb19867.x
936 Wright, D., Allen, E. R., Clark, M. H. A., Gitonga, J. N., Karanja, H. K., Hulswit, R. J. G., . . .
937   Warimwe, G. M. (2020). Naturally Acquired Rift Valley Fever Virus Neutralizing
938   Antibodies Predominantly Target the Gn Glycoprotein. *iScience, 23*(11), 101669.
939   doi:https://doi.org/10.1016/j.isci.2020.101669