Diel cycle of lanthanide-dependent methylotrophy by TMED127/Methylaecorales bacteria

in oligotrophic surface seawater

Jennifer B. Glass[1*], Leilani N. Warters[1], Abdulaziz M. Alajlan[1]

[1]School of Earth and Atmospheric Sciences, Georgia Institute of Technology, Atlanta, GA, USA

*Corresponding author: Jennifer.Glass@eas.gatech.edu

Abstract

Methanol, the simplest alcohol, has long been recognized as a key energy and carbon source for

soil and plant-associated bacteria and fungi, and is increasingly recognized as an important

substrate for marine bacteria. Lanthanide-dependent methanol dehydrogenases (encoded by the

gene *xoxF*) are now recognized as the key catalysts for methylotrophy in many environments, yet

the identity of the most transcriptionally active methylotrophs ("Clade X") has remained elusive.

Here we show that "Clade X" methylotrophs belong to the deep-branching alphaproteobacterial

order TMED127, which we propose be renamed 'Methylaecorales'; 'methyl' for 'methylotrophic

metabolism' and 'aecor' for 'ocean surface', as these bacteria are most transcriptionally active

near the sea surface. TMED127/Methylaecorales have small, streamlined genomes (~1.5 Mb)

and appear to be obligate methylotrophs that use the serine cycle for carbon assimilation.

TMED127/Methylaecorales showed a diel pattern of *xoxF5* and glucose dehydrogenase (*gdh*)

transcription, peaking in the late afternoon, in oligotrophic surface water of the Sargasso Sea.

Several other highly transcribed genes of unknown function had no homologs outside of

TMED127/Methylaecorales genomes. Our findings illuminate an overlooked marine

methylotrophic bacterium and predict a diel cycle of methanol production in surface seawater by

an unknown pathway.

Importance

Methanol metabolism is increasingly recognized as an important process in the marine carbon

cycle, yet the identity and metabolism of the microorganisms mediating methylotrophy in the

open ocean have remained unknown. This study reveals that bacteria in the TMED127 order of

Alphaproteobacteria, proposed new name of "Methylaecorales", abundantly transcribe the key

gene for lanthanide-dependent methylotrophy in oligotrophic surface waters of the world's

oceans. TMED127/Methylaecorales likely require methanol as a carbon and energy source, and

display a diel pattern of transcription of key genes for methylotrophy that peaks in the late

afternoon. These findings motivate future studies on the mechanisms of methanol production in

surface seawater.

Introduction

Methanol (CH_3OH) is a simple alcohol that is used by methylotrophic bacteria as an energy and

carbon source (1, 2). The first step of bacterial methylotrophy, oxidation of methanol to

formaldehyde, is mediated the periplasmic enzyme pyrrolo-quinoline quinone (PQQ) methanol

dehydrogenase, which requires either calcium or a light lanthanide elements (lanthanum, cerium,

praseodymium, and neodymium) as its cofactor (3). Lanthanide-dependent methanol

dehydrogenases are encoded by the gene *xoxF* and calcium-dependent methanol dehydrogenases

are encoded by the gene *mxaF*. Formaldehyde produced by methanol dehydrogenase is then

further oxidized to CO_2 via formate dehydrogenase, yielding energy for cellular growth, and/or

assimilated into cellular carbon via the ribulose monophosphate cycle or the serine cycle (4).

Methylotrophy appears to be widespread in terrestrial and marine ecosystems. Plants

make methanol from cell wall pectins using pectin methylesterase (5), and terrestrial plant-

associated methylotrophic bacteria have been extensively studied for their agricultural

importance (6). Methanol is also produced by marine phytoplankton via unknown mechanism(s)

(7) and can also be produced photochemically, particularly over tropical oceans, by reaction of

methyl peroxy radical with hydroxyl radicals (8). Most methanol emitted to the atmosphere is

rapidly deposited back to the ocean surface (9, 10), where it is rapidly consumed by bacteria for

energy and, to a lesser degree, for carbon assimilation (11-15).

Although calcium is nine orders of magnitude more abundant than lanthanides in

seawater, marine microbial methanol oxidation is primarily lanthanide dependent, based on the

prevalence and transcriptional activity of *xoxF* compared to *mxaF* (16). Selective drawdown of

light lanthanides relative to heavy lanthanides in seawater is associated with biological activity

(17-19). In coastal waters, methylotrophic bacteria are dominated by Methylophilaceae (also

known as OM43; 20, 21-26) and Methylophagaceae (15, 27-29). Omics studies have revealed a

much broader diversity of marine bacteria with the capacity for lanthanide-based methylotrophy

than previously realized (30-34). A recent study reported an unidentified alphaproteobacterial

"Clade X" of *xoxF* genes as having highest transcriptional activity in seawater (Voutsinos et al.,

2025). Here we investigated the identity, predicted metabolism, and transcriptional activity of

methylotrophic bacteria in oligotrophic waters representative of the vast majority of the ocean's

surface to resolve the identity and metabolism of the elusive "Clade X" methylotroph.

Results

Most highly transcribed xoxF5 genes in oligotrophic surface seawater belong to the

alphaproteobacterial order TMED127. While examining RNA-Seq datasets on JGI IMG, we

found extremely high transcription of lanthanide-dependent methanol dehydrogenase *xoxF5*

genes (up to 2056 RPKM) in oligotrophic surface seawater from three different ocean basins

(North Atlantic, South Atlantic, and Indian Ocean (32, 35-37); **Table S1**). The most highly

transcribed *xoxF5* genes had high identity (93-100%) to genes from the uncultured marine

alphaproteobacterial order TMED127 (GTDB taxonomy; 38), previously known as

"Ricksettsiales" in NCBI taxonomy, with the most abundant *xoxF5* transcripts from genus GCA-

2691245 **(Fig. 1A; Table S1)**. In a dataset from the Bermuda Atlantic Time-series Study (BATS;

Sargasso Sea, North Atlantic Ocean) with free-living (SX) and particulate (LSF) size fractions,

xoxF5 transcripts were almost exclusively present in free-living fractions (**Table S1)**, suggesting

that active methylotrophs are free living rather than particle associated.

TMED127 *xoxF5* genes form a monophyletic clade with other Alphaproteobacteria

xoxF5 genes transcribed in deeper seawater samples (10-150 RPKM; UBA8558, GCA-

002720895, UBA11136; **Fig. 1A; Table S1**). BLAST searches of metatranscriptomes from the

Eastern Tropical North Pacific oxygen deficient zone (39) revealed that UBA11136 *xoxF5*

transcripts are abundantly transcribed (up to 550 RPKM) in anoxic waters **(Table S2)**,

supporting the hypothesis that UBA11136 oxidizes methanol in oxygen deficient zones (40).

Other Alphaproteobacteria (GCA-002720895, UBA7887) and Gammaproteobacteria (UBA4486,

Ga0077536, Burkholderiales (UBA7377)) also transcribed *xoxF5* genes in globally distributed

samples and across a range of depths (10-500 RPKM; **Fig. 1A; Table S1**), consistent with

previous reports (30, 34). As predicted, Methylophilaceae/OM43 *xox4* transcripts (1-350 RPKM)

were present in coastal waters **(Table S1)**.

A

0.2

<100 m depth
>100 m depth

GOS2235 10351912
Ga0456549_000444_30_1862 ★
GIR25668 CM1_5m.P40
MFL2660763 LRMED-288
Ga0206688 109407361
PDH56613 MED-G10
MEL0124461 139539_COM_metabat_bins_2.tsv.238
MAI84053 MED657
Ga0456454_000930_29_1861
Ga0394919_003329_118_1950
MDC3176690 AH-716-D07

Methylaecorales (TMED127)

MAP33298 EAC1785
HAE75578 UBA8558

UBA8558

MEC8875952 XB_MAG762

GCA-002720895

MBT5373938 SI074 bin107
MBL6935190 BS150m-G7
MDP6304722 ETNP15_MAG_21
MDP6486191 Arabian12_MAG_22

UBA11136

META1p1740 *Methylorubrum extorquens* AM1
META1p2757 *Methylorubrum extorquens* AM1

Rhizobiales

QEW19709 *Marinibacterium anthonyi*
WP_108028715 *Rhodovulum kholense* DSM19783

Rhodobacterales

WP_010975084 *Sinorhizobium meliloti* 5A14

Rhizobiales

WP_011088953 *Bradyrhizobium diazoefficiens* USDA110
WP_085126452 *Tistlia consotensis* DSM21585

Kiloniellales

MBT2990921 *Thiodiazotropha taylori*
ODJ86441 *Thiodiazotropha endolucinida*

Chromatiales

Ga0138391 1435782
HIE19504 UWMA-0238

Planktothermales (UBA7887)
Rhodospirillales

MBH71000 SP38
Ga0456544_00244_35_1837
Ga0593112_004266_102_1943
Ga0567547_016405_47_1873
Ga0496913_003323_48_1850

GCA-002720895

OUW02738 TMED156
OUV02704 TMED82
GIR48069 CM1 5m P58
Ga0395878_010991_11_1834
Ga0395878_010653_56_1879
OUV31211 TMED100

Burkholderiales (UBA7377)

WP_024300827 *Methyloversatilis discipulorum* FAM1
WP_014751259 *Advenella kashmirensis*
WP_002539484 *Grimontia marina* CECT8713

Enterobacterales

WP_010959663 *Methylococcus capsulatus*
QBC28924 *Methylomonas* LW13

Methylococcales

KGM08091 *Methylophaga thiooxydans*
BDZ73483 *Methylophaga marina*

Nitrosococcales

MEE2983970 XB_MAG1169
Ga0138398_10565661
Ga0123370_11019921

Ga0077536

MAS81431 NAT241
MBK31765 SP289
Ga0593131_001334_2_1939
Ga0456486_00105_14_2014
MAS84057 NAT232
MCH9047714 B18D1T2

Riflewelliales (UBA4486)

xoxF5 100

xoxF4 100

Mmol_1770 *Methylotenera mobilis* JLW8 XoxF4-2
Mmol_2048 *Methylotenera mobilis* JLW8 XoxF4-1
Ga0453800_00330_46_1923
EAV46784 HTCC2181
Ga0453801_00305_93_1985
Ga0453770 0025617 51 1943
CAH7806198 *Methylopumilus planktonicus*

Burkholderiales (Methylophilaceae)

Bootstrap values: 99, 100, 100, 99, 91, 98, 72, 76, 88, 99, 100, 50, 87, 99, 100, 100, 100

Alphaproteobacteria
Gammaproteobacteria

B

3 kbp

Hopanoid biosynthesis   Tetrahydromethanopterin synthesis and C1 transfer   Methanol oxidation   *xoxF5*   PQQ biosynthesis   Lanthanide uptake

Figure 1. (A) Maximum likelihood phylogenetic tree of PQQ-methanol dehydrogenases XoxF5 sequences, with XoxF4 sequences as outgroup. Where available, sample depths are colored (light blue: 0-100 m depth; dark blue: >100 m depth) for marine samples. Bootstrap values are shown when ≥50. Orders are labeled according to GTDB taxonomy. The *xoxF5* gene GOS2235 10351912 is from a previous study (23). The star marks the JGI IMG seawater sample with highest transcription (2056 RPKM), from the Sargasso Sea (TMED127 genus GCA-

2691245). Characterized sequences (bold) are from refs. (31) and (41). The two Burkholderiales clades are designated by family (Methylophilaceae) and genus (UBA7377). All JGI IMG protein sequences (beginning with "Ga") are available in **Table S1**. **(B) "Methylotrophy island" (~45 kbp) of TMED127 genome.** The genome neighborhood is drawn based on TMED127 CM1_5m.P40 (GenBank: BOXE01000001) from the same genus as the most highly transcribed gene in oligotrophic surface seawater, with the following modules: hopanoid biosynthesis (CM15mP40_03190-03050), tetrahydromethanopterin synthesis and C1 transfer (CM15mP40_03030-02810), methanol oxidation (CM15mP40_02770-02720), PQQ biosynthesis (CM15mP40_02710-02680), and lanthanide uptake (CM15mP40_02670-02620). Gray indicates hypothetical proteins.

TMED127 genomes are small and contain "methylotrophy island". As of May 2025, there are 58 TMED127 SAG/MAGs in GTDB, originating from seawater samples of the Atlantic Ocean, Pacific Ocean, Mediterranean Sea, Red Sea, and Brisbane River **(Table S3)**. These include three MAGs comprising "Clade X" in ref. (16): MED715, MED657, and MED715 **(Table S3)**. TMED127 SAG/MAGs are small (~1.5 Mb estimated complete size) and contain a ~45 kbp "methylotrophy island" with genes for methanol oxidation, tetrahydromethanopterin synthesis and C1 transfer, lanthanide uptake (*lutABEFCG* (42)), and pyrroloquinoline quinone (PQQ) biosynthesis **(Fig. 1B).** Upstream of the "methylotrophy island" is a gene cluster encoding the hopanoid biosynthetic pathway implicated in structural ordering of the outer membrane (43, 44).

Diel rhythm of transcription of methanol and glucose dehydrogenases in TMED127. A previous metatranscriptomic time series at BATS found that ~10% of all genes had diel transcriptional patterns (36). To assess transcription of TMED127 genes, we mapped transcripts from surface seawater (5 m depth) from the previous study, sampled every 4 hours over 5 days **(Table S4)**, to most complete TMED127 SAG from genus GCA-2691245 (AG-892-F10; estimated 90% completeness; 0.01% contamination (45); **Fig. 1A; Table S1).** Throughout the time series, *xoxF5* was consistently the most abundant transcript, comprising 16 to 57% (1.6 to 5.7 x 10^5 TPM) of AG-892-F10 transcripts **(Table S5)**. The *xoxF5* gene displayed a diel

transcriptional cycle, consistently peaking in the late afternoon (16:00-20:00; **Fig. 2**). A PQQ-

dependent glucose dehydrogenase (*gdh*) showed the same transcriptional pattern at roughly an

order of magnitude lower abundance **(Fig. 2; Table S5)**. Five other genes were transcribed at

$>10^4$ TPM, but not did not show a temporal pattern; these included proteorhodopsin, two

hypothetical proteins (AG-892-F10.CDS.256 and AG-892-F10.CDS.257), the phosphate ABC

transport substrate-binding protein (*pstS*), and a DoxX-like family protein (**Table S5**). Homologs

of AG-892-F10.CDS.256 and AG-892-F10.CDS.257 (34% lysine), were only present in other

TMED127 SAGs/MAGs, whereas the DoxX-like integral membrane family protein had

homologs in other marine Alphaproteobacteria and Methylophilacaeae/OM43.

Figure 2. Diel transcription of TMED127 lanthanide-dependent methanol dehydrogenase
(*xoxF5*) and glucose dehydrogenase (*gdh*) at Bermuda Atlantic Time-series Study (BATS)
site, Sargasso Sea, North Atlantic Ocean. The asterisk represents a single anonymously high
sample. Error bars (standard deviation) are shown for the nine time points with multiple
metatranscriptomes. Missing lines are due to lack of samples for two timepoints.
143

TMED127 are likely obligate methylotrophs. Genomic analysis suggests that TMED127

are obligate methylotrophs. After methanol oxidation via XoxF, formaldehyde is either oxidized

to formate and carbon dioxide via the tetrahydromethanopterin C1 transfer pathway and

tungsten-containing formate dehydrogenase, respectively, or assimilated into biomass via the

serine and ethylmalonyl-CoA pathways **(Fig. 3; Table S5)**. TMED127's core methylotrophic

modules match those of the well-studied plant-associated facultative methylotroph

Methylorubrum extorquens AM1 (46, 47); roughly two-thirds (956/1535) of AG-892-F10's

genes have homologs in AM1 **(Tables S5)**. TMED127 genes without AM1 homologs include

several genes common in marine proteobacteria, including proteorhodopsin for light-driven

proton pumping and nickel-dependent superoxide dismutase **(Fig. 3)**. While TMED127's small

(~1.5 Mb) genome has only one copy of most key functional genes, AM1's large (6.9 Mb)

genome has multiple copies of many key functional genes. AM1 also has a full TCA cycle,

enabling its growth on multiple-carbon compounds. Although we cannot be certain of gene

absence in incomplete genomes, TMED127 SAGs/MAGs lack the 2-oxoglutarate dehydrogenase

complex, suggesting that they have incomplete TCA cycles characteristic of obligate

methylotrophs (48).

Proposed renaming of TMED127 to Methylaecorales. We propose a new name for

TMED127 (order: Methylaecorales; family Methyaecoralaceae) from "methyl" to denote

"methylotrophic metabolism", and "aecor" for "surface of the ocean".

Figure 3. Schematic of key metabolisms and functions in TMED127/Methylaecorales based on genomic reconstruction. Three of the most highly transcribed proteins are shown (PQQ-dependent methanol dehydrogenase (XoxF), PQQ-dependent glucose dehydrogenase (GDH), and proteorhopsin (PR)). Question marks represent hypothetical lanthanide deposits and lanthanide chelation by D-glucono-1,5-lactone, which remains unconfirmed.

Discussion

This study illuminates the identity, metabolism, and diel rhythms of TMED127 (proposed name: Methylaecorales (order); Methyaecoralaceae (family)), a deep-branching order of marine alphaproteobacteria (49). TMED127/Methylaecorales appear to be free-living obligate methylotrophs, although confirmation of their metabolism will require isolation and culture experiments. While cell counts of TMED127/Methylaecorales are not available, they appear to be a relatively minor bacterial population in surface seawater.

The source of methanol to TMED127/Methylaecorales is unknown. It may be produced as a waste product by eukaryotic phytoplankton and/or cyanobacteria, as is known to occur in culture experiments (7), and/or via photochemical degradation of other organic molecules

produced by phytoplankton in surface seawater (8). Peak *xoxF5* transcription in the late

afternoon may imply highest methanol supply at that time. Seawater carbohydrate concentrations

are highest in the afternoon, likely due to passive leakage of uncharged molecules during

"photosynthetic overflow" from phytoplankton (50). It is possible that similar methanol leakage

from phytoplankton also peaks in the afternoon, but further study of methanol production

mechanisms in seawater is required.

PQQ-dependent glucose dehydrogenase (GDH), a periplasmic protein that oxidizes D-

glucose to D-glucono-1,5-lactone using calcium as a cofactor (51), showed a synchronous

transcriptional pattern to *xoxF5* in TMED127/Methylaecorales, rising throughout the day and

falling at night. D-glucono-1,5-lactone is a strong metal chelator that can solubilize calcium

phosphate minerals (52) and may function as a lanthanide chelator in methylotrophic bacteria

(53). Crystalline accumulations of lanthanides are present in the periplasm (53, 54) and

cytoplasm (42) of terrestrial alphaproteobacterial methylotrophs. We hypothesize that D-

glucono-1,5-lactone production by GDH releases lanthanides from intracellular storage deposits

for incorporation into XoxF **(Fig. 3)**; this hypothesis will require future testing once bacterial

isolates are available.

The full complement of lanthanide acquisition and storage pathways in

TMED127/Methylaecorales also requires more study. We identified genes encoding the high

affinity lanthanide uptake transport system (LutABEFCG; 42), but did not find homologs of

genes for "lanthanophore" ligands (55) nor lanthanum storage proteins such as lanmodulin (56)

and lanpepsy (57). Given the extremely low (picomolar) lanthanide concentrations in seawater

(58, 59), it seems likely that these bacteria have additional lanthanide scavenging strategies,

possibly involving one or more of the highly transcribed genes of unknown function.

Our findings suggest that TMED127/Methylaecorales may be the key player in

lanthanide-dependent methylotrophy in oligotrophic seawater. More studies are needed to

determine whether a diel cycle of methanol production occurs in surface seawater and the

quantitative importance of TMED127/Methylaecorales in the global carbon and lanthanide

cycles.

Data Availability

The KBase bioinformatic pipeline is available at https://narrative.kbase.us/narrative/214786.

Methods and Materials

Identifying and quantifying xoxF gene transcription in seawater samples. A python code

(GeneSweeper; available at https://github.com/GlassLabGT/Python-scripts) was used to search

for all possible product names for the target protein, PQQ-dependent methanol dehydrogenase:

("PQQ-dependent dehydrogenase (methanol/ethanol family)", "methanol dehydrogenase

(cytochrome c) subunit 1", "methanol dehydrogenase (cytochrome)", "alcohol dehydrogenase

(cytochrome c)", "glucose dehydrogenase", "quinohemoprotein ethanol dehydrogenase", and

"quinoprotein glucose dehydrogenase") in CSV files (including gene ID, product name, scaffold

ID, gene sequence length, and read count) for all coastal and marine datasets in Joint Genome

Institute's (JGI) Integrated Microbial Genome (IMG) RNASeq Studies database as of May 2025.

Only unamended seawater samples were processed; enrichments or laboratory cultures were

removed. The total read count of each metatranscriptome was obtained from the JGI IMG

RNASeq Studies database. Gene transcription was then converted to reads per kilobase per

million mapped reads (RPKM). Amino acid sequences were obtained from reference

metagenomes using the JGI IMG "Find Genes" function with the Gene ID as input. Each amino

acid sequence was then queried against the NCBI nr database by BLASTP search (May 2025).

Only full-length gene sequences were kept for further processing. The top NCBI hit was matched

to its GTDB taxonomy (release 10-RS226) by searching GTDB (38) for the GCA assembly of

the SAG/MAG.

XoxF phylogeny and gene neighborhood. Sequences ($n = 63$; 552 gap-free sites) were aligned

using the MAFFT online server with the L-INS-i method (60). The fasta file alignment

(XoxF5_alignment.fasta) is available as a supplemental data set. A phylogeny of XoxF amino

acid sequences was generated with 1,000 bootstraps using model LG+F+I+G4 with W-IQ-Tree

(61). The phylogeny was visualized using FigTree v.1.4.4. The gene neighborhood was

generated using the EFI Gene Neighborhood Tool (48) with single sequence BLAST of the

UniProt database with NCBI accession GIR25668.1 as the query with 100 genes upstream and

downstream the gene of interest.

Transcription of *xoxF* in Eastern Tropical North Pacific oxygen minimum zone. Magic-

Basic Local Alignment Search Tool (Magic-BLAST) (62) was used with default parameters to

search ETNP ODZ metatranscriptomes (PRJNA727903; 39) for *xoxF* genes from UBA11136

MAGs using DNA sequence of *xoxF5* from Alphaproteobacteria bacterium isolate

ETNP15_MAG_21 (MDP6304722.1) as the query. Read hits were normalized to reads per

kilobase million (RPKM).

TMED127 SAG/MAG analysis and mapping to metatranscriptomes. All SAG/MAGs in the Alphaproteobacteria order TMED127 in GTDB (release 10-RS226) as of May 2025 were collected into a BV-BRC genome group, which was used to search for product names via "Proteins" tab and for metabolic pathways via "Pathways" tab (63). The TMED127 SAG AG-892-F10 (estimated 90% completeness; 0.01% contamination; 45) was chosen for further study because it was the most complete SAG/MAG from the genus GCA-2691245, which contained the most abundant *xoxF5* transcripts in surface seawater (see text). Additional analysis was performed in KBase (64). SAG AG-892-F10 (NCBI WGS Project CACMWI01) and *Methylorubrum extorquens* AM1 (NCBI Reference Sequence NC_012808.1) were imported into KBase, and the KBase program "Compare Two Proteomes" was used to find homologs. RNA-Seq data (JGI GOLD Study ID Gs0161320; **Table S4**) were imported into KBase and aligned to SAG AG-892-F10 using the Bowtie2 app (65). The aligned RNA-Seq data was then assembled into a set of transcripts quantified by transcripts per million (TPM) using the StringTie v.2.1.5 (66) app.

Acknowledgments

We thank Naomi Gilbert, Steven Wilhelm, Matthew Sullivan, and David Walsh for providing access to sequences and metadata. We thank Ann Pearson for helpful discussions.

References

1. Chistoserdova L, Kalyuzhnaya MG. 2018. Current trends in methylotrophy. Trends Microbiol 26:703-714.

2. Chistoserdova L, Kalyuzhnaya MG, Lidstrom ME. 2009. The expanding world of methylotrophic metabolism. Annu Rev Microbiol 63:477-99.

273   3.   Keltjens JT, Pol A, Reimann J, Op den Camp HJ. 2014. PQQ-dependent methanol
274   dehydrogenases: rare-earth elements make a difference. Appl Microbiol Biotechnol
275   98:6163-83.

276   4.   Chistoserdova L. 2011. Modularity of methylotrophy, revisited. Environ Microbiol
277   13:2603-2622.

278   5.   Dorokhov YL, Sheshukova EV, Komarova TV. 2018. Methanol in plant life. Front Plant
279   Sci 9:1623.

280   6.   Zhang C, Wang M-Y, Khan N, Tan L-L, Yang S. 2021. Potentials, utilization, and
281   bioengineering of plant growth-promoting *Methylobacterium* for sustainable agriculture.
282   Sustainability 13:3941.

283   7.   Mincer TJ, Aicher AC. 2016. Methanol production by a broad phylogenetic array of
284   marine phytoplankton. PLoS One 11:e0150820.

285   8.   Khan MAH, Cooke MC, Utembe SR, Xiao P, Derwent RG, Jenkin ME, Archibald AT,
286   Maxwell P, Morris WC, South N, Percival CJ, Shallcross DE. 2014. Reassessing the
287   photochemical production of methanol from peroxy radical self and cross reactions using
288   the STOCHEM-CRI global chemistry and transport model. Atmospheric Environment
289   99:77-84.

290   9.   Bates KH, Jacob DJ, Wang S, Hornbrook RS, Apel EC, Kim MJ, Millet DB, Wells KC,
291   Chen X, Brewer JF, Ray EA, Commane R, Diskin GS, Wofsy SC. 2021. The global
292   budget of atmospheric methanol: new constraints on secondary, oceanic, and terrestrial
293   sources. Journal of Geophysical Research: Atmospheres 126:e2020JD033439.

294   10.   Yang M, Nightingale PD, Beale R, Liss PS, Blomquist B, Fairall C. 2013. Atmospheric
295   deposition of methanol over the Atlantic Ocean. Proceedings of the National Academy of
296   Sciences 110:20034–20039

297   11.   Dixon JL, Beale R, Nightingale PD. 2011. Rapid biological oxidation of methanol in the
298   tropical Atlantic: significance as a microbial carbon source. Biogeosciences 8:2707-2716.

299   12.   Dixon JL, Beale R, Nightingale PD. 2011. Microbial methanol uptake in northeast
300   Atlantic waters. ISME J 5:704-16.

301   13.   Zhou Z, Zhuang GC, Mao SH, Liu J, Li XJ, Liu Q, Song GD, Zhang HH, Chen Z,
302   Montgomery A, Joye S, Yang GP. 2023. Methanol concentrations and biological
303   methanol consumption in the Northwest Pacific Ocean. Geophys Res Lett
304   50:e2022GL101605.

305   14.   Dixon JL, Nightingale PD. 2012. Fine-scale variability in methanol uptake and oxidation:
306   from the microlayer to 1000 m. Biogeosciences 9:2961-2972.

15. Dixon JL, Sargeant S, Nightingale PD, Colin Murrell J. 2013. Gradients in microbial
methanol uptake: productive coastal upwelling waters to oligotrophic gyres in the
Atlantic Ocean. ISME J 7:568-80.

16. Voutsinos MY, Banfield JF, McClelland HO. 2025. Extensive and diverse lanthanide-
dependent metabolism in the ocean. ISME J 19:wraf057.

17. Shiller AM, Chan EW, Joung DJ, Redmond MC, Kessler JD. 2017. Light rare earth
element depletion during Deepwater Horizon blowout methanotrophy. Sci Rep 7:10389.

18. Hara Y, Obata H, Doi T, Hongo Y, Gamo T, Takeda S, Tsuda A. 2009. Rare earth elements
in seawater during an iron-induced phytoplankton bloom of the western subarctic Pacific
(SEEDS-II). Deep Sea Research Part II: Topical Studies in Oceanography 56:2839-2851.

19. Meyer ACS, Grundle D, Cullen JT. 2021. Selective uptake of rare earth elements in
marine systems as an indicator of and control on aerobic bacterial methanotrophy. Earth
and Planetary Science Letters 558:116756.

20. Giovannoni SJ, Hayakawa DH, Tripp HJ, Stingl U, Givan SA, Cho JC, Oh HM, Kitner
JB, Vergin KL, Rappe MS. 2008. The small genome of an abundant coastal ocean
methylotroph. Environ Microbiol 10:1771-82.

21. Sowell SM, Abraham PE, Shah M, Verberkmoes NC, Smith DP, Barofsky DF,
Giovannoni SJ. 2011. Environmental proteomics of microbial plankton in a highly
productive coastal upwelling system. ISME J 5:856-65.

22. Rappé MS, Kemp PF, Giovannoni SJ. 2003. Phylogenetic diversity of marine coastal
picoplankton 16S rRNA genes cloned from the continental shelf off Cape Hatteras, North
Carolina. Limnol Oceanogr 42:811-826.

23. Ramachandran A, Walsh DA. 2015. Investigation of XoxF methanol dehydrogenases
reveals new methylotrophic bacteria in pelagic marine and freshwater ecosystems. FEMS
Microbiol Ecol 91:fiv105.

24. Taubert M, Grob C, Howat AM, Burns OJ, Dixon JL, Chen Y, Murrell JC. 2015. XoxF
encoding an alternative methanol dehydrogenase is widespread in coastal marine
environments. Environ Microbiol 17:3937-3948.

25. Georges AA, El-Swais H, Craig SE, Li WK, Walsh DA. 2014. Metaproteomic analysis of
a winter to spring succession in coastal northwest Atlantic Ocean microbial plankton.
ISME J 8:1301-13.

26. Morris RM, Longnecker K, Giovannoni SJ. 2006. Pirellula and OM43 are among the
dominant lineages identified in an Oregon coast diatom bloom. Environ Microbiol
8:1361-70.

27. Neufeld JD, Schafer H, Cox MJ, Boden R, McDonald IR, Murrell JC. 2007. Stable-
isotope probing implicates Methylophaga spp and novel Gammaproteobacteria in marine
methanol and methylamine metabolism. ISME J 1:480-91.

28. Janvier M. 1985. *Methylophaga marina* gen. nov. sp. nov. and *Methylophaga thalassica*
sp. nov., marine methylotrophs. Int J System Bacteriol 35:131-139.

29. Grob C, Taubert M, Howat AM, Burns OJ, Dixon JL, Richnow HH, Jehmlich N, von
Bergen M, Chen Y, Murrell JC. 2015. Combining metagenomics with metaproteomics
and stable isotope probing reveals metabolic pathways used by a naturally occurring
marine methylotroph. Environ Microbiol 17:4007-18.

30. Francis B, Urich T, Mikolasch A, Teeling H, Amann R. 2021. North Sea spring bloom-
associated Gammaproteobacteria fill diverse heterotrophic niches. Environ Microbiome
16:15.

31. Huang J, Yu Z, Groom J, Cheng J-F, Tarver A, Yoshikuni Y, Chistoserdova L. 2019. Rare
earth element alcohol dehydrogenases widely occur among globally distributed,
numerically abundant and environmentally important microbes. ISME J 13:2005-2017.

32. Cavaco MA, Bhatia MP, Hawley AK, Torres-Beltrán M, Johnson WM, Longnecker K,
Konwar K, Kujawinski EB, Hallam SJ. 2022. Pathway-centric analysis of microbial
metabolic potential and expression along nutrient and energy gradients in the Western
Atlantic Ocean. Front Mar Sci 9.

33. Hanson BT, Hewson I, Madsen EL. 2014. Metaproteomic survey of six aquatic habitats:
discovering the identities of microbial populations active in biogeochemical cycling.
Microb Ecol 67:520-39.

34. Zhou Z, Tran PQ, Kieft K, Anantharaman K. 2020. Genome diversification in globally
distributed novel marine Proteobacteria is linked to environmental adaptation. ISME J
14:2060-2077.

35. Gilbert NE, Muratore D, Shopen Gochev C, LeCleir GR, Cagle SM, Pound HL, Sun CL,
Carillo A, Ndlovu KS, Maidanik I, Coenen AR, Chittick L, DeBruyn JM, Buchan A,
Lindell D, Sullivan MB, Weitz JS, Wilhelm SW. 2025. Seasonal enhancement of the viral
shunt catalyzes a subsurface oxygen maximum in the Sargasso Sea. BioRxiv
doi:10.1101/2025.01.23.634377.

36. Muratore D, Gilbert NE, LeCleir GR, Wilhelm SW, Weitz JS. 2025. Diel partitioning in
microbial phosphorus acquisition in the Sargasso Sea. Proc Natl Acad Sci U S A
122:e2410268122.

37. Gilbert NE, LeCleir GR, Pound HL, Strzepek RF, Ellwood MJ, Twining BS, Roux S,
Boyd PW, Wilhelm SW. 2023. Giant virus infection signatures are modulated by euphotic
zone depth strata and iron regimes of the subantarctic Southern Ocean. mSystems 8:1-17.

38. Parks DH, Chuvochina M, Rinke C, Mussig AJ, Chaumeil PA, Hugenholtz P. 2022.
GTDB: an ongoing census of bacterial and archaeal diversity through a phylogenetically
consistent, rank normalized and complete genome-based taxonomy. Nucleic Acids Res
50:D785-D794.

39. Mattes TE, Burke S, Rocap G, Morris RM. 2022. Two metatranscriptomic profiles
through low-dissolved-oxygen waters (DO, 0 to 33 µM) in the Eastern Tropical North
Pacific Ocean. Microbiol Resour Announc 11:e01201-21.

40. Elbon CE, Stewart FJ, Glass JB. 2024. Novel Alphaproteobacteria transcribe genes for
nitric oxide transformation at high levels in a marine oxygen-deficient zone. Appl
Environ Microbiol 90:02099-23.

41. Huang J, Yu Z, Chistoserdova L. 2018. Lanthanide-dependent methanol dehydrogenases
of XoxF4 and XoxF5 clades are differentially distributed among methylotrophic bacteria
and they reveal different biochemical properties. Front Microbiol 9:1366.

42. Roszczenko-Jasińska P, Vu HN, Subuyuj GA, Crisostomo RV, Cai J, Lien NF, Clippard
EJ, Ayala EM, Ngo RT, Yarza F, Wingett JP, Raghuraman C, Hoeber CA, Martinez-
Gomez NC, Skovran E. 2020. Gene products and processes contributing to lanthanide
homeostasis and methanol metabolism in *Methylorubrum extorquens* AM1. Sci Rep 10:1-
15.

43. Bradley AS, Pearson A, Sáenz JP, Marx CJ. 2010. Adenosylhopane: The first
intermediate in hopanoid side chain biosynthesis. Organic Geochemistry 41:1075-1081.

44. Saenz JP, Grosser D, Bradley AS, Lagny TJ, Lavrynenko O, Broda M, Simons K. 2015.
Hopanoids as functional analogues of cholesterol in bacterial membranes. Proc Natl Acad
Sci U S A 112:11971-6.

45. Pachiadaki MG, Brown JM, Brown J, Bezuidt O, Berube PM, Biller SJ, Poulton NJ,
Burkart MD, La Clair JJ, Chisholm SW, Stepanauskas R. 2019. Charting the complexity
of the marine microbiome through single-cell genomics. Cell 179:1623-1635 e11.

46. Smejkalova H, Erb TJ, Fuchs G. 2010. Methanol assimilation in *Methylobacterium
extorquens* AM1: demonstration of all enzymes and their regulation. PLoS One 5:e13001.

47. Chistoserdova L, Chen SW, Lapidus A, Lidstrom ME. 2003. Methylotrophy in
Methylobacterium extorquens AM1 from a genomic point of view. J Bacteriol 185:2980-
7.

48. Wood AP, Aurikko JP, Kelly DP. 2004. A challenge for 21st century molecular biology
and biochemistry: what are the causes of obligate autotrophy and methanotrophy? FEMS
Microbiol Rev 28:335-52.

49. Geiger O, Sanchez-Flores A, Padilla-Gomez J, Degli Esposti M. 2023. Multiple
approaches of cellular metabolism define the bacterial ancestry of mitochondria. Sci Adv
9:eadh0066.

50.   Barthelmess T, Cristi A, Deppeler S, Safi K, Sellegri K, Law CS, Engel A. 2025.
Pronounced diel cycling of dissolved carbohydrates and amino acids in the surface ocean
and across diverse regimes. Environ Sci Technol 59:419-429.

51.   Southall SM, Doel JJ, Richardson DJ, Oubrie A. 2006. Soluble aldose sugar
dehydrogenase from Escherichia coli: a highly exposed active site conferring broad
substrate specificity. J Biol Chem 281:30650-9.

52.   Sashidhar B, Podile AR. 2010. Mineral phosphate solubilization by rhizosphere bacteria
and scope for manipulation of the direct oxidation pathway involving glucose
dehydrogenase. J Appl Microbiol 109:1-12.

53.   Gorniak L, Bechwar J, Westermann M, Steiniger F, Wegner C-E. 2023. Different
lanthanide elements induce strong gene expression changes in a lanthanide-accumulating
methylotroph. Microbiology Spectrum 11:e00867-23.

54.   Wegner CE, Westermann M, Steiniger F, Gorniak L, Budhraja R, Adrian L, Kusel K.
2021. Extracellular and intracellular lanthanide accumulation in the methylotrophic
Beijerinckiaceae bacterium RH AL1. Appl Environ Microbiol 87:e0314420.

55.   Zytnick AM, Gutenthaler-Tietze SM, Aron AT, Reitz ZL, Phi MT, Good NM, Petras D,
Daumann LJ, Martinez-Gomez NC. 2024. Identification and characterization of a small-
molecule metallophore involved in lanthanide metabolism. Proc Natl Acad Sci U S A
121:e2322096121.

56.   Cotruvo JA, Jr., Featherston ER, Mattocks JA, Ho JV, Laremore TN. 2018. Lanmodulin:
a highly selective lanthanide-binding protein from a lanthanide-utilizing bacterium. J Am
Chem Soc 140:15056-15061.

57.   Hemmann JL, Keller P, Hemmerle L, Vonderach T, Ochsner AM, Bortfeld-Miller M,
Gunther D, Vorholt JA. 2023. Lanpepsy is a novel lanthanide-binding protein involved in
the lanthanide response of the obligate methylotroph *Methylobacillus flagellatus*. J Biol
Chem 299:102940.

58.   Hatje V, Schijf J, Johannesson KH, Andrade R, Caetano M, Brito P, Haley BA, Lagarde
M, Jeandel C. 2024. The global biogeochemical cycle of the rare earth elements. Global
Biogeochemical Cycles 38.

59.   Noack CW, Dzombak DA, Karamalidis AK. 2014. Rare earth element distributions and
trends in natural waters with a focus on groundwater. Environ Sci Technol 48:4317-26.

60.   Katoh K, Rozewicki J, Yamada KD. 2019. MAFFT online service: multiple sequence
alignment, interactive sequence choice and visualization. Brief Bioinform 20:1160-1166.

61.   Trifinopoulos J, Nguyen LT, von Haeseler A, Minh BQ. 2016. W-IQ-TREE: a fast online
phylogenetic tool for maximum likelihood analysis. Nucleic Acids Res 44:W232-5.

62.   Boratyn GM, Thierry-Mieg J, Thierry-Mieg D, Busby B, Madden TL. 2019. Magic-
450   BLAST, an accurate RNA-seq aligner for long and short reads. BMC Bioinf 20:405.

451   63.   Olson RD, Assaf R, Brettin T, Conrad N, Cucinell C, Davis JJ, Dempsey DM, Dickerman
452   A, Dietrich EM, Kenyon RW, Kuscuoglu M, Lefkowitz EJ, Lu J, Machi D, Macken C,
453   Mao C, Niewiadomska A, Nguyen M, Olsen GJ, Overbeek JC, Parrello B, Parrello V,
454   Porter JS, Pusch GD, Shukla M, Singh I, Stewart L, Tan G, Thomas C, VanOeffelen M,
455   Vonstein V, Wallace ZS, Warren AS, Wattam AR, Xia F, Yoo H, Zhang Y, Zmasek CM,
456   Scheuermann RH, Stevens RL. 2023. Introducing the Bacterial and Viral Bioinformatics
457   Resource Center (BV-BRC): a resource combining PATRIC, IRD and ViPR. Nucleic
458   Acids Res 51:D678-D689.

459   64.   Arkin AP, Cottingham RW, Henry CS, Harris NL, Stevens RL, Maslov S, Dehal P, Ware
460   D, Perez F, Canon S, Sneddon MW, Henderson ML, Riehl WJ, Murphy-Olson D, Chan
461   SY, Kamimura RT, Kumari S, Drake MM, Brettin TS, Glass EM, Chivian D, Gunter D,
462   Weston DJ, Allen BH, Baumohl J, Best AA, Bowen B, Brenner SE, Bun CC, Chandonia
463   JM, Chia JM, Colasanti R, Conrad N, Davis JJ, Davison BH, DeJongh M, Devoid S,
464   Dietrich E, Dubchak I, Edirisinghe JN, Fang G, Faria JP, Frybarger PM, Gerlach W,
465   Gerstein M, Greiner A, Gurtowski J, Haun HL, He F, Jain R, et al. 2018. KBase: The
466   United States Department of Energy Systems Biology Knowledgebase. Nat Biotechnol
467   36:566-569.

468   65.   Langmead B, Salzberg SL. 2012. Fast gapped-read alignment with Bowtie 2. Nat
469   Methods 9:357-9.

470   66.   Pertea M, Pertea GM, Antonescu CM, Chang TC, Mendell JT, Salzberg SL. 2015.
471   StringTie enables improved reconstruction of a transcriptome from RNA-seq reads. Nat
472   Biotechnol 33:290-5.

A

0.2

<100 m depth
>100 m depth

GOS2235 10351912
Ga0456549_000444_30_1862 ★
GIR25668 CM1_5m.P40
MFL2660763 LRMED-288
Ga0206688 109407361
PDH56613 MED-G10
MEL0124461 139539_COM_metabat_bins_2.tsv.238
MAI84053 MED657
Ga0456454_000930_29_1861
Ga0394919_003329_118_1950
MDC3176690 AH-716-D07

99
100

MAP33298 EAC1785
HAE75578 UBA8558
MEC8875952 XB_MAG762

MBT5373938 SI074 bin107
MBL6935190 BS150m-G7
MDP6304722 ETNP15_MAG_21
MDP6486191 Arabian12_MAG_22

META1p1740 ***Methylorubrum extorquens*** AM1
META1p2757 ***Methylorubrum extorquens*** AM1
QEW19709 *Marinibacterium anthonyi*
WP_108028715 ***Rhodovulum kholense*** DSM19783
WP_010975084 ***Sinorhizobium meliloti*** 5A14
WP_011088953 ***Bradyrhizobium diazoefficiens*** USDA110
WP_085126452 ***Tistlia consotensis*** DSM21585

MBT2990921 *Thiodiazotropha taylori*
ODJ86441 *Thiodiazotropha endolucinida*

Ga0138391 1435782
HIE19504 UWMA-0238

MBH71000 SP38
Ga0456544_00244_35_1837
Ga0593112_004266_102_1943
Ga0567547_016405_47_1873
Ga0496913_003323_48_1850

OUW02738 TMED156
OUV02704 TMED82
GIR48069 CM1 5m P58
Ga0395878_010991_11_1834
Ga0395878_010653_56_1879
OUV31211 TMED100
WP_024300827 *Methyloversatilis discipulorum* FAM1
WP_014751259 *Advenella kashmirensis*

100
99
50

WP_002539484 *Grimontia marina* CECT8713
WP_010959663 *Methylococcus capsulatus*
QBC28924 *Methylomonas* LW13

KGM08091 *Methylophaga thiooxydans*
BDZ73483 *Methylophaga marina*

MEE2983970 XB_MAG1169
Ga0138398_10565661
Ga0123370_11019921

MAS81431 NAT241
MBK31765 SP289
Ga0593131_001334_2_1939
Ga0456486_00105_14_2014
MAS84057 NAT232
MCH9047714 B18D1T2

99
100

xoxF5
88
100

Mmol_1770 *Methylotenera mobilis* JLW8 XoxF4-2
Mmol_2048 *Methylotenera mobilis* JLW8 XoxF4-1
Ga0453800_00330_46_1923
EAV46784 HTCC2181
Ga0453801_00305_93_1985
Ga0453770 0025617 51 1943
CAH7806198 *Methylopumilus planktonicus*

xoxF4
100
100

Methylaecorales
(TMED127)

UBA8558
GCA-002720895

UBA11136

Rhizobiales

Rhodobacterales

Rhizobiales

Kiloniellales

Chromatiales

Planktothermales (UBA7887)
Rhodospirillales

GCA-002720895

Burkholderiales
(UBA7377)

Enterobacterales

Methylococcales

Nitrosococcales

Ga0077536

Riflewelliales
(UBA4486)

Burkholderiales
(Methylophilaceae)

Alphaproteobacteria

Gammaproteobacteria

B

3 kbp

Hopanoid biosynthesis   Tetrahydromethanopterin synthesis and C1 transfer   *xoxF5*   Methanol oxidation   PQQ biosynthesis   Lanthanide uptake