Mutation or deletion of the *Avr6* avirulence effector gene of *Fulvia fulva* leads to breakdown of the *Cf-6* leaf mold resistance locus in tomato

Christiaan R. Schol [1,2], Like Fokkens [1], Anan Hu [1], Ruifang Jia [1], Silvia de la Rosa [3,†], Bilal Ökmen [1,‡],

Anne M. Hilgers [2], Samuel L. van Zwoll [2], Luuk D.H. Veenendaal [2], Marie Turner [4], Claudie Monot [4],

Kazuya Maeda [5], Yuichiro Iida [5], Pierre J.G.M de Wit [1], Danny Esselink [2], Anne-Marie A. Wolters[2],

Yuling Bai [2*], Matthieu H.A.J. Joosten [1,*], Carl H. Mesarich [3,6*]

[1] Laboratory of Phytopathology, Wageningen University, Droevendaalsesteeg 1, 6708 PB Wageningen, The Netherlands.

[2] Plant Breeding, Wageningen University & Research, Droevendaalsesteeg 1, 6708 PB Wageningen, The Netherlands.

[3] Laboratory of Molecular Plant Pathology, School of Agriculture and Environment, Massey University, Palmerston North 4410, New Zealand.

[4] Vegenov BBV, 1040 Pen Ar Prat, 29250 Saint Pol de Léon, France.

[5] Laboratory of Plant Pathology, Faculty of Agriculture, Setsunan University, Hirakata, Osaka, 573-0101, Japan.

[6] Bioprotection Aotearoa, Massey University, Palmerston North 4410, New Zealand.

[†] Current address: BioLumic Limited, Palmerston North 4440, New Zealand.

[‡] Current address: Biotic Stress Department, KWS Group, 31 Grimsehlstraße, Mailbox 1463, 37555 Einbeck, Germany.

* Corresponding authors: Yuling Bai (bai.yuling@wur.nl), Matthieu H.A.J. Joosten (matthieu.joosten@wur.nl) and Carl H. Mesarich (c.mesarich@massey.ac.nz)

Summary

Genetic resistance is an important mitigation strategy in commercial tomato cultivation against leaf mold disease caused by the fungal pathogen *Fulvia fulva*. In recent years, the widely deployed *Cf-9* resistance locus has been overcome, which has prompted breeders to deploy the novel resistance locus *Cf-6*. However, *Cf-6*-mediated resistance has already been overcome in several locations worldwide. To date, the identity of *Cf-6*, as well as the gene encoding the matching Avr6 effector from *F. fulva* have remained unknown. Here, we report the identification of *Avr6* using a comparative genomics strategy.

Whole genome sequencing of *Cf-6*-breaking *F. fulva* strains revealed that the gene coding for the known

secreted effector Ecp5 is either deleted or mutated in these strains. *Ecp5* was subsequently confirmed

to be *Avr6* through potato virus X-mediated expression in *Cf-6* tomato plants, as well as by gene

knockout and complementation experiments. Using the Avr6-triggered hypersensitive response as a

marker for disease resistance, we mapped the *Cf-6* locus to the short arm of chromosome 12 in tomato.

In addition to identifying *Avr6* and setting the foundation for cloning of *Cf-6*, this work highlights the

urgent need for more research into durable gene-for-gene resistance in tomato against *F. fulva*.

Keywords

Fulvia fulva, *Cladosporium fulvum*, Tomato leaf mold disease, *Solanum lycopersicum* (tomato), *Avr6*

avirulence effector gene, *Cf-6* resistance gene, Breakdown of genetic resistance.

Introduction

Fulvia fulva, previously known as *Cladosporium fulvum*, is a Dothideomycete fungal pathogen that

causes leaf mold disease of tomato (*Solanum lycopersicum*) under humid greenhouse conditions

(Cooke, 1883; Thomma et al., 2005; Videira et al., 2017). It has no known sexual stage, although there

is growing evidence suggesting cryptic sexual reproduction (de la Rosa & Schol et al., 2024; Gül et al.,

2022; Stergiopoulos, Groenewald, et al., 2007). Introgression of genetic host resistance from wild

tomato relatives, mediated by cell-surface anchored, receptor-like proteins (RLPs) that recognize

specific *F. fulva*-secreted Avirulence (Avr) effector proteins (Dixon et al., 1998; Dixon et al., 1996; Jones

et al., 1994; Takken et al., 1999; Thomas et al., 1997), is an important breeding strategy against this

disease. These RLPs, encoded by *Cf* (resistance to *C. fulvum*) genes residing in clusters throughout the

tomato genome, mediate immune responses like the hypersensitive response (HR), associated with a

burst of reactive oxygen species (ROS) (Gust & Felix, 2014; Huang & Joosten, 2024; Kang & Yeom, 2018)

and the accumulation of pathogenesis-related (PR) proteins (Joosten & de Wit, 1989; Wubben et al.,

1993) and antimicrobial phytoalexins (de Wit & Flach, 1979). Our understanding of these RLPs and their

corresponding Avr effectors has positioned the *F. fulva*-tomato pathosystem as a long-standing model

for understanding molecular plant-pathogen interactions (de Wit, 2016; Mesarich et al., 2023; Thomma

et al., 2005).

Upon widespread deployment of *Cf* resistance genes in tomato cultivars, strong selection pressure is

exerted on *F. fulva* to circumvent Avr recognition to regain virulence. This is achieved through mutations

in, or complete loss of the corresponding *Avr* gene (de la Rosa & Schol et al., 2024; Iida et al., 2015;

Joosten et al., 1994; Mesarich et al., 2023; Stergiopoulos, De Kock, et al., 2007). To date, all known *F.*

fulva Avr genes encode small, secreted proteins (SSPs) that are highly expressed during infection

(Mesarich et al., 2023). Avr effectors are rich in cysteine residues and, often, through the mutation of

a cysteine residue and loss of the associated disulfide bond, recognition of an Avr effector by its

corresponding Cf receptor is lost (Joosten et al., 1997; Joosten et al., 1994). Most Avr effectors of *F.*

fulva show little homology to other proteins and lack known functional domains (Mesarich et al., 2018).

The only exceptions are Avr4 (chitin-binding domain) and Avr2 (protease inhibitor), which protect

against PR chitinases and proteases, respectively (Rooney et al., 2005; Van Esse et al., 2007). Recently,

a combined transcriptomics and proteomics approach was used to show that *F. fulva* secretes at least

75 SSPs, named extracellular proteins (Ecps), during infection, of which many are expected to be yet-

identified Avr effectors (Mesarich et al., 2018). In support of this, 10 Ecps were found to elicit an HR

upon expression in various wild tomato accessions (Mesarich et al., 2018).

We recently demonstrated that *F. fulva* overcame the widely deployed *Cf-9* resistance locus by

sequential loss of *Avr9* and loss or mutation of *Avr9B* (de la Rosa & Schol et al., 2024). Both of these

Avr genes encode SSPs and are recognized by the products of two RLP-encoding genes at the *Milky Way*

(MW) locus on the short arm of chromosome 1, *Cf-9C* and *Cf-9B*, respectively (Jones et al., 1994;

Parniske et al., 1997). This prompted breeders to deploy a novel resistance gene, *Cf-6*, which was

tentatively mapped to the short arm of chromosome 12 using the *S. lycopersicum* breeding line Ontario

7818 (CGN15839) and was reported to originate from *S. pimpinellifolium* PI211839 (Kanwar et al.,

1980). Recently, the International Seed Federation (ISF) updated the *F. fulva* and tomato differential

sets, used for determining virulence and resistance spectra, to include five new strains of *F. fulva* (races

9, 2.9, 4.9, 4.5.9 and 2.6.9) and Ontario 7818 as the differential line for *Cf-6* (Sangster, 2022). The

inclusion of a race 2.6.9 strain (breaking *Cf-2*, *Cf-6* and *Cf-9*-mediated resistance), indicates that the

resistance provided by *Cf-6* is race-specific. The first report of *Cf-6*-breaking *F. fulva* strains in Europe

came from Germany, where several race 2.6.9 strains were collected throughout the country in 2018

and 2019 (Meyer & Gärber, 2021). For the current study, we were provided with an *F. fulva* collection

containing several race 2.6.9 strains, collected between 2017 and 2021 in the Brittany region in France.

The emergence of these strains indicates that the recently deployed *Cf-6* locus has already been

overcome, which is vital information for breeders to adjust their breeding strategies accordingly.

To determine the genetic and molecular basis of *Cf-6* resistance breakdown, we set out to identify Avr6,

the Avr effector recognized by tomato plants carrying the *Cf-6* resistance locus, using comparative

genomics. Comparative transcriptomics and genomics have previously been successfully applied to

identify the *F. fulva Avr* genes *Avr5* and *Avr9B* but have been limited to comparison of a single

resistance-breaking strain with a single non-resistance breaking strain (de la Rosa & Schol et al., 2024;

Mesarich et al., 2014). To identify *Avr6*, we performed whole-genome sequencing and assembly of 15

F. fulva race 2, 2.9 and 2.6.9 strains, allowing us to compare polymorphisms in known SSP-encoding

genes, most of which are previously characterized *Ecp* genes (Laugé et al., 2000; Mesarich et al., 2018;
Van den Ackerveken et al., 1993).

Next, we validated the *Avr6-Cf-6* interaction through transient expression of *Avr6* in *Cf-6* tomato plants
and gene knockout and complementation assays in *F. fulva.* Then, we used Avr6 as a marker for
mapping the *Cf-6* gene in Ontario 7818 by means of bulked segregant analysis (BSA)-sequencing,
followed by a Comparative Subsequence Set Analysis (CoSSa) bioinformatics pipeline (Prodhomme et
al., 2019). Finally, we correlated resistance to *F. fulva* in the mapping population with the ability to
recognize Avr6. This work serves to alert breeders to the breakdown of the *Cf-6* gene, provide tools
that can be applied to quickly identify *Cf-6*-breaking *F. fulva* strains in the field, and to direct ongoing
resistance breeding efforts.

Results

Deletion or mutation of *Extracellular protein 5* (*Ecp5*) is associated with *Cf-6*-breaking *F. fulva* strains

As a starting point for identifying *Avr6*, we obtained 14 French *F. fulva* strains that were sampled from
diseased tomato plants in the French region of Brittany between 2017 and 2021, of which five were
reported to break *Cf-6,* and a strain from New Zealand that was reported to break *Cf-6* in 2012. We
confirmed that six of these strains indeed break *Cf-6*-mediated resistance with a disease assay on MM-
Cf-0 and Ontario 7818 (*Cf-6*) tomato plants (**Figure S1**). Subsequently, all 15 *F. fulva* strains were
subjected to Illumina sequencing and we generated whole-genome assemblies (**Table S1**), and
compiled these with publicly available *F. fulva* assemblies (**Table S2**) into a dataset of 32 genomes.
Then, we inferred a phylogeny based on a concatenated alignment of 3484 conserved, single-copy
orthologs and found that the strains fall into four distinct clades, with most (11) clustering into a single,
closely related clade (**Figure 1**). Then, we inferred mating types for all strains in the dataset and found
that 11 had *Mat1-1* (36.4%) and 21 had *Mat1-2* (63.6%) at their mating type (MAT) locus. Further
inspection of the distribution of genotypes for known *Avr* genes *Avr2*, *Avr4*, *Avr4E*, *Avr5*, *Avr9* and *Avr9B*
revealed that predicted mutations in Avr2 (p. del54-78, Tyr41fs*34, Lys23fs*42), Avr4 (Cys109*) and
Avr4E (Phe82Leu and Met93Thr) occur in both mating types, which suggests occurrence of
recombination and thus sexual reproduction within *F. fulva* populations.

Next, we queried these genomes to identify candidate genes for *Avr6*: i.e. putative effector genes that
have non-synonymous mutations or are (partially) absent in all race 2.6.9 strains, but not in others.
Hypothesizing that *Avr6* should be an SSP, we used BLAST to compare the sequences of a set of known
and predicted Ecp genes from the *F. fulva* race 5 reference strain (Mesarich et al., 2018; Zaccaron et al.,
2022) to our dataset, and searched for those that were absent or exclusively had one or more mutations
in the race 2.6.9 strains (**Figure S2**). As previously reported/predicted, the *Ecp* genes, although

displaying some minor sequence variation, are present in most or all strains (Mesarich et al., 2018;

Stergiopoulos, De Kock, et al., 2007), with *Ecp24-2* and *Ecp5* being notable exceptions. Strikingly, we

found that *Ecp5* (Laugé et al., 2000) was deleted in four of the six *Cf-6*-breaking strains (**Figure 1**, **Figure**

S2). In these strains, we observed a deletion of approximately 18 kb on chromosome 2, resulting in the

loss of *Ecp5* (*CLAFUR5_03423*) and three adjacent genes, *CLAFUR5_20157*, *CLAFUR5_03424*, and

CLAFUR5_03425 (**Figure S3**)*.* As previously shown, *Avr6* is surrounded by transposable elements (TEs)

of various types (de Wit et al., 2012), which are likely responsible for its deletion. The latter three are

predicted to code for uncharacterized, non-secreted proteins and are therefore unlikely candidates for

Avr6. In the other two *Cf-6*-breaking strains, non-synonymous mutations in *Ecp5*, resulting in amino

acid substitutions Tyr88His and Gly53Val, were observed. Strain IPO 2679, which was used for the

identification of *Avr9B* (de la Rosa & Schol et al., 2024), was the only other strain showing sequence

variation for *Ecp5*, resulting in a Cys30Tyr substitution. In all other strains, a wild-type *Ecp5* allele was

present. For *Ecp24-2*, we did not observe a clear correlation with the ability to overcome *Cf-6* (**Figure**

S2), and it was omitted as a candidate for *Avr6*. Together, these results suggest that *Ecp5* is *Avr6*.

0.0002

Figure 1: Phylogeny and predicted amino acid sequence comparisons of known Avrs, and Ecp5, across a panel of 32 *F. fulva* genomes.

Maximum-likelihood phylogeny inferred from a concatenated alignment of 3484 BUSCO genes, rooted with *Dothistroma septosporum* as outgroup (not shown in the tree). Only branches with bootstrap values < 95 are shown in the tree. Sequences generated for this study are shown in bold. Mating types are indicated with symbols at the tips of the tree (circles and triangles), and the country of origin is indicated with colored rectangles. Distinct predicted amino acid sequence mutations in known Avr proteins Avr2, Avr4, Avr4E, Avr5, Avr9 and Avr9B, and in Ecp5 are indicated with squares in different colors in the right panel, with wild-type (WT) sequences indicated in grey, and absence of an open reading frame (ORF) indicated in white. p., protein sequence numbering; del, deletion; fs, frame shift; *, stop codon.

Molecular validation confirms that Ecp5 is Avr6

To confirm that Ecp5 is Avr6, we systemically expressed it in *Cf-6* tomato (Ontario 7818) and *S. lycopersicum* G1.1161, the line carrying the *Cf-Ecp5* locus which was mapped to the short arm of chromosome 1 (Haanstra et al., 2000). Here, potato virus X (PVX)-based expression was employed, whereby plants are transiently transformed with a recombinant version of PVX containing *Ecp5* (PVX::*Ecp5*) using *Agrobacterium tumefaciens* (Haanstra et al., 2000; Laugé et al., 2000). As controls, *Ecp5* was also expressed in MM-*Cf-0* plants that are universally susceptible to *F. fulva* and in MM-*Cf-9* plants carrying the *Cf-9* locus. Concomitant with the systemic movement of the recombinant virus in the plant, Ecp5 is produced systemically. Should Ecp5 be recognized by Cf-6, a systemic, visible HR, resulting in chlorosis, necrosis, stunting, or death of the plant is expected to occur (Haanstra et al., 2000; Laugé et al., 2000).

An HR developed in Ontario 7818 and G1.1161 plants, but not in MM-*Cf-0* and MM-*Cf-9* plants, supporting that Ecp5 is Avr6 (**Figure 2**). To further validate that *Cf-6*-mediated resistance is solely based on Ecp5 recognition, we inoculated MM-*Cf-0* and Ontario 7818 plants with the wild-type race 0 *F. fulva* strain 0WU that produces Ecp5 (de Wit et al., 2012), and two independent *Ecp5* knockout strains in this background, generated previously by Ökmen (2013). As expected, the wild-type 0WU strain was virulent on MM-*Cf-0* plants, but avirulent on Ontario 7818 plants, whereas the two *Ecp5* knockout strains both had gained virulence on Ontario 7818 plants, with symptoms visually indistinguishable from those caused by the wild-type 0WU strain on *Cf-0* plants (**Figure 3**). This confirms that only Ecp5 is recognized by *Cf-6* tomato plants to mediate resistance to *F. fulva.* From here on, we will refer to Ecp5 as Avr6.

Figure 2: Ecp5 elicits a systemic HR in young tomato plants carrying *Cf-6* and *Cf-Ecp5*. Ecp5 was systemically expressed in seedlings of MM-*Cf-0* (no *Cf* genes), MM-*Cf-9* (*Cf-9B* and *Cf-9C*), Ontario 7818 (*Cf-6*), and G1.1161 (*Cf-Ecp5*), using the PVX-based expression system, with three replicates per tomato line. The recombinant virus PVX::*EV* (empty vector) was used as a negative control, while the recombinant virus PVX::*Avr9* was used as positive control. All recombinant viruses were delivered by *A. tumefaciens*-mediated transient transformation assays, through infiltration of the cotyledons of 10-day-old seedlings. Photographs were taken 10 days post agro-infiltration. White asterisks indicate plants undergoing systemic HR.

Figure 3: Deletion of *Ecp5* in the *F. fulva* race 0 strain 0WU confers virulence on Ontario 7818 (*Cf-6*) tomato plants. The 0WU wild-type strain, two independent *F. fulva Ecp5* knockout strains (*Δecp5*-2 and *Δecp5*-3) and an ectopic *green fluorescent protein* (*GFP*) transformant control were inoculated on 4-week-old Ontario 7818 (*Cf-6*) and MM-*Cf-0* (no *Cf* genes) tomato plants. The upper and lower surfaces of leaflets were photographed at 14 days post-inoculation.

Allelic variation in *Avr6* allows specific *F. fulva* strains to overcome *Cf-6*

Next, we validated that mutations in Avr6 observed in race 2.6.9 strains allow circumvention of *Cf-6*-mediated resistance. *F. fulva* strain IPO 2679 carries a Cys30Tyr mutation in Avr6, which we hypothesized would allow it to break *Cf-6*-mediated resistance. To test this, we inoculated transgenic IPO 2679 strains complemented with the wild-type *Avr6* gene from strain 0WU, generated by de la Rosa & Schol et al. (2024), onto MM-*Cf-0* and Ontario 7818 (*Cf-6*) plants. We observed that the wild-type IPO 2679 strain is virulent on both lines, whereas the two complemented transformants have lost the ability to infect Ontario 7818 (*Cf-6*), but not MM-*Cf-0* plants (**Figure 4A**). Similarly, when strains FR_269_1 (absent *Avr6* ORF) and NZ_269_1 (Avr6^Tyr88His) were complemented with wild-type *Avr6*, virulence was abolished on *Cf-6* plants, but not on MM-*Cf-0* (**Figure 4B**). We conclude that these three mutations in Avr6 each allow the fungus to break *Cf-6*-mediated resistance. This contrasts with previous results, in which no allelic variation had been described in *Avr6* across a large core collection of *F. fulva* strains (Iida et al., 2015; Stergiopoulos, De Kock, et al., 2007). To determine whether there is evidence for further selection pressure on *Avr6* in response to *Cf-6* deployment, we sequenced *Avr6* in 155 additional *F. fulva* strains that were predominantly collected over the past two decades and mostly originate from Europe and Japan. A wild-type *Avr6* gene was present in almost all strains (**Table S3**), suggesting that these are unable to cause disease in *Cf-6* plants. Only in one recently collected German strain (P3), we identified a predicted splice site mutation at the end of the second intron of *Avr6* which resulted in (partial) restoration of virulence on *Cf-6* plants (**Figure S4**). Although the observed

symptoms were not as pronounced as on MM-*Cf-0*, they were more severe compared to the symptoms caused by strain P30, which has a wild-type *Avr6* gene, indicating that this mutation impairs the recognition of Avr6 by Cf-6. In all 154 other strains, we found a wild-type *Avr6* gene, suggesting that *Cf-6*-breaking strains recently emerged.

Figure 4: Complementation of *Cf-6*-breaking *F. fulva* strains with wild-type *Avr6* from strain 0WU results in loss of virulence on Ontario 7818 (*Cf-6*) tomato plants. The IPO 2679 (Avr6[Cys30Tyr]) wild-type strain, two independently generated IPO 2679 *Avr6*-complemented *F. fulva* (*Avr6*-1 and *Avr6*-2) transformants, and an empty vector (EV) control transformant were inoculated on 4-week-old Ontario 7818 (*Cf-6*) and MM-*Cf-0* (no *Cf* genes) tomato plants (Panel A). Two independently generated NZ_269_1 and FR_269_1 *Avr6*-complemented *F. fulva* (*Avr6*-1 and *Avr6*-2) transformants, and an EV control, were inoculated on 4-week-old Ontario 7818 (*Cf-6*) and MM-*Cf-0* (no *Cf* genes) tomato plants (Panel B). Leaflets were photographed at 14 days post-inoculation.

Recently, however, non-synonymous and frame-shift mutations were reported in *Avr6* of several *F. fulva* strains collected in Argentina (Lucentini et al., 2021; Medina et al., 2015). To investigate whether these

mutations also result in virulence on *Cf-6* tomato, we inoculated strains expressing Avr6[Lys22*],
Avr6[Cys30fs*74], Avr6[Val40fs*101] and Avr6[Val98fs*104] (**Table S3**) onto MM-*Cf-0* and Ontario 7818 plants. This
resulted in visible disease symptoms on both genotypes for strains harboring these mutant alleles, but
not for the control strains 0WU or CIDEFI 305, which both contain wild-type *Avr6* and are therefore
unable to overcome *Cf-6* (**Figure 5**). As all mutations are predicted to result in either frameshifts or
cysteine mutations, it is not surprising that recognition by Cf-6 is abolished. Unfortunately, we could
not test strain CIDEFI 318 (Avr6[Cys48Tyr]) due to its poor growth and sporulation *in vitro*, but it is likely
that the Cys48Tyr mutation also abolishes recognition. In all, apart from deletion of the *Avr6* ORF and
the splice site mutation in P31, we identified eight different non-synonymous and frame-shift
mutations in *Avr6* across strains (**Figure 6**), of which seven were confirmed to result in breakdown of
Cf-6 by *F. fulva*.

238
Figure 5: **Non-synonymous and frame-shift mutations in *Avr6* of Argentinian *F. fulva* strains confer**
virulence on *Cf-6* tomato plants. Four-week-old MM-*Cf-0* (no *Cf* genes) and Ontario 7818 (*Cf-6*) tomato
plants were inoculated with several of the Argentinian *F. fulva* strains reported by Medina et al. (2015)
and Lucentini et al. (2021) to harbor mutations in *Avr6* (race 2.6 (initially reported as race 2): CIDEFI
300, 301, 307 and 329; race 6 (initially reported as race 0): CIDEFI 326). As controls, strain 0WU (race
0) and CIDEFI 305 (race 2), which can infect MM-*Cf-0* but not *Cf-6* plants, were also inoculated.
Symptoms were photographed at 14 days post-inoculation.

Figure 6: Schematic representation of the Avr6 protein showing the location of mutations that allow

F. fulva to overcome Cf-6-mediated resistance. The signal peptide (SP) is indicated in grey and yellow

bars indicate the positions of cysteine residues. The strain carrying the predicted Cys48Tyr mutation (in

red font) could not be tested.

Many plant-pathogenic Dothideomycete and Sordariomycete fungi are predicted to produce an Avr6-like protein

To investigate whether other fungi produce Avr6-like proteins, a BLAST search was performed against fungal genomes and their associated protein catalogues in publicly available sequence databases using *F. fulva* Avr6 as a query. This resulted in the identification of a large number of Avr6-like proteins, all belonging to plant-pathogenic Dothideomycetes and Sordariomycetes (**Figure S5**). One such protein was identified in *Pseudocercospora fuligena* (**Figure S5**), a Dothideomycete responsible for black leaf mold disease of tomato (Phengsintham et al., 1938). All identified Avr6-like proteins were predicted to possess an N-terminal signal peptide (SP) for extracellular targeting (**Notes S1**). Furthermore, based on a sequence alignment, the six cysteine residues present in Avr6 were conserved in all Avr6-like proteins (**Figure S5**). Aside from these conserved cysteine residues, several other amino acid residues were highly conserved, including Tyr88 and Gly53, which were mutated in *F. fulva* strains NZ_269_1 (Tyr88His) and FR_269_4 (Gly53Val), indicating that they may play a crucial role in the overall structure of Avr6 and/or its role in virulence (**Figure S5**).

The predicted tertiary structure of Avr6 shows similarity to LEPTOSPHAERIA AVIRULENCE AND SUPPRESSING (LARS) effectors

To gain insight into the potential function of Avr6, its tertiary structure, without its SP for extracellular targeting, was predicted using AlphaFold2 and compared to characterized tertiary structures of fungal proteins in the RCSB Protein Data Bank using Foldseek. Based on this, Avr6 is predicted to adopt a tertiary fold made up of six β-strands and one α-helix, stabilized by three disulfide bonds (**Figure 7**). Following the numbering of the mature Avr6 sequence, the disulfide connectivity pattern is predicted to be Cys13-Cys18, Cys31-Cys59 and Cys61-Cys94. Of note, AlphaFold2 did not predict the Cys61-Cys94 bond, but instead this association is anticipated based on the close proximity of these two residues in the predicted Avr6 tertiary structure (**Figure 7**).

Next, the mutated, conserved amino acid positions identified in Avr6 of *F. fulva* strains NZ_269_1

(Tyr88His) and FR_269_4 (Gly53Val) that are virulent on *Cf-6* tomato, were mapped onto the predicted

tertiary structure to examine whether they are likely to be surface-exposed and, thereby potentially

accessible to the Cf-6 receptor (should recognition be direct) (**Figure 7**). Interestingly, Tyr88 is predicted

to occupy an internal position within the Avr6 tertiary structure and therefore is not expected to be

surface-exposed. Gly53, on the other hand, is predicted to form part of the loop linking the sole α-helix

to β-strand 2 and is therefore predicted to be at least partially surface-exposed (**Figure 7**).

To investigate whether the predicted fold of Avr6 is found in other proteins, particularly those with a

characterized biological function, we used FoldSeek to compare its predicted tertiary structure to

experimentally determined tertiary structures of fungal proteins in the RCSB Protein Data Bank. Based

on this approach, we identified the top hits to Avr6 as AvrLm4-7 (PDB ID: 4FPR; Probability score: 0.77,

TM-Score: 0.549) and AvrLm5-9 (PDB ID: 7B76; Probability score: 0.75, TM-score: 0.0547), which are

both Avr effectors from the LEPTOSPHAERIA AVIRULENCE AND SUPPRESSING (LARS) family of the

Brassica blackleg pathogen *Leptosphaeria maculans* (Blondeau et al., 2015; Lazar et al., 2022). While

these proteins adopt a similar fold to Avr6, the structures of AvrLm4-7 and AvrLm5-9 are more

elongated and Avr6 possesses additional β-strands (**Figure 7**). Unfortunately, the specific host targets

of AvrLm4-7 and AvrLm5-9 are not yet known.

Figure 7: Avr6 is predicted to adopt a LARS-like fold. (a) Predicted tertiary structure of Avr6. The

tertiary structure is colored according to the predicted Local Distance Difference Test (pLDDT) score

(confidence: cyan = high; dark blue = very high). Gly53 and Tyr88 are shown as sticks and are colored

red. β-strands are numbered sequentially. Predicted disulfide bonds are colored yellow. N- and C-

termini are shown. **(b)** Alignment of Avr6 (blue) with AvrLm4-7 (magenta; RCSB Protein Data Bank (PDB)

ID: 4FPR) from *L. maculans.* **(c)** Alignment of Avr6 (blue) with AvrLm5-9 (green; PDB ID: 7B76) from *L.*

maculans. pTM: predicted Template Modelling score. RMSD: Root Mean Square Deviation.

Bulked segregant analysis maps *Cf-6* in Ontario 7818 to a 2 Mb interval on the short arm of chromosome 12

Currently, three distinct loci originating from different *S. pimpinellifolium* accessions which mediate a gene-for-gene relationship with *Avr6* have been reported: *Cf-Ecp5.1* on the top arm of chromosome 1 (Haanstra et al., 2000; Iakovidis et al., 2020), *Cf-Ecp5.7* on the bottom arm of chromosome 7, and *Cf-Ecp5.12* on the top arm of chromosome 12 (Iakovidis et al., 2020). To validate the location of *Cf-6* on the short arm of chromosome 12 in Ontario 7818 as reported by Kanwar et al. (1980), we applied a Bulked Segregant Analysis (BSA)-sequencing approach in an F2 population derived from a cross between Ontario 7818 (*Cf-6*) and MM-*Cf-0*. For this, 133 F2 seedlings were subjected to systemic PVX-based expression of *Avr6* and subsequently phenotyped based on presence or absence of a systemic HR (**Figure S6**, **Table S4**). Genomic DNA (gDNA) from 88 (out of 92) plants that underwent a systemic HR, indicating Avr6-responsiveness, was bulked into a responsive (R) pool, and from 40 (of 41) non-responders into a non-responsive (NR) pool (**Tables S4 and S5**). The expected 3R:1NR segregation pattern, under the hypothesis of *Cf-6* being a single dominant gene, was confirmed with a χ^2 test (χ^2=2.41, *p*-value=0.12) (**Table S6**). After short-read sequencing of the pools, we applied a Comparative Subsequence Sets analysis (CoSSa) bioinformatics pipeline (Prodhomme et al., 2019), to identify R-bulk-specific *k*-mers, which were subsequently mapped to the *S. lycopersicum* Heinz SL4.0 reference genome (Hosmani et al., 2019). We identified a small 2 Mb interval spanning from approximately 2.2 Mb to 4.2 Mb on the short arm of chromosome 12 that was significantly enriched for R-bulk-specific *k*-mers, indicating that presence of *Cf-6* (**Figure 8**). Potential resistance gene-encoding candidates for *Cf-6* within this region include eight RLP-encoding genes, divided over two distinct clusters, and two nucleotide-binding leucine-rich repeat (NB-LRR)-encoding genes. All of these genes show significant allelic variation in Ontario 7818 compared with the Heinz 1706 SL4.0 reference and MM-*Cf-0*, indicating that they could be involved in recognition of Avr6 in Ontario 7818 (**Figure S7**). The region coincides with previous findings mapping *Cf-6* to the short arm of chromosome 12 (Kanwar et al., 1980), and the later finding for the two *Cf-Ecp5.12* alleles (Iakovidis et al., 2020).

Figure 8: BSA-mapping of *Cf-6* (based on responsiveness to Avr6) in an Ontario 7818 x Moneymaker F2 population using Comparative Subsequence Sets analysis (CoSSa). Plots show R-pool-specific *k*-mers (resulting from intersection with Ontario 7818 specific *k-mers* and subtraction of NR-pool and MM-*Cf-0* specific *k*-mers), with respect to their position on the *S. lycopersicum* Heinz 1706 SL4.0 reference genome (Panel A). We found a strong correlation with a region spanning 2.2 to 4.2 Mb on the short arm of chromosome 12 (Panel B). One cluster containing two candidate NB-LRR-encoding genes (green lines) and two clusters both containing four RLP-encoding genes (blue and red lines, respectively) within this mapping interval are indicated (Panel C).

To determine which of these genes could be *Cf-6*, we predicted the domain structure of all ten encoded candidate Cf-6 proteins based on their predicted protein sequences from the Heinz 1706 ITAG4.0 reference annotation (**Figure S8, Tables S7 and S8**). Both NB-LRR proteins are predicted to encode for coiled-coil(CC)-NB-LRRs, whilst of the eight RLPs, only Solyc12g009510 and Solyc12g009520 are predicted to harbor an N-terminal SP for secretion, followed by a LRR domain and a transmembrane domain (TM), features that are deemed essential for RLP functionality. Interestingly, Solyc12g009510 is predicted to harbor a cytoplasmic LRR domain with a second C-terminal TM. The other RLPs do not show a complete predicted domain organization based on the reference, but because the structures of all of these proteins in Ontario 7818 remain unknown, they all remain candidates for Cf-6.

Finally, to verify the correlation between Avr6 recognition and resistance to *F. fulva* in the F2
population, we inoculated 52 new individuals from the same F2 population with PVX::*Avr6*, and
inoculated cuttings of these individuals with the *F. fulva* race 0 strain 0WU (**Figure S9**). For each
individual, we found a strict co-segregation of response to Avr6 with resistance to *F. fulva*, and of non-
responsiveness with susceptibility (**Table S9**). Again, we observed a 3R:1NR (40 R, 12 NR) segregation
of the response to Avr6 and of disease resistance (χ^2=0.103, *p*-value=0.75) (**Table S6**). This indicates
that *Cf-6* is the only resistance gene providing resistance to *F. fulva* in the Ontario 7818 F2 population
and further confirms that this gene confers resistance to *F. fulva* upon recognition of Avr6.

Discussion

F. fulva* overcomes *Cf-6* by loss or mutation of *Avr6

Avr6 codes for a 115-amino acid protein of (so-far) unknown function and was originally cloned as *Ecp5*
(Laugé et al., 2000). Upon removal of its 17-amino acid signal peptide, which is required for secretion,
Avr6 becomes a mature 98-amino acid SSP with six cysteine residues (Laugé et al., 2000). Expression
of *Avr6* is highly induced *in planta*, but not *in vitro* (Mesarich et al., 2014). Consistent with earlier
findings (Iida et al., 2015; Stergiopoulos, De Kock, et al., 2007), we observed a complete lack of allelic
variation in *Avr6* across a set of recently collected Japanese strains, as well as in all but the four French
race 2.6.9 strains and all but one (P31) of the other recently collected European strains we investigated.
This suggests that negative selection pressure on *Avr6* was only recently introduced, probably through
deployment of *Cf-6*, and it can be expected that more *Cf-6*-breaking strains will emerge when *Cf-6*
becomes more widespread in tomato cultivars. The Cys30Tyr mutation in Avr6 of strain IPO 2679 is
puzzling, as it was collected in New Zealand in the 1980s when, to our knowledge, *Cf-6* was not yet
commercially deployed (Iakovidis et al., 2020). Interestingly, a previous study reported an
uncompromised HR after expression of an Avr6[Cys30Ala] mutant (Avr6[Cys13Ala] in the mature protein) in *S.*
lycopersicum Cf-Ecp5 donor line G1.1161 (Luderer et al., 2002), suggesting either a different
recognition specificity of the putative Cf-Ecp5 receptor encoded by the G1.1161 chromosome 1 allele
compared to Cf-6 encoded by the Ontario 7818 chromosome 12 allele, or that Avr6[Cys30Tyr] but not
Avr6[Cys30Ala] abolishes recognition. The variety of polymorphisms in *Avr6* in the previously reported
Argentinian strains raises questions regarding their origin, as the tomato varieties from which they were
sampled are unlikely to harbor *Cf-6/Cf-Ecp5* loci, since they are susceptible to strains with wild-type
Avr6 (Lucentini et al., 2021; Medina et al., 2015). However, *Cf-6/Cf-Ecp5* is common and widespread
across wild tomato species (Iakovidis et al., 2020), so, given Argentina's close proximity to the tomato
center of origin, perhaps these mutations were acquired due to selection pressure exerted by wild
tomato. Subsequently, these strains could have spilled over into tomato cultivation areas. Potentially,

strain IPO 2679 also acquired Avr6^{Cys30Tyr} in South America before spilling over into New Zealand. Strains

originating from South America are so far underrepresented in current *F. fulva* collections, and

considering the variation that was found in the Argentinian strains of Lucentini et al. (2021) and Medina

et al. (2015), it would be interesting to determine the phylogeny and allelic variation of *Avr/Ecp* genes

in other South American strains by performing a sampling campaign for wild *F. fulva* strains (and

potentially their hosts) in the center of origin, followed by sequencing and population studies (Bartoli

& Roux, 2017; Lebeda & Burdon, 2023).

Stacked *Cf* genes can potentially be overcome by recombination between different mutant *Avr* genes

Surprisingly, the closely related clades consisting of five race 2.6.9 and six race 2.9 strains seem to have

overcome both *Cf-6* and *Cf-9B* by two distinct mutations in *Avr* and *Avr9B*: four race 2.6.9 strains show

a deletion of the *Avr6* ORF combined with Avr9B$^{Del129-133,\ L135V}$, whilst the race 2.9 strains have a wild-

type *Avr6* with Avr9B^{Cys107*}, and the last race 2.6.9 strain having Avr6^{Gly53Val} and Avr9B^{Cys107*} (**Figure 1**).

These observations show that single *Cf* gene-mediated resistance can rapidly be overcome

independently by closely related *F. fulva* lineages within a short time frame. It would be interesting to

determine which mutations in *Avr6* cause the recently identified German race 2.6.9 strains to overcome

Cf-6 (Meyer & Gärber, 2021), and to investigate their phylogenetic relationship to the French clade, as

this would shed more light on the epidemiology of tomato leaf mold and the rate at which *Cf* genes are

overcome. Interestingly, all 14 sequenced French strains have Avr2^{Tyr41fs*42}, despite the only two race 2

strains having an opposing mating type (*Mat1-2*) of the other strains (*Mat1-1*). So far, Avr2^{Tyr41fs*42} had

only been described in an Italian race 2 strain with *Mat1-1*, collected in 1993 (Stergiopoulos, De Kock,

et al., 2007). Alarmingly, we see more of the same mutations in Avr proteins occurring in strains of

opposing mating types, specifically Avr2^{Tyr41fs*42}, Avr2^{Lys23fs*34}, Avr2$^{Asp54-Gly78del}$, Avr4^{Cys109*} and

Avr4E^{Phe82Leu,Met93Thr}, suggesting the occurrence of (sporadic) sexual recombination within the previously

considered to be asexual *F. fulva* population. This finding would impact future resistance breeding

strategies, as *F. fulva* would be more readily able to accumulate mutations in *Avr* genes, allowing it to

overcome stacked *Cf* genes. This is supported by previous findings, like the occurrence of Avr9B^{Ser55*} in

both mating types of some Japanese strains (de la Rosa & Schol et al., 2024), and that the two mating

types occur in a 1:1 ratio in a collection of Turkish strains (Gül et al., 2022). In fact, a study on allelic

variation in *Avr* genes in this Turkish population resulted in similar observations to ours (Gül et al.,

2024). The publication of a high-quality chromosome-level reference genome (Zaccaron et al., 2022)

followed by five additional near-complete assemblies provides a solid basis for genomic analyses

(Zaccaron & Stergiopoulos, 2024). Further development of an *F. fulva* pangenome, potentially including

strains originating from the wild, can elucidate whether this species is indeed sexually reproducing.

Avr6 has sequence and structural similarity to putative effectors of other plant-pathogenic fungi

Sequence queries suggest that many plant-pathogenic Dothideomycete and Sordariomycete fungi

produce one or more Avr6-like proteins, of which one, *Pf*Avr6, was identified in *P. fuligena*, a pathogen

closely related to *F. fulva* that causes black leaf mold of tomato (Phengsintham et al., 1938). Given that

the mature version of *Pf*Avr6 has a high level of sequence identity to the mature version of Avr6

(~71.4%), including Gly53 and Tyr88 (**Figure S5**), and that there is evidence of resistance to *P. fuligena*

in cultivated and wild tomato (Hartman & Wang, 1993; Wang et al., 1995), it would be interesting to

determine if it is recognized by *Cf-6* tomato plants. If so, *Cf-6* could also provide protection against *P.*

fuligena. However, the emergence of *Cf-6*-breaking *F. fulva* strains suggests that any such protection

will likely be short-lived as *P. fuligena* may adapt in a similar way. Searches based on the predicted

tertiary structure of Avr6 revealed similarity to other LARS effectors, which have so far been identified

in at least 13 other plant-pathogenic fungi (Lazar et al., 2022). Examples include four Avr effectors from

L. maculans, as well as *F. fulva* Ecp11-1 (Lazar et al., 2022; Mesarich et al., 2018). In the *L. maculans*–

Brassica pathosystem, recognition of the LARS effectors AvrLm3 and AvrLm5-9 by their corresponding

R proteins can be masked by another LARS effector, AvrLm4-7 (Ghanbarnia et al., 2018; Plissonneau et

al., 2016). However, our experiments show no evidence to suggest that Avr6 masks the recognition of

other *F. fulva* effectors in susceptible tomato plants. The predicted tertiary structure of Avr6 displays

several notable differences to the LARS effectors described above, reflected in a lack of sequence

similarity. LARS-like effectors lacking such similarity or possessing subtle differences in their predicted

tertiary folds have been identified in other plant-pathogenic fungi including in *Venturia inaequalis*,

which causes apple scab disease (Rocafort et al., 2022).

431 *Cf-6* maps to the chromosome 12 allele of *Cf-Ecp5* at the *ANDROMEDA* locus

Initially, an HR was observed upon elicitation with the then-called Ecp5 protein (Laugé et al., 2000) in

the introgression line G1.1161, an F_3BC_1 resulting from a cross between *S. pimpinellifolium* CGN15529

and MM-*Cf-0* (Haanstra et al., 2000). The *Cf-Ecp5* gene in this accession was mapped to the *AURORA*

(*AU*) locus on the short arm of chromosome 1 (Haanstra et al., 2000), but a later study found that

rather, it was tightly linked with *Cf-9* at the *MW* locus (Iakovidis et al., 2020). This study also reported

two other independent *Cf-Ecp5* loci: *Cf-Ecp5.7* at the *CENTAURI* (*CE*) locus on the bottom arm of

chromosome 7 in *S. pimpinellifolium* LA0722, and *Cf-Ecp5.12* at the *ANDROMEDA* (*AN*) locus located

within an 888 kb region on the short arm of chromosome 12 spanning from approximately 2.1 to 3.0

Mb in *S. lycopersicum* Ontario 7522 and *S. pimpinellifolium* LA2852 (Iakovidis et al., 2020). Of these,

the mapping interval for *Cf-Ecp5.12* coincides largely with our results for *Cf-6* in Ontario 7818. The study

of Iakovidis *et al*. (2020) employed a marker at 3.014 Mb which showed 4.5% recombination with the

Cf-Ecp5.12 locus. Assuming that the *Cf-6* allele from Ontario 7818 and the *Cf-Ecp5* allele from Ontario

7522 are the same and that *Cf-6* is an RLP-encoding gene, it is more likely to reside within the "blue"

RLP-encoding gene cluster at 2.85 Mb, rather than at the "red" cluster at around 3.0 Mb (**Figure 9C**),

which is roughly at the same location as the mentioned marker of Iakovidis et al. (2020). Although

originating from the same breeding program, Ontario 7818 and 7522 do appear to have different

donors, namely *S. pimpinellifolium* PI 211839 and PI 124161, respectively. PI 124161 appears to

originate from Guatemala (**Table S10**), whilst the origin of PI 211839 is unknown. Since the three

mapped *Cf-6*/*Cf-Ecp5* alleles so far all originate from *S. pimpinellifolium*, it would be interesting to map

and clone additional alleles from different tomato species such as *S. peruvianum* or *S. chilense*, as some

accessions of these species also respond to Avr6 (Iakovidis et al., 2020). This could result in the

identification of more distinct *Cf-6/Cf-Ecp5* loci, shedding more light on the evolution and geographical

distribution of *Cf* genes in wild tomato. Finally, our study clarifies previous conflicting findings regarding

the location of *Cf-6*, as neither chromosome 6 (as reported by Grushetskaya et al. (2007)) nor 11 (as

reported by Wang et al. (2007)) seem to be involved in recognition of Avr6. (Iakovidis et al., 2020).

The identified candidate RLP-encoding genes within our mapping region in Ontario7818 are prime

candidates for *Cf-6*, particularly *Solyc12g009510* and *Solyc12g009520*, given their predicted complete

protein structures in Heinz 1706, although it is also possible that *Cf-6* of Ontario 7818 is not annotated

in the Heinz 1706 genome. This could be resolved by generating a *de novo* genome assembly of Ontario

7818, or by resistance gene enrichment sequencing (RENseq) (Lin et al., 2020; Witek et al., 2016),

revealing any structural differences that are often found across different accessions/species in RLP-

encoding genes (de Kock et al., 2005; Dixon et al., 1998; Dixon et al., 1996; Man et al., 2020). For

example, the recognition of Avr9 in wild *S. chilense* accessions is highly region-dependent and the *Cf-9*

locus is highly polymorphic amongst these accessions (Kahlon et al., 2020).

Future strategies for developing novel, more durable resistance to tomato leaf mold

Future breeding efforts should be directed towards achieving more durable resistance against *F. fulva*,

as breakdown of the recently deployed *Cf-6* locus, to our knowledge, occurred much more rapidly than

breakdown of the *Cf-9* locus (de la Rosa & Schol et al., 2024), demonstrating that single *Cf* gene

strategies against *F. fulva* are likely insufficiently durable. The wild tomato germplasm that showed a

specific HR when exposed to one or multiple of in total nine novel Ecp effectors (ten if Ecp17-1 is

included) identified by Mesarich et al. (2018), likely harbors an equal number of unused *Cf-Ecp* genes

that could be deployed in breeding programs. However, the potential for durability of these genes

should be carefully considered before deployment and could potentially be predicted by generating

Ecp knockout strains to determine their contribution to *F. fulva* virulence, as was recently performed

for *Ecp20-2* (Karimi-Jashni et al., 2022). In general, the paradigm that Ecp effectors are more conserved

and thus would be more essential for virulence than Avr effectors (Ökmen & de Wit, 2012;

Stergiopoulos, De Kock, et al., 2007), seems to be broken by our findings described here.

In summary, *Cf* gene pyramiding currently seems to be the most durable resistance breeding strategy

against *F. fulva*, as sequential loss of multiple effector genes seems to occurs less rapidly than loss of

single effector genes (de la Rosa & Schol et al., 2024). To truly achieve long-lasting, broad-spectrum

resistance in tomato against *F. fulva*, and to serve as an example for other economically important

fungal crop diseases, it is now time to study the dynamics of the *F. fulva*-tomato pathosystem in the

wild. Such an ecology-informed approach, potentially combined with pathogen-host interaction

modeling (Märkle et al., 2024; McMullan et al., 2018), will lead to a better understanding of which

effectors are more important, and thus which corresponding (combination of) *Cf* genes could

potentially be more effective and durable, as tomato accessions from locations with naturally high

levels of *F. fulva* disease pressure have likely adapted to possess more advanced/effective resistance

mechanisms. This should be supported by molecular studies into the function of *F. fulva* effectors and

their virulence targets, which could also reveal mechanisms, or, potentially, susceptibility factors

(Koseoglou et al., 2022; Sun KaiLe et al., 2014), that can be exploited to achieve durable resistance.

Materials and Methods

Fungal and plant material

Information on the *F. fulva* strains used in this study can be found in **Table S3**. Novel French strains

(sample IDs starting with FR_) were collected by VEGENOV and their virulence spectrum was

determined by inoculation onto a tomato differential set carrying different *Cf* genes (including *Cf-6*),

according to the ISF (Sangster, 2022). Argentinean *F. fulva* strains were previously collected and

characterized by Medina et al. (2015) and Lucentini et al. (2021). Their virulence spectrum was

determined by the authors using a tomato differential set (excluding *Cf-6*) and, in this study we assessed

their virulence on Ontario 7818 (CGN15839; carrying *Cf-6*) and Moneymaker (MM) plants (CGN14330).

Transgenic *Avr6* knockout strains in the 0WU background were previously generated by Ökmen (2013).

Strain 0WU (race 0) (de Wit et al., 2012) was used as a control strain in the inoculation studies and

strain IPO 2679 (de la Rosa & Schol et al., 2024) was included as it carries Avr6^{Cys30Tyr}.

For BSA-sequencing, an F2 population of 133 individuals derived from a cross between Ontario 7818

(*Cf-6*) and MM-*Cf-0* was utilized. *S. lycopersicum* Ontario 7818 (*Cf-6*), breeding line G1.1161 (*Cf-Ecp5*)

(Haanstra et al., 2000) and MM-*Cf-9* (CGN15338) were utilized for PVX-mediated expression of Avr6,

Avr9 and Avr9B. Detailed information regarding CGN accessions as provided by the Center of Genetic

Resources Netherlands (CGN) in Wageningen, The Netherlands, can be found in **Table S10.**

DNA isolation, whole-genome sequencing and assembly

Genomic DNA was isolated from *F. fulva* strains, MM-*Cf-0*, Ontario 7818 (*Cf-6*) and Ontario 7818 x MM-*Cf*-0 F2 plants using a cetyltrimethylammonium bromide (CTAB) DNA protocol (Murray & Thompson, 1980) and purified using a Genomic DNA Clean & Concentrator (Zymo Research). For BSA-sequencing, 200 ng of gDNA per F2 plant was pooled into the according R- or NR-pool and then purified (**Tables S4 and S5**). gDNA quality and concentration were measured using the Nanodrop™ One (Thermo Scientific) and Qubit 2.0 Fluorometer (Thermo-Scientific). All 150 bp paired-end (PE)-sequencing of *F. fulva* strains was performed by Novogene (https://www.novogene.com) on an Illumina NovaSeq6000 sequencing platform. All 150 bp PE-sequencing of (pooled) tomato gDNA was performed by BGI (https://www.bgi.com) on a DNBseq sequencing platform. For each of the 15 *F. fulva* strains, reads were trimmed with Trimmomatic v0.39 (Bolger et al., 2014) with parameters set to SLIDINGWINDOW:4:20, MINLEN:100 using Truseq2 adapter sequences, and quality of trimmed reads was checked with FASTQC (v0.12.1) (Andrews, 2019). Trimmed reads were assembled with Spades v4.0.0 (Bankevich et al., 2012) and contigs < 1 kb were removed post assembly. Assembly quality was assessed with Quast (v5.3.0) (Gurevich et al., 2013) and BUSCO (v5.7.1, genome-mode, with dothideomycetes_odb10 as reference BUSCO set and metaEUK as gene predictor) (Manni et al., 2021) (**Table S1**). All *F. fulva* sequencing data generated in this study is deposited in Genbank (PRJNA1271984).

Compiling a dataset of public genomes and inference of phylogeny

Publicly available genome sequences were downloaded from Genbank using the query '*Fulvia fulva*' (**Table S2**). The genome of *Dothistroma septosporum* NZE10 (GCA_000340195.1) was downloaded separately. We used BUSCO as described above to identify single-copy BUSCO genes in these assemblies and those generated in this study. For each BUSCO query gene, we created a fasta file with corresponding DNA sequences from each strain (33 *F. fulva* + 1 *D. septosporum*) if it was present as single-copy in all assemblies using a custom Python script. Next, we used Muscle (v 5.3.linux64) (Edgar, 2022) with default settings to infer a multiple sequence alignment for each fasta and trimmed alignments with TrimAl (v1.5.rev0) (Capella-Gutiérrez et al., 2009) with settings optimized for maximum likelihood inference (-automated1). Trimmed alignments were then concatenated with a second custom Python script into a single alignment with 6917 parsimony informative sites. We used IQtree (Minh et al., 2020) with ModelFinder (Kalyaanamoorthy et al., 2017) and UFBoot (-m MFP -B 1000 -bnni -alrt 1000) (Hoang et al., 2018) to identify an optimal substitution model (GTR+F+I+R10) and infer a consensus maximum likelihood phylogeny from this concatenated alignment with 1000 bootstrap replicates. For the figures, the outgroup (*Dothistroma septosporum* NZE10) and isolate CBS131901

542   (short-read assembly of 0WU) were removed. All custom scripts can be downloaded from

543   https://git.wur.nl/like.fokkens/clado_avr6.

Identification of *MAT* loci and sequence comparison of known and predicted effector genes

To infer mating types of *F. fulva* strains, we used megablast (v 2.13.0+) (Morgulis et al., 2008) with

default settings to align *Mat1-1* and *Mat1-2* sequences (accessions DQ659350.2 and DQ659351.2

respectively). For identification of effector genes in the assemblies, we used a custom python script

implementing bedtools getfasta (v2.30.0) (Quinlan & Hall, 2010) and a list of predicted effector genes

from Zaccaron et al. 2022 to obtain a fasta file with sequences from the assembly of the Race 5

reference strain (GCA_020509005.2) including introns. We then used megaBLAST (v 2.13.0+) with –

default settings to search for homologs of each of these genes. To obtain a similarity score for each

strain/assembly and each effector gene, relative to the Race 5 reference, we divided the maximum

Smith-Waterman score of the effector gene against the assembly with its maximum score against the

assembly of Race 5. We used a custom Python script to parse the BLAST output table and calculate

these scores (**Figure S1**). For known *Avr* genes we generated MSAs to identify distinct genotypes. To

confirm absence of predicted effectors (**Figure S1**) and to determine the size of the deletion resulting

in loss of *Avr6*, we used Bowtie2 (v2.5.4, with -I 100 -X 1000) (Langmead & Salzberg, 2012) to align

reads to the assembly of the Race 5 strain, and checked the regions where effectors were predicted to

be absent in IGV (Robinson et al., 2011). For some strains certain effectors were not present in the

assembly, while we did find reads mapped to the gene in the Race 5 genome, which is indicated in

Figure S1. We used custom R scripts to plot the phylogeny with different *Avr* genotypes (**Figure 1**), and

with a heatmap with effector gene sequence similarity (**Figure S1**).

Bioinformatics pipeline for BSA-sequencing by means of Comparative Subsequence Sets analysis (CoSSA)

A Comparative Subsequence Sets Analysis (CoSSA) bioinformatics pipeline was applied to find the

resistance-specific haplotype (Prodhomme et al., 2019). Sequence reads were trimmed and sequencing

adaptors were removed using fastp (version 0.23.4) (Chen, 2023) (**Table S11**). *K*-mer Counter (KMC)

and KMC tools (version 3.2.4) were used to produce *k*-mer (*k*=31) tables from the trimmed reads, and

k-mer table subtraction, intersection and coverage-based filtering (Kokot et al., 2017). GenomeScope

2.0 (Ranallo-Benavidez et al., 2020) was used to visualize the *k*-mer coverage profiles. The NR-bulk

specific *k*-mer table was subtracted from the R-bulk specific table, and a *k*-mer table of the susceptible

parent (MM-*Cf-0*) was subtracted from the resulting R-bulk specific *k*-mers, and intersected by a *k*-mer

table of the resistant parent (Ontario 7818). *K*-mer sequences with coverage outside a range of 26 to

43 were removed to reduce error-to-noise ratio. Cutoff values are calculated as 25% margins of the

expected coverage of the *R* haplotype. Filtered *k*-mers were mapped to the SL4.00 tomato reference

genome (Hosmani et al., 2019) using BWA aln (version 0.7.18) (Li & Durbin, 2009) and plotted based

on genomic location and coverage. Used scripts reflect the workflow of (Prodhomme et al., 2019) and

are available on Github (https://github.com/PBR/CoSSA). Structural predictions of Cf-6 candidate

proteins were performed using predicted protein sequences from the ITAG4.0 annotation of the Heinz

1706 SL4.0 genome using InterPro (Paysan-Lafosse et al., 2023).

Homolog searches and AlphaFold2 structural predictions

The Basic Local Alignment Search Tool for nucleotide (BLASTn), translated nucleotide (tBLASTn), and

protein (BLASTp) were used to identify *Avr6-/*Avr6-like sequences in the National Centre for

Biotechnology Information (NCBI) and Joint Genome Institute (JGI) Mycocosm sequence databases.

(Laugé et al., 2000) Signal peptides were predicted using SignalP v4.1 (Nielsen, 2017). The ColabFold

(Mirdita et al., 2022) implementation of AlphaFold2 (Jumper et al., 2021) was used to predict the

tertiary structure of Avr6 (without its endogenous signal peptide) and was assisted through use of a

custom MSA generated with Avr6-like proteins, using Geneious v9.1.8 (Kearse et al., 2012). Foldseek

(Van Kempen et al., 2024) was used to identify structural homologs of Avr6 in filamentous fungi in the

RCSB Protein Data Bank. The predicted tertiary structure of Avr6, along with those of proteins identified

using Foldseek, were visualized, rendered, and aligned (CEalign plugin) using PyMOL (DeLano, 2002).

F. fulva disease assays

F. fulva strains were inoculated from glycerol stocks on potato dextrose agar (PDA) plates supplemented

with 1.5% extra technical agar and 33 mg/L chloramphenicol and grown for approximately 2-3 weeks

at room temperature in the dark. Then, conidia were harvested and suspended in tap water to a

concentration of approximately 1×10^6 conidia/mL. This suspension was then spray-inoculated on the

abaxial side of 3- to 4-week-old tomato plants which were subsequently kept under high (90–100%)

humidity in a plastic tent for 3 days, after which the tent was opened. Plants were maintained in a

greenhouse compartment set to 21–23°C. Disease symptoms were scored at 14 days post-inoculation

(dpi).

PVX-mediated expression assays

603 Systemic PVX-based expression of Avr effectors was performed by means of agroinfiltration of the

604 cotyledons of 10-day (for the validation of Avr6) or 16-day (for BSA-sequencing of *Cf-6*) old tomato

605 seedlings, as previously described by Mesarich et al. (2018) and de la Rosa & Schol et al. (2024). The

606 *Agrobacterium* strain GV3101 carrying *pSfinx::Ecp5* was previously generated by Laugé et al. (2000) and
607 the strain carrying *pSfinx::Avr9* by Hammond-Kosack et al. (1995). Symptoms were scored at 10 dpi.

F. fulva gene complementation

609 Complementation of strains with the wild-type *Avr6* gene was performed by means of *Agrobacterium*-
610 mediated transformation using the same method and construct as described in (de la Rosa & Schol et
611 al., 2024).

Avr6 allelic variation analysis

Avr6 was amplified from the genomic DNA of 90 *F. fulva* strains using primers Avr6 Fw2 (5'-
CTGACATAGCATTACACGAGCAG-3') and Avr6 Rev (5'-CGCGTCGCCTGATAGATTTG-3) by PCR, as previously
described for the *Avr9B* gene (de la Rosa & Schol et al., 2024). Amplicons were purified using ExoSAP-
IT™ PCR Product Cleanup Reagent (Thermo Scientific) and sequenced by Macrogen™ Europe
(https://www.macrogeneurope.com) by means of Sanger Sequencing using an ABI 3730xl DNA
analyzer (Applied Biosystems). Consensus sequences were generated and imported in Benchling
(Biology Software, https://benchling.com) for end-trimming and sequence alignment to the *Ecp5*
coding sequence (CLAFUR5_03423) from the *F. fulva* strain Race 5 reference genome (Zaccaron et al.,
2022).

Acknowledgements

We would like to thank Bert Essenstam and colleagues of the Unifarm greenhouse facilities at
Wageningen University for excellent plant care, Dr Pedro Ballati of the Universidad Nacional de La Plata,
Buenos Aires, Argentina for kindly providing the Argentinian *F. fulva* strains, Dr Phillipe Nicot of INRAE
Centre de Recherche PACA, Montfavet, France for providing a selection of *F. fulva* strains from the
Southern region of France and the growers of the Cerafel association, St. Martin-des-Champs, France,
for providing *F. fulva* strains from the Brittany region. Finally, we thank Prof Ioannis Stergiopoulos and
Dr Alex Z. Zaccaron of The College of Agricultural and Environmental Sciences at UC Davis, California
for generating the public *F. fulva* sequencing data that was used in this study.

Funding statement

This research was largely funded by the EU Horizon 2020 HARNESSTOM project (Grant agreement no.:
101000716)

References

Andrews, S. F. (2019). V0. 11.9: A quality control tool for high throughput sequence data. *Babraham*
Bioinformatics: Cambridge, UK. Available online: http://www. bioinformatics. babraham. ac.
uk/projects/fastqc/(accessed on 18 January 2013).

Bankevich, A., Nurk, S., Antipov, D., Gurevich, A. A., Dvorkin, M., Kulikov, A. S., Lesin, V. M., Nikolenko,
S. I., Pham, S., & Prjibelski, A. D. (2012). SPAdes: a new genome assembly algorithm and its
applications to single-cell sequencing. *Journal of Computational Biology*, *19*(5), 455-477.

Bartoli, C., & Roux, F. (2017). Genome-wide association studies in plant pathosystems: toward an
ecological genomics approach. *Frontiers in Plant Science*, *8*, 763.

Blondeau, K., Blaise, F., Graille, M., Kale, S. D., Linglin, J., Ollivier, B., Labarde, A., Lazar, N., Daverdin,
G., & Balesdent, M. H. (2015). Crystal structure of the effector AvrLm4–7 of *Leptosphaeria*
maculans reveals insights into its translocation into plant cells and recognition by resistance
proteins. *The Plant Journal*, *83*(4), 610-624.

Bolger, A. M., Lohse, M., & Usadel, B. (2014). Trimmomatic: a flexible trimmer for Illumina sequence
data. *Bioinformatics*, *30*(15), 2114-2120.

Capella-Gutiérrez, S., Silla-Martínez, J. M., & Gabaldón, T. (2009). trimAl: a tool for automated
alignment trimming in large-scale phylogenetic analyses. *Bioinformatics*, *25*(15), 1972-1973.

Chen, S. (2023). Ultrafast one-pass FASTQ data preprocessing, quality control, and deduplication using
fastp. *Imeta*, *2*(2), e107.

Cooke, M. (1883). New american fungi. *Grevillea*, *12*, 22-33.

de Kock, M. J., Brandwagt, B. F., Bonnema, G., de Wit, P. J., & Lindhout, P. (2005). The tomato Orion
locus comprises a unique class of *Hcr9* genes. *Molecular Breeding*, *15*(4), 409-422.

de la Rosa, S., Schol, C. R., Ramos Peregrina, Á., Winter, D. J., Hilgers, A. M., Maeda, K., Iida, Y., Tarallo,
M., Jia, R., Beenen, H. G., Rocafort, M., de Wit, P. J. G. M., Bowen, J. K., Bradshaw, R. E.,
Joosten, M. H. A. J., Bai, Y., & Mesarich, C. H. (2024). Sequential breakdown of the *Cf-9* leaf
mould resistance locus in tomato by *Fulvia fulva*. *New Phytologist, 243*(4), 1522-1538.

de Wit, P., & Flach, W. (1979). Differential accumulation of phytoalexins in tomato leaves but not in
fruits after inoculation with virulent and avirulent races of *Cladosporium fulvum*.
Physiological Plant Pathology, *15*(3), 257-267.

de Wit, P. J. (2016). *Cladosporium fulvum* effectors: weapons in the arms race with tomato. *Annual*
Review of Phytopathology, *54*, 1-23.

de Wit, P. J., Van Der Burgt, A., Ökmen, B., Stergiopoulos, I., Abd-Elsalam, K. A., Aerts, A. L., Bahkali, A.
H., Beenen, H. G., Chettri, P., & Cox, M. P. et al. (2012). The genomes of the fungal plant

669   pathogens *Cladosporium fulvum* and *Dothistroma septosporum* reveal adaptation to different

670   hosts and lifestyles but also signatures of common ancestry. *PLoS Genetics*, *8*(11), e1003088.

671 DeLano, W. L. (2002). Pymol: An open-source molecular graphics tool. *CCP4 Newsletter on Protein*

672   *Crystallography*, *40*(1), 82-92.

673 Dixon, M. S., Hatzixanthis, K., Jones, D. A., Harrison, K., & Jones, J. D. (1998). The tomato *Cf-5* disease

674   resistance gene and six homologs show pronounced allelic variation in leucine-rich repeat

675   copy number. *The Plant Cell*, *10*(11), 1915-1925.

676 Dixon, M. S., Jones, D. A., Keddie, J. S., Thomas, C. M., Harrison, K., & Jones, J. D. (1996). The tomato

677   *Cf-2* disease resistance locus comprises two functional genes encoding leucine-rich repeat

678   proteins. *Cell*, *84*(3), 451-459.

679 Edgar, R. C. (2022). Muscle5: High-accuracy alignment ensembles enable unbiased assessments of

680   sequence homology and phylogeny. *Nature Communications*, *13*(1), 6968.

681 Ghanbarnia, K., Ma, L., Larkan, N. J., Haddadi, P., Fernando, W. G. D., & Borhan, M. H. (2018).

682   *Leptosphaeria maculans* AvrLm9: a new player in the game of hide and seek with AvrLm4-7.

683   *Molecular Plant Pathology*, *19*(7), 1754-1764.

684 Grushetskaya, Z., Lemesh, V., Poliksenova, V., & Khotyleva, L. (2007). Mapping of the *Cf-6* tomato leaf

685   mould resistance locus using SSR markers. *Russian Journal of Genetics*, *43*(11), 1266-1270.

686 Gül, E., Karakaya, A., & Ergül, A. (2022). Determination of mating types of *Fulvia fulva* isolates from

687   Turkey. *Tropical Plant Pathology*, *47*(3), 421-429.

688 Gül, E., Karakaya, A., & Ergül, A. (2024). The race distribution of the *Fulvia fulva* populations and

689   mutations in the Avr genes in Turkey. *Journal of Phytopathology*, *172*(1), e13273.

690 Gurevich, A., Saveliev, V., Vyahhi, N., & Tesler, G. (2013). QUAST: quality assessment tool for genome

691   assemblies. *Bioinformatics*, *29*(8), 1072-1075.

692 Gust, A. A., & Felix, G. (2014). Receptor like proteins associate with SOBIR1-type of adaptors to form

693   bimolecular receptor kinases. *Current Opinion in Plant Biology*, *21*, 104-111.

694 Haanstra, J., Meijer-Dekens, F., Lauge, R., Seetanah, D., Joosten, M. H., de Wit, P. J., & Lindhout, P.

695   (2000). Mapping strategy for resistance genes against *Cladosporium fulvum* on the short arm

696   of chromosome 1 of tomato*: Cf-ECP5* near the *Hcr9* Milky Way cluster. *Theoretical and*

697   *Applied Genetics*, *101*(4), 661-668.

698 Hammond-Kosack, K., Staskawicz, B., Jones, J., & Baulcombe, D. (1995). Functional expression of a

699   fungal avirulence gene from a modified potato virus X genome. *Molecular Plant-Microbe*

700   *interactions*, *8*(1), 181-185.

701 Hartman, G., & Wang, T. (1993). Resistance in Lycopersicon species to black leaf mold caused by

702   *Pseudocercospora fuligena*. *Euphytica*, *71*(1), 125-130.

Hoang, D. T., Chernomor, O., Von Haeseler, A., Minh, B. Q., & Vinh, L. S. (2018). UFBoot2: improving
the ultrafast bootstrap approximation. *Molecular Biology and Evolution*, *35*(2), 518-522.

Hosmani, P. S., Flores-Gonzalez, M., van de Geest, H., Maumus, F., Bakker, L. V., Schijlen, E., van
Haarst, J., Cordewener, J., Sanchez-Perez, G., & Peters, S. (2019). An improved de novo
assembly and annotation of the tomato reference genome using single-molecule sequencing,
Hi-C proximity ligation and optical maps. *bioRxiv*, 767764.

Huang, W. R., & Joosten, M. H. (2024). Immune signaling: receptor-like proteins make the difference.
Trends in Plant Science, *30*(1), 54-68.

Iakovidis, M., Soumpourou, E., Anderson, E., Etherington, G., Yourstone, S., & Thomas, C. (2020).
Genes encoding recognition of the *Cladosporium fulvum* effector protein Ecp5 are encoded at
several loci in the tomato genome. *G3 Genes|Genomes|Genetics*, *10*(5), 1753-1763.

Iida, Y., van 't Hof, P., Beenen, H., Mesarich, C., Kubota, M., Stergiopoulos, I., Mehrabi, R., Notsu, A.,
Fujiwara, K., & Bahkali, A. (2015). Novel mutations detected in avirulence genes overcoming
tomato *Cf* resistance genes in isolates of a Japanese population of *Cladosporium fulvum*. *PLoS*
ONE, *10*(4), e0123271.

Jones, D. A., Thomas, C. M., Hammond-Kosack, K. E., Balint-Kurti, P. J., & Jones, J. (1994). Isolation of
the tomato *Cf-9* gene for resistance to *Cladosporium fulvum* by transposon tagging. *Science*,
266(5186), 789-793.

Joosten, M., Vogelsang, R., Cozijnsen, T. J., Verberne, M. C., & de Wit, P. (1997). The biotrophic fungus
Cladosporium fulvum circumvents Cf-4-mediated resistance by producing unstable AVR4
elicitors. *The Plant Cell*, *9*(3), 367-379.

Joosten, M. H., Cozijnsen, T. J., & de Wit, P. J. (1994). Host resistance to a fungal tomato pathogen lost
by a single base-pair change in an avirulence gene. *Nature*, *367*(6461), 384-386.

Joosten, M. H., & de Wit, P. J. (1989). Identification of several pathogenesis-related proteins in tomato
leaves inoculated with *Cladosporium fulvum* (syn. *Fulvia fulva*) as 1, 3-β-glucanases and
chitinases. *Plant Physiology*, *89*(3), 945-951.

Jumper, J., Evans, R., Pritzel, A., Green, T., Figurnov, M., Ronneberger, O., Tunyasuvunakool, K., Bates,
R., Žídek, A., & Potapenko, A. (2021). Highly accurate protein structure prediction with
AlphaFold. *Nature*, *596*(7873), 583-589.

Kahlon, P. S., Seta, S. M., Zander, G., Scheikl, D., Hückelhoven, R., Joosten, M. H., & Stam, R. (2020).
Population studies of the wild tomato species *Solanum chilense* reveal geographically
structured major gene-mediated pathogen resistance. *Proceedings of the Royal Society B*,
287(1941), 20202723.

Kalyaanamoorthy, S., Minh, B. Q., Wong, T. K., Von Haeseler, A., & Jermin, L. S. (2017). ModelFinder:
fast model selection for accurate phylogenetic estimates. *Nature Methods*, *14*(6), 587-589.

Kang, W.-H., & Yeom, S.-I. (2018). Genome-wide identification, classification, and expression analysis
of the receptor-like protein family in tomato. *The Plant Pathology Journal*, *34*(5), 435.

Kanwar, J., Kerr, E., & Harney, P. (1980). Linkage of the *Cf-12* to *Cf-24* genes for resistance to tomato
leaf mold *Cladosporium fulvum* Cke. *Report of the Tomato Genetics Cooperative*, *30*, 22-23.

Karimi-Jashni, M., Maeda, K., Yazdanpanah, F., de Wit, P. J., & Iida, Y. (2022). An integrated omics
approach uncovers the novel effector Ecp20-2 required for full virulence of *Cladosporium*
fulvum on tomato. *Frontiers in Microbiology*, *13*, 919809.

Kearse, M., Moir, R., Wilson, A., Stones-Havas, S., Cheung, M., Sturrock, S., Buxton, S., Cooper, A.,
Markowitz, S., & Duran, C. (2012). Geneious Basic: an integrated and extendable desktop
software platform for the organization and analysis of sequence data. *Bioinformatics*, *28*(12),
1647-1649.

Kokot, M., Długosz, M., & Deorowicz, S. (2017). KMC 3: counting and manipulating *k*-mer statistics.
Bioinformatics, *33*(17), 2759-2761.

Koseoglou, E., van der Wolf, J. M., Visser, R. G., & Bai, Y. (2022). Susceptibility reversed: modified plant
susceptibility genes for resistance to bacteria. *Trends in Plant Science*, *27*(1), 69-79.

Langmead, B., & Salzberg, S. L. (2012). Fast gapped-read alignment with Bowtie 2. *Nature Methods*,
9(4), 357-359.

Laugé, R., Goodwin, P. H., de Wit, P. J., & Joosten, M. H. (2000). Specific HR-associated recognition of
secreted proteins from *Cladosporium fulvum* occurs in both host and non-host plants. *The*
Plant Journal, *23*(6), 735-745.

Lazar, N., Mesarich, C. H., Petit-Houdenot, Y., Talbi, N., Li de la Sierra-Gallay, I., Zélie, E., Blondeau, K.,
Gracy, J., Ollivier, B., & Blaise, F. (2022). A new family of structurally conserved fungal
effectors displays epistatic interactions with plant resistance proteins. *PLoS Pathogens*, *18*(7),
e1010664.

Lebeda, A., & Burdon, J. J. (2023). Studying wild plant pathosystems to understand crop plant
pathosystems: status, gaps, challenges, and perspectives. *Phytopathology*, *113*(3), 365-380.

Li, H., & Durbin, R. (2009). Fast and accurate short read alignment with Burrows–Wheeler transform.
Bioinformatics, *25*(14), 1754-1760.

Lin, X., Armstrong, M., Baker, K., Wouters, D., Visser, R. G., Wolters, P. J., Hein, I., & Vleeshouwers, V.
G. (2020). *RLP/K* enrichment sequencing; a novel method to identify *receptor-like protein*
(*RLP*) and *receptor-like kinase* (*RLK*) genes. *New Phytologist*, *227*(4), 1264-1276.

Lucentini, C. G., Medina, R., Franco, M. E., Saparrat, M. C., & Balatti, P. A. (2021). *Fulvia fulva* [syn.
Cladosporium fulvum, Passalora fulva] races in Argentina are evolving through genetic
changes and carry polymorphic *avr* and *ecp* gene sequences. *European Journal of Plant*
Pathology, *159*(3), 525-542.

Luderer, R., De Kock, M. J., Dees, R. H., de Wit, P. J., & Joosten, M. H. (2002). Functional analysis of cysteine residues of ECP elicitor proteins of the fungal tomato pathogen *Cladosporium fulvum*. *Molecular Plant Pathology*, *3*(2), 91-95.

Man, J., Gallagher, J. P., & Bartlett, M. (2020). Structural evolution drives diversification of the large LRR-RLK gene family. *New Phytologist*, *226*(5), 1492-1505.

Manni, M., Berkeley, M. R., Seppey, M., Simão, F. A., & Zdobnov, E. M. (2021). BUSCO update: novel and streamlined workflows along with broader and deeper phylogenetic coverage for scoring of eukaryotic, prokaryotic, and viral genomes. *Molecular Biology and Evolution*, *38*(10), 4647-4654.

Märkle, H., John, S., Metzger, L., Ansari, M. A., Pedergnana, V., & Tellier, A. (2024). Inference of host–pathogen interaction matrices from genome-wide polymorphism data. *Molecular Biology and Evolution*, *41*(9), msae176.

McMullan, M., Rafiqi, M., Kaithakottil, G., Clavijo, B. J., Bilham, L., Orton, E., Percival-Alwyn, L., Ward, B. J., Edwards, A., & Saunders, D. G. (2018). The ash dieback invasion of Europe was founded by two genetically divergent individuals. *Nature Ecology & Evolution*, *2*(6), 1000-1008.

Medina, R., López, S. M., Franco, M. E., Rollan, C., Ronco, B. L., Saparrat, M. C., de Wit, P. J., & Balatti, P. A. (2015). A survey on occurrence of *Cladosporium fulvum* identifies race 0 and race 2 in tomato-growing areas of Argentina. *Plant Disease*, *99*(12), 1732-1737.

Mesarich, C. H., Barnes, I., Bradley, E. L., de la Rosa, S., de Wit, P. J., Guo, Y., Griffiths, S. A., Hamelin, R. C., Joosten, M. H., Lu, M., McCarthy, H. M., Schol, C. R., Stergiopoulos, I., Tarallo, M., Zaccaron, A. Z., & Bradshaw, R. E. (2023). Beyond the genomes of *Fulvia fulva* (syn. *Cladosporium fulvum*) and *Dothistroma septosporum*: New insights into how these fungal pathogens interact with their host plants. *Molecular Plant Pathology*, *24*(5), 474-494.

Mesarich, C. H., Griffiths, S. A., van der Burgt, A., Ökmen, B., Beenen, H. G., Etalo, D. W., Joosten, M. H. A. J., & de Wit, P. J. G. M. (2014). Transcriptome sequencing uncovers the *Avr5* avirulence gene of the tomato leaf mold pathogen *Cladosporium fulvum*. *Molecular Plant-Microbe Interactions*, *27*(8), 846-857.

Mesarich, C. H., Ökmen, B., Rovenich, H., Griffiths, S. A., Wang, C., Karimi Jashni, M., Mihajlovski, A., Collemare, J., Hunziker, L., Deng, C. H., ven der Burgt, A., Beened, H. G., Templeton, M. D., Bradshaw, R.E., & det Wit, P. J. G. M. (2018). Specific hypersensitive response–associated recognition of new apoplastic effectors from *Cladosporium fulvum* in wild tomato. *Molecular Plant-Microbe Interactions*, *31*(1), 145-162.

Meyer, U., & Gärber, U. (2021). Bestimmung der in Deutschland vorkommenden Pathotypen des Erregers der Samtfleckenkrankheit an Tomate, *Fulvia fulva*. *Journal of Cultivated Plants/Journal für Kulturpflanzen*, *73*.

Minh, B. Q., Schmidt, H. A., Chernomor, O., Schrempf, D., Woodhams, M. D., Von Haeseler, A., &
Lanfear, R. (2020). IQ-TREE 2: new models and efficient methods for phylogenetic inference in
the genomic era. *Molecular Biology and Evolution*, *37*(5), 1530-1534.

Mirdita, M., Schütze, K., Moriwaki, Y., Heo, L., Ovchinnikov, S., & Steinegger, M. (2022). ColabFold:
making protein folding accessible to all. *Nature Methods*, *19*(6), 679-682.

Morgulis, A., Coulouris, G., Raytselis, Y., Madden, T. L., Agarwala, R., & Schäffer, A. A. (2008). Database
indexing for production MegaBLAST searches. *Bioinformatics*, *24*(16), 1757-1764.

Murray, M., & Thompson, W. (1980). Rapid isolation of high molecular weight plant DNA. *Nucleic
Acids Research*, *8*(19), 4321-4326.

Nielsen, H. (2017). Predicting secretory proteins with SignalP. *Protein Function Prediction: Methods
and Protocols*, 59-73.

Ökmen, B. (2013). Identification and characterization of novel effectors of *Cladosporium fulvum*. PhD
Thesis, Wageningen University and Research.

Ökmen, B., & de Wit, P. J. (2012). *Cladosporium fulvum*–tomato pathosystem: Fungal infection
strategy and plant responses. *Molecular Plant Immunity*, 211-224.

Parniske, M., Hammond-Kosack, K. E., Golstein, C., Thomas, C. M., Jones, D. A., Harrison, K., Wulff, B.
B., & Jones, J. D. (1997). Novel disease resistance specificities result from sequence exchange
between tandemly repeated genes at the *Cf-4/9* locus of tomato. *Cell*, *91*(6), 821-832.

Paysan-Lafosse, T., Blum, M., Chuguransky, S., Grego, T., Pinto, B. L., Salazar, G. A., Bileschi, M. L., Bork,
P., Bridge, A., & Colwell, L. (2023). InterPro in 2022. *Nucleic Acids Research*, *51*(D1), D418-
D427.

Phengsintham, P., Chukeatirote, E., McKenzie, E., Hyde, K., & Braun, U. (1938). Tropical
phytopathogens 2: *Pseudocercospora fuligena*. *J. Sci*, *66*(8).

Plissonneau, C., Daverdin, G., Ollivier, B., Blaise, F., Degrave, A., Fudal, I., Rouxel, T., & Balesdent, M. H.
(2016). A game of hide and seek between avirulence genes *AvrLm4-7* and *AvrLm3* in
Leptosphaeria maculans. *New Phytologist*, *209*(4), 1613-1624.

Prodhomme, C., Esselink, D., Borm, T., Visser, R. G., Van Eck, H. J., & Vossen, J. H. (2019). Comparative
Subsequence Sets Analysis (CoSSA) is a robust approach to identify haplotype specific SNPs;
mapping and pedigree analysis of a potato wart disease resistance gene *Sen3*. *Plant Methods*,
15, 1-20.

Quinlan, A. R., & Hall, I. M. (2010). BEDTools: a flexible suite of utilities for comparing genomic
features. *Bioinformatics*, *26*(6), 841-842.

Ranallo-Benavidez, T. R., Jaron, K. S., & Schatz, M. C. (2020). GenomeScope 2.0 and Smudgeplot for
reference-free profiling of polyploid genomes. *Nature Communications*, *11*(1), 1432.

Robinson, J. T., Thorvaldsdóttir, H., Winckler, W., Guttman, M., Lander, E. S., Getz, G., & Mesirov, J. P.

(2011). Integrative genomics viewer. *Nature Biotechnology*, *29*(1), 24-26.

Rocafort, M., Bowen, J. K., Hassing, B., Cox, M. P., McGreal, B., de la Rosa, S., Plummer, K. M.,

Bradshaw, R. E., & Mesarich, C. H. (2022). The *Venturia inaequalis* effector repertoire is

dominated by expanded families with predicted structural similarity, but unrelated sequence,

to avirulence proteins from other plant-pathogenic fungi. *BMC biology*, *20*(1), 246.

Rooney, H. C., Van't Klooster, J. W., van der Hoorn, R. A., Joosten, M. H., Jones, J. D., & de Wit, P. J.

(2005). *Cladosporium* Avr2 inhibits tomato Rcr3 protease required for Cf-2-dependent

disease resistance. *Science*, *308*(5729), 1783-1786.

Sangster, W. (2022). *ISF Passalora fulva project tomato*. https://worldseed.org/document/isf-

passalora-fulva-project-tomato-final-report-2019-2022/

Stergiopoulos, I., De Kock, M. J., Lindhout, P., & de Wit, P. J. (2007). Allelic variation in the effector

genes of the tomato pathogen *Cladosporium fulvum* reveals different modes of adaptive

evolution. *Molecular Plant-Microbe Interactions*, *20*(10), 1271-1283.

Stergiopoulos, I., Groenewald, M., Staats, M., Lindhout, P., Crous, P. W., & de Wit, P. J. (2007). Mating-

type genes and the genetic structure of a world-wide collection of the tomato pathogen

Cladosporium fulvum. *Fungal Genetics and Biology*, *44*(5), 415-429.

Sun KaiLe, S. K., Zheng Zheng, Z. Z., Jacobsen, E., Visser, R. G., & Bai YuLing, B. Y. (2014). Breeding for

disease resistance by editing plant susceptibility genes. *CABI Reviews,* (2014), 1-5.

Takken, F. L., Thomas, C. M., Joosten, M. H., Golstein, C., Westerink, N., Hille, J., Nijkamp, H. J. J., de

Wit, P. J., & Jones, J. D. (1999). A second gene at the tomato *Cf-4* locus confers resistance to

Cladosporium fulvum through recognition of a novel avirulence determinant. *The Plant*

Journal, *20*(3), 279-288.

Thomas, C. M., Jones, D. A., Parniske, M., Harrison, K., Balint-Kurti, P. J., Hatzixanthis, K., & Jones, J.

(1997). Characterization of the tomato *Cf-4* gene for resistance to *Cladosporium fulvum*

identifies sequences that determine recognitional specificity in Cf-4 and Cf-9. *The Plant Cell*,

9(12), 2209-2224.

Thomma, B. P., Van Esse, H. P., Crous, P. W., & de Wit, P. J. (2005). *Cladosporium fulvum* (syn.

Passalora fulva), a highly specialized plant pathogen as a model for functional studies on

plant pathogenic Mycosphaerellaceae. *Molecular Plant Pathology*, *6*(4), 379-393.

Van den Ackerveken, G., Van Kan, J. A., Joosten, M., Muisers, J. M., Verbakel, H. M., & de Wit, P.

(1993). Characterization of two putative pathogenicity genes of the fungal tomato pathogen

Cladosporium fulvum. *Molecular Plant-Microbe Interactions*, *6*, 210-215.

Van Esse, H. P., Bolton, M. D., Stergiopoulos, I., de Wit, P. J., & Thomma, B. P. (2007). The chitin-
binding *Cladosporium fulvum* effector protein Avr4 is a virulence factor. *Molecular Plant-*
Microbe Interactions, *20*(9), 1092-1101.

Van Kempen, M., Kim, S. S., Tumescheit, C., Mirdita, M., Lee, J., Gilchrist, C. L., Söding, J., &
Steinegger, M. (2024). Fast and accurate protein structure search with Foldseek. *Nature*
Biotechnology, *42*(2), 243-246.

Videira, S., Groenewald, J., Nakashima, C., Braun, U., Barreto, R. W., de Wit, P. J., & Crous, P. (2017).
Mycosphaerellaceae–chaos or clarity? *Studies in Mycology*, *87*, 257-421.

Wang, A., Meng, F., Xu, X., Wang, Y., & Li, J. (2007). Development of molecular markers linked to
Cladosporium fulvum resistant gene *Cf-6* in tomato by RAPD and SSR methods. *HortScience*,
42(1), 11-15.

Wang, T., Black, L., Hsieh, W., & Hanson, P. (1995). Inheritance of black leaf mold resistance in tomato.
Euphytica, *86*, 111-115.

Witek, K., Jupe, F., Witek, A. I., Baker, D., Clark, M. D., & Jones, J. D. (2016). Accelerated cloning of a
potato late blight–resistance gene using RenSeq and SMRT sequencing. *Nature*
Biotechnology, *34*(6), 656-660.

Wubben, J. P., Eijkelboom, C. A., & de Wit, P. J. (1993). Accumulation of pathogenesis-related proteins
in the epidermis of tomato leaves infected by *Cladosporium fulvum*. *Netherlands Journal of*
Plant Pathology, *99*, 231-239.

Zaccaron, A. Z., Chen, L.-H., Samaras, A., & Stergiopoulos, I. (2022). A chromosome-scale genome
assembly of the tomato pathogen *Cladosporium fulvum* reveals a compartmentalized
genome architecture and the presence of a dispensable chromosome. *Microbial Genomics*,
8(4).

Zaccaron, A. Z., & Stergiopoulos, I. (2024). Analysis of five near-complete genome assemblies of the
tomato pathogen *Cladosporium fulvum* uncovers additional accessory chromosomes and
structural variations induced by transposable elements effecting the loss of avirulence genes.
BMC Biology, *22*(1), 25.

902