Murine peritoneal macrophages undergo female-specific remodeling with aging

Ryan J. Lu[1,2], Shiqi Chen[3], Minhoo Kim[1], Nirmal K. Sampathkumar[1,4], Evelyn H. Lee[1], Amy

Christensen[1], Eric E. Wang[1,5], Isabella Y. Lau[1], Shivay Parihar[1,6], Charan K. Ravikumar[1,5], Jane

Jung[1], Shelby Brown[1], Alan Xu[1,7], Julio L. Alvarenga[1], Kristen Mehalko[1], Changhan D. Lee[1,8],

Helen S. Goodridge[9,10], Bérénice A. Benayoun[1,3,5,8,*]

[1] Leonard Davis School of Gerontology, University of Southern California, Los Angeles, CA

90089, USA.

[2] Graduate Program in the Biology of Aging, University of Southern California, Los Angeles,

CA 90089, USA.

[3] Cancer Biology Department, USC Keck School of Medicine, Los Angeles, CA 90089, USA.

[4] Present address: Samsara Therapeutics, Headington, Oxford, OX3 8SB, UK.

[5] Molecular and Computational Biology Department, USC Dornsife College of Letters, Arts and

Sciences, Los Angeles, CA 90089, USA.

[6] Graduate Program in Geroscience, University of Southern California, Los Angeles, CA 90089,

USA.

[7] Thomas Lord Department of Computer Science, USC Viterbi School of Engineering, Los

Angeles, CA 90089, USA.

[8] USC Norris Comprehensive Cancer Center, Epigenetics and Gene Regulation, Los Angeles,

CA 90089, USA.

[9] Board of Governors Regenerative Medicine Institute, Cedars-Sinai Medical Center, 8700

Beverly Boulevard, Los Angeles, CA 90048, USA

[10] Research Division of Immunology in the Department of Biomedical Sciences, Cedars-Sinai

Medical Center, 8700 Beverly Boulevard, Los Angeles, CA 90048, USA

* correspondence: berenice.benayoun@usc.edu

Abstract

Aging is a complex process characterized by a progressive decline in physiological functions driven by both biological and environmental factors, with notable differences between sexes. Immune function is strongly influenced by biological sex, affecting both innate and adaptive immune responses, including macrophage behavior. In this study, we investigated the effects of age and sex on the immune cell composition within the peritoneal cavity niche and identified macrophages as the most affected cell type. Macrophages, as central components of the innate immune system, play critical roles in maintaining tissue homeostasis and responding to infections. Here, we find that aging induces sex-specific remodeling of peritoneal macrophage transcriptomic and epigenomic landscapes. Consistently, peritoneal macrophages undergo sex-specific functional remodeling with aging (*i.e.* female-specific phagocytic decline and metabolic rewiring). Modulation of gonadal hormone signaling showed that changes in circulating estrogen levels likely contribute to aspects of female-specific macrophage age-related changes. Importantly, multi-omic analysis identified candidate transcription factors whose sex-specific age-regulated expression may drive aspects of sex-specific 'omic' remodeling with aging. Specifically, Irf2 downregulation in female macrophages recapitulates distinct transcriptomic and metabolic aspects of macrophage female aging phenotypes. These findings suggest that female-specific age-related functional remodeling arises through hormone-dependent and -independent mechanisms in peritoneal macrophages.

Introduction

Aging is a complex process characterized by a progressive decline in physiological

functions, with notable differences between sexes (Shi et al., 2024). This decline is driven by a

combination of biological factors, and extrinsic factors (Lopez-Otin et al., 2023). Life

expectancy of females are on average longer than males (Ostan et al., 2016). However, despite

increased longevity, females tend to experience higher levels of comorbidity and disability

compared to males (*i.e.* the mortality-morbidity paradox) (2014; Gordon et al., 2017; Kulminski

et al., 2008). This trend is reflected in healthspan studies, which consistently show that females

have higher frailty scores than males at any given age (Gordon et al., 2017). Additionally,

females face an elevated risk of developing age-related chronic conditions (*e.g.* coronary heart

disease, osteoarthritis, Type 2 diabetes, and neurodegenerative disorders) (Appt and Ethun,

2010). Chronic sterile inflammation, a mechanism dubbed "inflamm-aging", is a hallmark of

aging and is associated to more general immune system dysfunction or "immunosenescence"

(Franceschi and Campisi, 2014; Santoro et al., 2021). Despite well-documented sex-differences

in aging and disease susceptibility, underlying mechanisms driving sex-based aging disparities

remain poorly understood.

Immune function is heavily shaped by biological sex, influencing both innate and

adaptive immune cells (Dunn et al., 2024; Klein and Flanagan, 2016), including macrophages

(McGill et al., 2024). Females generally exhibit a more robust immune response compared to

males, such as quicker pathogen clearance, higher antibody response, higher cytokine secretion,

and reduction of mortality-related infections (Dunn et al., 2024; Flanagan et al., 2017; Klein and

Flanagan, 2016; Perrotta et al., 2020; Thompson et al., 2004). However, females are also more

susceptible to autoimmune diseases (Fairweather et al., 2024). During aging, hematopoietic stem

cells (HSC) skew production toward myeloid cells (*e.g.* monocytes, neutrophils) at the expense

of lymphoid cells (*e.g.* B- and T-cells), a shift that is more pronounced in males than females,

thereby impacting leukemogenesis and hematopoiesis (Pang et al., 2011; Zhang et al., 2025).

Aged males exhibit greater myeloid bias and inverted $CD4^+/CD8^+$ T-cell ratios (Lee et al., 2014),

along with functional impairments such as a sharp decline in T-cell proliferation and cytokine

secretion compared to females (Hirokawa et al., 2013; Strindhall et al., 2013). Importantly,

lifelong sex differences in immune function could stem from differential 'omic' regulation across

immune cell types (Barman et al., 2022; Gal-Oz et al., 2019; Huang et al., 2021; Lu et al., 2021;

Marquez et al., 2020; Qu et al., 2015). Despite the potential significance of such sex differences
in immune aging, how biological sex shapes the impact of age on the 'omic' and functional
landscape of innate immune cells remain largely unknown.

Macrophages are key components of the innate immune system, functioning as both
regulatory and effector cells in maintaining tissue homeostasis and responding to infections.
Tissue-resident macrophages (*e.g.* large peritoneal macrophages), which originate primarily from
embryonic precursors, exhibit specialized, organ-specific functions shaped by the local
microenvironment. In contrast, bone marrow-derived macrophages differentiate from circulating
monocytes and typically adopt more pro-inflammatory phenotypes during immune responses
(Ginhoux and Guilliams, 2016; Lavin et al., 2014). Macrophages have a repertoire of functions:
phagocytosis of pathogens and other foreign material, clearance of dead, dying, and senescent
cells, chemotaxis and migration through tissues and cytokine production (Ross et al., 2021).
During aging, macrophages exhibit altered phagocytosis, metabolic changes, and cytokine
secretion, leading to impaired clearance of apoptotic cells and poor inflammation control (Barrett
et al., 2015; Dube et al., 2022; Fix et al., 2021; Kim et al., 2017; Li et al., 2017; Linehan et al.,
2014; Lu et al., 2022; Peradinovic et al., 2023; Wong et al., 2017). In addition to aging
differences, sex-differences in macrophages have been observed in young adult mice (Klein and
Flanagan, 2016), where young females exhibit greater phagocytic activity and more efficient
antigen presentation (Mondal and Rai, 1999; Scotland et al., 2011). Interestingly, microglia, the
brain's resident macrophages, undergo a female-specific shift in glycolytic function with age that
is dependent on C3a signaling (Kang et al., 2024). While some pioneering studies have revealed
sex-differences in aging macrophages (Baker et al., 2025), the molecular mechanisms driving
sex-dimorphic aging phenotypes in macrophages have not been elucidated.

In this study, we employed an unbiased multi-omics approach to profile the transcriptome
and epigenome of primary peritoneal macrophages in female *vs*. male mice with aging. Our
analyses revealed that macrophages are regulated in a sex-dimorphic manner with aging,
potentially influencing numerous macrophage functions. Indeed, single cell transcriptomic
profiling of the peritoneal cavity revealed that macrophages are the most impacted cell type in
this niche as a function of aging and sex. Consistently, we identified divergent regulation of the
macrophage 'omic' landscape using RNA-seq, ATAC-seq, and H3K4me3 CUT&RUN profiling.
These analyses revealed that aging significantly alters immune function, with distinct patterns

observed across multiple immune parameters, some of which are modulated by sex. For instance,
our analyses showed that aging macrophages follow sex-dimorphic trajectories, with key
phenotypes related to 'omic' pathway changes—namely phagocytosis and metabolic
preference—exhibiting female-specific age-related remodeling. Our experiments suggest that
female-specific age-related remodeling of peritoneal macrophages likely occurs through both
gonadal hormone-dependent and independent mechanisms.
116
117

Results

Macrophages are the immune cell type most impacted by aging in the peritoneal cavity niche

The peritoneal cavity hosts a rich and accessible immune niche, housing B-cells, T-cells macrophages, neutrophils, and other cell types (Liu et al., 2021). This immune niche is easily accessible through post-mortem peritoneal lavage (Okabe, 2021; Saris et al., 2025), making it tractable for studies without the need for mechanical tissue dissociation, which can compromise both cell viability and phenotypes (Aliaghaei and Haun, 2022). We leveraged this niche to better understand how biological sex may impact the aging immune system using mice as a model system. Specifically, we obtained peritoneal lavage cells from young adult (4-5 months) and old (20-21 months) female and male C57BL/6JNia and C57BL/6J mice (**Figure 1A**). Isolated cells were then used for unbiased profiling by single-cell RNA sequencing (scRNA-seq) using the 10xGenomics v2 and v3.1 3' platforms (**Figure 1A**) in 3 independent sets of aging mice (one v2 dataset on C57BL/6JNia mice [NIA 10x v2], one v3.1 dataset on C57BL/6JNia mice [NIA 10x v3], and one v3.1 dataset on C57BL/6J mice [JAX]; see **Methods**).

After stringent quality control and cell annotation, we identified expected cell types within this niche in our rich dataset: B-cells, macrophages, monocytes, natural killer (NK) cells, and T-cells (**Figure 1B**; 18,805 cells, 10,758 cells, and 14,595 cells for the NIA 10x v2, NIA 10x v3 and JAX datasets, respectively). To determine whether any of these cell types were especially impacted by age and/or sex, we leveraged Augur, an algorithm that helps prioritize cell types based on their transcriptional response to biological conditions using a machine-learning approach(Skinnider et al., 2021) (**Figure 1C**; **Supplementary Figure S1A-D**). Using area under the curve (AUC) scores from Augur, we found that the transcriptome of macrophages was the most impacted among peritoneal cell types in the combined dataset (0.632; **Supplementary Figure S1E**), as well as each individual dataset ($\geq$ 0.567; **Supplementary Figure S1F-H**)**.**

Another important aspect that can be inferred from scRNA-seq corresponds to relative cell type abundances in each sample, especially in the absence of potential dissociation artifacts. Thus, we asked if there were any age- or sex-related changes in the relative abundances of peritoneal cell types, using both sample-specific determination and the scProportionTest algorithm (Miller et al., 2021) (**Figure 1D; Supplementary Figure S1I-L**). Interestingly, both methods revealed a significant increase in the relative proportion of B-cells and a significant

decrease in the relative proportion of macrophages in the peritoneal cavity with aging, regardless

of sex (**Figure 1D; Supplementary Figure S1I-L**), a finding consistent with previous studies

(Gautier et al., 2014; Mogilenko et al., 2022). In addition, we also observed an increase in the

relative proportion of the T-cell population with aging specifically in males (**Figure 1D;**

Supplementary Figure S1I-L). To confirm these results, we then used flow cytometry staining

of peritoneal cells to validate whether these changes in relative cell proportions also occurred *in*

vivo, using cell surface markers $CD19^+$ (B-cells), $F4/80^+$ (macrophages), and $CD3^+$ (T-cells)

(**Figure 1A, E**; **Supplementary Figure S1M**). Our data confirmed a relative increase in the

proportion of B-cells and a relative decrease in the proportion of macrophages in the peritoneal

cavity with aging, regardless of sex (**Figure 1E**). However, changes in relative T-cell abundance

were not validated with in flow cytometry, potentially due to high animal-to-animal variability

and a smaller effect size (**Figure 1E**). Leveraging this data, we then investigated whether these

changes in relative abundances were accompanied by changes in absolute immune cell type

quantities within the aging peritoneal cavity. Interestingly, we observed that regardless of sex,

there were increases of the total number of all 3 immune cell types within the peritoneal cavity

with aging (**Figure 1F**).

Tissue-resident peritoneal macrophages can self-renew after acute inflammation (Davies

et al., 2011; Louwe et al., 2021), and we posited that loss of self-renewal properties may

contribute to decreased macrophage proportion with age. To evaluate macrophage proliferative

capacity with aging, we leveraged our scRNA-seq data to predict the cell cycle stage of

macrophages across our samples, using the Seurat 'CellCycleScoring' module (Kowalczyk et al.,

2015). Intriguingly, the proportion of macrophages predicted to be in a resting non-cycling state

(*i.e*. in G1 phase) was significantly reduced with aging irrespective of sex, whereas the

proportion of macrophages predicted to be actively proliferating (*i.e*. in S, G2 or M phases)

tended to increase (**Supplementary Figure S1N**). Our data suggests that despite their reduced

proportion, macrophages are predicted to be more proliferative in the aged peritoneal cavity.

Thus, differences in macrophage abundance with aging are unlikely to stem from self-renewal

defects. However, changes in the peritoneal cavity niche with aging are likely to lead to

functional remodeling of these cells.

Together, our analyses suggest that macrophages are most significantly impacted by
aging and sex in the peritoneal cavity. Thus, we decided to focus on the peritoneal macrophage
compartment to understand potential sex-dimorphic molecular and functional effects with aging.
181

Peritoneal macrophages show sex-dimorphic 'omic' phenotypes with aging

To better understand the impact of aging and sex on aging peritoneal macrophages, we
leveraged a bulk multi-omic profiling approach using MACS-purified macrophages from young
adults (4-5 months) and old (20-21 months) female and male C57BL/6JNia mice. We opted to
use a bulk approach due to its increased sensitivity and robustness, especially in the context of
purified cells, compared to single cell level profiling. We chose to profile both transcriptional
and epigenomic features, specifically (i) transcriptome profiling by bulk sequencing of
ribosomal-RNA depleted RNA (RNA-seq), (ii) chromatin accessibility landscape through the
Assay for Transposon-Accessible Chromatin followed by sequencing (ATAC-seq) (Buenrostro
et al., 2013; Corces et al., 2017), and (iii) promoter activity/priming using CUT&RUN profiling
for the H3K4me3 histone post-translational modification(Skene and Henikoff, 2017) (**Figure
2A**). These 'omic' assays were each performed on an independent cohort of C57BL/6JNia
animals (n = 5 animals per group; **Supplementary Table S1A**). After accounting for premature
death before the experimental day (**Supplementary Table S1A**) and retaining only samples with
> 90% macrophage purity (**Supplementary Figure S2A, B**), final effective sample number
range from 3-5 per group and 'omic' layer.
We first leveraged Multi-Dimensional Scaling [MDS] to evaluate the main axes of
separation of our samples across the 3 'omic' layers (**Figure 2B-D**). Interestingly, MDS revealed
clear separation of samples by biological sex of the mice, regardless of age (**Figure 2B-D**). Thus,
MDS analysis supports the existence of strong sex-dimorphism of the molecular aging landscape
of primary peritoneal macrophages. Interestingly, we also observed a secondary separation of
samples by age of the mice, albeit weaker than that by sex (**Figure 2B-D**). Thus, we investigated
whether, in addition to baseline differences, aging may impact the 'omic' landscapes of
peritoneal macrophage in a sex-dimorphic manner. For this purpose, we compared DESeq2 $\log_2$
fold changes in gene expression (**Figure 2E, Supplementary Figure S2C**), chromatin
accessibility (**Figure 2F, Supplementary Figure S2D**) and H3K4me3 signal (**Figure 2G,
Supplementary Figure S2E**) for every detected gene/peak in aging female *vs*. male peritoneal

macrophages. We reasoned that a high rank correlation value would correspond to largely
colinear effects of aging on 'omic' landscapes regardless of sex and that divergent aging
trajectories would result in weaker rank correlation values. Using this method, we observed only
a relatively weak correlation between 'omic' aging trajectories of female and male peritoneal
macrophages (Genome-wide Spearman Rho = 0.232-0.329; **Figure 2E-G, Supplementary**
Figure S2C-E). To further investigate these divergent omic remodeling patterns with aging, we
next performed analyzed 'omic' changes with aging separately in females vs. males for each of
the 'omic' layers. Specifically, we labelled genes with FDR < 5% in one sex but FDR > 10% in
the other sex as significant in one sex only (sex-divergent regulation) and genes with FDR < 5%
in both sexes as consistently regulated (sex-independent regulation; **Supplementary Table S2A-**
C). Importantly, although a core set of genes and peaks significantly changed similarly in both
sexes with aging (respectively 61, 420 and 244; **Supplementary Figure S2C-E**), we identified
many more genes and/or peaks with significant age-related changes in only one sex (respectively
284/234, 1044/2825 and 131/1699 [male-specific/female-specific]; **Figure 2E-M**). Together,
these analyses are consistent with the notion of broadly divergent aging 'omic' trajectories in
peritoneal macrophages as a function of biological sex.
Sex differences in aging may result from (i) downstream effects of sex hormone signaling
(*e.g.* lifelong exposure to an estrogen- or androgen-rich milieu, or (ii) differential impact of sex
chromosome karyotypes (*e.g.* XX vs. XY), including through X chromosome inactivation [XCI]
escape(Baker et al., 2025). Genes and/or peaks with sex-dimorphic aging regulation were
distributed across the genome, indicating that autosomal gene dysregulation is a major feature of
sex-dimorphism in the aging 'omic' landscapes of macrophages (**Supplementary Figure S2F-**
H). To note, gene set enrichment analysis [GSEA] revealed a slight, but significant, trend for
downregulation of X-linked genes in females (FDR = 1.4 x10^{-2}; **Supplementary Figure S2I**),
consistent with a potential role of X chromosome dysregulation in aging female macrophages
(Ferraye et al., 2025). Consistently, we observed that one of the top female-specific regulated
genes with aging in our bulk RNA-seq dataset was *Xist* (FDR = 5.4x10^{-5}; **Supplementary**
Figure S2J), which encodes a non-coding RNA critical to XCI initiation and maintenance(Loda
and Heard, 2019; Zylicz et al., 2019). A trend for increased *Xist* expression in peritoneal
macrophages was also observed in our 10xGenomics v3 single cell RNA-seq datasets
(**Supplementary Figure S2K**). Interestingly, upregulation of *Xist* has been observed in an

scRNA-seq study of female brain aging (Hajdarovic et al., 2022), and there is evidence for

degraded XCI and increased escape in aged female brains across cell types (Gadek et al., 2025;

Hajdarovic et al., 2022). To determine whether a similar mechanism was at play in the context of

aging peritoneal macrophages, we compiled a list of previously identified XCI escapee genes in

mice from various cellular/tissue contexts (Berletch et al., 2015; Li et al., 2012; Sierra et al.,

2023; Wu et al., 2014; Yang et al., 2010) and used a GSEA approach to detect any potential XCI

deficiency in aged female macrophages. Our analysis showed that known XCI escapees were

significantly downregulated in aged female macrophages (FDR = 3.6×10^{-3}; **Supplementary**

Figure S2I), consistent with *Xist* upregulation but inconsistent with observations in the aging

brain (Gadek et al., 2025; Hajdarovic et al., 2022).

Together, our data reveals strong sex differences in the 'omic' landscapes of primary

peritoneal macrophages throughout life and suggest that aging may differentially impact the

function of these cells in female *vs.* male mice.

Functional enrichment analysis reveals sex-dimorphic regulation of key immune function

pathways with aging

We next asked whether sex-specific age-related changes in peritoneal macrophage gene

expression regulation, at the transcriptomic and epigenomic levels, were expected to impact

downstream functions using protein-protein interaction network construction and functional

enrichment analyses (**Figure 3, Supplementary Figure S3; Supplementary Table S3**).

First, predicted networks for female and male age-related changes were constructed with

NetworkAnalyst 3.0 (Zhou et al., 2019) and protein-protein interaction [PPI] information from

IMEx/InnateDB data (Breuer et al., 2013), a knowledgebase specifically geared for analyses of

innate immune networks (**Figure 3A-B**). Our analysis revealed that genes regulated with aging

in female vs. male macrophages are part of clear, interconnected PPI networks (**Figure 3A, B**).

Interestingly, although some predicted nodes were identified in both sexes (Irf7, Il1r1), there was

a majority of sex-specific nodes (*e.g.* Irf2 and Mef2c in females, Irf1 and Rela in males; **Figure**

3A, B). The tight gene interconnectivity within the female and male predicted PPI networks

supports the idea that macrophage transcriptional changes with aging are coordinated within each

sex, likely leading to differential aging related functional remodeling.

Next, we performed GSEA analysis using high quality gene sets from Gene Ontology
(GO; all domains; ENSEMBL v108) and Reactome (MSigDB v2022.1) to evaluate differentially
regulated processes in aging peritoneal macrophages at the levels of transcription, chromatin
accessibility and H3K4me3 signal, both in a sex-independent (**Supplementary Figure S3A-F;**
Supplementary Table S3A-F) and sex-dimorphic manner (**Figure 3C-E, Supplementary**
Figure S3G-I; Supplementary Table S3A-F). Importantly, top gene sets that were regulated in
a sex-independent manner were consistent with expected immune activation with aging (*e.g.*
Gene ontology terms "innate immune response" or "immune system process", Reactome gene
sets "complement pathway" and "neutrophil degranulation"; **Supplementary Figure S3A-C;**
Supplementary Table S3A-F). These observations are consistent with widespread observations
of age-related immune remodeling or "inflamm-aging"(Lee et al., 2022), regardless of sex. In
addition, we also observed strong sex-dimorphic regulation of GO and Reactome gene sets in
aging peritoneal macrophages (**Figure 3C-E, Supplementary Figure S3G-I; Supplementary**
Table S3A-F). Most notably, top significant sex-dimorphic aging gene sets revealed several
phagocytosis-related gene sets (*e.g.* GO "phagocytosis, recognition", "endosomes", Reactome
"cross-presentation of soluble exogenous antigens endosomes"), metabolism-related gene sets
(*e.g.* GO "proton transmembrane transport", "mitochondrial respiratory chain complex IV",
Reactome "respiratory electron transport ATP synthesis by chemiosmotic coupling and heat
production by uncoupling proteins") and inflammation-related gene sets (*e.g.* GO "positive
regulation of cytokine production involved in inflammatory response", "arachidonic acid
epoxygenase activity", Reactome "interleukin 2 family signaling", "defensins"; **Figure 3C-E,**
Supplementary Figure S3G-I; Supplementary Table S3A-F).

Since these types of terms reflect potential changes in key pathways relating to
macrophage function(Russell et al., 2019), we decided to complement our analysis using curated
gene sets related to these functions to independently validate their sex-specific regulation in the
transcriptome of aging peritoneal macrophages via GSEA (**Figure 3F-H, Supplementary Table**
S4). Specifically, we (i) curated a list of phagocytosis regulator genes from two phagocytosis
CRISPR/Cas9 screens (Haney et al., 2018; Pluvinage et al., 2019), (ii) utilized curated lists of
genes involved in oxidative phosphorylation and glycolysis from the MSigDB Hallmark gene
sets(Liberzon et al., 2011), and (iii) identified genes involved in M1-like and M2-like
polarization based on a publicly available macrophage RNA-seq dataset (Das et al., 2018).

Consistent with our unbiased GO and Reactome GSEA analyses, we observed female-specific downregulation of phagocytosis regulators from CRISPR/Cas9 screens (**Figure 3F; Supplementary Table S2G**) and a male-specific upregulation of curated Hallmarks metabolism related pathways (**Figure 3G; Supplementary Table S2H**). In contrast, both female and male macrophages showed significant upregulation of M1-enriched genes, whereas only male macrophages showed significant upregulation of M2-enriched genes as well (**Figure 3H; Supplementary Table S2I**).

Together, our functional enrichment analyses suggest that several processes key to macrophage function may be regulated in a sex-specific manner: (i) phagocytosis capacity, which is required for pathogen and/or dead cell clearance (Hirayama et al., 2017; Martin et al., 2014), (ii) metabolic preference, which reflects macrophage inflammatory status — with greater reliance on glycolysis being associated to more pro-inflammatory macrophages (Russell et al., 2019) and (iii) macrophage polarization trends, whereby distinct pro- and anti-inflammatory programs are activated (Murray, 2017).

Peritoneal macrophages undergo a female-specific age-related functional decline

Based on our 'omic' analyses (**Figure 3**), we hypothesized that peritoneal macrophages undergo sex-specific functional remodeling in phagocytosis capacity, metabolic preferences and polarization state with age. Thus, we selected *ex-vivo* assays to evaluate these phenotypes using primary peritoneal macrophages from young and old, female and male mice (**Figure 4A**).

To evaluate potential sex-differences in macrophages phagocytic abilities, we used Alexa Fluor 488-Zymosan bioparticles as cargo to evaluate phagocytosis capacity as a function of age and sex. Specifically, we evaluated the "phagocytosis index", defined as the proportion of macrophages internalizing the labelled cargo in a set amount of time, using 2 complementary approaches based on (i) microscopy, after 1 day of cell culture (Lu et al., 2020) and (ii) flow cytometry, on freshly isolated peritoneal macrophages (**Supplementary Figure S4A**; see **Methods**). In line with computational predictions, we observed a significant reduction in phagocytosis capacity specifically in aged female peritoneal macrophages, as demonstrated by both our microscopy (p-value = 2.0×10^{-2}; **Figure 4B, C**) and flow cytometry (p-value = 4.7×10^{-2}; **Figure 4D**) protocols. Importantly, we observed a consistent trend of decreased phagocytosis in female bone marrow-derived macrophages (BMDMs), a distinct population of primary

macrophages (p-value = 0.07; **Supplementary Figure S4B**). This suggests that female-specific phagocytic remodeling in aging macrophages may not be niche-specific.

Within the peritoneal cavity, the peritoneal macrophage population has been described to be made of both large peritoneal macrophages (LPM) expressing high levels of F4/80 and small peritoneal macrophages (SPM) expressing lower levels of F4/80 (Ghosn et al., 2010). LPMs are the tissue-resident macrophages of the peritoneal cavity, are long-lived cells derived from the fetal yolk sac, and constitute the overwhelming majority of peritoneal macrophages in naïve mice (Cassado Ados et al., 2015; Goswami et al., 2021; Jayakumar et al., 2022). In contrast, SPMs are derived from bone marrow progenitors through differentiation of blood monocytes constantly throughout life, are recruited to the peritoneal cavity in response to inflammatory signals, and represent a minority of peritoneal macrophages in healthy naïve mice (Jayakumar et al., 2022; Louwe et al., 2021). Thus, to rule out the possibility that female-specific changes in phagocytosis with aging could be a trivial byproduct of shifts in LPM/SPM proportions with aging (potentially in a sex-specific manner), we analyzed surface levels of macrophage marker F4/80 in our MACS-purified peritoneal macrophages using flow cytometry quantification (**Supplementary Figure S4C-E**). Intriguingly, although there were no aging differences in LPM/SPM abundance for macrophages purified from C57BL/6JNia mice, shifts were observed in aged female C57BL/6J mice (**Supplementary Figure S4D, E**). Despite provider/substrain differences in LPM/SPM proportion with aging, female-specific phagocytosis decreases were observed consistently, ruling out the potential impact of macrophage subtype changes to explain female-specific age-related remodeling in phagocytic capacity (**Supplementary Figure S4D, E**).

In addition, we also examined whether differential expression of pattern recognition receptors involved in Zymosan recognition may explain decreased phagocytosis in aging females specifically. Zymosan is recognized by both Tlr2 and Dectin-1 (encoded by *Clec7a*)(Dillon et al., 2006). There was no evidence of differential regulation of Zymosan receptors *Tlr2* and *Clec7a* in our RNA-seq datasets (**Supplementary Figure S4F**). Using flow cytometry quantification, we observed no change in Tlr2 protein expression with aging or sex (**Supplementary Figure S4G, H**). Dectin-1 abundance showed a significant reduction during aging in female macrophages, although Dectin-1 levels in aged females matched those of males regardless of age (**Supplementary Figure S4G**), despite comparable phagocytosis in young animals in our microscopy protocol (**Figure 4C**).

To identify potential mechanistic mediators of the specific phagocytosis phenotype in aged female macrophages, we overlapped our list of female-specific aging downregulated genes (RNA-seq) with the list of phagocytosis regulators from CRISPR/Cas9 phagocytosis screens (Haney et al., 2018; Pluvinage et al., 2019) (**Figure 4E, Supplementary Table S4A-B**). This revealed 9 candidate mediator genes of female-specific phagocytosis remodeling phenotypes (6 at FDR < 5%: *B4galt1*, *Dpy19l3*, *Man1a2*, *Rps6ka3*, *Tbl1xr1*, and *Tm9sf3;* 3 at FDR < 10%: *Gna13*, *Hmgcs1*, *Magt1*; **Figure 4E**). To validate whether these 9 candidate regulators could indeed impact Zymosan phagocytosis in murine macrophages, we performed a mini lentiviral short hairpin RNA (shRNA)-mediated knockdown screen in the RAW264.7 murine macrophage cell line, using 2 independent hairpins per candidate gene (**Figure 4F**). Lentivirus-transduced cells were selected using puromycin resistance, and successful knockdown of target genes was confirmed using RT-qPCR (**Supplementary Figure S4I**). Interestingly, only knockdown of *Rps6ka3* and *Magt1* resulted in significantly decreased Zymosan phagocytosis capability (**Figure 4G**), consistent with their potential mechanistic role in the female-specific age-related decreased phagocytosis.

Next, we evaluated the metabolic preference of primary macrophages with respect to aging and sex. Macrophage function in both homeostasis and disease conditions relies on metabolic reprogramming(Viola et al., 2019), whereby increased reliance on glycolysis is characteristic of pro-inflammatory states and increased preference of mitochondrial respiration is more associated to tissue remodeling states(Russell et al., 2019). We took advantage of the unique capabilities of the Agilent Seahorse platform to metabolically phenotype primary macrophages as a function of sex and age(Van den Bossche et al., 2015). Since this platform requires cell numbers that are difficult to achieve with naïve peritoneal macrophages, we leverage BMDMs to measure oxygen consumption rate [OCR] and extracellular acidification rate [ECAR], using a protocol designed to evaluate glycolytic and mitochondrial activity(Van den Bossche et al., 2015) (**Figure 4H-J, Supplementary Figure S4J, K**). Contrary to expectations from our 'omic' analysis, no differences in glycolytic activity or mitochondrial respiration were observed in aging male BMDMs (**Figure 4I, J, Supplementary Figure S4J, K**). However, we observed a significant increase in glycolytic activity in aging female BMDMs (**Figure 4I, J**). To independently validate this phenotype in peritoneal macrophages, we employed a luminescence-based assay to measure a central byproduct of glycolysis (lactate), an

assay with much lower cell number requirements. Consistent with our BMDM findings, we also
observed a female-specific glycolytic shift in aging peritoneal macrophages (**Figure 4K**).

Finally, we set out to validate our 'omic' findings of sex-dimorphic activation of
polarization pathways (Figure 3H) using flow cytometry quantification of cell surface markers
known to be associated to M1-like (CD80, CD86) and M2-like (CD163, CD206) macrophage
phenotypes (Chavez-Galan et al., 2015; Furgiuele et al., 2022; Yao et al., 2019). We found that
cell surface markers CD80 and CD86 increased with age in both sexes (**Supplementary Figure
S4L**), indicating a general shift toward a more inflammatory phenotype with aging regardless of
sex, consistent with previous studies in the field. In contrast, markers CD163 and CD206 showed
inconsistent age-related changes; notably, CD206—but not CD163—significantly increased with
age specifically in female macrophages (**Supplementary Figure S4M**).

Our functional validation analyses revealed robust female-specific age-related
remodeling related to primary macrophage phagocytosis and metabolic preference.

Impact of ovarian hormone signaling on macrophage phenotypes with female-specific age-
related remodeling

Our functional analyses revealed two phenotypes with strong female-specific age-related
remodeling: (i) phagocytosis and (ii) glycolytic capacity. Notably, female mammals are known
to undergo drastic age-related decreases in circulating ovarian hormones (*e.g.* estradiol,
progesterone, etc.)(Kim et al., 2016). In addition, previous work suggested that ovarian
hormones can impact macrophage function, including phagocytosis (Loiola et al., 2019; Sun et
al., 2021) and metabolic reprogramming (Ribas et al., 2011). Hence, we asked whether the
known age-related decrease in circulating ovarian hormones in female mice (Kim et al., 2016)
may underlie female-specific age-related remodeling of peritoneal macrophages. Consistent with
the ability to respond to variations in circulating sex hormone levels, our RNA-seq dataset
showed that primary peritoneal macrophages expressed sex hormone receptors, with *Esr1*
(encoding estrogen receptor α) being the most robustly expressed across samples
(**Supplementary Figure S5A**). Therefore, we reasoned that decreased circulating 17β-estradiol
[E2] levels (the major form of estrogen in females) with aging may drive aspects of female-
specific macrophage remodeling.

We used a multi-pronged approach, combining analyses of macrophages (i) from young

females across the estrus cycle (thus experiencing endogenous changes in circulating E2) and (ii)

from young female ovariectomized [OVX] vs. sham control mice. First, we characterized cohorts

of 4 month old C57BL/6NTac mice and identified animals actively cycling using vaginal

cytology (Cora et al., 2015) to obtain peritoneal macrophages exposed to different endogenous

levels of E2 (Zysow et al., 1997) (proestrus/estrus: high E2, metestrus/diestrus: low E2;

Supplementary Figure S5C; Supplementary Table S1B, C). Interestingly, peritoneal

macrophages isolated from females in low E2 phases of the estrus cycle showed significantly

lower phagocytosis rates compared to those from females in high E2 phases (p-value = 1.3×10^{-2};

Supplementary Figure S5D). However, metabolic profiling of BMDMs from naturally cycling

mice did not reveal E2-dependent differences in metabolic preference (**Supplementary Figure**

S5E, F). To more directly test the causality of ovarian hormone depletion, we next used an OVX

vs. sham surgery approach on young C57BL/6J females at 3 months of age and allowed the mice

to recover for one month before euthanasia at 4 months (**Figure 5A; Supplementary Table**

S1D). We then evaluated 4 phenotypes: (i) peritoneal lavage composition, (ii) peritoneal

macrophage transcriptome profiling, (iii) peritoneal macrophage phagocytosis, and (iv)

peritoneal macrophage glycolytic activity (**Figure 5A**). Relative abundances of peritoneal

immune cell types were unaffected in OVX animals (**Supplementary Figure 5G**), suggesting

that B-cells and macrophages proportions are not regulated by ovarian hormone levels. Total

numbers of peritoneal immune cells were also not affected (**Supplementary Figure S5H**).

Importantly, the transcriptional landscape of peritoneal macrophages was strongly remodeled in

OVX vs. sham animals based on MDS analysis and differential gene expression analysis (**Figure**

5B; samples with > 90% purity; **Supplementary Figure S5I; Supplementary Table S5A**). To

evaluate whether OVX impacts macrophage transcriptional landscapes similarly to female aging,

we next performed GSEA of genes differentially regulated upon OVX in macrophages (FDR <

5%) in the aging female peritoneal macrophage transcriptome (**Figure 5C**). Importantly, genes

upregulated/downregulated upon OVX were significantly upregulated/downregulated in aging

female macrophages (**Figure 5C**), supporting that OVX mimics transcriptional effects of aging

in female macrophages. Next, we evaluated the functional impact of OVX on peritoneal

macrophages. Our flow cytometry-based phagocytosis assay showed that peritoneal

macrophages from OVX mice had significantly decreased phagocytosis capacity when compared

to sham mice (p-value = 2.2×10^{-3}; **Figure 5D**), similar to the effect of aging. However, lactate secretion (and by proxy glycolysis) was unaffected by OVX (**Figure 5E**). Together, our findings support the notion that lower levels of E2 are sufficient to recapitulate aspects of female macrophage aging (*i.e.* transcriptome, phagocytosis), but not others (i.e. metabolic preference).

Next, we reasoned that because decreased E2 conditions phenocopied aspects of female macrophage aging, supplementation of E2 in post-estropausal female mice may rescue aspects of aging. For this purpose, we treated 3 months and 19 months old female C57BL/6J mice with E2 (440ng/mL) or vehicle (*i.e.* ethanol 0.1%) in their drinking water (Gordon et al., 1986; Kohama et al., 1989) for one month before euthanasia (**Figure 5F; Supplementary Table S1E**). This delivery method provides a minimally invasive framework for hormone supplementation and was confirmed to work in our hands, as uterine dry weight was significantly increased in aged E2-treated animals (p-value = 1.1×10^{-2} in Wilcoxon rank-sum test; **Supplementary Table S1E**). We then phenotyped the peritoneal cavity and purified peritoneal macrophages as above. While we observed expected age-related changes in B-cells and macrophages relative abundances, E2 supplementation did not restore these abundances to youthful levels (**Supplementary Figure S5J**). To note, although E2 was previously reported to upregulate cell cycle genes in young macrophages(Pepe et al., 2017), there were no changes in the total number of peritoneal immune cells isolated from old females supplemented with E2 (**Supplementary Figure S5K**), consistent with post-estropausal peritoneal immune cells being refractory to E2 signaling. Interestingly, the transcriptional landscape of peritoneal macrophages was strongly remodeled in response to aging and E2 supplementation, as assessed by MDS and differential expression analysis (**Figure 5G**; samples with > 90% purity; **Supplementary Figure S5L; Supplementary Table S5B-C**). Interestingly, the transcriptomes of macrophages from E2-supplemented aged mice clustered in between those of young vehicle-treated mice and old vehicle-treated mice (**Figure 5G**), suggesting a potentially "younger" transcriptomic signature. Indeed, GSEA using genes significantly regulated in response to E2 supplementation in old mice (FDR < 5%) and female macrophage aging revealed that genes upregulated in response to E2 were downregulated during aging, whereas genes downregulated in response to E2 were significantly during aging (**Figure 5H**), consistent with partial transcriptional rejuvenation. We next evaluated whether E2 supplementation would also functionally reprogram female macrophage toward a more youthful state. Using our flow-cytometry-based assay, we observed an expected age-related decrease in

phagocytosis with vehicle treated animals (p-value = 1.9×10^{-2}), but no significant rescue upon
E2 supplementation (p-value = 0.56; **Figure 5I**). Glycolytic output, as measured by lactate
production, was increased with aging as expected (p-value = 1.9×10^{-2}) and showed a non-
significant trend for decrease upon E2 treatment (p-value = 8.5×10^{-2}; **Figure 5J**). Therefore, we
found that short-term *in vivo* E2 supplementation of post-estropausal mice, while showing signs
of rescuing markers of transcriptional aging in female macrophages, did not significantly impact
functional aging phenotypes. Since post-menopausal hormonal therapy is known to have a
limited window for efficacy in humans (Rossouw et al., 2002) and C57BL/6 mice undergo
estropause between 12-15 months of age (Frick, 2009), it is possible that short-term exposure at
19 months of age cannot fully rescue the effects of age-related E2 loss.

Our results suggest that age-related decline in circulating E2 in females may explain
aspects of female-specific macrophage aging. Since *Esr1* was the highest expressed receptor for
E2 in our peritoneal macrophage bulk RNA-seq data (**Supplementary Figure S5A**), we next
evaluated potential changes in expression or activity aging peritoneal macrophages. To note,
there was a trend for increased *Esr1* expression with aging in female (but not male) macrophages
in bulk RNA-seq (p-value = 7.1×10^{-2}; **Supplementary Figure S5B**), consistent with the known
negative feedback loop of Esr1 regulation (Lung et al., 2020), but expression levels were too low
for robust detection in our scRNA-seq datasets (as expected for genes encoding transcription
factors (Saliba et al., 2014)). Thus, to determine whether Esr1 may mediate the impact of
reduced E2 levels with aging in peritoneal macrophages, we leveraged an existing *Esr1* full body
knockout model (Hewitt et al., 2010) (**Figure 5K**). Specifically, we isolated and characterized
cells from 3-4 months old *Esr1* knockout and wild-type littermate female mice, similarly as
above (**Figure 5K; Supplementary Table S1F**). By using surface marker analysis through flow
cytometry, we found that *Esr1* knockout mice showed a significantly increased B-cell proportion
(p-value = 4.1×10^{-2}) and a trend for decreased proportion of macrophages within the peritoneal
cavity (p-value = 0.056; **Supplementary Figure S5M**), consistent with a prematurely aged
immune phenotype. There was a change in the numbers of peritoneal macrophages, but not other
cell types (p-value = 4.9×10^{-3}; **Supplementary Figure S5N**), which could stem from the impact
of E2 on cell cycle genes in young animals (Pepe et al., 2017). Next, we evaluated the
transcriptomic impact of constitutive *Esr1* knockout. Although *Esr1* knockout led to substantial
transcriptional remodeling according to MDS and differential expression analysis (**Figure 5L;**

samples with > 90% purity; **Supplementary Figure S5O; Supplementary Table S5D**), GSEA analysis of *Esr1* knockout regulated genes surprisingly showed a pattern consistent with transcriptional rejuvenation of macrophages (**Figure 5M**). However, consistent with cycling and OVX functional results, *Esr1* knockout peritoneal macrophages exhibited significantly decreased ability to perform phagocytosis, as determined by flow-cytometry (p-value = 3.7×10^{-3}; **Figure 5N**), but showed no difference in glycolytic output (p-value = 0.23; **Figure 5O**).

Together, our results support the notion that sustained E2 exposure, at least in part through Esr1 signaling, is required to maintain youthful transcriptomic signatures and high levels of phagocytosis, but not low glycolytic dependence, in female peritoneal macrophages throughout life.

Identifying non hormonal mechanisms driving aspects of female-specific macrophage aging

Since decreased E2 signaling could only account for limited aspects of female-specific age-related remodeling in peritoneal macrophages (*i.e.* phagocytosis, transcriptional landscapes), we next sought to identify additional potential molecular mediators contributing to these female-specific macrophage remodeling with aging. To do so, we leveraged the comprehensive 'omic' datasets we generated for this study (**Figure 1, 2**) as tools to pinpointing candidate regulators (**Figure 6A; Supplementary Table S6B**). Among the most promising candidates are genes encoding transcription factors [TFs], given their broad regulatory influence over the expression of hundreds to thousands of other genes(Lee and Young, 2013).

We leveraged our bulk RNA-seq dataset to perform an *in-silico* screen to identify candidate TF-encoding genes that significantly changed in expression with aging in female, but not male, peritoneal macrophages (**Figure 6A; Supplementary Table S6B**). We then examined evidence for disrupted TF signaling specifically in aging female macrophages using: (i) TF binding motif analysis in ATAC-seq regions remodeled specifically in aging female macrophages and (ii) TF binding motif analysis in H3K4me3 CUT&RUN regions remodeled specifically in aging female macrophages (**Figure 6A; Supplementary Table S6B**). Mouse TF-encoding genes were obtained from the AnimalTFDB3.0 database(Hu et al., 2019) and overlapped to genes with significant age-regulation in females only, revealing 20 candidate TF-encoding genes (**Figure 6A, B; Supplementary Table S6B**). To note, 14 of these TFs were upregulated with age, and 6 of them were downregulated with age (**Figure 6A, B,**

Supplementary Figure S6A; Supplementary Table S6B). We then performed TF motif analysis in the genomic regions with ATAC-seq and H3K4me3 CUT&RUN signal remodeled specifically in female macrophages with aging using HOMER(Heinz et al., 2010). Significantly enriched motifs in peaks with increased or reduced signals were then filtered based on our 20 candidate TFs for ATAC-seq female-specific aging differential peaks (**Figure 6C**) and for H3K4me3 CUT&RUN female-specific aging differential peaks (**Figure 6D**). This filtering led to the identification of 4 TF candidates with age-related <u>decreased</u> expression (*i.e. Irf2, Mef2C, Meis1, Tal1*), and 3 TF candidates with age-related <u>increased</u> expression (*i.e. Nr2f2, Sox7, Tead1*) whose motifs where enriched in chromatin regions specifically remodeled with aging in female macrophages. For simplicity, we next decided to focus on the set of TFs specifically downregulated with aging in female macrophages (*i.e. Irf2, Mef2C, Meis1, Tal1*). Interestingly, these candidate TFs were also nodes of the predicted interconnected PPI network identified in aging female macrophages (**Figure 3A**). Finally, we leveraged our scRNA-seq datasets to predict differential TF activity across sex and ages in macrophage single cell transcriptomes using the SCENIC algorithm(Aibar et al., 2017). This analysis revealed a predicted significant decrease in Irf2 activity and significant increase in Mef2c activity in aging female macrophages (**Supplementary Figure S6B-C; Supplementary Table S6A**), providing another layer of evidence that age-related disruption of these TFs may impact female macrophages. Additionally, we noted that one of our previously evaluated candidate phagocytosis regulators (*Tbl1xr1*; **Figure 4F-G**) functions as a transcriptional co-regulator(Mastrototaro et al., 2021). Intriguingly, *Tbl1xr1* was also identified in our SCENIC analysis as showing a significant age-related decrease in predicted activity (**Supplementary Figure S6B, C; Supplementary Table S6A**). In our analyses hereafter, we considered a set of 4 TFs and 1 TF co-regulator (*i.e. Irf2, Mef2c, Meis1, Tal1, Tbl1xr1*; hereafter terms as "TFs") as candidate TF regulators of female macrophage aging (**Figure 6A; Supplementary Table S6B**). The impact of candidate TFs on macrophage phenotypes (*i.e.* transcriptome, phagocytic capacity and metabolic preference) was evaluated using the RAW264.7 macrophage cell line through a mini lentiviral shRNA knockdown screen, using 2 independent hairpins per candidate TF gene (**Figure 6F**; see **Methods**).

        After confirming successful knockdown of TF candidates by RT-qPCR (**Supplementary Figure S6D**), we analyzed the impact of decreased TF expression on RAW264.7 macrophage

transcriptomes by mRNA-seq (**Supplementary Figure S6E-G**). We found that TF knockdown,

regardless of specific hairpin, led to clear remodeling of macrophage transcriptional landscapes

as assessed by MDS (**Supplementary Figure S6F**). Using a stringent filter of combined FDR

from both hairpins $< 10^{-4}$, we were able to identify sets of robust transcriptional targets of

candidate TFs (**Supplementary Figure S6G; Supplementary Table S6C-G**). We next used this

set of robust targets as gene sets for GSEA, in order to evaluate whether the transcriptional

impact of decreased TF expression was consistent with that of female macrophage aging (**Figure**

6G). Interestingly, knockdown of *Irf2* and *Tal1* showed gene regulation patterns colinear with

female macrophage aging, with upregulated genes being significantly upregulated in aging

female macrophages (and downregulated genes being downregulated, respectively). Importantly,

Irf2 targets showed substantially stronger enrichment than those of *Tal1* (**Figure 6G**). (**Figure**

6G). In the case of *Mef2c* and *Meis1* knockdown, downregulated genes showed strong

downregulation in aging female macrophages, whereas upregulated genes showed weak

downregulation opposite of aging (**Figure 6G**). However, knockdown of *Tbl1xr1* revealed a

transcriptional pattern inconsistent with aging, with targets genes skewing in the opposite

direction of aging female macrophages (**Figure 6G**). Intriguingly, when investigating whether

candidate TFs may belong to a regulatory network of their own, we found that *Mef2c* knockdown

led to significantly reduced *Irf2* expression (**Supplementary Figure S6H, I**), suggesting that

concomitant downregulation of *Irf2* and *Mef2c* may act synergistically to shape the aging female

macrophage transcriptome.

We next evaluated whether decreased expression of candidate TFs may recapitulate

female-specific age-related macrophage functional remodeling in phagocytosis and metabolic

preferences (**Figure 6H-J**). First, with the notable exception of *Meis1* knockdown, we observed

no other significant recapitulation of age-related phagocytosis decreases in response to TF

knockdown on macrophage phagocytic capacity (**Figure 6H**). Thus, age-related downregulation

of *Meis1* may contribute to female-specific aging macrophage phagocytosis decreases.

Interestingly, *Rps6ka3*, our top hit from our mini-shRNA knockdown screen for candidate

regulators of phagocytosis was significantly downregulated in RAW264.7 macrophages after

Meis1 knockdown (FDR = 0.004; **Supplementary Figure S6J; Supplementary Table S6E**).

This finding supports the notion that *Meis1* downregulation in aged female macrophages occurs

upstream of *Rps6ka3* downregulation to promote reduced phagocytosis. In contrast to our

phagocytosis assay, using both the Agilent Seahorse and Promega Lactate-Glo assays, we

observed a more widespread impact of TF candidate knockdown on glycolytic reliance (**Figure**

6I, J). Based on these 2 assays, we observed that that knockdown of *Irf2*, *Tal1*, and *Tb11xr1* led

to significantly increased macrophage glycolytic activity, consistent to the female macrophage

aging phenotype (**Figure 6I, J**).

Together, our mini-shRNA screen showed that downregulation of *Irf2* led to the strongest

(i) transcriptional and (ii) metabolic remodeling consistent with female macrophage aging.

Consequently, its age-related downregulation represents the best candidate mechanism to explain

observations of female-specific glycolytic shift in aging macrophages, which could not be

explained by age-related estrogen deprivation (**Figure 5, Supplementary Figure S5**).

Age-related downregulation of Irf2 expression may promote female macrophage metabolic

reprogramming through derepression of *Hk3* expression

Since Irf2 was our top candidate from our mini-shRNA TF candidate screen, we next set

out to independently validate its regulation in aging primary peritoneal macrophages (**Figure**

7A-D, **Supplementary Figure S7A-C**). First, RT-qPCR analysis of *Irf2* RNA expression in

peritoneal macrophages in independent aging mouse cohorts confirmed significant

transcriptional downregulation of *Irf2* with aging, albeit in both sexes (**Figure 7B**). Interestingly,

Irf2 transcriptional levels, as assessed from our previously generated RNA-seq datasets, was

unaffected by ovarian hormone signaling, with no significant change in expression in response to

OVX (DESeq2 FDR = 0.69; **Figure S7A upper panel**), late-life E2 supplementation (DESeq2

FDR = 1; **Figure S7A middle panel**) or *Esr1* knock-out (DESeq2 FDR = 0.91; **Figure S7A**

lower panel), consistent with *Irf2* downregulation occurring independently of estrogen signaling

and/or ovarian aging. Next, we asked whether protein levels of Irf2 were impacted in aging

peritoneal macrophages using intracellular immunolabeling and quantification by flow cytometry

(**Figure 7C-D**; **Supplementary Figure S7B-C**). Consistent with RNA-seq observations, we

observed a significant female-specific age-related decrease in Irf2 protein labeling in peritoneal

macrophages (**Figure 7D**). Importantly, this labeling was specific, as RAW264.7 and J774.1

macrophage cell lines expressing shRNA against *Irf2* showed significantly decreased labeling

(**Supplementary Figure S7D, E**). Thus, our results confirm that Irf2 is downregulated in aging

female peritoneal macrophages at both the transcriptional and protein levels.

To better understand the impact of Irf2 in young female macrophages (and thus the

impact of its decreased expression with aging), we next evaluated the Irf2 chromatin binding

landscape in primary female macrophages from young female C57BL/6JNia mice (**Figure 7E**).

To note, since TF ChIP-seq profiling requires 10 million cells per replicate, we leveraged

BMDMs (rather than rare naïve peritoneal macrophages) to map the Irf2 macrophage regulome.

We identified 339 robust peaks across 2 biological replicates (**Supplementary Table S7A**).

ChIP-seq peaks were significantly enriched for DNA binding motifs recognized by interferon

regulatory factors (IRFs), consistent with high specificity of the immunoprecipitation

(**Supplementary Figure S7F; Supplementary Table S7B**). Consistent with its role as a TF,

Irf2-bound regions were mostly found near promoters ($\leq$ 3kb from annotated transcription start

sites [TSSs]) and in distal intergenic regions, likely enhancers; **Supplementary Figure S7G**). To

gain further insights about the regulatory network of Irf2 in primary macrophages, we leveraged

the Genomic Regions Enrichment of Annotations Tool (GREAT) tool(McLean et al., 2010),

which revealed enrichments of GO terms related to immune response and metabolism (*e.g*,

GO:0006955 "immune response", GO:0003950 "NAD+ ADP-ribosyltransferase activity";

Figure 7F; Supplementary Table S7C, D). Similar terms were found to be enriched among Irf2

targets when using the alternative gene-centric ChIPseeker enrichment tool(Wang et al., 2022)

(*e.g*, GO:0045088 "regulation of innate immune response", GO:0046320 "regulation of fatty

acid oxidation"; **Supplementary Table S7E, F**).

We next investigated how Irf2 *bona fide* target genes were regulated with aging at the

transcriptional and epigenomic levels using our previous datasets (**Figure 2**). Importantly, Irf2

ChIP-seq <u>direct</u> targets (regardless of distance to TSSs) were significantly upregulated in aging

female macrophages based on GSEA statistics (**Supplementary Figure S7H**), consistent with its

primary role as a transcriptional repressor(Nguyen et al., 1995; Yamamoto et al., 1994). Notably,

even with a less powered overlap-based approach, the majority of direct Irf2 targets that

overlapped with genes significantly regulated in aging female macrophages (FDR $<$ 5%) were

found to be upregulated (**Supplementary Figure S7H**). Interestingly, 3 of the upregulated genes

in this overlap, *Tm6sf1*, *Usp18*, and *Ifit3*, were previously reported to be linked to cell

metabolism(Chen et al., 2024; Huang et al., 2024; Lai et al., 2024). Consistent with increased

activation of Irf2 targets in aging female peritoneal macrophages, Irf2-bound chromatin regions

showed increased chromatin accessibility by ATAC-seq (**Supplementary Figure S7I-J,** left

panel) and increased H3K4me3 CUT&RUN signal (**Supplementary Figure S7I-J**, right panel).

Together, these results support the notion that decreased Irf2 expression with aging may lead to

reactivation of its target gene network, potentially leading to immune and metabolic remodeling.

Next, we manually inspected the list of Irf2 *bona fide* direct targets to gain further

insights into the impact of Irf2 downregulation in aging female macrophages (**Supplementary**

Table S7A). Intriguingly, we noted that there was strong binding of Irf2 in the promoter of the

Hk3 gene, which encodes an isoform of hexokinase, the enzyme that catalyzes the initial rate-

limiting step of glycolysis (**Figure 7H**). There was a non-significant trend for increased *Hk3*

expression with aging in our peritoneal macrophage RNA-seq dataset, likely due to an outlier

value in the young female group (**Supplementary Figure S7K**). However, our RT-qPCR

analysis of peritoneal macrophages from additional 3 independent cohorts of animals showed a

significant increase in *Hk3* RNA levels (p-value = 1.6×10^{-2}; **Figure 7I**). Thus, age-related

downregulation of Irf2 in female peritoneal macrophages is likely to lead to *Hk3* upregulation,

and thus increased glycolytic activity.

To complement mechanistic insights downstream of Irf2 downregulation from our screen

in RAW264.7 macrophages (**Figure 6**), we performed shRNA knockdown in the independent

J774A.1 macrophage cell line (**Figure 7J, Supplementary Figure S7L-N**). There was good

agreement in the transcriptional remodeling following *Irf2* knockdown across both macrophage

cell lines as measured by mRNAseq (**Supplementary Figure S7L, M; Supplementary Table**

S6G). Consistent our screen results in RAW264.7 macrophages, *Irf2* knockdown did not impact

phagocytic capacity in J774A.1 macrophages (p-value = 0.51; **Supplementary Figure S7N**).

However, as expected from the status of *Hk3* as a strong direct target of Irf2 and observations in

female aging macrophages, *Irf2* knockdown led to robust upregulation of *Hk3* expression in both

macrophage cell lines according to RNA-seq (RAW264.7 p-value = 2.6×10^{-37}; J774A.1 p-value

= 3.9×10^{-9}; **Figure 7K**). Consistent with *Hk3* expression acting as a limiting factor for glycolytic

potential of macrophages, upregulation of *Hk3* associated significantly with increased glycolytic

activity according to measurements using both the Agilent Seahorse and Promega Lactate-Glo

platforms in both the RAW264.7 (p-value = 1.2×10^{-3}, 7.3×10^{-3} respectively) and J774A.1

macrophage cell lines (p-value = 1.7×10^{-3}, 2.6×10^{-2} respectively; **Figure 6I-J, 7L-M**).

Together, our data suggests that decreased *Irf2* expression with aging in female peritoneal
macrophages leads to an upregulation of *Hk3*, which then leads to an aberrant age-related
increase in glycolytic output (**Figure 7N**).
704
705

Discussion

A multi-omic resource to study murine tissue-resident macrophages as a function of age and biological sex.

To understand how sex may influence aging of the immune system, specifically macrophages, we generated a multi-omic resource focusing on primary resident peritoneal macrophages from young and old, female and male C57BL/6J[Nia] mice, with high quality transcriptomic (single-cell and bulk RNA-seq) and epigenomic (chromatin accessibility by ATAC-seq and H3K4me3 genomic distribution by CUT&RUN) maps. To our knowledge, our dataset is the largest multi-omic dataset for the study of primary macrophages during aging, and we believe it will be a rich resource to the scientific community in understanding the influence of biological sex on age-related immune remodeling. Analysis of these datasets revealed that, not only is there a baseline difference in genomic regulation of primary macrophages at young age, but also that aging differentially impacts 'omic' profiles in female *vs*. male animals. Specifically, although there were shared features of molecular aging in female and male macrophages, we observed much larger sex-dimorphic age-related changes in gene expression and chromatin organization (**Figure 2; Supplementary Figure S2**).

Importantly, genes and genomic regions showing age-related sex-dimorphic regulation were enriched for processes key to macrophage function, including phagocytosis regulation, metabolic wiring and inflammatory phenotypes (**Figure 3; Supplementary Figure S3**). Although aging phenotypes of macrophages across various niches have been reported (*e.g*. visceral adipose macrophages, alveolar macrophages, Kupffer cells, *etc*.), previous studies largely focused on using animals of one sex, or mostly reported results combining sexes (reviewed in (Lu et al., 2022)). A notable exception includes a series of recent studies on aging murine microglia, the brain resident macrophage, which also reported large sex-differences in molecular landscapes with aging(Kang et al., 2024; Ocanas et al., 2023).

Peritoneal macrophages show female-specific age-related remodeling in phagocytosis and metabolic wiring.

To determine whether sex-differential regulation of molecular landscapes with aging yielded functional consequences, we assessed phagocytic capacity, metabolic preference, and polarization surface markers in isolated peritoneal macrophages (**Figure 4; Supplementary**

Figure S4). Intriguingly, this revealed two significant female-specific age-related remodeling in
primary peritoneal macrophages: (i) a decrease in phagocytic capacity, and (ii) a metabolic shift
to increased glycolytic output. Different populations of tissue-resident macrophages have
previously been observed to show age-related decreases in phagocytosis across a range of cargo
types, but with no insights on the potential influence of sex on these phenotypes (Lu et al., 2022).
Our data suggests that age-related remodeling in macrophage functions is sex-dimorphic.

To understand the molecular mechanism driving female-specific regulation of
phagocytosis, we tested candidate regulator genes of phagocytosis from CRISPR/Cas9 screens
that were specifically downregulated in aged female macrophages through a mini shRNA
knockdown screen (**Figure 4G**). Our results identified two phagocytosis regulators (*Magt1* and
Rps6ka3) whose knockdown mimicked the age-related decrease in phagocytosis of female
macrophages, consistent with the notion that their downregulation with aging may explain
mechanistically our observed female-specific age-related decrease in macrophage phagocytic
capacity. Although its impact during aging has not been evaluated before, *Magt1* is a Magnesium
transporter located to the endoplasmic reticulum involved in protein glycosylation with critical
function in immune cells(Blommaert et al., 2019; Matsuda-Lennikov et al., 2019), which was
identified as required for proper phagocytosis of Zymosan in a whole genome CRISPR/Cas9
screen of human phagocytes(Haney et al., 2018). Interestingly, *Rps6ka3* encodes a member of
the S6 kinase family, whose activity is regulated by mTOR(He et al., 2019), which is a key
regulator of aging(He et al., 2025), autophagy(Panwar et al., 2023), phagosome
maturation(Krajcovic et al., 2013), and macrophage polarization(Byles et al., 2013). *Rps6ka3*
was also shown to play a direct role in innate immune responses(Kakugawa et al., 2009) and was
found to be required for phagocytosis of Zymosan in human phagocytes(Haney et al., 2018).
Thus, downregulation of *Magt1* and *Rps6ka3* in aging female peritoneal macrophages is likely to
mechanistically underlie the female-specific age-related phagocytosis defect that we observed.

Interestingly, despite 'omic'-based predictions of male specific metabolic rewiring, we
identified a metabolic shift specifically in aging female macrophages (both peritoneal
macrophages and BMDMs), leading to increased glycolytic output. Increased glycolytic output is
known to support inflammatory phenotypes and ROS-production in macrophages and is a
characteristic of disease-associated macrophages(Wculek et al., 2022). Furthermore, a recent
study of aging murine microglia also identified a female-specific increase in glycolytic output,

which was driven by increased C3a – C3aR signaling(Kang et al., 2024). Thus, our observations

are consistent with a potential shift to a more dysfunctional state in female macrophages with

aging.

Sustained estrogen signaling is necessary for youthful macrophage phagocytosis.

Since the age-related functional remodeling we uncovered were female specific, we reasoned

age-related decreases in circulating ovarian hormones may explain these phenotypes (**Figure 5;**

Supplementary Figure S5). Indeed, previous work found that E2 signaling can shape

macrophage phenotypes(Loiola et al., 2019; Sun et al., 2021). Importantly, decreased exposure to

youthful E2 levels by ovariectomy recapitulated aspects of female-specific macrophage aging,

including transcriptional remodeling and decreased phagocytosis, but had no impact on

glycolytic output of female macrophages (**Figure 5C-F**). Similar functional outcomes were

observed in macrophages from mice lacking estrogen receptor gene *Esr1*, the most abundantly

expressed estrogen receptor in peritoneal macrophages, suggesting the functional impact of

youthful E2 may be mediated through ERα (**Figure 5Q-R**). In contrast, late-life short-term E2 *in*

vivo supplementation partially rejuvenated the transcriptional landscape of peritoneal

macrophages (**Figure 5I-J**), although there was no significant rescue of the phagocytic or

metabolic remodeling (**Figure 5K-L**). Accumulating data has shown that delayed onset of

hormone therapy in post-menopausal women leads to decreased efficacy (or even increased risk

of adverse side effects)(Baker and Benayoun, 2023; Cho et al., 2023; Lobo, 2007), suggesting

that late-life supplementation, months after the onset of estropause (9-12 months of age in

C57BL/6 mice)(Finch, 2014), is likely too late to observe substantial rescue of female-specific

age-related macrophage functional remodeling. Thus, our results suggest that youthful E2 levels

are necessary (although not sufficient) to maintain (i) aspects of the youthful female macrophage

transcriptome, and (ii) youthful phagocytosis capacity in peritoneal macrophages. However,

youthful E2 exposures had no influence on glycolytic output, suggesting that the female-specific

metabolic shift of peritoneal macrophages occurs independent of circulating levels of sex

hormones. Future work should determine whether hormone treatment initiated earlier in life,

closer to estropause onset, will lead to at least partial functional rescue of aging female

macrophage phenotypes.

TF-encoding genes are involved in female-specific age-related macrophage phenotypes.

We also identified a set of putative TF encoding genes that were specifically downregulated in aging female macrophages and may shape female-specific aging phenotypic remodeling: *Irf2*, *Mef2c*, *Meis1*, *Tal1*, and *Tbl1xr1* (**Figure 6**; **Supplementary Figure S6**). Interestingly, with the exception of *Tbl1xr1*, knockdown of candidate TF genes recapitulated phenotypic aspects of aging female macrophages, with *Irf2* and *Tal1* knockdown most closely resembling the effect of aging for both upregulated and downregulated genes, while *Mef2c* and *Meis1* knockdown led to strong repression of aging-related downregulated genes (but not induction of upregulated genes; **Figure 6G**).

Among candidates, only knockdown of *Meis1* led to aged-like decreased phagocytic capacity, likely due to downstream repression of phagocytosis regulator *Rps6ka3*. To note, *Meis1* expression is unaffected in macrophages from OVX mice or in response to late-life short term E2 supplementation (**Supplementary Table S5A, C**), suggesting that its impacy on phagocytosis occurs independently of E2 signaling. In contrast, knockdown of *Irf2*, *Tal1*, and *Tbl1xr1* promoted increased glycolytic output, suggesting that their age-related expression downregulation may underlie the female-specific glycolytic shift we observed in female peritoneal macrophages. How age-related decreased expression of these TFs in female macrophages may synergize together (and with decreased E2 signaling) to shape female-specific aging phenotypes is an interesting avenue for future studies.

Female-specific age-related downregulation of Irf2 leads to increased *Hk3* expression and glycolytic output.

Since *Irf2* knockdown best mimicked the aging female macrophage transcriptome and led to the strongest increase in glycolytic output, we focused on elucidating how it may drive aspects of female-specific peritoneal macrophage phenotypes (**Figure 7**; **Supplementary Figure S7**). Interestingly, Irf2 shares a similar DNA binding domain as Irf1, but works as a competitive antagonist of Irf1-mediated gene expression(Wang et al., 2024). Consistent with decreased Irf2 activity driving aspects of aging female macrophage molecular landscapes, ChIP-seq analysis revealed that *bona fide* Irf2 target genes in primary macrophages were significantly upregulated and overlapped with remodeled chromatin regions in aging female macrophages. Most notably, ChIP-seq identified a strong region of Irf2 binding in the promoter of *Hk3*, the gene encoding

Hexokinase 3. Both female macrophage aging and *Irf2* knockdown led to significant
upregulation of *Hk3*, suggesting that decreased *Irf2* expression with aging is responsible for
increased expression of *Hk3*. Importantly, hexokinases catalyze the rate-limiting first step of
glycolysis (the phosphorylation of glucose on C6)(Zuo et al., 2021), thus gatekeeping cellular
glycolytic potential. Thus, our data suggests that the female-specific increase in glycolytic output
during peritoneal macrophage aging is driven by downregulation of *Irf2* and derepression of its
downstream target *Hk3*. Interestingly, this mechanism that we identified in peritoneal
macrophages is distinct from the one previously reported in microglia(Kang et al., 2024),
suggesting that metabolic rewiring of female macrophages is driven by niche-specific
mechanisms.

Together, our study provides the first systematic characterization of sex-differences in
immune aging in the context of macrophages and provides mechanistic insights into how sex-
specific aging phenotypes can be achieved. Intriguingly, our data suggests that both mechanisms
involving age-related changes in gonadal hormone levels and mechanisms independent of
gonadal hormones can drive sex-dimorphism in the aging trajectories of immune cells.
Ultimately, such mechanisms could be leveraged as potential targets to rejuvenate immune
responses in a sex-specific manner, the first and easy step towards personalized geroscience.
847

Material and Methods

Mouse husbandry

848
849
All animals were treated and housed in accordance with the Guide for Care and Use of
Laboratory Animals. All experimental procedures were approved by the University of Southern
California's Institutional Animal Care and Use Committee (IACUC) in accordance with
institutional and national guidelines. Experiments described in this manuscript were performed
on approved IACUC protocols 20770, 20804, 21004 and 21454. All animals were euthanized
between 8-11 am using a "snaking order" to minimize batch-processing confounding factors
related to circadian effects and post-mortem interval, by using by CO_2 asphyxiation followed by
cervical dislocation.

For intact aging analyses, male and female C57BL/6JNia mice were obtained from the
National Institute on Aging (NIA) colony at Charles Rivers, or male and female C57BL/6J mice
were obtained from the Jackson Laboratories (3 and 19 months, acclimated at USC for 4-6
weeks) **(Supplementary Table S1A).**

For estrus cycle tracking experiments male and female C57BL/6NTac were obtained
from Taconic Biosciences (3 months, acclimated at USC for 2 week;**Supplementary Table
S1B**). The cycle stage of adult female mice was characterized by vaginal lavage daily for 3
weeks until euthanasia on 3-4 months old mice. Mice that seen at least a full cycle or seen of the
different phases during the three weeks of vaginal smear were taken for euthanasia
(**Supplementary Table S1C**). To reduce animal waste, non-cycling mice were housed with a
mixture of clean and used male bedding to elicit additional cycle to stimulate cycling. Mice were
cycled for another two weeks and mice that were cycling were taken for euthanasia.

For ovariectomy (OVX) *vs*. sham surgery experiments, female C57BL/6J mice were
obtained from Jackson Laboratories (8 weeks old animals, acclimated at USC for 4 weeks)
(**Supplementary Table S1D**). C57BL6/J female mice gone through sham or OVX surgery at 12
weeks of age. All surgery procedures and post-operative care were conducted by University of
Southern California Department of Animal Resources veterinarians and veterinarian technicians.
Surgery entry point was from the back to preserve the peritoneal cavity as much as possible for
post-mortem peritoneal cell collection. Mice went through a recovery period of 4 to 5 weeks
before euthanasia.

For late-life short-term E2 supplementation experiments, female C57BL/6J mice were

obtained from Jackson Laboratories at age 10 and 80 weeks and acclimated at USC for 2 weeks

before treatment initiation (**Supplementary Table S1E**). For supplementation, 17β-estradiol

(E2) was used in the purest chemical grade that could be obtained (≥98% purity; Sigma-Aldrich

#E8875). We generated our E2 stock solution of 5 ng/mL in pure 100% ethanol (Sigma-Aldrich

#E7023). E2 stock solution and 100% ethanol were filtered sterilized on 0.22μm PES filters

(VWR #76479-024) and stored at -20°C for no longer than 1 month. Drinking water of

C57BL6/J mice was supplemented with either vehicle (0.1% EtOH) or 440 ng/mL of E2 for 4

weeks, corresponding to ~94μg/kg of body weight/day(Gordon et al., 1986; Kohama et al.,

1989). Treatment water was changed every 2-3 days (3 times a week) to maximize treatment

potency and limit potential E2 degradation. Water levels and mice were monitored daily. Water

supplementation of E2 in this manner was found to be a safe and effective alternative to daily

injections (Gordon et al., 1986; Kohama et al., 1989; Kohama et al., 1992). Mice were

euthanized ~4 weeks after treatment initiation for experimental phenotyping.

For B6N(Cg)-*Esr1$^{tm4.2Ksk}$*/J [*Esr1* KO](Hewitt et al., 2010) (JAX #026176) experiments,

matched female knockout [KO] and wild-type [WT] littermates were obtained from Jackson

Laboratories (**Supplementary Table S1F**). After receiving mice at USC, *Esr1* KO/WT mice

were acclimated and aged at USC until they reached 12-16 weeks before euthanasia.

Charles Rivers, Jackson Laboratories, and Taconic Biosciences have specific-pathogen-

free (SPF) facilities. All animals were acclimated at the SPF animal facility at USC for >2 weeks

before any processing or intervention to minimize the impact of travel stress. Some animals had

to be euthanized prior to experimental endpoints upon recommendation of USC veterinarians due

to welfare concerns and could thus not be included in the study.

Cell hashing, single-cell RNA-seq library preparation & sequencing of peritoneal cells

Single-cell libraries were prepared either using Chromium Single Cell 3' Library & Gel

Bead Kit v2 (10X Genomics #PN-120237) [NIA pooled cohorts 1 & 2], or the Chromium Next

GEM Single Cell 3□ GEM, Library & Gel Bead Kit v3.1 (10X Genomics #PN-1000121) [JAX

and NIA single animal cohorts with HTO labeling] according to manufacturer's instructions (10x

Genomics User Guide Chromium Next GEM Single-cell 3′ Reagent Kits v3.1 (CG000204, Rev

D)). For the two experiments using the v2 system, isolated peritoneal cells were pooled from

each biological group with equal cell number to ensure equal representation of all animals, based
on cell concentration estimates from a COUNTESS cell counter (Thermo Fisher Scientific #
AMQAF1000). This resulted in 4 libraries each (*i.e.* young female, young male, old female, old
male) for each of 2 independent cohorts. In this case, cell suspensions were loaded for a targeted
cell recovery of 3,000 cells per sample. For the two experiments using the v3.1 system, cell
hashing was performed on isolated peritoneal cells from each individual using Biolegend
TotalSeq^{TM}-A Antibodies according to the manufacturer's instructions (Biolegend #155801,
#155803, #155805, #155807, #155809, #155811, #155813, #155815, #155817, #155819).
Specifically, 0.5 million peritoneal cells were first incubated with mouse FcR-blocking Reagent
(Miltenyi Biotec #130-092-575) at 4°C for 10 minutes. Then, each blocked sample was
incubated with 1μg of specific TotalSeq Cell Hashing antibody (HTOs 1-10) at 4°C for 30
minutes. After incubation, stained cells were washed 3 times, and assessed for viability and yield
using the COUNTESS cell counter (JAX) or the DeNovix CellDrop (Denovix #CellDrop FL-
UNLTD) (NIA). Making sure no overlapping HTOs were included in a pool, samples were
pooled in equicellular mixes for 3 reactions (JAX) or 2 reactions (NIA), with targeted cell
recoveries of 14,000 and 30,000 cells, respectively. Samples were run on Chromium Single Cell
A Chips (10X Genomics #PN-120236) [v2 reactions] or Chromium Next GEM Chip G (10X
Genomics #PN-2000177) [v3.1 reactions] according to the manufacturer's instructions. For the
v3.1 reactions, HTO Additive primer v2 (Integrated DNA Technologies) was added to the cDNA
amplification reaction to amplify the HTO molecules. The amplified HTOs were recovered in the
cleanup step for the cDNA amplification, and HTO libraries were built using the 2x NEBNext
PCR Master Mix and Truseq indexing oligos (Integrated DNA Technologies). Partially amplified
and final single-cell RNA-seq libraries were quantified by a dsDNA High Sensitivity assay
(Invitrogen #Q32854) on a Qubit 3.0 Fluorometer (Thermo Fisher Scientific #Q33216). Quality
and size distribution of single-cell libraries were assessed on the 4200 TapeStation system
(Agilent Technologies #G2991A) with High Sensitivity D1000 DNA ScreenTapes (Agilent
Technologies #5067-5582) and D1000 reagent (Agilent Technologies #5067-5583). Libraries
were sequenced on an Illumina Novaseq 6000 generating 150 bp paired-end reads at Novogene
USA. Raw FASTQ reads have been deposited to the sequence read archive under accession
PRJNA1094925.

Computational processing of 10xGenomics sequencing libraries

Mapping, cell filtering and demultiplexing

For all datasets, raw reads were aligned to the mm10 mouse genome and processed using cellranger 6.0.2(Zheng et al., 2017) to generate gene-barcode matrices. For our hashed v3.1 datasets, HTO quantification was performed using CITE-seq-Count v.1.4.4 (Roelli et al., 2019) with parameters "-cbf 1 -cbl 16 -umif 17 -umil 26" to parse UMI/HTO sequences. Gene-barcode matrices and HTO counts were loaded in R 3.6.3(Team, 2022) for downstream processing using Seurat 3.2.2 (Hao et al., 2021; Satija et al., 2015). We used SoupX v1.6.1(Young and Behjati, 2020) to correct counts for any interference by ambient RNA in our libraries. Cells that were detected in the cellranger data (NIA v2 library) or in both cellranger and HTO data (NIA v3 and JAX v3 libraries) were filtered for downstream analysis. Dead and low-quality cells were filtered out by selecting for cells that contained between 300-6000 genes and < 25% mitochondrial reads. After removal of low-quality unwanted cells, UMI expression data was normalized for downstream processing using SCTransform(), regressing out the impact of number of genes and mitochondrial read load ("vars.to.regress = c("nFeature_RNA", "percent.mito")").

Cell cycle prediction

A list of mouse cell cycle genes was obtained from the Seurat Vignettes (https://www.dropbox.com/s/3dby3bjsaf5arrw/cell_cycle_vignette_files.zip?dl=1), derived from a mouse study of cell cycling (Kowalczyk et al., 2015). Cell cycle phase was predicted by these reference cell cycle genes using the function CellCycleSorting() to assign cell cycle scores and most likely phase to each cell. This data was used to determine whether there were age-related changes in proliferation patterns of peritoneal macrophages (**Supplementary Figure S1N**).

Singlet identification, doublet removal and cell demultiplexing

Dataset dimensionality was determined using (i) the dimension where the principal components only contribute 5% of standard deviation, (ii) cumulative PCs contribute 90% of the standard deviation, and (iii) the dimension where the percent change in variation between the consecutive PCs is less than 0.1%, using the smallest value. This was determined to be 16 (NIA v2), 21 (NIA v3) and 16 (JAX v3), respectively.

For all our datasets, potential doublets were annotated using Doubletfinder v2.0.3

(McGinnis et al., 2019) in each particular library, using the predicted doublet rate based on cell

capture provided by 10x Genomics. For the NIA v2 dataset, all cells annotated as singlets by

Doubletfinder were retained for downstream analyses.

In addition, for our NIA v3 and JAX v3 libraries, we took advantage of HTO libraries to

add a second layer of doublet identification and removal. Specifically, HTO counts were added

to the Seurat object as an independent assay and normalized using centered log-ratio (CLR)

transformation. Cells were demultiplexed according to their HTO barcode of origin using the

MULTIseqDemux() function into their component samples. For these datasets, cells were

considered to be singlets if and only if they were annotated as singlets by Doubletfinder and

MULTIseqDemux.

Dimensional reduction and cell clustering

Singlet UMI expression data was normalized and scaled for downstream analyses using

SCTransform(), regressing out the impact of number of genes, mitochondrial read load and

predicted cell cycle scores ("vars.to.regress = c("nFeature_RNA", "percent.mito", S.Score",

"G2M.Score)"). The first N principal components (PCs), where N is the estimated

dimensionality of the dataset (see above) were used to compute the UMAP transformation with

RunUMAP() and cluster cells using FindNeighbors() and FindClusters().

Cell type annotation using SingleR

To identify the most likely cell type annotation for each cell, singlets were annotated

using SingleR v.1.0.1(Aran et al., 2019), together with the reference ImmGen(Heng et al., 2008)

and the mouse RNAseq(Benayoun et al., 2019) expression datasets. To guarantee annotation

robustness, only cells annotated to the same cell type using both references were kept and

annotated with the consensus prediction for all downstream analyses.

Seurat object merging

After independent pre-processing for each dataset, the 3 cohort datasets were also merged

into a single Seurat object for analysis. Reciprocal PCA was used to integrate data from the 3

datasets and mitigate batch effects, using the top 3000 most variable genes and k = 10 anchors.

1001

Cell type prioritization using AUGUR and cell type proportion analysis

We used the R package AUGUR v1.0.3(Squair et al., 2021) to determine which

peritoneal cell types were most transcriptionally impacted by age and sex, in the combined and

separated objects (**Figure 1C, Supplementary Figure S1A-H**).

We calculated a per library cell type proportion using annotations and analyzed changes

in proportions for the most abundant cell types (B-cells, Macrophages and T-cells) across

biological groups (**Figure 1D**). In addition, as an orthogonal approach, we computed changes in

cell type proportions from the combined and dataset-specific Seurat objects using

'scProportionTest' v0.0.0.9000(Miller et al., 2021), which runs a permutation test on two

different single-cell groups for each cell type and returns the relative change in cell type

abundance between groups with a confidence interval (**Supplementary Figure S1I-L**).

1013

Peritoneal lavage immune cell proportion

Flow cytometry analysis was used to validate changes in proportions of immune cell

types within the peritoneal cavity from at least two independent cohorts of aged C57BL/6JNia

mice, aged C57BL/6J, OVX *vs*. sham C57BL/6J mice, E2-supplemented C57BL/6J mice, and

Esr1 KO mice. A panel of antibodies was used to evaluate differences in the proportion of B-

cells, macrophages, and T cells (see below). Specifically, among live cell singlets, B-cells were

defined as CD19$^+$, T-cells were defined as CD3$^+$, and macrophages were defined as F4/80$^+$.

Cells were isolated from the peritoneal cavity using a peritoneal lavage. Then, 200,000

peritoneal cells were washed once by adding 1mL of PBS and 2mM EDTA pH 7.2 (Miltenyi

Biotec #130-091-222) supplemented with 0.5% BSA (Miltenyi Biotec #130-091-376; MACS

resuspension buffer) in a 5mL polystyrene round bottom tube (Falcon #352054), then

centrifuged 300g for 10 minutes at 4°C. Cells were resuspended in MACS resuspension buffer in

the presence of FcR-blocking solution (concentration following manufacturer recommended

dilution) for 10 minutes at 4°C. Afterwards, cells were stained with antibodies for 10 minutes at

4°C protected from light, and excess antibody was washed away using MACS resuspension

buffer. Stained cells were then analyzed by flow cytometry on a MACSQuant Analyzer 10

(Miltenyi Biotec #130-096-343), and flow cytometry data was analyzed using FlowLogic V8.

Gating was determined using fluorescence-minus-one [FMO] controls for each color used in the

experiment to ensure that positive populations were solely associated with the antibody for that specific marker. Raw cytometry data, including FMO controls, were deposited for naïve aging, sham vs. OVX, E2 supplementation, and *Esr1* KO to Figshare under 10.6084/m9.figshare.28934213, 10.6084/m9.figshare.29169674, 10.6084/m9.figshare.29169737, and 10.6084/m9.figshare.29169707, respectively.

Peritoneal lavage immune cell yield

To assess the age-related impact to immune cell number within the peritoneal cavity, we took total peritoneal lavage cell counts from the DeNovix CellDrop and the identified the fraction of each immune cell types (B-cell, macrophage, T-cell; **Figure 1E**) to calculate original immune cell numbers in the peritoneal cavity. To calculate the number of immune cell types (B-cell, macrophage, and T-cell), the total peritoneal lavage cell counts were multiplied by the fraction of immune cell obtained from flow cytometry (**Figure 1F**).

Isolation of primary peritoneal macrophages

Isolation of primary peritoneal macrophages was performed on aged C57BL/6JNia, aged C57BL/6J, estrus cycle tracking C57BL/6NTac, OVX *vs*. sham C57BL/6J, E2-supplemented C57BL/6J, and *Esr1* KO mice following a protocol from Lu et al. 2020(Lu et al., 2020) with modifications to remove excess T-cells and B-cells in old animals. Briefly, peritoneal cells were obtained by using a 25G needle (BD #305125) to inject 10mL of filtered sterilized 3% bovine serum albumin (BSA) (VWR #0332-500G) in DPBS (Corning, 21-031-CV) into the peritoneal cavity of euthanized mice. Using a 20-gauge needle (BD #305175), the BSA solution was then harvested back from the peritoneum, now referred to as "peritoneal lavage". Peritoneal lavages were centrifuged 300g for 10 minutes at 4°C, and cells were resuspended in 1mL of MACS resuspension buffer. To remove any potential contaminating red blood cells, samples were treated with 10mL of 1x Red Blood Cell [RBC] lysis solution (Biolegend #420301) inverted to mix and incubated at 2 minutes at room temperature, then centrifuged 300g for 10minutes at 4°C. To obtain single cell suspension and removal of cell clumps from RBC lysis, macrophages were filtered on 30µm MACS SmartStrainers (Miltenyi Biotec #130-110-915).

Peritoneal macrophages were then isolated using Miltenyi Biotec peritoneal isolation kit (Miltenti Biotec #130-110-434), with modifications to the magnetic labeling step improve purity

of macrophage isolation regardless of age, treatment, and genotype. Indeed, while using the unmodified manufacturer's protocol, we found that aged mice exhibited poor macrophage purity, with large contamination of B-cells (likely due to B-cell population expansion with aging; **Figure 1D, E**). To improve peritoneal macrophage purity, we routinely added additional biotin labelled antibodies during the "macrophage (peritoneum) biotin-antibody cocktail" labeling step to help deplete potential excess B-cells, T-cells and monocytes from peritoneal lavages. Specifically, 5μL of biotinylated anti-CD19 (Miltenyi Biotec #130-119-757), anti-CD79b (Miltenyi Biotec #130-105-832), anti-CD3e (Miltenyi Biotec #130-093-179) and anti-Ly6c (Miltenyi Biotec #130-111-776) antibodies were added during incubation for 10 minutes at 4°C. After primary antibody incubation, all other steps in isolating peritoneal macrophages were performed according to the manufacturer's instructions. This allowed us to routinely achieve high purity for isolated peritoneal macrophages. Magnetic Activated Cell Sorting (MACS)-purified peritoneal macrophages were used for downstream experiments, including 'omic' and functional profiling.

Purity and identify of MACS-purified peritoneal macrophages

Potential phenotypic differences in MACS-purified peritoneal macrophage purity were assessed from independent cohorts of aged C57BL/6JNia, aged C57BL/6J, OVX vs. sham C57BL/6J, E2-supplemented C57BL/6J, and *Esr1* KO mice. Briefly, 100,000 MACS-purified peritoneal macrophages were used for flow cytometry labeling. After FcR-blocking, macrophages were stained using F4/80-APC (1:50; Miltenyi Biotec #130-116-547) and CD11b-Vioblue (1:50; Miltenyi Biotec #130-113-238) antibodies according to the manufacturer's instructions. Stained cells were analyzed by flow cytometry on the Miltenyi Biotec MACSQuant Analyzer 10. Flow cytometry results were processed using the FlowLogic V8 software. Pure macrophages were designated as double positive F4/80$^+$ CD11b$^+$ cells. From these, large peritoneal macrophages were defined as F4/80^{+high}, and small peritoneal macrophages were defined as F4/80^{+low}. The raw flow cytometry data for naïve aging, sham vs. OVX, E2 supplementation, and *Esr1* KO analyses has been deposited to Figshare under 10.6084/m9.figshare.28934198, 10.6084/m9.figshare.29169578, 10.6084/m9.figshare.29169581, and 10.6084/m9.figshare.29169587, respectively.

RNA purification from primary macrophages and macrophage cell lines

For RNA isolation of primary peritoneal macrophages, frozen cell pellets were resuspended in 500μL of TRIzol reagent (Thermo-Fisher #15596018), and total RNA was purified following the manufacturer's instructions. RNA quality was assessed using the Agilent Technologies 4200 TapeStation system using High Sensitivity RNA ScreenTapes (Agilent Technologies #5067-5579) together with RNA ScreenTape Sample Buffer (Agilent Technologies #5067-5577).

For RNA isolation of modified RAW264.7 and J774A.1 cells (see below), 60-80% confluency in a 6-well plate were lysed using 1mL of TRIzol reagent, and RNA was extracted using DirectZol columns (Zymo #R2052) following manufacturer's protocol.

RNA-seq library preparations for primary peritoneal macrophages and macrophage cell lines

For primary peritoneal macrophages, we constructed ribosomal RNA depleted strand-specific bulk RNA-sequencing libraries in house as previously described Lu et al. 2021 (Lu et al., 2021). Briefly, 450ng of total RNA was subjected to ribosomal-RNA depletion using the NEBNext rRNA Depletion Kit v2 (New England Biolabs #E705X), according to the manufacturer's protocol. Strand specific RNA-seq libraries were then constructed using the SMARTer Stranded RNA-seq Kit (Clontech #634841, #634863), according to the manufacturer's protocol. Libraries were quality controlled on the Agilent Technologies 4200 TapeStation system with Agilent Technologies D1000 ScreenTape and Agilent Technologies D1000 reagent, before multiplexing the libraries for sequencing. Paired-end 150bp reads were generated on the Illumina Novaseq X or Novaseq 6000 platforms at Novogene USA. Raw FASTQ sequencing reads were deposited to SRA under Bioprojects accession PRJNA1095555 and PRJNA1101936.

For macrophage cell line derived samples, total RNA samples were sent to the Novogene corporation for unstranded mRNA-sequencing. Briefly, the company performed mRNA selection using poly (A) capture, RNA fragmentation, cDNA reverse transcription and then unstranded RNA sequencing library preparation. RNA libraries were sequenced as paired-end 150bp reads on the Illumina Novaseq X or Novaseq 6000 platforms. Raw FASTQ sequencing reads were deposited to SRA under Bioprojects accession PRJNA1096052 and PRJNA1222371.

1125

RNA-seq analysis pipeline

1126

Paired end 150bp FASTQ reads were hard-trimmed to 100bp at their 3' end to improve

mapping rates and the first 9bp at the 5' end were trimmed to eliminate priming biases (Hansen

et al., 2010). This was done using Fastx Trimmer 0.0.13

(https://github.com/agordon/fastx_toolkit), or Trimgalore 0.6.6

(github.com/FelixKrueger/TrimGalore) after a system upgrade made it impossible to use Fastx

Trimmer. For Trimgalore, additional required stringency filters were added to eliminate any

bases with phred score < 15 (corresponding to $< 1\%$ of bases). Trimmed reads were mapped to

the mm10 genome reference using STAR 2.7.7a (Dobin et al., 2013). Read counts were assigned

to genes from the UCSC mm10 reference using featureCounts from subread 1.5.3 (Liao et al.,

2014), using either the stranded or unstranded option based on library construction (see above).

Read count tables were imported into R 3.6.3 (Team, 2022) for downstream processing. Based

on general RNA-seq processing guidelines, only genes with mapped reads in at least half of the

samples in each analysis were considered to be expressed and retained for analysis.

The 'DESeq2' R package v1.26.0 was used for normalization and differential gene

expression analysis of the RNA-seq data in R (Love et al., 2014) using appropriate analytical

covariates (*e.g.* sex, age, treatment, etc.). Genes with a false discovery rate $< 5\%$ in any analysis

were considered statistically significant. For analysis of sex-specific age-regulation of genes, we

considered a gene to be specifically regulated with age in females (males) if the DESeq2 FDR

was $< 5\%$ in females (males) and $> 10\%$ in males (females). DESeq2 results are reported in

Supplementary Table S2A.

1147

ATAC-seq library generation for primary peritoneal macrophages

1148

A second independent cohort of C57BL/6JNia mice was used to assess the impact of age

and sex on peritoneal macrophage chromatin accessibility (**Supplementary Table S1A**). We

constructed ATAC-seq libraries in house as described Lu et al. 2021 (Lu et al., 2021), using the

omni-ATAC protocol and starting from 50,000 MACS-purified peritoneal macrophages (Corces

et al., 2017). Libraries were quality controlled using the Agilent Technologies 4200 TapeStation

system with Agilent Technologies High Sensitivity D1000 DNA ScreenTape and reagent before

multiplexing the libraries for sequencing. Paired-end 150bp reads were generated on the Illumina

Novaseq 6000 platform at Novogene USA. Raw FASTQ sequencing reads were deposited to

SRA under Bioproject accession PRJNA1095555.

1158

ATAC-seq preprocessing pipeline

Paired-end ATAC-seq reads were adapter-trimmed using NGmerge 0.3 (Gaspar, 2018b),

a tool recommended for use in ATAC-seq preprocessing which clips overhanging reads in a

sequence-independent fashion. Trimmed paired-end were mapped to the mm10 genome

reference using bowtie2 v2.5.1 (Langmead and Salzberg, 2012). After alignment, PCR duplicates

were removed using the 'rmdup' function of samtools 0.1.19 (Li et al., 2009). Reads from all

samples were pooled define "meta-peaks" for robust differential accessibility analysis using

samtools 'merge'. We used MACS2 v2.2.7.1 to identify ATAC-seq accessible regions (Gaspar,

2018a). A normalized read count matrix was then obtained using our mapped read bam files over

MACS2 meta-peaks using R package 'Diffbind' 2.14 (Ross-Innes et al., 2012), which was used

for downstream analyses. 'DESeq2' 1.26.0 was used for normalization and differential gene

expression analysis of ATAC-seq data in R (Love et al., 2014) using appropriate analytical

covariates (*e.g*. sex, age). Peaks with a false discovery rate < 5% in any analysis were considered

statistically significant. For analysis of sex-specific age-regulation of chromatin region

accessibility, we considered a chromatin region to be specifically remodeled with age in females

(males) if the DESeq2 FDR was < 5% in females (males) and > 10% in males (females).

DESeq2 results are reported in **Supplementary Table S2B**.

1176

H4K4me3 CUT&RUN library generation for primary peritoneal macrophages

A third independent cohort of C57BL/6JNia mice was used to assess the impact of age

and sex on peritoneal macrophage H3K4me3 landscapes (**Supplementary Table S1A**). We used

a CUT&RUN protocol adapted from the kit manual from EpiCypher, with modifications to

account for the use of immune cells (*i.e*. to avoid unscheduled immune activation from ConA

signaling on cells). All buffers were made in-house.

1183

1184 *Nuclei isolation*

1185 250,000 MACS-purified peritoneal macrophages were harvested and resuspended in

1186 100µL of 1X DPBS. Samples were centrifuged for 3 min at 600g at room temperature and

supernatant was discarded. Cells were resuspended in 100µL of ice-cold Nuclear Extraction Buffer (20mM HEPES [Sigma-Aldrich #H3375], pH 7.9, 10mM KCl [Sigma-Aldrich # P9541], 0,1% Triton X-100 [VWR # M143], 20% Glycerol [VWR # BDH1172], 0.5mM Spermidine [EMD Millipore # 56766] and 1x Protease Inhibitor [Roche #11873580001]) and incubated on ice for 10 min. Samples were centrifuged for 3 min at 600g at 4 °C and supernatant was removed. Samples were resuspended in 100µL of ice-cold Nuclear Extraction Buffer and kept on ice until further processing.

Concanavalin A bead activation

11µL per sample of gently resuspended Concanavalin A (ConA) beads were transferred to fresh 1.5mL tubes. Tubes were placed on a magnetic separation rack until slurry cleared and supernatant was removed. 100µL of ice-cold Bead Activation Buffer (20 mM HEPES, pH 7.9, 10 mM KCl, 1 mM $CaCl_2$ and 1 mM $ManCl_2$) was added to each tube and pipetted 10 times to mix and wash the beads for a total of two washes. Washed beads were resuspended in 11µL of ice-cold Bead Activation Buffer. 10µL of activated bead slurry was transferred into fresh 1.5mL tubes. Beads were kept on ice until further processing.

Immunolabeling of chromatin

Isolated nuclei were collected by centrifugation at for 3 min at 600g at 4 °C and supernatant was discarded. Nuclei were resuspended in 100µL of Wash Buffer (20mM HEPES, pH 7.5, 150mM NaCl, 0.5mM Spermidine and 1x Protease Inhibitor) and transferred to the tubes containing the activated Concanavalin A beads. Samples were incubated at room temperature for 10 min. Tubes were placed on a magnetic separation rack until slurry cleared and supernatant was removed. 50µL of ice-cold Antibody Buffer (20mM HEPES, pH 7.5, 150mM NaCl, 0.5mM Spermidine, 1x Protease Inhibitor, 0.01% Digitonin and 2mM EDTA) was added to each sample. 0.5µL of anti-H3K4me3 antibody (Active Motif #39159) was added to each sample. Samples were incubated on a nutator overnight at 4°C. Tubes were placed on a magnetic separator rack until slurry cleared and supernatant was removed. While keeping the tubes on the magnetic separator, 250µL of ice-cold Digitonin Buffer (20mM HEPES, pH 7.5, 150mM NaCl, 0.5mM Spermidine and 1x Protease Inhibitor and 0.01% Digitonin) was added to the tubes and

1217   supernatant was removed without disturbing the tubes. Digitonin Buffer wash step was repeated

1218   once more for a total of two washes. 50µL of ice-cold Digitonin Buffer was added.

1219

pAG-MNase binding and DNA purification

2.5µL of CUTANA pAG-MNase (EpiCypher # 15-1016) was added to each sample and

gently mixed by vortexing. Samples were incubated at room temperature for 10 min and placed

on a magnetic separator. Supernatant was removed by pipetting. While the tubes were on the

magnet separator, 250µL of ice-cold Digitonin Buffer was added to each sample and removed.

Digitonin Buffer wash step was repeated once more for a total of two washes. 50µL of ice-cold

Digitonin Buffer was added to the tubes and gently mixed by vortexing. 1µL of 100mM $CaCl_2$

was added to each sample and gently mixed by vortexing. Samples were incubated on a nutator

for 2 hours at 4°C. 33µL of Stop Buffer (340mM NaCl, 20mM EDTA, 4mM EGTA, 50µg/mL

RNaseA, 50µg/mL GlycoBlue Coprecipitant) was added to each sample. Samples were

incubated at 37 °C for 10 min in a thermocycler. Samples were placed on a magnetic separator

until slurry cleared and the supernatant was collected to new 1.5 ml tubes. DNA purification was

performed using the DNA Clean & Concentration-5 kit (Zymo Research #D4013) according to

the manufacturer's instructions.

1234

Sequencing library preparation and sequencing

CUT&RUN sequencing libraries were prepared using the NEBNext Ultra II Library Prep

Kit for Illumina kit (NEBNext #E7645), according to the manufacturer's instructions. H4K4me3

CUT&RUN libraries were multiplexed and sent for sequencing on the Illumina NovaSeq 6000

platform to Novogene USA. De-multiplexed FASTQ files were provided by Novogene for data

analysis. Raw FASTQ sequencing reads were deposited to SRA under Bioproject accession

PRJNA1095555.

1242

1243

H4K4me3 CUT&RUN preprocessing pipeline

Paired-end H4K4me3 CUT&RUN reads were adapter-trimmed using NGmerge 0.3

(Gaspar, 2018b). Trimmed paired-end were mapped to the mm10 genome reference using

bowtie2 v2.5.1 (Langmead and Salzberg, 2012). After alignment, PCR duplicates were removed

using the 'rmdup' function of samtools 0.1.19 (Li et al., 2009). Reads from sequenced samples

were pooled in order to define "meta-peaks" for robust differential accessibility analysis using

samtools 'merge'. We used MACS2 v2.2.7.1 to identify H4K4me3 modified regions (Gaspar,

2018a). A normalized read count matrix was then obtained using our mapped read bam files over

MACS2 meta-peaks using R package 'Diffbind' 2.14 (Ross-Innes et al., 2012), which was used

for downstream analyses. 'DESeq2' 1.26.0 was used for normalization and differential gene

expression analysis of H4K4me3 CUT&RUN data in R (Love et al., 2014) using appropriate

analytical covariates (*e.g*. sex, age). Peaks with a FDR < 5% in any analysis were considered

statistically significant. For analysis of sex-specific age-regulation of H4K4me3 signal intensity,

we considered a chromatin region to be specifically remodeled with age in females (males) if the

DESeq2 FDR was < 5% in females (males) and > 10% in males (females). DESeq2 results are

reported in **Supplementary Table S2C**.

1260

1261   **Multidimensional Scaling analysis**

1262   To perform Multidimensional Scaling (MDS) analysis, we used a simple distance metric

1263   between samples based on the Spearman's rank correlation value (1-Rho), which was then

1264   provided to the core R command 'cmdscale'. Dimensionality reduction was applied to DESeq2-

1265   normalized data, obtained as described in relevant sections above.

1266

Putative Protein-Protein Interaction [PPI] network analysis with NetworkAnalyst

Genes with significant regulation with aging in female or male peritoneal macrophages

(DESeq2 FDR < 5%; **Supplementary Table S2A**) were used as input for network analysis with

NetworkAnalyst 3.0(Zhou et al., 2019). Significant age-regulated genes in each sex and their

corresponding DESeq2 computed $\log_2$(fold-change) were used as input. In both cases, we had

NetworkAnalyst use PPI information derived from IMEx/InnateDB data, a knowledgebase

specifically geared for analyses of innate immune networks(Breuer et al., 2013), to construct

putative PPI networks. In each case, the largest subnetwork determined by NetworkAnalyst was

used in our analyses (**Figure 3A, B**).

1276

1277   **Functional enrichment analysis by Gene Set Enrichment Analysis**

1278   We used the Gene Set Enrichment Analysis (GSEA) paradigm(Subramanian et al., 2005)
1279   through the 'phenotest' v1.34.0 package. Significance was calculated using 10,000 permutations.
1280

GSEA analysis using Gene Ontology and Reactome gene sets

Gene Ontology term annotation were obtained from ENSEMBL through Biomart
(Ensembl 108; downloaded on 2022-12-21), and gene-term association with only author
statement support (GO evidence codes 'TAS' and 'NAS') or unclear evidence (GO evidence
code 'ND') was filtered out. The Reactome annotations were obtained from the Molecular
Signature Database mouse v2022.1(Liberzon et al., 2011). For RNA-seq GSEA analysis,
differential gene regulation information and gene-centric statistics were obtained directly from
the DESeq2 result table without further processing. For ATAC-seq and H3K4me3 CUT&RUN
GSEA analyses, the peak-centric DESeq2 result tables were first processed to generate gene-
centric results compatible with GSEA. Specifically, only peaks with 10kb of annotated TSSs
according to HOMER v5.1 'annotatePeaks.pl' script(Heinz et al., 2010) were retained. For
ATAC-seq, to capture the most strongly remodeled regulatory elements, the gene-centric data
was obtained by capturing the DESeq2 statistics for the peak with the most significant DESeq2
FDR, and in the case of peaks tied for FDR, with the largest absolute $\log_2$(fold change) with
aging among tied peaks for each annotated gene. For H3K4me3 CUT&RUN, to capture the peak
most representative of gene promoters (since H3K4me3 is a modification typical of promoters),
the gene-centric data was obtained by capturing the DESeq2 statistics for the peak closest to each
annotated gene.

In all cases, from gene-centric result tables, the DESeq2 t-statistic was used to generate
the ranked list of genes for functional enrichment analysis with aging in female and male
macrophages separately. The 'gsea' function from phenotest' was run on gene sets with a size of
20 to 5000 genes, filtering too small or too large gene sets, so as to improve biological
robustness. For ease of reading, only the top 5 most significant sex-specific gene sets or top 10
most significant sex independent gene sets are shown in figures, with all shown gene sets passing
FDR < 5%. All significantly regulated gene sets are reported in **Supplementary Tables S3**.
1306

1307   *GSEA analysis of curated gene sets for top functional processes*

For curated gene set analyses (**Figure 3F-H**), we used curated gene sets for validation of important top curated processes. Specifically, we curated candidate regulators of phagocytosis from CRISPR/Cas9 screens based on results reported by Haney et al. 2018 and Pluvinage et al. 2019 (Haney et al., 2018; Pluvinage et al., 2019). We obtained mouse Hallmark curated gene sets "Glycolysis" and "Oxidative Phosphorylation" from the Molecular Signature Database mouse v2022.1 (Liberzon et al., 2011). We obtained a set of genes upregulated in M1 and M2 *in vitro* polarized BMDMs, by reanalyzing a publicly available RNA-seq dataset (accession number GSE103958)(Das et al., 2018). Briefly, downloaded FASTQ reads were mapped using STAR, a count table generated using featureCounts from subread, and differential gene expression analysis was performed using DESeq2 comparing M0 (untreated) vs. M1 (IFNg LPS treated) BMDMs, and M0 (untreated) vs. M1 (IL4 or IL13 treated) BMDMs. Genes with FDR < 5% and upregulated upon M1/M2 treatments were used as signatures of polarization for analysis (M1: 1392 genes; M2: 760 genes; **Supplementary Table S4C**).

GSEA analysis of macrophage-regulated gene sets

To compare the impact of various interventions on macrophage transcriptomes to that of aging in peritoneal macrophage from female mice, we also leveraged a GSEA approach. Specifically, we used our various RNA-seq experiments to identify genes significantly regulated in response to *in vivo* interventions (*i.e.* OVX vs. Sham, E2-supplementation vs. Vehicle, *Esr1* KO) or *in vitro* modulation (*i.e.* TF gene shRNA knockdown; see below).

In all cases, we performed DESeq2 differential gene expression analysis of our datasets to identify genes with significant regulation across our conditions (see above). For RNA-seq datasets measuring the impact of *in vivo* interventions on primary peritoneal macrophage transcriptomes, we obtained gene sets corresponding to genes significantly upregulated or downregulation in response to the intervention at FDR < 5%. These lists were then used as gene sets for GSEA analysis with phenotest (**Figure 5D, J, P**). For *in vitro* modulation of TF gene expression datasets, we used a more stringent definition for target gene sets to limit the impact of hairpin-specific off-target effects. Specifically, for each candidate TF, we performed differential expression analysis comparing control (i.e. non mammalian and luciferase hairpins) to each hairpin separately. To identify robust and consistent transcriptional targets, we then performed Fisher combination of adjusted p-values across both hairpins (analytical tables provided in

Supplementary Table S6C-G). Genes were considered to be significant transcriptional targets
of a candidate TF if and only if (i) they showed consistent direction of transcriptional regulation
across both hairpins, and (ii) if their Fisher combined adjusted p-values were $< 10^{-4}$ as a stringent
filter. Heatmaps showing consistent regulation for the chosen gene sets to be used for GSEA
analysis are reported in **Supplementary Figure S6G**, showing that our approach identifies
consistent transcriptional targets. These lists were then used as gene sets for GSEA analysis with
phenotest (**Figure 6G**).

1346

Growth and maintenance of cell lines

All cell lines were cultured in a humidified incubator under 5% CO_2-95% air at 37°C.
L929 cell line (ATCC CCL-1) were cultured with Dulbecco's Modified Eagle's Medium/ Hams
F-12 50/50 mix (DMEM/F-12 50/50; Corning #10-090-CV) supplemented with 10% fetal bovine
serum (FBS) (Sigma-Aldrich #F0926), and 1x penicillin streptomycin L-glutamine (Corning #
30-009-Cl). L929 cells were seeded in vented T175 tissue culture flasks (VWR #10062-864) at
10% confluency. Once cells reached around 80% confluency, conditioned media was collected
and filtered through a 0.45µM filter (GenClone #25-246). Conditioned media was stored at -
20°C until use (see below). HEK293T (ATCC CRL-3216), RAW264.7 (ATCC TIB71), and
J774A.1 (ATCC TIB-67) were cultured with Dulbecco's Modified Eagle's Medium (DMEM)
(Corning #15-013-CV) supplemented with 10% FBS, and 1x penicillin streptomycin L-
glutamine. Media was changed every 2-3 days. To passage cell lines, L929, HEK293T, and
RAW264.7 cells were detached using 0.05% trypsin (Coring #25-052-Cl) for 5 minutes at 37°C.
After incubation, trypsin was neutralized by adding double the volume of complete media to the
trypsin used. Cells were pelleted by centrifugation at 300g for 5 minutes at room temperature.
For passaging of J774A.1 cells, cells were detached using a cell lifter (VWR #76036-006). Cells
were collected and pelleted at 300g for 5 minutes at room temperature. Supernatant was
discarded and cells were resuspended with 1mL of complete media. Cells were seeded at lower
densities in a new tissue culture plate as needed based on experimental needs. All cell lines were
re-authenticated after establishing culture in the laboratory using the ATCC cell line STR
profiling service to confirm their identity in presented experiments.

1368

Culture of primary peritoneal macrophages and BMDMs

All primary macrophages were cultured in a humidified incubator under 5% CO_2-95% air

at 37°C. MACS-purified peritoneal macrophages were cultured in macrophage media

(DMEM/F-12 50/50 supplemented with 10% FBS, 1x antibiotic antimycotic (Gibco #15240-

062), 10% L929 conditioned media and 2ng/mL M-CSF (Miltenyi Biotec #130-101-706)) in

non-treated tissue culture plates. Bone marrow cells were obtained from the hind legs of aged

C57BL/6JNia, aged C57BL/6J and estrus cycle tracked C57BL/6NTac mice following as before

(Lu et al., 2020). Briefly, after bone marrow harvest following Amend et al. 2016

protocol(Amend et al., 2016) (Lu et al., 2020), cells were resuspended in 1mL of MACS

resuspension buffer and treated with 1x RBC lysis solution for 2 minutes at room temperature,

then centrifuged at 300g for 10minutes at 4°C. Resuspended RBC-lysed bone marrow cells were

then filtered on a 70µM filter to remove debris and clumps (Miltenyi Biotec #130-110-916).

Cells were then transferred to macrophage media for culture in T75 vented untreated tissue

culture flasks (Greiner Bio-One #658195) for differentiation into bone marrow derived

macrophages. After 3 days, the media was replaced with fresh macrophage media and

differentiation continued for a total of seven days after euthanasia and bone marrow harvest.

Obtained BMDMs were then used for downstream experiments (*i.e.* phagocytosis assay

(**Supplementary Figure S4B**) and Agilent Seahorse metabolic profiling (**Figure 4I, J,**

Supplementary Figure S4J, K).

1388

Analysis of macrophage phagocytic capacity

We used our previously described microscopy-based phagocytosis assay protocol(Lu et

al., 2020) and a new streamlined flow cytometry-based phagocytosis assay protocol to quantify

the phagocytic capacity of primary macrophages and macrophage cell lines.

Briefly, for the microscopy protocol, 75,000 macrophages were seeded in a 24-well plate

with a 12mm glass coverslip (Electron Microscopy Sciences #72231-01), with technical

duplicates for each animal. Macrophages were allowed to attach overnight, before treatment for

1 hour with 1mg/mL Alexa Fluor 488-labelled Zymosan bioparticles (Invitrogen #Z23373)

diluted in macrophage media. After 1-hour, Zymosan particles were washed three times with ice-

cold DPBS, and cells were fixed with 4% formaldehyde (Macron #H121-05) for 10 minutes at

room temperature. After fixation, coverslips were washed three times with DPBS and mounted

on microscope slides (VWR #48311-703) with prolong diamond antifade mounting medium with

DAPI (Invitrogen #P36962). Images of phagocytosis were acquired using the Keyence microscope (Keyence #BZ-X710) with the 10x objective. Stitched images of a 3 by 3 field were then counted by blinded experimenters. To quantify Zymosan-internalized (green fluorescent) cells and total cell number, images were analyzed using ImageJ software (Schneider et al., 2012). Total cell counts were determined by identifying DAPI-stained nuclei using the "Analyze Particles" function following appropriate thresholding. Due to the possibility of multiple phagosomes within a single cell, Zymosan-positive cells were manually counted to ensure accurate identification of individual phagocytic cells The phagocytosis index was determined by calculating the ratio of Alexa Fluor 488–positive cells to the total number of cells. Phagocytosis images for aging and estrus cycling has been deposited to Figshare under 10.6084/m9.figshare.29286194 and 10.6084/m9.figshare.29286206, respectively.

For our new flow cytometry-based protocol, 100,000 macrophages were first blocked with FcR-blocking reagent for 10 minutes at 37°C. For primary MACS-purified peritoneal macrophages, we also included an additional staining step with F4/80-APC (Miltenyi Biotec #130-116-547) for downstream data filtering. For both primary and immortalized macrophages, cells were then treated with Alexa Fluor 488-labelled Zymosan bioparticles as above, targeting with a multiplicity of infection (MOI) of 2 per macrophage for 30 minutes at 37°C. After treatment, macrophages were fixed with a final concentration of ice cold 2% formaldehyde solution (Macron #H121-05) at 4°C protected from light. Macrophages were washed twice and resuspended in 500uL of MACS resuspension buffer, then analyzed on the MACSQuant Analyzer 10 to phagocytosis capacity. Gating was determined using fluorescence-minus-one [FMO] controls for each color used in the experiment to ensure that positive populations were solely associated with the antibody/particle Analysis was performed in Flowlogic V8 software. Flow cytometry data for aging, sham vs. OVX, E2 supplementation, *Esr1* KO, RAW264.7 phagocytosis regulator knockdown, RAW264.7 transcription factor candidate knockdown, and J774A.1 Irf2 knockdown phagocytosis has been deposited to Figshare under 10.6084/m9.figshare.29161955, 10.6084/m9.figshare.29169836, 10.6084/m9.figshare.29169821, 10.6084/m9.figshare.29169860, 10.6084/m9.figshare.29169821, 10.6084/m9.figshare.29169872, 10.6084/m9.figshare.29169899, and 10.6084/m9.figshare.29286278, respectively.

Cell surface marker protein level quantification by flow cytometry

To estimate potential differences in peritoneal macrophage surface markers (*i.e*. Zymosan
pattern recognition receptors, M1/M2 markers) with respect to sex and age, 100,000 MACS-
purified peritoneal macrophages were subjected to Fc receptor blocking (Miltenyi Biotec #130-
092-575) and stained with corresponding antibodies according to the manufacturer's instructions.
For analysis of pattern recognition receptors, cells were stained with F4/80-Brilliant Violet 421
(1:20) (Biolegend #123131), Dectin-1-PE (1:10) (Miltenyi Biotec #130-102-284), and CD282
(Tlr2)-APC (1:50) (Miltenyi Biotec #130-120-052). For M1 marker detection, cells were stained
with CD80-FITC (1:20) (Biolegend #104705) and CD86-APC (1:20) (Biolegend #105011). For
M2 marker detection, cells were stained with CD163-PE (1:20) (Biolegend #111803) and
CD206-APC (1:20) (Biolegend #141707). After gating on F4/80 positive cells, we obtained
median fluorescent intensity (MFI) of stained cells for surface proteins of interest using flow
cytometry quantification on the Milteni Biotec MACSQuant Analyzer 10. Flow cytometry
results were processed using the FlowLogic V8 software. Flow cytometry data has been
deposited to Figshare under 10.6084/m9.figshare.29162144.

1446

1447 Metabolic profiling of macrophages using the Agilent Seahorse platform

50,000 BMDMs or transduced macrophage cell lines (RAW264.7, J774A.1) cells were
seeded in a 96-well Seahorse assay plate with macrophage media (or appropriate culture media)
and incubated overnight for cells to attach. To characterize the bioenergetics of cells, a XF–96
Flux Analyzer (Agilent Technologies) was used. Growth media was replaced with 175μL of XF
assay medium (specially formulated, unbuffered Dulbecco's modified Eagle's medium for XF
assay; Agilent Technologies #103575-100) supplemented with 1mM sodium pyruvate at pH 7.4
(Agilent Technologies #103578-100). The cells were kept in a non-CO_2 incubator for 60 minutes
at 37°C before placing in the analyzer. Oxygen consumption rate [OCR] and extracellular
acidification rate [ECAR] were measured using 3 cycles of 3 minutes wait and 4 minutes
measure time. Each biological replicate had three technical replicates and the results obtained
were averaged. Injection of compounds and calculating metabolic parameters were obtained by
following an established protocols for macrophages (Van den Bossche et al., 2015).

To normalize OCR and ECAR values to actual live/seeded cell number, the cell media
was removed from 96-well plates at the end of the assay and air died for around 10 minutes
before storage at -80°C at least overnight to initiate cell lysis. To finalize cell lysis, 100μL of

nuclease-free water was added to each well and incubated at 37°C for 1 hour, and plates were
refrozen for > 1 hour at -80°C. Plates were thawed at room temperature, and 200μg/mL Hoechst
(Thermo Scientific #H1399) in TNE buffer (10mM Tris [Thermo scientific #J22675-A1], 1mM
EDTA [VWR #BDH7830-1], 2M NaCl [Sigma-Aldrich #S5150-1L], pH 7.4) was added to each
well. Plates were read at excitation 346 and emission 460 on Molecular Devices SpectraMAX
M3. Fluorescent values were then used to divide the OCR and ECAR values to obtain
normalized, comparable values. Seahorse data for aging, estrus cycling, transcription factor
candidate knockdown in RAW264.7, and *Irf2* knockdown in J774A.1 has been deposited under
10.6084/m9.figshare.29169911, 10.6084/m9.figshare.29169920, 10.6084/m9.figshare.29169926
, and 10.6084/m9.figshare.29169938, respectively.
1473

Measurement of macrophage glycolytic function using Lactate-Glo
30,000 MACS-purified peritoneal macrophages or RAW264.7/J774A.1 cells were seeded
into a 96-well plate tissue culture plate (GenClone #25-109) with 100μL of fresh media. Then,
48 hours after seeding (primary peritoneal macrophages) or 24 hours (macrophage cell lines), the
conditioned media was collected and stored at -20°C until further processing. To note, for this
experiment, transduced RAW264.7/J774A.1 cells were cultured without puromycin (see below)
to avoid metabolic abnormalities due to the impact of puromycin on mitochondrial function
(Price and Verner, 1993). To measure the amount of lactate in the condition media, Lactate-Glo
(Promega #J5021) was used following to the manufacturer's protocol and luminescence values
were collected using Promega GLOMAX (Promega #GM3500) plate reader. Luminescence
values were normalized to actual live/seeded cell numbers using the same method as for the
Seahorse assay (see above).
1486

Lentiviral particle production for shRNA-mediated gene knockdown
In all cases, plasmid DNA was isolated from overnight cultures using mini-preps from
the Macherey-Nagel NucleoSpin Plasmid kit (Macherey-Nagel #740588.250) according to the
manufacturer's protocol, and DNA was sent to GENEWIZ or Plasmidsaurus for sequence
validation before use in any experiment.
We used a lentiviral shRNA particle approach to mediate gene-specific RNA interference
and evaluate the impact of decreased gene expression of candidate phagocytosis regulators or

TF-mediators on macrophage phenotypes. We obtained 3rd generation of lentiviral vector

packaging backbones from Addgene: pMDLg/pRRE (Addgene #12251), pRSV-Rev (Addgene

#12253), VSV.G (Addgene #14888). To guarantee single clone origin of laboratory isolates,

bacteria were re-streaked on a 100μg/mL carbenicillin (Sigma-Aldrich #C3416) Luria broth (LB;

Thermo Scientific #J75852-A1) 1% agar (Invitrogen #30391-023) plates and incubated overnight

at 37°C. For plasmid preparation, single colonies were isolated and inoculated into culture tube

containing 100μg/mL carbenicillin in LB media for culture at 37 °C with shaking for 16-20

hours for each construct. Plasmid DNAs were obtained using the NucleoBond Xtra midi plus EF

kit (Macherey-Nagel #740422.50) following manufacturer's recommendation.

MISSION® shRNA control and target bacterial glycerol stocks were obtained from

Sigma-Aldrich (**Supplementary Table S1G-H**). To obtain single bacteria colonies, frozen

glycerol stocks were inoculated into sterile culture tubes (Genesee Scientific #21-129) containing

0.5mL terrific broth (TB) (Sigma-Aldrich #T0918-1KG) without antibiotics and incubated at 37

°C with shaking for 30 minutes. Next, 50μL of bacteria pre-cultures was spread on prewarmed

100μg/mL carbenicillin TB 1% agar plate, and incubated for 16 hours at 37 °C. For plasmid

preparation, a single colony was isolated from the plate and inoculated into a culture tube

containing 100μg/mL TB, incubated at 37 °C with shaking for 16-20 hours for each construct.

To produce a large quantity and endotoxin free plasmid DNA, 200mL of 100μg/mL TB bacteria

culture was inoculated and incubated at 37°C with shaking for 20 hours. Plasmid DNAs were

obtained using the NucleoBond Xtra midi plus EF kit (Macherey-Nagel #740422.50) following

manufacturer's recommendation.

Lentiviral particles were produced by transfection of HEK293T cells using the calcium

phosphate transfection method (Green and Sambrook, 2012). Transfection mixtures were

prepared by combining plasmid DNA (contained pMDLg/pRRE (1,000 ng), VSV.G (500 ng),

pRSV-Rev (500 ng), and the pLKO shRNA plasmid of interest (1000ng)), with 2.5 M calcium

chloride (Sigma-Aldrich #C1016-500G) and mixing with 0.22μM filter sterilized 2xHEPES-

buffered saline ([HBS] 140mM NaCl, 1.5mM Na$_2$HPO$_4$·2H$_2$O (VWR #0348-500G), 50mM

HEPES)) to form DNA-calcium phosphate complexes. The DNA-calcium phosphate complexes

were formed for 2 minutes at room temperature and then added dropwise on HEK293T cells at

60-80% confluency. HEK293T were incubated with DNA-calcium phosphate complexes for 16

hours, then media was changed to fresh complete media. After 24 hours, supernatants containing viral particles were collected and filtered using a 0.45µM filter to remove cells.

To concentrate lentiviral particles before use, supernatant was carefully layered on top of a layer of 20% sucrose (Sigma-Aldrich #S0389-1KG) in DPBS solution in a Beckman Coulter polypropylene quick-seal centrifuge tube (Beckman Coulter #343414). Viral preps were centrifuged at 70,000 x g for 2 hours at 4°C in an Optima XPN-100 Ultracentrifuge (Beckman Coulter #B10053). After centrifugation, the layered supernatant was removed, and viral pellets were resuspended and concentrated 100x in ice cold DPBS. Lentiviral particles were aliquoted and stored at -80°C until experiments.

Lentiviral transduction of macrophage cell lines

Macrophage cell lines were seeded at a density of 30,000 cells per well in a 96-well tissue culture plate (GenClone #25-109) and incubated for 48 hours. During this period, cells were treated with various concentrations of puromycin in complete media (10, 8, 6, 4, 2,1µg/mL) (Sigma-Aldrich #P8833), to assess cytotoxicity. On the second day, 100µL of the culture medium in each well was replaced with fresh medium containing the corresponding concentration of puromycin. Following the manufacturer's instructions, CyQUANT™ XTT Cell Viability (Invitrogen #X12223) assay reagents were added to each well to assess cell viability. The plates were then incubated for 4 hours at 37□°C. Absorbance was measured at 450□nm and 660□nm using a microplate reader. Specific absorbance values were calculated by subtracting the 660□nm reference readings from the 450□nm readings.

For transduction, 200,000 cells from macrophage cell lines (RAW264.7, J774A.1) were seeded in a 12-well tissue culture plate (GenClone #25-109) one day before infection. The next day, cells were infected with viral particles at a MOI of 1 in the presence of 8µg/mL polybrene (Sigma-Aldrich #TR-1003-G). After 24 hours, the media was replaced with fresh complete media. 48 hours after infection, cells were selected using complete media with addition of 10µg/mL of puromycin (Sigma-Aldrich #P8833), as determined with our in house kill curve for these cells (see above). Molecular and/or functional profiling of modified cells was performed on puromycin-selected macrophage cells cultured in complete media supplemented with 10µg/mL puromycin, expect for metabolic assays where 24 hours prior, media was changed to complete media without puromycin.

1555

Reverse transcription quantified PCR verification of shRNA gene knockdown

Total RNA was reverse transcribed into complementary DNA (cDNA) using the Thermo
Scientific™ Maxima H Minus First Strand cDNA Synthesis Kit (Thermo scientific #K1682),
according to the manufacturer's instructions. Quantitative PCR was performed using the
SensiFAST SYBR No-ROX Kit (Bioline #BIO-98020) with gene-specific primers (For
sequences see **Supplementary Table S1I**) to assess relative gene expression. cDNA, primers,
and SYBR master mix were first combined in a PCR tube to ensure a homogeneous mixture,
then transferred into Mic qPCR-specific tubes (QuantaBio #95910-20) for amplification using
the Mic qPCR Cycler (Bio Molecular Systems #MIC-2). Gene expression levels were quantified
using the ΔΔCt method and normalized to the geometric mean of three housekeeping genes (*18s*,
Hprt, and *Ubc*).

1567

Vaginal cytology determination of estrus cycling

Vaginal lavages were obtained using a standard protocol (McLean et al., 2012). Lavages
were deposited to glass slides (VWR #48311-703) and left to dry for at least 24 hours. Vaginal
cytology slides were stained for 30 seconds 1% Thionin (Alfa Aesar #A18192), in 20mM acetic
acid (Sigma-Aldrich #695092-2.5L-GL), 0.36mM Sodium Hydroxide (Sigma-Aldrich #221465-
500G) at pH 4.0. Excess thionin staining was washed twice in MilliQ water. Slides were
dehydrated for 30 seconds in 70%, 95%, and 100% ethanol, then air dried overnight before
microscopic analysis. Slides were viewed under a microscope at 20x magnification identify the
estrus phase each mouse was in at the time of sample collection (**Supplementary Table S1C**).

1577

Motif analysis in remodeled chromatin regions

To identify potential TFs regulating sex-specific aging phenotypes in macrophages,
ATAC-seq and H3K4me3 peaks with age-related remodeling in only one sex were obtained (see
above for criteria). Motif finding was performed using HOMER v5.1 'findMotifsGenome.pl'
script(Heinz et al., 2010), using all robustly detected metapeaks as enrichment background and
the mm10 reference genome. We focused on results for enrichment of known motifs reported in
"knownResults.txt". Top TF motifs corresponding to TF genes significantly regulated with
female aging are shown in **Figure 6C, D**. All enriched motifs from our analyses are reported in

Supplementary Table S7A. To confirm enrichment of Irf2-related binding motifs, Irf2 peaks (see below) were used with the same pipeline, and selected motifs are reported in **Supplementary Figure S6F**.

SCENIC analysis of regulon activity in single cell RNA-seq analysis

As an orthogonal method to evaluate candidate TF activity in aging female and male macrophages, we took advantage of macrophages identified in our scRNA-seq datasets (see above) and applied the SCENIC algorithm to our data(Aibar et al., 2017). We applied the SCENIC pipeline using R packages SCENIC v1.1.2-01, RcisTarget v1.6.0, GENIE3 v1.8.0 and AUCell v1.8.0 following the package tutorial. Specifically, SCT normalized UMI counts from a combined Seurat object with macrophages transcriptomes from our 3 datasets were used as the input expression matrix. Mouse feather datasets for the mm10 assembly were downloaded from the SCENIC website (*i.e.* mm10__refseq-r80__500bp_up_and_100bp_down_tss.mc9nr.feather and mm10__refseq-r80__10kb_up_and_down_tss.mc9nr.feather). Computed TF regulon activities were obtained for 82 TFs in every cell, and we used a Wilcoxon rank-sum test approach to determine whether regulon activity was significantly impacted by aging in female or male macrophages (**Supplementary Table S6A**). Dot plots and violin plots were generated for TFs also identified from our bulk 'omic' data (**Supplementary Figure S6B, C**).

Irf2 protein level quantification using intracellular labeling and flow cytometry quantification

Cells were resuspended in 90μL of staining buffer and incubated with 10μL of FcR blocking reagent (Miltenyi Biotec #130-092-575) at 4□°C for 20 minutes. For MACS-purified peritoneal macrophage analysis, surface F4/80 staining was performed (to guarantee quantification of Irf2 expression solely on macrophages) by adding 2μL of anti-F4/80 antibody per sample, followed by a 10-minute incubation at 4□°C. Then, 2mL of resuspension buffer was added, and cells were centrifuged at 300g for 10 minutes to remove unbound antibodies. Supernatant were then removed, and cells were resuspended in 1mL of Foxp3 Fixation/Permeabilization working solution (Invitrogen #00-5523-00), pulse-vortexed, and incubated at 4□°C for 1 hour protected from light. Following fixation, 2□mL of 1x Permeabilization Buffer (Invitrogen #00-5523-00) was added, and samples were centrifuged at

600g for 5 minutes at room temperature. Supernatants were carefully decanted, leaving

approximately 200μL, and cells were gently resuspended. To block nonspecific binding, 10μL

normal rat serum was added, and samples were incubated at room temperature for 15 minutes.

Each sample was split into 98μL aliquots for intracellular staining with 2μL of directly

conjugated PE-IgG isotype control (Abcam #ab209478) or PE-Irf2 (Abcam #ab318372)

antibodies, and incubated at 4□°C for 1 hour, protected from light. After staining, cells were

washed twice with 2mL of 1x Permeabilization Buffer by centrifugation at 600g for 5 minutes at

room temperature. Finally, cells were resuspended in eBioscience Flow Cytometry Staining

Buffer (Invitrogen #00-4222-26). Stained cells median fluorescent intensity (MFI) was analyzed

by flow cytometry on the Miltenyi Biotec MACSQuant Analyzer 10, and flow cytometry results

were processed using the FlowLogic V8 software. To obtain the TF expression level, non-

specific background signal was removed by subtracting the MFI of IgG isotype control from the

MFI of Irf2 for each specific cell sample (i.e. isolate for a single animal, or transduced cell

aliquot)(Manso and Medina, 2021). TF levels were then normalized dividing the median entire

cohort per value to mitigate the impact of cohort-specific variation in fluorescence intensity. Raw

flow cytometry data for aging, RAW264.7 *Irf2* KD, J774A.1 has been deposited to Figshare

under 10.6084/m9.figshare.29285708, 10.6084/m9.figshare.29169947, and

10.6084/m9.figshare.29169959 respectively.

1635

Active Motif FactorPath Irf2 ChIP-seq

Red-blood cell lysed bone marrow cells from 4 months old female C57BL/6JNia mice

were differentiated into BMDMs in two independent biological replicates (pool of 2 animals per

replicate) as above. For each replicate ChIP experiment, we plated 10 million BMDMs from an

independent primary culture, which were fixed by adding 1:20 (v/v) ratio of freshly prepared

formaldehyde solution (11% formaldehyde (Sigma-Aldrich #F8775-25ML), 100□mM

NaCl,1□mM EDTA pH 8.0 (EMD Millipore #324506), and 50□mM HEPES pH = 8) to the cell

culture media, followed by incubation for 15□min at room temperature. Fixation was stopped by

adding 2.5□M glycine (Sigma-Aldrich #G7403-100G) solution at a 1:20 (v/v) ratio and

incubating for 5□min at room temperature. Cells were collected by scraping on ice and spun

down at 4°C 800g for 10 minutes. All subsequent steps were performed on ice. Cells were

collected and washed twice with chilled PBS (Sigma-Aldrich #P4474-500ML) with 0.5% Igepal

(Sigma-Aldrich #I8896-50ML). 1mM of phenylmethylsulfonyl fluoride (PMSF) (Sigma-Aldrich

#P7626-250MG) was added to the second wash before pelleting. Cell pellets were snap-frozen

on dry ice and sent to Active Motif for further processing.

Subsequent processing steps including chromatin extraction, fragmentation, antibody

precipitation, input, and library preparation were performed by Active Motif using a ChIP

validated antibody to IRF2 (Abcam #ab124744). Active motif generated ChIP-seq libraries from

the ChIP-DNA and a pooled Input control library using a custom Illumina library type protocol

on an automated system (Apollo 342, Wafergen Biosystems/Takara). ChIP-seq libraries were

sequenced on Illumina NovaSeq 6000 as 75 nucleotides single-end reads. Demultiplexed FASTQ

files were provided by Active Motif, and the ChIP-seq FASTQ reads were deposited to SRA

under Bioproject PRJNA1126274.

1659

BMDM Irf2 ChIP-seq analysis

Single-end Irf2 ChIP-seq reads were mapped to the mm10 genome reference using

bowtie1 v1.2.3(Langmead et al., 2009). After alignment, PCR duplicates were removed using the

FIXSeq algorithm(Hashimoto et al., 2014), as previously described(Benayoun et al., 2019). We

used MACS2 v2.2.7.1 to identify Irf2-bound regions in each of our 2 biological

replicates(Gaspar, 2018a). Consensus peaks were then calculated using MSPC v5.5.0(Jalili et al.,

2023), which identified 339 reproducibly bound regions (**Supplementary Table S7A**). Peaks

were annotated to the closest TSS using the HOMER 'annotatePeaks.pl' script(Heinz et al.,

2010) (**Supplementary Table S7B**). To compute functional enrichment for Irf2 target genes, we

leveraged the 'GREAT' 4.0.4 tool (McLean et al., 2010), a tool specifically optimized to identify

functional enrichment starting from genomic regions. We used the mm10 genome as the

background for enrichment and all other options were left to default parameters. Results were

exported as 'tsv', processed in R to filter significant enrichments with FDR < 5% and are

reported in **Supplementary Table S7C, D**. We also used R packages ChIPSeeker v1.22.1(Yu et

al., 2015) and TxDb.Mmusculus.UCSC.mm10.knownGene v3.10.0 to perform genomic

distribution annotation of peaks (**Supplementary Figure S7G**), together with clusterProfiler

v3.14.3 as an orthogonal approach for functional enrichment analysis of Irf2-bound regions (see

results in **Supplementary Table S7E, F**).

To determine whether chromatin remodeling (*i.e.* changes in ATAC-seq or H3K4me3

CUT&RUN signal) may occur in female peritoneal macrophages with aging at Irf2 bound sites,

we used R packages ChIPpeakAnno v3.20.1 (Zhu et al., 2010) to perform overlap analyses

(**Supplementary Figure S7H**) and NGSplot v2.61(Shen et al., 2014) to generate signal plots

(**Figure 7I**).

1683

Data availability

Sequencing data has been deposited to the Sequence Read Archive, under Bioprojects

PRJNA1094925, PRJNA1095555, PRJNA1096052, PRJNA1101936, PRJNA1126274,

PRJNA1222371.

Raw   flow   cytometry   data   (10.6084/m9.figshare.28934213,

10.6084/m9.figshare.29169674, 10.6084/m9.figshare.29169737, 10.6084/m9.figshare.29169707,

10.6084/m9.figshare.28934198, 10.6084/m9.figshare.29169578, 10.6084/m9.figshare.29169581,

10.6084/m9.figshare.29169587, 10.6084/m9.figshare.29161955, 10.6084/m9.figshare.29169836,

10.6084/m9.figshare.29169821, 10.6084/m9.figshare.29169860, 10.6084/m9.figshare.29169821,

10.6084/m9.figshare.29169872, 10.6084/m9.figshare.29169899, 10.6084/m9.figshare.29285708,

10.6084/m9.figshare.29169947,10.6084/m9.figshare.29169959,   and

10.6084/m9.figshare.29286278),   microscopy   images   (10.6084/m9.figshare.29286194,

10.6084/m9.figshare.29286206), and Agilent Seahorse data (10.6084/m9.figshare.29169911,

10.6084/m9.figshare.29169920, 10.6084/m9.figshare.29169926) have been deposited to

Figshare.

1699

1700   **Code Availability**

1701   All R code for this study was run using R version 3.6.3. Genomic data processing scripts

1702   have   been   made   available   on   the   Benayoun   lab   GitHub   repository   at

1703   https://github.com/BenayounLaboratory/ Macrophage_SexDim_Aging.

Acknowledgements

We are grateful to Dr. Caleb Finch, Dr. Raphael Williams, Clayton Baker, Dr. Marc Vermulst, Sarah Shemtov, Allison Chae, Courtney Shen, Wyatt Web, Evelyn Navar, Emily K. Wang, Prakroothi S. Danthi, Amy Huang, Rochelle W. Lai, Guan J. Lai and Claire Chung for scientific insights and assistance during the preparation of this manuscript. We thank USC DAR veterinarians Dr. Lynlee M. Stevey-Rindenow and Dr. Bradley Ahrens for performing the ovariectomy and sham surgeries. We thank Hemal Mehta at the USC Leonard Davis school of Gerontology Seahorse core facility for helping run Seahorse experiments. We thank Dr. Didier Trono for providing the pMDLg/pRRE (Addgene #12251) and pRSV-Rev (Addgene #12253) plasmids. We thank Dr. Tannishtha Reya for providing VSV.G (Addgene #14888) plasmid. We thank Dr. Kenneth S. Korach for providing the B6N(Cg)-Esr1$^{tm4.2Ksk}$/J (*Esr1* KO) mice(Hewitt et al., 2010) (JAX #026176).

This work was supported by a Diana Jacobs Kalman/AFAR Scholarships for Research in the Biology of Aging, NIA T32 AG052374, and AthenaDAO student leader award to R.J.L.; USC Provost's Undergrad Research Fellowships to E.E.W., E.H.L., and C.R.; GCRLE-2020 post-doctoral fellowship from the Global Consortium for Reproductive Longevity and Equality at the Buck Institute, made possible by the Bia-Echo Foundation to M.K.; NIAID R01 AI134987 to H.S.G.; core vouchers from the USC Leonard Davis School of Gerontology, NIA R00 AG049934, Pew Biomedical Scholar award #00034120, an innovator grant from the Rose Hills foundation, the Kathleen Gilmore Biology of Aging research award, a generous gift from Dr. Eric Hennigan, and Hevolution grant HF-GRO-23-1199072-28 to B.A.B. The authors acknowledge the Center for Advanced Research Computing [CARC] at the University of Southern California for providing computing resources that contributed to the research results reported within this publication (URL: https://carc.usc.edu). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Author contributions

R.J.L., N.S.K., and B.A.B. designed the study. R.J.L., N.S.K, A.P., performed live animal experiments. A.P. read vaginal cytology slides. R.J.L., S.C., M.K., L.A., I.Y.L., K.M., N.S.K., and B.A.B. dissected mouse tissues. R.J.L. and M.K. generated omics libraries. B.A.B. performed computational analyses. R.J.L. performed flow cytometry experiments. R.J.L.,

1735 E.E.W., I.Y.L., C.R., J.J., and S.B. imaged and analyzed microscopy phagocytosis data. R.J.L.,

1736 S.C., and A.X. performed phagocytosis flow cytometry and quantification analysis. R.J.L., S.C.,

1737 and E.H.L. performed metabolic profiling experiments. R.J.L., A.X., and B.A.B analyzed

1738 Seahorse experiments. R.J.L., E.H.L, S.P., C.R., I.Y.L. performed RT-qPCR analyses. R.J.L.,

1739 and S.C. perform intracellular Irf2 quantification. C.D.L. and H.S.G. contributed scientific

1740 insights to the project. R.J.L., S.C., and B.A.B. wrote the manuscript with input from all authors.

1741 All authors edited and commented on the final manuscript.

1742

References

(2014). In Sociality, Hierarchy, Health: Comparative Biodemography: A Collection of Papers, M. Weinstein, and M.A. Lane, eds. (Washington (DC)).

Aibar, S., González-Blas, C.B., Moerman, T., Huynh-Thu, V.A., Imrichova, H., Hulselmans, G., Rambow, F., Marine, J.-C., Geurts, P., Aerts, J.*, et al.* (2017). SCENIC: single-cell regulatory network inference and clustering. Nature Methods *14*, 1083-1086.

Aliaghaei, M., and Haun, J.B. (2022). Optimization of Mechanical Tissue Dissociation Using an Integrated Microfluidic Device for Improved Generation of Single Cells Following Digestion. Front Bioeng Biotechnol *10*, 841046.

Amend, S.R., Valkenburg, K.C., and Pienta, K.J. (2016). Murine Hind Limb Long Bone Dissection and Bone Marrow Isolation. J Vis Exp.

Appt, S.E., and Ethun, K.F. (2010). Reproductive aging and risk for chronic disease: Insights from studies of nonhuman primates. Maturitas *67*, 7-14.

Aran, D., Looney, A.P., Liu, L., Wu, E., Fong, V., Hsu, A., Chak, S., Naikawadi, R.P., Wolters, P.J., Abate, A.R.*, et al.* (2019). Reference-based analysis of lung single-cell sequencing reveals a transitional profibrotic macrophage. Nat Immunol *20*, 163-172.

Baker, C., and Benayoun, B.A. (2023). Menopause Is More Than Just Loss of Fertility. Public Policy Aging Rep *33*, 113-119.

Baker, C., Kim, M., and Benayoun, B.A. (2025). A researcher's guide to studying sex differences in immune aging. Trends Mol Med.

Barman, P.K., Shin, J.E., Lewis, S.A., Kang, S., Wu, D., Wang, Y., Yang, X., Nagarkatti, P.S., Nagarkatti, M., Messaoudi, I.*, et al.* (2022). Production of MHCII-expressing classical monocytes increases during aging in mice and humans. Aging Cell *21*, e13701.

Barrett, J.P., Costello, D.A., O'Sullivan, J., Cowley, T.R., and Lynch, M.A. (2015). Bone marrow-derived macrophages from aged rats are more responsive to inflammatory stimuli. J Neuroinflammation *12*, 67.

Benayoun, B.A., Pollina, E.A., Singh, P.P., Mahmoudi, S., Harel, I., Casey, K.M., Dulken, B.W., Kundaje, A., and Brunet, A. (2019). Remodeling of epigenome and transcriptome landscapes with aging in mice reveals widespread induction of inflammatory responses. Genome Res *29*, 697-709.

Berletch, J.B., Ma, W., Yang, F., Shendure, J., Noble, W.S., Disteche, C.M., and Deng, X.

(2015). Escape from X inactivation varies in mouse tissues. PLoS Genet *11*, e1005079.

Blommaert, E., Peanne, R., Cherepanova, N.A., Rymen, D., Staels, F., Jaeken, J., Race, V.,

Keldermans, L., Souche, E., Corveleyn, A*., et al.* (2019). Mutations in MAGT1 lead to a

glycosylation disorder with a variable phenotype. Proc Natl Acad Sci U S A *116*, 9865-9870.

Breuer, K., Foroushani, A.K., Laird, M.R., Chen, C., Sribnaia, A., Lo, R., Winsor, G.L.,

Hancock, R.E., Brinkman, F.S., and Lynn, D.J. (2013). InnateDB: systems biology of innate

immunity and beyond--recent updates and continuing curation. Nucleic Acids Res *41*, D1228-

1233.

Buenrostro, J.D., Giresi, P.G., Zaba, L.C., Chang, H.Y., and Greenleaf, W.J. (2013).

Transposition of native chromatin for fast and sensitive epigenomic profiling of open chromatin,

DNA-binding proteins and nucleosome position. Nat Methods *10*, 1213-1218.

Byles, V., Covarrubias, A.J., Ben-Sahra, I., Lamming, D.W., Sabatini, D.M., Manning, B.D., and

Horng, T. (2013). The TSC-mTOR pathway regulates macrophage polarization. Nat Commun *4*,

2834.

Cassado Ados, A., D'Imperio Lima, M.R., and Bortoluci, K.R. (2015). Revisiting mouse

peritoneal macrophages: heterogeneity, development, and function. Front Immunol *6*, 225.

Chavez-Galan, L., Olleros, M.L., Vesin, D., and Garcia, I. (2015). Much More than M1 and M2

Macrophages, There are also CD169(+) and TCR(+) Macrophages. Front Immunol *6*, 263.

Chen, G., Chen, X., Duan, X., Zhang, R., and Bai, C. (2024). Unraveling the roles of IFIT3 gene

and immune-metabolic pathways in psoriasis: a bioinformatics exploration for diagnostic

markers and therapeutic targets. Front Mol Biosci *11*, 1439837.

Cho, L., Kaunitz, A.M., Faubion, S.S., Hayes, S.N., Lau, E.S., Pristera, N., Scott, N., Shifren,

J.L., Shufelt, C.L., Stuenkel, C.A*., et al.* (2023). Rethinking Menopausal Hormone Therapy: For

Whom, What, When, and How Long? Circulation *147*, 597-610.

Cora, M.C., Kooistra, L., and Travlos, G. (2015). Vaginal Cytology of the Laboratory Rat and

Mouse: Review and Criteria for the Staging of the Estrous Cycle Using Stained Vaginal Smears.

Toxicol Pathol *43*, 776-793.

Corces, M.R., Trevino, A.E., Hamilton, E.G., Greenside, P.G., Sinnott-Armstrong, N.A.,

Vesuna, S., Satpathy, A.T., Rubin, A.J., Montine, K.S., Wu, B*., et al.* (2017). An improved

ATAC-seq protocol reduces background and enables interrogation of frozen tissues. Nat
Methods *14*, 959-962.

Das, A., Yang, C.S., Arifuzzaman, S., Kim, S., Kim, S.Y., Jung, K.H., Lee, Y.S., and Chai, Y.G.
(2018). High-Resolution Mapping and Dynamics of the Transcriptome, Transcription Factors,
and Transcription Co-Factor Networks in Classically and Alternatively Activated Macrophages.
Front Immunol *9*, 22.

Davies, L.C., Rosas, M., Smith, P.J., Fraser, D.J., Jones, S.A., and Taylor, P.R. (2011). A
quantifiable proliferative burst of tissue macrophages restores homeostatic macrophage
populations after acute inflammation. Eur J Immunol *41*, 2155-2164.

Dillon, S., Agrawal, S., Banerjee, K., Letterio, J., Denning, T.L., Oswald-Richter, K.,
Kasprowicz, D.J., Kellar, K., Pare, J., van Dyke, T.*, et al.* (2006). Yeast zymosan, a stimulus for
TLR2 and dectin-1, induces regulatory antigen-presenting cells and immunological tolerance. J
Clin Invest *116*, 916-928.

Dobin, A., Davis, C.A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P., Chaisson,
M., and Gingeras, T.R. (2013). STAR: ultrafast universal RNA-seq aligner. Bioinformatics *29*,
15-21.

Dube, C.T., Ong, Y.H.B., Wemyss, K., Krishnan, S., Tan, T.J., Janela, B., Grainger, J.R.,
Ronshaugen, M., Mace, K.A., and Lim, C.Y. (2022). Age-Related Alterations in Macrophage
Distribution and Function Are Associated With Delayed Cutaneous Wound Healing. Front
Immunol *13*, 943159.

Dunn, S.E., Perry, W.A., and Klein, S.L. (2024). Mechanisms and consequences of sex
differences in immune responses. Nat Rev Nephrol *20*, 37-55.

Fairweather, D., Beetler, D.J., McCabe, E.J., and Lieberman, S.M. (2024). Mechanisms
underlying sex differences in autoimmunity. J Clin Invest *134*.

Ferraye, L., Nieucel, M., Savignac, M., Chaumeil, J., and Guery, J.C. (2025). The Inactive X
Chromosome: A Genetic Driver of Female-Biased Rheumatic Autoimmune Disorders? Eur J
Immunol *55*, e202451331.

Finch, C.E. (2014). The menopause and aging, a comparative perspective. J Steroid Biochem
Mol Biol *142*, 132-141.

Fix, D.K., Ekiz, H.A., Petrocelli, J.J., McKenzie, A.M., Mahmassani, Z.S., O'Connell, R.M., and

Drummond, M.J. (2021). Disrupted macrophage metabolic reprogramming in aged soleus

muscle during early recovery following disuse atrophy. Aging Cell *20*, e13448.

Flanagan, K.L., Fink, A.L., Plebanski, M., and Klein, S.L. (2017). Sex and Gender Differences

in the Outcomes of Vaccination over the Life Course. Annu Rev Cell Dev Biol *33*, 577-599.

Franceschi, C., and Campisi, J. (2014). Chronic inflammation (inflammaging) and its potential

contribution to age-associated diseases. J Gerontol A Biol Sci Med Sci *69 Suppl 1*, S4-9.

Frick, K.M. (2009). Estrogens and age-related memory decline in rodents: what have we learned

and where do we go from here? Horm Behav *55*, 2-23.

Furgiuele, S., Descamps, G., Cascarano, L., Boucq, A., Dubois, C., Journe, F., and Saussez, S.

(2022). Dealing with Macrophage Plasticity to Address Therapeutic Challenges in Head and

Neck Cancers. Int J Mol Sci *23*.

Gadek, M., Shaw, C.K., Abdulai-Saiku, S., Saloner, R., Marino, F., Wang, D., Bonham, L.W.,

Yokoyama, J.S., Panning, B., Benayoun, B.A.*, et al.* (2025). Aging activates escape of the silent

X chromosome in the female mouse hippocampus. Sci Adv *11*, eads8169.

Gal-Oz, S.T., Maier, B., Yoshida, H., Seddu, K., Elbaz, N., Czysz, C., Zuk, O., Stranger, B.E.,

Ner-Gaon, H., and Shay, T. (2019). ImmGen report: sexual dimorphism in the immune system

transcriptome. Nat Commun *10*, 4295.

Gaspar, J.M. (2018a). Improved peak-calling with MACS2. bioRxiv, 496521.

Gaspar, J.M. (2018b). NGmerge: merging paired-end reads via novel empirically-derived models

of sequencing errors. BMC Bioinformatics *19*, 536.

Gautier, E.L., Ivanov, S., Williams, J.W., Huang, S.C., Marcelin, G., Fairfax, K., Wang, P.L.,

Francis, J.S., Leone, P., Wilson, D.B.*, et al.* (2014). Gata6 regulates aspartoacylase expression in

resident peritoneal macrophages and controls their survival. J Exp Med *211*, 1525-1531.

Ghosn, E.E., Cassado, A.A., Govoni, G.R., Fukuhara, T., Yang, Y., Monack, D.M., Bortoluci,

K.R., Almeida, S.R., Herzenberg, L.A., and Herzenberg, L.A. (2010). Two physically,

functionally, and developmentally distinct peritoneal macrophage subsets. Proc Natl Acad Sci U

S A *107*, 2568-2573.

Ginhoux, F., and Guilliams, M. (2016). Tissue-Resident Macrophage Ontogeny and

Homeostasis. Immunity *44*, 439-449.

Gordon, E.H., Peel, N.M., Samanta, M., Theou, O., Howlett, S.E., and Hubbard, R.E. (2017).

Sex differences in frailty: A systematic review and meta-analysis. Exp Gerontol *89*, 30-40.

Gordon, M.N., Osterburg, H.H., May, P.C., and Finch, C.E. (1986). Effective oral administration

of 17 beta-estradiol to female C57BL/6J mice through the drinking water. Biol Reprod *35*, 1088-

1095.

Goswami, D.G., Rubio, A.J., Mata, J., Munoz, S., Gallegos, A., and Walker, W.E. (2021). Large

Peritoneal Macrophages and Transitional Premonocytes Promote Survival during Abdominal

Sepsis. Immunohorizons *5*, 994-1007.

Green, M.R., and Sambrook, J. (2012). Molecular cloning : a laboratory manual, 4th ed. edn

(Cold Spring Harbor, N.Y: Cold Spring Harbor Laboratory Press).

Hajdarovic, K.H., Yu, D., Hassell, L.A., Evans, S., Packer, S., Neretti, N., and Webb, A.E.

(2022). Single-cell analysis of the aging female mouse hypothalamus. Nat Aging *2*, 662-678.

Haney, M.S., Bohlen, C.J., Morgens, D.W., Ousey, J.A., Barkal, A.A., Tsui, C.K., Ego, B.K.,

Levin, R., Kamber, R.A., Collins, H.*, et al.* (2018). Identification of phagocytosis regulators

using magnetic genome-wide CRISPR screens. Nat Genet *50*, 1716-1727.

Hansen, K.D., Brenner, S.E., and Dudoit, S. (2010). Biases in Illumina transcriptome sequencing

caused by random hexamer priming. Nucleic Acids Res *38*, e131.

Hao, Y., Hao, S., Andersen-Nissen, E., Mauck, W.M., 3rd, Zheng, S., Butler, A., Lee, M.J.,

Wilk, A.J., Darby, C., Zager, M.*, et al.* (2021). Integrated analysis of multimodal single-cell data.

Cell *184*, 3573-3587 e3529.

Hashimoto, T.B., Edwards, M.D., and Gifford, D.K. (2014). Universal count correction for high-

throughput sequencing. PLoS Comput Biol *10*, e1003494.

He, J., McLaughlin, R.P., van der Noord, V., Foekens, J.A., Martens, J.W.M., van Westen, G.,

Zhang, Y., and van de Water, B. (2019). Multi-targeted kinase inhibition alleviates mTOR

inhibitor resistance in triple-negative breast cancer. Breast Cancer Res Treat *178*, 263-274.

He, L., Cho, S., and Blenis, J. (2025). mTORC1, the maestro of cell metabolism and growth.

Genes Dev *39*, 109-131.

Heinz, S., Benner, C., Spann, N., Bertolino, E., Lin, Y.C., Laslo, P., Cheng, J.X., Murre, C.,

Singh, H., and Glass, C.K. (2010). Simple combinations of lineage-determining transcription

factors prime cis-regulatory elements required for macrophage and B cell identities. Mol Cell *38*,

576-589.

Heng, T.S., Painter, M.W., and Immunological Genome Project, C. (2008). The Immunological

Genome Project: networks of gene expression in immune cells. Nat Immunol *9*, 1091-1094.

Hewitt, S.C., Kissling, G.E., Fieselman, K.E., Jayes, F.L., Gerrish, K.E., and Korach, K.S.

(2010). Biological and biochemical consequences of global deletion of exon 3 from the ER alpha

gene. FASEB J *24*, 4660-4667.

Hirayama, D., Iida, T., and Nakase, H. (2017). The Phagocytic Function of Macrophage-

Enforcing Innate Immunity and Tissue Homeostasis. Int J Mol Sci *19*.

Hirokawa, K., Utsuyama, M., Hayashi, Y., Kitagawa, M., Makinodan, T., and Fulop, T. (2013).

Slower immune system aging in women versus men in the Japanese population. Immun Ageing

10, 19.

Hu, H., Miao, Y.R., Jia, L.H., Yu, Q.Y., Zhang, Q., and Guo, A.Y. (2019). AnimalTFDB 3.0: a

comprehensive resource for annotation and prediction of animal transcription factors. Nucleic

Acids Res *47*, D33-D38.

Huang, S., Zhao, H., Lou, X., Chen, D., Shi, C., and Ren, Z. (2024). TM6SF1 suppresses the

progression of lung adenocarcinoma and M2 macrophage polarization by inactivating the

PI3K/AKT/mtor pathway. Biochem Biophys Res Commun *718*, 149983.

Huang, Z., Chen, B., Liu, X., Li, H., Xie, L., Gao, Y., Duan, R., Li, Z., Zhang, J., Zheng, Y.*, et

al.* (2021). Effects of sex and aging on the immune cell landscape as assessed by single-cell

transcriptomic analysis. Proc Natl Acad Sci U S A *118*.

Jalili, V., Cremona, M.A., and Palluzzi, F. (2023). Rescuing biologically relevant consensus

regions across replicated samples. BMC Bioinformatics *24*, 240.

Jayakumar, P., Laganson, A., and Deng, M. (2022). GATA6(+) Peritoneal Resident Macrophage:

The Immune Custodian in the Peritoneal Cavity. Front Pharmacol *13*, 866993.

Kakugawa, S., Shimojima, M., Goto, H., Horimoto, T., Oshimori, N., Neumann, G., Yamamoto,

T., and Kawaoka, Y. (2009). Mitogen-activated protein kinase-activated kinase RSK2 plays a

role in innate immune responses to influenza virus infection. J Virol *83*, 2510-2517.

Kang, S., Ko, E.Y., Andrews, A.E., Shin, J.E., Nance, K.J., Barman, P.K., Heeger, P.S.,

Freeman, W.M., Benayoun, B.A., and Goodridge, H.S. (2024). Microglia undergo sex-dimorphic

transcriptional and metabolic rewiring during aging. J Neuroinflammation *21*, 150.

Kim, O.H., Kim, H., Kang, J., Yang, D., Kang, Y.H., Lee, D.H., Cheon, G.J., Park, S.C., and Oh,

B.C. (2017). Impaired phagocytosis of apoptotic cells causes accumulation of bone marrow-

derived macrophages in aged mice. BMB Rep *50*, 43-48.

Kim, Y.J., Tamadon, A., Park, H.T., Kim, H., and Ku, S.Y. (2016). The role of sex steroid

hormones in the pathophysiology and treatment of sarcopenia. Osteoporos Sarcopenia *2*, 140-

155.

Klein, S.L., and Flanagan, K.L. (2016). Sex differences in immune responses. Nat Rev Immunol

16, 626-638.

Kohama, S.G., Anderson, C.P., Osterburg, H.H., May, P.C., and Finch, C.E. (1989). Oral

administration of estradiol to young C57BL/6J mice induces age-like neuroendocrine

dysfunctions in the regulation of estrous cycles. Biol Reprod *41*, 227-232.

Kohama, S.G., Brown, S.A., Finch, C.E., and McNeill, T.H. (1992). Chronic estradiol

administration did not cause loss of hypothalamic LHRH or TIDA neurons in young or middle-

aged C57BL/6J mice. Brain Res *574*, 341-344.

Kowalczyk, M.S., Tirosh, I., Heckl, D., Rao, T.N., Dixit, A., Haas, B.J., Schneider, R.K.,

Wagers, A.J., Ebert, B.L., and Regev, A. (2015). Single-cell RNA-seq reveals changes in cell

cycle and differentiation programs upon aging of hematopoietic stem cells. Genome Res *25*,

1860-1872.

Krajcovic, M., Krishna, S., Akkari, L., Joyce, J.A., and Overholtzer, M. (2013). mTOR regulates

phagosome and entotic vacuole fission. Mol Biol Cell *24*, 3736-3745.

Kulminski, A.M., Culminskaya, I.V., Ukraintseva, S.V., Arbeev, K.G., Land, K.C., and Yashin,

A.I. (2008). Sex-specific health deterioration and mortality: the morbidity-mortality paradox

over age and time. Exp Gerontol *43*, 1052-1057.

Lai, J.H., Wu, D.W., Huang, C.Y., Hung, L.F., Wu, C.H., and Ho, L.J. (2024). USP18 induction

regulates immunometabolism to attenuate M1 signal-polarized macrophages and enhance IL-4-

polarized macrophages in systemic lupus erythematosus. Clin Immunol *265*, 110285.

Langmead, B., and Salzberg, S.L. (2012). Fast gapped-read alignment with Bowtie 2. Nat

Methods *9*, 357-359.

Langmead, B., Trapnell, C., Pop, M., and Salzberg, S.L. (2009). Ultrafast and memory-efficient

alignment of short DNA sequences to the human genome. Genome Biology *10*, R25.

Lavin, Y., Winter, D., Blecher-Gonen, R., David, E., Keren-Shaul, H., Merad, M., Jung, S., and

Amit, I. (2014). Tissue-resident macrophage enhancer landscapes are shaped by the local

microenvironment. Cell *159*, 1312-1326.

Lee, K.A., Flores, R.R., Jang, I.H., Saathoff, A., and Robbins, P.D. (2022). Immune Senescence,

Immunosenescence and Aging. Front Aging *3*, 900028.

Lee, O.J., Cho, Y.N., Kee, S.J., Kim, M.J., Jin, H.M., Lee, S.J., Park, K.J., Kim, T.J., Lee, S.S.,

Kwon, Y.S.*, et al.* (2014). Circulating mucosal-associated invariant T cell levels and their

cytokine levels in healthy adults. Exp Gerontol *49*, 47-54.

Lee, T.I., and Young, R.A. (2013). Transcriptional regulation and its misregulation in disease.

Cell *152*, 1237-1251.

Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis, G.,

Durbin, R., and Subgroup, G.P.D.P. (2009). The Sequence Alignment/Map format and

SAMtools. Bioinformatics *25*, 2078-2079.

Li, S.M., Valo, Z., Wang, J., Gao, H., Bowers, C.W., and Singer-Sam, J. (2012). Transcriptome-

wide survey of mouse CNS-derived cells reveals monoallelic expression within novel gene

families. PLoS One *7*, e31751.

Li, Z., Jiao, Y., Fan, E.K., Scott, M.J., Li, Y., Li, S., Billiar, T.R., Wilson, M.A., Shi, X., and

Fan, J. (2017). Aging-Impaired Filamentous Actin Polymerization Signaling Reduces Alveolar

Macrophage Phagocytosis of Bacteria. J Immunol *199*, 3176-3186.

Liao, Y., Smyth, G.K., and Shi, W. (2014). featureCounts: an efficient general purpose program

for assigning sequence reads to genomic features. Bioinformatics *30*, 923-930.

Liberzon, A., Subramanian, A., Pinchback, R., Thorvaldsdottir, H., Tamayo, P., and Mesirov,

J.P. (2011). Molecular signatures database (MSigDB) 3.0. Bioinformatics *27*, 1739-1740.

Linehan, E., Dombrowski, Y., Snoddy, R., Fallon, P.G., Kissenpfennig, A., and Fitzgerald, D.C.

(2014). Aging impairs peritoneal but not bone marrow-derived macrophage phagocytosis. Aging

Cell *13*, 699-708.

Liu, M., Silva-Sanchez, A., Randall, T.D., and Meza-Perez, S. (2021). Specialized immune

responses in the peritoneal cavity and omentum. J Leukoc Biol *109*, 717-729.

Lobo, R.A. (2007). Menopause and stroke and the effects of hormonal therapy. Climacteric *10*

Suppl 2, 27-31.

Loda, A., and Heard, E. (2019). Xist RNA in action: Past, present, and future. PLoS Genet *15*,
e1008333.

Loiola, R.A., Wickstead, E.S., Solito, E., and McArthur, S. (2019). Estrogen Promotes Pro-
resolving Microglial Behavior and Phagocytic Cell Clearance Through the Actions of Annexin
A1. Front Endocrinol (Lausanne) *10*, 420.

Lopez-Otin, C., Blasco, M.A., Partridge, L., Serrano, M., and Kroemer, G. (2023). Hallmarks of
aging: An expanding universe. Cell *186*, 243-278.

Louwe, P.A., Badiola Gomez, L., Webster, H., Perona-Wright, G., Bain, C.C., Forbes, S.J., and
Jenkins, S.J. (2021). Recruited macrophages that colonize the post-inflammatory peritoneal niche
convert into functionally divergent resident cells. Nat Commun *12*, 1770.

Love, M.I., Huber, W., and Anders, S. (2014). Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. Genome Biol *15*, 550.

Lu, R., Sampathkumar, N.K., and Benayoun, B.A. (2020). Measuring Phagocytosis in Bone
Marrow-Derived Macrophages and Peritoneal Macrophages with Aging. Methods Mol Biol
2144, 161-170.

Lu, R.J., Taylor, S., Contrepois, K., Kim, M., Bravo, J.I., Ellenberger, M., Sampathkumar, N.K.,
and Benayoun, B.A. (2021). Multi-omic profiling of primary mouse neutrophils predicts a
pattern of sex and age-related functional regulation. Nat Aging *1*, 715-733.

Lu, R.J., Wang, E.K., and Benayoun, B.A. (2022). Functional genomics of inflamm-aging and
immunosenescence. Brief Funct Genomics *21*, 43-55.

Lung, D.K., Reese, R.M., and Alarid, E.T. (2020). Intrinsic and Extrinsic Factors Governing the
Transcriptional Regulation of ESR1. Horm Cancer *11*, 129-147.

Manso, B.A., and Medina, K.L. (2021). Standardized flow-cytometry-based protocol to
simultaneously measure transcription factor levels. STAR Protoc *2*, 100485.

Marquez, E.J., Chung, C.H., Marches, R., Rossi, R.J., Nehar-Belaid, D., Eroglu, A., Mellert,
D.J., Kuchel, G.A., Banchereau, J., and Ucar, D. (2020). Sexual-dimorphism in human immune
system aging. Nat Commun *11*, 751.

Martin, C.J., Peters, K.N., and Behar, S.M. (2014). Macrophages clean up: efferocytosis and
microbial control. Curr Opin Microbiol *17*, 17-23.

Mastrototaro, G., Zaghi, M., Massimino, L., Moneta, M., Mohammadi, N., Banfi, F., Bellini, E.,
Indrigo, M., Fagnocchi, G., Bagliani, A., *et al.* (2021). TBL1XR1 Ensures Balanced Neural

Development Through NCOR Complex-Mediated Regulation of the MAPK Pathway. Front Cell Dev Biol *9*, 641410.

Matsuda-Lennikov, M., Biancalana, M., Zou, J., Ravell, J.C., Zheng, L., Kanellopoulou, C., Jiang, P., Notarangelo, G., Jing, H., Masutani, E.*, et al.* (2019). Magnesium transporter 1 (MAGT1) deficiency causes selective defects in N-linked glycosylation and expression of immune-response genes. J Biol Chem *294*, 13638-13656.

McGill, C.J., White, O.S., Lu, R.J., Sampathkumar, N.K., and Benayoun, B.A. (2024). Sex-dimorphic gene regulation in murine macrophages across niches. bioRxiv, 2024.2012.2020.629843.

McGinnis, C.S., Murrow, L.M., and Gartner, Z.J. (2019). DoubletFinder: Doublet Detection in Single-Cell RNA Sequencing Data Using Artificial Nearest Neighbors. Cell Syst *8*, 329-337 e324.

McLean, A.C., Valenzuela, N., Fai, S., and Bennett, S.A. (2012). Performing vaginal lavage, crystal violet staining, and vaginal cytological evaluation for mouse estrous cycle staging identification. J Vis Exp, e4389.

McLean, C.Y., Bristor, D., Hiller, M., Clarke, S.L., Schaar, B.T., Lowe, C.B., Wenger, A.M., and Bejerano, G. (2010). GREAT improves functional interpretation of cis-regulatory regions. Nat Biotechnol *28*, 495-501.

Miller, S.A., Policastro, R.A., Sriramkumar, S., Lai, T., Huntington, T.D., Ladaika, C.A., Kim, D., Hao, C., Zentner, G.E., and O'Hagan, H.M. (2021). LSD1 and Aberrant DNA Methylation Mediate Persistence of Enteroendocrine Progenitors That Support BRAF-Mutant Colorectal Cancer. Cancer Res *81*, 3791-3805.

Mogilenko, D.A., Shchukina, I., and Artyomov, M.N. (2022). Immune ageing at single-cell resolution. Nat Rev Immunol *22*, 484-498.

Mondal, S., and Rai, U. (1999). Sexual dimorphism in phagocytic activity of wall lizard's splenic macrophages and its control by sex steroids. Gen Comp Endocrinol *116*, 291-298.

Murray, P.J. (2017). Macrophage Polarization. Annu Rev Physiol *79*, 541-566.

Nguyen, H., Mustafa, A., Hiscott, J., and Lin, R. (1995). Transcription factor IRF-2 exerts its oncogenic phenotype through the DNA binding/transcription repression domain. Oncogene *11*, 537-544.

Ocanas, S.R., Pham, K.D., Cox, J.E.J., Keck, A.W., Ko, S., Ampadu, F.A., Porter, H.L., Ansere,

V.A., Kulpa, A., Kellogg, C.M.*, et al.* (2023). Microglial senescence contributes to female-

biased neuroinflammation in the aging mouse hippocampus: implications for Alzheimer's

disease. J Neuroinflammation *20*, 188.

Okabe, Y. (2021). Immune Niche Within the Peritoneal Cavity. Curr Top Microbiol Immunol

434, 123-134.

Ostan, R., Monti, D., Gueresi, P., Bussolotto, M., Franceschi, C., and Baggio, G. (2016). Gender,

aging and longevity in humans: an update of an intriguing/neglected scenario paving the way to a

gender-specific medicine. Clin Sci (Lond) *130*, 1711-1725.

Pang, W.W., Price, E.A., Sahoo, D., Beerman, I., Maloney, W.J., Rossi, D.J., Schrier, S.L., and

Weissman, I.L. (2011). Human bone marrow hematopoietic stem cells are increased in frequency

and myeloid-biased with age. Proc Natl Acad Sci U S A *108*, 20012-20017.

Panwar, V., Singh, A., Bhatt, M., Tonk, R.K., Azizov, S., Raza, A.S., Sengupta, S., Kumar, D.,

and Garg, M. (2023). Multifaceted role of mTOR (mammalian target of rapamycin) signaling

pathway in human health and disease. Signal Transduct Target Ther *8*, 375.

Pepe, G., Braga, D., Renzi, T.A., Villa, A., Bolego, C., D'Avila, F., Barlassina, C., Maggi, A.,

Locati, M., and Vegeto, E. (2017). Self-renewal and phenotypic conversion are the main

physiological responses of macrophages to the endogenous estrogen surge. Sci Rep *7*, 44270.

Peradinovic, J., Mohovic, N., Bulic, K., Markovinovic, A., Cimbro, R., and Munitic, I. (2023).

Ageing-Induced Decline in Primary Myeloid Cell Phagocytosis Is Unaffected by Optineurin

Insufficiency. Biology (Basel) *12*.

Perrotta, F., Corbi, G., Mazzeo, G., Boccia, M., Aronne, L., D'Agnano, V., Komici, K.,

Mazzarella, G., Parrella, R., and Bianco, A. (2020). COVID-19 and the elderly: insights into

pathogenesis and clinical decision-making. Aging Clin Exp Res *32*, 1599-1608.

Pluvinage, J.V., Haney, M.S., Smith, B.A.H., Sun, J., Iram, T., Bonanno, L., Li, L., Lee, D.P.,

Morgens, D.W., Yang, A.C.*, et al.* (2019). CD22 blockade restores homeostatic microglial

phagocytosis in ageing brains. Nature *568*, 187-192.

Price, J., and Verner, K. (1993). Puromycin inhibits protein import into mitochondria by

interfering with an intramitochondrial ATP-dependent reaction. Biochim Biophys Acta *1150*, 89-

97.

Qu, K., Zaba, L.C., Giresi, P.G., Li, R., Longmire, M., Kim, Y.H., Greenleaf, W.J., and Chang,
H.Y. (2015). Individuality and variation of personal regulomes in primary human T cells. Cell
Syst *1*, 51-61.

Ribas, V., Drew, B.G., Le, J.A., Soleymani, T., Daraei, P., Sitz, D., Mohammad, L., Henstridge,
D.C., Febbraio, M.A., Hewitt, S.C.*, et al.* (2011). Myeloid-specific estrogen receptor alpha
deficiency impairs metabolic homeostasis and accelerates atherosclerotic lesion development.
Proc Natl Acad Sci U S A *108*, 16457-16462.

Roelli, P., Flynn, B., and Gui, G. (2019). Hoohm/CITE-seq-Count: 1.4.2. Zenodo.

Ross, E.A., Devitt, A., and Johnson, J.R. (2021). Macrophages: The Good, the Bad, and the
Gluttony. Front Immunol *12*, 708186.

Ross-Innes, C.S., Stark, R., Teschendorff, A.E., Holmes, K.A., Ali, H.R., Dunning, M.J., Brown,
G.D., Gojis, O., Ellis, I.O., Green, A.R.*, et al.* (2012). Differential oestrogen receptor binding is
associated with clinical outcome in breast cancer. Nature *481*, 389-393.

Rossouw, J.E., Anderson, G.L., Prentice, R.L., LaCroix, A.Z., Kooperberg, C., Stefanick, M.L.,
Jackson, R.D., Beresford, S.A., Howard, B.V., Johnson, K.C.*, et al.* (2002). Risks and benefits of
estrogen plus progestin in healthy postmenopausal women: principal results From the Women's
Health Initiative randomized controlled trial. JAMA *288*, 321-333.

Russell, D.G., Huang, L., and VanderVen, B.C. (2019). Immunometabolism at the interface
between macrophages and pathogens. Nat Rev Immunol *19*, 291-304.

Saliba, A.E., Westermann, A.J., Gorski, S.A., and Vogel, J. (2014). Single-cell RNA-seq:
advances and future challenges. Nucleic Acids Res *42*, 8845-8860.

Santoro, A., Bientinesi, E., and Monti, D. (2021). Immunosenescence and inflammaging in the
aging process: age-related diseases or longevity? Ageing Res Rev *71*, 101422.

Saris, J., Li Yim, A.Y.F., Bootsma, S., Lenos, K.J., Franco Fernandez, R., Khan, H.N., Verhoeff,
J., Poel, D., Mrzlikar, N.M., Xiong, L.*, et al.* (2025). Peritoneal resident macrophages constitute
an immunosuppressive environment in peritoneal metastasized colorectal cancer. Nat Commun
16, 3669.

Satija, R., Farrell, J.A., Gennert, D., Schier, A.F., and Regev, A. (2015). Spatial reconstruction of
single-cell gene expression data. Nat Biotechnol *33*, 495-502.

Schneider, C.A., Rasband, W.S., and Eliceiri, K.W. (2012). NIH Image to ImageJ: 25 years of
image analysis. Nat Methods *9*, 671-675.

Scotland, R.S., Stables, M.J., Madalli, S., Watson, P., and Gilroy, D.W. (2011). Sex differences
in resident immune cell phenotype underlie more efficient acute inflammatory responses in
female mice. Blood *118*, 5918-5927.

Shen, L., Shao, N., Liu, X., and Nestler, E. (2014). ngs.plot: Quick mining and visualization of
next-generation sequencing data by integrating genomic databases. BMC Genomics *15*, 284.

Shi, Y., Ma, J., Li, S., Liu, C., Liu, Y., Chen, J., Liu, N., Liu, S., and Huang, H. (2024). Sex
difference in human diseases: mechanistic insights and clinical implications. Signal Transduct
Target Ther *9*, 238.

Sierra, I., Pyfrom, S., Weiner, A., Zhao, G., Driscoll, A., Yu, X., Gregory, B.D., Vaughan, A.E.,
and Anguera, M.C. (2023). Unusual X chromosome inactivation maintenance in female alveolar
type 2 cells is correlated with increased numbers of X-linked escape genes and sex-biased gene
expression. Stem Cell Reports *18*, 489-502.

Skene, P.J., and Henikoff, S. (2017). An efficient targeted nuclease strategy for high-resolution
mapping of DNA binding sites. Elife *6*.

Skinnider, M.A., Squair, J.W., Kathe, C., Anderson, M.A., Gautier, M., Matson, K.J.E., Milano,
M., Hutson, T.H., Barraud, Q., Phillips, A.A.*, et al.* (2021). Cell type prioritization in single-cell
data. Nat Biotechnol *39*, 30-34.

Squair, J.W., Skinnider, M.A., Gautier, M., Foster, L.J., and Courtine, G. (2021). Prioritization
of cell types responsive to biological perturbations in single-cell data with Augur. Nat Protoc *16*,
3836-3873.

Strindhall, J., Skog, M., Ernerudh, J., Bengner, M., Lofgren, S., Matussek, A., Nilsson, B.O., and
Wikby, A. (2013). The inverted CD4/CD8 ratio and associated parameters in 66-year-old
individuals: the Swedish HEXA immune study. Age (Dordr) *35*, 985-991.

Subramanian, A., Tamayo, P., Mootha, V.K., Mukherjee, S., Ebert, B.L., Gillette, M.A.,
Paulovich, A., Pomeroy, S.L., Golub, T.R., Lander, E.S.*, et al.* (2005). Gene set enrichment
analysis: A knowledge-based approach for interpreting genome-wide expression profiles.
Proceedings of the National Academy of Sciences *102*, 15545.

Sun, Z., Qu, J., Xia, X., Pan, Y., Liu, X., Liang, H., Dou, H., and Hou, Y. (2021). 17beta-
Estradiol promotes LC3B-associated phagocytosis in trained immunity of female mice against
sepsis. Int J Biol Sci *17*, 460-474.

Team, R.C. (2022). R: A language and environment for statistical computing. (Vienna, Austria:

R Foundation for Statistical Computing).

Thompson, W.W., Shay, D.K., Weintraub, E., Brammer, L., Bridges, C.B., Cox, N.J., and

Fukuda, K. (2004). Influenza-associated hospitalizations in the United States. JAMA *292*, 1333-

1340.

Van den Bossche, J., Baardman, J., and de Winther, M.P. (2015). Metabolic Characterization of

Polarized M1 and M2 Bone Marrow-derived Macrophages Using Real-time Extracellular Flux

Analysis. J Vis Exp.

Viola, A., Munari, F., Sanchez-Rodriguez, R., Scolaro, T., and Castegna, A. (2019). The

Metabolic Signature of Macrophage Responses. Front Immunol *10*, 1462.

Wang, L., Zhu, Y., Zhang, N., Xian, Y., Tang, Y., Ye, J., Reza, F., He, G., Wen, X., and Jiang,

X. (2024). The multiple roles of interferon regulatory factor family in health and disease. Signal

Transduct Target Ther *9*, 282.

Wang, Q., Li, M., Wu, T., Zhan, L., Li, L., Chen, M., Xie, W., Xie, Z., Hu, E., Xu, S., *et al.*

(2022). Exploring Epigenomic Datasets by ChIPseeker. Curr Protoc *2*, e585.

Wculek, S.K., Dunphy, G., Heras-Murillo, I., Mastrangelo, A., and Sancho, D. (2022).

Metabolism of tissue macrophages in homeostasis and pathology. Cell Mol Immunol *19*, 384-

408.

Wong, C.K., Smith, C.A., Sakamoto, K., Kaminski, N., Koff, J.L., and Goldstein, D.R. (2017).

Aging Impairs Alveolar Macrophage Phagocytosis and Increases Influenza-Induced Mortality in

Mice. J Immunol *199*, 1060-1068.

Wu, H., Luo, J., Yu, H., Rattner, A., Mo, A., Wang, Y., Smallwood, P.M., Erlanger, B.,

Wheelan, S.J., and Nathans, J. (2014). Cellular resolution maps of X chromosome inactivation:

implications for neural development, function, and disease. Neuron *81*, 103-119.

Yamamoto, H., Lamphier, M.S., Fujita, T., Taniguchi, T., and Harada, H. (1994). The oncogenic

transcription factor IRF-2 possesses a transcriptional repression and a latent activation domain.

Oncogene *9*, 1423-1428.

Yang, F., Babak, T., Shendure, J., and Disteche, C.M. (2010). Global survey of escape from X

inactivation by RNA-sequencing in mouse. Genome Res *20*, 614-622.

Yao, Y., Xu, X.H., and Jin, L. (2019). Macrophage Polarization in Physiological and

Pathological Pregnancy. Front Immunol *10*, 792.

Young, M.D., and Behjati, S. (2020). SoupX removes ambient RNA contamination from droplet-

based single-cell RNA sequencing data. Gigascience *9*.

Yu, G., Wang, L.-G., and He, Q.-Y. (2015). ChIPseeker: an R/Bioconductor package for ChIP

peak annotation, comparison and visualization. Bioinformatics *31*, 2382-2383.

Zhang, C., Hao, T., Bortoluzzi, A., Chen, M.H., Wu, X., Wang, J., Ermel, R., Kim, Y., Chen, S.,

and Chen, W. (2025). Sex-dependent differences in hematopoietic stem cell aging and

leukemogenic potential. Oncogene *44*, 64-78.

Zheng, G.X., Terry, J.M., Belgrader, P., Ryvkin, P., Bent, Z.W., Wilson, R., Ziraldo, S.B.,

Wheeler, T.D., McDermott, G.P., Zhu, J.*, et al.* (2017). Massively parallel digital transcriptional

profiling of single cells. Nat Commun *8*, 14049.

Zhou, G., Soufan, O., Ewald, J., Hancock, R.E.W., Basu, N., and Xia, J. (2019). NetworkAnalyst

3.0: a visual analytics platform for comprehensive gene expression profiling and meta-analysis.

Nucleic Acids Res *47*, W234-W241.

Zhu, L.J., Gazin, C., Lawson, N.D., Pagès, H., Lin, S.M., Lapointe, D.S., and Green, M.R.

(2010). ChIPpeakAnno: a Bioconductor package to annotate ChIP-seq and ChIP-chip data. BMC

Bioinformatics *11*, 237.

Zuo, J., Tang, J., Lu, M., Zhou, Z., Li, Y., Tian, H., Liu, E., Gao, B., Liu, T., and Shao, P.

(2021). Glycolysis Rate-Limiting Enzymes: Novel Potential Regulators of Rheumatoid Arthritis

Pathogenesis. Front Immunol *12*, 779787.

Zylicz, J.J., Bousard, A., Zumer, K., Dossin, F., Mohammad, E., da Rocha, S.T., Schwalb, B.,

Syx, L., Dingli, F., Loew, D.*, et al.* (2019). The Implication of Early Chromatin Changes in X

Chromosome Inactivation. Cell *176*, 182-197 e123.

Zysow, B.R., Kauser, K., Lawn, R.M., and Rubanyi, G.M. (1997). Effects of estrus cycle,

ovariectomy, and treatment with estrogen, tamoxifen, and progesterone on apolipoprotein(a)

gene expression in transgenic mice. Arterioscler Thromb Vasc Biol *17*, 1741-1745.

2190

2191

2192

2193

2194

2195

Legends to figures

Figure 1. Peritoneal macrophages are the most perturbed cell type within the peritoneal cavity immune niche as a function of age and biological sex.

(**A**) Experimental scheme illustrating age-based cell profiling in mouse peritoneal lavage. (**B**) Uniform Manifold Approximation and Projection (UMAP) visualization of unsupervised clustering of peritoneal lavage cells. (**C**) UMAP representation of AUGUR area under the curve (AUC) scores for peritoneal lavage immune cell populations. (**D**) Boxplots of peritoneal cell composition quantified by scRNA-seq; n = 8 libraries per sex and age; 4 independent cohorts in 3 independent datasets. (**E**) Boxplots of peritoneal cell composition quantified by flow cytometry; n ≥ 25 per sex and age (6 independent cohorts). (**F**) Boxplots of peritoneal immune lavage cell number calculated by the proportion obtained from flow cytometry and total peritoneal lavage cell count; n ≥ 25 per sex and age (6 independent cohorts). For boxplots in panels (**D-F**), circles/squares represent NIA/JAX mice, respectively. Significance in non-parametric two-sided Wilcoxon rank-sum tests are reported in **D-F**. The center line of the box plots represents the sample median, the box limits consist of the 25th and 75th percentiles, the whiskers span 1.5x the interquartile range.

Figure 2. Peritoneal macrophage transcriptome and epigenome are highly sex-dimorphic with age.

(**A**) Experimental scheme of murine peritoneal macrophage 'omics' profiling. (**B**) Multidimensional scaling (MDS) plot of transcriptomic profiles, indicating distinct transcriptomic aging patterns between sexes. (**C**) MDS plot of chromatin accessibility profiles by ATAC-seq. (**D**) MDS plot of H3K4me3 profiles. (**E**) Scatterplot of age-related changes in peritoneal macrophage transcriptomes in female vs. males, reported as DESeq2 log$_2$ fold changes per month. 284 Male-specific (blue) and 234 female-specific (pink) aging-regulated genes (Transcriptome-wide Spearman Rho = 0.288). (**F**) Scatterplot of age-related changes in peritoneal macrophage ATAC-seq accessible regions in female vs. males, reported as DESeq2 log$_2$ fold changes per month. 1044 Male-specific (blue) and 2825 female-specific (pink) regions (Genome-wide Spearman Rho = 0.232). (**G**) Scatterplot of age-related changes in peritoneal macrophage H3K4me3 signal in female vs. males, reported as DESeq2 log$_2$ fold changes per

month. 131 male-specific (blue) and 1699 female-specific (pink) marks (Genome-wide

Spearman Rho = 0.329). (**H**) Heatmap of male-specific aging gene expression changes by RNA-

seq. (**I**) Heatmap of female-specific aging gene expression changes by RNA-seq. (**J**) Heatmap of

male-specific aging chromatin accessibility changes by ATAC-seq. (**K**) Heatmap of female-

specific aging chromatin accessibility changes by ATAC-seq. (**L**) Heatmap of male-specific

aging H3K4me3 peak intensity by H3K4me3 CUT&RUN. (**M**) Heatmap of female-specific

H3K4me3 peak intensity by H3K4me3 CUT&RUN.

2234

Figure 3. Functional enrichment analyses reveal convergent and sex-dimorphic phenotypes

in aging peritoneal macrophages.

(**A**, **B**) NetworkAnalyst putative PPI network of genes changing with age in female (**A**) and male

(**B**) peritoneal macrophages using IMEx network information. (**C**, **D**, **E**) Top 5 sex-dimorphic

GSEA enrichments for aging in each sex for RNA-seq (**C**), ATAC-seq (**D**), and H3K4me3

CUT&RUN (**E**). (**F**) GSEA of CRISPR-Cas9 phagocytosis regulators from screens in aging

female and male peritoneal macrophages. (**G**) GSEA of metabolic gene sets from MSigDB

Hallmarks in aging female and male peritoneal macrophages. (**H**) GSEA of M1/M2 polarization

genes in aging female and male peritoneal macrophages. NES: Normalized Enrichment Score.

FDR: False Discovery Rate.

2245

Figure 4. Female-specific macrophage remodeling in phagocytosis and metabolism.

(**A**) Experimental scheme of murine peritoneal macrophage purification and functional analysis.

(**B**) Representative microscopy image showing a phagocytosis event by young old male and

female MACS-purified peritoneal macrophages engulfing Zymosan bioparticles. Blue = DAPI,

Green = Zymosan bioparticles, Scale bar: 200μm. (**C**) Phagocytosis quantification of

phagocytosis images from young old male and female peritoneal macrophages. The phagocytosis

index is calculated by dividing phagocytosing cells by the total number of imaged cells. Due to

cohort-to-cohort variation, values are normalized to the median value of each independent

experiment. Animals from 5 independent cohorts (3 from NIA, 2 from JAX). Young females n =

21, old females n = 18, young males n = 22, and old males n = 20 (variation due to animal death

prior to experimental day). (**D**) Flow cytometry phagocytosis quantification of peritoneal

macrophage. Each point represents one animal. Animals from 6 independent cohorts (3 from

NIA, 3 from JAX). Circles/squares represent NIA/JAX mice, respectively Samples with < 5000

F4/80$^+$ events or < 75% F4/80$^+$ purity were excluded from the analysis. (**E**) Venn diagram for

CRISPR-Cas9 phagocytosis regulators and female-specific down-regulated genes according to

DESeq2 at FDR < 5% and < 10%. (**F**) Lentiviral shRNA phagocytosis regulator screen

experimental scheme. (**G**) Mini shRNA screen of RAW264.7 phagocytosis capacity using flow

cytometry. The horizontal red line in the panel shows the median control value in the relative

phagocytosis index of the macrophage. Dark green/light green for sh1/sh2 samples. Significance

in non-parametric one-sided Wilcoxon rank-sum tests were reported (Hypothesis: same

regulation direction as aging). (**H**) Experimental scheme of aging bone marrow-derived

macrophage seahorse. (**I**) Representative normalized extracellular acidification rate (ECAR)

mpH/min/DNA from 1 cohort of animals. (**J**) Glycolysis quantification using Seahorse (n = 15

for young old females and males; animals from 3 independent NIA cohorts). (**K**) Glycolysis

quantification using Lactate-Glo (n = 18 for young males, n = 18 for young females, n = 17 for

old males, and n = 17 for old females; animals from 5 independent cohorts (3 from NIA, 2 from

JAX). For boxplots in panels (**C-D**, **J-H**), circles/squares represent NIA/JAX mice, respectively.

Significance in non-parametric two-sided Wilcoxon rank-sum tests were reported. (**C-D**, **G**, **J-H**)

The center line of the box plots represents the sample median, the box limits consist of the 25th

and 75th percentiles, the whiskers span 1.5x the interquartile range.

2276

Figure 5. Female-specific macrophage phagocytosis declines due to decreased estrogen

signaling through the Esr1 receptor.

(**A**) Experimental scheme outlining the analysis of OVX mice. (**B**) Multi-dimensional scaling

(MDS) analysis of gene expression profiles from Sham and OVX mice. (**C**) GSEA of OVX-

regulated genes in the aging female peritoneal macrophage transcriptome. (**D**) Boxplot of

phagocytosis quantification of peritoneal macrophages in Sham and OVX mice using flow

cytometry (n = 11 for Sham, n = 12 for OVX; animals from 3 independent cohorts). (**E**) Boxplot

of glycolysis quantification of peritoneal macrophages in Sham and OVX mice (n = 12 for Sham,

n = 11 for OVX; 2 independent cohorts). (**F**) Experimental scheme describing late-life short-term

17β-estradiol (E2) supplementation in female mice. (**G**) MDS analysis of gene expression

profiles in young females, old females, and E2-treated old females; n = 5 for young females, n =

5 for old females, n = 4 for E2-treated old females. (**H**) GSEA of E2-regulated genes in old

female macrophages in the aging female peritoneal macrophage transcriptome. (**I**) Boxplot of phagocytosis of peritoneal macrophages in young females, old females, and E2-treated old females (n = 13 for young females, n = 13 for old females, n = 14 for E2-treated old females; animals from 3 independent cohorts). (**J**) Boxplot of glycolysis quantification of peritoneal macrophages in young females, old females, and E2-treated old females (n = 14 for young females, old females, and E2-treated old females; animals from 3 independent cohorts). (**K**) Experimental scheme for *Esr1* knockout (KO) mice. (**L**) MDS analysis of gene expression profiles in WT and *Esr1* KO mice (n = 5 for WT, n = 6 for *Esr1* KO). (**M**) GSEA of *Esr1* KO-regulated genes in the aging female peritoneal macrophage transcriptome. (**N**) Boxplot of phagocytosis of peritoneal macrophages in WT and *Esr1* KO mice (n = 15 for WT, n = 13 for *Esr1* KO; animals from 4 independent cohorts). (**O**) Boxplot of glycolysis quantification of peritoneal macrophages in WT and Esr1 KO mice (n = 15 for WT and *Esr1* KO; animals from 4 independent cohorts). For boxplots in panels (**D**, **E**, **I**, **J**, **N**, **O**), circles/squares represent NIA/JAX mice, respectively. Significance in non-parametric two-sided Wilcoxon rank-sum tests were reported. The center line represents the sample median, the box limits consist of the 25th and 75th percentiles, and the whiskers span 1.5x the interquartile range.

Figure 6. A mini shRNA screen for transcription factors driving in female-specific macrophage remodeling with aging.

(**A**) Flow chart outlining the steps for the selection of candidate transcription factors (TFs) for screening. (**B**) Heatmap displaying female-specific differential transcription factor gene expression with aging (FDR < 5% in aging female macrophages, FDR > 10% in aging male macrophages based on DESeq2 analysis). (**C**) Selected enriched motifs identified at female-specific differentially accessible ATAC-seq peaks with aging (HOMER; FDR < 5%). (**D**) Selected enriched motifs identified at female-specific differentially marked H3K4me3 peaks with aging (HOMER; FDR < 5%). (**E**) Experimental scheme of shRNA-mediated TF knockdown in RAW264.7 macrophages. (**F**) GSEA of TF shRNA knockdown regulated genes in the aging female peritoneal macrophage transcriptome. (**G**, **H**, **I**) Boxplots showing phagocytosis quantification using flow cytometry (**G**), glycolysis quantification using Agilent Seahorse assays (**H**), and lactate levels measured with Lactate-Glo in RAW264.7 macrophages transduced with lentiviral TF shRNA against (**I**) *Irf2*, *Mef2c*, *Meis1*, *Tal1*, and *Tbl1xr1* (n = 6 independent

infections for each hairpin). Dark green/light green for sh1/sh2 samples. Significance in non-
parametric one-sided Wilcoxon rank-sum tests were reported (Hypothesis: same direction as
aging). The horizontal red line in the panel shows the median control value in the relative
phagocytosis index (**G**), relative glycolysis (**H**), and relative lactate secretion (**I**) of macrophages.
The center line represents the sample median, the box limits show the 25^{th} and 75^{th} percentiles,
and the whiskers span 1.5 times the interquartile range.

2326

Figure 7. Irf2 impacts Hk3 leading to an increase in glycolytic dependence.
(**A**) Analysis of Irf2 expression at the RNA and protein levels in aging peritoneal macrophages.
(**B**) Boxplot of *Irf2* gene expression in mouse peritoneal macrophages using RT-qPCR (young
female n = 12, old female n = 8, young male n = 12, and old male n = 9; animals from 4
independent cohorts). (**C**) Histogram of female and male Irf2 protein expression between young
and old peritoneal macrophages based on intracellular immunolabelling and flow cytometry
quantification. (**D**) Boxplot of Irf2 protein quantification in peritoneal macrophages between
young and old female and male mice (young female n = 13, old female n = 14, young male n =
14, and old male n = 13; animals from 3 independent cohorts). (**E**) Experimental scheme for Irf2
ChIP-seq in young female BMDMs. (**F**) Top GREAT enriched terms of Irf2-bound genomic
regions. **(G)** GSEA of direct Irf2 target genes in the aging female peritoneal macrophage
transcriptome. Peaks were assigned to target genes based on their absolute distance to
transcriptional start sites (500bp-5kb). (**H**) Integrated Genomics Viewer (IGV) snapshot of Irf2
ChIP-seq signal at the *Hk3* gene locus on chromosome 13. (**I**) Quantitative RT-qPCR analysis of
Hk3 mRNA levels in young and old female peritoneal macrophages samples (n = 12 young, n =
8 old female; animals from 4 independent cohorts). (**J**) Experimental design for lentiviral
knockdown of *Irf2* in J774A.1 cells. (**K**) Boxplot of *Hk3* expression from RNA-seq after
knockdown of *Irf2* in macrophage cells. Circles/squares for RAW264.7/J774A.1 samples;
shLuciferase (goldenrod), non-targeting control (light goldenrod), sh1 (dark green), and sh2
(light green); n = 3 infections per hairpin. Significance as FDR from DESeq2. (**L**) Boxplot of
glycolysis quantification using Seahorse assays on lentiviral knockdown of *Irf2* RAW264.7 and
J774A.1 cell line (n = 6 infections per hairpin). Note that the RAW264.7 data is repeated from
Figure 6I for comparison convenience. (**M**) Boxplot of glycolysis quantification using Lactate-
Glo on lentiviral knockdown *Irf2* in RAW264.7 and J774A.1 cells (n = 6 infections per hairpin).

Note that the RAW264.7 data is repeated from **Figure 6J** for comparison convenience. (**N**)

Proposed mechanism for Irf2-driven age-related glycolytic shift in aging female peritoneal

macrophages. For boxplots in panels (**B**, **D**, **I**), circles/squares represent NIA/JAX mice,

respectively. Significance in non-parametric two-sided Wilcoxon rank-sum tests is reported. For

boxplots in panels (**L**, **M**) circles/squares for RAW264.7/J774A.1 samples; dark green/light

green for sh1/sh2 samples. Significance in non-parametric one-sided Wilcoxon rank-sum tests

were reported (Hypothesis: same regulation direction as aging). The horizontal red line in the

panel shows the median control value of the normalized glycolysis (**L**) and normalized lactate

(**M**).

Figure 2

A Experimental scheme

Young (4-5 months)
Old (20-21 months)

Peritoneal lavage cells → MACS purification → Macrophages > 90% Cd11b+ F4/80+ → RNA-seq / ATAC-seq / H3K4me3 CUT&RUN

B RNA-seq

C ATAC-seq

D H3K4me3

E Sex-dimorphic aging changes (RNA-seq)
- Male-specific aging (284)
- Female-specific aging (234)
Rho = 0.288

F Sex-dimorphic aging changes (ATAC-seq)
- Male-specific aging (1044)
- Female-specific aging (2825)
Rho = 0.232

G Sex-dimorphic aging changes (H3K4me3)
- Male-specific aging (131)
- Female-specific aging (1699)
Rho = 0.329

H Male-specific aging genes (RNA-seq)

J Male-specific aging peaks (ATAC-seq)

L Male-specific aging peaks (H3K4me3)

I Female-specific aging genes (RNA-seq)

K Female-specific aging peaks (ATAC-seq)

M Female-specific aging peaks (H3K4me3)

● Young females (4-5 months) ● Old females (20-21 months) ● Young males (4-5 months) ● Old males (20-21 months)

Figure 3

A Putative PPI network of genes changing with age in females (IMEx)

B Putative PPI network of genes changing with age in males (IMEx)

Up with age

Down with age

C Top GSEA enrichments for sex-dimorphic aging from RNA-seq (GO)

- **GO:0006910 phagocytosis, recognition**
- GO:0061077 chaperone-mediated protein folding
- GO:0042470 melanosome
- GO:0042310 vasoconstriction
- GO:0019373 epoxygenase P450 pathway
- GO:0015035 protein-disulfide reductase activity
- **GO:0015078 proton transmembrane transporter activity**
- **GO:1902600 proton transmembrane transport**
- **GO:0005751 mitochondrial respiratory chain complex IV**
- **GO:1900017 pos. regulation of cytokine production involved in inflammatory response**

Female | Male

D Top GSEA enrichments for sex-dimorphic aging from ATAC-seq (GO)

- GO:0005540 hyaluronic acid binding
- GO:0098656 anion transmembrane transport
- GO:0031683 G-protein beta/gamma-subunit complex binding
- GO:0060088 auditory receptor cell stereocilium organization
- GO:0031641 regulation of myelination
- **GO:0045104 intermediate filament cytoskeleton organization**
- GO:0034364 high-density lipoprotein particle
- **GO:0050919 negative chemotaxis**
- GO:0007187 G protein-coupled receptor signaling pathway coupled to cyclic nucleotide second messenger
- GO:0032809 neuronal cell body membrane

Female | Male

E Top GSEA enrichments for aging sex differences from H3K4me3 (GO)

- GO:0005691 U6atac snRNP
- **GO:0006910 phagocytosis, recognition**
- GO:0080008 Cul4-RING E3 ubiquitin ligase complex
- GO:0008401 retinoic acid 4-hydroxylase activity
- GO:0036342 post-anal tail morphogenesis
- **GO:0050766 positive regulation of phagocytosis**
- GO:0033038 bitter taste receptor activity
- **GO:0038187 pattern recognition receptor activity**
- **GO:0008392 arachidonic acid epoxygenase activity**

Female | Male

NES: 3, 2, 1, 0, -1, -2, -3

-log₁₀(FDR): 10, 20, 30

F GSEA of CRISPR-Cas9 phagocytosis regulators

-log₁₀(FDR): 1, 3, 5

- Haney M.S., et al. screen
- Pluvinage J.V., et al. screen
- Combined screen

Female | Male

G GSEA of metabolic genes

-log₁₀(FDR): 5, 15, 20

- Hallmark Glycolysis
- Hallmark Oxidative Phosphorylation

Female | Male

H GSEA of polarization genes

-log₁₀(FDR): 5, 15, 20

NES: 2, 0, -2

- M1 enriched
- M2 enriched

Female | Male

Figure 4

A Experimental scheme

Young (4-5 months) ♀ ♂

Old (20-21 months) ♀ ♂

→ Peritoneal lavage cells → MACS purification → Macrophages →

Phagocytosis
Microscopy
Flow cytometry

Metabolism
Seahorse (BMDM)
Lactate-Glo

Polarization
Flow cytometry

B Representative phagocytosis images

Female / Male

Young / Old

C Phagocytosis quantification (Microscopy)

Relative phagocytosis index (A.U.)

2.0E-2   0.16

Young Old Young Old
Female   Male

D Phagocytosis quantification (Flow cytometry)

Relative phagocytosis index (A.U.)

4.7E-2   0.68

Young Old Young Old
Female   Male

E Candidate phagocytosis regulators specifically regulated in female aging

Female-specific down-regulated genes (FDR < 5%, FDR < 10%)   Phagocytosis regulators

128   9   515

B4galt1   *Gna13*
Dpy19l3   *Hmgcs1*
Man1a2   *Magt1*
Rps6ka3
Tbl1xr1
Tm9sf3

F shRNA mini screen of phagocytosis regulators experimental scheme

Lentiviral shRNA
● sh1
● sh2

RAW264.7 → 1-2 week puromycin selection → Selected RAW264.7 → 24-72hr functional assay → RT-qPCR / Phagocytosis

G Mini shRNA screen for phagocytosis capacity (Flow cytometry)

Relative phagocytosis index (A.U.)

Inconsistent with aging
Consistent with aging

0.88 0.69 0.23 1.6E-2 0.92 0.58 0.31 0.40 1.7E-2

B4galt1 Dpy19l3 Man1a2 Rps6ka3 Tbl1xr1 Tm9sf3 Gna13 Hmgcs1 Magt1

H BMDM Seahorse scheme

Young Old Male Female

Plate cells → Bone marrow cells

Differentiate 7 days +M-CSF

BMDM → Seahorse

I Representative normalized ECAR (Seahorse)

Normalized ECAR (mpH/min/DNA)

Glucose Oligomycin FCCP Rot/AA

Time (minutes)

● Young females (4-5 months)   ● Young males (4-5 months)
● Old females (20-21 months)   ● Old males (20-21 months)

J Glycolysis quantification (Seahorse)

Normalized glycolysis (A.U.)

4.5E-2   0.88

Young Old Young Old
Female   Male

K Glycolysis quantification (Lactate-Glo)

Relative luminescence (A.U.)

1.6E-2   0.46

Young Old Young Old
Female   Male

Figure 5

A OVX experimental scheme

B Multi-dimensional scaling

C GSEA of OVX-regulated genes (FDR < 5%)

D Phagocytosis quantification (Flow cytometry)

E Glycolysis quantification (Lactate-Glo)

F Short term E2 supplementation (1 month) scheme

G Multi-dimensional scaling

H GSEA of E2 supplemented regulated genes (FDR < 5%)

I Phagocytosis quantification (Flow cytometry)

J Glycolysis quantification (Lactate-Glo)

K *Esr1* KO scheme

L Multi-dimensional scaling

M GSEA of *Esr1* knockout regulated genes (FDR < 5%)

N Phagocytosis quantification (Flow cytometry)

O Glycolysis quantification (Lactate-Glo)

Figure 6

A Flowchart for identifying transcription factors (TFs) for screening

Differential expressed genes specifically in **females**

234 total genes

149 ↑
85 ↓

→ Animal TFDB 3.0 →

Transcription factors with **female**-specific age-regulation

TFs

↑ 14
↓ 6

TF motifs enriched at **female**-specific remodeled chromatin region

8 ATAC motifs
5 H3K4me3 motifs

→

Downregulated TFs and co-regulators

Irf2
↓ Mef2c
Meis1
Tal1

↓ Tbl1xr1

B **Female**-specific differential TFs gene expression with aging

Row Z-score

Plag1
Mef2c
Meis1
Jdp2
Tal1
Irf2
Tfec
Nr1h3
Nr2f2
Hoxb3
Hoxb6
Hoxb5
Hoxc6
Hoxc4
Sp110
Zfp697
Tead1
Sox7
Rfx2
Hsf4

● Young females (4-5 months)   ● Young males (4-5 months)
● Old females (20-21 months)   ● Old males (20-21 months)

C Selected enriched motifs at **female**-specific differential ATAC peaks with aging (FDR < 5%)

Irf2
Tal1 (Scl)
Nr2f2 (COUP–TFII)
Tead1
Meis1
Sox7
Mef2c
Tal1 (Scl)

Enrichment
2
1
0
-1

-log$_{10}$(p-val)
● 20
● 40
● 60
● 80

D Selected enriched motifs at **female**-specific differential H3K4me3 peaks with aging (FDR < 5%)

Irf2
Tead1
Meis1
Tal1 (Scl)
Tal1 (Scl)

Enrichment
2
1
0
-1

-log$_{10}$(p-val)
● 10
● 30
● 50

E shRNA mini screen for candidate TF regulators

Lentiviral shRNA
● sh1
● sh2

→ RAW264.7

1-2 week puromycin selection →

Selected RAW264.7

24-72hr functional assay →

→ RT-qPCR

→ Omics
  RNA-seq

→ Functional profiling
  Phagocytosis
  Metabolism

F GSEA of shRNA knockdown compared to **female** aging

shRNA up
shRNA down

shIrf2 *shMef2c* *shMeis1* *shTal1* *shTbl1xr1*

NES
2
1
0
-1
-2

-log$_{10}$(FDR)
● 5
● 10
● 15
● 20

G Phagocytosis quantification (Flow cytometry)

Normalized phagocytosis index to control median (A.U.)

Inconsistent with aging

Consistent with aging

0.99   0.74   4.6E-2   1   0.95

Irf2   *Mef2c*   *Meis1*   *Tal1*   *Tbl1xr1*

H Glycolysis quantification (Seahorse)

Normalized glycolysis to control median (A.U.)

Consistent with aging

1.2E-3   0.38   0.14   1.2E-3   2.7E-2

Inconsistent with aging

Irf2   *Mef2c*   *Meis1*   *Tal1*   *Tbl1xr1*

I Glycolysis quantification (Lactate-Glo)

Normalized lactate secretion (A.U.)

Consistent with aging

7.3E-3   7.7E-2   7.7E-2   2.3E-2   2.7E-2

Inconsistent with aging

Irf2   *Mef2c*   *Meis1*   *Tal1*   *Tbl1xr1*

Figure 7

A Irf2 expression validation scheme

B *Irf2* gene expression (RT-qPCR)

C Irf2 protein expression (Flow cytometry)

D Irf2 protein quantification

E ChIP-seq scheme

F Great enrichment ChIP-seq (Gene ontology)

G Irf2 ChIP-seq targets

H Genome browser snapshot at *Hk3* locus

I *Hk3* gene expression (RT-qPCR)

J Irf2 shRNA knockdown experimental scheme (J774A.1)

K *Hk3* gene expression (RNA-seq)

L Glycolysis quantification (Seahorse)

M Glycolysis quantification (Lactate-Glo)

N Proposed mechanism for female macrophage age-related glycolytic shift

Figure 1

A Experimental scheme

B Peritoneal lavage cell UMAP

- B-cells
- Macrophages
- Monocytes
- NK cells
- T-cells

C UMAP of Augur AUC scores

D Peritoneal cell composition (scRNA-seq)

E Peritoneal cell composition (Flow cytometry)

F Peritoneal lavage immune cell yield (Flow cytometry)