Group-level matching behavior in phototaxis of acoel flatworm *Praesagittifera naikaiensis*

Hiroshi Matsui[1] & Yumi Hata[2]

[1] Graduate School of Human Sciences, The University of Osaka

[2] Institute for Protein Research, The University of Osaka

Author Note

Correspondence concerning this article should be addressed to Hiroshi Matsui.

Department of Human Sciences, Osaka University 1-2, Yamada-oka, Suita-shi, Osaka, Japan, +81-90-8515-5258

We have no known conflict of interest to disclose.

Email: matsui.h1004@gmail.com

1 Abstract

The matching law, which posits that animals allocate their responses in proportion to the rate of

reinforcement, has been supported across diverse animal taxa. Although originally formulated in the

context of operant choice, matching also applies to time allocation in foraging and to Pavlovian

responses, indicating its generality across behavioral domains. However, empirical evidence has

thus far been largely limited to vertebrates and arthropods. Addressing the broader applicability of

this principle requires extending investigations beyond these taxonomic groups, across a wider

phylogenetic spectrum. Here, we examined phototactic behavior in the acoel flatworm

Praesagittifera naikaiensis, a species that acquires nutrients through photosynthesis by symbiotic

algae and exhibits positive phototaxis. Using a custom-built T-maze in which the number of

illuminated LEDs varied across arms, we found that the animals distributed themselves in

proportion to relative brightness, consistent with matching behavior. Moreover, prior exposure to

light for 24 hours attenuated this pattern. This manipulation was intended to induce a state of

nutritional sufficiency, and the resulting decline in phototactic responses suggests that internal

physiological states can modulate even seemingly reflexive locomotor behaviors.

Keywords, matching law, acoelomorpha, phototaxis, choice, ideal free distribution

19 Introduction

All behavior takes time. Thus, animals must constantly allocate their finite time across possible

actions (Baum, 2013). How do animals determine the distribution of their own behavior by

allocating their investment over time?

In addressing this question, the matching law (Herrnstein, 1961) has proven empirically reliable

across a wide range of circumstances and species. Put simply, it states that the rate of responding is

proportional to the rate of reinforcement for each option. Among vertebrates, matching behavior has

been observed across numerous taxa, including pigeons (Herrnstein, 1961), rats (Belke & Belliveau,

2001), humans (Kollins et al., 1997), zebrafish (Kuroda et al., 2021), horses (Dougherty & Lewis,

1992), coyotes (Gilbert-Norton et al., 2009), and possums (Bron, 2003). Although most studies

have focused on vertebrates, matching behavior has also been reported in invertebrates, for

example, in place learning of fruit flies (Zars & Zars, 2009), avoidance learning of cockroaches

(Longo, 1964), and discrete-trial choice in honeybees (Couvillon & Bitterman, 1985). The natural

foraging behavior of bees has been theoretically shown to conform to the matching law, and to

constitute an optimal strategy under certain ecological conditions (Thuijsman et al., 1995). While

matching behavior has been primarily examined in operant choice settings, it also holds under non-

contingent conditions, such as concurrent variable-time (VT) schedules (DeCarlo, 1985). Similarly,

matching under VT schedules has been observed in carpenter ants (DeCarlo & Abramson, 1989).

Moreover, in Pavlovian conditioning, strength of conditioned response has been shown to match the

relative amounts of unconditioned stimulus presentations between two conditioned stimuli (Harris

& Carpenter, 2011).

The observation that matching occurs across a wide range of behavioral domains suggests that it

may represent a 'default pattern' of action selection in animals. Indeed, Gallistel et al. (2007)

demonstrated that even naïve mice exhibited matching in foraging behavior, independent of prior

training. In *Caenorhabditis elegans*, a member of the phylum Nematoda, solitary strains are known

to distribute themselves across food patches in proportion to bacterial density, effectively

maintaining a ratio-based foraging pattern (Gloria-Soria & Azevedo, 2008). Haley and Chalasani

(2024) have argued that this distribution is consistent with the matching law, albeit not exactly the

same. Notably, the behavior in *C. elegans* appears to be elicited through chemosensory processing

and driven by the salience of environmental stimuli, rather than being the result of operant

reinforcement. This suggests that matching can emerge even in stimulus-driven behaviors.

Likewise, evidence for the universality of learning has been accumulating across even more

diverse taxa (Loy et al., 2021; Perry et al., 2013). Habituation has been observed across cnidarians,

protostomes, and deuterostomes. Associative learning, particularly classical conditioning, has been

widely documented in bilaterians. The origin of associative learning remains debated. Because early

learning experiments in cnidarians have been criticized for insufficient experimental control, this

has led to the view that they possess only non-associative learning (Cheng, 2021). However, recent

studies employing more rigorous procedures have demonstrated associative learning in sea

anemones (Botton-Amiot et al., 2023) and box jellyfish (Bielecki et al., 2023). It has also been

documented in brittle stars, echinoderms with poorly centralized nervous systems, after 10 months

of extended training (Notar et al., 2023). Together, these studies suggest that basic learning abilities

were acquired early in animal evolution, and raise a fundamental question: through which

evolutionary changes in nervous systems, body plans, or behavioral repertoires did this principle

emerge? Recent perspectives redefine animal cognition as a basal capacity that emerged once life

developed mechanisms for extracting environmental information (Lyon & Cheng, 2023).

Investigating matching behavior across a broader taxonomic spectrum may illuminate the

evolution of nervous systems and behavior. To evaluate the generality of matching as a behavioral

principle, it is critical to examine whether similar patterns emerge in organisms typically not used as

animal models with diverse body plans and nervous systems.

Acoelomorphs provide a unique opportunity to test the generality of laws derived from behavioral

psychology, particularly in the context of early bilaterian evolution. These animals possess a

bilaterally symmetrical, worm-like body plan, yet they lack a through-gut or anus, reflecting a

structurally simple organization. Notably, the degree of neural centralization varies across species,

ranging from forms with little centralization to those possessing a 'commissural brain' structure at

the anterior pole (Gavilán et al., 2016; Perea-Atienza et al., 2015). Despite acoelomorphs'

anatomical simplicity, genomic analyses have revealed the near-complete presence of major

bilaterian signaling pathways, including Wnt, TGF-β, Notch, Hedgehog, and FGF. This suggests

that simplicity in morphology does not equate to genetic primitiveness (Schiffer et al., 2024). These

features indicate that, although acoelomorphs retain traits shared with basal metazoans—Porifera,

Placozoa, Cnidaria, and Ctenophora—their body plan has undergone a profound shift from radial to

bilateral organization. Consequently, acoelomorphs provide a valuable opportunity to identify the

behavioral repertoires that emerged with the evolution of a bilaterian body plan and its associated

nervous system.

The growing interest in acoelomorphs was initially driven by debates surrounding their

phylogenetic position. While they were once thought to be closely related to flatworms, molecular

phylogenetic analyses have revealed that they are more distantly related, leading to the

establishment of a new phylum (Baguñà & Riutort, 2004). The phylogenetic placement of

acoelomorphs remains contentious. Some studies suggest that they represent a basal lineage of all

bilaterians (Cannon et al., 2016; Hejnol et al., 2009; Ryan, 2013), whereas others argue that they are

a derived clade within Deuterostomia, forming a sister group to Xenoturbellida (Mertes et al., 2019;

Philippe et al., 2011, 2019). In either case, given their unique nervous system organization and body

plan, examining behavioral complexity in acoelomorphs may offer unparalleled insights into the

generality of behavioral principles across evolutionary lineages.

Praesagittifera naikaiensis, a species of acoel flatworm, is endemic to the Seto Inland Sea of

Japan (Yamasu, 1982). It inhabits shallow marine environments, ranging from intertidal zones to

depths of approximately one meter, and can be readily collected in the field (Hikosaka-Katayama et

al., 2020). Owing to these advantages, *P. naikaiensis*, together with other species in the family

Convoluta, has attracted attention as a suitable model organism in studies of acoela. Indeed,

numerous research has been conducted on this species, including its neural architecture (Sakagami

et al., 2021), sensory receptor (Sakagami et al., 2024), and genome sequence (Arimoto et al., 2019).

P. naikaiensis exhibits positive phototaxis, which is mediated by light detection through its ocelli.

Because it derives nutrients from photosynthetic symbiotic algae, approaching a light source is

likely to facilitate nutrient acquisition through enhanced photosynthesis. However, aspects such as

the quantitative adjustment of behavioral choices and modifications in response to internal states

shaped by prior experience remain largely unexplored in this species.

Flexible adjustment in phototactic behavior has also been reported in *S. roscoffensis*, a species in

the same family as *P. naikaiensis*. In intertidal environments, individuals face the risk of excessive

sunlight exposure when lying on surface substrates, which can cause damage to their symbiotic

algae (Straka & Rittmann, 2018). However, it remained unclear whether *S. roscoffensis* actively

avoids harmful light by inhibiting phototaxis (Nissen et al., 2015, Serôdio et al., 2011). Recently,

Thomas et al. (2024) conducted behavioral experiments in which *S. roscoffensis* was exposed to

excessive light levels. The animals were able to escape by burrowing into the sand, and indeed, they

did so. Accordingly, it is reasonable to expect that *P. naikaiensis* likewise flexibly adjusts its choice

behavior in response to prior light exposure.

In this study, we investigated whether the phototactic behavior of *P. naikaiensis* follows the

matching law with respect to light intensity. Using a custom-built T-maze, we varied the number of

lit LEDs (0 to 3) on each of the two choice arms and examined whether individuals preferentially

moved toward the brighter side. In Experiment 1, the proportion of choices closely tracked the

proportion of lit LEDs, consistent with matching behavior. In Experiment 2, we tested whether this

matching was modulated by prior exposure to light. Individuals were continuously exposed to LED

illumination for 24 hours prior to the experiment, a treatment intended to induce a 'satiated 'state,

given that *P. naikaiensis* obtains nutrients through the photosynthesis of its symbiotic algae. By

examining whether such exposure altered choice behavior, we assessed whether internal states—

such as nutritional status or algal activity—influence phototactic decision-making. These two

experiments allowed us to test whether phototactic responses in *P. naikaiensis* follow the

quantitative pattern predicted by the matching law, and whether this behavior is modulated by prior

exposure to light.

Materials and Method

Subjects and housing: Acoel flatworms, *P. naikaiensis*, were used (Figure 1a). They were collected

from public beaches in Hiroshima Prefecture (winter 2024) and Okayama Prefecture (fall 2024),

Japan, according to the collecting spots described in Hikosaka-Katayama et al. (2020)[1]. The rearing

system was based on the method described by Hikosaka (2015). The animals were group-housed in

an acrylic aquarium (25 cm W × 20 cm D × 20 cm H), initially filled with seawater from the

collection sites and gradually replaced with artificial seawater (Delphis, Japan). The salinity was

maintained at 30–35‰, based on that of the natural habitat. The algal culture medium KW21 (ISC

Inc., Japan) was diluted at a ratio of 1:4000 and added. The bottom of the aquarium was covered

with coral sand. The water temperature was kept at 20–24°C using an air conditioner in the room.

The light cycle was controlled at 12L:12D. Illuminance ranged from approximately 2,000 to 10,000

lx, depending on the aquarium location. No additional food was provided, and the animals were

acclimated for at least 7 days prior to experimentation.

Apparatus: a custom-built T-maze was built using a 3D printer (Figure 1b). The maze was printed

with translucent PETG filament to facilitate phototaxis behavior, and it was filled with artificial

seawater during experiments. The initial arm was 10 cm long and 3 cm wide, leading to two

branching choice arms, each 5 cm in length. The maze was 3 cm deep and was filled with artificial

seawater to a depth of approximately 2.5 cm. At the end of each choice arm, three white bullet-

shaped LEDs (6500-7000K) were mounted using utility wax, and they were arranged adjacently

with a 2 mm spacing between each unit. The LEDs were controlled via an Arduino microcontroller

(Arduino LLC, USA). The apparatus is freely available and may be modified or reused without

restriction.

Design and Procedure: two experiments were conducted in a dark room during the light phase of

the light–dark cycle. Each session began by gently placing individuals of *P. naikaiensis* into the

initial arm of a T-maze using a 1 mL dropper to avoid injury. The number of individuals per session

---

[1] Although we could not evaluate site effects, *P. naikaiensis* exhibits gene flow among regions, making population genetic differentiation unlikely (Hikosaka-Katayama et al., 2020).

ranged from 41 to 137, depending on how many could be collected with the dropper (see

Supplementary Table). Several LEDs were illuminated on each side of the maze, and *P. naikaiensis*

were expected to move within the maze according to the relative light intensity on the left and right

arms. Each session was terminated 1 hour after the last individual had been placed in the maze. At

the end of the session, the final positions of all individuals were recorded using a smartphone

camera. Each session in both experiments was conducted with a different set of individuals.

The x–y coordinates of *P. naikaiensis* were manually digitized using ImageJ (NIH, USA).

Coordinates were obtained based on the final positions of individuals. Each individual was

categorized into one of three groups: left choice, right choice, or no-choice. A left or right choice

was defined as movement beyond the width of the initial arm (3 cm), measured from the central

axis, into either side of the choice zone. Individuals that did not cross this threshold were classified

as no-choice. Choice rates were calculated after excluding no-choice individuals. If matching

behavior is present, the proportion of left and right choices should correspond to the relative number

of lit LEDs on each arm. All statistical analyses and data visualizations were conducted using R

v.4.1.0.

Experiment 1:

P. naikaiensis individuals that had gathered at the bottom of the aquarium were gently collected

using a 1 mL dropper and immediately transferred to the experimental room for testing.

LED lighting conditions included all 16 possible combinations of illumination on the left and right

choice arms, ranging from (0,0) to (3,3), representing the number of LEDs on the left and right

arms, respectively. The (0, 0) condition was excluded from the analysis of choice rates (Fig. 2a)

because no choice is required when both arms are unlit. Experiments were conducted 8–10 h into

the light cycle. This window was chosen based on unpublished preliminary observations indicating

that *P. naikaiensis* exhibits its typical phototactic behavior during this period. Experimental

sessions for each condition were conducted on different days.

Experiment 2:

Prior to the experiment, individuals of *P. naikaiensis* were continuously exposed to white LED light

(LTC-LC08U-KN 06-0910, OHM, Japan) for 24 hours. This treatment was intended to induce a

satiated state, as the animals primarily obtain nutrients through photosynthesis by their symbiotic

algae. The satiation procedure was conducted in a 200 mL glass cup covered with transparent

plastic wrap to prevent salinity increase due to evaporation. The light was positioned approximately

10 cm directly above the cup. Illuminance was 10,000–12,000 lx, approximately equivalent to the

maximum brightness within the aquarium.

  A subset of the LED lighting conditions used in Experiment 1 was employed in this experiment.

Specifically, eight combinations were selected to include all possible ratios of lit LEDs between the

left and right arms: (2,3), (3,2), (3,1), (1,3), (1,2), (2,1), (0,3), and (3,0). These combinations were

selected based on the results of Experiment 1, which indicated that they were sufficient to observe

matching behavior based on visual inspection.

Transparency and Openness Statement

The code and data are available upon request to the authors. The 3D-printable file of the T-maze is

shared on GitHub (https://github.com/HeathRossie/stlfiles). The sample size was not pre-

determined and fluctuated due to the procedure of collecting subjects; however, the influence of

varying sample size was assessed (see Results). The analysis plan and study design were not

preregistered.

Results

Experiment 1: Group-level matching in phototactic locomotion

The final choice proportions closely matched the proportion of lit LEDs, indicating matching

behavior (Figure 2a). The Pearson correlation between the proportion of lit LEDs and the choice

rates was significant (t(14) = 9.74, $p < .001$), with an explained variance of $R^2 = .87$, indicating the

strong linear relationship between choice distribution and light intensity.

To further examine the determinants of choice behavior, we conducted a multiple linear regression

with the proportion of right-arm choices as the dependent variable. The independent variables

included the proportion of lit LEDs in the right arm, the total number of lit LEDs, and their

interaction term. The model revealed a significant interaction (F(1,12) = 39.48, $p < .001$), with a

positive regression coefficient for the interaction term ($\beta = 0.22$). This indicates that the degree of

matching increased as the total number of lit LEDs increased. In other words, matching behavior

approached a more complete form under high-luminance conditions, whereas slight under-matching

was observed when fewer LEDs were lit (Figure 2a).

To evaluate whether stimulus intensity affected no-choice behavior, we performed a regression

analysis to examine whether the total number of lit LEDs predicted the proportion of no-choice

responses (Figure 2b). The rationale for this analysis was that a lower number of lit LEDs might

have failed to provide sufficient control to elicit phototaxis. However, the analysis revealed no

significant relationship between the total number of lit LEDs and the proportion of no-choice

responses (F(1,14) = 3.72, $p = 0.07$), suggesting that no-choice behavior was not simply due to

insufficient light intensity to elicit phototaxis.

Because the number of subjects varied widely (41 to 99), we examined whether it affected

phototactic choice. The correlation between the number of subjects and the no-choice rate was R

= .03 (t(14) = .10, $p = .92$), indicating that group size did not affect choice behavior.

Experiment 2: Prior light exposure reduces matching behavior

The matching behavior observed in *P. naikaiensis* arises from phototactic movement. However,

does this behavior reflect the behavioral history of being exposed to light? To address this question,

we tested whether pre-exposure to light for 24 hours would attenuate the matching response.

As predicted, matching behavior was attenuated by light pre-exposure, resulting in an under-

matching pattern (Figure 3a). While the choice rate remained positively correlated with the

proportion of lit LEDs, the strength of this relationship was weaker than in Experiment 1. Indeed,

the linear regression revealed that the correlation coefficient was not statistically significant ($t(6) =$

$2.150, p = .075$). The explained variance was $R^2 = .44$ indicating a reduced correlation compared to

the Experiment 1($R^2 = .87$).

As in Experiment 1, the no-choice rate was not related to the total number of lit LEDs (Figure 3b).

This suggests again that a lower number of LEDs was sufficient to elicit phototaxis. However, the

overall no-choice rate was higher than in Experiment 1 ($t(11.04) = 5.22, p < .001$). This indicates

that light pre-exposure reduced the animals 'sensitivity to the stimulus, leading to a general decline

in phototactic behavior.

As in Experiment 1, the number of subjects differed across conditions (77-137). The correlation

between the number of subjects and the no-choice rate was again not significant ($R = -.33; t(6)$

$= .87, p = .42$).

Discussion

Taxis has traditionally been regarded as a reflexive and mechanistic form of behavior, typically

characterized by rigid control with little capacity for adjustment (Loeb, 1900; Pisula & Pisula,

2024). In this study, we aimed to test whether phototactic locomotion in *P. naikaiensis* results in

matching behavior, in which the distribution of choices is proportional to stimulus intensity.

Furthermore, we examined whether this behavioral pattern could be modulated by animals' prior

exposure to light.

The distribution of choices was clearly proportional to the relative number of lit LEDs, indicating

the presence of matching behavior. The degree of matching became more complete as the total

number of lit LEDs increased, whereas a slight under-matching pattern was observed under

conditions with fewer lit LEDs. This trend may reflect increased discriminability under high-

luminance conditions, in contrast to what would be predicted by the Weber–Fechner law, which

claims that the discriminability is proportional to the ratio of stimulus strengths. Importantly, this

pattern was not attributable to a failure in light detection under low-luminance conditions, as the

total number of lit LEDs did not affect the rate of no-choice responses. In sum, *P. naikaiensis*

exhibited matching behavior in its phototactic choices under the T-maze condition. This pattern was

modulated by prior light exposure, which resulted in under-matching and increased no-choice

behavior. These findings suggest that phototaxis in *P. naikaiensis* is not a rigid reflex, but rather a

flexible behavior influenced by internal physiological states.

The ecological significance of matching behavior is typically discussed in relation to optimality in

behavioral strategies. In operant choice, formal models have derived the matching law from the

optimization of feedback functions linking reinforcement rates to response rates, and from its

connection to optimal foraging theory (Baum, 1981; Staddon, 2016). Empirically, matching has

been observed as a near-optimal behavioral pattern in pigeons (Hinson & Staddon, 1983), monkeys

(Iigaya et al., 2019), and human visual foraging (Kobayashi et al., 2024a, 2024b). In our study, *P.*

naikaiensis exhibited choice patterns that were consistent with the matching law. Despite their

phylogenetic uniqueness, their behavioral allocation followed a lawful distribution similar to those

observed in more extensively studied taxa.

Matching behavior can deviate depending on environmental conditions or internal states. For

example, uncertainty regarding food availability can lead to over-matching in pigeons (Anselme et

al., 2022). In humans and mice, under-matching has been interpreted as a consequence of

"compressed" decision-making policies, which serve to reduce cognitive costs (Bari & Gershman,

2023). Theoretically, deviations from perfect matching have been explained in terms of over- or

underestimation of reinforcer utility (Baum, 1974; Kubanek, 2024). In this context, the under-

matching observed in *P. naikaiensis* may also reflect a state-dependent deviation, modulated by pre-

exposure to light and the resulting changes in internal physiological conditions.

Given the less centralized nervous system of Xenacoelomorpha, deviations from matching

behavior are unlikely to be achieved through 'complex' computational processes. Instead,

reductions in matching behavior may be mediated by bodily mechanisms. In *S. roscoffensis*, a

species in the same family as *P. naikaiensis*, symbiotic algae are not located within host cells but

are embedded in the intercellular spaces of subepidermal tissues (Arboleda et al., 2018; Bailly et al.,

2014). Although the precise routes of nutrient transfer remain unknown, it is presumed that the

algae release photosynthetically derived amino acids and products into these intercellular spaces.

Phototaxis in *P. naikaiensis* is likely driven by a pair of rhabdomeric eyes, where photoreceptor

cells are concentrated and presumably connected to motor circuits that produce directed movement

toward light. The phototactic behavior shown in our study suggests an interaction between the

metabolic system supporting nutrient uptake via algal photosynthesis and the sensory-motor system.

Understanding how such integration is achieved in the nervous system of acoelomorphs is a key

issue for elucidating the evolution of behavior in early bilaterians.

Our study suggests that taxis is not merely a mechanical reflex. As noted in the Introduction, the

closely related *S. roscoffensis* can avoid damaging levels of light by burrowing (Thomas et al.,

2024). Consistent with that observation, we found that prior light exposure attenuated phototactic

movement toward the light source. Taken together, these findings warrant renewed attention to the

flexible modulation of taxis by environmental conditions and behavioral history.

It is important to acknowledge a limitation of the present study. Standard formulations of the

matching law are typically applied at the level of individual behavior, where an organism's choice

distribution is shown to vary proportionally with reinforcement ratios across conditions (e.g.,

Herrnstein, 1961). In contrast, our experiments with *P. naikaiensis* revealed matching behavior at

the group level. Extending this line of research to examine the individual-level decision making

within this taxon would be a valuable direction for future studies.

This limitation, however, also points to an intriguing feature: matching behavior appears to

operate across scales. Collective matching behavior is reminiscent of the classical ideal free

distribution (IFD), under which the ratio of individuals among resource patches matches the ratio of

resource availability, yielding equal intake rates for all individuals (Harper, 1982). In practice, the

spatial distributions of many animal groups during foraging typically show undermatching yet

remain qualitatively close to the IFD (reviewed by Kennedy & Gray, 1993). The IFD, however,

presupposes consumptive foraging. In *P. naikaiensis*, sunlight is the primary nutrient source; thus,

aggregation within a patch does not result in feeding competition (apart from occlusion due to body

overlap), and patch "depletion" is independent of the animals' actions (sunlight is not exhausted by

photosynthesis). These features place *P. naikaiensis* outside the canonical equilibria of patch use in

behavioral ecology. The persistence of collective matching behavior may therefore reflect other

factors, such as increased predation risk associated with aggregation. Disentangling the

determinants that produce phenomenologically similar patterns remains an important direction for

future work.

An influential view holds that cognitive function is a basic biological capacity that emerged early

in evolution and extends beyond the animal kingdom to non-neural organisms such as slime molds

and unicellular life (Lyon & Cheng, 2023). Specifically, adaptive functions necessary for an

organism's persistence—perception, thought, memory, and action—are regarded as the minimal

requirements for generating cognitive phenomena and are referred to as "minimal cognition" or

"basal cognition" (van Duijn et al., 2006; Lyon, 2020; Lyon et al., 2021). On this view, cognition is

not, tacitly or as an inviolable assumption, treated as a human-centered concept. Rather, it should be

understood as continuous across living systems, as part of life itself. The background to this

perspective is the recognized need to connect classes of behavioral phenomena long not considered

"cognitive" to cognitive research in a way that is more than metaphorical. For example, even in *E.*

coli, chemotaxis is mediated by a sensorimotor system that switches between running and tumbling

(van Dujin et al., 2006).

This perspective is not confined to the philosophy of biology; it also gives rise to questions of

substantive significance for comparative psychology. For example: if the body plan present when

nervous systems first evolved resembled that of extant basal metazoans, what behavioral

innovations emerged with the acquisition of bilateral symmetry? Ultimately answering this question

requires accumulating comparative behavioral evidence across a broad range of animals. The

modifiability of behavior based on internal states that we have demonstrated suggests that

mechanisms enabling "goal-directed" activity were already in place during the early evolution of

bilateral symmetry. As we hypothesized above, this involves experience-dependent regulation

within metabolic and sensorimotor systems. Interactions among such biological systems also play a

decisive role in producing adaptive behavior in vertebrates (a typical example is feeding behavior;

Ginane et al., 2015). Whether similar mechanisms are implemented in Cnidarians or in non-neural

organisms, which are phylogenetically more distant from us, will be an interesting topic for

investigation. A perspective that integrates biologically informed cognitive research within a

common framework, rather than leaving it as a mere collection of facts, has been growing as a

powerful complement to traditional cognitive research (Barton & Barrett, 2025; Matsui & Hata,

2025). Our behavioral research on acoels likewise follows this line.

Placing our findings in a broader context, our results resonate with early behavioral studies by

Charles Darwin on earthworms (Darwin, 1892) and Herbert Spencer Jennings on protists (Jennings,

1906). Darwin observed that earthworms pulled leaves and small stones into their burrows.

Interestingly, he noted that under warmer conditions, this behavior became sloppier, whereas under

colder conditions, the burrow entrances were covered more carefully, presumably to block cold air.

He interpreted this variation not as a failure of reflex but as evidence that the worms adjusted their

actions according to necessity. Similarly, Jennings described how even unicellular organisms such

as Paramecium and Amoeba exhibited state-dependent modulation of taxis behavior. In line with

these observations, *P. naikaiensis* also exhibited matching behavior that was attenuated following

light pre-exposure, suggesting that internal states can modulate even seemingly reflexive

phototactic behavior. Reed (1982) later highlighted the significance of Darwin's work, arguing that

identifying the external and internal conditions under which behavior is adaptively modulated is a

central agenda for evolutionary psychology.

CRediT Statement

Author 1: Conceptualization, Methodology, Investigation, Software, Data curation, Formal analysis,

Visualization, Writing – Original draft, Writing – Reviewing and Editing, Project administration;

Author 2: Conceptualization, Investigation, Writing - Original draft, Writing – Reviewing and

Editing.

Funding

No funding support was used in this study.

Declaration of generative AI

The authors used ChatGPT o3 for language-proofing to refine the manuscript written by non-native

English speakers. After using this tool/service, the authors reviewed and edited the content as

needed and takes full responsibility for the content of the publication.

Reference

Anselme, P., Wittek, N., Oeksuez, F., & Güntürkün, O. (2022). Overmatching under food

uncertainty in foraging pigeons. *Behavioural Processes*, *201*, 104728.

Arimoto, A., Hikosaka-Katayama, T., Hikosaka, A., Tagawa, K., Inoue, T., Ueki, T., ... & Satoh,

N. (2019). A draft nuclear-genome assembly of the acoel flatworm *Praesagittifera*

naikaiensis. *GigaScience*, *8*, giz023.

Arboleda, E., Hartenstein, V., Martinez, P., Reichert, H., Sen, S., Sprecher, S., & Bailly, X. (2018).

An emerging system to study photosymbiosis, brain regeneration, chronobiology, and behavior:

the marine acoel S*ymsagittifera roscoffensis*. *BioEssays*, *40*, 1800107.

Bailly, X., Laguerre, L., Correc, G., Dupont, S., Kurth, T., Pfannkuchen, A., Entzeroth, R., Probert,

I., Vinogradov, S., Lechauve, C., Garet-Delmas, M., Reichert, H., & Hartenstein, V. (2014). The

chimerical and multifaceted marine acoel *Symsagittifera roscoffensis*: from photosymbiosis to

brain regeneration. *Frontiers in microbiology*, *5*, 498.

Baguñà, J., & Riutort, M. (2004). Molecular phylogeny of the Platyhelminthes. Canadian Journal of

Zoology, 82(2), 168–193.

Bari, B. A., & Gershman, S. J. (2023). Undermatching is a consequence of policy

compression. *Journal of Neuroscience*, *43*, 447-457.

Barron, A. B., & Cheng, K. (2013). Invertebrate learning and cognition: Relating phenomena to

neural substrate. *WIREs: Cognitive Science*, *4*, 561-582.

Barton, R., & Barrett, L. (2025). Embodied cognitive evolution and the limits of convergence.

Philosophical Transactions B, 380, 20240255.

Baum, W. M. (1974). On two types of deviation from the matching law: bias and undermatching.

Journal of the Experimental Analysis of Behavior, 22, 231-242.

Baum, W. M. (1981). Optimization and the matching law as accounts of instrumental

behavior. Journal of the experimental analysis of behavior, 36(3), 387-403.

Baum, W. M. (2013). What counts as behavior? The molar multiscale view. *The Behavior*

Analyst, *36*, 283-293.

Belke, T. W., & Belliveau, J. (2001). The generalized matching law describes choice on concurrent

variable-interval schedules of wheel-running reinforcement. *Journal of the Experimental*

Analysis of Behavior, *75*, 299-310.

Bielecki, J., Nielsen, S. K. D., Nachman, G., & Garm, A. (2023). Associative learning in the box

jellyfish Tripedalia cystophora. *Current Biology*, *33*, 4150-4159.

Botton-Amiot, G., Martinez, P., & Sprecher, S. G. (2023). Associative learning in the cnidarian

Nematostella vectensis. *Proceedings of the National Academy of Sciences*, *120*, e2220685120.

Bron, A., Sumpter, C. E., Foster, T. M., & Temple, W. (2003). Contingency discriminability,

matching, and bias in the concurrent-schedule responding of possums (Trichosurus

vulpecula). *Journal of the Experimental Analysis of Behavior*, *79*, 289-306.

Cannon, J. T., Vellutini, B. C., Smith, J., Ronquist, F., Jondelius, U., & Hejnol, A. (2016).

Xenacoelomorpha is the sister group to Nephrozoa. *Nature*, *530*, 89-93.

Cheng, K. (2021). Learning in Cnidaria: A systematic review. *Learning & Behavior*, *49*, 175-189.

Couvillon, P. A., & Bitterman, M. E. (1985). Analysis of choice in honeybees. *Animal Learning &*

Behavior, *13*, 246-252.

Darwin, C. (1892). *The formation of vegetable mould, through the action of worms, with*

observations on their habits. J. Murray.

DeCarlo, L. T. (1985). Matching and maximizing with variable-time schedules. J*ournal of the*

Experimental Analysis of Behavior, *43*, 75-81.

DeCarlo, L. T., & Abramson, C. I. (1989). Time allocation in carpenter ants (*Componotus*

herculeanus). *Journal of Comparative Psychology*, *103(4)*, 389-400.

Dougherty, D. M., & Lewis, P. (1992). Matching by horses on several concurrent variable-interval

schedules. *Behavioural Processes*, *26*, 69-76.

Gallistel, C. R., King, A. P., Gottlieb, D., Balci, F., Papachristos, E. B., Szalecki, M., & Carbone,

K. S. (2007). Is matching innate?. *Journal of the Experimental Analysis of Behavior*, *87*, 161-

199.

Gavilán, B., Perea-Atienza, E., & Martínez, P. (2016). Xenacoelomorpha: a case of independent

nervous system centralization? *Philosophical Transactions of the Royal Society B: Biological*

Sciences, *371*, 20150039.

Gilbert-Norton, L. B., Shahan, T. A., & Shivik, J. A. (2009). Coyotes (*Canis latrans*) and the

matching law. *Behavioural Processes*, *82*, 178-183.

Ginane, C., Bonnet, M., Baumont, R., & Revell, D. K. (2015). Feeding behaviour in ruminants: a

consequence of interactions between a reward system and the regulation of metabolic

homeostasis. *Animal Production Science*, *55*, 247-260.

Harper, D. G. C. (1982). Competitive foraging in mallards:"Ideal free'ducks. *Animal Behaviour*, *30*,

575-584.

Harris, J. A., & Carpenter, J. S. (2011). Response rate and reinforcement rate in Pavlovian

conditioning. *Journal of Experimental Psychology: Animal Behavior Processes*, *37*, 375–384.

Hejnol, A. (2015). Acoelomorpha. In Schmidt-Rhaesa, A., Harzsch, S., Purschke, G. (Eds)

Structure and Evolution of Invertebrate Nervous Systems, 56-61.

Hejnol, A., Obst, M., Stamatakis, A., Ott, M., Rouse, G. W., Edgecombe, G. D., Martinez, P.,

Baguñà, J., Bailly, X., Jondelius, U., Wiens, M., Müller, W. E. G., Seaver, E., Wheeler, W.,

Martindale, M. Q., Giribet, G., & Dunn, C. W. (2009). Assessing the root of bilaterian animals

with scalable phylogenomic methods. Proceedings of the Royal Society B: Biological Sciences,

276(1677), 4261-4270.

Herrnstein, R. J. (1961). Relative and absolute strength of response as a function of frequency of

reinforcement. *Journal of the Experimental Analysis of Behavior*, *4,* 267-272.

Hikosaka, A. (2015). Artifi cial Rearing System for *Praesagittifera naikaiensis*

(Acoela, Acoelomorpha) [in Japanese with English abstract]. *Bulletin of the Graduate School of*

Integrated Arts and Sciences, Hiroshima University. I, Studies in Human Sciences, *10*, 17–23.

Hinson, J. M., & Staddon, J. E. R. (1983). Matching, maximizing, and hill-climbing. *Journal of the*

Experimental Analysis of Behavior, *40*, 321-331.

Hikosaka-Katayama, T., Watanuki, N., Niiho, S., & Hikosaka, A. (2020). Geographical distribution

and genetic diversity of *Praesagittifera naikaiensis* (Acoelomorpha) in the Seto Inland Sea,

Japan. *Zoological Science*, *37*, 314-322.

Iigaya, K., Ahmadian, Y., Sugrue, L. P., Corrado, G. S., Loewenstein, Y., Newsome, W. T., & Fusi,

S. (2019). Deviation from the matching law reflects an optimal strategy involving learning over

multiple timescales. *Nature Communications*, *10*, 1466.

Jennings, H. S. (1906). *Behavior of the lower organisms*. Columbia University Press.

Kennedy, M., & Gray, R. D. (1993). Can ecological theory predict the distribution of foraging

animals? A critical analysis of experiments on the ideal free distribution. *Oikos*, *68*, 158-166.

Kobayashi, H., Matsui, H., & Ogawa, H. (2024a). Disrupting optimal decision making in visual

foraging: The impact of search experience. *Journal of Experimental Psychology: Human*

Perception and Performance, *50*, 233–248.

Kobayashi, H., Matsui, H., & Ogawa, H. (2024b). The influence of blocking on feature learning and

optimal decision in the visual foraging task. *Journal of Experimental Psychology: Learning,*

Memory, and Cognition, *50*, 400–417.

Kollins, S. H., Newland, M. C., & Critchfield, T. S. (1997). Human sensitivity to reinforcement in

operant choice: How much do consequences matter? *Psychonomic Bulletin & Review*, *4*, 208-

220.

Kubanek, J. (2024). Matching provides efficient decisions. *bioRxiv*, 2024.02.15.580481

Kuroda, T., Ritchey, C. M., & Podlesnik, C. A. (2021). Zebrafish choice behavior is sensitive to

reinforcer rate, immediacy, and magnitude ratios. *Journal of the Experimental Analysis of*

Behavior, *116*, 182-207.

Loeb, J. (1900). *Comparative physiology of the brain and comparative psychology*. G. P. Putnam's

Sons.

Longo, N. (1964). Probability-learning and habit-reversal in the cockroach. *American Journal of*

Psychology, *77*, 49–51.

Loy, I., Carnero-Sierra, S., Acebes, F., Muñiz-Moreno, J., Muñiz-Diez, C., & Sánchez-González, J.

C. (2021). Where association ends. A review of associative learning in invertebrates, plants and

protista, and a reflection on its limits. *Journal of Experimental Psychology: Animal Learning and*

Cognition, *47*, 234–251.

Lyon, P. (2020). Of what is "minimal cognition" the half-baked version?. *Adaptive Behavior*, *28*,

407-424.

Lyon, P., & Cheng, K. (2023). Basal cognition: shifting the center of gravity (again). *Animal*

Cognition, *26*, 1743-1750.

Lyon, P., Keijzer, F., Arendt, D., & Levin, M. (2021). Reframing cognition: getting down to

biological basics. *Philosophical Transactions of the Royal Society B*, *376*, 20190750.

Matsui, H., & Hata, Y. (2025). On the significance of biogenic approach in comparative cognition.

New Ideas in Psychology, *79*, 101186.

Nissen, M., Shcherbakov, D., Heyer, A., Brümmer, F., & Schill, R. O. (2015). Behaviour of the

plathelminth *Symsagittifera roscoffensis* under different light conditions and the consequences

for the symbiotic algae Tetraselmis convolutae. *The Journal of Experimental Biology*, *218*, 1693-

1698.

Notar, J. C., Go, M. C., & Johnsen, S. (2023). Learning without a brain: classical conditioning in

the ophiuroid Ophiocoma echinata. *Behavioral Ecology and Sociobiology*, *77*, 126.

Perea-Atienza, E., Gavilán, B., Chiodin, M., Abril, J. F., Hoff, K. J., Poustka, A. J., & Martinez, P.

(2015). The nervous system of Xenacoelomorpha: a genomic perspective. *Journal of*

Experimental Biology, *218*, 618-628. Perry, C. J.,

Philippe, H., Brinkmann, H., Copley, R. R., Moroz, L. L., Nakano, H., Poustka, A. J., Wallberg, A.,

Peterson, K.J., & Telford, M. J. (2011). Acoelomorph flatworms are deuterostomes related to

Xenoturbella. Nature, 470(7333), 255-258.

Philippe, H., Poustka, A. J., Chiodin, M., Hoff, K. J., Dessimoz, C., Tomiczek, B., Schiffer, P. H.,

Müller, S., Domman, D., Horn, M., Kuhl, H., Timmermann, B., Satoh, N., Hikosaka-Katayama,

T., Nakano, H., Rowe, M. L., Elphick, M. R., Thomas-Chollier, M., Hankeln, T., Mertes, F.,

Wallberg, A., Rast, J. P., Copley, R. R., Martinez, P., & Telford, M. J. (2019). Mitigating

anticipated effects of systematic errors supports sister-group relationship between

Xenacoelomorpha and Ambulacraria. *Current Biology*, *29*, 1818–1826.

Pisula, E., & Pisula, W. (2024). Back to Actual Behavior–A Modest Proposal on the Example of

Exploratory Behavior in Children on the Autism Spectrum. *Integrative Psychological and*

Behavioral Science, *58*, 693-710.

Reed, E. S. (1982). Darwin's earthworms: A case study in evolutionary

psychology. *Behaviorism*, *10*, 165-185.

Ryan, J. F., Pang, K., Schnitzler, C. E., Nguyen, A. D., Moreland, R. T., Simmons, D. K., Koch, B.

J., Francis, W. R., Havlak, P., NISC Comparative Sequencing Program, Smith, S., Putnam, N.,

Haddock, S. H. D., Dunn, C. W., Wolfsberg, T. G., Mullikin, J. C., Marindale, M. Q., &

Baxevanis, A. D. (2013). The genome of the ctenophore Mnemiopsis leidyi and its implications

for cell type evolution. *Science*, *342*, 1242592.

Sakagami, T., Watanabe, K., Hamada, M., Sakamoto, T., Hatabu, T., & Ando, M. (2024). Structure

of putative epidermal sensory receptors in an acoel flatworm, *Praesagittifera naikaiensis*. *Cell*

and Tissue Research, *395*, 299-311.

Sakagami, T., Watanabe, K., Ikeda, R., & Ando, M. (2021). Structural analysis of the statocyst and

nervous system of *Praesagittifera naikaiensis*, an acoel flatworm, during development after

hatching. *Zoomorphology*, *140*, 183-192.

Schiffer, P. H., Natsidis, P., Leite, D. J., Robertson, H. E., Lapraz, F., Marlétaz, F., Fromm, B.,

Baudry, L., Simpson, F., Høye, E., Zakrzewski, A. C., Kapli, P., Hoff, K. J., Müller, S.,

Marbouty, M., Marlow, H., Copley, R. R., Koszul, R., Starkies, P., & Telford, M. J. (2024).

Insights into early animal evolution from the genome of the xenacoelomorph worm *Xenoturbella*

bocki. *Elife*, *13*, e94948.

Serôdio, J., Silva, R., Ezequiel, J., & Calado, R. (2011). Photobiology of the symbiotic acoel

flatworm *Symsagittifera roscoffensis*: algal symbiont photoacclimation and host photobehaviour.

Journal of the Marine Biological Association of the United Kingdom, *91*, 163-171.

Staddon, J. E. R. (2016). *Adaptive Behavior and Learning*. Cambridge University Press.

Thomas, N. J., Tang, K. W., & Coates, C. J. (2024). To move or not to move: taxis responses of the

marine acoel *Symsagittifera roscoffensis* to different stimuli. *Marine and Freshwater Behaviour*

and Physiology, *57*, 17-31.

Thuijsman, F., Peleg, B., Amitai, M., & Shmida, A. (1995). Automata, matching and foraging

behavior of bees. *Journal of Theoretical Biology*, *175*, 305-316.

van Duijn, M., Keijzer, F., & Franken, D. (2006). Principles of minimal cognition: Casting

cognition as sensorimotor coordination. *Adaptive Behavior*, *14*, 157-170.

Yamasu, T. (1982). Five new species of acoel flat worms from Japan. Galaxea, 1, 29-43.

Zars, M., & Zars, T. (2009). Rapid matching in Drosophila place learning. *Naturwissenschaften*, *96*,

927-931.

(a)

(b)

Figure 1: (a) Photographs of acoel flatworms, *Praesagittifera naikaiensis*. Top: a single individual

(×240). Bottom: several individuals attached to the inner wall of the aquarium. (b) Photograph of

the custom-built T-maze apparatus used in the experiments. Individual P. naikaiensis were

introduced into the initial arm at the start of each session.

Figure 2: (a) Choice behavior of *P. naikaiensis* showed a proportional relationship with the relative number of lit LEDs on each arm, indicating matching behavior. (b) The total number of lit LEDs had no significant effect on the rate of no-choice responses.

(a)

(b)

Figure 3: (a) Choice behavior deviated from strict matching and exhibited under-matching

following 24-hour light pre-exposure, suggesting a modulatory effect of internal state. (b) The rate

of no-choice responses increased after light pre-exposure.