# Bacterial species and surface structures shape gene transfer and the transcriptional landscape during early conjugation

Galain C Williams[1], Jillian C Danne[2], Simon Crawford[2], Jihane Homman-Ludiye[3], Xenia Kostoulias[4], Bailey Heron[1], Tasnia Islam[1], Melvin Yong[5], Yunn-Hwen Gan[5], Yogitha N Srikhanta[1,6] and Dena Lyras[1,6†]

[1]Infection Program, Monash Biomedicine Discovery Institute and Department of Microbiology, Monash University, Clayton, Victoria Australia, 3800.

[2]Monash Ramaciotti Centre for Cryo-Electron Microscopy, A Node of Microscopy Australia, Monash University, Clayton, Victoria, Australia, 3800.

[3]Monash Micro Imaging, Monash University, Clayton, Victoria, Australia, 3800.

[4]Department of Infectious Diseases, The Alfred Hospital and School of Translational Medicine, Monash University, VIC, Australia, 3004

[5]Infectious Diseases Translational Research Program, Department of Biochemistry, Yong Loo Lin School of Medicine, National University of Singapore, Singapore, Singapore, 117596.

[6]Joint senior author; these authors contributed equally to this article

Corresponding Author: Professor Dena Lyras

**Email:** Dena.Lyras@monash.edu

**Author Contributions:** G.W. co-designed experiments, performed the majority of the experimental work and co-wrote the manuscript. J.D. and S.C. performed the TEM and assisted with SEM imaging. J.H. assisted with fluorescent microscopy imaging and analysis. X.K. assisted with RNA sequencing. B. H. assisted with fluorescent microscopy. T. I. assisted with conjugation time course experiments. M.Y. and Y.H.G assisted with the *K. pneumoniae* conjugation experiments and experimental design. Y.N.S. co-wrote manuscript & co-designed the majority of experiments. D.L. co-designed all experiments and co-wrote the manuscript.

**Competing Interest Statement:** No Competing Interests

**Keywords:** Bacterial Conjugation; Plasmid Transfer; Antibiotic Resistance; Broad host range plasmids; mobile genetic elements

**Abstract**

Conjugative plasmids drive bacterial evolution and niche adaptation. Their carriage often corresponds with changes to the host transcriptome, but it is unknown during which stage of the conjugation process these transcriptional changes are initiated, and how gene transfer is modulated as they occur. This is because they are normally studied after their full acquisition by conjugation has already occurred, not during the process. Here, active RP4 conjugation revealed that the transcriptional response to plasmid acquisition is almost immediate and is host and surface factor dependent. Responses included activation of non-SOS stress pathways, motility, exopolysaccharide production, anaerobic respiration, and metabolic adaptation. These responses were not a barrier to conjugation, indicating that their significance relates to host homeostasis rather than gene transfer. The recipient could prevent these responses by blocking conjugation through capsule expression. RP4 expression was also inhibited by recipient capsule, since capsule physically blocked donor-recipient contact. Unexpectedly, within a population, RP4 was found to exploit single cell variation in recipient capsule thickness and to successfully conjugate with recipient bacteria with reduced capsule. These results show a novel interplay between plasmid and surface architecture across diverse species that controls the conjugation transcriptional landscape, with important ramifications for plasmid dissemination and evolution.

**Main Text**

**Introduction**

Plasmid dissemination by bacterial conjugation contributes to rapid change in bacteria, exemplified by the continuing emergence of highly virulent antibiotic resistant bacterial pathogens (1). During conjugation plasmid DNA is translocated into an apparently passive recipient cell from an active donor cell via a type 4 secretion system (T4SS) (2). Plasmid carriage can have a profound impact on the transcriptional profile of both host and plasmid (3), bacteria isolated following plasmid acquisition often exhibit altered expression of metabolic, respiration, transport, virulence, and other pathways (4-7). Apart from the SOS response which is induced by ssDNA entry into the recipient (8), it is unclear what broader

transcriptional changes occur within the recipient during the active conjugation process as
plasmid-host responses are typically studied 20-40 generations after the plasmid was
acquired. Similarly, whether these responses act as barriers or facilitators of conjugation
remains unknown.

Indeed, while no recipient factors have yet been identified as truly essential for conjugation
(9, 10), recent studies have highlighted an increasingly important role for the recipient in
conjugation. The composition of the recipient plasma membrane can impact conjugation
frequencies (11), as can the presence of capsules (12) and cognate accessory receptors
(13). Recipient bacteria respond to conjugation through the expression of SOS pathways,
and consequently plasmids have evolved suppressors of the recipient SOS response (14,
15). These suppressors are likely to protect recipient bacteria from deleterious SOS-
enhanced recombination (16). Although a picture of an indirect role for the recipient during
the conjugation process is beginning to emerge, a deeper understanding of how recipient
bacteria respond to, and govern, foreign DNA acquisition is needed because this process
directly facilitates the evolution of traits problematic for human, animal and environmental
health.

In this study, we examined how recipient bacteria respond to and modulate the active
conjugation of plasmid RP4 using a transcriptomic and imaging-based approach.
Transcriptional analysis of a recipient that was proficient for RP4 conjugation (conjugation-
permissive), *E. coli*, revealed that conjugation greatly alters the recipient transcriptional
profile almost immediately upon exposure to the donor. This response included the
expression of mixed stress adaptations, anerobic respiratory pathways, motility, colanic
acid production, and metabolic pathways. However, this pronounced response did not
seem to ultimately impact RP4 acquisition, suggesting it is either not a barrier to
conjugation or that RP4 encodes mechanisms to overcome it. We also observed that RP4
gene expression was globally activated during *E. coli* conjugation, which resembled past
transcriptional studies of RP4 during *E. coli* conjugation (17).

When we repeated these experiments with *K. pneumoniae,* a poor recipient due to its
capsule (non-conjugation-permissive), we observed only a small transcriptional response,
and no activation of RP4 genes within the donor. Conversion of *K. pneumoniae* to a state

that was proficient for conjugation (conjugation-permissive) by capsule deletion restored RP4 conjugation, leading to a greater recipient response that whilst mostly unique to *K. pneumoniae*, did involve genes shared with *E. coli* that are known to impact conjugation when deleted in the recipient. RP4 gene expression was also activated by the presence of unencapsulated *K. pneumoniae*, with DNA mobilisation genes in particular increased in expression. We resolved the mechanism through which capsule inhibits conjugation using electron and fluorescence microscopy. This analysis revealed that capsule is a physical barrier to donor-recipient outer membrane contact, preventing mating pair formation and RP4 gene activation in the donor. Unexpectedly, we discovered that single cell variation in capsule thickness within *K. pneumoniae* populations gave rise to the small number of transconjugants typically observed in RP4 conjugations with capsulated *K. pneumoniae*. Our findings highlight that recipient species, and their surface structures directly shape both conjugation, and the transcriptional landscape of the recipient and plasmid during actively occurring conjugation.

**Results**

**RP4 conjugation elicits the rapid activation of non-SOS stress, respiratory and metabolic adaptation pathways in a conjugation-permissive recipient**

To assess recipient and plasmid transcriptional activity with a conjugation-permissive recipient, RNA-seq was performed on actively conjugating donor *E. coli* LT101(pVS520) (phylogroup A), carrying an RP4 derivative pVS520, and the genetically distant recipient *E. coli* DLL7555 (phylogroup B2). Based on a conjugation time course, we extracted RNA 15 minutes after donor-recipient mixing, when 15% of the recipient population had received RP4, but conjugation was still actively occurring (Fig 1a). RNA was also extracted from mock-conjugations between the same strains without pVS520, and from the donor LT101(pVS520) alone without a recipient.

To determine how recipient *E. coli* DLL7555 responded to RP4 conjugation, we compared gene expression of DLL7555 between the conjugation and mock conjugation conditions. The many nucleotide differences between DLL7555 and LT101 (>78,000 genetic variations within core genome) enabled effective recovery of strain-specific reads by

alignment against a concatenated LT101-DLL7555 reference sequence in which reads

aligning to both strains were discarded. We identified 130 differentially expressed genes,

88 upregulated and 42 downregulated (Fig 1b). These involved genes associated with

adaption and aversion to external stressors, which we have termed a mixed-stress

response, including temperature shock (*cspG*, *hslJ, deaD*), pH regulation (*yagU, ygaC,*

*ybay, speF*), osmotic shock (*osmB, osmY, nhaA*) and others (*mliC, soxS, rcsA, phoR*).

This response was paired with increased expression of colanic acid production (*wzc, wzb,*

*cpsB, wcaJ, wcaI, wcaE, wcaC, wcaA, gmd, fcl, wza, wcaD*), exopolyscharide production

(*yjbG, yjbF, yjbH*), and flagella production (*fliN, fliM, flgG, flgF, glgI, flgH, flhA, fliO*) genes,

responses associated with aversion to stressors and biofilm formation. Curiously, despite

an increase in flagella expression, the master flagella transcriptional regulator *flhD* was

decreased in expression, whilst the flagella anti-sigma factor *flgM* was increased. As some

conjugative plasmids induce biofilm formation following acquisition, we predicted that the

genes identified were involved with the induction of a recipient biofilm in the conjugation

mixture. Unexpectedly, however, we determined that DLL7555 co-culture with

LT101(pVS520) had a slight reduction in biofilm formation compared to the LT101 co-

culture, or monoculture (Supplementary Fig 1). This minor reduction may result from the

decreased expression of key biofilm formation factors encoded by *csgD* and *csgG* during

conjugation.

Unexpectedly, conjugation was also associated with increased expression of anaerobic

respiration and energy generation pathways (*fdnI, dmsB, napC, narG, narH, narI, hycF,*

*nrfA, nrfD, nrfC, nirB, fumB, dcuC*) which overlapped with inorganic nutrient metabolism

and  nitrogen cycle metabolic processes (*napG, napH, narG, narH, narI, nrfA, narJ, nirB,*

*nirD, narK, napC*). Curiously, multiple siderophore, iron transport and iron homeostasis

pathways were downregulated, including (*fiu, cirA, fepA, fyuA, efeO, efeU, efeB, fhuE,*

*entD, entC, entF, chuA, chuS, chuT*), though the master iron uptake regulator *fur* was not

differentially expressed.

GO term analysis reflected the multi-factorial recipient response to RP4 conjugation with

terms associated with stressors, respiratory pathways, and metabolic adaption being

significant (Fig 1c). Collectively, our findings suggest that early conjugation is both highly

stressful, and energetically intensive, for the recipient, requiring the activation of
respiratory pathways not typically present during aerobic growth whilst modulating iron
acquisition and homeostasis for an unknown reason. Our findings may explain the prior
observation that nascent RP4 transconjugants are growth impaired (18).

Finally, although our experiment was not designed to resolve donor-recipient specific
differences in RP4 expression, by comparison against the donor alone we reasoned we
could gain insight into the global expression pattern of RP4 across the entire conjugating
population. This comparison revealed increased expression of 47 pVS520-encoded genes
associated with DNA mobilisation (MOB), mating pair formation (MPF), Regulation and
replication, host-kill functions (*kil*), and other processes, a response entirely consistent
with previous reports of RP4 gene expression during *E. coli* conjugation (17) (Fig 1d).

**The recipient conjugation response is diminished by capsule but expressed in its**
**absence.**

To determine how a non-conjugation-permissive recipient reacts to an RP4 donor and
whether making it conjugation-permissive would produce an *E. coli*-like recipient
response, we examined conjugation in strains producing or not producing a capsule.
Capsule was used as the basis for non-permissiveness because it strongly blocks
conjugation but its removal through genetic deletion allows conjugation to proceed with
high efficiency (19). Capsule was also chosen because it differs from other mechanisms
like restriction-modification systems, which do not prevent conjugation, but instead prevent
plasmid establishment.

To identify a suitable capsulated recipient, we compared conjugation frequencies between
*Klebsiella pneumoniae* B5055, *Acinetobacter baumannii* AB5075, and *Pasteurella*
*multocida* VP161, against their respective acapsular derivatives (Supplementary Fig 2).
Whilst capsule hindered conjugation in all three species, only the acapsular *K.*
*pneumoniae* B5055Δ*wzbc* achieved conjugation frequencies comparable to *E. coli*
DLL7555. As a high frequency is essential for effectively resolving the recipient's
transcriptional response, we therefore performed RNA-seq on conjugation mixtures using

donor LT101(pVS520) with *K. pneumoniae* B5055 or B5055Δ*wzbc* recipients as before
with *E. coli* DLL7555.

The non-conjugation-permissive B5055 response to donor co-culture involved 10
differentially expressed genes (*carB, carA, lysS, clbS, glnH,* 08025*, glnP,* 08030, 08050,
08045*)*, with mostly decreased expression (Fig 2a). GO-term analysis identified terms
associated with sucrose catabolism, L-arginine biosynthesis and nucleotide biosynthesis
(Fig 2b). Conversion of B5055 to a conjugation-permissive state greatly bolstered the
response, with 54 B5055Δ*wzbc* genes being differentially expressed (Fig 2a). We
observed Increased expression of mixed stress adaptations that were seen in *E. coli* (*cspD*
and *nhaA*), but also those unique to B5055Δ*wzbc* (*loiP*, *ychH, hcp*). We also observed
increased expression of genes for carbohydrate transport (*araH, fucP, galP, manY, manZ,*
*manX, bglF*) and catabolism (*treC, fruK pck, bglB, opbG*). Broader transmembrane
transport genes (*psuT, sdaC, ompC, ttdT, nhaA, yebQ, malE, gntT, ompW*) were also
differentially expressed, which together suggest that increased substrate efflux across the
cell envelope may occur in B5055Δ*wzbc* during conjugation, along with altered
carbohydrate metabolism. Some of these genes were associated with growth under
anaerobic conditions (*ompW*, *ttdT*) alongside anaerobic respiration genes (*fdnG, fdhF,*
*narG*). GO-term analysis also identified terms for mixed stress responses and for altered
carbohydrate transport and metabolism (Fig 2b).

To determine how many genes were common between DLL7555 and B5055Δ*wzbc*, blast
was used to identify homologues within respective gene lists (Fig 2c). This revealed that
13 differentially expressed genes were shared between DLL7555 and B5055Δ*wzbc,*
indicating that approximately 10% of the permissive-recipient response is shared. These
included *ysab, galP, ypfG, ypeC, ompW, narG, cspG, citT, nhaA,* ACPEGL_23855*, ytjA*
with increased expression and *csgD* and *ompF* with decreased expression. Notably,
previous studies have shown that *nhaA* and *ompW* reduce conjugation when disrupted in
recipient bacteria (10, 13). Finally, modified *lysS* expression was also identified as
common between the B5055 and B5055Δ*wzbc* response. Our findings reveal that
recipient bacteria exhibit a response to conjugation that involves thematically similar

processes and common genes whilst also involving genes and pathways unique to each
recipient species.

**Recipient capsule inhibits donor RP4 transfer gene activation and conjugation by**
**blocking donor and recipient cell contact**

As described here and by others (19) capsules are potent inhibitors of conjugation, yet the
precise mechanism underlying this phenomenon remains only partially described. Our
earlier observations showed that capsule blocked the transcriptional response of the
recipient to conjugation. To determine if this effect extended to blocking the activation of
RP4 conjugation gene expression within the donor, additional analysis of the
transcriptome datasets was performed.

No RP4 genes were differentially expressed during conjugation with B5055 compared to
in the donor alone condition (Fig 2d), whilst 27 were activated during conjugation with
B5055Δ*wzbc* (Fig 2e), suggesting that capsule blocks RP4 gene expression in the donor
cell. Because the expression of RP4 genes in the B5055Δ*wzbc* mating could have
originated within transconjugant cells, a donor-specific fluorescent reporter assay was
performed to examine activity arising from the MOB promoter P*traJ* using rifampicin to
silence recipient transcription. This assay confirmed that MOB gene activation occurred
within the donor and could be blocked by recipient capsule (Supplementary Fig 3). These
experiments indicate that exposure to a conjugation-permissive recipient triggers RP4
gene expression within the donor, and can be inhibited by recipient capsule production.

The analysis described above also revealed that RP4 gene expression differed between
the conjugation-permissive recipients B5055Δ*wzbc* and DLL7555. The RP4 response to
DLL7555 included all genes activated by B5055Δ*wzbc* (Fig 2f), but was also broader, with
activation of the MPF *trb* operon detected (Fig 1d), which was not observed during
conjugation with B5055Δ*wzbc* (Fig 2e). Thus, RP4 gene expression is shaped by both
recipient species and surface structures, such as capsule.

Finally, to gain mechanistic insight into how capsule exerted its anti-conjugation effect,
imaging studies were performed to visualize the impact of *K. pneumoniae* capsule on
donor and recipient cell-to-cell interactions. Transmission and scanning electron

microscopy showed capsulated recipients localized within the donor periphery, but they were physically prevented from direct outer membrane contact with the donor due to a barrier formed by their capsules (Fig 2g, top panels). In contrast, direct donor and recipient outer membrane contact was evident with the unencapsulated B5055Δ*wzb* (Fig 2g, bottom panels). Unexpectedly, the electron microscopy imaging also revealed variation in the thickness and architecture of the B5055 capsule at the single cell level, which is described in detail below (Fig 3).

The reduced cell-to-cell contact caused by capsule appeared to block the conjugation process, which was confirmed *via* fluorescence microscopy of conjugations on agarose pads (Fig 2h). In these experiments, a mobilizable plasmid encoding GFP was used to track conjugation to recipient cells counter-labelled with BFP on a non-mobilizable plasmid. With the unencapsulated B5055Δ*wzbc* recipient, 70% of donor-contacting recipient cells became transconjugants (369 transconjugants from 527 potential mating pairs), indicating that direct cell-to-cell contact was highly productive for conjugation (Fig 2h, left panel). Conversely, with B5055 only 0.58% of donor-contacting recipient cells formed transconjugants (2 transconjugants from 345 potential mating pairs), and mating pairs were observed less frequently because capsule prevented direct donor and recipient contact (Fig 2h, right panel). Notably, B5055 cells also showed limited contact with each other, indicating that capsule reduces both intra- and inter-cellular contact.

Taken together, our transcriptomic and imaging data reveal that recipient capsule blocks conjugation by preventing donor and recipient outer membrane contact, thereby inhibiting mating pair formation and the activation of conjugation genes within the donor.

**RP4 exploits variation in capsule thickness and expression to overcome capsule as a barrier to conjugation**

During the earlier imaging experiments, variation in *K. pneumoniae* capsular thickness at the single cell level was unexpectedly observed, which prompted further characterization of the *K. pneumoniae* capsule by both SEM and TEM (Fig 3).

SEM imaging showed capsule formed a complex, diffuse covering of the cell surface which connected adjacent cells through thick projections (Fig 3a). These projections were

between 0.5 μm and 4 μm in length, and present within 86% of cells (581/676). The majority of *K. pneumoniae* cells exhibited a complex capsular architecture, but occasionally individual cells with a simple and smooth surface were observed. Under TEM, capsule appeared as a diffuse, occasionally non-uniform layer which varied considerably in thickness between individual cells (Fig 3b-c).

Total capsular diameter, i.e. the combined thickness on both sides of the cell, was assessed both within and between biological replicates (Fig 3d). Across replicates, the mean total capsular diameter was 287 nm and ranged from 132.8 nm to 678.8 nm, reflecting considerable variation at the single cell level. Unexpectedly, capsule thickness differed significantly between replicates grown and treated identically, with mean values of 420.2 nm, 192.7 nm, and 263.3 nm (Fig 3d). Overall, these results demonstrate that *K. pneumoniae* has considerable phenotypic capsular heterogeneity within clonal populations.

Further experiments were performed to determine if variation in recipient capsule thickness was the mechanism through which the small number of B5055 transconjugants that normally arise during RP4 conjugation were formed. Since capsule inhibited conjugation by blocking cell-to-cell contact (Fig 2g-h), it was hypothesized that recipients with thin capsules could form tighter contact with donor cells, leading to effective conjugation.

In support of this hypothesis, TEM and SEM visualisation of conjugations revealed that thin-capsulated B5055 recipients formed much closer associations with the donor than did thick-capsulated ones (Fig 3 e-f). Indeed, capsule thickness correlated strongly with the distance between donor and recipient outer membranes (Fig 3g, R2=0.72, N=47, F test P<0.0001), meaning that thicker capsules corresponded to a greater distance between donor and recipient. Importantly, live cell fluorescence microscopy confirmed that B5055 transconjugants arose from recipients which could form intimate direct contact with donor cells (Fig 3h). These B5055 transconjugant cells were arranged in tightly packed clusters that bore a striking morphological resemblance to B5055Δ*wzbc* capsule mutants (Fig 3h, inset), supporting that they were thin-capsulated.

Earlier, significant variation in capsule thickness was shown to occur between replicates
grown exactly the same way (Fig 3d). It was therefore hypothesized that variation in
capsule thickness between replicates would occasionally yield higher conjugation
frequencies for the B5055 recipient. To examine this hypothesis, numerous independent
conjugation experiments were performed, with the expectation that variation in capsule
thickness would vary between replicates, thus influencing conjugation efficiency. Of 19
conjugations, 4 had a mean transfer frequency of $4.2 \times 10^{-2}$ (Fig 4A). This value was ~50-
fold higher than the median value for the combined conjugations, but still 20-fold lower
than the B5055Δ*wzbc* recipient (Fig 4a), indicating that the strains were not completely
free of capsule. The resulting transconjugants were mucoid and string-test positive,
confirming that they were not spontaneous acapsular mutants.

RNA-seq was used to examine RP4 gene expression and recipient capsule synthesis
expression in the high-frequency conjugations compared to the earlier, low-frequency
dataset (See Fig 2). RP4 gene expression in the high frequency conjugations resembled
that observed with unencapsulated B5055Δ*wzbc,* in which genes for DNA mobilisation but
not mating pair formation were increased in expression (Fig 4b), suggesting that this is a
common RP4 expression pattern in conjugation-permissive *K. pneumoniae* matings.

Importantly, there was greatly reduced expression of principle capsule synthesis genes
under the control of P*wzi in* the samples that had a high conjugation frequency (Fig 4c).
Genes from this operon with reduced expression included *wza, wzb, wzc,* J6T75_19265*,*
J6T75_19270, J6T75_19275, *wzy, wzx, orf12, orf13,* and *wcaJ*. No differentially
expressed genes from the minor capsule synthesis operons controlled by P*galF* and
P*manC* were identified. A large number of additional genes contribute to capsule synthesis
in *K. pneumoniae* (20) and 13 of these were differentially expressed including the key
capsule synthesis regulator *kvrA* (21), and a variety of other regulatory and structural
genes, including *hha, ompR, pitA, miaA, arnD, ackA, wabH, rcsF, wabN, wcaG, fis, rfaH*
and *mlaF* (Fig 4d)*.* Many other genes of diverse function were also differentially
expressed.

Lastly, because capsule serotypes have variable capsule thickness and expression, the
impact of this natural capsule variation on conjugation was examined. Capsule caused a

31-fold reduction in conjugation frequency with a hypermucoid K1 recipient, but a 3.3 and
2.2-fold reduction for two other hypermucoid K5 recipients compared to their parental
wildtype, suggesting that differences in capsule between serotypes modulate the effect of
capsule on conjugation at a strain level (Fig 4e).

Collectively, these experiments reveal an unexpected role for dynamic variation in the
production of recipient capsule in determining the expression of RP4 encoded genes and
conjugation frequency. These findings indicate that while recipient surface factors can
greatly modulate the rate of conjugation, RP4 donor cells can exploit natural variation in
these factors to permit the entry of plasmid DNA into otherwise non-permissive recipients.

**Discussion**
We investigated how recipient species and surface structures impact conjugation and the
early transcriptional dynamics of the recipient and plasmid. Transcriptomic analysis of
actively conjugating populations revealed a substantial recipient response in conjugation-
permissive *E. coli* involving stress, motility, colanic acid production, respiratory, and
metabolic pathways, alongside global RP4 gene activation (Fig 1). In contrast, *K.
pneumoniae* exhibited a markedly attenuated response, and notably failed to trigger donor
RP4 gene expression. Capsule disruption restored both RP4 gene activation and
conjugation, leading to a recipient response with partial overlap to that of *E. coli* (Fig 2).
Mechanistically, imaging revealed that capsule prevented conjugation by acting as a
physical barrier to donor-recipient contact (Fig 2gh). Surprisingly, we discovered that
single-cell variation in recipient capsule thickness could be exploited by the donor to
facilitate RP4 entry into otherwise recalcitrant populations (Fig 3), shaping both RP4 gene
expression and conjugation frequency (Fig 4).

An important finding of this work is that active conjugation triggers an immediate reshaping
of the recipient transcriptome to an extent much greater than previously thought. Previous
work had shown minor induction of the SOS response well after RP4 conjugation had
finalized (8), but this was not detectable in two distinct recipients during early conjugation.
Instead, we found that recipients express mixed stress adaptations in response to RP4
conjugation, suggesting that conjugation is detected as a potentially harmful stimulus.

Unexpectedly, we discovered that conjugation triggered the activation of anaerobic
respiration genes in both *E. coli* and *K. pneumoniae*, which were particularly upregulated
by *E. coli*. Both strains also exhibited differing metabolic pathway activation that in *K.*
*pneumoniae* was dominated by carbohydrate metabolism and in *E. coli* involved inorganic
nutrient metabolism. The variable impact of RP4 acquisition on metabolic pathway
adaptation is consistent with prior studies on plasmid metabolic reprogramming which
shows a high degree of dependence on host and plasmid identity (15, 22, 23). These
differences may arise from the unique metabolic programs of each host and reflect
different adaptations to meet increased energetic and stress demands post-conjugation.
364
10% of the recipient response was common between *E. coli* and *K. pneumoniae*, and two
shared genes, *ompW* and *nhaA*, reduce conjugation when deleted in recipient bacteria
(10, 13). OmpW is a porin involved in the transition from aerobic to anaerobic growth (24),
and NhaA plays a central role in maintaining sodium ion, pH, and osmotic homeostasis
(25). OmpW contributes to conjugation by serving as a receptor for mating pair
stabilization proteins, which RP4 notably does not encode. However, it is possible that the
common induction of *ompW* expression across two distinct recipient species might have
favored the evolution of mating pair stabilization systems that target OmpW. The function
of NhaA in conjugation is unknown, but it may protect the integrity of the recipient
membrane following the cell wall shock caused by conjugation as NhaA plays an essential
role in maintaining cellular envelope integrity (26).
376
Aspects of the *E. coli* response involving colanic acid and flagella production resemble the
establishment of a biofilm, which was perhaps not surprising as conjugation can trigger
biofilm formation (27). Unexpectedly, however, co-culture with the donor caused a minor
reduction in biofilm formation compared to the mock donor or monoculture
(Supplementary Fig 1), which was supported by decreased expression of a key biofilm
regulator *csgD*. This result suggests that the mixed-stress response induced by RP4 is
not a biofilm formation signal, but a distinct, and novel, process.
384
Our approach of examining the recipient response as conjugation actively occurred
enabled new insights into flagella expression following plasmid acquisition to be gained.

Plasmid carriage often corresponds with silencing of flagella expression (7, 23), which our
results suggest may occur as a compensatory measure to control flagella biosynthesis
triggered by conjugation itself. This suggestion explains the contradictory expression
pattern observed, whereby increased expression of flagella biosynthesis genes was
paired with decreased expression of the flagellar master activator *flhD*, and increased
expression of the flagella anti-sigma factor *flgM*. This may be caused by RP4 encoded
factors that suppress flagella production, perhaps to reduce movement of the recipient
away from the donor. Alternatively, RP4 may increases host cell motility to overcome
population structures which are barriers to RP4 dissemination (28), which the host
attempts to suppress through altered FlhD and FlgM production.

While conjugation can cause substantial alteration to the recipient transcriptome,
recipients can be shielded from this reprogramming by the production of capsule (Fig 2).
This protective mechanism could be beneficial in environmental contexts where a sudden
change in transcriptional expenditure might be detrimental to the cell, such as in a
mammalian host. Perhaps other surface structures serve a similar role during conjugation;
indeed, it is possible that the increased expression of colanic acid in DLL7555 occurred to
achieve a similar outcome, albeit unsuccessfully.

Although capsules have previously been reported to block conjugation (12, 29), the
mechanism underpinning this effect has not been fully described, nor has the significance
of capsule for the recipient response. Prior genetic evidence had suggested capsule was
too thick for the IncP pilus to cross (19), though the physical effect of capsule on
conjugation was not examined. By imaging conjugations through both electron and
fluorescence microscopy, we show that capsule forms a physical barrier that prevents
essential contact between donor and recipient, that in turn inhibits productive mating pair
formation (Fig 2). Indeed, with an average thickness of 143 nm, capsule was rarely thicker
than the well-characterized length the RP4 pilus, 0.2-1 µm (30-32).

Importantly, this finding was built upon further through the discovery that capsule prevents
RP4 gene activation in the donor (Fig 2, supplementary Fig 3). This suggests that capsule
obscures a component of the recipient outer membrane that triggers RP4 to begin

conjugation. Because RP4 can conjugate across all domains of life (33), and no non-
essential recipient genes are known to be necessary for conjugation, the signal is likely to
be a common membrane lipid that cannot be deleted. This hypothesis aligns well with the
fact that the lumen of conjugative pili is lined with lipids (34), and perhaps changes in pilus
lipid composition following their merger with the recipient outer membrane provide the
signal for DNA transfer.

Although recipient contact can trigger the expression of the pilus in plasmid R388 (35), the
pilus of RP4 is constitutively expressed (36), and we did not identify increased expression
of pilus-related genes during conjugation with unencapsulated *K. pneumoniae*. However,
pilus-related genes were activated with *E. coli*, suggesting that the recipient host
influences the expression of RP4 encoded genes. The basis of this phenotype is unclear
but could be due to the presence of co-resident plasmids within the strain of *K.*
*pneumoniae* used in these studies. The presence of these resident plasmids may also
explain why *K. pneumoniae* exhibited a smaller response to RP4 conjugation, as their
presence might modulate how the host responds to the acquisition of additional plasmids,
as for example *psiB* of the F plasmid which inhibits SOS activation (14).

An unexpected finding of this work was that stochastic variation in recipient capsule
thickness enabled RP4 to overcome capsule as a barrier to conjugation (Fig 3-4). This
was observed both at the single cell level (Fig 3) and at the level of independently grown
bacterial cultures (Fig 4). This finding indicates that dynamic variation in capsule can
greatly alter the entry of genes into recipient bacterial populations. Although these findings
agree with previous reports that artificially swapping capsule serotypes can change
conjugation frequencies (19), they also present a new perspective because they reveal
how natural and dynamic changes in capsule expression can allow RP4 to overcome
structural barriers to conjugation. The impact of single cell heterogeneity on gene transfer
is not yet well described but heterogeneity has been described for donor conjugation gene
expression, showing that MPF expression differs between individual donor cells (37). Our
study adds to this growing understanding of how conjugation functions between individual
cells within a population by showing how recipient single-cell heterogeneity in surface
architecture can shape gene transfer.

The cause of capsular heterogeneity studied here is unknown, but capsular heterogeneity

is common within many bacterial species (38-40). One mechanism through which capsule

variation occurs, phase variation (40), has long been associated with bacterial adaptation

to new environments such as during host colonisation by *Neisseria meningitidis* (41).

However, single cell and population heterogeneity in capsule thickness can occur by

mechanisms other than phase variation, and only recently has the impact of this form of

variation on natural bacterial populations begun to be understood (42). Capsule

expression in *K. pneumoniae* is coordinated by multiple complex regulatory networks

involving more than 70 genes (20, 43), and is mostly not phase variable (20, 44, 45). Our

data suggest that the capsule variation we observe is stochastic, as there was no

discernible pattern associated with its expression. However, our data raise the hypothesis

that environments which alter capsule expression can shift the frequency of conjugation

by modulating recipient susceptibly to conjugative plasmids.

Collectively, our findings provide new insights into how conjugation impacts recipient cell

populations, and in turn how they shape conjugation. RP4 plasmid conjugation rapidly

modifies the transcriptional landscape of the recipient, but its gene expression is also

influenced by recipient identity and, critically, surface structures. Indeed, capsule functions

not only as a modulator of physical contact but also of the early transcriptional response

of the recipient. Furthermore, the unexpected discovery of stochastic capsule variation

within *K. pneumoniae* reveals a novel mechanism by which gene transfer can occur in

otherwise recalcitrant recipients. We determined that the early stages of conjugation are

transcriptionally intensive and stressful for recipients beyond the SOS response, and that

seemingly minor variations in recipient surface structures can have profound

consequences for plasmid establishment within bacterial communities. Our findings have

important implications for the evolution of traits which are widely disseminated by

conjugation, including antibiotic resistance, virulence determinants, and metabolic

pathways.

## Materials and Methods

### Bacterial strains, plasmids, and growth conditions

Table 1 lists the bacterial strains and plasmids used in this work. For routine growth, *E. coli* and *K. pneumoniae* strains were cultured in low-salt LB at 37°C shaking in liquid at 200 rpm or on solid media. Media was supplemented with antibiotics where necessary: Ampicillin (100 μg ml$^{-1}$), rifampicin (150 μg ml$^{-1}$), erythromycin (150 μg/ml), spectinomycin (50 μg/ml) and tetracycline (10 μg ml$^{-1}$).

### Strain and plasmid construction

Prior to use as recipients, wildtype and Δ*wzbc K. pneumoniae*, along with *E. coli* DLL7555, were electroporated with the low copy number vector pWSK29 to provide an ampicillin counter selection maker for conjugation. Fluorescent reporter plasmids were made as follows. *GFPmut3* or *mTagBFP2* were fused to a synthetic *em7* promoter (46) and synthesised as gBlocks (IDT). These were cloned into the XbaI and BamHI site of pMMB67EH or pSU18 respectively through restriction digestion cloning. pWSK29P*traJ*GFPx3 was similarly constructed, except that three translationally coupled copies of *mGreenLantern* were fused to a 367 bp region upstream of the *traJ* start codon. Each *mGreenLantern* subunit was independently codon optimised until each shared < 80% sequence homology.

### Conjugation assays

Overnight liquid cultures of donor and recipient were diluted 1:50 in fresh LB and grown for 4 hours before being washed once in PBS and normalized to an OD$_{600}$ 1.0. 500 μl of donor and recipient culture was passed through a 0.44 μm nitrocellulose membrane filter (Whattman) and incubated on LB agar at 37°C for 15 minutes before extraction in 2 ml of PBS by vigorous vortexing. 15 minutes was chosen as a timepoint to prevent secondary transmission of plasmid RP4, and to minimize growth of the recipient which can skew transfer frequencies. For routine assays, recipient and transconjugant CFU/ml were quantified by plating onto ampicillin, and ampicillin + tetracycline respectively. For *A. baumannii*, transconjugants were quantified on ampicillin + erythromycin, and for *P. multocida* kanamycin + spectinomycin using mobilizable plasmid pAL99S-*oriT* as a surrogate for RP4 transfer. Conjugation frequency was calculated as the transconjugant CFU/ml divided by recipient CFU/ml.

**RNA extraction and sequencing**

Conjugations were fixed after 15 minutes of incubation by vortexing mating filters in RNAprotect (Qiagen). Fixed samples were pelleted at 13,300 RPM for 5 minutes, the RNAprotect decanted and samples stored at -80°C for later processing. RNA was extracted from the DLL7555 samples using an RNeasy RNA extraction kit (Qiagen) as per the manufacturer's instructions with an on-column DNase treatment (Qiagen). Due to its thick capsule, RNA from *K. pneumoniae* was isolated with hot-phenol as previously described (47) and further purified using an RNeasy kit with on column-DNase treatment. RNA from DLL7555-related experiments was sequenced at the MHTP Medical Genomics Facility at the Monash Translational Facility. Samples were quality checked by Bioanalyzer (Agilent), and libraries generated using the Illumina stranded total RNA-seq protocol with ribosomal RNA depletion performed using the bacterial Ribo-Zero rRNA removal kit (Illumina). Sequencing was performed on an Illumina NextSeq550 on high output mode with chemistry version 2.5, using the Illumina protocol 15046563 version 2. RNA from *K. pneumoniae* related experiments was sequenced at Micromon Genomics. Samples were quality checked by Fragment Analyzer (AATI) and RNA library construction performed using MGI Tech MGIEasy RNA Directional Library Prep Kit V2 (1000006386). Ribosomal RNA was depleted using the rRNA depletion kit bacteria (NEB). Resulting libraries were sequenced on an MGIEasy MGISEQ2000RS using V2 chemistry. For all experiments, samples were sequenced to a depth that would ensure least 10 million reads per isolate in the sample.

**Transcriptomic analysis**

Prior to analysis, we assessed the suitability of our two genetically distant *E. coli* donor and recipient strains for RNA-seq analysis. Genetic variation between the donor and recipient was assessed using snippy version 4.6 (48) which detected >75,000 genetic variants including >68,000 SNPs which were equally distributed across the recipient genome (Supplementary Fig 4). This number was in line with previous single species co-culture RNA-seq experiments (49). Although the two strains were sufficiently genetically distinct, 53 genes that were identical between DLL7555 and HB101 were excluded from the final analysis. RNA reads were aligned against a concatenated sequence of the donor

(accession, CP184671.1), recipient (DLL7555, accession CP186931.1 or B5055,

assembly GCF_024138975.1), and pVS520 sequences (CP184672.1). The sequence for

HB101 (Rif$^S$ version of LT101) was used as the donor sequence owing to its availability.

Read alignment was performed with subread version v1.5.1 (50) against the concatenated

sequence which discarded reads of ambiguous donor/recipient origin. >99.8% of donor

originating reads were filtered by this strategy for both B5055 and DLL755 matings, which

was assessed by alignment of RNA isolated from the donor only condition against the

concatenated sequence. Read counting and assignment were performed via

FeatureCounts (51). Differential gene expression was analysed with the Voom/Limma

method in Degust v4.1 (https://drpowell.github.io/degust/) with a minimum of 3 counts per

million reads across 3 samples. At this stage, an additional 5 genes were subsequently

excluded from the DLL7555 analysis due to the detection of donor originating reads which

were not successfully filtered earlier. Genes were considered differentially expressed if

they had a Log2 fold change ≥ 0.585, a P value ≤ 0.05, and an FDR ≤ 0.05. GO term

analysis was performed using the TopGO R package v2.58 (52), with the weight_01

algorithm using raw P-values due to the structured dependency of GO terms as per the

package guidelines. eggNOG mapper v2.1.8 (53) was used to assign GO-terms to the

GenBank recipient proteomes. Gene and pathway functions were also assessed using the

ecocyc database (54). Heatmaps and GO term plots were generated using ggplot2 v3.5.2

(55).

**Scanning and Transmission Electron Microscopy**

SEM was performed on *K. pneumoniae* cells during conjugation, or in stationary phase

when capsule is highly expressed(20). 500 µl of cells were collected onto a 0.4 µm

polycarbonate membrane, and the capsular ultrastructure preserved by lysine acetate,

ruthenium red and osmium (LRR) fixation (56), followed by graduated dehydration in

ethanol, critical point drying, and gold sputter coating as previously described (57).

Samples were imaged on a field emission NovaSEM 450 SEM with an acceleration

voltage of 10 kV in secondary electron mode with a working distance of ~4.5mm. Images

were taken under immersion mode through the lens detector.

For TEM imaging, cells were collected on polycarbonate membranes and prepared with
LRR fixation as for SEM. Fixed membranes were then cut into small rectangles no longer
than 1 mm cubed, dehydrated and then infiltrated with Epon resin (58). Each membrane
was placed on a glass slide cell-side up, and a BEEM® capsule containing 100% resin
positioned over the flat membrane before polymerisation at 60°C for 48 hours. The resin-
filled BEEM® capsule and slide were cooled in liquid nitrogen and then immediately
separated, leaving the flat membrane with cells in a longitudinal orientation on the surface
of the resin block. Cells were ultrathin sectioned using a Leica UC7 ultramicrotome and
Diatome diamond knife and then contrasted with uranyl acetate and lead citrate. Cells
were imaged on a JEOL JEM-1400 Plus TEM at 80 KeV, equipped with a high sensitivity
bottom mount CMOS 'Flash' camera. Total capsule diameter was measured from the
average of 5 total cell widths from which the cell body width was subtracted, as described
previously (59).
590

**Fluorescent Microscopy and Image analysis**
Conjugation was analysed by fluorescent microscopy by inoculation of 1 µl of $OD_{600}$ 0.1
recipient and donor onto LB 1% agarose pads. Pads were incubated in parafilm sealed 25
mm fluorodish chamber (WPI) for 2 hours at 37°C. Small strips of moist KimWipe® were
included in the chamber to prevent pad dehydration. Images were acquired using a
63x/1.30 oil objective on a Leica DMI-8 widefield microscope equipped with a Lecia K8
sCMOS monochromatic camera using Lecia LasX software. Mating pair productivity was
assessed by counting transconjugants that were in direct contact with a donor and dividing
that by the total number of recipients in contact with a donor.

**Fluorescent reporter assays**
Fluorescent measurements were taken on a Tecan infinite pro 200 plate reader. For
measurement of P*traJ*, 15 µl of *K. pneumoniae* was inoculated onto a 200 µl LB 1%
agarose pad with rifampicin in a 96-well special optics microtiter plate (Corning). After 5
minutes of incubation and air drying, 15 µl of donor culture was spotted on top and
immediately afterwards the OD600 and fluorescent intensity (488 nm / 520 nm) were
measured in 5-minute intervals at 37°C. Fluorescent intensity was then normalised to

OD600 and analysed as change over time by calculating fold-change against the 0-time
point value.

**Acknowledgments:** We would like to acknowledge Dr Francesca Short for providing
strains and protocols for RNA isolation, and Dr Tom Smallman and Prof John Boyce for
providing strains. The authors acknowledge use of the facilities and the assistance of Alex
Fulcher from Monash Micro Imaging. The authors acknowledge the use of instruments
and assistance at the Monash Ramaciotti Centre for Cryo-Electron Microscopy, a Node of
Microscopy Australia. The authors acknowledge the MHTP Medical Genomics Facility for
assistance with RNA sequencing.

**Funding:** This work was supported by the Australian Research Council Grant
DP250103521, and Australian Research Council Laureate Fellowship FL210100258
awarded to D.L., and funding for Y.H. Gan from the National Medical Research Council,
grant number OFIRG20NOV-0045.

## References

1.    S. Castañeda-Barba, E. M. Top, T. Stalder, Plasmids, a molecular cornerstone of antimicrobial resistance in the One Health era. *Nature Reviews Microbiology* **22**, 18-32 (2024).

2.    C. Virolle, K. Goldlust, S. Djermoun, S. Bigot, C. Lesterlin, Plasmid Transfer by Conjugation in Gram-Negative Bacteria: From the Cellular to the Community Level. *Genes (Basel)* **11** (2020).

3.    K. Billane, E. Harrison, D. Cameron, M. A. Brockhurst, Why do plasmids manipulate the expression of bacterial phenotypes? *Philosophical Transactions of the Royal Society B: Biological Sciences* **377**, 20200461 (2022).

4.    H. Lee, K. S. Ko, Effect of multiple, compatible plasmids on the fitness of the bacterial host by inducing transcriptional changes. *Journal of Antimicrobial Chemotherapy* **76**, 2528-2537 (2021).

5.    C. Huang *et al.*, Comparative Analysis of Transcriptome and Proteome Revealed the Common Metabolic Pathways Induced by Prevalent ESBL Plasmids in Escherichia coli. *International Journal of Molecular Sciences* **24**, 14009 (2023).

6.    A. San Millan, M. Toll-Riera, Q. Qi, R. C. MacLean, Interactions between horizontally acquired genes create a fitness cost in Pseudomonas aeruginosa. *Nature Communications* **6**, 6845 (2015).

7.    K. Schaufler *et al.*, Carriage of Extended-Spectrum Beta-Lactamase-Plasmids Does Not Reduce Fitness but Enhances Virulence in Some Strains of Pandemic E. coli Lineages. *Front Microbiol* **7**, 336 (2016).

8.    Z. Baharoglu, D. Bikard, D. Mazel, Conjugative DNA Transfer Induces the Bacterial SOS Response and Promotes Antibiotic Resistance Development through Integron Activation. *PLOS Genetics* **6**, e1001165 (2010).

9.    K. Moriguchi *et al.*, Targeting Antibiotic Resistance Genes Is a Better Approach to Block Acquisition of Antibiotic Resistance Than Blocking Conjugal Transfer by Recipient Cells: A Genome-Wide Screening in Escherichia coli. *Front Microbiol* **10**, 2939 (2019).

10.   D. Pérez-Mendoza, F. de la Cruz, Escherichia coli genes affecting recipient ability in plasmid conjugation: Are there any? *BMC Genomics* **10**, 71 (2009).

11.   C. M. Johnson, A. D. Grossman, The Composition of the Cell Envelope Affects Conjugation in Bacillus subtilis. *J Bacteriol* **198**, 1241-1249 (2016).

12.   M. Haudiquet, A. Buffet, O. Rendueles, E. P. C. Rocha, Interplay between the cell envelope and mobile genetic elements shapes gene flow in populations of the nosocomial pathogen Klebsiella pneumoniae. *PLOS Biology* **19**, e3001276 (2021).

13.   W. W. Low *et al.*, Mating pair stabilization mediates bacterial conjugation species specificity. *Nature Microbiology* **7**, 1016-1027 (2022).

14.   M. Al Mamun Abu Amar, K. Kishida, J. Christie Peter, Protein Transfer through an F Plasmid-Encoded Type IV Secretion System Suppresses the Mating-Induced SOS Response. *mBio* **12**, 10.1128/mbio.01629-01621 (2021).

15.   L. González-Montes, I. Del Campo, M. P. Garcillán-Barcia, F. de la Cruz, G. Moncalián, ArdC, a ssDNA-binding protein with a metalloprotease domain, overpasses the recipient

hsdRMS restriction system broadening conjugation host range. *PLoS Genet* **16**, e1008750
(2020).

16.  K. H. Maslowska, K. Makiela-Dzbenska, I. J. Fijalkowska, The SOS system: A complex and
tightly regulated response to DNA damage. *Environ Mol Mutagen* **60**, 368-384 (2019).

17.  M. Miyakoshi, Y. Ohtsubo, Y. Nagata, M. Tsuda, Transcriptome Analysis of Zygotic
Induction During Conjugative Transfer of Plasmid RP4. *Frontiers in Microbiology* **Volume
11 - 2020** (2020).

18.  H. Prensky, A. Gomez-Simmonds, A. C. Uhlemann, A. J. Lopatkin, Conjugation dynamics
depend on both the plasmid acquisition cost and the fitness cost. *Molecular Systems
Biology* **17**, e9913 (2021).

19.  M. Haudiquet *et al.*, Capsules and their traits shape phage susceptibility and plasmid
conjugation efficiency. *Nature Communications* **15**, 2032 (2024).

20.  M. J. Dorman, T. Feltwell, D. A. Goulding, J. Parkhill, F. L. Short, The Capsule Regulatory
Network of Klebsiella pneumoniae Defined by density-TraDISort. *mBio* **9** (2018).

21.  M. Palacios *et al.*, Identification of Two Regulators of Virulence That Are Conserved in
Klebsiella pneumoniae Classical and Hypervirulent Strains. *mBio* **9** (2018).

22.  J. Hall Rebecca, E. Snaith Ann, J. N. Thomas Matthew, A. Brockhurst Michael, A. McNally,
Multidrug resistance plasmids commonly reprogram the expression of metabolic genes
in Escherichia coli. *mSystems* **9**, e01193-01123 (2024).

23.  L. Toribio-Celestino *et al.*, A plasmid-chromosome crosstalk in multidrug resistant
enterobacteria. *Nature Communications* **15**, 10859 (2024).

24.  M. Xiao, Y. Lai, J. Sun, G. Chen, A. Yan, Transcriptional Regulation of the Outer Membrane
Porin Gene ompW Reveals its Physiological Role during the Transition from the Aerobic
to the Anaerobic Lifestyle of Escherichia coli. *Front Microbiol* **7**, 799 (2016).

25.  T. A. Krulwich, G. Sachs, E. Padan, Molecular aspects of bacterial pH sensing and
homeostasis. *Nat Rev Microbiol* **9**, 330-343 (2011).

26.  Z. Mao *et al.*, NhaA facilitates the maintenance of bacterial envelope integrity and the
evasion of complement attack contributing to extraintestinal pathogenic Escherichia coli
virulence. *Infect Immun* **91**, e0003923 (2023).

27.  J. M. Ghigo, Natural conjugative plasmids induce bacterial biofilm development. *Nature*
**412**, 442-445 (2001).

28.  S. Djermoun *et al.*, Biofilm architecture determines the dissemination of conjugative
plasmids. *Proceedings of the National Academy of Sciences* **122**, e2417452122 (2025).

29.  M. Yong *et al.*, Dominant Carbapenemase-Encoding Plasmids in Clinical Enterobacterales
Isolates and Hypervirulent Klebsiella pneumoniae, Singapore. *Emerg Infect Dis* **28**, 1578-
1588 (2022).

30.  G. Schröder, E. Lanka, The mating pair formation system of conjugative plasmids—A
versatile secretion machinery for transfer of proteins and DNA. *Plasmid* **54**, 1-25 (2005).

31.  W. Pansegrau, E. Lanka, Enzymology of DNA Transfer by Conjugative Mechanisms.
*Progress in Nucleic Acid Research and Molecular Biology* **54**, 197-251 (1996).

32.  J. Haase, R. Lurz, A. M. Grahn, D. H. Bamford, E. Lanka, Bacterial conjugation mediated by
plasmid RP4: RSF1010 mobilization, donor-specific phage propagation, and pilus
production require the same Tra2 core components of a proposed DNA transport
complex. *Journal of Bacteriology* **177**, 4779-4791 (1995).

33.    S. Bates, A. M. Cashmore, B. M. Wilkins, IncP plasmids are unusually effective in mediating
727            conjugation of Escherichia coli and Saccharomyces cerevisiae: involvement of the tra2
728            mating system. *J Bacteriol* **180**, 6538-6543 (1998).
729    34.    T. R. D. Costa, J. B. Patkowski, K. Macé, P. J. Christie, G. Waksman, Structural and
730            functional diversity of type IV secretion systems. *Nature Reviews Microbiology* **22**, 170-
731            185 (2024).
732    35.    A. K. Vadakkepat *et al.*, Cryo-EM structure of the R388 plasmid conjugative pilus reveals
733            a helical polymer characterized by an unusual pilin/phospholipid binary complex.
734            *Structure* 10.1016/j.str.2024.06.009.
735    36.    D. E. Bradley, Morphological and serological relationships of conjugative pili. *Plasmid* **4**,
736            155-169 (1980).
737    37.    V. Cyriaque *et al.*, Single-cell RNA sequencing reveals plasmid constrains bacterial
738            population heterogeneity and identifies a non-conjugating subpopulation. *Nature
739            Communications* **15**, 5853 (2024).
740    38.    S. E. George *et al.*, Phenotypic heterogeneity and temporal expression of the capsular
741            polysaccharide in Staphylococcus aureus. *Molecular Microbiology* **98**, 1073-1088 (2015).
742    39.    J. E. King, H. A. Aal Owaif, J. Jia, I. S. Roberts, Phenotypic Heterogeneity in Expression of
743            the K1 Polysaccharide Capsule of Uropathogenic Escherichia coli and Downregulation of
744            the Capsule Genes during Growth in Urine. *Infect Immun* **83**, 2605-2613 (2015).
745    40.    L. M. Reyes Ruiz, C. L. Williams, R. Tamayo, Enhancing bacterial survival through
746            phenotypic heterogeneity. *PLOS Pathogens* **16**, e1008439 (2020).
747    41.    S. Hammerschmidt *et al.*, Capsule phase variation in Neisseria meningitidis serogroup B
748            by slipped-strand mispairing in the polysialyltransferase gene (siaD): correlation with
749            bacterial invasion and the outbreak of meningococcal disease. *Mol Microbiol* **20**, 1211-
750            1220 (1996).
751    42.    J. Zhu *et al.*, Effects of Capsular Polysaccharide amount on Pneumococcal-Host
752            interactions. *PLOS Pathogens* **19**, e1011509 (2023).
753    43.    K. A. Walker, V. L. Miller, The intersection of capsule gene expression, hypermucoviscosity
754            and hypervirulence in Klebsiella pneumoniae. *Current opinion in microbiology* **54**, 95-102
755            (2020).
756    44.    M. A. Schembri, J. Blom, K. A. Krogfelt, P. Klemm, Capsule and fimbria interaction in
757            Klebsiella pneumoniae. *Infect Immun* **73**, 4626-4633 (2005).
758    45.    D.-W. Wei *et al.*, Insertion sequences accelerate genomic convergence of multidrug
759            resistance and hypervirulence in Klebsiella pneumoniae via capsular phase variation.
760            *Genome Medicine* **17**, 45 (2025).
761    46.    S. Zobel *et al.*, Tn7-Based Device for Calibrated Heterologous Gene Expression in
762            Pseudomonas putida. *ACS Synthetic Biology* **4**, 1341-1351 (2015).
763    47.    N. R. Mattatall, K. E. Sanderson, Salmonella typhimurium LT2 possesses three distinct 23S
764            rRNA intervening sequences. *Journal of Bacteriology* **178**, 2272-2278 (1996).
765    48.    T. Seemann, Snippy: rapid haploid variant calling and core SNP phylogeny. *GitHub.
766            Available at: github. com/tseemann/snippy* (2015).
767    49.    T. A. Gray *et al.*, Intercellular communication and conjugation are mediated by ESX
768            secretion systems in mycobacteria. *Science* **354**, 347-350 (2016).

50. Y. Liao, G. K. Smyth, W. Shi, The Subread aligner: fast, accurate and scalable read mapping
by seed-and-vote. *Nucleic Acids Res* **41**, e108 (2013).
51. Y. Liao, G. K. Smyth, W. Shi, featureCounts: an efficient general purpose program for
assigning sequence reads to genomic features. *Bioinformatics* **30**, 923-930 (2014).
52. A. Alexa, J. Rahnenführer, Gene set enrichment analysis with topGO. *Bioconductor Improv*
**27**, 776 (2009).
53. C. P. Cantalapiedra, A. Hernández-Plaza, I. Letunic, P. Bork, J. Huerta-Cepas, eggNOG-
mapper v2: Functional Annotation, Orthology Assignments, and Domain Prediction at the
Metagenomic Scale. *Molecular Biology and Evolution* **38**, 5825-5829 (2021).
54. P. D. Karp *et al.*, The EcoCyc Database (2023). *EcoSal Plus* **11**, eesp-0002-2023 (2023).
55. H. Wickham, C. Sievert, *ggplot2: elegant graphics for data analysis* (springer New York,
2009), vol. 10.
56. S. Hammerschmidt, M. Rohde, "Electron Microscopy to Study the Fine Structure of the
Pneumococcal Cell" in Streptococcus pneumoniae: Methods and Protocols*, F. Iovino, Ed.
(Springer New York, New York, NY, 2019), 10.1007/978-1-4939-9199-0_2, pp. 13-33.
57. T. R. Smallman, G. C. Williams, M. Harper, J. D. Boyce, Genome-Wide Investigation of
Pasteurella multocida Identifies the Stringent Response as a Negative Regulator of
Hyaluronic Acid Capsule Production. *Microbiology Spectrum* **10**, e00195-00122 (2022).
58. S. A. C. Jillian C Danne and Jillian C Danne, Rachel Templin, Joan Clark, Viola Oorschot,
Georg Ramm, Preparation of cells for transmission electron microscopy ultrastructural
analysis. *protocols.io* dx.doi.org/10.17504/protocols.io.j8nlk47ewg5r/v1 (2022).
59. T. Guess *et al.*, Size Matters: Measurement of Capsule Diameter in Cryptococcus
neoformans. *JoVE* doi:10.3791/57171, e57171 (2018).
60. B. M. Coffey, G. G. Anderson, Biofilm formation in the 96-well microtiter plate. *Methods*
*Mol Biol* **1149**, 631-641 (2014).
61. E. A. Palombo, K. Yusoff, V. A. Stanisich, V. Krishnapillai, N. S. Willetts, Cloning and genetic
analysis of tra cistrons of the Tra 2/Tra 3 region of plasmid RP1. *Plasmid* **22**, 59-69 (1989).
62. Y. Fu *et al.*, Mechanisms of blaIMP-4 dissemination across diverse carbapenem-resistant
clinical isolates. *Journal of Global Antimicrobial Resistance* **41**, 189-194 (2025).
63. M. M. C. Lam *et al.*, Population genomics of hypervirulent Klebsiella pneumoniae clonal-
group 23 reveals early emergence and rapid global dissemination. *Nature*
*Communications* **9**, 2703 (2018).
64. I. R. Lee *et al.*, Differential host susceptibility and bacterial virulence factors driving
Klebsiella liver abscess in an ethnically diverse population. *Scientific Reports* **6**, 29316
(2016).
65. A. C. Jacobs *et al.*, AB5075, a Highly Virulent Isolate of Acinetobacter baumannii, as a
Model Strain for the Evaluation of Pathogenesis and Antimicrobial Treatments. *mBio* **5**,
e01076-01014 (2014).
66. L. A. Gallagher *et al.*, Resources for Genetic and Genomic Analysis of Emerging Pathogen
Acinetobacter baumannii. *J Bacteriol* **197**, 2027-2035 (2015).
67. R. F. Wang, S. R. Kushner, Construction of versatile low-copy-number vectors for cloning,
sequencing and gene expression in Escherichia coli. *Gene* **100**, 195-199 (1991).

68.   B. Bartolomé, Y. Jubete, E. Martínez, F. de la Cruz, Construction and properties of a family
of pACYC184-derived cloning vectors compatible with pBR322 and its derivatives. *Gene*
**102**, 75-78 (1991).
69.   J. P. Fürste *et al.*, Molecular cloning of the plasmid RP4 primase region in a multi-host-
range tacP expression vector. *Gene* **48**, 119-131 (1986).

**Figures and Tables**

**Figure 1. The transcriptional response of an *E. coli* recipient and plasmid RP4 during active**
**conjugation**. **a)** *E. coli* DLL7555 conjugation frequency over time shows that RP4 conjugation is
active at 15 minutes and completed by 30 minutes. Arrow indicates time for transcriptional
sampling. **b)** Differentially expressed DLL7555 genes during active RP4 conjugation. **c)** GO term
analysis of enriched genes with the R TopGo package with the weight_01 algorithm. **d)** Differential
expression of RP4 encoded genes during active conjugation with *E. coli* DLL7555. Numbers in *E.*
*coli* gene list represent locus tags trimmed of the prefix ACPEGL. Error bars = mean ± SD, N = 4.
862
863

**Figure 2. The effect and mechanism of capsule on conjugation and the recipient and plasmid response. a)** Differentially expressed genes (DEGS) in *K. pneumoniae* B5055 (top) and B5055Δ*wzbc* (bottom) genes during active conjugation or co-culture with an RP4 donor. **b)** GO term analysis of DEG for B5055 (top) and B5055Δ*wzbc*, with raw P-values reported as per TopGO package guidelines for the weight_01 algorithm. **c)** Common DEGS between recipient responses (top), that are increased in expression (middle) or decreased (bottom). **d)** No RP4 DEGS were detected in the conjugation mixture with B5055. **e)** Differentially expressed RP4 genes during active conjugation with B5055Δ*wzbc*. **f)** DEGs shared between RP4 during conjugation across different recipients. **g)** TEM, and SEM images of conjugation between *K. pneumoniae* B5055 or B5055Δ*wzbc* recipients with *E. coli* LT101. Recipients are distinguished based on the presence of capsule, and shape. **h)** *in situ* visualisation of conjugation between either wildtype or Δ*wzbc* recipients by fluorescent microscopy. All images are representative of ≥2 independent experiments. Abbreviations: D=Donor, R=Recipient, T=Transconjugant. Numbers in *K. pneumoniae* gene lists represent locus tags trimmed of the prefix J6T75_RS. Scale bars: SEM 3 μm, TEM 1 μm, Fluorescence microscopy 20 μm, inset 5 μm.

**Figure 3. Recipient *K. pneumoniae* capsule thickness variation impacts donor-recipient interactions and conjugation. a)** SEM imaging of capsular variation and projections. **b-c)** TEM imaging of capsule thickness variation between individual cells. **d)** Variation in capsule thickness within, and between biological replicates (n = ≥50 cells per sample, one-way ANOVA with sidak's multiple comparison testing P-values = ** ≤ 0.01, **** ≤ 0.0001.). **e-f)** Effect of capsule thickness variation on donor-recipient interactions. **g)** linear regression analysis of distance between outer membrane of donor and recipient and total capsular diameter ($R^2$=.72, N=47, F-test P≤0.0001). **h)**

905   live cell imaging of B5055 transconjugant formation, showing that B5055 transconjugants arise
906   from recipients   intimately contacting the donor and morphologically resemble acapsular
907   B5055Δ*wzbc*. Error bars = mean ± SD. Scale bars: a = 5 μm, bc = 200 nm, ef = 500 nm, f inset =
908   200 nm, h = 20 μm, h insert = 5 μm. Images are representative of ≥ 2 independent experiments,
909   except for h which depicts the rare formation of a B5055 transconjugant.

**Figure 4. Population level capsule variation enables RP4 conjugation in capsulated *K. pneumoniae*.**

**a)** Variation in conjugation frequency of *K. pneumoniae* B5055. The conjugation frequency of B5055Δ*wzbc* is shown for context. **b)** Differential expression of RP4 encoded genes between conjugation high and low samples. **c)** differential expression of genes in the capsule synthesis locus between conjugation high (top) and low (bottom) samples. **d)** Genes associated with capsule production between samples with high or low conjugation frequencies. **e)** The

923    conjugation frequency of RP4 to hypermucoid K1 or K5 *K. pneumoniae* (N=9, P>.05 Mann-

924    Whitney U test). Data are presented as the median (**a**), or the mean+SEM (**d**).

927
928

**Supplementary Figure 1: Donor LT101(pVS520) and Recipient DLL7555 co-culture causes a**
**minor reduction in biofilm formation.**

Biofilms were grown as in (60). Briefly, 1:100 dilutions of overnight broth cultures of donor, recipient
and mock donor in fresh LB medium were mixed in 1:1 in a flat bottomed 96 well tray and grown
for 24 hours at 37°C. Outer wells were not used to avoid evaporation effects. Following incubation,
the liquid portion of the culture was removed and plated to confirm transconjugants were present
and therefore conjugation had occurred. The biofilms were washed in water, and quantitated by
staining with 0.1% crystal violet, which after washing was solubilized with 30% acetic acid and
measured by absorbance reading at 550 nm standardized against 30% acetic acid.  Error bars =
mean ± SD, N = 4 biological replicates repeated 3 times). Significance testing was performed with
a Kruskal Wallis one way ANOVA with Dunn's multiple comparison testing P-values = *** ≤ 0.01,
**** ≤ 0.0001.).

**Supplementary Figure 2: Capsule is a barrier to conjugation in diverse species.**

The conjugation frequency of RP4 or RP4-mobilizable plasmids from *E. coli* LT101 to K.

*pneumoniae* B5055, *A. baumannii* AB5075, or *P. multocida* VP161 and their unencapsulated

derivatives (Mean+SD, N=4, P>.05 Mann-Whitney U test).

**Supplementary Figure 3: Capsule inhibits activation of the RP4 MOB promoter P*traJ* within**

**the donor.** We constructed a fluorescent reporter for the RP4 MOB promoter P*traJ* by fusing it to

three tandem low sequence-homology GFP reporters. Three tandem GFPs were used because

fluorescence from 1 GFP reporter could not be detected, likely because P*traJ* is a weak promoter

**a)** we confirmed our reporter behaved as expected by comparing expression in the presence or

absence of RP4, which is known to repress P*traJ* activity (N = 4, $P<0.05$, Mann-Whitney U-Test).

**b)** Measurement of P*traJ*x3GFP in the donor cell was achieved by performing conjugations in a 96-

well tray on agar containing rifampicin which inhibits transcription only in the recipient cell.

Significance testing was performed with a one-way ANOVA with Sidak's multiple comparison. Error

bars = mean ± SD

**Supplementary Figure 4: SNP and RNA-seq read alignment for HB101(pVS520) x DLL7555.**
a) schematic overview of RNA-seq alignment strategy. b-c) IGV plots depicting the coverage of
SNPs, and the alignment of mating, pure recipient and pure donor read coverage against a recipient
portion of the concatenated reference sequence at a global (b) and local (c) level. Gaps indicated
by black arrows in (b) represent genome segments entirely unique to DLL7555. The reads are log
scaled in (b).

**Table 1.** Bacterial strains and plasmids used in this work.

| Strain | Genotype/Characteristics | Reference |
|---|---|---|
| *E. coli* | | |
| DH5α | F- φ80*lac*ZΔM15Δ(*lac*ZYA-*arg*F)U169 *rec*A1 *end*A1 *hsd*R17(r$_K$ ,m$_K^+$) *pho*A *sup*E44λ⁻*thi*-1 *gyr*A96 *rel*A1 | Invitrogen |
| LT101 | Rif$^R$ derivative of HB101 | (61) |
| DLL7555 | ST73 Rif$^R$ clinical isolate of *E. coli* | (62) |
| *K. pneumoniae* | | |
| B5055 | Wild type *K. pneumoniae* O1:K2 | Provided by Richard Strugnell |
| B5055Δ*wzbc* | Non-encapsulated *wzbc* deletant mutant of B5055, Km$^R$ | Provided by Richard Strugnell, Adam Jenney, and Tania Uren |
| SGH10 | Hospital isolate of *K. pneumoniae* K1, ST23 | (63) |
| NUH26 | Hospital isolate of *K. pneumoniae* K5, ST1049 | (64) |
| NUH40 | Hospital isolate of *K. pneumoniae* K5, ST1049 | (29) |
| *A. baumannii* | | |
| AB5075 | Wild type *A. baumanii* KL25, Ap$^R$, Erm$^S$ | (65) |
| AB10132 | Capsular transposon mutant of AB5075, *wzC*::Tn*5*, Tc$^R$ | (66) |
| *Pasteurella multocida* | | |
| AL3145 | Km$^R$ derivative of VPI161 | (57) |
| AL2234 | VP161 capsular transposon mutant *hyaD*-Tn*7,* Km$^R$ | (57) |
| Plasmids | | |
| pWSK29 | Ap$^R$, low copy number cloning vector | (67) |
| pWSK29 P*traJ*GFPx3 | P*traJ*::*mGreenLantern*x3 | |
| pSU18 | Cm$^R$, medium copy number cloning vector | (68) |
| pSU18*mTagBFP2* | P*em7*::*mTagBFP2* | This work |

| | | |
|---|---|---|
| pVS520 | $Tc^R$, $Ap^S$, $Km^S$, derivative of RP4; $Tra^+$ $Mob^+$ | (61) |
| pMMB67EH | RSF1010 derived expression vector. $Ap^R$, $Mob^+$ | (69) |
| RFS1010-*erm* | $Erm^R$ derivative of RSF1010, $Mob^+$ | This work |
| pAL99S-*oriT* | $Spec^R$, $Mob^+$ derivative of pAL99S | This work |
| pMMB67EH*gfpmut3* | P*em7*::*gfpmut3* | This work |

**Note:** rif = rifampicin, km = kanamycin, tc = tetracycline, ap = ampicillin, cm = chloramphenicol, erm = erythromycin, spec = spectinomycin