A short linear motif, conserved from yeast to human, binds to members of the Spa2/GIT1 family of cortical scaffold proteins

by Lara Bareis, Annika Siewert, Benjamin Grupp, Tim Bergner*, Clarissa Read*, Steffi Timmermann, Nicole Schmid, and Nils Johnsson

Institute of Molecular Genetics and Cell Biology, Department of Biology, Ulm University, James-Franck-Ring N27, D-89081 Ulm, Germany

*Electron microscopy facility, Ulm University

Contact: N. Johnsson (e-mail: nils.johnsson@uni-ulm.de; orcid number: 0000-0003-1416-835X), Department of Biology, Ulm University, James-Franck-Ring N27, D-89081 Ulm, Germany. Phone: + 49 731 50 36300. Fax: + 49 731 50 36302.

Running title: A conserved Spa2-binding motif

Abstract

Tip growth is closely tied to fungal pathogenicity. Spa2, a multi-domain protein and member of the polarisome, orchestrates tip growth in yeast and other fungi. We identified a conserved short linear motif in the RabGAPs Msb3 and Msb4, and the MAP kinase kinase Ste7 and Mkk1, which mediates their interaction with Spa2. AlphaFold predictions suggest that these initially unstructured motifs adopt an alpha-helical conformation upon binding to the hydrophobic cleft of Spa2's N-terminal domain. Altering the predicted key contact residues in either Spa2 or the motif reduces complex stability. Such mutations also cause mis-localization of Msb3, Msb4, and Ste7 within the cell. Deleting the motif in Msb3 or Msb4 abolishes tip-directed growth of the yeast bud. Protein assemblies that spatially confine secretion to specific membrane regions are a common feature of eukaryotic cells. Accordingly, Spa2-motif complexes were predicted in orthologs and paralogs across selected Opisthokonta, including pathogenic fungi and humans. A search for functional motifs in conformationally flexible regions of all yeast proteins identified Dse3 as a novel Spa2-binding partner.

Introduction

Budding yeast, like other walled fungi, shapes its growth by directing secretory vesicles to specific plasma membrane regions during the cell cycle. The polarisome, a multi-protein assembly, orchestrates this process by selecting target sites for vesicle fusion (Sheu et al., 1998; Sheu et al., 2000; Dünkler et al., 2021). Its primary scaffold protein, Spa2, interacts with two Rab-GAPs, Msb3 and Msb4, which convert the Rab-GTPase Sec4 from its active GTP-bound state to its inactive GDP-bound state (Snyder, 1989; Gao et al., 2003; Tcheperegine et al., 2005). GTP-bound Sec4 recruits the motor protein Myo2 and the exocyst component Sec15 to post-Golgi vesicles (Guo et al., 1999; Jin et al., 2011). The vesicle then moves on actin cables to the sites of the bud membrane where fusion occurs. Deletion of both Msb3 and Msb4 impairs vesicle fusion with the plasma membrane, leading to vesicle accumulation in the bud (Tcheperegine et al., 2005). The GAP activity not only recycles Msb3/4 but also catalyzes a critical, yet poorly understood, step in vesicle targeting or fusion (Gingras et al., 2022). Thus, a key function of the polarisome during tip growth is to localize Msb3/4 to prospective vesicle fusion sites. The interaction between Spa2 and the GAPs occurs via the N-terminal Spa2 homology domain 1 (SHD1) and potentially unstructured N-terminal regions of Msb3/4 (Tcheperegine et al., 2005).

Intrinsically disordered regions, often less conserved evolutionarily, may contain short linear motifs that regulate protein localization, stability, or interactions (Holehouse and Kragelund, 2024; Mughal and Caetano-Anollés, 2025). These motifs are typically recognized by folded domains of their binding partners. Such

interactions are often weak or transient, making them challenging to detect. As a consequence, many motifs likely remain undiscovered within the disordered regions of an organism's proteome (Tompa et al., 2014). However, identifying a motif's sequence can reveal its presence in other proteins across species, enabling predictions of binding partners, localization, and even cellular roles (Neduva et al., 2005).

Here, we identify a short linear motif (SLiM) in the disordered regions of Msb3, Msb4, the MAP kinase kinases Ste7 and Mkk1 that binds the SHD1 domain of Spa2. AlphaFold predictions reveal SHD1-motif interactions across the Opisthokonta, demonstrating that the motif's sequence, its interaction with SHD1, and its role in linking the polarisome to vesicle fusion are evolutionarily conserved.

Results

A conserved five-residue motif mediates binding of Msb3, Msb4, and Ste7 to Spa2

The Spa2 homology domain 1 of Spa2 (SHD1) is a conserved protein interaction module that is essential for the scaffold functions of Spa2 during bud growth and mating of yeast cells (Sheu et al., 1998; Lawson et al., 2022). A Split-Ubiquitin screen of SHD1CRU against an array of 530 N_{ub}-fusion proteins, enriched for polar growth regulators, identified Msb3, Msb4 and Ste7 as interaction partners of SHD1 (Fig. 1A) (Hruby et al., 2011). Ste7 is the MAP kinase kinase in the kinase signaling pathway of the mating response of yeast cells (Courchesne et al., 1989; Teague et al., 1986). The GAP domains of Msb3, Msb4 and the kinase domain of Ste7 are preceded by

intrinsically unstructured regions of 151, 59 and 187 residues in length respectively. All three regions were shown to be critical for the interaction with Spa2 (Sheu et al., 1998). Sequence analysis reveals a shared five-residue motif (V/I)-(I/L)-D-(L/A)-Y (from here on FRM) within the N-terminal regions of all three proteins (Fig. 1B). A similar motif is also found in the N-terminal regions of Mkk1 and Mkk2, MAP kinases of the cell wall integrity pathway, previously reported to interact with SHD1 (Fig. 1B) (Sheu et al., 1998).

Deletion of the FRM in N_{ub}-Msb3$_{\Delta84-88}$, N_{ub}-Msb4$_{\Delta35-39}$, and N_{ub}-Ste7$_{\Delta119-123}$ abolished their binding to SHD1CRU (Fig. 1C). Ste7's N-terminal region also binds the mating scaffold Ste5 (Fig. 1D). This interaction is unaffected by the FRM deletion, as N_{ub}-Ste7$_{\Delta119-123}$ retains similar affinity for Ste5CRU as wild-type N_{ub}-Ste7 (Fig. 1D). The FRM does not influence Msb3 or Msb4 binding to their substrate, the Rab-GTPase Sec4, as CRU fusions of Msb3, Msb4, and their FRM-deficient mutants (Msb3$_{\Delta76-93}$CRU, Msb4$_{\Delta27-44}$CRU) bind N_{ub}-Sec4 with equal affinity (Fig. 1E). Conversely, N_{ub} fusions to Spa2 and the polarisome subunit Pea2 preferentially interact with CRU fusions of the wild-type GAPs (Fig. 1E) (Valtz and Herskowitz, 1996).

N-terminal deletion analysis of N_{ub}-fusions to Spa2 confirmed SHD1 as the sole FRM-binding site for Msb3CRU (Fig. 1F). In pull-down assays, GST-Msb3$_{1-150}$ precipitated 6xHis-SHD1, whereas GST-Msb3$_{1-150\Delta76-93}$ lacking the FRM did not (Fig. 1G, S1A). Short FRM-containing regions of Msb3 (N_{ub}-Msb3$_{71-100}$) and Msb4 (N_{ub}-Msb4$_{21-50}$), but not Ste7 (N_{ub}-Ste7$_{100-135}$), were sufficient for SHD1CRU interaction in the Split-Ubiquitin assay (Fig. 1H). Consistently, bacterially expressed GST-Msb3$_{71-98}$ and GST-Msb4$_{22-49}$ precipitated 6xHis-SHD1 *in vitro* (Fig. 1I, S1B).

Alpha-Fold predicts the FRM-SHD1 complex with high confidence

Alpha-Fold predicts a compact globular fold for SHD1 where five helices create a groove lined by hydrophobic side chains (Fig. S1C)(Abramson et al., 2024). The rim of the groove exposes positively charged residues. The prediction is in good agreement with the solved structures of the SHD1s of the Spa2 homologues from *Neurospora crassa* and from mouse (GIT1) (Zheng et al., 2020; Zhu et al., 2020). The FRMs of Msb3, Msb4, Ste7 and Mkk1 reside in unstructured regions of the proteins. When confronted with the sequence of SHD1, AlphaFold converts these unstructured, FRM-containing regions into alpha helices and grafts them into the hydrophobic pocket of SHD1 (Fig. 2A, S2). The model of the complex predicts that Tyr at position +2 of the motif and the hydrophobic residues at position -2, -1 and to a lesser extent +1 contact the hydrophobic face of the SHD1 groove. Asp at position 0 of Msb3/4 and Ste7 is oriented toward K97 of SHD1 (Fig. 2A). The hydroxyl group of Y+2 is hydrogen-bonded to D112 of Spa2. The sequence alignment of the Msb3 homologues of different yeasts underscores the importance of the hydrophobic character of the residues at positions -2, -1, and the Tyrosine at +2 (Fig. 2B). To independently test the model of the complex, we exchanged each residue of the FRM of Msb3 individually against an Ala, and in addition the conserved Leu at position -4 against Tyr. Split-Ub analysis of interactions of the N_{ub}-fusions showed that the exchanges D0A, and L-4Y outside of the motif do not measurably decrease the strength of the interaction between N_{ub}-Msb3 and SHD1CRU whereas the Ala-exchanges at positions -2, -1 and +2 disrupt the complex, and the exchange at position +1A only slightly impairs the interaction (Fig. 2C). The replacement Y+2F

strongly reduces complex stability (Fig. 2C). The low sensitivity of position +1 against the Ala-exchange is in line with the observation that the FRM of Ste7 displays an Ala and Msb3 of *Candida albicans* a Glu at this position (Fig. 1B, 2B). All Msb3 alleles were still predicted by AlphaFold to interact with SHD1. However, the scores of the interface predicted template modelling (ipTM) vary among the complexes and loosely correlate with the binding strength of the measured Split-Ub interactions. Mutations that impair the binding reduce the ipTM scores accordingly (Fig. 2C, lower panel). The Y+2A and the D0A exchanges had comparable effects on the complex stability of Msb4-SHD1 (Fig. 2D). This observation permitted us to generalize the findings on the FRM$_{Msb3}$-SHD1 interaction to the motifs of the other Spa2 binding partners. The Leu at positions 48, 51, 59, 98, 101, and 109 of Spa2 are predicted to contribute to the hydrophobic patch of the binding pocket of SHD1 (Fig. 2E, lower panel). We exchanged positions 48, 51, 101 and 109 individually or in combination against Ala and measured the interactions of the mutants against the N$_{ub}$-fusions of Msb4, Msb3, and Ste7. Each single residue exchange abolished the Split-Ub measured interaction between SHD1 and N$_{ub}$-Ste7 or N$_{ub}$-Msb3 (Fig. 2E). The dissolution of Split-Ub-measured interaction between Msb4 and SDH1 required two simultaneous exchanges in the SHD1 domain (L48A and L51A; L101A and L109A) (Fig. 2E). Exchanging K97 against D abolished the interaction between SHD1$_{K97D}$ and Msb3 but hardly affected the interaction between SHD1$_{K97D}$ and Msb4 or Ste7 (Fig. 2E).

Ste7, Msb3 and Msb4 compete for binding to Spa2

The similarity of the AlphaFold-structures and the comparable influences of the mutations within SHD1 on the interactions with its ligands strongly suggest but do not

prove that Msb3, Msb4 and Ste7 compete for the same interface of Spa2. AlphaFold prediction shows that FRMs once fused to the N-terminus of Spa2 will fold back into the groove of the SHD1 and should consequently block any intermolecular binding of those proteins that use the same site on Spa2 (Fig. 3A). We devised a competition assay where we extended the N-terminus of the full length Spa2 with residues 22-81 of Msb4 ($Msb4_{22-81}$-Spa2) (foldback competition assay, Fig. 3A). The N-terminal extension eliminated the binding of $Msb4_{22-81}$-Spa2CRU with N_{ub}-Msb3, and N_{ub}-Msb4 but did not affect the interactions with the N_{ub}-fusions of the known Spa2-binding partners Pea2, Epo1, and Aip5 (Fig. 3B) (Neller et al., 2015; Glomb et al., 2019). Deleting the FRM in the Msb4 extension ($Msb4_{22-81\Delta35-39}$-Spa2) recovered the interactions of $Msb4_{22-81\Delta35-39}$-Spa2CRU with N_{ub}-Msb4 and N_{ub}-Msb3 (Fig. 3B). The interaction between the full length Spa2CRU and N_{ub}-Ste7 is not detected by the Split-Ub assay. To confirm that FRM_{Ste7} shares with $FRM_{Msb3/Msb4}$ the same binding site on SHD1, we fused residues 105-165 of Ste7 in front of Spa2 ($Ste7_{105-162}$-Spa2CRU). As for $Msb4_{22-81}$-Spa2, $Ste7_{105-162}$ blocked the interaction between $Ste7_{105-162}$-Spa2CRU and N_{ub}-Msb3 and N_{ub}-Msb4 (Fig. 3B). Alternatively, we extended the N-terminus of the isolated SHD1 by the unstructured FRM-containing regions of Msb3, Msb4, Ste7 and tested their influence on the interaction between the SHD1CRU and N_{ub}-Msb3. The N-terminal extensions of Msb3/4 and Ste7 impaired the interaction between N_{ub}-Msb3 and SHD1CRU whereas those extensions lacking a functional FRM did not (Fig. 3C).

Mkk1 contains a FRM with a week affinity to SHD1

Sequence alignment reveals that Mkk1 possesses an FRM, yet N_{ub}-Mkk1 was not identified as a binding partner of SHD1CRU in our initial screen (Mkk2 is absent from the N_{ub}-fusion collection) (Fig. 1B). This could result from low affinity between SHD1 and Mkk1, low N_{ub}-Mkk1 expression, or differing cellular localization of CRU- and N_{ub}-fusion proteins. To address these factors, we employed the foldback competition assay, which is less sensitive to such variables (Fig. 3A, C). We fused residues 143–202 of Mkk1 to SHD1CRU (Mkk1$_{143-202}$SHD1CRU). Similar to the corresponding extensions of Msb3, Msb4 or Ste7, Mkk1$_{143-202}$ blocked the interaction of Mkk1$_{143-202}$SHD1CRU with N_{ub}-Msb3 (Fig. 3C). Deleting the FRM in Mkk1$_{143-202}$SHD1CRU (Mkk1$_{160-204\Delta171-175}$SHD1CRU) restored this interaction (Fig. 3C). The foldback competition assay reliably detects FRM-SHD1 interactions even for Mkk1 that binds SHD1 with lower affinity than Msb3, Msb4, or Ste7 (see also Fig.6B). This finding aligns with AlphaFold predictions, which assign the Mkk1-SHD1 complex the lowest ipTM score among the four FRM-SHD1 complexes (Fig. S2; see also Fig. 7).

To confirm FRM$_{Mkk1}$ binding to SHD1, we repeated the intermolecular Split-Ub assay with reduced SHD1CRU expression to enhance sensitivity. This revealed a weak but specific interaction between SHD1CRU and N_{ub}-Mkk1 (Fig. 3D). Inspection of the FRM sequence in Mkk1 and Mkk2 identified a serine at position 0, unlike the more common aspartate in Msb3/4 and Ste7 (Fig. 1B). Substituting Ser0 with Asp slightly enhanced the interaction signal, whereas replacing Leu2 with the more prevalent Ile had no effect. Deleting the FRM in N_{ub}-Mkk1 (N_{ub}-Mkk1$\Delta171-175$) reduced the interaction with SHD1CRU to background levels (Fig. 3E).

The SHD1$_{Spa2}$-FRM interaction targets the complex to sites of polar growth

Spa2 recruits Msb3 and Msb4 to the bud tip during bud growth, to the bud neck during mitosis and to the tip of mating projections (Tcheperegine et al., 2005; Lawson et al., 2022). Ste7-GFP localizes visibly only at the tip of mating projections (Maeder et al., 2007). To compare the influence of the FRM on the localization of each of the three proteins, we measured the cellular distributions of the respective GFP-fusions and their mutants upon exposure to mating hormone. Removing the FRMs from Msb3/4 and Ste7, or the SHD1 from Spa2 (Spa2$_{\Delta SHD1}$) abolishes the shmoo tip localization of Msb3-GFP, Msb4-GFP, and Ste7-GFP (Fig. 4A-D). In addition, tip and bud neck signals of the GFP-fusions to Msb3 and Msb4 in mitotic cells are lost or very much reduced when the FRMs of Msb3, Msb4 or the SHD1 of Spa2 are deleted (Fig. 4E).

The SHD1-FRM$_{Msb3/4}$ complex directs vesicle fusion to the tip of the cell.

The simultaneous loss of Msb3 and Msb4 induce the accumulation of post Golgi vesicles in the buds of *msb3Δmsb4Δ*-cells (Fig. 5A) (Gao et al., 2003; Tcheperegine et al., 2005). To investigate the influence of the Spa2-Msb3/4 interaction on the fusion of post-Golgi vesicles, we deleted the FRM of Msb3 in a *msb4Δ*-strain (*msb4Δ msb3$_{\Delta 76-93}$*), the FRM of Msb4 in a *msb3Δ* strain (*msb3Δ msb4$_{\Delta 27-44}$*), or the SHD1 of Spa2 (*spa2$_{\Delta SHD1}$*). Electron microscopy of these cells did not reveal the characteristic accumulation of 50-70 nm vesicles that are prominently seen in *msb3Δmsb4Δ*-cells (Fig. 5A, B). Accordingly, no vesicle accumulation occurs in cells that lack the SHD1 of Spa2 (*spa2$_{\Delta 2-150}$*), or lack Spa2 (*spa2Δ*) completely (Fig. 5A, B).

A tip-directed fusion of vesicles leads to ellipsoid bud shapes of the yeast, whereas random fusion leads to round bud shapes (Dünkler et al., 2021). The extent of tip-

directed vesicle fusion can be approximated by the length to width ratio of the bud (L/W ratio). *spa2Δ-* and *msb3Δmsb4Δ-*cells have a very similar L/W ratio that is significantly smaller than those of wild type cells (Fig.5C). Cells expressing $msb3_{\Delta 76-93}$ in *msb4Δ-*cells, or $msb4_{\Delta 27-44}$ in *msb3Δ-*cells display L/W ratios that are close to those of *spa2Δ-* or *msb3Δmsb4Δ-* cells. Deleting SHD1, or introducing the L48A L51A mutations in the hydrophobic pocket of SHD1 reduces the L/W ratio to a similar extent (Fig.5C). Accordingly, cells expressing $Ste7_{105-162}$-Spa2, or $Msb4_{22-81}$-Spa2, showed a significant reduced L/W ratio compared to cells expressing $Msb4_{22-81\Delta 35-39}$-Spa2 (Fig.5C). We conclude that the SHD1-Msb3/Msb4 interaction restricts vesicle fusion to the bud tip by recruiting the Sec4-GAP-activity to this site. Although the interaction between Ste7 and Spa2 is required to attach Ste7 to the shmoo tip (Fig. 4C, D), cells lacking this interaction arrest the cell cycle upon pheromone treatment like wildtype and not like *ste7Δ-*cells (Fig. 5D).

The FRM of human β-PIX binds the SHD1 of Spa2

GIT1/2 are the mammalian homologues of Spa2 (Zhao et al., 2000). Their SHD1s are preceded by ankyrin repeats and an ARF GAP domain. The SHD1 domain of GIT1/2 binds to β-PIX and recruits the Cdc42-GEF to focal adhesions (Zhao et al., 2000). The solved crystal structure of the complex shows how the hydrophobic groove of $SHD1_{GIT1}$ encloses a short helix of β-PIX (Zhu et al., 2020). Closer inspection of the β-PIX helix revealed the essential features of the yeast FRM (Fig. 6A). This observation encouraged us to test whether $FRM_{\beta-PIX}$ binds to SHD1 of yeast. Probed on a blot by bacterially expressed 6xHis-$SHD1_{Spa2}$, the overlay assay of the different GST-FRM fusion proteins demonstrates that $FRM_{\beta-PIX}$ binds to SHD1

as strongly as FRM_{Ste7} but less strongly than $FRM_{Msb3/4}$ (Fig. 6B). The N-terminal regions of Mkk1 or Mkk2 did not measurably interacted with 6xHis-SHD1 under these conditions (Fig. 6B).

Predicting SHD1-FRM complexes in other members of the Opisthokonta

The similarity between the predicted SHD1-FRM complex of yeast and the GIT1-β-PIX complex of mammals suggests, that SHD1s might generally bind sequence-conserved FRMs. Spa2 was discovered in polarisome-like structures of other yeasts and filamentous fungi (Knechtle et al., 2003; Jones and Sudbery, 2010). The architectures and compositions of these protein assemblies are not fully worked out and evidence for a physical interaction between Spa2 and Msb3/4 is only provided for the polarisome of *Fusarium graminearum* (FG), an ascomyceteous fungus that causes head blight in wheat (Zheng et al., 2021). The interaction between $Spa2_{FG}$ and $Msb3_{FG}$ were mapped to the SHD1 of $Spa2_{FG}$ and to residues 168-399 in the supposedly unstructured region of $Msb3_{FG}$ (Zheng et al., 2021). Supplied with the sequences of the SHD1 of $Spa2_{FG}$ and the sequence of full-length $Msb3_{FG}$, AlphaFold models with high confidence a short helix of $Msb3_{FG}$ into the groove of $Spa2_{FG}$ (Fig. 6C). The sequence of the helix matches the characteristic FRM of the budding yeast, and, by stretching from residues 362 to 365, falls exactly into the experimentally determined binding region of $Msb3_{FG}$ (Fig. 6A, C, G).

By using AlphaFold as a tool to predict homologous SHD1-FRM-complexes, we explored whether RabGAPs are integral polarisome components in other species as well. The SHD1 of the Spa2 homologue from *Neurospora crassa* (NC) was shown to bind to the Calponin-like protein CCP-1 (Zheng et al. 2020). The binding region of

Spa2$_{NC}$ was assigned to the hydrophobic groove of SHD1, whereas the corresponding binding site on CCP-1 remained unspecified. AlphaFold predicts with high confidence a complex between both proteins. The interface of this complex is formed by the hydrophobic groove of SHD1$_{NC}$ and a short helix from residue 497-507 of CCP-1 that matches the consensus of the FRM but carries a Gly instead of a hydrophobic residue in position +1 (Fig. 6A, D, G). Furthermore, indirect evidence suggest that Spa2$_{NC}$ might also interact with Gyp3$_{NC}$, the Msb3-homologue of *Neurospora crassa* (Callejas-Negrete and Castro-Longoria, 2019; Lichius et al., 2012). The predicted complex between SHD1$_{NC}$ and full length Gyp3$_{NC}$ confirms this suggestion. SHD1$_{NC}$ captures a short helix with strong sequence similarity to the FRM. (Fig. 6A, E, G).

Magnaporthe oryzae (MO), is a pathogenic, filamentous fungi that causes rice blast. The polarsiome of *Magnaporthe oryzae* was shown to form at the tip of the hyphae (Li et al., 2014). A complex between Spa2$_{MO}$ and Gyp3$_{MO}$ was not described, but is predicted by AlphaFold with high confidence. Again, the helix of Gyp3$_{MO}$ that fits into the groove of SHD1$_{MO}$ is situated in the unstructured N-terminal region of Gyp3$_{MO}$ and aligns well with the consensus sequence of the FRM (Fig. 6A, F, G).

The mammalian GIT1 displays an internal Rab-GAP activity at its N-terminus thus freeing the SHD1 from its obligation to bind an external GAP. Instead, SHD1$_{GIT1}$ binds besides β-PIX the scaffold protein Piccolo (Kim et al., 2003). The complex is part of the active zone below the presynaptic plasma membrane that might be considered to be functionally and structurally related to the cortical organization underlying the membrane of the yeast bud tip. The binding region of Piccolo from rat (RN) was experimentally localized within a fragment of 150 residues of the more than

5000 residue long protein (Kim et al., 2003). AlphaFold proposes a complex between SHD1$_{GIT1}$ and a peptide within the binding fragment of Piccolo (Fig. S3A). The sequence of this peptide folds into a helix upon contact with the SHD1 binding cleft and matches the FRM consensus sequence (Fig. 6A, S3A-D).

A proteome-wide search predicts Dse3 as novel interactor of Spa2

The relaxed stringency of the FRM's consensus sequence increases the likelihood of its frequent, random occurrence within the yeast proteome. To distinguish functional motifs that directly interact with the SHD1 domain of Spa2 from these occurrences, we implemented the following strategy:

1. Identify all sequences conforming to the pattern (V/I)-(I/L)-(D/S)-(L/A)-Y.
2. Discard motifs not located in unfolded, or helical conformations.
3. Exclude non-cytosolic proteins.
4. Exclude proteins that, according to AlphaFold, do not bind the hydrophobic SHD1 groove through the motif's helical conformation.
5. Exclude proteins as potential binding partners of SHD1 if the ipTM of the predicted complex does not scores equal to or higher than that of Mkk1.
6. Use the foldback competition assay to prove the motif-dependent binding of the candidate protein.

The strategy reduced the number of FRM containing proteins from 124 in Step1 to 95 in Step2, to nine in Step 4 (Fig. 7A). In Step 5, only the prediction of complexes between SHD1 and Msb3, Msb4, Ste7, or Dse3 sored equal or higher than the Mkk1-SHD1 complex (Fig. 7A, Table S4). Similar to the motif-containing regions of Msb3,

Msb4, Ste7 or Mkk1, Dse3$_{181-240}$ inhibited binding of N$_{ub}$-Msb3 to SHD1CRU in the Dse3$_{181-240}$-SHD1CRU fusion protein (Step 6, Fig. 7B, C, Table S4). The inhibition was strictly dependent on the presence of the FRM as N$_{ub}$-Msb3 interacted with Dse3$_{181-240\Delta208}$-212-SHD1CRU as strongly as with SHD1CRU alone (Fig 7C). The FRM-containing helix of Dse3, unlike the other yeast complexes, is predicted to adopt a flipped orientation within the binding groove. The Tyr residue at the +2 position of the motif forms a hydrogen bond with Asp62 of Spa2 (Fig. 7D, E).

Discussion

The N-terminal domain of Spa2 binds to a conserved motif in Msb3, Msb4, Ste7, and Mkk1. Complex formation requires the initially unstructured motifs to adopt an alpha-helical conformation upon binding to the SHD1 domain. AlphaFold predictions, supported by mutational analysis of potential contact sites in Msb3-SHD1, Msb4-SHD1, Ste7-SHD1 and Mkk1-SHD1 complexes, and *in vivo* competition assays among these FRMs, confirm this model. Independent validation comes from the solved structure of the mammalian GIT1 SHD1 domain bound to its partner, β-PIX (Zhu et al., 2020). The structures of the different complexes explain why the tyrosine at position +2 in the FRM is conserved across proteins and species. The tyrosine's phenyl group forms hydrophobic contacts with the SHD1 groove, while its hydroxyl group forms a hydrogen bond with the carboxyl group of Asp112 in yeast Spa2 or Asp348 in mammalian GIT1, orienting the helix within the groove. Substituting tyrosine with phenylalanine, which lacks the hydroxyl group, significantly reduce the stabilities of both complexes (Fig. 2C and (Zhu et al., 2020)).

Surprisingly, the FRM orientation within the SHD1 hydrophobic groove is reversed in the predicted complexes from *Neurospora crassa*, *Magnaporthe oryzae*, *Fusarium graminearum*, and the complex between SHD1 and Dse3. Here, the tyrosine at position +2 forms a hydrogen bond with an aspartate corresponding to Asp62 in *Saccharomyces cerevisiae*. Both Asp112 and Asp62 are conserved in the SHD1 domains of yeast Spa2 and mammalian GIT1/2, occupying equivalent positions in SDR repeats 1 and 2, respectively, thus reflecting the internal symmetry of the SHD1 fold. Using Asp112 or Asp62 as the hydrogen bond acceptor requires reshuffling hydrogen bonds within SHD1 and reversing the helix orientation in the groove. Further experimental validation is needed to distinguish between these alternative arrangements of the helices in the binding groove of SHD1.

Our findings indicate that an FRM in an unstructured protein region strongly suggests binding to a member of the Spa2/GIT1 family of cortical scaffold proteins. Using AlphaFold to identify potential binding sites, rather than relying solely on sequence alignments, enables detection of sites with partial matches to a not fully defined consensus sequence, and checks in addition the spatial accessibility of the sequence at the same time. However, the weak correlation between interaction strength (measured by the Split-Ub assay) and ipTM scores in complexes with single-residue substitutions underscores the limitations of AlphaFold for these applications. A low ipTM score may still reflect an FRM with functionally relevant, though weaker affinity. If the strength of interactions affects the accuracy of complex predictions, establishing computational criteria to differentiate non-functional FRMs from those engaged in weak but functional interactions would be highly beneficial. This is

especially crucial when translating insights from protein interaction networks in model organisms to species less amenable to experimental study.

Our assays demonstrate that Msb4, Msb3, Ste7, and Mkk1 bind Spa2 with decreasing affinity. These differences are physiologically relevant for interactions involving Spa2 with Msb3, Msb4, and Ste7. The stronger binding of Msb3 and Msb4 ensures their localization to the cell tip and bud neck during mitotic growth and mating, whereas Ste7's weaker affinity for SHD1 likely requires additional interactions with proteins like Ste5, which co-assemble at the cell tip exclusively during mating to stabilize Ste7 alongside Spa2 (Choi et al., 1994;˙Maeder et al., 2007). Similarly, Mkk1's polar distribution depends on Spa2, despite its interaction with SHD1 being near the detection limit of our assays (Van Drogen and Peter, 2002). Mkk1 stabilization at these sites may require Slt2, the MAP kinase in this pathway, which binds to both Mkk1 and Spa2 (Hruby et al., 2011; Breitkreutz et al., 2010).

Our homology search for functional FRMs in the yeast proteome identified Dse3 as a novel FRM-containing protein interacting with Spa2. *DSE3* is specifically transcribed in daughter cells (Colman-Lerner et al., 2001). Nothing is known about the function of the expressed protein. The functional relevance of its interaction with Spa2 thus remains unclear and requires further exploration. The fold-back competition assay's simplicity, sensitivity, and reliability make it an effective new tool for analyzing SLiMs, complementing existing methods (Subbanna et al., 2025). It enables efficient screening of previously overlooked candidates, such as Age2, Rpg1, Ybr200w-A,

and Mkk2, for direct FRM-dependent interactions with Spa2. Additionally, it supports revisiting homology searches with less stringent FRM consensus sequences.

Material and Methods

Growth conditions, cultivation of yeast strains, and genetic methods

All yeast strains were derivatives of JD47, a segregant from a cross of the strains YPH500 and BBY45. Yeast strains were cultivated in synthetic defined (SD) or yeast extract peptone dextrose (YPD) media at the indicated temperatures. Media preparation followed standard protocols (Glomb et al., 2020). SD medium for Split-Ub assays contained in addition 1 mg/ml 5-fluoro-orotic acid (5-FOA; Formedium). Gene deletions and promoter replacements by *PMET17* were performed by homologous integration of the cassettes derived by PCR from the plasmids pFA6a-hphNT1, pFA6a- natNT2, pFA6a-kanMX6, pFA6a-CmLEU2, or pYM-N35 (Bähler et al., 1998; Janke et al., 2004). *E. coli* XL1 blue cells were used for plasmid amplification and grown at 37°C in lysogeny broth (LB) medium containing antibiotics. Proteins were expressed in the *E. coli* strain BL21 DE3, grown in LB or super broth (SB) medium at 37°C.

Generation of plasmids and yeast strains

Detailed lists of all yeast strains and plasmids used in this study are provided in Table S1 and S2, sequence information on oligos is available upon request. Genomic deletions and modifications of the SHD1 or FRM motifs were generated by CRISPR/Cas9 manipulation using plasmid pML104 or pML107 containing specific 20mer guide RNA sequences, and template oligonucleotides for exchanging the

information on the genomic DNA (Laughery et al., 2015). The correct insertions of PCR fragments and mutations were verified by PCR amplification and sequencing.

Gene deletions were obtained by replacing the ORF through single-step homologous recombination with an antibiotic resistance cassette derived by PCR from the plasmids pFA6a-hphNT1, pFA6a- natNT2, pFA6a-kanMX6, pFA6a-CmLEU2, or pYM-N3(Janke et al., 2004) (Bähler et al., 1998). Genomic gene fusions to the GFP- or CRU-modules were obtained as described (Müller et al., 2024). The fusion of *GFP* or *CRU* to *SPA2*, *MSB3*, MSB4, Ste7, Mkk1, and derivates of those genes with deleted or modified FRM or SHD1 regions were constructed by PCR amplification of the respective C-terminal ORFs without stop codon from genomic DNA. The obtained DNA fragments were cloned via *Eag*I and *Sal*I restriction sites in front of the *CRU* or *GFP* module on pRS303, pRS304, or pRS306 vectors (Wittke et al., 1999). The plasmids were linearized using a single restriction site within the C-terminal genomic DNA sequence and transformed into yeast. Successful integration was verified by PCR of single yeast colonies with diagnostic primer combinations using a forward primer annealing in the target ORF but upstream of the linearization site, and a reverse primer annealing in the C-terminal module.

Fusions between the N-terminal regions of Msb4, or Ste7, and full length Spa2 were generated *via* CRISPR/Cas9-mediated insertion of a PCR-fragment encoding the respective regions of *MSB4* or *STE7* flanked by 45bp identical to the region upstream of the start codon of *SPA2* and the 5'-end of its open reading frame. The plasmid-borne fusions between SHD1 and the N-terminal regions of Msb4, Ste7, Mkk1 or Msb3 were

obtained by integrating the PCR products of the respective regions in front of SHD1 in the pMET17-Spa2(1-124)CRU313 vector.

Fragments of *SPA2, Msb3 or Msb4* were expressed as GST- or 6xHis- fusions in *E.coli* strains BL21 or BL21 Gold. Fragments for the GST-fusions were amplified from yeast genomic DNA using primers containing *Bam*HI/*Eco*RI restriction sites. The PCR products were fused in-frame behind GST on a pGex2T vector (GE Healthcare, Buckinghamshire, UK). 6xHis-tagged fragments were amplified from genomic DNA of wild type yeast or the strain lacking residues 76-93 of Msb3 using primers containing *Sfi*I restriction sites. The products were inserted into the pES plasmid, downstream and in frame of a 6xHis-tag. GST-FRM fusions of MSB3, MSB4, STE7, MKK1, MKK2 and β-PIX were obtained by inserting double stranded oligonucleotides harboring the sequences of the FRMs and containing matching *Sal*I/EcoRI restriction sites at their ends in-frame behind the ORF of GST on a pGEX2T vector.

In vivo Split-Ub interaction analysis

Large-scale Split-Ub assays were performed as described (Hruby et al., 2011). A library of 540 different α-strains, each expressing a different N_{ub} fusion, was mated with a *PMET17SHD1*-CRU–expressing a-strain. Diploids were transferred as independent quadruplets on SD media lacking methionine and containing 1 mg/ml 5-FOA, and different concentrations of copper sulfate to adjust the expression of the N_{ub} fusions. For small-scale interaction analysis, a- and α-strains expressing N_{ub} or C_{ub} fusion constructs were mated. The diploid cells were spotted onto SD-FOA medium in four 10-fold serial dilutions starting from $OD_{600} = 1$. Growth was recorded at 30°C every day for 2–5 d.

Binding assay

All incubation steps were carried out under rotation at 4°C. GST or GST-tagged proteins were immobilized from *E. coli* extracts on 50 μl Glutathione–Sepharose beads in PBS (GE Healthcare). After incubation for 1 hour at 4°C, with either *E. coli* extracts or purified proteins, the beads were washed three times, the bound material was eluted with GST elution buffer (50 mM Tris, 20 mM reduced glutathione), and subjected to SDS-PAGE followed by Ponceau S staining and western blot analysis using anti-His (H1029), or anti-GST (G1160) antibodies (Sigma-Aldrich). To prevent re-binding of detached 6×His-SDH1 to immobilized GST fusions during antibody incubation (Fig.1I), Ponceau-stained membranes were cut horizontally between the ~16 kDa and ~30 kDa regions immediately after transfer. The resulting strips were processed in parallel in separate trays with identical solutions and reassembled for chemiluminescent detection.

Far-western blot

Enriched GST and GST-FRM-fusions were transferred onto nitrocellulose after SDS-PAGE. The membrane was blocked for 1 h in 20mM Tris, 150 NaCl, 0.1% Tween 20 (TBST) 2% milk powder (w/v), washed for 5 min in 10 ml TBST, and incubated for 1 h at 4°C with enriched 6xHis-SHD1. After washing three times for 5 min with 10 ml TBST, the membrane was sequentially incubated at room temperature for 1 h with anti-His (H1029) and conjugated anti-mouse antibodies (A4416; Sigma-Aldrich) (Wu et al., 2007). The source data of the western and far-western blots are shown in Fig. S4.

Halo Assay

Exponentially growing yeast strains were diluted to OD_{600} = 0.5 and 300 µL of the suspension were plated evenly on SD-full agar plates. Round filter papers with a diameter of 5 mm were placed on the cell lawn and 10 µL of alpha-factor was applied to the filter papers in serial dilutions of 10 µg, 5 µg, 2.5 µg, and 1.25 µg. Plates were incubated overnight (o/n) at 30°C. Images were taken on day one and two. Sensitivity to mating factor was determined by measuring the halo of growth inhibition. All strains were measured four times, and the values were normalised to the wild type.

Fluorescence microscopy

Yeast cultures were grown overnight in SD medium, diluted in 3–4 ml fresh SD medium, and grown for 2–3 h at 30°C to mid-log phase. Microscopic observations were performed with the Axio Observer spinning disc confocal microscope (Zeiss) equipped with an Evolve512 electron-multiplying charge-coupled device camera (Photometrics), a Plan-Apochromat 63×/1.4 oil differential interference contrast (DIC) objective, and a 100×/1.4 oil DIC objective. Fluorescence was excited with 488- diode lasers (Zeiss) and detected with a high-efficiency filter set 38 (GFP). Time-lapse experiments were carried out at RT over the course of 4h. Operations were performed with the ZEN2.6 (2012) software package (Zeiss). Exposure times were adjusted to the respective GFP -labeled proteins to reduce bleaching and phototoxicity (Müller et al., 2024).

Standard time-lapse experiments were carried out with the 63× objective in a Sarstedt 1-well on cover glass II incubation chamber. 100 µl culture was pipetted onto the glass bottom of the chamber and covered with a slice of standard solid SD medium. Images from a series of 14 z-slices over a total range of 2.43 µm were taken in intervals of 3

min. Special glass slides with a ground indentation in the centre were used for monitoring shmoo formation of alpha-factor-treated yeast cells (custom-made by the University's glassblowing workshop). α-mating factor was added to a concentration of 10 µg/mL to 500 µL in 2x SD full media, mixed 1:1 with 3.4% agarose (heated until liquid), and immediately poured into the indentation. A coverslip was placed on the top to remove excess agarose and to match the surface of the glass. After cooling, the coverslip was removed, 3.6 µL of cell suspension were applied, covered with a new coverslip and sealed with parafilm to prevent evaporation and drying of the agarose patch. The slide was then incubated at 30°C for 15 minutes to allow the cells to settle.

Analysis of microscopic data

Image analysis and signal quantifications were carried out with the open source software platform Fiji (ImageJ) (Schneider et al., 2012; Schindelin et al., 2012). Plot profiles of fluorescence intensities were performed in FIJI to detect differences in GFP distributions across the cell. Circular regions of interest (ROI) of the same size were placed at the shmoo tips of cells to quantify the GFP intensity relative to signal of the cell body. The background signal (I[background]) was measured with ROIs placed outside but in proximity of the cell and subtracted from the intracellular signals. The relative GFP intensities at the shmoo tip were then calculated:

$$\text{relative fluorescence intensity} = \frac{\text{I[shmoo tip]} - \text{I[background]}}{\text{I[cytoplasm]} - \text{I[background]}}$$

L/W ratio determination

Yeast strains were grown overnight in SD medium and diluted in YPD to measure cell- and bud sizes. Images were taken on the Zeiss Observer Z1 using the Axio camera at

100x magnification with 14 z-layers over 3 µm in the DIC channel. Cells were randomly selected, and strain identities were blinded to avoid bias. The images were analysed in ImageJ (FIJI) after z-projection. To determine the bud morphology, the length (L) and width (W) of the buds were measured. The region of interest (ROI) for length was chosen along the axis of growth, and the ROI for width was set perpendicular to the length at the widest point of the bud. Only non-blurred, living cells with visibly attached buds were measured and strain names were kept blinded for evaluation. All values were transferred to Excel, sorted by length and the L/W ratio was calculated. Each strain was measured in triplicate, with a minimum of 150 cells measured. Buds smaller than 1.5 µm were excluded from the analysis.

Transmission electron microscopy

Exponentially grown yeast cells were harvested and vitrified through high-pressure freezing on carbon coated sapphire discs in a Wohlwend HPF compact 01 high-pressure freezer. Freeze-substitution was performed with 0.2% (v/v) osmium tetroxide, 0.1% (w/v) uranyl acetate and 5% (v/v) H_2O in acetone by gradually increasing the temperature over 17 hours from -90 °C to 0 °C. Temperature was kept at 0 °C for one hour and then increased within one hour to room temperature. Samples were washed with acetone, followed by embedding in Epon 812 in four consecutive steps at room temperature: 33% Epon 812/67% acetone for 1 hour, followed by 67% Epon 812/33% acetone for 3 hours, 90% Epon 812/10% acetone overnight, and pure Epon 812 overnight (v/v). Samples were transferred into fresh Epon 812 and polymerized at 60 °C for 72 hours. Embedded cells were sectioned in ~70 nm steps with a Leica EM

UC7 ultramicrotome. Sections were captured on a formvar film on glow-discharged copper grids. Images were acquired with a 120 kV Jeol JEM-1400 Transmission electron microscope and a Veleta CCD camera (Olympus).

Statistical evaluation

All statistical evaluations were based on data sets of at least three independent experiments. Depending on the result of the normality test, parametric (ANOVA) or non-parametric (Kruskal-Wallis) tests were chosen. The Kruskal-Wallis test was followed by a Dunn's Test for multiple comparisons. The p values and the sample size (n) are given in the caption or in the supplementary data. Unless otherwise stated, the red bars represent the mean and standard deviation (s.d.) of each data set, the p values are given in GP standard values.

Structure prediction

Structure prediction was performed with the AlphaFold 3 (Abramson et al., 2024). The models were ranked by their pTM/ipTM scores, and the model with the highest score is shown. The PAE scores were obtained from the AlphaFold3 data by the PAE viewer tool (Elfmann and Stülke, 2023). Processing and visualization of the predicted models were performed with ChimeraX (Meng et al., 2023).

Proteome-Wide Motif Search

To identify amino acid sequences in Saccharomyces cerevisiae matching the motif (V/I)-(I/L)-(D/S)-(L/A)-Y, we analyzed the proteome-wide structure prediction dataset (n = 6,039) from the AlphaFold Protein Structure Database (version 4; available at: https://ftp.ebi.ac.uk/pub/databases/alphafold/v4/UP000002311_559292_YEAST_v4.t

ar). A custom Bash script iteratively executed PyMOL (http://www.pymol.org/pymol.) (Schrödinger and DeLano, 2020) to scan each protein structure for the motif and document the secondary structure assignments of the corresponding five residues. Protein identifiers containing the motif were compiled into a comprehensive list of all matches. A separate list was generated for matches where the motif was exclusively located in residues annotated by PyMOL as helical or loop regions (including cases with both annotations).

Use of artificial intelligence tools

Grok3 was used during manuscript preparation to provide suggestions for translation, grammar, and stylistic improvements.

Data and Resource Availability

All strains and plasmids are available upon request.

Acknowledgement

We thank C.M. Pfeiffer for helpful advice.

Funding

The work was funded by grants from the Deutsche Forschungsgemeinschaft (DFG) to N.J. (Jo 187/8-1).

Competing interest

The authors declare no competing financial interests.

Author contributions

Conceptualization: N. Johnsson. Investigation: L. Bareis, A. Siewert, S. Timmermann, N. Schmid, T. Bergner, Benjamin Grupp. Writing - original draft: N. Johnsson; Writing - review & editing: N. Johnsson, L. Bareis. Supervision: N. Johnsson, C Read. Funding acquisition: N. Johnsson.

References

Abramson, J., Adler, J., Dunger, J., Evans, R., Green, T., Pritzel, A., Ronneberger, O., Willmore, L., Ballard, A. J., Bambrick, J., et al. (2024). Accurate structure prediction of biomolecular interactions with AlphaFold 3. *Nature* **630**, 493–500.

Bähler, J., Wu, J. Q., Longtine, M. S., Shah, N. G., McKenzie, A., Steever, A. B., Wach, A., Philippsen, P. and Pringle, J. R. (1998). Heterologous modules for efficient and versatile PCR-based gene targeting in Schizosaccharomyces pombe. *Yeast* **14**, 943–951.

Breitkreutz, A., Choi, H., Sharom, J. R., Boucher, L., Neduva, V., Larsen, B., Lin, Z. Y., Breitkreutz, B. J., Stark, C., Liu, G., et al. (2010). A global protein kinase and phosphatase interaction network in yeast. *Science* **328**, 1043–1046.

Callejas-Negrete, O. A. and Castro-Longoria, E. (2019). The role of GYP-3 in cellular morphogenesis of Neurospora crassa: Analyzing its relationship with the polarisome. *Fungal Genetics and Biology* **128**, 49–59.

Choi, K.-Y., Satterberg, B., Lyons, D. M. and Elion, E. A. (1994). Ste5 Tethers Multiple Protein Kinases in the MAP Kinase Cascade Required for Mating in S. cerevisiae. *Cell* **78**, 499-512

Colman-Lerner, A., Chin, T. E. and Brent, R. (2001). Yeast Cbk1 and Mob2 Activate Daughter-Specific Genetic Programs to Induce Asymmetric Cell Fates. *Cell* **107**, 739-750

Courchesne, W. E., Kunisawa, R. and Thorner, J. (1989). A putative protein kinase overcomes pheromone-induced arrest of cell cycling in S. cerevisiae. *Cell* **58**, 1107–1119.

Dünkler, A., Leda, M., Kromer, J. M., Neller, J., Gronemeyer, T., Goryachev, A. B. and Johnsson, N. (2021). Type V myosin focuses the polarisome and shapes the tip of yeast cells. *Journal of Cell Biology* **220**,

Elfmann, C. and Stülke, J. (2023). PAE viewer: A webserver for the interactive visualization of the predicted aligned error for multimer structure predictions and crosslinks. *Nucleic Acids Res* **51**, W404–W410.

Gao, X. D., Albert, S., Tcheperegine, S. E., Burd, C. G., Gallwitz, D. and Bi, E. (2003). The GAP activity of Msb3p and Msb4p for the Rab GTPase Sec4p is required for efficient exocytosis and actin organization. *Journal of Cell Biology* **162**, 635–646.

Gingras, R. M., Sulpizio, A. M., Park, J. and Bretscher, A. (2022). High-resolution secretory timeline from vesicle formation at the Golgi to fusion at the plasma membrane in S. cerevisiae. *Elife* **11**,.

Glomb, O., Bareis, L. and Johnsson, N. (2019). YFR016c/Aip5 is part of an actin nucleation complex in yeast. *Biol Open*. **8**,

Glomb, O., Wu, Y., Rieger, L., Rüthnick, D., Mulaw, M. A. and Johnsson, N. (2020). The cell polarity proteins Boi1 and Boi2 direct an actin nucleation complex to sites of exocytosis in Saccharomyces cerevisiae. *J Cell Sci* **133**,

Guo, W., Roth, D., Walch-Solimena, C. and Novick, P. (1999). *The exocyst is an effector for Sec4p, targeting secretory vesicles to sites of exocytosis. The EMBO J.* **18,** 1071-1080

Holehouse, A. S. and Kragelund, B. B. (2024). The molecular basis for cellular function of intrinsically disordered protein regions. *Nat Rev Mol Cell Biol* **25**, 187–211.

Hruby, A., Zapatka, M., Heucke, S., Rieger, L., Wu, Y., Nussbaumer, U., Timmermann, S., Dünkler, A. and Johnsson, N. (2011). A constraint network of interactions: protein-protein interaction analysis of the yeast type II phosphatase Ptc1p and its adaptor protein Nbp2p. *J Cell Sci* **124**, 1603.

Janke, C., Magiera, M. M., Rathfelder, N., Taxis, C., Reber, S., Maekawa, H., Moreno-Borchart, A., Doenges, G., Schwob, E., Schiebel, E., et al. (2004). A versatile toolbox for PCR-based tagging of yeast genes: New fluorescent proteins, more markers and promoter substitution cassettes. *Yeast* **21**, 947–962.

Jin, Y., Sultana, A., Gandhi, P., Franklin, E., Hamamoto, S., Khan, A. R., Munson, M., Schekman, R. and Weisman, L. S. (2011). Myosin V transports secretory vesicles via a Rab GTPase cascade and interaction with the exocyst complex. *Dev Cell* **21**, 1156–1170.

Jones, L. A. and Sudbery, P. E. (2010). Spitzenkörper, exocyst, and polarisome components in Candida albicans hyphae show different patterns of localization and have distinct dynamic properties. *Eukaryot Cell* **9**, 1455–1465.

Kim, S., Ko, J., Shin, H., Lee, J. R., Lim, C., Han, J. H., Altrock, W. D., Garner, C. C., Gundelfinger, E. D., Premont, R. T., et al. (2003). The GIT family of proteins forms multimers and associates with the presynaptic cytomatrix protein piccolo. *Journal of Biological Chemistry* **278**, 6291–6300.

Knechtle, P., Dietrich, F. and Philippsen, P. (2003). Maximal polar growth potential depends on the polarisome component AgSpa2 in the filamentous fungus Ashbya gossypii. *Mol Biol Cell* **14**, 4140–4154.

Laughery, M. F., Hunter, T., Brown, A., Hoopes, J., Ostbye, T., Shumaker, T. and Wyrick, J. J. (2015). New vectors for simple and streamlined CRISPR-Cas9 genome editing in Saccharomyces cerevisiae. *Yeast* **32**, 711–720.

Lawson, M. J., Drawert, B., PetzoldID, L. and YiID, T.-M. (2022). A positive feedback loop involving the Spa2 SHD domain contributes to focal polarization. *PLoS One.* **17**

Li, C., Yang, Jun, Zhou, W., Chen, X.-L., Huang, J.-G., Cheng, Z.-H., Zhao, W.-S., Zhang, Y. and Peng, Y.-L. (2014). A spindle pole antigen gene MoSPA2 is important for polar cell growth of vegetative hyphae and conidia, but is dispensable for pathogenicity in Magnaporthe oryzae. *Curr Genet* **60**, 255–263.

Lichius, A., Yáñez-Gutiérrez, M. E., Read, N. D. and Castro-Longoria, E. (2012). Comparative live-cell imaging analyses of SPA-2, BUD-6 and BNI-1 in Neurospora crassa reveal novel features of the filamentous fungal polarisome. *PLoS One* **7**,.

Maeder, C. I., Hink, M. A., Kinkhabwala, A., Mayr, R., Bastiaens, P. I. H. and Knop, M. (2007). Spatial regulation of Fus3 MAP kinase activity through a reaction-diffusion mechanism in yeast pheromone signalling. *Nat Cell Biol* **9**, 1319–1326.

Meng, E. C., Goddard, T. D., Pettersen, E. F., Couch, G. S., Pearson, Z. J., Morris, J. H. and Ferrin, T. E. (2023). UCSF ChimeraX: Tools for structure building and analysis. *Protein Science* **32**,.

Mughal, F. and Caetano-Anollés, G. (2025). Evolution of intrinsic disorder in the structural domains of viral and cellular proteomes. *Sci Rep* **15**, 2878.

Müller, J., Furlan, M., Settele, D., Grupp, B. and Johnsson, N. (2024). Transient septin sumoylation steers a Fir1-Skt5 protein complex between the split septin ring. *Journal of Cell Biology* **223**,.

Neduva, V., Linding, R., Su-Angrand, I., Stark, A., De Masi, F., Gibson, T. J., Lewis, J., Serrano, L. and Russell, R. B. (2005). Systematic discovery of new recognition peptides mediating protein interaction networks. *PLoS Biol* **3**, 1–10.

Neller, J., Dünkler, A., Rösler, R. and Johnsson, N. (2015). A protein complex containing Epo1p anchors the cortical endoplasmic reticulum to the yeast bud tip. *Journal of Cell Biology* **208**, 71–87.

Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B., et al. (2012). Fiji: An open-source platform for biological-image analysis. *Nat Methods* **9**, 676–682.

Schneider, C. A., Rasband, W. S. and Eliceiri, K. W. (2012). NIH Image to ImageJ: 25 years of image analysis. *Nat Methods* **9**, 671–675.

Schrödinger, L. and DeLano, W. (2020). "PyMOL." http://www.pymol.org/pymol

Sheu, Y.-J., Santos, B., Fortin, N., Costigan, C. and Snyder, M. (1998). Spa2p Interacts with Cell Polarity Proteins and Signaling Components Involved in Yeast Cell Morphogenesis. *Mol Cell Biol.* **18**, 4053-4069

Sheu, Y.-J., Barral, Y. and Snyder, M. (2000). *Polarized Growth Controls Cell Shape and Bipolar Bud Site Selection in Saccharomyces cerevisiae. Mol Cell Biol.* **20,** 5235-5247

Snyder, M. (1989). The SPA2 Protein of Yeast Localizes to Sites of Cell Growth. *Journal of Cell Biology* **108**, 1419–1429.

Subbanna, M. S., Winters, M. J., Örd, M., Davey, N. E. and Pryciak, P. M. (2025). A quantitative intracellular peptide-binding assay reveals recognition determinants and context dependence of short linear motifs. *Journal of Biological Chemistry* **301**,.

Tcheperegine, S. E., Gao, X.-D. and Bi, E. (2005). Regulation of Cell Polarity by Interactions of Msb3 and Msb4 with Cdc42 and Polarisome Components. *Mol Cell Biol* **25**, 8567–8580.

Teague, M. A., Chalefft, D. T. and Errede, B. (1986). Nucleotide sequence of the yeast regulatory gene STE7 predicts a protein homologous to protein kinases. *Proc.Natl.Acad.Sci.USA* **83**, 7371–7375.

Tompa, P., Davey, N. E., Gibson, T. J. and Babu, M. M. (2014). A Million peptide motifs for the molecular biologist. *Mol Cell* **55**, 161–169.

Van Drogen, F. and Peter, M. (2002). Spa2p Functions as a Scaffold-like Protein to Recruit the Mpk1p MAP Kinase Module to Sites of Polarized Growth. *Current Biol.* **12**, 1698-1703

Valtz, N., Herskowitz,I. (1996). Pea2 Protein of Yeast Is Localized to Sites of Polarized Growth and Is Required for Efficient Mating and Bipolar Budding. *Journal of Cell Biology* **135,** 725-739

Wittke, S., Lewke, N. and Johnsson, N. (1999). Probing the Molecular Environment of Membrane Proteins In Vivo. *Mol Biol Cell* **10**, 2519–2530.

Wu, Y., Li, Q. and Chen, X. Z. (2007). Detecting protein-protein interactions by far western blotting. *Nat Protoc* **2**, 3278–3284.**Zhao, Z.-S., Manser, E., Loo, T.-H.**

and Lim, L. (2000). Coupling of PAK-Interacting Exchange Factor PIX to GIT1 Promotes Focal Complex Disassembly. *Mol. Cell Biol*. **20**, 6354-6363.

Zheng, P., Nguyen, T. A., Wong, J. Y., Lee, M., Nguyen, T. A., Fan, J. S., Yang, D. and Jedd, G. (2020). Spitzenkörper assembly mechanisms reveal conserved features of fungal and metazoan polarity scaffolds. *Nat Commun* **11**,.

Zheng, H., Li, L., Yu, Z., Yuan, Y., Zheng, Q., Xie, Q., Li, G., Abubakar, Y. S., Zhou, J., Wang, Z., et al. (2021). FgSpa2 recruits FgMsb3, a Rab8 GAP, to the polarisome to regulate polarized trafficking, growth and pathogenicity in Fusarium graminearum. *New Phytologist* **229**, 1665–1683.

Zhu, J., Zhou, Q., Xia, Y., Lin, L., Li, J., Peng, M., Zhang, R. and Zhang, M. (2020). GIT/PIX Condensates Are Modular and Ideal for Distinct Compartmentalized Cell Signaling. *Mol Cell* **79**, 782-796.

Legends to the Figures

Figure 1:

(A) Cut-outs of a Split-Ub array of diploid yeast cells each expressing SHD1CRU (CRU: C-terminal half of Ubiquitin (C_{ub})-R-Ura3) together with a different N_{ub} fusion (N_{ub}: N-terminal half of Ubiquitin). Identities of the N_{ub} fusions are given next to the respective cut outs. Mated N_{ub}- and C_{ub} expressing cells were arrayed as quadruplets on media containing 5-FOA. Colony growth indicates interaction between the fusion proteins. (B) Sequence alignment of putative binding sites of interaction partners of SHD1. FRM= Five residue motif. (C) Split-Ub assay of cells co-expressing SHD1CRU with N_{ub}-fusions to SHD1-binding partners, their mutants lacking the respective FRM, or to Guk1 as negative control. Cells were grown to OD_{600}=1 and 4 µl, or 4 µl of a 10-fold serial dilution were spotted on media containing 5-FOA and selecting for the presence of the N_{ub}- and C_{ub} fusions. (D) As in (C) but with cells co-expressing Ste5CRU and N_{ub}-fusions to Ste7, Ste11 and their mutants. (E) Split-Ub assay of cells co-expressing CRU-fusions to Msb3, Msb4 or their mutants lacking the FRM together with N_{ub}-Spa2, N_{ub}-Pea2, or N_{ub}-Sec4. Shown are four independent transformants each, directly transferred from selective media onto 5-FOA media. (F) As in (C) but with cells co-expressing Msb3CRU together with N_{ub} fusions to Spa2 and mutant of Spa2 lacking the indicated regions of the protein. (G) Enriched 6xHis-Msb3$_{1-150}$ (lanes 1-2) or 6xHis-Msb3$_{1-150\Delta76-93}$ (lanes 3, 4) were incubated with GST-SHD1 (lanes 1, 3) or GST-coupled beads (lanes 2, 4). Glutathione-eluates were separated by SDS-PAGE, transferred onto nitrocellulose, and stained with anti-His antibody (lanes 1-4) (Fig. S1A). (H) As in (C) but with cells co-expressing SHD1CRU together with N_{ub} fusions to Ms3, Msb4, Ste7 or to their

peptides carrying the FRMs in central positions. (I) Enriched 6xHis-SHD1 (lanes 1-3) was incubated with bead-coupled GST-Msb3$_{71-89}$, GST-Msb4$_{22-49}$ or GST (lanes 1, 2, 3). Glutathione-eluates were separated by SDS-PAGE, transferred onto nitrocellulose, and stained with anti-His antibody (lanes 1-3) (Fig. S1B). The arrow indicates the position where the membrane was horizontally cut before antibody incubation to prevent rebinding of the detached 6xHis-SHD1to the immobilized GST-Msb3$_{71-89}$ or GST-Msb4$_{22-49}$. Split-Ub experiments were performed twice, pulldown analysis was performed four (1G) or three (1I) times.

Figure 2:

(A) Upper and lower panel: AlphaFold prediction of the Msb3-SHD1 complex. The FRM containing helix (grey) of Msb3 docks into the binding pocked of SHD1 (ochre). Y88 and D86 of Msb3 are indicated in red and D112 and K97 of Spa2 in green. The PAE scores of the predictions of the complexes between SHD1 and Msb3, Msb4, Ste7, and Mkk1 and the structures of the respective FRM-containing helices are shown in Fig. S2A-F. (B) Alignment of the sequences around the FRM of the Msb3 homologues from different yeast species. (C) Upper panel: Split-Ub analysis as in Fig.1C but with cells co-expressing SHD1CRU together with N$_{ub}$ fusions of Msb3 or its mutants containing the indicated residue exchanges. Lower panel: Correlation between the ipTM scores of the predicted Msb3-SHD1 complexes carrying different residue exchanges in the FRM, and the estimated interaction strengths according to the Split-Ub assay. (D) Split-Ub analysis as in Fig.1C but with cells co-expressing Shd1CRU together with N$_{ub}$ fusions of Msb4, mutants of Msb4 containing the indicated residue exchanges, and Ste7. (E) Upper panel: Split-Ub analysis as in

Fig.1C but with cells co-expressing N_{ub} fusions to Msb3, Msb4 or Ste7 together with

SHD1CRU or mutants of SHD1, carrying the indicated residue exchanges. Lower

panel: Structural model of SHD1 by AlphaFold, highlighting the positions of the

residues that form the hydrophobic patch of the binding groove. Split-Ub analysis

was perfomed twice (2C, D), or four times (2E).

Figure 3

(A) A novel foldback competition assay shows that Mkk1 binds Spa2. Left panel:

AlphaFold-predicted structure of SHD1(blue) extended at its N-terminus by residues

22-81 of Msb4 (ochre). Right panel: Foldback competition assay: Spa2 or its SHD1

domain (blue) are artificially extended at their N-termini by an FRM-containing amino

acid stretch (ochre) and at their C-termini by the C_{ub}RUra3 module (red). The FRM

occupies intramolecularly the binding pocket of SHD1. As a consequence, an

additionally expressed N_{ub}-fusion to an FRM-containing protein (green) will not bind

to SHD1, N_{ub} and C_{ub} will not reconstitute the native-like ubiquitin, and RUra3 will not

be cleaved from C_{ub}. The cells remain 5-FOA sensitive. Introducing an interaction-

interfering mutations (FRM*) or deleting the FRM in the Spa2/SHD1 fusion protein

will promote the intermolecular binding of the N_{ub}-FRM fusion and the subsequent

cleavage and degradation of the RUra3 module. The cell become 5-FOA resistant

and grow in media containing 5-FOA. (B) Split-Ub assay as in Fig. 1E but with cells

co-expressing Spa2CRU extended at its N-terminus by the N-terminal regions of

Msb4, Msb4$_{\Delta35-39}$, or Ste7 together with N_{ub} fusions of the indicated proteins. (C)

Split-Ub analysis as in Fig.1C but with cells co-expressing N_{ub}-Msb3 together with

SHD1CRU extended at its N-terminus by the FRM-containing regions of Msb4,

Msb3, Ste7, or Mkk1. (D) Split-Ub analysis as in Fig.3D but with cells co-expressing

SHD1CRU together with N_{ub} fusions of Mkk1, Mkk1$_{\Delta171\text{-}175}$, or its mutants containing

the indicated residue exchanges on media containing 10mM methionine and 150 µm

copper. (E) Split-Ub analysis as in (C) but with cells co-expressing SHD1CRU

together with N_{ub} fusions of Mkk1, or Mkk1$_{\Delta171\text{-}175}$. Split-Ub analyses in (B)–(D) were

performed twice.

Figure 4

(A) Fluorescence microscopy of cells expressing Msb3-GFP (upper left panel) or

Msb3$_{\Delta76\text{-}93}$-GFP (lower left panel) after treatment with mating factor. Red lines

indicate sites of the fluorescence intensity measurements. Right panels:

Fluorescence intensity profiles across the indicated cells. Green bars indicate the

borders of the cell. Shmoo-tip ends at the right bar. (B) As in (A) but with cells

expressing Msb4-GFP, or Msb4$_{\Delta27\text{-}44}$-GFP. (C) As in (A) but with cells expressing

Ste7-GFP, or Ste7$_{\Delta119\text{-}123}$-GFP. (D) Shmoo-tip to cytosol fluorescence intensity ratios

of cells expressing GFP fusions to Msb3 (n=10), Msb3$_{\Delta76\text{-}93}$ (n=10), Msb4 (n=10),

Msb4$_{\Delta27\text{-}44}$ (n=10), Ste7 (n=10) or Ste7$_{\Delta119\text{-}123}$ (n=10) (Table S3). Whiskers of the box

plots defined by the Tukey method. A-C, E representative images from at least two

independent clones.

Figure 5

(A) Electron microscopical images of buds of cells of the indicated genotypes. Red

arrowheads point at post-Golgi vesicles. (B) Quantification of 50-70 nm vesicles in 30

buds each of wt-, *spa2Δ, spa2$_{\Delta2\text{-}150}$-, msb3$_{\Delta76\text{-}93}$ msb4Δ-, msb3Δ msb4$_{\Delta27\text{-}44}$-,* and

msb3Δ msb4Δ-cells. Whiskers of the box plots defined by the Tukey method. (C) Length/width ratios of buds of cells of wt (n=400), *spa2Δ* (n=240)*, spa2$_{L48,51}$* (n=224)*, msb3Δ msb4Δ* (n=353)*, msb3$_{Δ76-93}$ msb4Δ* (n=342)*, msb3Δ msb4$_{Δ27-44}$* (n=291), *spa2$_{Δ2-150}$* (n=264)*, ste7$_{105-165}$spa2* (n=168)*, msb4$_{22-81}$spa2* (n=149), *msb4$_{22-81Δ35-39}$spa2* (n=233). The value for wt-cells is shown twice for clarity. (D) Diameters of zones of growth inhibition after central application of alpha-factor on a lawn of cells of the indicated genotypes (n=4; error bars=s.d.). EM images and quantifications (5A, B) from one clone each. Analysis in 5C from two independent clones each.

Figure 6

(A) Sequence alignment of the FRMs from different members of the *Ophistokonta*. (B) Extracts of *E.coli* cells expressing GST (lane 1), GST-Msb3$_{71-98}$ (lane 2), GST-Msb4$_{22-49}$ (lane 3), GST-Ste7$_{105-135}$ (lane 4), GST-Mkk1$_{160-185}$ (lane 5), GST-Mkk2$_{153-178}$ (lane 6), and GST-β-PIX $_{703-727}$ (lane 7), were separated by SDS PAGE, transferred on nitrocellulose, stained with Ponceau S (upper panel) and then incubated with enriched 6xHis-SHD1 before being incubated with anti-His and peroxidase-coupled secondary antibody (lower panel). Experiment performed three times. (C-F) Predicted Aligned Error (PAE) matrices from the AlphaFold predictions of the complexes between the SHD1s of *Fusarium graminearum* (C), *Neurospora crassa* (D, E), and *Magnaporthe oryzae* (F) with the indicated binding partners. Blue arrows points to the sequences around the FRMs that display a high score of interaction with the SHD1. (G) Predicted structures of the FRM-containing sequences of the complexes of C-F.

Figure 7

(A) List of the FRM-containing yeast proteins displaying the highest ipTM scores for the AlphaFold prediction of their complexes with the SHD1 of Spa2. (B) Sequence alignment of the FRMs of Msb3 and Dse3. (C) Foldback competition analysis as in Fig.1C but with cells co-expressing N_{ub}-Msb3 together with SHD1CRU extended at its N-terminus by the FRM-containing regions of Msb3 ($Msb3_{71-130}$-SHD1CRU), Dse3 ($Dse3_{181-240}$-SHD1CRU), or its mutant lacking the FRM ($Dse3_{181-240\Delta208-212}$-SHD1CRU) (performed twice, with two independent clones each). (D) Predicted Aligned Error (PAE) matrix from the AlphaFold prediction of the complex between the SHD1 of Spa2 and Dse3. (E) Structural model of the interaction between the FRM-containing helix of Dse3 (grey) and SHD1 (ochre) derived from the AlphaFold prediction of the complex. E207 and Y212 of Dse3 highlighted in red; D62 and K97 of Spa2 highlighted in green. Split-Ub analysis (C) was performed twice.

A

N$_{ub}$‑

Msb4	Yel023c
Msb3	Bud2
Ste7	Bud3

SHD1CRU

SD 5-FOA

B

FRM

Msb3	80	G N L S V I D L Y G D E V
Msb4	31	A N Y S V L D L Y D D D D
Ste7	115	S T P C I I D A Y S N N F
Mkk1	167	D F S N L I S A Y E Q S S
Mkk2	160	Y P S T L L S A Y D F S S

-2 0 +2

C

SHD1CRU

N$_{ub}$‑

Msb3
Msb3$_{\Delta 84-88}$
Msb4
Msb4$_{\Delta 35-39}$
Ste7
Ste7$_{\Delta 119-123}$
Guk1
-

SD 5-FOA

D

Ste5CRU

N$_{ub}$‑

Ste11
Ste11$_{106-end}$
Ste11$_{411-end}$
Ste7
Ste7$_{\Delta 119-123}$
Ste7$_{165-end}$
Guk1

SD 5-FOA

E

N$_{ub}$‑

	Spa2	Pea2	Sec4
Msb3CRU			
Msb3CRU$_{\Delta 76-93}$			
Msb4CRU			
Msb4CRU$_{\Delta 27-44}$			

SD 5-FOA

F

Msb3CRU

N$_{ub}$‑

Spa2
Spa2$_{\Delta 2-188}$
Spa2$_{\Delta 2-150}$
Spa2$_{\Delta 168-188}$
-
Guk1

SD 5-FOA

G

GST-SHD1	+	-	+	-
GST	-	+	-	+
6xHis-Msb3$_{1-150}$	+	+	-	-
6xHis-Msb3$_{1-150\ \Delta 76-93}$	-	-	+	+

kDa

45 —

27 —
                                        — 6xHis-Msb3$_{1-150}$
15 —

1   2   3   4

anti-His

H

SHD1CRU

N$_{ub}$‑

Msb3
Msb3$_{71-100}$
Msb4
Msb4$_{21-50}$
Ste7
Ste7$_{105-135}$
Guk1
-

SD 5-FOA

I

GST-Msb3$_{71-98}$	+	-	-
GST-Msb4$_{22-49}$	-	+	-
GST	-	-	+
6xHis-SHD1	+	+	+

kDa

27 —
                                        ←
17 —   — 6His-SHD1

1   2   3

anti-His

A

Y-D

D-K

FRM-Msb3

SHD1

SHD1

FRM-Msb3

B

						FRM-Msb3						
S. cerevisiae	80	G	N	L	S	V	I	D	L	Y	G	D
K. lactis	82	A	G	L	S	L	I	N	L	Y	Y	D
Y. lipolytica	133	D	E	I	G	I	I	D	L	Y	S	K
S. bayanus	80	G	N	L	S	V	I	D	L	Y	G	D
S. stipitis	114	H	N	Y	S	L	L	D	I	Y	D	S
S. kluyveri	76	T	N	L	S	F	I	E	L	Y	N	E
N. castellii	74	G	S	L	S	V	I	D	L	Y	D	D
C. albicans	45	Q	N	F	S	V	L	R	E	Y	D	I
						-2		0		+2		

E

X-CRU

SHD1   SHD1$_{K97D}$ SHD1$_{L48,51A}$

N$_{ub}$-

Msb3
Msb4
Guk1
-

Ste7
Guk1
-

SHD1$_{L101A}$ SHD1$_{L109A}$ SHD1$_{L101,109A}$

Msb3
Msb4
Guk1
-

Ste7
Guk1
-

SD 5-FOA

C

SHD1CRU

N$_{ub}$-

Msb3
Msb3$_{\Delta84-88}$
Msb3$_{L82Y}$
Msb3$_{V84A}$
Msb3$_{I85A}$
Msb3$_{D86A}$
Msb3$_{L87A}$
Msb3$_{Y88A}$
Msb3$_{Y88F}$
-
Guk1

SD 5-FOA

	Split-Ub	ipTM
Msb3	+++	0.85
Msb3$_{D86A}$	+++	0.86
Msb3$_{L87A}$	++	0.82
Msb3$_{Y88F}$	+	0.74
Msb3$_{V84A}$	-	0.72
Msb3$_{I85A}$	-	0.59
Msb3$_{Y88A}$	-	0.32

D

SHD1CRU

N$_{ub}$-

Msb4
Msb4$_{\Delta35-39}$
Msb4$_{D37S}$
Msb4$_{D37K}$
Msb4$_{Y39A}$
Ste7
-
Guk1

SD 5-FOA

L109   L48   L51

L59

L101

L98

K97

SHD1

A

SHD1

Msb4$_{22-81}$

C$_{ub}$ RUra3

N

FRM

FRM

C

N$_{ub}$

5-FOA^S

N$_{ub}$

C$_{ub}$

RUra3

FRM

C

N

FRM*

5-FOA^R

B

N$_{ub}$-

Msb4 Msb3 Epo1 Aip5 Pea2

X- Spa2CRU

-

Msb4$_{22-81}$

-

Msb4$_{22-81Δ35-39}$

-

Ste7$_{105-165}$

SD 5-FOA

C

N$_{ub}$-Msb3

X- SHD1CRU

Msb4$_{22-81Δ35-39}$

Msb4$_{22-81Y39A}$

Msb4$_{22-81}$

Msb3$_{71-130}$

Ste7$_{105-165}$

Mkk1$_{143-202}$

Mkk1$_{143-202Δ171-175}$

-

SD 5-FOA

D

SHD1CRU

N$_{ub}$-

Mkk1

Mkk1$_{L171V}$

Mkk1$_{S173D}$

Ste7

-

Guk1

SD 5-FOA

E

SHD1CRU

N$_{ub}$-

Mkk1

Mkk1$_{Δ171-175}$

-

Guk1

SD 5-FOA

A

			FRM					
S.c. Msb3	80	G N L S	V I D L	Y	G D
N.c. GYP-3	502	S I E D	V M V M	Y	E R
N.c. CCP-1	498	L V Q D	I L D G	Y	L D
M.o. Gyp3	585	S V E D	V M G M	Y	E K
F.g. FgMsb3	358	S L E D	V M G M	Y	E Q
H.s. βPIX	712	Q I L K	V I E A	Y	C T
R.n. Piccolo	2258	I I L E	V L D A	Y	K D

−2   0   +2

A

Protein	ipTM
Dse3	0.89
Msb3	0.85
Ste7	0.85
Msb4	0.77
Mkk1	0.75
Ybr200w	0,73
Age2	0,69
Rpg1	0,67
Mkk2	0,61

B

FRM

S.c. Msb3   80   G N L S V I D L Y G D
S.c. Dse3   204   L F D E I L S A Y E N

C

N$_{ub}$-Msb3

X-SHD1CRU

Msb3$_{71-130}$
Dse3$_{181-240}$
Dse3$_{181-240\Delta208-212}$
-

SD 5-FOA

D

Scored residue / atom

Dse3 *SC*   SHD1 *SC*$_{123}$

Aligned residue / atom

Dse3 *SC*

SHD1 *SC*

FRM

FRM

E

FRM-Dse3

SHD1

Supplemental Material

Supplemental Figures

Figure S1

(A) Inputs for Fig.1 G. Coomassie staining of SDS-PAGE of extracts of *E.coli* cells expressing GST-SHD1 (lanes 1, 6) or GST (lanes 5, 10), the enriched 6xHis-Msb3$_{1-150}$ (lane 3), or 6xHis-Msb3$_{1-150\Delta76-93}$ (lanes 8) and the eluates of GST-SHD1- (lane 4, 9) or GST- (lanes 5, 10) coupled beads after being incubated with the enriched 6xHis-Msb3$_{1-150}$ (lanes 4, 5), or 6xHis-Msb3$_{1-150\Delta76-93}$ (lanes 9, 10).

(B) Full analysis of the GST pull-down experiment shown in Figure 1I. Lanes 1–4: input; lanes 5–7: SDS eluate after glutathione eluate; lanes 8–10: glutathione eluate. Lanes 8–10 are identical to lanes 1–3 in Figure 1I. Upper panel: anti-His immunoblot. Arrow indicates position of the cut. Lower panel: Ponceau S stain of the same membrane before antibody incubation. (C) AlphaFold structure prediction of SHD1 from yeast.

Figure S2

(A-E) Predicted Aligned Error (PAE) matrices from the AlphaFold predictions of the complexes between the SHD1 of budding yeast and the full length Msb3 (A), Msb4 (B), Ste7 (C), Mkk1 (D) and Mkk2 (E). Blue arrows point to the sequences around the FRMs. Dark green indicates a high probability score of interaction with the SHD1. (G) Predicted structures of the FRM-containing sequences of the complexes of A-E.

Figure S3

(A) Predicted Aligned Error (PAE) matrices from the AlphaFold predictions of the complexes between the GIT1$_{1-366}$ of rat and a fragment of Piccolo. Blue arrows points to the sequences around the FRMs that display a high score of interaction with the SHD1 of GIT1. (B) AlphaFold structure of the complex between GIT$_{1-366}$ and the FRM of Piccolo indicated in grey and highlighting the Y and D at the central positions. (C) Structure of the complex between the SHD1 of GIT1 and the FRM of Piccolo.
(D) Structure of the helix containing the FRM of Piccolo.

Figure S4

Source File: Figure 1G

1   2   3   4

106 kDa

65 kDa

45 kDa

27 kDa
17 kDa
15 kDa

Source File: Figure S1B   Figure 1I

1   2   3   4   1   2   3

106 kDa

65 kDa

45 kDa

27 kDa

17 kDa
15 kDa

Source File: Figure 6B

1   2   3   4   5   6   7

106 kDa

65 kDa

45 kDa

27 kDa
17 kDa
15 kDa

Supplemental Tables

Table S1

List of *S. cerevisiae* strains used and created in this study.

Identifier	Strain	Genotype	Source
JD47	JD47	*MATa; his3-Δ200; leu2-3, 112; lys2-801; trp1Δ63; ura3-52*	(Dohmen et al., 1995)
JD53	JD53	*MATα; his3-Δ200; leu2-3, 112; lys2-801; trp1Δ63; ura3-52*	(Dohmen et al., 1995)
YAD248	spa2Δ	*SPA2::hphNT1, JD47*	Johnsson Lab
YAD2703	msb3Δ; msb4Δ	*MSB3::hphNT1, msb4::kanMX, JD47*	Johnsson Lab
YAD2586	msb3Δ	*MSB3::natNT2, JD47*	Johnsson Lab
YAD2588	msb4Δ	*MSB4::kanMX4, JD47*	Johnsson Lab
YAD3097	$spa2_{\Delta 2\text{-}150}$	*$spa2_{\Delta 2\text{-}150}$, JD47*	Johnsson Lab
YLB53	$msb3_{\Delta 76\text{-}93}$	*$msb3_{\Delta 76\text{-}93}$, JD47*	This study
YAD4313	$msb4_{\Delta 27\text{-}44}$	*$msb4_{\Delta 27\text{-}44}$, JD47*	This study
YLB55	$msb3\Delta; msb4_{\Delta 27\text{-}44}$	*$MSB3::natNT2, msb4_{\Delta 27\text{-}44}, JD47$*	This study
YLB61	$msb4\Delta; msb3_{\Delta 76\text{-}93}$	*$MSB4::kanMX4, msb3_{\Delta 76\text{-}93}, JD47$*	This study
YAD4580	ste7Δ	*STE7::natNT2, JD47*	Johnsson Lab
YAD4570	$ste7_{\Delta 119\text{-}123}$	*$ste7_{\Delta 119\text{-}123}$, JD47*	Johnsson Lab
NBY167	N_{ub}-Msb3	*$P_{MSB3}::kanMX4\ P_{CUP1}\ N_{Ub}\text{-}HA, JD53$*	Johnsson Lab
NBY176	N_{ub}-Msb4	*$P_{MSB4}::kanMX4\ P_{CUP1}\ N_{Ub}\text{-}HA, JD53$*	Johnsson Lab
NBY22	N_{ub}-Bud2	*$P_{BUD2}::kanMX4\ P_{CUP1}\ N_{Ub}\text{-}HA, JD53$*	Johnsson Lab
NBY31	N_{ub}-Bud3	*$P_{BUD3}::kanMX4\ P_{CUP1}\ N_{Ub}\text{-}HA, JD53$*	Johnsson Lab
NJY194	N_{ub}-Ste7	*$P_{STE7}::kanMX4\ P_{CUP1}\ N_{Ub}\text{-}HA, JD53$*	Johnsson Lab
NBY270	N_{ub}-YEL023C	*$P_{YEL023C}::kanMX4\ P_{CUP1}\ N_{Ub}\text{-}HA, JD53$*	Johnsson Lab
YAD4299	N_{ub}-Msb3$_{\Delta 84\text{-}88}$	*$P_{MSB3}::kanMX4\ P_{CUP1}\ N_{Ub}\text{-}HA, MSB3::msb3_{\Delta 84\text{-}88}, JD53$*	This study
YAD4300	N_{ub}-Msb4$_{\Delta 35\text{-}39}$	*$P_{MSB4}::kanMX4\ P_{CUP1}\ N_{Ub}\text{-}HA, MSB4::msb4_{\Delta 35\text{-}39}, JD53$*	This study
YAD4303	N_{ub}-Ste7$_{\Delta 119\text{-}123}$	*$P_{STE7}::kanMX4\ P_{CUP1}\ N_{Ub}\text{-}HA, STE7::ste7_{\Delta 119\text{-}123}, JD53$*	This study
NBY339	N_{ub}-Guk1	*$P_{GUK1}::kanMX4\ P_{CUP1}\ N_{Ub}\text{-}HA, JD53$*	Johnsson Lab

MZY16	N_{ub} empty	P_{CUP1} N_{Ub}-HA, JD53	Johnsson Lab
NBY156	N_{ub}-Spa2	P_{SPA2}::kanMX4 P_{CUP1} N_{Ub}-HA, JD53	Johnsson Lab
NBY673	N_{ub}-Spa2$_{\Delta2-188}$	$P_{SPA2\Delta2-188}$::kanMX4 P_{CUP1} N_{Ub}-HA, JD53	Johnsson Lab
NBY674	N_{ub}-Spa2$_{\Delta2-150}$	$P_{SPA2\Delta2-150}$::kanMX4 P_{CUP1} N_{Ub}-HA, JD53	Johnsson Lab
NBY665	N_{ub}-Spa2$_{\Delta2-188}$	$P_{SPA2\Delta2-188}$::kanMX4 P_{CUP1} N_{Ub}-HA, JD53	Johnsson Lab
NBY787	N_{ub}-Msb3$_{L82Y}$	$P_{MSB3L82Y}$::kanMX4 P_{CUP1} N_{Ub}-HA, JD53	This study
NBY785	N_{ub}-Msb3$_{V84A}$	$P_{MSB3V84A}$::kanMX4 P_{CUP1} N_{Ub}-HA, JD53	This study
NBY774	N_{ub}-Msb3$_{I85A}$	$P_{MSB3I85A}$::kanMX4 P_{CUP1} N_{Ub}-HA, JD53	This study
NBY776	N_{ub}-Msb3$_{D86A}$	$P_{MSB3D86A}$::kanMX4 P_{CUP1} N_{Ub}-HA, JD53	This study
NBY782	N_{ub}-Msb3$_{L87A}$	$P_{MSB3L87A}$::kanMX4 P_{CUP1} N_{Ub}-HA, JD53	This study
NBY778	N_{ub}-Msb3$_{Y88A}$	$P_{MSB3Y88A}$::kanMX4 P_{CUP1} N_{Ub}-HA, JD53	This study
NBY780	N_{ub} -Msb3$_{Y88F}$	$P_{MSB3Y88F}$::kanMX4 P_{CUP1} N_{Ub}-HA, JD53	This study
NBY703	N_{ub}-Msb4$_{D37S}$	$P_{MSB4D37S}$::kanMX4 P_{CUP1} N_{Ub}-HA, JD53	This study
NBY704	N_{ub}-Msb4$_{D37K}$	$P_{MSB4D37K}$::kanMX4 P_{CUP1} N_{Ub}-HA, JD53	This study
NBY715	N_{ub}-Msb4$_{Y39A}$	$P_{MSB4Y39A}$::kanMX4 P_{CUP1} N_{Ub}-HA, JD53	This study
NBY107	N_{ub}-Epo1	P_{EPO1}::kanMX4 P_{CUP1} N_{Ub}-HA, JD53	Johnsson Lab
NBY595	N_{ub}-Aip5	P_{AIP5}::kanMX4 P_{CUP1} N_{Ub}-HA, JD53	Johnsson Lab
NBY450	N_{ub}-Pea2	P_{PEA2}::kanMX4 P_{CUP1} N_{Ub}-HA, JD53	Johnsson Lab
NBY211	N_{ub}-Mkk1	P_{MKK1}::kanMX4 P_{CUP1} N_{Ub}-HA, JD53	Johnsson Lab
NBY734	N_{ub}-Mkk1$_{L171V}$	$P_{MKK1L171V}$::kanMX4 P_{CUP1} N_{Ub}-HA, JD53	This study
NBY705	N_{ub}-Mkk1 S173D	$P_{MKK1173D}$::kanMX4 P_{CUP1} N_{Ub}-HA, JD53	This study
NBY139	N_{ub}-Ste11	P_{STE11}::kanMX4 P_{CUP1} N_{Ub}-HA, JD53	This study
NBY767	N_{ub}-Ste11$_{\Delta1-105}$	$P_{STE11\Delta1-105}$::kanMX4 P_{CUP1} N_{Ub}-HA, JD53	This study
NBY754	N_{ub}-Ste11$_{\Delta1-410}$	$P_{STE11\Delta1-410}$::kanMX4 P_{CUP1} N_{Ub}-HA, JD53	This study
N0BY73	N_{ub}-Ste7Δ1-164	$P_{STE7\Delta1-164}$::kanMX4 P_{CUP1} N_{Ub}-HA, JD53	This study
STY165	Msb3$_{\Delta76-93}$-CRU 303	$MSB3_{\Delta76-93}$::msb3-C_{ub}-R-URA3 HIS3, JD47	Johnsson Lab
YLB58	Msb3$_{\Delta76-93}$-CRU 303	msb3$_{\Delta76-93}$::msb3-C_{ub}-R-URA3 HIS3, JD47	This study

YAD4440	pMet17 Msb4-CRU 303	$P_{MSB4}::P_{MET17}$, $MSB4::msb4-C_{ub}-R-URA3$ HIS3, JD47	Johnsson Lab
YAD4441	pMet17 Msb4$_{\Delta27-44}$-CRU 303	$P_{MSB4}::P_{MET17}$, $msb4_{\Delta27-44}::msb4-C_{ub}-R-URA3$ HIS3, JD47	This study
UNY7	Ste5-CRU303	$ste5::ste5-C_{ub}-R-URA3$ HIS3, JD47	Johnsson Lab
STY1659	Ste7$_{105-165}$-Spa2	$P_{SPA2}::ste7(105-165)-spa2::spa2-C_{ub}-R-URA3$ HIS3, JD47	Johnsson Lab
STY1657	Msb4$_{22-81}$-Spa2	$P_{SPA2}::msb4(22-81)-spa2::spa2-C_{ub}-R-URA3$ HIS3, JD47	Johnsson Lab
STY1658	Msb4$_{22-81\Delta35-39}$-Spa2	$P_{SPA2}::msb4(22-81)_{\Delta35-39}-spa2::spa2-C_{ub}-R-URA3$ HIS3, JD47	Johnsson Lab
YAD2998	Msb3-GFP304	$MSB3:MSB3-GFP$ TRP1	Johnsson Lab
YAD2999	Msb4-GFP304	$MSB3:MSB4-GFP$ TRP1	Johnsson Lab
YAD4318	Msb3$_{\Delta76-93}$-GFP 304	$MSB3::msb3_{\Delta76-93}-GFP$ TRP1	This study
YLB83	Msb4$_{\Delta27-44}$-GFP 304	$MSB4::msb4_{\Delta27-44}-GFP$ TRP1	This study
YLB86	Msb3-GFP 304; Spa2$_{\Delta2-150}$	$MSB3::msb3-GFP$ TRP1, $SPA2::spa2_{\Delta2-150}$, JD47	This study
YLB87	Msb4-GFP 304; Spa2$_{\Delta2-150}$	$MSB4::msb4-GFP$ TRP1, $SPA2::spa2_{\Delta2-150}$, JD47	This study
YAD4500	Ste7-GFP 304	$STE7::ste7-GFP$ TRP1, JD47	Johnsson Lab
YAD4501	Ste7$_{\Delta119-123}$-GFP 304	$STE7::ste7_{\Delta119-123}-GFP$ TRP1, JD47	Johnsson Lab

Table S2
List of plasmids used and created in this study.

Identifier	Plasmid	Genotype	Source
BS4075	PMET17-Spa2(1-124)-CRU pRS313	*PMET17-spa2(1-124)- Cub-R-URA3, AmpR, HIS3, CEN*	Johnsson Lab
BA2009	PMET17-Spa2 L48,51A (1-124) - CRU pRS313	*PMET17-spa2 L48,51A (1-124)- Cub-R-URA3, AmpR, HIS3, CEN*	Johnsson Lab
BL1001	PMET17-Spa2 L101A (1-124)-CRU pRS313	*PMET17-spa2 L101A (1-124)- Cub-R-URA3, AmpR, HIS3, CEN*	This study
BL1002	PMET17-Spa2 L109A (1-124)-CRU pRS313	*PMET17-spa2 L109A (1-124)- Cub-R-URA3, AmpR, HIS3, CEN*	This study
BL1003	PMET17-Spa2 L101,109A (1-124)- CRU pRS313	*PMET17-spa2 L101,109A (1-124)- Cub-R-URA3, AmpR, HIS3, CEN*	This study
BL1004	PMET17-Spa2 K97D (1-124)-CRU pRS313	*PMET17-spa2 K97D (1-124)- Cub-R-URA3, AmpR, HIS3, CEN*	This study
BA2075	Msb4(22-81)Δ35-39-Spa(1-124)-CRU	$P_{MET17}MSB4(22-81)\Delta35-39-SPA2(1-124)-CRU313$	This study
BA2080	Msb4(22-81)Y39A-Spa2(1-124)-CRU	$P_{MET17}MSB4(22-81)Y39A-SPA2(1-124)-CRU313$	This study
BA2072	Mkk1(105-164)-Spa2(1-124)-CRU	$P_{MET17}MKK1(105-164)-SPA2(1-124)-CRU313$	This study
BA2045	Ste7(105-165)-Spa2(1-124)-CRU	$P_{MET17}STE7(105-165)-SPA2(1-124)-CRU313$	This study
BA2044	Msb3(71-130)-Spa2(1-124)-CRU	$P_{MET17}MSB3(71-130)-SPA2(1-124)-CRU313$	This study
BA2075	Msb4(22-81)-Spa2(1-124)-CRU	$P_{MET17}MSB4(22-81)-SPA2(1-124)-CRU313$	This study
BA884	N_{ub}-Spa2$_{1-150}$	$P_{CUP}NubHA-Spa2(1-150) kanMX CEN$	Johnsson Lab
BA2006	N_{ub}-Msb3$_{71-100}$	$P_{Cup}NubHAkanMX Msb3_{71-100} CEN$	Johnsson Lab
BA2007	N_{ub}-Msb4$_{21-50}$	$P_{Cup}NubHAkanMX Msb4_{21-50} CEN$	Johnsson Lab
BA2008	N_{ub}-Ste7$_{105-135}$	$P_{Cup}NuiHAkanMX Ste7_{105-135} CEN$	Johnsson Lab

Table S3
Raw fluorescence intensities of GFP fusions for calculating the fluorescence ratios in Figure 4D.

Msb3

Shmoo tip	Cytoplasm	background
964,583	643,375	512,833
801,583	614,292	525,167
778,875	643,25	525,625
878,458	668,333	524,833
769,5	612,625	529,375
973,875	689,583	528
838,917	625,917	525,75
1035,167	694,375	531,292
966,708	676,083	531,792
858,708	658,833	544,333

Msb3$_{\Delta76\text{-}93}$

Shmoo tip	Cytoplasm	Background
829,5	882,417	587,083
758,182	725,909	578,909
746,727	758	576,273
752,818	742,818	595,727
732,818	711,909	567,636
727,545	807	564,455
732,636	726	586,455
755,545	762	541,727
813,727	820,091	576,818
831,545	801,727	545,273

Msb4

Shmoo tip	Cytoplasm	background
1363,364	1094,364	788,818
1162,545	966,455	775
1442	1051,545	769,455
1279,818	1089	751,909
1435,364	1096,091	727
1454,273	1088,182	773,182
1343,125	991,625	801,75
1159,438	858,455	751,545
1642	1226,364	902,818
1667,455	1453,818	1069,909

Msb4$_{\Delta27\text{-}44}$

Shmoo tip	Cytoplasm	background
1294,636	1343	835,273
1317,273	1264	824,818
1112	1096,727	866,273
1212,909	1306,273	776,818
1212,818	1291,636	801,636
1257	1258,909	847,273
1268,455	1251,727	826,818
1246,091	1397,273	834,091
1209,75	1203,182	843,545
1209,545	1104	856,75

Ste7

Shmoo tip	Cytoplasm	background
958,792	706,042	533,167
885,75	661,375	542
840,083	696,917	542,581
909	666,25	540,292
785,125	649,083	553,708
721,167	606,792	530,625
981,208	672,833	541,458
926,333	627,292	545,875
908,167	651,583	544,708
862,125	671,875	529,833

Ste7$_{\Delta112\text{-}133}$

Shmoo tip	Cytoplasm	background
763,333	640,417	541,708
661,625	683,417	530,583
646,708	635,042	543,208
702,167	699,667	538,5
748,625	663,333	524,458
685,667	664,583	529,458
658,083	668,5	531,042
687,75	638,792	537,333
696,708	657,25	518,958
640,125	666,083	523

Table S4
List of potential new FRM-containing binding partners of Spa2.

UniProt Accession	Yeast Gene	Matching Step
P04710	ADT1	Step3
P06784	STE7	Step6
P07264	LEU1	Step4
P07866	LTE1	Step4
P09032	GCD1	Step2
P0C5P0	YJL197C	Step4
P0CX94	YER190C	Step2
P0CX95	YEL077W-A	Step2
P0CX96	YGR296C-B	Step2
P0CX97	YPL283W-B	Step2
P0CX98	YPR204C-A	Step2
P0CX99	YFL068W	Step2
P0CY00	YLL066W-A	Step2
P0CY01	YLL067W-A	Step2
P13433	RPO41	Step3
P14772	BPT1	Step4
P14908	MTF1	Step2
P15274	AMD1	Step4
P18239	ADT2	Step3
P18412	*TEC1*	Step4

P19097	*FAS2*	Step4
P19263	*MED14*	Step4
P19658	*EXO70*	Step4
P20967	*ODO1*	Step3
P21595	*CP56*	Step3
P23255	*TAF2*	Step4
P23642	*VAN1*	Step4
P25046	*MED19*	Step4
P25351	*YCR3*	Step4
P25389	*KCC4*	Step4
P25569	*GID7*	Step2
P26725	*YUR1*	Step2
P27654	*TIP1*	Step4
P28273	*OXP1*	Step4
P32461	*DPH2*	Step4
P32490	*MKK1*	Step6
P32491	*MKK2*	Step5
P32496	*RPN12*	Step4
P32522	*PET309*	Step3
P32523	*PRP19*	Step2
P32862	*RGT1*	Step4
P32906	*MNS1*	Step4

P32907	ATO2	Step4
P35189	TAF14	Step2
P36051	MCD4	Step4
P36083	YKH5	Step4
P38009	ADE17	Step2
P38041	BOB1	Step4
P38073	EDS1	Step4
P38207	APN2	Step2
P38249	RPG1	Step5
P38696	ARP1	Step4
P39079	CCT6	Step4
P39931	SSP120	Step4
P39985	DPO5	Step4
P39986	SPF1	Step4
P40004	YEA4	Step4
P40350	ALG5	Step4
P40509	SEC28	Step4
P40529	AGE2	Step5
P40552	TIR3	Step4
P40857	PHS1	Step4
P40986	CDC1	Step2
P43571	BST1	Step4

P43583	BLM10	Step4
P46682	AP3B	Step4
P46954	SIP4	Step4
P46956	PHO86	Step2
P47069	MPS3	Step2
P47101	RBH2	Step4
P47167	MCM22	Step4
P48566	MSB3	Step6
P49704	PRP31	Step4
P50273	ATP22	Step3
P53108	YIP5	Step4
P53170	PDK2	Step3
P53239	COX18	Step3
P53829	CAF40	Step4
P53901	INN1	Step2
P53909	AAH1	Step4
P53952	YNL050C	Step4
P54113	ADE16	Step2
P60010	ACT1	Step4
Q00764	TPS1	Step4
Q02457	TBF1	Step4
Q02732	CFT19	Step4

Q03210	NGL3	Step2
Q03233	ADD37	Step4
Q03305	RMT2	Step4
Q03920	MTQ2	Step2
Q04002	SCC2	Step4
Q04178	HPT1	Step2
Q04301	VBA1	Step4
Q04322	GYL1	Step4
Q04372	TAP42	Step4
Q04511	UFO1	Step4
Q04585	YDR109C	Step4
Q04602	VBA4	Step4
Q05791	CDC123	Step2
Q06058	SEI1	Step2
Q06142	IMB1	Step4
Q06598	ARR3	Step4
Q06632	CFT1	Step2
Q06704	IMH1	Step4
Q07376	MCH1	Step4
Q07527	TRM3	Step4
Q07830	GPI13	Step4
Q08179	MDM38	Step4

Q08236	*AVO1*	Step4
Q08729	*DSE3*	Step6
Q12029	*FSF1*	Step4
Q12038	*SRO7*	Step2
Q12109	*SYWC*	Step4
Q12167	*RRG1*	Step3
Q12200	*NPC1*	Step4
Q12205	*MNL2*	Step4
Q12297	*TAF3*	Step4
Q12317	*MSB4*	Step6
Q12321	*MED1*	Step4
Q12511	*PDP1*	Step2
Q3E755	*YBR200W-A*	Step5
Q6Q560	*ISD11*	Step4
Q99207	*NOP14*	Step4
Q99222	*AFI1*	Step4