Ultrasound neuromodulation reveals distinct roles of the dorsal anterior cingulate cortex and anterior insula in learning

Nomiki Koutsoumpari *[1,2], Johannes Algermissen *[3,4], Siti Yaakub[1,2],

Hanneke E. M. den Ouden[4], Nadege Bault[1,2] *, Elsa Fouragnan[1,2] *

[1]School of Psychology, University of Plymouth

[2]Brain Research Imaging Center (BRIC), University of Plymouth

[3]University of Oxford, Department of Experimental Psychology, Oxford OX1 3SR, United Kingdom

[4]Radboud University, Donders Institute for Brain, Cognition and Behaviour, Nijmegen, The Netherlands

* Authors contributed equally to this paper

Abstract

Pavlovian biases reflect the notorious influence of hard-wired, evolutionarily conserved cue-
response tendencies on instrumental action selection: people show automatic action
invigoration in face of potential rewards, but action suppression in face of potential
punishments. The neural origin of these biases is unclear. Past evidence suggests dorsal
anterior cingulate cortex (dACC) and anterior insula (aIns) as part of a "reset network" that
rapidly responds to salient information and might contribute to these biases. We used
transcranial ultrasonic stimulation (TUS) in 29 healthy participants to interfere with neural
activity in these regions and test their causal role in a within-subject, counter-balanced design
across three sessions (sham, TUS-dACC, TUS-aIns). Computational modelling revealed a
double dissociation, with distinct roles of both regions in Pavlovian biases: while TUS to the
aIns decreased people's tendency to overly take credit for rewards following action and to
ignore punishments following inaction, TUS to dACC increased participants' tendency to take
the cue valence as a reinforcer signal. Although the dACC and aIns are part of the same
network and often co-activate during decision-making tasks, TUS interference reveals their
distinct roles: the dACC mediates cue-dependence persistence while the aIns is critical for
inferring whether outcomes are self-caused.

Introduction

Some of our everyday decisions are remarkably fast: in the blink of an eye, we paralyze before our hand reaches a potentially venomous spider, or we instinctively reach for the last remaining cookie before our siblings can grab it. Psychology and neuroscience have explained our ability for such fast, instinctive decisions by postulating the existence of multiple decision-making systems that trade-off speed against deliberation and sophistication (Daw et al., 2005; Kahneman, 2011; Keramati et al., 2011). One class of especially fast, but also rigid and seemingly hardwired decisions has been proposed as being driven by a Pavlovian control system (Algermissen and den Ouden, 2023; Dayan et al., 2006; Guitart-Masip et al., 2014; O'Doherty et al., 2017). This system rigidly responds with invigoration to any chance to gain rewards ("Go"), but with inhibition to any looming threat of punishment ("NoGo"). This bias has classically been called the Pavlovian *response bias* (Guitart-Masip et al., 2012; Swart et al., 2017).

The Pavlovian control system gives rise to two other subtle, yet well documented Pavlovian biases that influence learning. One the one hand, a Pavlovian *learning bias* describes people's tendency to overly claim rewards as caused by their actions, but ignore punishments that follow inaction. For example, a football coach might credit himself for his team's win after having changed the player lineup, but fail to acknowledge how not changing it led to repeated defeats in the past. One the other hand, a Pavlovian cue-valence dependent *persistence bias* describes people's tendency to interpret cues that signal the chance for rewards/punishments as reinforcer signals in themselves. In case of no (or an ambiguous) feedback, people are likely to persist in situations in which a reward could be obtained, but likely to change their behaviour in situations in which a negative outcome might occur (Algermissen et al., 2024). Pavlovian biases such as these become evident in tasks that orthogonalize action requirement (Go/NoGo) against the valence of potential outcomes (rewards/punishments) (Guitart-Masip et al., 2012; Swart et al., 2017) and appear to be evolutionarily ancient and shared across the animal realm (Breland and Breland, 1961). By shaping how we assign value to actions and learn from outcomes, these biases play a fundamental role in adaptive behaviour, and their dysregulation may underlie core symptoms across a range of psychiatric disorders (Huys et al., 2016; Scholz et al., 2025).

Pavlovian biases have been linked to neural activity across various cortical and subcortical regions (Algermissen et al., 2024, 2022; Guitart-Masip et al., 2012), yet the precise causal contribution of each of these regions remain unclear. Pinpointing when and where in the brain these biases emerge is essential for uncovering their underlying mechanisms. We have recently observed that the mere anticipation of potential punishment elicits a rapid freezing of gaze within 200–300ms of cue onset (Algermissen and den Ouden, 2024a), coinciding with early value-related components in electroencephalography (EEG) recordings (Algermissen et al., 2024, 2022). These findings suggest that value-related information is processed remarkably early, allowing it to exert a fast and automatic influence on action selection. EEG-informed functional magnetic resonance imaging (fMRI) analyses have further revealed that such early neural responses recruit a network involving the dorsal anterior cingulate cortex (dACC) and anterior insula (aIns), which are selectively engaged by salient/negative information and appear to support rapid behavioural adaptation (Fouragnan et al., 2015). In contrast, later feedback processing stages engage the striatum to support more deliberate reinforcement learning. Together, these findings position the dACC and aIns as strong candidates for causal contributors to Pavlovian biases observed in behaviour.

Selective interference with neural processing in these deep cortical regions to test their role in Pavlovian biases is exceedingly challenging with conventional non-invasive brain stimulation techniques which are limited to superficial cortical regions. However, recently, TUS has emerged as a promising non-invasive tool for neuromodulation (Bault et al., 2024; Murphy and Fouragnan, 2024) that can stimulate deep brain regions with unprecedented spatial precision when careful precautions are taken to limit the transmission loss caused by the skull. TUS has been successfully used to change choice behaviour and neural activity in non-human primates (Folloni et al., 2021; Fouragnan et al., 2019) and humans (Yaakub et al., 2024, 2023b). We

thus tested the role of dACC and aIns in Pavlovian biases using a single-blinded within-subject
counterbalanced 3-session (dACC/aIns/sham) offline TUS experiment. We expected TUS to
interfere with neural processing in these regions and thus lead to different expression of
Pavlovian biases in behaviour (Guitart-Masip et al., 2012; Swart et al., 2017) since both regions
have been implicated in different aspects of feedback-based learning, including biased credit
assignment (Algermissen et al., 2024; de Boer et al., 2019; Swart et al., 2017). We thus also
tested for effects on several Pavlovian biases, including a *persistence bias* and a *learning bias,*
using computational reinforcement learning models designed to isolate latent cognitive
processes.

To foreshadow our results, we found that the stimulation of both dACC and aIns altered
Pavlovian biases. Using computational modelling, we found a double dissociation in which
dACC and aIns affected distinct learning and decision processes. TUS-aIns selectively
reduced participants' tendency to overly attribute rewards to their own actions and their
reluctance to attribute punishments to their inactions, captured by a learning bias parameter
(Swart et al., 2017). In contrast, TUS-dACC impaired participants' ability to tell apart the
valence of the feedback from the valence of the cue they responded to, captured by a
persistence bias parameter (Algermissen et al., 2024). These results indicate that the dACC
plays a role in selecting which cues in the environment to use as a reinforcer signal, whereas
the aIns is critical for inferring whether outcomes are self-caused. These two mechanisms —
reliance on cue valence as a reinforcer signal and assigning credit to one's own actions —may
be altered in psychiatric disorders (Korn et al., 2014; Shahar et al., 2021) potentially
contributing to symptoms such as pathological guilt, externalized blame, and compulsive
repetition of actions despite unhelpful outcomes. Disentangling their specific neural
contributions is therefore essential for advancing both cognitive neuroscience and clinical
psychiatry.

This study also demonstrates the transformative potential of TUS for causally probing decision-
making processes in deep cortical regions typically beyond the reach of conventional non-
invasive techniques such as transcranial magnetic stimulation (TMS). By targeting the dACC
and aIns, two regions often grouped within the salience network and previously implicated in
rapid valence-action processing, we show that they play dissociable roles in biased learning.
The ability to modulate them independently with TUS represents not only a significant
methodological advance but also lays the foundation for mechanistically precise interventions
targeting maladaptive decision-making in psychiatric disorders.

Results

Participants and procedure

29 participants, screened for any counter indications to TUS and magnetic resonance imaging
(MRI) [Contraindications section in Supplementary Material] completed a four-session study
protocol. The first session involved collecting MR images for neuronavigation and generating
a pseudo-computed tomography (pseudo-CT) image for personalised acoustic planning
(Yaakub et al., 2023a) as well as a task practice (Fig. 1A). Participants were then assigned to
one of three TUS conditions: Sham, dACC, or aIns (Fig. 1B), using a counterbalanced design.
Each of these subsequent sessions included an 80 sec long 5 Hz-patterned TUS intervention,
with a spatial peak pulse average intensity (ISPPA) of approximately 50 W/cm² in water,
immediately followed by a motivational Go/NoGo learning task (Algermissen et al., 2024;
Guitart-Masip et al., 2012; Swart et al., 2017). While the task structure remained consistent
across sessions, the cue sets were different and individually randomized to prevent learning
effects. Each session lasted approximately 1.5 hours.

During the motivational Go/NoGo learning task, on each trial, participants were presented with
one of several cues. They had to learn from probabilistic feedback via trial-and-error to either
perform a Go (button press) or a NoGo action (no button press) to four cues per block. Half of
these cues were Win cues for which correct responses (mostly) returned a reward (increase

in point score), while incorrect responses (mostly) led to neutral feedback (no change in point
score). For the other half of cues, called Avoid cues, correct responses (mostly) led to neutral
feedback, while incorrect responses (mostly) led to punishments (decrease in point score).
The task timeline is presented in Fig. 1D. Feedback was probabilistic such that correct
responses resulted in the desired outcome (reward or omitted punishment) 80% of the time,
whereas incorrect actions led to the desired outcome in only 20% of trials. On each session,
participants completed 320 trials, split into four blocks of 80 trials. In each block, a distinct set
of four cues was presented, each repeated 20 times. The order of cue presentations was
randomized for each participant within each block (Fig. 1D).

Targeting of the bilateral aIns and dACC was individually adjusted using T1-weighted MRI
(structural scans) during planning for target and transducer placement, as well as acoustic and
thermal simulation. The dACC was defined anatomically as the region anterior to the anterior
commissure, dorsal to the genu of the corpus callosum, and bounded dorsally by the cingulate
sulcus. The aIns was identified as the anterior portion of the insular cortex, located deep within
the Sylvian fissure, anterior to the central insular sulcus, and bordering the frontal operculum.
Repetitive TUS followed a 5 Hz-patterned protocol with a 10% duty cycle (Fundamental
Frequency = 500 KHz, Pulse Duration = 20 ms, pulsed every 200 ms), applied for 80 seconds
(400 pulses in total). The ISPPA in water was maintained at 50 W/cm² across participants
(validated in a hydrophone tank (Yaakub et al., 2023b)). We took care to remain within
guidelines for human ultrasound exposure as defined the International Thermal and
Radiological Ultrasound Safety Standards and Thresholds (ITRUSST; Aubry et al., 2024).
Importantly the transcranial Mechanical Index (MItc) was way below 1.9 and the results of our
acoustic simulations predominantly indicated a maximum soft tissue temperature rise below
2°C [Supplementary Table 1]. We also calculated the Cumulative Equivalent Minutes at 43°C
(CEM43), a metric reflecting both duration and intensity of heating relative to 43°C, the critical
threshold for thermal cell damage. The CEM43 values were always below 0.1, i.e. well below
the safety threshold of 0.25 (Aubry et al., 2024).

Sham was administered in the same manner as active aIns-TUS, placing the transducer over
the site but without actual stimulation. Instead, participants listened to a sound that mimicked
the TUS protocol's acoustic output (validated across 6 participants independent from this
study), played through bone conduction headphones. Acoustic simulations for each participant
were used to plan the stimulation to ensure accurate pressure delivery at the target area and
adherence to safety protocols (an example participant's targeted areas are shown in Fig. 1E).
Detailed acoustic simulation parameters and outputs for all study participants can be found in
Fig. 1F, Supplementary Materials [Table 1] and summarised in Table 1.

Side effects

Participants completed a post-TUS questionnaire for side effects twice: once at the end of each
session, and a second time 12 to 48 hours after the session. They were presented with a list
of symptoms for which they had to rate the intensity of the symptom and the degree to which
they believe that symptom was related to the stimulation (refer to the Methods section under
TUS protocol and procedure for further safety details; see also Supplementary Material 1 for
the full safety questionnaire). Immediately after stimulation of the insula, participants reported
the following side effects: anxiety (N = 1), difficulty paying attention (N = 2), forgetfulness (N =
1), headache (N = 1) and sleepiness (N = 3). All side effects were rated as mild and unrelated
to the stimulation. Following stimulation of the dACC, moderate anxiety (N = 1), mild
forgetfulness (N = 1), moderate neck pain (N = 1), mild to moderate sleepiness (N = 2) and
mild vision problem (N = 1) were reported. Only the participant reporting vision problems
considered those as possibly related to the stimulation. Following sham stimulation, one
participant reported mild itchiness (N = 1) and another moderate sleepiness (N = 1), again both
problems were rated as unrelated to the stimulation. Overall, no qualitative difference in the
number and quality of side effects was observed between sham and active stimulation. In most
cases of side effects, participants did not attribute them to the stimulation.

Target	Tissue	Temperature (°C)		Pressure (MPa)		ISPPA (W/cm2)		MItc	
		M	SD	M	SD	M	SD	M	SD
dACC	Target ROI	37.18	0.05	0.45	0.04	6.69	1.16	0.63	0.06
	Soft tissue	37.96	0.28	0.45	0.04	6.72	1.18	0.63	0.06
	Skull	38.73	0.41	0.32	0.08	3.57	2.29	0.45	0.12
l-aIns	Target ROI	37.19	0.04	0.47	0.08	7.60	2.26	0.67	0.11
	Soft tissue	38.14	0.34	0.48	0.06	7.74	1.98	0.68	0.09
	Skull	38.99	0.64	0.33	0.05	3.63	1.09	0.46	0.07
r-aIns	Target ROI	37.17	0.03	0.48	0.03	7.53	1.06	0.67	0.05
	Soft tissue	38.20	0.38	0.48	0.03	7.54	1.05	0.67	0.05
	Skull	39.10	0.62	0.35	0.06	4.06	1.39	0.49	0.08

Table 1. Ultrasound Parameters Across Different Target Tissues for dACC and aIns
Stimulation. A sphere (Target ROI) with a 3mm radius centred on the target coordinates is
used to reliably extract peak values from the brain region where the focus is located.

Figure 1. A. Study Design: In the first session, a T1-weighted image was acquired for
neuronavigation and acoustic planning. Participants also practised a short version of the task.
In sessions two to four, participants were randomly allocated to one of three conditions: Sham,
dACC, or bilateral aIns. Post-stimulation safety questionnaires were completed on the same
day and the following day. **B. TUS positioning:** Visualisation of the positioning of the
transducer on the head for the active TUS sessions. **C. Task design:** On each trial,
participants saw an abstract cue and needed to learn from trial-and-error to respond with either
a Go or NoGo response to that cue. The cue disappeared at the response onset, followed by
a jittered interval and finally response-dependent feedback. Each cue had one correct action
(Go or NoGo) and was either a Win cue (leading to rewards or neutral outcomes) or an Avoid
cue (leading to neutral outcomes or punishments). Rewards and punishments were

represented as money falling into or out of a bucket. **D. Feedback validity and cue types:** Each block of session included 4 novel cues, varying in valence (Win/Avoid) and required action (Go/NoGo). Incongruent cues, where the required action was in opposition to the Pavlovian bias (NoGo to Win, Go to Avoid punishment), are highlighted in grey. Feedback was probabilistic. Correct responses to Win cues resulted in rewards 80% of the time and neutral outcomes 20% of the time. Correct actions to Avoid cues led to neutral outcomes in 80% of cases and punishments in 20%. Incorrect responses led to feedback with the reverse probabilities. **E. Regions:** Targeted brain regions for TUS (dACC and bilateral aIns) are shown (top), with post-stimulation pressure simulated through k-Plan software (BrainBox, Inc.) expressed in megapascals on a representative participant (bottom). **E. Regions:** Targeted brain regions for TUS (dACC and bilateral aIns) are shown (top), with post-stimulation pressure simulated through k-Plan software (BrainBox, Inc.) expressed in megapascals on a representative participant (bottom). **F. TUS simulations:** Maximum pressure in the brain target volume (top panel) and maximum temperature rise in the brain target volume (bottom panel), (n=29). Box plots show the mean, and the standard error (bounds of the box). Data from each individual participant are presented as small black circles.

Pavlovian biases in responding and learning in the sham session

We first focus on the data from the sham session to replicate Pavlovian biases in responding and learning as typically observed in this task (Algermissen et al., 2024; Guitart-Masip et al., 2012; Swart et al., 2017).

First, in line with previous work (Algermissen et al., 2024; Scholz et al., 2022; Swart et al., 2017; van Nuland et al., 2020), we used mixed-effects logistic regression to fit Go/NoGo responses as a function of the required action of a given cue (Go vs. NoGo), the cue valence (Win vs. Avoid cues), and their interaction. Participants successfully learned the task, as they made more Go responses to Go than NoGo cues (main effect of required action: b = 2.132, 95%-CI [1.797, 2.467], $\chi^2(1)$ = 53.812, p < .001; Fig. 2D). Furthermore, there was a Pavlovian bias in behaviour, indexed by significantly more Go responses to Win than Avoid cues (main effect of cue valence: b = 0.689, 95%-CI [0.421, 0.966], $\chi^2(1)$ = 17.173, p < .001). Lastly, the interaction between both terms was significant (b = 0.269, 95%-CI [0.104, 0.434], $\chi^2(1)$ = 9.646, p = .002), reflecting a stronger Pavlovian bias for Go than for NoGo cues. Similar analyses of the reaction times (RTs) also revealed a bias (Supplementary Fig. 2). These analyses confirm that participants' behaviour was influenced by Pavlovian biases in the sham session, which constitutes a prerequisite for modulating them in the other two active sonication sessions.

Second, we used mixed-effects logistic regression to analyse participants' propensity to repeat vs. switch a response on the next trial as a function of the obtained outcome and performed response on the current trial. We have previously found that Pavlovian biases are present not only in action selection, but also in learning, with a larger propensity to take valenced feedback into account (i.e., repeat actions after rewards, switch actions after punishments) for Go actions compared to NoGo actions. Specifically, learning is increased for rewarded Go (relative to rewarded NoGo) actions and decreased for punished NoGo (relative to punished Go) actions (see detailed explanation in the computational modelling) (Algermissen et al., 2024; Swart et al., 2017). Because this bias should only occur for valenced outcomes (rewards, punishments), but not for neutral outcomes, we only included valenced outcomes in this analysis. We observed a significant main effect of the obtained outcome (reward vs. punishment), b = 1.436, 95%-CI [1.190, 1.683], $\chi^2(1)$ = 50.170, p < .001, with more response repetitions after rewards than after punishments, and a main effect of performed response, b = 0.211, 95%-CI [0.081, 0.341], $\chi^2(1)$ = 6.422, p = .001, with more response repetitions for Go than NoGo responses (likely due to the fact that participants showed overall more Go than NoGo responses in this task). Crucially, the interaction between the obtained outcome and performed response was significant, b = 0.423, 95%-CI [0.195, 0.650], $\chi^2(1)$ = 10.051, p < .001, with a stronger outcome effect after Go than after NoGo responses (Fig. 2E). This result confirmed the presence of Pavlovian biases also in learning from feedback.

Thirdly, we used mixed-effects logistic regression to analyse participants' propensity to repeat
vs. switch a response on the next trial as a function of the cue valence. We have previously
found that participants show a propensity to repeat responses for Win cues than for Avoid
cues, suggesting that they also partly interpret the cue valence as a feedback signal
(Algermissen et al., 2024). This leads to response repetition (persistence) tendencies that
depend on the cue valence, a tendency that was not well captured by previous computational
models (Supplementary Fig. 4), but can only be accounted for by a third, *persistence bias*.
Indeed, we found a strong main effect of cue valence on response repetitions, $b = 0.351$, 95%-
CI [0.262, 0.440], $\chi^2(1) = 60.117$, $p < .001$, with more repetitions for Win than Avoid cues (Fig.
2F). This result indicated the presence of a third Pavlovian bias in which the cue valence
affected persistence tendencies. Taken together, these results motivated us to further
interrogate these biases using computational reinforcement learning models.

Computational modelling of the sham data

We next fit a series of increasingly complex reinforcement learning models to participants'
responses in the sham sessions. These models follow previous publications on this task
(Algermissen et al., 2024; Guitart-Masip et al., 2012; Swart et al., 2017). These models
incorporate three previously described forms of Pavlovian biases: (a) a response bias, (b) a
learning bias, and (c) a persistence bias.

The first bias, called *response bias*, is a bonus/malus added to the learned action values (Fig.
2A): For Win cues, a bonus is added to the value of Go actions, and for Avoid cues, a malus
is subtracted from the value of Go actions (Guitart-Masip et al., 2012). This bias is present at
the onset of the task and constant throughout learning. Since action values and the bias jointly
determine which action is selected, and since action values will increasingly separate with
learning over the course of the task, the role of this response bias will, in net, decrease over
the time (Swart et al., 2017).

The second bias, called *learning bias*, captures two tendencies well described in the literature:
people ascribe positive outcomes to their own actions (which in other contexts can lead to
"illusions of control" (Brown and Jenkins, 1968; Wegner, 2002)), but tend to ignore negative
outcomes after having remained passive (a form of "omission bias" (Ritov and Baron, 1990;
Zeelenberg et al., 2000); Fig. 2B). These tendencies are captured by a boost in the learning
rate for rewarded Go actions and a decrement in the learning rate for punished NoGo actions
(Swart et al., 2017). The impact of this bias on behaviour only develops over time, where the
action values become increasingly distorted relative to unbiased learning.

The third bias, called *persistence bias*, captures the fact that the cue valence (Win or Avoid
cue) can act as a reinforcer (Fig. 2C) (Algermissen et al., 2024): For Win cues, rewards signal
positive feedback, and neutral outcomes signal negative feedback. However, the Win cue itself
can erroneously be interpreted as a feedback signal and reinforce behaviour irrespective of
the eventual outcome. Thus, while a neutral outcome for a Win cue signals negative feedback,
which suggests switching behaviour, the cue valence itself might be taken for positive
feedback, which usually translates into a response repetition. Vice versa, for Avoid cues,
neutral outcomes signal positive feedback and punishments signal negative feedback. The
Avoid cue itself can erroneously be interpreted as negative feedback. Thus, while a neutral
outcome for an Avoid cue signals positive feedback, which suggests repeating behaviour, the
Avoid cue itself can be taken for negative feedback, which usually translates into a response
switch. We have previously observed that this bias can dominate repeat/switch behaviour after
neutral outcomes, resulting in the paradoxical observation of more response repetition after
negative (i.e. no reward) feedback for Win cues than positive (i.e. no punishment) feedback to
Avoid cues (Algermissen et al., 2024). Such a bias should be invisible at the beginning of the
task and only arise with learning, leading to increasingly distorted learning curves over time.

We tested the presence of these three biases in behaviour using a series of seven
reinforcement learning models. The base model (M1) was a simple Q-learning model that

learned the values of Go and NoGo actions for each cue. To generate choices, action values
were turned into action probabilities via a softmax transform. Action values were subsequently
updated based on the obtained outcomes via reward prediction errors, which were scaled by
a free learning rate parameter ε and then added to the old action values. In line with previous
implementations (Algermissen et al., 2024; Guitart-Masip et al., 2012; Swart et al., 2017),
objective outcomes were scaled by a free feedback sensitivity parameter ρ. High values on
this parameter lead to learned action values developing further apart and thus more
deterministic choices, similar to a high inverse temperature parameter used in alternative
models. We extended this model with a constant Go bias parameter b to capture overall
tendencies towards Go or NoGo responses (M2), a response bias parameter π that added a
bonus/malus to the action value of Go depending on the valence of a cue (M3), and finally, a
learning bias parameter κ capturing increased learning rates for rewarded Go responses and
decreased learning rates for punished NoGo responses (M4). M5 combined all three
parameters and fitted the data better than any of the simpler models, suggesting that both a
response bias and learning bias were present in the data. Finally, we extended M5 by adding
a persistence parameter φ_{INT} that captured overall tendencies to repeat vs. switch actions (M6),
and a persistence bias φ_{DIFF} that captured the higher tendency to repeat actions for Win cues
than for Avoid cues (M7).

We compared these seven models using Bayesian model selection (Stephan et al., 2009). In
line with our previous work (Algermissen et al., 2024), M7, implementing all three Pavlovian
biases, was the best-fitting model according to the log-model evidence (Fig. 2G), the model
frequency (M7: 61%; M6: 14%; M5: 25%), and crucially, according to the protected
exceedance probability (M7: 98%; M5: 2%; Fig. 2K). We performed posterior-predictive checks
by simulating the action probabilities based on the fitted parameters combined with
participants' actions and outcomes (one-step ahead predictions), which captured the empirical
data patterns very well (Fig. 2H, I, J). Simpler models failed to capture core features of the
empirical data patterns. Model recovery showed that all models were distinguishable from each
other, and parameter recovery on M7 showed that all parameters were well identifiable
(Supplementary Fig. 6 & 7). We also explored alternative models in which cue valence did not
affect persistence directly, but biased the outcomes in the prediction error updating equation.
Such models predicted, qualitatively, a very similar pattern as M7, but showed slightly inferior
fit (Supplementary Fig. 5). In sum, these results confirmed the presence of all three Pavlovian
biases in participants' behaviour in the sham sessions.

Figure 2. Behavioural performance in the sham dataset. **A. Response bias:** Humans exhibit a tendency to show actions (i.e. button presses) for Win cues (blue), aligning reward seeking with action invigoration, and to withhold actions for Avoid cues (red), aligning punishment avoidance with action suppression/inaction. **B. Learning bias:** Learning is enhanced for rewarded Go actions compared to rewarded NoGo actions, promoting faster acquisition of Go actions from rewards. Conversely, learning is reduced for punished NoGo actions compared to punished Go actions, slowing the extinction of NoGo responses via punishments. **C. Persistence bias:** Seeing a Win cue partly acts as a reward in itself, leading to a higher tendency to repeat responses to such cues, even when a neutral outcome (negative feedback) suggests that one should switch the response. In contrast, seeing an Avoid cue partly acts as a punishment in itself, leading to a higher tendency to switch responses, even if a neutral outcome (positive feedback) suggests repeating the response. **D.** Trial-by-trial proportion of Go responses (error bands = ± SEM, n = 29) for Go cues (solid lines) and NoGo cues (dashed lines). A clear response bias is evident from the start, with participants making more Go responses to Win (green lines) than Avoid (red lines) cues. Participants show a general bias towards Go responses, with initial response rates around 80%. Additionally, instrumental learning is apparent in Go responses increasing for Go cues and decreasing for NoGo cues over time. **E.** Probability to repeat a response on the next encounter of the same cue as a function of the response and outcome for valenced outcomes (rewards, punishments) only (error bars are ±SEM across participants, n = 29, dots indicate individual participants). Learning was reflected in the higher probability of staying after rewarded responses than after being punished (main effect of outcome valence). Biased learning was evident in the stronger outcome effect after Go responses than after NoGo responses, indicating the presence of a

learning bias. **F.** Probability to repeat a response on the next encounter of the same cue as a function of cue valence (error bars are ± SEM across participants, n = 29, dots indicate individual participants). Participants showed a higher tendency to repeat responses for Win than Avoid cues, indicative of a *persistence* bias. **G.** Laplace approximation to the log-model evidence favours the model incorporating all three Pavlovian biases—response bias, learning bias, and persistence bias (M7)—over simpler models (M1–M6); error bars are ± SEM across participants, n = 29. **H-J.** Posterior predictive checks: one-step-ahead predictions based on the best fitting parameters (estimated with hierarchical Bayesian inference) for M7. **K.** Bayesian model selection. The winning model is M7 as determined by model frequency and protected exceedance probability, in line with log model evidence.

Sonication effects on Go/NoGo choices

We next analysed the effect of TUS on Pavlovian biases in responding and learning. We again first used regression models based on participants' Go/NoGo responses and their propensity to repeat/switch responses, and then subsequently contrasted parameters obtained from the best-fitting reinforcement learning model across all sonication conditions.

We used a mixed-effects logistic regression model to test for effects of TUS on (biased) responding, i.e., their Go/NoGo responses. This model included the required action, cue valence, the TUS sonication condition, as well as the block half (first vs. second half of a block) to account for the possibility that TUS might affect learning processes, which might only become visible in the second half of each block. Beyond the main effects of required action and cue valence and their interaction, this model yielded a significant 3-way interaction between required action, cue valence, and sonication condition, $\chi^2(2) = 14.277$, $p < .001$, a significant 3-way interaction between cue valence, sonication condition, and block half, $\chi^2(2) = 9.191$, $p = .010$, and a significant 4-way interaction between required action, cue valence, sonication condition, and block half, $\chi^2(2) = 21.548$, $p < .001$ (Fig. 3A, B, C). See Supplementary Results for results when analysing each cue condition (Go-to-Win, Go-to-Avoid, NoGo-to-Win, NoGo-to-Avoid) separately. To better understand these effects, we followed them up with the regression analyses of response repetitions vs. switches.

Figure 3. A-B Empirical Probability of Go. Trial-by-trial proportion of Go responses across sonication conditions (TUS-dACC, and TUS-aIns, grey lines represent the sham condition). TUS-dACC decreased Go responses in the Go-to-Win condition but increased them in the NoGo-to-Win condition. TUS-aIns decreased Go responses in the Go-to-Win condition but

increased them in the Go-to-Avoid and NoGo-to-Win conditions. Coloured lines display the
active sonication condition. **C-D Probability of Stay.** TUS effects on response repetitions
based on outcomes (rewarded Go vs. punished NoGo) and cue valence (Win vs. Avoid). TUS-
aIns (but not TUS-dACC) attenuated the tendency to repeat rewarded Go responses, but
increased the tendency to repeat punished NoGo responses compared to sham. TUS-dACC
(but not TUS-aIns) increased the cue valence effect on response repetitions, with stronger
persistence for Win cues relative to Avoid cues. **E-H** Posterior predictive checks reproduced
the empirical data patterns for the probability of Going and staying.

Sonication effects on response repetitions/switches

Analyses of Go/NoGo choices suggested effects of TUS on learning (i.e., changes in Go/NoGo
responses over time). To more directly investigate learning, i.e., the impact of outcomes on
subsequent choices, we analysed participants' repeat/switch choices as a function of
outcomes and responses on the current trial. We focused on the signatures of biased learning.

First, we contrasted sonication conditions selectively on those trials on which biased learning
should occur, namely trials with rewarded Go responses or with punished NoGo responses. If
the learning bias became stronger, participants would repeat rewarded Go responses more
and punished NoGo responses less, yielding a stronger difference between both conditions
(with no change in any other condition). Vice versa, if the bias became weaker, the difference
between both conditions would decrease. In a mixed-effects logistic regression model fit to the
response repetitions vs. switches, we observed a significant main effect of outcome (reward
vs. punishment) qualified by a significant outcome × sonication interaction, $\chi^2(2) = 6.162$, $p =$
.046: participants more often repeated rewarded Go responses than punished NoGo
responses and this tendency was significantly attenuated after TUS-aIns relative to sham, $b =$
-0.150, 95%-CI [-0.293, -0.006], $\chi^2(1) = 4.197$, $p = .040$, and not after TUS-dACC compared to
sham, $b = 0.001$, 95%-CI [-0.058, 0.249], $\chi^2(1) = 0.001$, $p = .984$ (Fig. 3C). These results
suggest that TUS-aIns, but not TUS-dACC, reduced the learning bias.

Second, we tested whether sonications changed the effect of cue valence on response
repetitions vs. switches, reflecting a bias in persistence. Participants were more likely to repeat
responses for Win than Avoid cues (main effect of valence: $b = 0.428$, 95%-CI [0.342, 0.514],
$\chi^2(1) = 94.620$, $p < .001$], which was qualified by a cue valence × sonication interaction, $\chi^2(2)$
= 7.617, $p = .022$: the cue valence effect was significantly stronger after TUS-dACC compared
to sham, $b = 0.113$, 95%-CI [0.031, 0.194], $\chi^2(1) = 7.368$, $p = .007$, with no difference between
TUS-aIns and sham, $b = 0.006$, 95%-CI [-0.063, 0.074], $\chi^2(1) = 0.025$, $p = .873$ (Fig. 3D). These
results suggest that TUS-dACC, but not TUS-aIns, affects the persistence bias. Next, we
directly confirmed these observations by fitting these parameters using computational
reinforcement learning models and comparing the fitted parameters across TUS sonication
conditions.

Effects of TUS on computational modelling parameters

We fit the best-fitting model from the sham session, M7, comprising all three Pavlovian biases,
namely a response bias π, a learning bias κ, and a persistence bias φ_{DIFF}, to the other two
sonication conditions. Model comparisons suggested that, in the TUS-dACC sessions, M7 was
again the best-fitting model according to both the model frequency (M7: 89%; M6: 11%) and
the protected exceedance probability (M7: 100%). In the TUS-aIns sessions, a slightly different
outcome occurred, with M6 being the best fitting model according to both the model frequency
(M6: 68%; M7: 32%) and protected exceedance probability (M6: 97%; M7: 3%). Given that M6
is a nested version of M7 (i.e., M7 with the φ_{DIFF} parameter fixed to zero is equivalent to M6),
we proceeded with comparing the parameter values in M7 across all three sonication
conditions.

We focused on the three parameters reflecting the three different bias parameters: the
response bias π, the learning bias κ (Fig. 4A), and the persistence bias φ_{DIFF} (Fig. 4D). We
compared the parameter values between all three conditions with a one-way repeated-
measures ANOVA, following up any significant effects with paired *t*-tests. There was no
significant difference in the response bias π, $F(2, 56) = 1.719$, $p = .189$, generalized $\eta^2 = 0.018$.
However, there was a significant difference in the learning bias κ, $F(2, 56) = 7.680$, $p = .001$,
$\eta^2 = 0.134$: this bias was significantly attenuated after TUS-aIns compared to sham, $t(28) =$
4.381, $p < .001$, Cohen's *d* = -1.072, 95%-CI [-1.687, -0.457], but not after TUS-dACC
compared to sham, $t(28) = 0.489$, $p = .629$, Cohen's *d* = -0.135, 95%-CI [-0.688, 0.419] (Fig.
4B). This observation is in line with the regression results showing an attenuated difference in
repetitions of rewarded Go vs. punished NoGo responses after TUS-aIns.

Furthermore, there was a significant difference in the persistence bias φ_{DIFF}, $F(2, 56) = 9.836$,
$p < .001$, $\eta^2 = 0.157$: this bias was significantly stronger after TUS-dACC compared to sham,
$t(28) = 4.028$, $p < .001$, Cohen's *d* = 1.031, 95%-CI [0.397, 1.667], but not after TUS-aIns
compared to sham, $t(28) = 1.058$, $p = .299$, Cohen's *d* = 0.214, 95%-CI [-0.624, 0.196] (Fig.
4F). This effect was in line with the increased cue valence effect on response repetitions vs.
switches after TUS-dACC. In sum, TUS-aIns attenuated the learning bias, while TUS-dACC
increased the persistence bias.

Beyond the bias parameters of interest, both TUS conditions also unspecifically increased the
Go bias parameter, $F(2, 56) = 4.958$, $p = .010$, $\eta^2 = 0.071$ (Fig. 4C), and the persistence
parameter φ_{INT}, $F(2, 56) = 3.517$, $p = .036$, $\eta^2 = 0.055$ (Fig. 4E; neither significant when
correcting for multiple comparisons): Relative to sham, the overall Go bias *b* was significantly
increased after both TUS-dACC, $t(28) = 3.148$, $p = .004$, Cohen's *d* = 0.720, 95%-CI [0.206,
1.233], and TUS-aIns, $t(28) = 2.130$, $p = .042$, Cohen's *d* = 0.438, 95%-CI [0.007, 0.869].
Similarly, the overall persistence parameter φ_{INT} was significantly increased after TUS-dACC,
$t(28) = 3.070$, $p = .005$, Cohen's *d* = 0.639, 95%-CI [0.181, 1.096], and marginally significantly
increased after TUS-aIns, $t(28) = 1.972$, $p = .059$, Cohen's *d* = 0.429, 95%-CI [0.026, 0.884]
(not significant when correcting for multiple comparisons). There was no significant difference
in the feedback sensitivity parameter ρ, $F(2, 56) = 0.928$, $p = .401$, $\eta^2 = 0.015$, or the learning
rate ε, $F(2, 56) = 1.047$, $p = .358$, $\eta^2 = 0.018$.

Taken together, the effects of dACC and aIns sonication were best captured by a decrease in
the learning bias after TUS-aIns, an increase in the persistence bias after TUS-dACC, and
increases in overall Go biases and persistence after both types of sonications. These effects
were well captured by posterior predictive checks: Simulated learning curves closely followed
the empirical learning curves (Fig. 3E, F). These observations also match the regression
results of response repetitions/switches, with an attenuated difference between rewarded Go
and punished NoGo responses after TUS-aIns (Fig. 3G) and a stronger cue valence effect
after TUS-dACC (Fig. 3H). In sum, in this study, we found a double dissociation between the
effects of TUS-aIns and TUS-dACC, with both altering different Pavlovian biases.

Figure 4. Effects of TUS on instrumental learning biases, as reflected by computational model parameters. A. Learning bias illustration: Learning is stronger for rewarded Go actions, and weaker for punished NoGo actions. **B**. Learning bias (κ): TUS-aIns (but not TUS-dACC) significantly diminished the learning bias, i.e., the tendency to attribute rewards to actions and the reluctance to attribute punishments to inactions. **C.** Go bias (b): both sonication conditions significantly increased participants' overall tendency for Go responses regardless of the cue valence. **D.** Persistence bias illustration: participants show an overall higher tendency to repeat actions for Win cues than for Avoid cues. For Win cues, this implies that they sometimes ignore neutral outcomes (negative feedback, which suggests to switch the response) and instead take the Win cue itself as positive feedback (which suggests to repeat the response). **E.** Persistence parameter (φ_{INT}): both sonication conditions increased participants' overall tendency to repeat previous responses. **F.** Persistence bias (φ_{DIFF}): TUS-dACC (but not TUS-aIns) significantly increased the persistence bias, leading to a stronger tendency to repeat responses relatively more for Win cues than for Avoid cues. Asterisks indicate significance levels: ***$p < 0.001$; **$p < 0.01$; *$p < 0.05$; † $p < 0.10$.

Discussion

This study investigated the effects of TUS to the dACC and aIns on Pavlovian biases using a Motivational Go/NoGo task. Participants showed three forms of Pavlovian biases: a response bias, learning bias, and persistence bias (Swart et al., 2017; Algermissen et al., 2024). TUS revealed a double dissociation, with dACC and aIns affecting distinct learning processes: TUS-aIns attenuated a learning bias, reducing the tendency to attribute rewards to actions and reluctance in attributing punishments to inactions, while TUS-dACC increased a persistence bias, strengthening the tendency to use cue valence as a reinforcer signal and persist after positive (compared to negative) cues. Given that dACC and aIns are part of the same resting-state network, the saliency network (Seeley, 2019; Taylor et al., 2008), and regularly co-activate in cognitive tasks (Kurth et al., 2010; Medford and Critchley, 2010), previous research has assigned largely similar roles to these regions. Our study contributes to understanding their different roles in learning from feedback: While the dACC appear important in selecting which environmental signals to use as feedback, the aIns seems important in credit assignment whether these signals were caused by an agent's own actions.

We set out this study under the hypothesis that perturbing neural activity in dACC and aIns, which are part of an early "reset" network that rapidly responds to negative/salient stimuli (Fouragnan et al., 2015), could alter the strength of Pavlovian biases. In particular, several previous studies have observed a selective role of dACC and aIns in choices to avoid losses (Cecchi et al., 2024; Palminteri et al., 2012), learning from negative feedback (Billeke et al., 2020; Horing and Büchel, 2022), and adjusting behaviour (McGuire and Kable, 2015; van Geen et al., 2025). However, in our results, TUS did not selectively affect responses to negative (Avoid) cues. Specifically, TUS did not change the cue valence effect (i.e., response bias), the required action effect (i.e., overall accuracy), or their interaction (i.e., a change in accuracy for only Avoid cues). Similarly, in our computational models, (baseline) learning rates and the response bias were unaffected by TUS. Instead, the effects of TUS only became apparent over time, indicative of changes in learning processes, and were spread across all four cue conditions. These findings reflect more intricate changes in learning that were well captured by a computational model (Algermissen et al., 2024). This finding might not be surprising given a broad literature implicating dACC and aIns in learning from feedback and adjusting action policies (Alexander and Brown, 2015; Behrens et al., 2007; Fouragnan et al., 2019).

We found that stimulating the aIns with TUS led to a reduction in a learning bias: participants showed a reduced tendency (i.e., lower learning rate) to ascribe rewards to their own actions, as well as a reduced reluctance to ignore punishments of their own inactions. Such biases have been interpreted as global "priors" on which action-outcome relationships likely hold in an organism's environment (Algermissen et al., 2024; Swart et al., 2017). Using such priors, individuals can trade-off noisy outcomes of single actions against prior hypotheses on what is generally the best response strategy. Such biases will facilitate the acquisition of actions that are conducive to rewards but slow down the acquisition of actions in contexts in which punishments are involved. While such biases will usually be adaptive and lead to more robust learning, the tendency to attribute rewards to own actions can lead to the false inference of "illusionary control" over outcomes that are in fact random (Aarts et al., 2005; Ebert and Wegner, 2011). Similarly, these biases could explain why humans judge actions that lead to negative outcomes more harshly than omissions leading to such outcomes ("omission bias") (Spranca et al., 1991), a phenomenon cited to explain some humans' reluctance to vaccinate (Ritov and Baron, 1995, 1990).

We have previously found that neural signals associated with such biased learning are first visible in the blood-oxygen-level-dependent (BOLD) signal from cortical regions including the perigenual anterior cingulate cortex (pgACC) and posterior cingulate cortex (PCC), prior to subcortical regions (striatum) (Algermissen et al., 2024). Previous research has indeed suggested cortical regions, most prominently the dACC, to control the speed of learning (learning rates) in subcortical regions (Alexander and Brown, 2015; Behrens et al., 2007; Meder et al., 2017). Although related bias phenomena, such as auto-shaping and negative self-maintenance, have been shown in animals such as rodents or pigeons (Brown and

Jenkins, 1968; Williams and Williams, 1969), we have previously speculated that a prefrontal,
cortical basis of this bias in humans might hint at a more recent innovation in primates that is
responsible for this particular learning bias.

Given the previous literature on the role of the dACC in scaling learning rates (Behrens et al.,
2007; O'Reilly et al., 2013; Piray et al., 2019b), we would have a priori expected that TUS
dACC should have affected the learning bias, and a selective reduction under TUS-aIns might
at first appear puzzling. The aIns has previously been found to encode the uncertainty/risk
under which a choice is made (Hsu et al., 2005; Preuschoff et al., 2008; Vilares et al., 2012)
as well as the average reward rate in an environment (Wittmann et al., 2020), which are key
environmental variables that should scale learning rates according to normative models (Cazé
and van der Meer, 2013; Piray and Daw, 2021; Yu and Dayan, 2005). Overall, these findings
suggest that aIns plays a crucial role in using higher-order beliefs about the environment to
decide how much to update lower order beliefs based on reward/punishment feedback. This
supports a role of the aIns in scaling learning rates, which can give rise to learning biases.

An alternative possibility is that the TUS-aIns effect reducing biased credit assignment arises
from its role in social cognition and self-other attribution. Increased aIns activity has been
shown both when an agent is directly affected by an outcome (e.g., pain, unpleasant taste)
and when they observe such outcomes in another individual (Jabbi et al., 2008; Lamm and
Singer, 2010; Singer et al., 2004). Hence, the computational challenge arises for the aIns to
tell apart which outcomes were caused by oneself and which were merely observed in others
(Jabbi et al., 2008). TUS-aIns might blur this boundary, leading to changes in credit assignment
as observed in the current study. Some studies have speculated that this role is subserved by
von Economo and Fork neurons, which, within the primate brain, occur almost exclusively in
the aIns (Allman et al., 2010, 2005) and have only been observed in animal species that live
in large social groups (Evrard, 2019). This speculative interpretation suggests a dedicated role
of the aIns in credit assignment, following an evolutionary route that might be special in
primates. Investigating the causal contribution of such deep cortical circuits to learning from
feedback might greatly benefit from the use of non-invasive brain stimulation tools such as
TUS. Taken together, our results suggest that aIns plays an important role in credit assignment
and deciding whether environmental feedback signals were caused by personal actions.

In contrast to TUS-aIns, TUS-dACC selectively affected the persistence bias, with an
increased tendency to repeat actions for Win cues than Avoid cues, regardless of the actions
and outcomes they had been followed by. This bias reflects participants' tendency to use the
cue valence instead of the outcome valence as a feedback signal (Algermissen et al., 2024).
Knowledge of the cue valence is important for disambiguating neutral outcomes, which can
only be used as a teaching signal when contrasted against the counterfactual that could
otherwise have been obtained (i.e., a neutral outcome signalling a missed reward is negative
feedback; a neutral outcome signalling a missed punishment is positive feedback). However,
human participants show limited ability to tell apart counterfactual and real outcomes,
sometimes taking a neutral outcome obtained instead of a reward as a positive outcome, and
a neutral outcome obtained instead of a punishment as a negative outcome (Algermissen et
al., 2024), a tendency exacerbated by TUS-dACC. Previous studies have implicated a causal
role of the dACC in learning from counterfactual feedback (Fouragnan et al., 2019) and
deciding when it is worth switching an action policy (Hayden et al., 2011; Karlsson et al., 2012;
Kolling et al., 2012). When perturbed with TUS, the dACC might mediate a confusion between
real and counterfactual feedback, leading to a decreased ability to switch behaviour after a
missed reward, which gives rise to larger persistence bias. In sum, our results suggest that the
dACC is important in selecting which environmental signals to use as reinforcer to adjust
behaviour.

Our findings also highlight the promise of TUS as a neuromodulation tool in cognitive
neuroscience. Compared to other methods, such as TMS or transcranial direct current
stimulation (tDCS), TUS allows for deeper, more focal targeting of subcortical and
midline/lateralised deep cortical structures, making it particularly suited to probing the causal

roles of regions like the aIns and dACC. As such, this method opens new possibilities for
investigating the neural mechanisms of learning and decision-making.

In summary, our study reveals a double dissociation in the effects of TUS on learning biases,
with TUS-aIns selectively attenuating the learning bias and TUS-dACC increasing the
persistence bias. Hence, while dACC and aIns often co-activate in cognitive roles, they appear
to have distinct roles in learning from feedback: while TUS to the aIns decreased people's
tendency to overly take credit for rewards following action and to ignore punishments following
inaction, TUS to dACC increased participants' tendency to take the cue valence as a reinforcer
signal. These findings advance our understanding of the differential roles of deep cortical
regions in instrumental learning and highlight the potential of TUS as a tool for modulating
cognitive and affective processes. Our findings challenge the traditional view that learning
biases are exclusively driven by subcortical regions associated with rigid, habit-like behaviours.
Instead, these results emphasize the role of frontal inputs, which contribute to counterfactual
reasoning and behavioural flexibility (Behrens et al., 2007; Fouragnan et al., 2019). The
involvement of these two areas in biased learning suggests that these biases may not be rigid
constraints, but rather adaptive priors that guide how individuals integrate past experiences.
By balancing prior beliefs about action-outcome associations with flexible learning from
rewards and punishments, these biases may enhance decision-making in dynamic
environments (Algermissen et al., 2024; Algermissen and den Ouden, 2023).

Methods

Participants

34 healthy participants (18 females, 16 males) aged 19 to 59 years (mean age = 29.73, SD = 11.16) took part in this study. One participant was excluded because they did not understand the task and three because of missing sessions. 29 participants (16 females, 13 males), aged 19 to 59 years (mean age = 30.24, SD = 11.63) were included in the analysis. Participants had normal or corrected-to-normal vision and were carefully screened for any counter indications to TUS and MRI, with none reporting a current diagnosis of neurological or psychiatric disorders or use of psychoactive medications. All participants provided written informed consent following a full explanation of the study procedures. Ethical approval was granted by the University of Plymouth Faculty of Health Staff Research Ethics and Integrity Committee (reference ID: 3394; date: 28/06/22). Participants were compensated an average of £85, which included a performance bonus of up to £10 based on points collected over three sessions in the Motivational Go/NoGo task. Travel expenses were reimbursed up to £5 per session. All study sessions took place at the Brain Research & Imaging Centre (BRIC) in Plymouth, UK.

Study design

The study design is summarised in Fig. 1A. Participants began with an MRI-only session, which included a series of scans, including a T1-weighted MRI scan. This imaging data set was used to create a personalised head model for neuronavigation and acoustic and thermal simulations, enabling precise planning of the TUS target and transducer placement for the upcoming TUS sessions. In the first session, participants completed a practice phase to become familiar with the Motivational Go/NoGo learning task.

For the TUS sessions, participants received a randomised sequence of TUS conditions. Each session involved an 80-second TUS application with three different conditions: active stimulation of the bilateral anterior insula (TUS-aIns), active stimulation targeting the dorsal anterior cingulate cortex (TUS-dACC), and a sham condition. In the sham condition, the transducer was positioned over the temporal region bilaterally, similar to the TUS-aIns setup, but no ultrasound was delivered. To maintain a consistent procedure, both sonication and sham conditions were always applied first to the left hemisphere, followed by the right.

MRI data acquisition

For TUS planning, including neuronavigation and acoustic simulations, MRI scans were acquired using a Siemens MAGNETOM Prisma 3T scanner (syngo MR V11E, Siemens Healthineers, Erlangen, Germany) equipped with a 64-channel head coil. The scanning protocol for this study included a T1-weighted MPRAGE sequence in the sagittal plane (TR = 2100 ms, TE = 2.26 ms, inversion time = 900 ms, flip angle = 8°, GRAPPA acceleration factor = 2, matrix size = 256 × 256, 176 slices, voxel size = 1 × 1 × 1 mm³).

TUS protocol and procedure

Targeting of the bilateral aIns and dACC was guided by personalised anatomical landmarks using each participant's T1-weighted MRI scans during the planning of target and transducer placement (see Supplementary Table 1). Acoustic field simulations, generated from pseudo-CT skull models derived from participants' T1-weighted images using a convolutional neural network (CNN) (Yaakub et al., 2023), ensured precise targeting and safety by keeping both on-target and off-target pressure within acceptable ranges.

A bespoke CTX-500 NeuroFUS TPO system (Brainbox Ltd., Cardiff, UK) with a four-element annular transducer (diameter = 64 mm, central frequency = 500 kHz, and steering range

between 27.3 and 82.6mm) (Supplementary Fig. 1) was used to deliver the TUS protocol. The
TUS protocol consisted of repetitive TUS applied using a 5 Hz-patterned protocol with a 10%
duty cycle, lasting 80 seconds and delivering a total of 400 pulses (pulse duration of 20 ms
and pulse interval of 200 ms). The initial spatial-peak pulse-average intensity (ISPPA) in free-
field conditions was set at 50 W/cm² for all participants. We took care to remain within the
guidelines for human ultrasound exposure as defined by ITRUSST (Aubry et al., 2024). The
results of our acoustic simulations predominantly indicated a maximum skull temperature rise
below 2°C (Supplementary Table 1). In cases where this threshold was exceeded, we
calculated the CEM43, a metric reflecting both duration and intensity of heating relative to
43°C, the critical threshold for thermal cell damage. We ensured that CEM43 values remained
well below 0 (Aubry et al., 2024). In our study, CEM43 was always below 0.1.

To improve ultrasound transmission, a layer of ultrasound gel (Aquasonic 100, Parker
Laboratories Inc.) was applied at the transducer placement site, with a 2 cm gel pad (Aquaflex,
Parker Laboratories Inc.) positioned between the transducer and the participant's head.
Participants' heads remained unshaved, and any air bubbles were carefully eliminated by
smoothing and combing the hair. Neuronavigation was performed using Brainsight v2.5
(Rogue Research Inc., Montréal, Québec, Canada) with T1-weighted anatomical MR images.
During each session, focal depth measurements obtained from Brainsight were entered into
the NeuroFUS TPO system (Brainbox Ltd., Cardiff, UK) before stimulation, and the trajectory
was sampled to support confirmatory acoustic simulations following each session.

Blinding procedure

713 Sham TUS was administered identically to active TUS, except that no actual stimulation was
714 applied. Instead, participants listened to a sound mimicking the transducer's pulse repetition
715 frequency through bone conduction headphones, positioned approximately 2 cm posterior and
716 superior to the temples. This sound was only played during the Sham condition to ensure
717 blinding, while headphones were worn in both active and Sham sessions.

Task

The task consisted of four blocks, presented approximately 15, 25, 35, and 45 minutes after
TUS (Fig. 1B, C). Each block featured four cues (gems), with each trial requiring participants
to determine, through trial and error, the correct action (Go or NoGo) and the potential outcome
(reward or punishment) associated with each stimulus. The task design involved four different
cues that varied by cue valence (Win vs. Avoid) and required action (Go vs. NoGo),
counterbalanced across participants in three sessions of 320 trials each. Motivationally
incongruent cues were included, where participants' natural action tendencies (e.g., to act for
a reward) conflicted with the task requirements (e.g., to refrain from action when a reward was
expected). For each cue, correct responses resulted in positive outcomes (80% chance of
reward or avoiding punishment), whereas incorrect actions had only a 20% chance of yielding
positive outcomes. The valence (Win or Avoid) of each cue was not explicitly signalled and
had to be learned over time.

This probabilistic task was designed to capture learning and decision-making under
uncertainty, similar to real-world scenarios where outcomes are not guaranteed. The structure
of probabilistic feedback allowed us to observe participants' flexibility in adjusting strategies,
sensitivity to feedback, and the cognitive and neural processes underlying decisions when
faced with motivational conflicts, such as balancing exploration versus exploitation in uncertain
environments.

Stimuli display

The probabilistic motivational learning task was displayed on an LCD screen (1920 × 1080
resolution, 60 Hz refresh rate, 8-bit colour depth, RGB, standard dynamic range (SDR)),
positioned 0.5 m in front of participants. The experiment was programmed using the
Presentation software (Neurobehavioral Systems Inc., Berkeley, CA, USA) and run on a
Windows-based computer. Participants responded using their preferred hand by pressing the
SPACE bar on a keyboard.

Post-TUS questionnaire

After each TUS session, as well as on the following day, participants were asked to report any
adverse effects. They were presented with a list of symptoms and asked to rate the intensity
to which they experienced each of them using a 4-point scale (absent, mild, moderate, severe)
and whether they thought their experience was related to the stimulation on a 5-point scale
(unrelated, unlikely, possible, probable, definite). Symptoms included those reported in (Legon
et al., 2020), to which we added speech problems, vision problems, and muscle tightness of
the face or arm. Respondents had the possibility of listing and rating up to two additional
symptoms and to provide details for their experience.

For further insights, Supplementary Table 1 present detailed information regarding the acoustic
simulation parameters and the outcomes observed across participants. This includes data on
mechanical indices, derated ISPPA values at the focal point, and the incidence of side effects
associated with both aIns and dACC stimulation.

Regression analyses

762 We fit mixed-effects logistic regression models using the lme4 package in R. We always
763 included a random intercept per subject and all possible random slopes and random
764 correlations, achieving a maximal random effects structure (Barr et al., 2013). We used sum-
765 to-zero coding for all categorical predictors. We obtained p-values from Wald chi-square tests
766 using Type-3 sums-of-squares via the Anova() function from the car package.

Computational modelling and model comparison

We fit a series of increasingly complex computational reinforcement learning models to
participants' choices, following previous studies (Algermissen et al., 2024; Guitart-Masip et al.,
2012; Swart et al., 2017). All models are Q-learners that learn the expected value of performing
an action for a given cue using reward prediction errors and select actions with higher expected
value. On a given trial t, the base model **M1** computes choice probabilities for Go and NoGo
actions (a) given a cue (s) using action weights (w; modified Q-values) turned into probabilities
via a softmax transform:

$$p(a_t|s_t) = \frac{\exp(w(a_t, s_t))}{\sum_a \exp(w(a\prime, s_t))} \tag{1}$$

For each action, the model obtains an outcome r and computes a reward prediction error as
the difference between the outcome r and the expected value Q(a, s). It then uses prediction
errors to update Q-values via the delta learning rule:

$$Q_t(a_t, s_t) = Q_{t-1}(a_t, s_t) + \varepsilon(\rho r - Q_{t-1}(a_t, s_t)) \tag{2}$$

Outcomes can be +1 (reward) or 0 (no reward) for Win cues, or 0 (no punishment) or -1
(punishment) for Avoid cues. In this learning rule, prediction errors are scaled by the learning
rate ε, with large learning rates leading to faster updating (and a stronger recency bias in value
estimates) and small learning rates leading to slower updating (and a more long-lasting impact
of outcomes obtained far in the past). Furthermore, outcomes themselves are scaled with a
participant-specific feedback sensitivity parameter ρ, which plays a similar role as an inverse

temperature parameter: high values of ρ lead to Q-values for Go and NoGo actions developing
further apart with learning and leading to more deterministic choices, while small values of ρ
lead to Q-values closer together and thus more stochastic choices. We initialized Q-values to
the midpoint between the two possible outcomes of each cue (+0.5 for Win cues, -0.5 for Avoid
cues) multiplied with participants' respective feedback sensitivity parameter such that starting
Q-values are the midpoint of each participant's subjective value space. Given that the valence
of each cue (Win/Avoid) was not instructed but had to be inferred from the attainment of
valenced outcomes (rewards/punishments), participants could not interpret neutral outcomes
as positive/negative feedback before having observed such a valence outcome. During this
phase, Q-values were updated based on feedback, but "muted" in the action selection process
(multiplied with zero when computing choice probabilities), assuming that participants could
retrospectively infer the meaning of neutral outcomes once they determined the valence of a
given cue. Upon the first encounter of a valenced outcome (reward, punishment), Q-values for
this cue were "unmuted" and subsequently used for action selection.

We extended this base model using several bias terms. In **M2**, we added a Go bias term b to
the Q-values of making a Go response for each cue, capturing participants' overall propensity
of making a Go response:

806
$$w(a_t, s_t) = \begin{cases} Q_t(a_t, s_t) + b & if\ a = Go \\ Q_t(a_t, s_t) & else \end{cases} \qquad (3)$$

808 In **M3**, in addition to the Go bias, we added the response bias π multiplied with the cue valence
809 V (+0.5 for Win; -0.5 for Avoid; value of 0.5 arbitrarily chosen for scaling) to the Q-values of
810 making a Go response:

811
$$w(a_t, s_t) = \begin{cases} Q_t(a_t, s_t) + b + \pi V(s) & if\ a = Go \\ Q_t(a_t, s_t) & else \end{cases} \qquad (4)$$

Note that the cue valence was fixed and participants instructed that each cue was either a Win
or an Avoid cue. Hence, no incremental learning of the cue valence was required, but the
valence could be inferred as soon the first valenced outcome was encountered. Similar to early
learning from neutral outcomes, the response bias was muted (multiplied with zero) up until
the first encounter of a valenced outcome and only subsequently unmuted.

In **M4**, instead of a response bias, we added a learning bias κ, capturing participants' tendency
to use an increased learning rate for rewarded Go actions (i.e., attribute rewards to their own
actions) and a decreased learning rate for punished NoGo actions (i.e., an unwillingness to
attribute punishments to own inactions):

822
$$\varepsilon = \begin{cases} \varepsilon_0 + \kappa & if\ r_t = 1\ and\ a = go \\ \varepsilon_0 - \kappa & if\ r_t = -1\ and\ a = nogo \\ \varepsilon_0 & else \end{cases} \qquad (5)$$

Since learning rates were sampled in a continuous space and later inverse-logit transformed
to constrain them to the range of [0, 1], the impact of κ might be asymmetric after the
transformation. To achieve a symmetric impact of κ, we first determined whether the base
learning rate ε_0 was smaller or bigger than 0.5, then computed one half of the bias (i.e., the
side closer to the range limits), took the difference between the base and the biased learning
rate, and used this difference to compute the other half of the bias:

830
$$\varepsilon = \begin{cases} \varepsilon_0 = inv.logit(\varepsilon) \\ \varepsilon_{punished\ NoGo} = inv.logit(\varepsilon - \kappa) & if\ \varepsilon_0 < 0.5 \\ \varepsilon_{rewarded\ Go} = \varepsilon_0 + (\varepsilon_0 - \varepsilon_{punished\ NoGo}) & if\ \varepsilon_0 < 0.5 \end{cases} \qquad (6)$$

$$\varepsilon = \begin{cases} \varepsilon_0 = inv.\, logit(\varepsilon) \\ \varepsilon_{rewarded\ Go} = inv.\, logit(\varepsilon + \kappa) & if\ \varepsilon_0 > 0.5 \\ \varepsilon_{punished\ NoGo} = \varepsilon_0 - (\varepsilon_{rewarded\ Go} - \varepsilon_0) & if\ \varepsilon_0 > 0.5 \end{cases} \tag{7}$$

In **M5**, we added both the response bias π and the learning bias κ, which provided a better fit than each bias in isolation.

In **M6**, to capture participants' overall tendency to repeat responses irrespective of the received outcome, we extended M5 by adding an intercept persistence parameter φ_{INT} to the weight of the action performed on the last encounter of the same cue:

$$w(a_i, s_t) = \begin{cases} w(a_i, s_t) + \varphi_{INT} & if\ last\ action\ to\ same\ cue\ s\ was\ a_i \\ w(a_i, s_t) & else \end{cases} \tag{8}$$

Finally, in **M7**, to capture participants' higher tendency to repeat responses for Win than for Avoid cues, we added the persistence bias φ_{DIFF}, which was added to the weight of the last performed action in case of a Win cue and subtracted from this weight in case of an Avoid cue:

$$w(a_i, s_t) = \begin{cases} w(a_i, s_t) + \varphi_{INT} + \varphi_{DIFF} & if\ Win\ cue\ and\ last\ action\ to\ same\ cue\ s\ was\ a_i \\ w(a_i, s_t) + \varphi_{INT} - \varphi_{DIFF} & if\ Avoid\ cue\ and\ last\ action\ to\ same\ cue\ s\ was\ a_i \\ w(a_i, s_t) & else \end{cases} \tag{9}$$

We fitted models using weekly informative hyperpriors to $X_\rho \sim \mathcal{N}(2,3)$, $X_\varepsilon \sim \mathcal{N}(0,2)$, $X_{b,\pi,\kappa} \sim \mathcal{N}(0,3)$ in line with previous studies (Algermissen et al., 2024; Swart et al., 2017). We constrained feedback sensitivities ρ to be positive using the exponential transform and learning rates ε to the range [0, 1] using an inverse the inverse-logit transform. Furthermore, we constrained κ and φ_{DIFF} to be positive in line with the hypothesized direction of the learning and persistence biases (Fig. 2B, C) using the $y = log(1 + exp(x))$ transform, which is y = 0 for negative numbers, smoothly asymptotes 0 for small positive numbers, and is roughly y = x for large positive numbers. Leaving these parameters unconstrained led to the same qualitative conclusions, but a slightly inferior fit. In line with recent recommendations (Valton et al., 2020), we fitted the data of each sonication condition separately.

We fitted all models using the CBM toolbox in MATLAB (Piray et al., 2019a) using three steps: In the first step, we fitted the respective model to each participants' data individually using an iterative expectation-maximization algorithm that uses the Laplacian approximation for estimating the parameter posteriors. Chances of getting stuck in a local minimum were mitigated by using multiple parameter initialisations. In a second step, we used the individual fits of all participants as the starting point in a hierarchical Bayesian inference procedure. Group-level parameters were iteratively fitted using mean-field variational Bayes and served as hyperpriors for the individual participants' parameters. Thus, the parameter values of each participant were constrained/informed by the parameter values of all other participants, leading to more robust estimates and a lower chance of overfitting. We used these parameter values for all our posterior predictive checks. Finally, in a third step, we fitted all candidate models for all participants in one single step by iteratively (a) determining the probability that a given participant X was best fitted by a given model Y and (b) fitting the group-level parameters based on (only) those participants whose behaviour was in fact best characterized by model Y (i.e., parameters weighted proportionally to participants' latent "model responsibility" weights). This approach allows for simultaneous hierarchical Bayesian inference (with participants constraining each other's parameter values) and random-effects model selection (allowing for the possibility that different participants are best characterized by different models). Based on this last step, we computed the model frequency and protected exceedance probability of each model and performed Bayesian model selection (Stephan et al., 2009), selecting the model with the highest exceedance probability as the winning model.

We compared parameters across conditions using one-way repeated-measures ANOVAs as
implemented in the ezAnova() function of the ez package in R. We computed paired *t*-tests
using the t.test() function from the stats package in R and computed paired-samples Cohen's
d using the Cohen.d() function from the effsize package in R.

To validate that our best fitting model M7 could reproduce key patterns observed in the
empirical data, we performed posterior predictive checks (Nassar and Frank, 2016; Palminteri
et al., 2017) using one-step-ahead predictions (Steingroever et al., 2014). We used each
participant's best fitting parameters from the hierarchical Bayesian fit and their actual choices
and outcomes from the empirical data to simulate the action probabilities of each participant
on each trial, which we used as synthetic data to then plot the proportions of Go
response/response repetitions the same way as the empirical data.

To confirm that we could reliably measure individual differences in model parameters, we
performed parameter recovery (Algermissen and den Ouden, 2024b; Wilson and Collins,
2019). We first fitted a multivariate normal distribution to the best fitting parameter values of
M7, sampled 1,000 new combinations of parameter values from this distribution, simulated
new data based on this parameter combinations, and fitted M7 to each simulated data set. We
then correlated the sampled "ground-truth" parameters to the fitted "recovered" parameters.
To test whether these correlations were significantly higher than expectable by chance, we
compared them against a permutation null distribution obtained by randomly permuting the
assignment of ground-truth to recovered parameter values 10,000 times and computing the
95^{th} percentile of this distribution as the upper bound of a one-sided confidence interval.

To confirm that we could reliably detect the best fitting model for a given data set, we performed
model recovery (Algermissen and den Ouden, 2024b; Wilson and Collins, 2019). For each of
the seven candidate models, we fitted a multivariate normal distribution to the best fitting
parameter values and then sampled 1,000 new parameter value combinations for each model
(in total 7,000 parameter sets). We applied the following parameter constraints: $\rho < 400$
(otherwise possibility of infinitely large numbers), $\varepsilon > 0.05$ (otherwise too little learning), and ρ,
π, κ, φ_{INT} and φ_{DIFF} being far enough away from zero (discarding the lowest 10% of absolute
sampled parameter values). These constraints were important to ensure that each data set
expressed characteristics of the respective generating model; otherwise, if one of the bias
parameters was too close to zero, a more complex model would effectively reduce to a simpler
model. We then simulated a new data set for each parameter combination (in total 7,000
simulated data sets) and fitted each of the seven models to each simulated data set (in total
49,000 fits). For each data set, we used the log model evidence to determine which model
fitted the data best. We then computed the forward confusion matrix with the conditional
probabilities (relative frequencies) of model Y emerging as a best fitting model for a data set
generated by model X. We also computed the inverse confusion matrix with the conditional
probabilities of model X being the generative model for a data set best fitted by model Y. To
test whether the on-diagonal probabilities of these matrices were significantly higher than
expectable by chance, we created a permutation null distribution by randomly permuting the
log model evidences for the different fits to a given data set, computing the respective
confusion matrices, and saving the on-diagonal probabilities. We again computed the 95^{th}
percentile of these distributions as upper bounds of one-sided confidence intervals.

Data availability

The data files are available as .csv and .mat files on the OSF repository https://osf.io/pux6s/?view_only=c7dab129b4f246ed937e16e5941f04da.

Code availability statement

Analyses code to reproduce the presented acoustic simulations as well as behavioural, RS-fMRI, MRS, and self-report questionnaire results are in a GitHub repository under https://github.com/johalgermissen/mgng_tus_dacc_ains.

Competing interests

All authors declare that they have no conflicts of interest to disclose.

Acknowledgement

Nomi Koutsoumpari has received pump priming funds from the School of Psychology, and from the Bain Research Imaging Center, University of Plymouth for this study. Elsa Fouragnan is funded by a UKRI FLF (MR/Y034368/1), a BBSRC (BB/Y001494/1), a Neuromod+ grant (EP/W035057/1) and an ARIA grant (SCNI-PR01-P15).

References

Aarts H, Custers R, Wegner DM. 2005. On the inference of personal authorship: Enhancing experienced agency by priming effect information. *Consciousness and Cognition* **14**:439–458. doi:10.1016/j.concog.2004.11.001

Alexander WH, Brown JW. 2015. Hierarchical error representation: A computational model of anterior cingulate and dorsolateral prefrontal cortex. *Neural Computation* **27**:2354–2410. doi:10.1162/NECO_a_00779

Algermissen J, den Ouden HEM. 2024a. Pupil dilation reflects effortful action invigoration in overcoming aversive Pavlovian biases. *Cogn Affect Behav Neurosci* **24**:720–739. doi:10.3758/s13415-024-01191-y

Algermissen J, den Ouden HEM. 2024b. High stakes slow responding, but do not help overcome Pavlovian biases in humans. *Learn Mem* **31**:a054017. doi:10.1101/lm.054017.124

Algermissen J, den Ouden HEM. 2023. Goal-directed recruitment of Pavlovian biases through selective visual attention. *Journal of Experimental Psychology: General* **152**:2941–2956. doi:10.1037/xge0001425

Algermissen J, Swart JC, Scheeringa R, Cools R, den Ouden HEM. 2024. Prefrontal signals precede striatal signals for biased credit assignment in motivational learning biases. *Nat Commun* **15**:19. doi:10.1038/s41467-023-44632-x

Algermissen J, Swart JC, Scheeringa R, Cools R, den Ouden HEM. 2022. Striatal BOLD and midfrontal theta power express motivation for action. *Cerebral Cortex* **32**:2924–2942. doi:10.1093/cercor/bhab391

Allman JM, Tetreault NA, Hakeem AY, Manaye KF, Semendeferi K, Erwin JM, Park S, Goubert V, Hof PR. 2010. The von Economo neurons in frontoinsular and anterior cingulate cortex in great apes and humans. *Brain Struct Funct* **214**:495–517. doi:10.1007/s00429-010-0254-0

Allman JM, Watson KK, Tetreault NA, Hakeem AY. 2005. Intuition and autism: a possible role for Von Economo neurons. *Trends in Cognitive Sciences* **9**:367–373. doi:10.1016/j.tics.2005.06.008

Aubry J-F, Attali D, Schafer M, Fouragnan E, Caskey C, Chen R, Darmani G, Bubrick EJ, Sallet J, Butler C, Stagg C, Klein-Flügge M, Yoo S-S, Treeby B, Verhagen L, Pauly KB. 2024. ITRUSST consensus on biophysical safety for transcranial ultrasonic stimulation. doi:10.48550/arXiv.2311.05359

Barr DJ, Levy R, Scheepers C, Tily HJ. 2013. Random effects structure for confirmatory hypothesis testing: Keep it maximal. *Journal of Memory and Language* **68**:255–278. doi:10.1016/j.jml.2012.11.001

Bault N, Yaakub SN, Fouragnan E. 2024. Early-phase neuroplasticity induced by offline transcranial ultrasound stimulation in primates. *Current Opinion in Behavioral Sciences* **56**:101370. doi:10.1016/j.cobeha.2024.101370

Behrens TEJ, Woolrich MW, Walton ME, Rushworth MFS. 2007. Learning the value of information in an uncertain world. *Nature Neuroscience* **10**:1214–1221. doi:10.1038/nn1954

Billeke P, Ossandon T, Perrone-Bertolotti M, Kahane P, Bastin J, Jerbi K, Lachaux J-P, Fuentealba P. 2020. Human anterior insula encodes performance feedback and relays prediction error to the medial prefrontal cortex. *Cerebral Cortex* **30**:4011–4025. doi:10.1093/cercor/bhaa017

Breland K, Breland M. 1961. The misbehavior of organisms. *American Psychologist* **16**:681–684. doi:10.1037/h0040090

Brown PL, Jenkins HM. 1968. Autoshaping of pigeon's key-peck. *Journal of the Experimental Analysis of Behavior* **11**:1–8. doi:10.1901/jeab.1968.11-1

Cazé RD, van der Meer MAA. 2013. Adaptive properties of differential learning rates for positive and negative outcomes. *Biol Cybern* **107**:711–719. doi:10.1007/s00422-013-0571-5

992 Cecchi R, Collomb-Clerc A, Rachidi I, Minotti L, Kahane P, Pessiglione M, Bastin J. 2024.
993   Direct stimulation of anterior insula and ventromedial prefrontal cortex disrupts
994   economic choices. *Nat Commun* **15**:7508. doi:10.1038/s41467-024-51822-8

995 Daw ND, Niv Y, Dayan P. 2005. Uncertainty-based competition between prefrontal and
996   dorsolateral striatal systems for behavioral control. *Nature Neuroscience* **8**:1704–
997   1711. doi:10.1038/nn1560

998 Dayan P, Niv Y, Seymour B, Daw N. 2006. The misbehavior of value and the discipline of the
999   will. *Neural Networks* **19**:1153–1160. doi:10.1016/j.neunet.2006.03.002

1000 de Boer L, Axelsson J, Chowdhury R, Riklund K, Dolan RJ, Nyberg L, Bäckman L, Guitart-
1001   Masip M. 2019. Dorsal striatal dopamine D1 receptor availability predicts an
1002   instrumental bias in action learning. *Proceedings of the National Academy of
1003   Sciences* **116**:261–270. doi:10.1073/pnas.1816704116

1004 Ebert JP, Wegner DM. 2011. Mistaking randomness for free will. *Consciousness and
1005   Cognition* **20**:965–971. doi:10.1016/j.concog.2010.12.012

1006 Evrard HC. 2019. The organization of the primate insular cortex. *Front Neuroanat* **13**.
1007   doi:10.3389/fnana.2019.00043

1008 Folloni D, Fouragnan E, Wittmann MK, Roumazeilles L, Tankelevitch L, Verhagen L, Attali D,
1009   Aubry J-F, Sallet J, Rushworth MFS. 2021. Ultrasound modulation of macaque
1010   prefrontal cortex selectively alters credit assignment–related activity and behavior.
1011   *Science Advances* **7**:eabg7700. doi:10.1126/sciadv.abg7700

1012 Fouragnan E, Retzler C, Mullinger K, Philiastides MG. 2015. Two spatiotemporally distinct
1013   value systems shape reward-based learning in the human brain. *Nat Commun
1014   ***6**:8107. doi:10.1038/ncomms9107

1015 Fouragnan EF, Chau BKH, Folloni D, Kolling N, Verhagen L, Klein-Flügge M, Tankelevitch L,
1016   Papageorgiou GK, Aubry J-F, Sallet J, Rushworth MFS. 2019. The macaque anterior
1017   cingulate cortex translates counterfactual choice value into actual behavioral change.
1018   *Nat Neurosci* **22**:797–808. doi:10.1038/s41593-019-0375-6

1019 Guitart-Masip M, Duzel E, Dolan R, Dayan P. 2014. Action versus valence in decision
1020   making. *Trends in Cognitive Sciences* **18**:194–202. doi:10.1016/j.tics.2014.01.003

1021 Guitart-Masip M, Huys QJM, Fuentemilla L, Dayan P, Duzel E, Dolan RJ. 2012. Go and no-
1022   go learning in reward and punishment: Interactions between affect and effect.
1023   *NeuroImage* **62**:154–166. doi:10.1016/j.neuroimage.2012.04.024

1024 Hayden BY, Pearson JM, Platt ML. 2011. Neuronal basis of sequential foraging decisions in
1025   a patchy environment. *Nature Neuroscience* **14**:933–939. doi:10.1038/nn.2856

1026 Horing B, Büchel C. 2022. The human insula processes both modality-independent and pain-
1027   selective learning signals. *PLOS Biology* **20**:e3001540.
1028   doi:10.1371/journal.pbio.3001540

1029 Hsu M, Bhatt M, Adolphs R, Tranel D, Camerer CF. 2005. Neural systems responding to
1030   degrees of uncertainty in human decision-making. *Science* **310**:1680–1683.
1031   doi:10.1126/science.1115327

1032 Huys QJM, Gölzer M, Friedel E, Heinz A, Cools R, Dayan P, Dolan RJ. 2016. The specificity
1033   of Pavlovian regulation is associated with recovery from depression. *Psychological
1034   Medicine* **46**:1027–1035. doi:10.1017/S0033291715002597

1035 Jabbi M, Bastiaansen J, Keysers C. 2008. A common anterior insula representation of
1036   disgust observation, experience and imagination shows divergent functional
1037   connectivity pathways. *PLOS ONE* **3**:e2939. doi:10.1371/journal.pone.0002939

1038 Kahneman D. 2011. Thinking, fast and slow. New York, NY: Farrar, Strauss, and Giroux.

1039 Karlsson MP, Tervo DGR, Karpova AY. 2012. Network resets in medial prefrontal cortex
1040   mark the onset of behavioral uncertainty. *Science* **338**:135–139.
1041   doi:10.1126/science.1226518

1042 Keramati M, Dezfouli A, Piray P. 2011. Speed/accuracy trade-off between the habitual and
1043   the goal-directed processes. *PLoS Computational Biology* **7**:e1002055.
1044   doi:10.1371/journal.pcbi.1002055

1045 Kolling N, Behrens TEJ, Mars RB, Rushworth MFS. 2012. Neural mechanisms of foraging.
1046   *Science* **336**:95–98. doi:10.1126/science.1216930

Korn CW, Sharot T, Walter H, Heekeren HR, Dolan RJ. 2014. Depression is related to an
absence of optimistically biased belief updating about future life events. *Psychol Med*
44:579–592. doi:10.1017/S0033291713001074

Kurth F, Zilles K, Fox PT, Laird AR, Eickhoff SB. 2010. A link between the systems:
functional differentiation and integration within the human insula revealed by meta-
analysis. *Brain Struct Funct* **214**:519–534. doi:10.1007/s00429-010-0255-z

Lamm C, Singer T. 2010. The role of anterior insular cortex in social emotions. *Brain Struct*
Funct **214**:579–591. doi:10.1007/s00429-010-0251-3

Legon W, Adams S, Bansal P, Patel PD, Hobbs L, Ai L, Mueller JK, Meekins G, Gillick BT.
2020. A retrospective qualitative report of symptoms and safety from transcranial
focused ultrasound for neuromodulation in humans. *Sci Rep* **10**:5573.
doi:10.1038/s41598-020-62265-8

McGuire JT, Kable JW. 2015. Medial prefrontal cortical activity reflects dynamic re-evaluation
during voluntary persistence. *Nat Neurosci* **18**:760–766. doi:10.1038/nn.3994

Meder D, Kolling N, Verhagen L, Wittmann MK, Scholl J, Madsen KH, Hulme OJ, Behrens
TEJ, Rushworth MFS. 2017. Simultaneous representation of a spectrum of
dynamically changing value estimates during decision making. *Nature*
Communications **8**:1942. doi:10.1038/s41467-017-02169-w

Medford N, Critchley HD. 2010. Conjoint activity of anterior insular and anterior cingulate
cortex: awareness and response. *Brain Struct Funct* **214**:535–549.
doi:10.1007/s00429-010-0265-x

Murphy K, Fouragnan E. 2024. The future of transcranial ultrasound as a precision brain
interface. *PLOS Biology* **22**:e3002884. doi:10.1371/journal.pbio.3002884

Nassar MR, Frank MJ. 2016. Taming the beast: Extracting generalizable knowledge from
computational models of cognition. *Current Opinion in Behavioral Sciences* **11**:49–54.
doi:10.1016/j.cobeha.2016.04.003

O'Doherty JP, Cockburn J, Pauli WM. 2017. Learning, reward, and decision making. *Annual*
Review of Psychology **68**:73–100. doi:10.1146/annurev-psych-010416-044216

O'Reilly JX, Schüffelgen U, Cuell SF, Behrens TEJ, Mars RB, Rushworth MFS. 2013.
Dissociable effects of surprise and model update in parietal and anterior cingulate
cortex. *Proceedings of the National Academy of Sciences* **110**:E3660–E3669.
doi:10.1073/pnas.1305373110

Palminteri S, Justo D, Jauffret C, Pavlicek B, Dauta A, Delmaire C, Czernecki V, Karachi C,
Capelle L, Durr A, Pessiglione M. 2012. Critical roles for anterior insula and dorsal
striatum in punishment-based avoidance learning. *Neuron* **76**:998–1009.
doi:10.1016/j.neuron.2012.10.017

Palminteri S, Wyart V, Koechlin E. 2017. The importance of falsification in computational
cognitive modeling. *Trends in Cognitive Sciences* **21**:425–433.
doi:10.1016/j.tics.2017.03.011

Piray P, Daw ND. 2021. A model for learning based on the joint estimation of stochasticity
and volatility. *Nature Communications* **12**:6587. doi:10.1038/s41467-021-26731-9

Piray P, Dezfouli A, Heskes T, Frank MJ, Daw ND. 2019a. Hierarchical Bayesian inference
for concurrent model fitting and comparison for group studies. *PLOS Computational*
Biology **15**:e1007043. doi:10.1371/journal.pcbi.1007043

Piray P, Ly V, Roelofs K, Cools R, Toni I. 2019b. Emotionally aversive cues suppress neural
systems underlying optimal learning in socially anxious individuals. *Journal of*
Neuroscience **39**:1445–1456. doi:10.1523/JNEUROSCI.1394-18.2018

Preuschoff K, Quartz SR, Bossaerts P. 2008. Human insula activation reflects risk prediction
errors as well as risk. *Journal of Neuroscience* **28**:2745–2752.
doi:10.1523/JNEUROSCI.4286-07.2008

Ritov I, Baron J. 1995. Outcome knowledge, regret, and omission bias. *Organizational*
Behavior and Human Decision Processes **64**:119–127. doi:10.1006/obhd.1995.1094

Ritov I, Baron J. 1990. Reluctance to vaccinate: Omission bias and ambiguity. *Journal of*
Behavioral Decision Making **3**:263–277. doi:10.1002/bdm.3960030404

Scholz V, Algermissen J, Kandroodi MR, Gillan C, Ouden H den. 2025. Adaptive
suppression of motivational biases increases with mood/anxiety traits.
doi:10.31234/osf.io/v3ndf_v1

Scholz V, Hook RW, Kandroodi MR, Algermissen J, Ioannidis K, Christmas D, Valle S,
Robbins TW, Grant JE, Chamberlain SR, den Ouden HEM. 2022. Cortical dopamine
reduces the impact of motivational biases governing automated behaviour.
Neuropsychopharmacology. doi:10.1038/s41386-022-01291-8

Seeley WW. 2019. The salience network: a neural system for perceiving and responding to
homeostatic demands. *J Neurosci* **39**:9878–9882. doi:10.1523/JNEUROSCI.1138-
17.2019

Shahar N, Hauser TU, Moran R, Moutoussis M, Bullmore E, Dolan RJ, Goodyer I, Fonagy P,
Jones P, Moutoussis M, Hauser T, Neufeld S, Romero-Garcia R, Clair MS, Vértes P,
Whitaker K, Inkster B, Prabhu G, Ooi C, Toseeb U, Widmer B, Bhatti J, Villis L,
Alrumaithi A, Birt S, Bowler A, Cleridou K, Dadabhoy H, Davies E, Firkins A, Granville
S, Harding E, Hopkins A, Isaacs D, King J, Kokorikou D, Maurice C, McIntosh C,
Memarzia J, Mills H, O'Donnell C, Pantaleone S, Scott J, Kiddle B, Polek E, Fearon
P, Suckling J, van Harmelen A-L, Kievit R, Chamberlain S, Bullmore ET, Dolan RJ,
NSPN consortium, Principal investigators, NSPN (funded) staff, Affiliated scientists.
2021. Assigning the right credit to the wrong action: compulsivity in the general
population is associated with augmented outcome-irrelevant value-based learning.
Translational Psychiatry **11**:564. doi:10.1038/s41398-021-01642-x

Singer T, Seymour B, O'Doherty J, Kaube H, Dolan RJ, Frith CD. 2004. Empathy for pain
involves the affective but not sensory components of pain. *Science* **303**:1157–1162.
doi:10.1126/science.1093535

Spranca M, Minsk E, Baron J. 1991. Omission and commission in judgment and choice.
Journal of Experimental Social Psychology **27**:76–105. doi:10.1016/0022-
1031(91)90011-T

Steingroever H, Wetzels R, Wagenmakers E-J. 2014. Absolute performance of
reinforcement-learning models for the Iowa Gambling Task. *Decision* **1**:161–183.
doi:10.1037/dec0000005

Stephan KE, Penny WD, Daunizeau J, Moran RJ, Friston KJ. 2009. Bayesian model
selection for group studies. *NeuroImage* **46**:1004–1017.
doi:10.1016/j.neuroimage.2009.03.025

Swart JC, Froböse MI, Cook JL, Geurts DE, Frank MJ, Cools R, den Ouden HE. 2017.
Catecholaminergic challenge uncovers distinct Pavlovian and instrumental
mechanisms of motivated (in)action. *eLife* **6**:e22169. doi:10.7554/eLife.22169

Taylor KS, Seminowicz DA, Davis KD. 2008. Two systems of resting state connectivity
between the insula and cingulate cortex. *Hum Brain Mapp* **30**:2731–2745.
doi:10.1002/hbm.20705

Valton V, Wise T, Robinson OJ. 2020. Recommendations for Bayesian hierarchical model
specifications for case-control studies in mental health.
doi:10.48550/arXiv.2011.01725

van Geen C, Chen Y, Kazinka R, Vaidya AR, Kable JW, McGuire JT. 2025. Lesions to
different regions of the frontal cortex have dissociable effects on voluntary
persistence in humans. *J Neurosci* **45**. doi:10.1523/JNEUROSCI.0068-24.2024

van Nuland AJ, Helmich RC, Dirkx MF, Zach H, Toni I, Cools R, den Ouden HEM. 2020.
Effects of dopamine on reinforcement learning in Parkinson's disease depend on
motor phenotype. *Brain* **143**:3422–3434. doi:10.1093/brain/awaa335

Vilares I, Howard JD, Fernandes HL, Gottfried JA, Kording KP. 2012. Differential
representations of prior and likelihood uncertainty in the human brain. *Current Biology*
22:1641–1648. doi:10.1016/j.cub.2012.07.010

Wegner DM. 2002. The illusion of free will. Cambridge, MA: MIT Press.

Williams DR, Williams H. 1969. Auto-maintenance in the pigeon: Sustained pecking despite
contingent non-reinforcement. *Journal of the Experimental Analysis of Behavior*
12:511–520. doi:10.1901/jeab.1969.12-511

Wilson RC, Collins AG. 2019. Ten simple rules for the computational modeling of behavioral
data. *eLife* **8**:1–35. doi:10.7554/eLife.49547
Wittmann MK, Fouragnan E, Folloni D, Klein-Flügge MC, Chau BKH, Khamassi M,
Rushworth MFS. 2020. Global reward state affects learning and activity in raphe
nucleus and anterior insula in monkeys. *Nature Communications* **11**:3771.
doi:10.1038/s41467-020-17343-w
Yaakub SN, Bault N, Lojkiewiez M, Bellec E, Roberts J, Philip N, Rushworth MF, Fouragnan
EF. 2024. Non-invasive ultrasound deep neuromodulation of the human nucleus
accumbens increases win-stay behaviour. doi:10.1101/2024.07.25.605068
Yaakub SN, White TA, Kerfoot E, Verhagen L, Hammers A, Fouragnan EF. 2023a. Pseudo-
CTs from T1-weighted MRI for planning of low-intensity transcranial focused
ultrasound neuromodulation: An open-source tool. *Brain Stimulation: Basic,*
Translational, and Clinical Research in Neuromodulation **0**.
doi:10.1016/j.brs.2023.01.838
Yaakub SN, White TA, Roberts J, Martin E, Verhagen L, Stagg CJ, Hall S, Fouragnan EF.
2023b. Transcranial focused ultrasound-mediated neurochemical and functional
connectivity changes in deep cortical regions in humans. *Nat Commun* **14**:5318.
doi:10.1038/s41467-023-40998-0
Yu AJ, Dayan P. 2005. Uncertainty, neuromodulation, and attention. *Neuron* **46**:681–692.
doi:10.1016/j.neuron.2005.04.026
Zeelenberg M, Pligt J van der, de Vries NK. 2000. Attributions of responsibility and affective
reactions to decision outcomes. *Acta Psychologica* **104**:303–315. doi:10.1016/S0001-
6918(00)00034-2

1179

1180