Multi-level broad-yet-sparse input organization of LC-NE neurons revealed by multiplexed whole-brain EM reconstruction

Fu-Ning Li[1,6], Jia-Zheng Liu[2,6], Can Shi[1,4,6], Jing-Bin Yuan[2,3,6], Ya-Nan Lv[2,3,6], Jing Liu[2,6], Li-Na Zhang[2,6], Lin-Lin Li[2,6], Li-Jun Shen[2,6], Xi Chen[2,6], Hao Zhai[2,3], Yan-Chao Zhang[2,3], Rong-Kun Tao[1], Han-Yang Hu[1], Fang-Xu Zhou[2], Tong Xin[2], Bo-Hao Chen[2,3], Hao-Ran Chen[2,3], Sheng Chang[2], Hong-Tu Ma[2], Hai-Yang Yan[2,3], Jia-Liang Zu[1,3], Jin-Yue Guo[2,3], Zheng-Huan Fang[1], Xiao-Hui Dong[2], Li-Mei Lin[2], Xing-Hui Zhao[2], Chen-Ying Qian[1], Yan-Song Wang[1,3], Shi-Rong Jin[1], Chang-Mei Zhang[1], Pei-Shan Xiang[1,4], Yong-Xin Yang[1,4], Yu Qian[1,3], Yu-Chen Gong[1,3], Xu-Fei Du[1], Qi-Wei Xie[2], Hua Han[2,3,*], Jiu-Lin Du[1,3,4,5,*]

[1] Institute of Neuroscience, State Key Laboratory of Brain Cognition and Brain-Inspired Intelligence Technology, Center for Excellence in Brain Science and Intelligence Technology, Chinese Academy of Sciences, 320 Yue-Yang Road, Shanghai 200031, China

[2] State Key Laboratory of Brain Cognition and Brain-Inspired Intelligence Technology, Institute of Automation, Chinese Academy of Sciences, Beijing 100190, China

[3] School of Future Technology, School of Artificial Intelligence, University of Chinese Academy of Sciences, Beijing 101408, China

[4] University of Chinese Academy of Sciences, 19A Yu-Quan Road, Beijing 100049, China

[5] School of Life Science and Technology, ShanghaiTech University, 319 Yue-Yang Road, Shanghai 200031, China

[6] These authors contributed equally to this work.

[*] These authors jointly supervised this work: Hua Han (hua.han@ia.ac.cn),
Jiu-Lin Du (forestdu@ion.ac.cn)

Diverse neuromodulatory systems confer functional flexibility upon the

hardwired sensorimotor pathways of the brain[1-6]. Among these, the

evolutionarily conserved locus coeruleus (LC)-norepinephrinergic (NE) system

integrates and broadcasts information globally[7-9]. While previous studies have

primarily focused on its axonal outputs and effects on neural processing, the

organizational logic of its synaptic inputs, enabling it to sense global brain states

and in turn shape its activity dynamics for appropriate neuromodulation[10,11],

remains poorly explored. To address this, we mapped the synaptic input

architecture of individual LC-NE neurons by establishing Fish-X, a whole-brain

microscale reconstruction of larval zebrafish with neuron-type annotations. This

dataset encompasses the retina, brain and anterior spinal cord,

capturing >240,000 cells and >25 million synapses. Monoaminergic (including

NE, dopaminergic, and serotonergic), hypocretinergic, and glycinergic neurons

were resolved by multiplexed subcellular APEX2 labeling, while glutamatergic

and GABAergic identities were inferred via morphology comparison with a

zebrafish mesoscopic atlas[12,13]. Compared with other neuronal populations,

LC-NE neurons display distinct perisomatic features and high dendritic

indegrees. Reconstruction of near-complete dendritic inputs to individual LC-NE

neurons reveals their broad yet sparse synaptic convergence across the brain.

These synaptic inputs are not randomly distributed but instead organized

according to sensory/motor modality, excitatory/inhibitory identity, and synaptic

strength. Individual LC-NE neurons share common inputs, a feature conserved

within and across monoaminergic systems, suggesting a co-innervation

mechanism for coordinated neuromodulation. Thus, our study uncovers

multi-level principles governing the spatial organization of LC-NE neurons'

broad-yet-sparse inputs and provides a pivotal resource for deciphering the

microscale architecture of neuromodulatory systems.

The brain is organized across scales, from precisely wired synapses, through

morphologically and functionally heterogeneous neurons, to specific neural circuits

and brain-wide networks[14-16]. A simplified framework for deconstructing its

organization is the concept of a bi-pathway architecture, in which inter-connected

sensorimotor and neuromodulatory pathways function complementarily[1-6].

Neuromodulatory systems orchestrate sensorimotor circuits, shaping their state,

dynamics and output without directly driving behavioral outputs. While previous

studies mainly focused on their axonal outputs and influence on sensorimotor

processing[7,10,17-21], a critical question remains: how do these systems, through their

synaptic inputs, perceive external environmental changes as well as internal brain

states and in turn generate dynamic activities enabling spatiotemporally specific
modulation[1-3,5,6,10,11]? To address this issue, it is essential to elucidate the spatial
organization of synaptic inputs onto neuromodulatory neurons, including cellular
origin, functional modality, and excitatory (E)/inhibitory (I) identity.

Microscale connectomics based on electron microscopy (EM) is a powerful
approach for synaptic input mapping[22-27], yielding wiring diagrams for the central
nervous system (CNS) of small invertebrates (including *Caenorhabditis elegans* and
Drosophila melanogaster)[28-33] and cubic millimeter-scale reconstructions of the
mouse and human cortices[34-38]. The larval zebrafish (*Danio rerio*) with its manageable
brain scale has emerged as a tractable vertebrate model for whole-brain
reconstruction[39]. Previous studies have provided foundational datasets, allowing the
synaptic-resolution mapping of sensorimotor circuits[40-45]. The functional complexity
of neural networks arises from both the physical wiring and molecular diversity of
their constituent neurons. However, the absence of molecular annotations, particularly
for neuromodulatory neurons, limits functional insights from most connectomes.
While traditional EM connectomics identifies neuron types using structural landmarks,
mapping cell types that lack distinct anatomical signatures requires correlative
light-electron microscopy (CLEM)[45], immunogold labeling[46], or genetically encoded
tags for EM (GETEM)[47]. Among these, GETEM offers a highly feasible approach, as
it leverages genetically encoded tags to embed molecular identity directly within the
ultrastructural map, sidestepping computational registration challenges and ensuring
sample preservation.

We therefore built Fish-X, a whole-brain, synaptic-resolution serial-section EM
(ssEM) dataset with multiplexed molecular annotations from a 6-day post-fertilization
(dpf) larval zebrafish, spanning the CNS from the retina to the anterior spinal cord
(aSC). Within this dataset, diverse neuromodulatory (norepinephrinergic, NE;
dopaminergic, DA; serotonergic, 5HT; hypocretinergic, Hcrt) and inhibitory
glycinergic (Gly) neurons were precisely identified using enhanced ascorbate
peroxidase 2 (APEX2)[48], which was targeted to different subcellular compartments of
distinct neuron types using signal peptides and neuron type-specific promoters.
Furthermore, we inferred glutamatergic (Glu) and GABAergic (GABA) types of
neurons through the zebrafish mesoscopic atlas[12]-assisted morphology comparison. A
whole-brain algorithmic reconstruction was then performed, identifying 176,810 cells
and 25.25 million synapses, thereby providing a foundation for systematic analysis of
the organization of synaptic inputs to these neuromodulatory neuronal populations.

We focused on NE neurons in the evolutionarily conserved locus coeruleus (LC),
a key nuclei for brain-wide regulation[49-53]. Viral tracing revealed cellular-level input
convergence onto LC-NE neurons[7,8], yet the spatial organization of synaptic-level
inputs remains unknown. To address this, we performed somatodendritic

reconstructions and synaptic mapping of APEX2-labeled LC-NE neurons. These
reconstructions enabled us to define their characteristic perisomatic ultrastructural
features, establishing structural criteria for the precise, label-free identification of
LC-NE neurons[54,55]. Brain-wide, synaptic-resolution input reconstruction revealed
that individual LC-NE neurons integrate widely distributed inputs. These inputs were
spatially organized by functional modality and E/I type, with sensory-related I inputs
biasing to distal dendrites and motor-related E inputs to proximal ones. While most
inputs are unique to individual neurons, a subset constitutes intra-system common
inputs that innervate multiple LC-NE neurons. Moreover, this co-innervation principle
extends across different monoaminergic systems, suggesting an inter-system
organization for orchestrating NE, DA, and 5HT release, which may collectively
modulate neural network activities. Take together, leveraging the Fish-X resource, a
whole-brain, synaptic-resolution dataset with molecularly defined neuron-type
annotations, we have quantified the ultrastructural features of diverse
neuromodulatory neurons and delineated the organizational principles governing
synaptic inputs to the LC-NE system. Integrated with the zebrafish mesoscopic
atlases[12,13], the Fish-X can serve as a valuable resource for deciphering how, at the
level of synaptic architecture, neuromodulatory systems interact with sensorimotor
pathways to shape neurocomputing in the larval zebrafish brain.
125
Multiplexed APEX2-labeled whole-brain ssEM of larval zebrafish
We achieved simultaneous EM labeling of multiple molecularly defined neuron types
within a single brain preparation using APEX2, an enzyme that catalyzes the
hydrogen peroxide (H_2O_2)-dependent localized polymerization of diaminobenzidine
(DAB)[48]. APEX2 fused with signal peptides and fluorescent proteins was expressed
via cell type-specific promoters for labeling distinct subcellular compartments in
different types of neurons. After the DAB reaction, the localized APEX2 generated
EM-visible staining patterns, thereby enabling the identification of different neuron
types (Fig. 1a; see Methods). We achieved EM labeling of multiple subcellular
compartments including the nucleus, cytosol, plasma membrane, mitochondria, Golgi
apparatus, and presynaptic sites (Fig. 1b, six leftmost images, and Extended Data Fig.
1a). In addition, by fusing APEX2 with self-assembling proteins, Polymer King-size
Unit (PKU) tags[56], we achieved fibrous and spherical staining patterns localized in the
nucleus or cytosol (Fig. 1b，three rightmost images, and Extended Data Fig. 1a). Thus
we generated a versatile genetic toolbox to target a wide range of neuromodulatory
types, including NE, DA, 5HT, and Hcrt neurons, as well as Glu, GABA, and Gly
neurons across the CNS (Supplementary Table 1,2).
To examine the inputs and their spatial organization of multiple neuromodulatory
neuronal populations within the same animal, we made a quadruple transgenic

zebrafish   line,
Gt(vmat2:GAL4FF);Tg(dbh:GAL4FF;UAS:cytosol-APEX2-TagRFP);Tg(hcrt:APEX2
-TagRFP-NLS);Tg(glyt2:MITO-APEX2-TagRFP), in which we were able to
simultaneously annotate cytosol-labeled NE neurons in the LC and medulla oblongata
(MO), cytosol-labeled DA neurons in the olfactory bulb (OB), posterior tuberculum
(PT) and caudal hypothalamus (Hc), cytosol-labeled 5HT neurons in the pretectum
(Pr) and raphe nucleus (RN), nucleus-labeled Hcrt neurons in the hypothalamus, and
mitochondria-labeled Gly neurons in the hindbrain and aSC (Fig. 1c and Extended
Data Fig. 1b). The specificity of labeling was confirmed through
immunohistochemical staining or crossing with neuron type-specific reporter fish
lines, albeit with variations in labeling efficiency across neuronal populations
(Extended Data Fig. 1c).

We acquired a whole-brain ssEM dataset (named as Fish-X) from a 6-dpf
multiplexed APEX2-labeled larva using the automated tape-collecting ultramicrotome
– scanning electron microscopy (ATUM-SEM) approach[40]. With a voxel resolution of
$4\times4\times33$ nm^3, Fish-X extended from the olfactory epithelium to the aSC (the vertebra
1) and included the right retina and optic nerve (Fig. 1d, Extended Data Fig. 2; see
Methods). The imaged volume, generated from 22,887 consecutive coronal-view
ultrathin sections, encompassed $\sim3.1160\times10^7$ μm^3 for the entire brain and aSC,
$\sim5.5620\times10^4$ μm^3 for the right optic nerve, and $\sim6.0877\times10^6$ μm^3 for the right retina.
Utilizing an automated three-dimensional (3D) convolutional neural network
(CNN)-based reconstruction of the brain and aSC (Fig. 1e; see Methods), we
performed multidimensional structural annotations (Supplementary Video 1)
encompassing nuclei (Fig. 1f and Extended Data Fig. 3), neuronal morphologies (Fig.
1g), brain regions (Fig. 1h, Supplementary Table 3, and Supplementary Video 2),
neural tracts and vessels (Fig. 1i).

Based on the results of nucleus segmentation, a total of 176,810 cells, including
66 NE, 105 DA, 126 5HT, 12 Hcrt, and 2,689 Gly neurons, were identified in the
brain and aSC, and 70,910 cells in the right retina (see Fig. 1d). We quantified nuclear
morphology by integrating deep-learned and traditional features (see Methods) and
discovered that morphological diversity formed a continuous spectrum, though
vasculature-associated cell types, such as endothelial cells (ECs) and pericytes (PCs),
clustered distinctly within the morphological space (Extended Data Fig. 3a,b). Spatial
mapping revealed heterogeneity in cell density across brain regions (Extended Data
Fig. 3c, Supplementary Table 4, and Supplementary Video 2).

In a parallel analysis of synaptic structures (Extended Data Fig. 4a; see Methods),
automatic segmentation identified 25.25 million postsynaptic densities (PSDs), 15.77
million synaptic vesicle clouds (SVCs), and 13.73 million mitochondria (Mito) in the
brain and aSC. While 93.5% (56.6% with SVC and Mito + 36.9% with SVC only) of

PSDs colocalized with SVCs, only 59.2% (56.6% with SVC and Mito + 2.6% with Mito only) colocalized with presynaptic mitochondria (Extended Data Fig. 4b), suggesting heterogeneity in synaptic energy supply. Meanwhile, 84.3% of SVCs and 61.3% of mitochondria co-localized with PSDs (Extended Data Fig. 4c,d). All three structures exhibited heterogeneous yet highly concerted distribution patterns across regions (Extended Data Fig. 4e-j, Supplementary Tables 5,6,7, and Supplementary Video 2).

Across regions, the density of typical synapses with colocalized PSDs and SVCs was inversely correlated with cell density, which was consistently observed across forebrain, midbrain, and hindbrain, highlighting the spatial segregation of synapses and somata (Fig. 1j). When regions were grouped into broader anatomical (i.e., forebrain, midbrain, and hindbrain) or functional (i.e., sensory-related, and motor-related) categories, no significant difference in density was found (Fig. 1k; Kruskal-Wallis test for the anatomical category, Mann Whitney test for the functional category). Similarly, the volumes of all three components followed comparable distributions across regions, with the exception of a slight difference in SVC volume between the forebrain and hindbrain (Fig. 1l; Kruskal-Wallis test with Dunn's multiple comparison test for the anatomical category, Mann Whitney test for the functional category). Regions from the same anatomical categories generally exhibited similar lognormal distributions of synaptic structure volumes (Extended Data Fig. 4k-n).

Collectively, we developed a whole-brain, synaptic-resolution dataset for larval zebrafish, annotated with neuromodulatory and glycinergic neurons via multiplexed subcellular APEX2 labeling.

Unique perisomatic features and high dendritic indegrees of LC-NE neurons

Different neuromodulatory systems vary in the degree of axonal innervation patterns[7,10,17-21]. As evidenced by light microscopy (LM)-based single-cell morphological tracing in 6-dpf larval zebrafish, we found that LC-NE neurons exhibited more widespread axonal projections throughout the brain in comparison with other types of neuromodulatory neurons, including MO-NE, OB-DA, PT-DA, Hc-DA, Pr-5HT, RN-5HT, and Hcrt neurons (Extended Data Fig. 5). To characterize the ultrastructural properties of these neurons' synaptic inputs, we reconstructed the somatodendritic morphology and synaptic input sites of these types of neurons at the right hemisphere, along with Gly neurons, and proofread and quantified their perisomatic ultrastructures (Fig. 2a-g, Extended Data Figs. 6,7, and Supplementary Table 8).

LC-NE neurons exhibited not only larger nuclear and somatic volumes but also featured extensive endoplasmic reticulum (ER) networks, numerous mitochondria,

and a specialized cilium (Extended Data Fig. 7a and Supplementary Table 8).
Compared with a 10-fold pool of randomly selected adjacent cells, LC-NE neurons
had a lower mean nucleus/soma volume ratio (LC-NE neurons *vs* adjacent cells:
0.4123 ± 0.0808 *vs* 0.6530 ± 0.0113, $P < 0.05$, Mann Whitney test) and a higher
somatic mitochondrial density (LC-NE neurons *vs* adjacent cells: 0.5449 ± 0.0496 *vs*
0.0843 ± 0.0088 per 1 μm^3, $P < 0.001$, Mann Whitney test) (Extended Data Fig. 7b,
left). Clustering analyses integrating both nuclear and somatic characteristics
validated the above observation (Extended Data Fig. 7b, right). Based on these
distinct perisomatic features, we identified an additional unlabeled NE neuron in the
right LC region (Extended Data Fig. 7b, blue arrows), whose morphology closely
matched those of APEX2-labeled LC-NE neurons (Extended Data Fig. 7c). We
extended these analyses to other APEX2-labeled neuronal populations and found that,
with the exception of PT-DA neurons (Extended Data Fig. 7f), other types of
neuromodulatory neurons lacked perisomatic features that clearly distinguished them
from adjacent cells (Extended Data Fig. 7d-k and Supplementary Table 8). This
finding further highlights the value of APEX2 labeling for precise identification of
neuron types. In companion with other types of neuromodulatory neurons, LC-NE
neurons exhibited higher values in terms of nucleus/soma volume ratio (Fig. 2f;
LC-NE *vs* PT-DA, $P = 0.1255$, Mann Whitney test; LC-NE *vs* all other types except
PT-DA, $P < 0.05$, Mann Whitney test) and somatic mitochondrial density (Fig. 2g;
LC-NE *vs* all other types, $P < 0.05$, Mann Whitney test), suggesting unique
perisomatic properties of LC-NE neurons.

We then reconstructed and quantified all synaptic input sites across 27
APEX2-labeled neuromodulatory neurons (including 5 LC-NE, 3 MO-NE, 3 OB-DA,
4 PT-DA, 3 Hc-DA, 3 Pr-5HT, 4 RN-5HT, and 2 Hcrt) and 17 labeled Gly neurons at
somatodendritic regions (Fig. 2h,i and Supplementary Videos 3,4). LC-NE, PT-DA,
and RN-5HT neurons collectively displayed a higher mean synaptic input number and
density compared with other types, with the LC-NE neurons exhibiting the highest
synaptic input number (Fig. 2j,k; Input density among LC-NE, PT-DA, and RN-5HT,
$P = 0.9533$, Kruskal-Wallis test; Input density of LC-NE/PT-DA/RN-5HT *vs* all other
types, $P < 0.0001$, unpaired *t* test; Input number, $P < 0.0001$, Kruskal-Wallis test).
Moreover, the synaptic input number showed a positive correlation with the total
projection length of respective neuron types (Fig. 2l). Notably, these neuromodulatory
neurons had sparse synaptic inputs on their somata (Supplementary Video 4), unlike
the well-studied Mauthner cell, which received substantial somatic innervation
(Extended Data Fig. 8 and Supplementary Video 5).

Thus these results show that LC-NE neurons constitute a structurally distinct
system with unique perisomatic properties and high dendritic indegrees. Furthermore,
the Fish-X dataset provides a structural ground truth for multiple neuromodulatory

neuron types, serving as an essential reference for their identification in other EM
datasets lacking molecular annotations[54,55].

Spatial organization of broad-yet-sparse inputs to individual LC-NE neurons
To examine the spatial organization of synaptic inputs on LC-NE neurons, we focused
on LC-NE neuron #3 due to its possession of the longest dendrites (~1.812 mm) and
the highest dendritic indegree (560 synapses) among the five APEX2-labeled LC-NE
neurons in the right hemisphere (Fig. 3a,b and Extended Data Fig. 9).

Presynaptic neurons (PNs) for this LC neuron were traced from all input sites
and categorized by putative functional modality (i.e., sensory-related *vs* motor-related)
based on PN soma location. We successfully traced 418 inputs to the somata of 374
PNs, averaging 1.12 inputs per PN. Among these, 182 synapses originated from 160
PNs in sensory-related regions (including sensory-relay regions), and 236 synapses
from 214 PNs in motor-related regions (Fig. 3c). Spanning the forebrain, midbrain,
and hindbrain, these PNs were found in 32 out of 52 total brain regions (Fig. 3d and
Supplementary Video 6), indicative of a broad-yet-sparse connectivity pattern. The
mean input number per PN did not differ significantly between sensory- and
motor-related regions (Fig. 3d; Sensory *vs* Motor, *P* = 0.5192, Mann Whitney test),
suggesting that both modalities conform to this connectivity pattern. Through a
combinatorial analysis of dendritic position (relative to the LC-NE neuron's soma)
against PN functional modality, we found a modality-dependent spatial organization
on the LC-NE neuron's dendrites (Fig. 3e,f). Specifically, for LC-NE neuron #3,
sensory-related inputs preferentially targeted distal dendritic segments, whereas
motor-related ones were biased toward proximal segments (Fig. 3e; path length to the
soma: sensory-related, 122.8 ± 3.2 μm *vs* motor-related, 103.3 ± 2.5 μm; *P* < 0.0001,
Mann Whitney test). The similar organizational profiles were observed for the other
four LC-NE neurons, though the synaptic sampling of these neurons was
unintentionally biased toward the proximal dendrites, resulting in a shorter average
path length (Fig. 3f; path length to the soma: sensory-related, 76.7 ± 3.6 μm *vs*
motor-related, 57.8 ± 2.5 μm; *P* < 0.0001, Mann Whitney test). Furthermore, both
sensory- and motor-related inputs exhibited a spatial gradient of synaptic strength,
with a comparable trend of increasing PSD and SVC volumes when approaching the
soma (Fig. 3g,h; *P* < 0.001, Mann Whitney test).

We next investigated the E/I type of these PNs. In mammals, Glu synapses are
typically asymmetrical and characterized by thick PSDs, whereas GABA synapses are
symmetrical with thin PSDs[57]. However, an analysis of additional EM datasets from
Glu and GABA neurons with mitochondria-localized APEX2 labeling showed no
significant difference in PSD thickness or volume between the two types of synapses,
with only minor exceptions in rare cases (Extended Data Fig. 10a-d), suggesting that

synapses in zebrafish may not follow the criterion observed in mammals. Then we

inferred the neuron type by comparing the morphology of the PNs with a reference

dataset of LM-based fully-reconstructed neurons from the zebrafish mesoscopic

atlas[12]. This dataset comprised 12,219 Glu, 5,371 GABA, and 2,193 Gly neurons. We

pre-registered the EM query neurons to the mesoscopic brain template via a CLEM

pipeline and extracted mesoscopically traced neurons within a 25-μm radius sphere

centered at each query neuron's centroid. Skeleton matching against these references

yielded penalty scores, with lower scores indicating higher morphological similarity

(see Methods). Manual verification of top-ranked matches enabled the classification

of all 467 EM query neurons as matched or unmatched. Among them, 389 queries

were successfully matched to morphologically similar E or I reference neurons (Fig.

3i and Extended Data Fig. 10e,f), while 78 showed morphological discrepancies from

their top references (Extended Data Fig. 10g), suggesting the need for more complete

reconstructions and broader reference coverage. Matched queries had lower penalty

scores than unmatched ones (Extended Data Fig. 10h; $P < 0.0001$, Mann Whitney

test), supporting the efficacy of morphology-based matching. Of the 389 matched

queries, 56% (219) were E/I-specific match, with top references predominantly of a

single type, and the remaining 170 were E/I-ambiguous match, showing top matches

of mixed types. For instance, query neuron #41's top references included Glu, GABA,

and Gly neurons, preventing reliable E/I typing by morphology match (Extended Data

Fig. 10i). Its identity as a Gly neuron was, however, confirmed by mitochondrial

APEX2 labeling (Extended Data Fig. 10j), again demonstrating the value of APEX2

labeling for definitive identification of neuron types.

Based on the morphology matching, we performed E/I inference for PNs of all

five APEX2-labeled LC-NE neurons (Fig. 3j,k and Extended Data Fig. 10i-l). For the

LC-NE neuron #3, we inferred the E/I identity for 188 of the 374 PNs, with Glu being

the predominant type (Fig. 3j; Sensory-related, 60 Glu, 15 GABA, 7 Gly;

Motor-related, 75 Glu, 18 GABA, 13 Gly). Spatial analysis along the left-right

dendritic axis revealed that the spatial distribution of synaptic inputs showed both E/I

type- and sensory/motor modality-dependency. In comparison with I-type inputs,

E-type inputs relatively preferred to proximal dendritic segments, in particular for

sensory-related inputs (Fig. 3l,m; $P < 0.05$ for sensory-related inputs, $P = 0.1072$ for

motor-related inputs, two-sample Kolmogorov-Smirnov test). I-type sensory-related

inputs were distributed more distally than I-type motor-related inputs ($P < 0.05$,

two-sample Kolmogorov-Smirnov test), while E-type motor-related inputs were

biased more proximally than E-type sensory-related inputs ($P < 0.001$, two-sample

Kolmogorov-Smirnov test) (Fig. 3l,m).

These results together show that LC-NE neurons receive a broad and diverse

range of synaptic inputs, organized by sensory/motor modality, E/I identity, and

synaptic strength at their dendrites (Supplementary Video 7).

Intra- and inter-system common synaptic inputs among monoaminergic systems

Among the 374 PNs of the LC-NE neuron #3, 69 of them also targeted at least one

other neuron among the five APEX2-labeled LC-NE neurons (Fig. 4a), thereby

serving as common presynaptic neurons (CPNs) within the LC-NE system

("intra-CPNs"). These intra-CPNs were widely distributed across sensory- and

motor-related regions (Fig. 4a,b), and the majority of identified intra-CPNs were E

neurons (Fig. 4c). Moreover, these intra-CPNs displayed diverse innervation patterns,

with the majority synapsing onto two LC-NE neurons (Fig. 4d). To assess whether

intra-CPNs co-innervate LC-NE neurons randomly or selectively, we performed

theoretical simulations using identical cell numbers and innervation distributions. The

simulation revealed that the LC-NE neuron #2 was co-innervated with #3 more

frequently than with other LC-NE neurons (Fig. 4e; $P < 0.05$, chi-square

goodness-of-fit test), suggesting the potential existence of LC-NE subpopulations

with preferential co-innervation. These intra-CPNs may represent as a structural

substrate for synchronous activity within LC-NE neuron populations, as reported

previously[9,58,59], and help coordinate functional outputs of the LC system.

In contrast, 305 of the 374 PNs (81.28%) reconstructed for the LC-NE neuron #3

did not target any of the other four LC-NE neurons and accounted for 330 of 418

(78.95%) synapses onto this neuron. We also observed that the two reconstructed Hcrt

neurons exclusively innervated the LC-NE neuron #4 (see Extended Data Fig. 10k).

Together, these findings indicate that individual LC-NE neurons receive both unique

and shared synaptic inputs, aligning with recent studies supporting the heterogeneous

and modular organization of the LC-NE system[8,9,60-63,].

We then extended the analysis to other neuromodulatory systems. Among the 374

PNs of the LC-NE neuron #3, we identified 26 widely distributed CPNs that

co-innervated PT-DA or RN-5HT neurons ("inter-CPNs"; Fig. 4f,g).

Morphology-based assessment of E/I type indicated that inter-CPNs connecting

LC-NE and PT-DA neurons were predominantly Glu (5/8), with the remainders being

E/I-ambiguous (3/8) (Fig. 4h). Meanwhile, inter-CPNs linking LC-NE and RN-5HT

neurons showed a balanced E/I characteristics (Glu, 5/18; GABA, 4/18; Gly, 2/18;

E/I-ambiguous, 7/18) (Fig. 4h). We thus hypothesized that LC-NE and PT-DA

neurons may receive shared feedforward driving signals, whereas LC-NE and

RN-5HT neurons may be subject to common inhibitory control. To support this

hypothesis, we performed *in vivo* calcium imaging across these monoaminergic nuclei

in the same larval zebrafish, and observed that LC-NE neurons exhibited spontaneous

activities synchronized with PT-DA neurons, but not with RN-5HT neurons (Fig. 4i,j).

These inter-CPNs could provide a structural basis for coordinated activity across

distinct monoaminergic systems, enabling more diversified regulatory functions.
Some inter-CPNs also innervated multiple PT-DA or RN-5HT neurons, serving as
intra-CPNs for these monoaminergic systems (Fig. 4k, left and right). Similar to the
LC-NE system, PNs targeting PT-DA and RN-5HT neurons were also
widely-distributed, which contrasts with the spatially restricted input profiles of
OB-DA neurons (Fig. 4k, top-middle), suggesting a conserved organizational
principle across monoaminergic systems. Notably, synaptic inputs from intra- and
inter-CPNs preferentially targeted proximal dendritic regions, whereas those from
non-CPN sources showed more dispersed innervation patterns (Fig. 4l).

Taken together, these results offer a multi-level view on the spatial organization
of broad-yet-sparse synaptic inputs onto LC-NE neurons' dendrites, characterized by:
1) broad-yet-sparse synaptic inputs; 2) proximity-dependent synaptic strength scaling;
3) modality and E/I type-based distribution; 4) dual input architecture (private *vs.*
shared); 5) Cross-system shared inputs with proximal bias (Fig. 5 and Supplementary
Video 7).

Discussion
Fish-X represents the first successful implementation of multiplexed APEX2 labeling
within an intact vertebrate brain, supported by a suite of genetic tools including
relevant stable zebrafish lines and plasmid constructs. The APEX2-based GETEM
strategy provides distinct advantages: (i) registration-free neuron identification, (ii)
multiplexed annotation via orthogonal compartmental labeling that bypasses the
spectral limits of fluorescent proteins[64,65], and (iii) reliable labeling of long-range
projecting axons in partial EM reconstructions where full neuronal morphologies are
unavailable[66]. In comparison, the traditional CLEM approach offers an alternative by
transferring neuronal identities from fluorescently labeled LM data into EM volumes
via image registration[37,41,42,44,45]. By leveraging existing transgenic lines and utilizing
an H_2O_2-free sample preparation, this strategy is generally compatible with
high-quality EM preservation. Importantly, the two strategies can be synergistically
combined—the fluorescent signals from APEX2-fused proteins precisely match their
DAB products, which serve as intrinsic fiducial landmarks that streamline and
enhance the accuracy of CLEM registration. Furthermore, by implementing the
CLEM framework beyond a single specimen[67], our mesoscopic atlas-based strategy
operates as a computational CLEM variant that matches EM neuronal morphologies
to a mesoscopic reference atlas. Collectively, these complementary strategies
constitute a versatile toolkit for comprehensive molecular annotation of zebrafish EM
connectomes.

Using this whole-brain, molecularly annotated EM dataset encompassing multiple
neuromodulatory types, we investigated the fine-scale spatial organization of synaptic

inputs to neuromodulatory systems through the lens of the LC-NE system, an
evolutionarily conserved hub for brain-wide regulation. Our analysis reveals a
structural blueprint for global integration in the LC-NE system, which receives broad,
sparse inputs from sensory-, sensory relay-, and motor-related regions across the brain.
With individual PNs contributing very few synapses—on average, fewer than
two—this "broad-yet-sparse" connectivity pattern is ideally suited for monitoring
global brain state through consensus[68]. Individual LC-NE neurons therefore detect
when widespread, coincident activity across brain regions reaches a threshold
signaling a salient event, rather than responding to a single, localized signal. Within
this architecture, dominant proximal excitatory inputs from motor-related regions
create a low-threshold channel for motor commands or feedback, ensuring rapid and
reliable NE release to facilitate swift behavioral state transitions[69-71]. In contrast, distal
sensory inputs, particularly inhibitory ones, likely exert modulatory control by setting
global neuronal excitability through dendritic calcium spike modulation or shunting
inhibition, rather than directly vetoing action potentials[72-75]. This fine-tuning enables
context-dependent NE neuromodulation. The contrast between this organizational
logic and that of cortical pyramidal neurons[11,76-81] underscores the fundamental
diversity in dendritic integration mechanisms across cell types and highlights the
necessity of building neuron models grounded in cell-type-specific connectomes.

At the population level, accumulating molecular, morphological, and functional
evidence indicates that the LC-NE system exhibits modular rather than homogeneous
organization[8,9,60-63]. Our study provides new insights into this organization from the
perspective of synaptic input architecture. We found that individual LC-NE neurons
receive both "private" (i.e., non-CPNs) and shared (i.e., intra-CPNs) inputs,
collectively forming an architectural basis for population-level activity. Private inputs
likely drive sparse, asynchronous NE release from single neurons. In contrast, shared
inputs with varying innervation numbers can recruit specific subsets or even the entire
population of LC-NE neurons. This recruitment, potentially further synchronized via
gap junctions among LC-NE neurons, enables regionally distinct and intensity-graded
NE release. An intriguing question for future investigation is whether this
private–shared input dichotomy corresponds to the well-established tonic and phasic
firing modes of LC-NE neurons, potentially encoding distinct presynaptic information
streams[82,83]. Furthermore, by extending this modularity concept beyond the LC-NE
system, we identified shared inputs (inter-CPNs) that co-innervate both LC-NE and
other monoaminergic neurons, suggesting the existence of higher-order, inter-system
modules that provide a structural basis for the coordinated neuromodulation observed
in diverse physiological and pathological brain states.

The observed co-innervation architecture between distinct neuromodulatory
systems not only demonstrates the value of our multiplexed molecularly annotated

EM resource, but also underscores the need for more complete reconstructions to fully resolve the inputome of each neuromodulatory system and the organizational principles governing their interactions. Notably, the anatomical comprehensiveness of Fish-X, encompassing the retina, optic nerve, brain, and aSC, enables the systematic investigation of both neuromodulatory and sensorimotor pathways across the entire CNS. To fully realize this potential, the dataset can be computationally refined through advanced alignment and segmentation algorithms combined with online collaborative proofreading to improve reconstruction efficiency[84-88]. Furthermore, the ATUM-SEM approach has preserved all sections, enabling targeted re-imaging of both suboptimal regions and previously unmapped areas. In parallel, there is a pressing need for semi- or fully-automated reconstruction algorithms specifically designed for GETEM-labeled neurons, particularly those with challenging cytosol labeling[89]. Our dataset, with its rich APEX2-labeled resource encompassing a wide range of staining intensities, provides an ideal benchmark for developing and refining such algorithms. Beyond enhancing the current resource, this study has established a versatile APEX2-based genetic toolbox and optimized procedures—including real-time quality control, image repair, and image denoising—that enable the creation of next-generation EM datasets with expanded simultaneous neuron type annotation and higher image quality.

In summary, we have established a feasible pipeline for generating molecularly annotated whole-brain EM-reconstructions in larval zebrafish and provided a valuable resource for the community. Based on this dataset, we have revealed the principles of synaptic input architecture of the LC-NE system. Further investigation building upon this work will not only expand the ultrastructural mapping of input – output connectomes for diverse neuromodulatory neuronal populations—providing a substrate for their realistic neuronal modeling—but also deepen our understanding of how neuromodulatory neurons are synaptically integrated within the bi-pathway brain architecture.

References

1. Marder, E. Neuromodulation of neuronal circuits: back to the future. *Neuron* **76**, 1-11 (2012).2

2. Shang, C.-F., Wang, Y.-F. & Du, J.-L. Du, Information integration for motor generation. *Curr. Opin. Physiol.* **8**, 116-120 (2019).3

3. McCormick, D. A., Nestvogel, D. B. & He, B.-J. Neuromodulation of brain state and behavior. *Annu. Rev. Neurosci.* **43**, 391-415 (2020).4

4. Witvliet, D. et al. Connectomes across development reveal principles of brain maturation. *Nature* **596**, 257-261 (2021).5

5. Mei, J., Muller, E. & Ramaswamy, S. Informing deep neural networks by multiscale principles of neuromodulatory systems. *Trends. Neurosci.* **45**, 237-250

497   (2022).6

6. Li, F.-N., Zhang, C.-M. & Du, J.-L. Neuromodulatory processing in the bi-pathway brain architecture. *Curr. Opin. Neurobiol.* **93**, 103055 (2025).1

7. Schwarz, L. A. et al. Viral-genetic tracing of the input-output organization of a central noradrenaline circuit. *Nature* **524**, 88-92 (2015).7

8. Schwarz, L. A. & Luo, L.-Q. Organization of the locus coeruleus-norepinephrine system. *Curr. Biol.* **25**, R1051-R1056 (2015).8

9. Zhang, C.-M. et al. Multiple-level organizational logic of the locus coeruleus-norepinephrine system in structure and function. *bioRxiv* (2025). doi:10.1101/2025.06.11.659024 9

10. Yee, A. G., Liao, Y., Muntean, B. S. & Ford, C. P. Discrete spatiotemporal encoding of striatal dopamine transmission. *Science* **389**, 200-206 (2025).10

11. Cupolillo, D., Regio, V. & Barberis, A. Synaptic microarchitecture: the role of spatial interplay between excitatory and inhibitory inputs in shaping dendritic plasticity and neuronal output. *Front. Cell. Neurosci.* **18**, 1513602 (2024).11

12. Du, X.-F. et al. Central nervous system atlas of larval zebrafish constructed using the morphology of single excitatory and inhibitory neurons. *bioRxiv* (2025). doi:10.1101/2025.06.06.658008 13

13. Wang, Y.-F. et al. BrAVe: a unified framework for 3D interactive and integrative exploration and analysis of cross-modal brain atlas data. *bioRxiv* (2025). doi:10.1101/2025.10.04.680445 14

14. Swanson, L. W. & Lichtman, J. W. From Cajal to connectome and beyond. *Annu. Rev. Neurosci.* **39**, 197-216 (2016).15

15. Brittin, C. A., Cook, S. J., Hall, D. H., Emmons, S. W. & Cohen, N. A multi-scale brain map derived from whole-brain volumetric reconstructions. *Nature* **591**, 105-110 (2021).16

16. Luo, L.-Q. Architectures of neuronal circuits. *Science* **373**, eabg7285 (2021).17

17. Tay, T. L., Ronneberger, O., Ryu, S., Nitschke, R. & Driever, W. Comprehensive catecholaminergic projectome analysis reveals single-neuron integration of zebrafish ascending and descending dopaminergic systems. *Nat. Commun.* **2**, 171 (2011).18

18. Kim, H. R. et al. A unified framework for dopamine signals across timescales. *Cell* **183**, 1600–1616.e25 (2020).19

19. Dag, U. et al. Dissecting the functional organization of the C.elegans serotonergic system at whole-brain scale. *Cell* **186**, 2574-2592.e20 (2023).20

20. Mondoloni, S. et al. Serotonin release in the habenula during emotional contagion promotes resilience. *Science* **385**, 1081-1086 (2024).21

21. Jiao, Z.-L. et al. Projectome-based characterization of hypothalamic peptidergic neurons in male mice. *Nat. Neurosci.* **28**, 1073-1088 (2025).22

22. Wanner, A. A., Genoud, C., Masudi, T., Siksou, L. & Friedrich, R. W. Dense EM-based reconstruction of the interglomerular projectome in the zebrafish olfactory bulb. *Nat. Neurosci.* **19**, 816-825 (2016).23

23. Liang, L. et al. A fine-scale functional logic to convergence from retina to thalamus. *Cell* **173**, 1343-1355.e24 (2018).24

24. Motta, A. et al. Dense connectomic reconstruction in layer 4 of the somatosensory
cortex. *Science* **366**, eaay3134 (2019).25

25. Morgan, J. L. & Lichtman, J. W. An individual interneuron participates in many
kinds of inhibition and innervates much of the mouse visual thalamus. *Neuron*
106, 468-481.e2 (2020).26

26. Gour, A. et al. Postnatal connectomic development of inhibition in mouse barrel
cortex. *Science* **371**, eabb4534 (2021).27

27. Uytiepo, M. et al. Synaptic architecture of a memory engram in the mouse
hippocampus. *Science* **387**, eado8316 (2025).28

28. White, J. G., Southgate, E., Thomson, J. N. & Brenner, S. The structure of the
nervous system of the nematode Caenorhabditis elegans. *Philos. Trans. R. Soc.*
Lond. B Biol. Sci. **314**, 1-340 (1986).29

29. Zheng, Z.-H. et al. A complete electron microscopy volume of the brain of adult
drosophila melanogaster. *Cell* **174**, 730-743.e22 (2018).30

30. Cook, S. J. et al. Whole-animal connectomes of both Caenorhabditis elegans
sexes. *Nature* **571**, 63-71 (2019).31

31. Scheffer, L. K. et al. A connectome and analysis of the adult Drosophila central
brain. *Elife* **9**, e57443 (2020).32

32. Winding, M. et al. The connectome of an insect brain. *Science* **379**, eadd9330
(2023).33

33. Dorkenwald, S. et al. Neuronal wiring diagram of an adult brain. *Nature* **634**,
124-138 (2024).34

34. Kasthuri, N. et al. Saturated reconstruction of a volume of neocortex. *Cell* **162**,
648-661 (2015).35

35. Loomba, S. et al. Connectomic comparison of mouse and human cortex. *Science*
377, eabo0924 (2022).36

36. Shapson-Coe, A. et al. A petavoxel fragment of human cerebral cortex
reconstructed at nanoscale resolution. *Science* **384**, eadk4858 (2024).37

37. MICrONS Consortium. Functional connectomics spanning multiple areas of
mouse visual cortex. *Nature* **640**, 435-447 (2025).38

38. Ding, Z.-K. et al. Functional connectomics reveals general wiring rule in mouse
visual cortex. *Nature* **640**, 459-469 (2025).39

39. Helmstaedter, M. Synaptic-resolution connectomics: towards large brains and
connectomic screening. *Nat. Rev. Neurosci.* **27**, 101-120 (2026).40

40. Hildebrand, D. G. C. et al. Whole-brain serial-section electron microscopy in
larval zebrafish. *Nature* **545**, 345-349 (2017).41

41. Wanner, A. A. & Friedrich, R. W. Whitening of odor representations by the wiring
diagram of the olfactory bulb. *Nat. Neurosci.* **23**, 433-442 (2020).42

42. Svara, F. et al. Automated synapse-level reconstruction of neural circuits in the
larval zebrafish brain. *Nat. Methods* **19**, 1357-1366 (2022).43

43. Vishwanathan, A. et al. Predicting modular functions and neural coding of
behavior from a synaptic wiring diagram. *Nat. Neurosci.* **27**, 2443-2454
(2024).44

44. Boulanger-Weill, J. et al. Correlative light and electron microscopy reveals the

fine circuit structure underlying evidence accumulation in larval zebrafish. *bioRxiv* (2025). doi:10.1101/2025.03.14.643363 45

45. Petkova, M. D. et al. A connectomic resource for neural cataloguing and circuit dissection of the larval zebrafish brain. *bioRxiv* (2025). doi:10.1101/2025.06.10.658982 46

46. Amiry-Moghaddam, M. & Ottersen, O. P. Immunogold cytochemistry in neuroscience. *Nat. Neurosci.* **16**, 798-804 (2013).47

47. Shigemoto, R. & Joesch, M. The genetic encoded toolbox for electron microscopy and connectomics. *Wiley Interdiscip. Rev. Dev. Biol.* **6** (2017).48

48. Lam, S. S. et al. Directed evolution of APEX2 for electron microscopy and proximity labeling. *Nat. Methods* **12**, 51-54 (2015).12

49. Usher, M., Cohen, J. D., Servan-Schreiber, D., Rajkowski, J. & Aston-Jones, G. Aston-Jones, The role of locus coeruleus in the regulation of cognitive performance. *Science* **283**, 549-554 (1999).

50. Berridge, C. W. & Waterhouse, B. D. The locus coeruleus-noradrenergic system: modulation of behavioral state and state-dependent cognitive processes. *Brain Res. Brain Res. Rev.* **42**, 33-84 (2003).

51. Aston-Jones, G. & Cohen, J. D. An integrative theory of locus coeruleus-norepinephrine function: adaptive gain and optimal performance. *Annu. Rev. Neurosci.* **28**, 403-450 (2005).

52. Zerbi, V. et al. Rapid reconfiguration of the functional connectome after chemogenetic locus coeruleus activation. *Neuron* **103**, 702-718.e5 (2019).

53. Poe, G. R. et al. Locus coeruleus: a new look at the blue spot. *Nat. Rev. Neurosci.* **21**, 644-659 (2020).

54. Eckstein, N. et al. Neurotransmitter classification from electron microscopy images at synaptic sites in Drosophila melanogaster. *Cell* **187**, 2574-2594.e23 (2024).

55. Elabbady, L. et al. Perisomatic ultrastructure efficiently classifies cells in mouse cortex. *Nature* **640**, 478-486 (2025).

56. Sun, R.-B. et al. PKU tags, novel genetically encoded shape tags for cell labeling in light and electron microscopy. *bioRxiv* (2025). doi:10.1101/2025.11.25.690457

57. Tao, C.-L. et al. Differentiation and characterization of excitatory and inhibitory synapses by cryo-electron tomography and correlative microscopy. *J. Neurosci.* **38**, 1493-1510 (2018).

58. Christie, M. J., Williams, J. T. & North, R. A. Electrical coupling synchronizes subthreshold activity in locus coeruleus neurons in vitro from neonatal rats. *J. Neurosci.* **9**, 3584-3589 (1989).

59. Ishimatsu, M. & Williams, J. T. Synchronous activity in locus coeruleus results from dendritic interactions in pericoerulear regions. *J. Neurosci.* **16**, 5196-5204 (1996).

60. Uematsu, A. et al. Modular organization of the brainstem noradrenaline system coordinates opposing learning states. *Nat. Neurosci.* **20**, 1602-1611 (2017).

61. Totah, N. K., Neves, R. M., Panzeri, S., Logothetis, N. K. & Eschenko, O. The locus coeruleus is a complex and differentiated neuromodulatory system. *Neuron* **99**, 1055-1068.e6 (2018).

62. Chandler, D. J. et al. Redefining noradrenergic neuromodulation of behavior: impacts of a modular locus coeruleus architecture. *J. Neurosci.* **39**, 8239-8249 (2019).

63. Luskin, A. T. et al. Heterogeneous pericoerulear neurons tune arousal and exploratory behaviours. *Nature* **643**, 437-447 (2025).

64. Cruz-Lopez, D., Ramos, D., Castilloveitia, G. & Schikorski, T. Quintuple labeling in the electron microscope with genetically encoded enhanced horseradish peroxidase. *PLoS One* **13**, e0200693 (2018).

65. Zhang, Q.-Y., Lee, W. A., Paul, D. L. & Ginty, D. D. Multiplexed peroxidase-based electron microscopy labeling enables simultaneous visualization of multiple cell types. *Nat. Neurosci.* **22**, 828-839 (2019).

66. Atasoy, D. et al. A genetically specified connectomics approach applied to long-range feeding regulatory circuits. *Nat. Neurosci.* **17**, 1830-1839 (2014).

67. Costa, M., Manton, J. D., Ostrovsky, A. D., Prohaska, S. & Jefferis, G. S. NBLAST: rapid, sensitive comparison of neuronal structure and construction of neuron family databases. *Neuron* **91**, 293-311 (2016).

68. Suzuki, M., Pennartz, C. M. A. & Aru, J. How deep is the brain? The shallow brain hypothesis. *Nat. Rev. Neurosci* **24**, 778-791 (2023).

69. Magee, J. C. Dendritic integration of excitatory synaptic input. *Nat. Rev. Neurosci.* **1**, 181-190 (2000).

70. Jarsky, T., Roxin, A., Kath, W. L. & Spruston, N. Conditional dendritic spike propagation following distal synaptic activation of hippocampal CA1 pyramidal neurons. *Nat. Neurosci.* **8**, 1667-1676 (2005).

71. Takahashi, H. & Magee, J. C. Pathway interactions and synaptic plasticity in the dendritic tuft regions of CA1 pyramidal neurons. *Neuron* **62**, 102-111 (2009).

72. Isaacson, J. S. & Scanziani, M. How inhibition shapes cortical activity. *Neuron* **72**, 231-243 (2011).

73. Gidon, A. & Segev, I. Principles governing the operation of synaptic inhibition in dendrites. *Neuron* **75**, 330-341 (2012).

74. Boivin, J. R. & Nedivi, E. Functional implications of inhibitory synapse placement on signal processing in pyramidal neuron dendrites. *Curr. Opin .Neurobiol.* **51**, 16-22 (2018).

75. Suzuki, M. & Larkum, M. E. General anesthesia decouples cortical pyramidal neurons. *Cell* **180**, 666-676.e13 (2020).

76. Megías, M., Emri, Z., Freund, T. F. & Gulyás, A. I. Total number and distribution of inhibitory and excitatory synapses on hippocampal CA1 pyramidal cells. *Neuroscience* **102**, 527-540 (2001).

77. Beaulieu-Laroche, L. et al. Enhanced dendritic compartmentalization in human cortical neurons. *Cell* **175**, 643-651.e14 (2018).

78. Ferguson, K. A. & Cardin, J. A. Mechanisms underlying gain modulation in the cortex. *Nat. Rev. Neurosci.* **21**, 80-92 (2020).

79. Iascone, D. M. et al. Whole-neuron synaptic mapping reveals spatially precise excitatory/inhibitory balance limiting dendritic and somatic spiking. *Neuron* **106**, 566-578.e8 (2020).

80. Wright, W. J., Hedrick, N. G. & Komiyama, T. Distinct synaptic plasticity rules operate across dendritic compartments in vivo during learning. *Science* **388**, 322-328 (2025).

81. Robert, V. et al. Cortical glutamatergic and GABAergic inputs support learning-driven hippocampal stability. *Science* **390**, eadn0623 (2025).

82. Ghosh, S. & Maunsell, J. H. R. Maunsell, Locus coeruleus norepinephrine contributes to visual-spatial attention by selectively enhancing perceptual sensitivity. *Neuron* **112**, 2231-2240.e5 (2024).

83. Grimm, C. et al. Tonic and burst-like locus coeruleus stimulation distinctly shift network activity across the cortical hierarchy. *Nat. Neurosci.* **27**, 2167-2177 (2024).

84. Boergens, K. M. et al. webKnossos: efficient online 3D data annotation for connectomics. *Nat. Methods* **14**, 691-694 (2017).

85. Dorkenwald, S. et al. FlyWire: online community for whole-brain connectomics. *Nat. Methods* **19**, 119-128 (2022).

86. Schmidt, M., Motta, A., Sievers, M. & Helmstaedter, M. RoboEM: automated 3D flight tracing for synaptic-resolution connectomics. *Nat. Methods* **21**, 908-913 (2024).

87. Celii, B. et al. NEURD offers automated proofreading and feature extraction for connectomics. *Nature* **640**, 487-496 (2025).

88. Dorkenwald, S. et al. CAVE: Connectome Annotation Versioning Engine. *Nat. Methods* **22**, 1112-1120 (2025).

89. Joesch, M. et al. Reconstruction of genetically identified neurons imaged by serial-section electron microscopy. *Elife* **5**, e15015 (2016).

Methods

Zebrafish husbandry

Adult Zebrafish (*Danio rerio)* were kept in a circulating system at 28℃ under a 14:10 hour light-dark cycle. All animal protocols were reviewed and approved by the Animal Care and Use Committee of the Center for Excellence in Brain Science and Intelligence Technology (CEBSIT), Chinese Academy of Sciences (NA-046-2023). Larvae were fed with *Brachionid* rotifers from $4.5 - 9.0$ dpf, then switched to freshly hatched *Artemia* nauplii thereafter. The sex of embryonic and larval zebrafish was undetermined.

Zebrafish lines

All transgenic zebrafish larvae were in *nacre* background. The *vmat2*, *th1*, *dbh*, *tph2*, *hcrt*, *glyt2*, *vglut2a*, *gad1b* promoters were used to label NE/DA/5HT, NE/DA, NE, 5HT, Hcrt, Gly, Glu, and GABA neurons, respectively. Published fish lines used in this study include *Tg(slc6a5:GFP)cf3* (*abbreviation*: *Tg(glyt2:GFP))*[90], *Tg(elavl3:Hsa.H2B-GCaMP6f)jf7* (*Tg(HuC:H2B-GCaMP6f))*[91], *Tg(5×UAS-hsp70l:GCaMP6s)nkUAShspzGCaMP6s13a* (*Tg(UAS:GCaMP6s))*[92], *Ki(th:GAL4-VP16)ion32d* (*Ki(th:GAL4-VP16))*[93], *Ki(tph2:EGFP)ion39d* (*Ki(tph2:EGFP))*[93], *Tg(slc17a6:GAL4FF)ion177d* (*Tg(vglut2a:GAL4FF))*[94]. The unpublished fish lines *TgBAC(gad1b:GAL4-VP16)ion19h*[12] and *gSAIzGFFM242A* (*Gt(vmat2:GAL4FF))* were provided by Dr. Jie He and Dr. Koichi Kawakami, respectively. *Ki(tph2:GAL4FF,myl7:EGFP)ion127d* (*Ki(tph2:GAL4FF,cmlc2:EGFP))* was constructed using a CRISPR/Cas9-based knockin method[93]. To generate the larva for Fish-X dataset, *Tg(dbh:GAL4FF;5×UAS-hsp:cytosol-APEX2-TagRFP)ion142d* (*Tg(dbh:GAL4FF;UAS:cytosol-APEX2-TagRFP))*, *Tg(hcrt:APEX2-TagRFP-NLS)ion100d* (*Tg(hcrt:APEX2-TagRFP-NLS))*, and *Tg(slc6a5:Hsa.Cox8a-APEX2-TagRFP)ion97d* (*Tg(glyt2:MITO-APEX2-TagRFP))* were constructed using the Tol2-based transgenic system[95]. All other APEX2-based transgenic lines and plasmids were listed in Supplementary Table 1. The *APEX2* sequence was cloned from the commercial plasmid *Bact2-APEX2-GBP*[96] (Addgene, Cat# 67668). The *dbh*, *hcrt*, and *glyt2* promoter sequences were cloned from plasmids *pTol2-dbh (short)-GAL4FF-5XUAS-hsp-TRPV1-RFPT*, *pTol2-hcrt-GAL4-VP16*, and *pTol2-glyt2-P2A-ECFP*, respectively. Signal peptide sequences were listed in Supplementary Table 2. Original PKU constructs were provided by Dr. Yu-Long Li.

Immunostaining

Larvae were fixed overnight on a shaker at 4℃ using 4% paraformaldehyde (PFA, Sigma, Cat# 158127) prepared in phosphate-buffered saline (PBS). Following three 5-min washes with PBS containing 1% Tween-20 (PBST), brains were carefully

dissected using fine forceps and subjected to another three PBST washes. Samples then underwent graded dehydration and rehydration through sequential incubation in (v/v): 25% methanol/75% PBST (5 min, RT), 50% methanol/50% PBST (5 min, RT), 75% methanol/25% PBST (5 min, RT), 100% methanol (4 h, -20℃), followed by reverse gradient to 25% methanol, then three 5-min PBST washes at RT. After ddH$_2$O rinse (5 min, RT) and acetone treatment (10 min, -20℃), samples were rehydrated through ddH$_2$O and three final PBST washes. Blocking was performed overnight at 4℃ with gentle shaking, followed by incubation with primary antibodies (1:200 in blocking buffer) at 4℃ for 24 – 48 h. After primary antibody recovery and three quick PBST rinses, extended washing was performed (2 × 1 h, 1 × 20 min, 1 × 10 min). Secondary antibodies (1:500 in blocking buffer) were applied at 4℃ for 36 h, followed by removal, three quick PBST washes, and final rinses (3 × 10 min, 1 × 30 min) before light-protected storage on ice for fluorescence imaging. Commercial antibodies included Anti-Orexin-A antibody (Millipore, Cat# AB3704), Anti-Tyrosine Hydroxylase antibody (Millipore, Cat# MAB318), Alexa Fluor® 488 Goat anti-Rabbit secondary antibody (Abcam, Cat# AB150077), and Alexa Fluor® 568 Goat anti-Mouse secondary antibody (Invitrogen, Cat# A-11004).

In vivo structural and functional imaging

At 3 dpf, multiplexed APEX2-labeled larvae were initially screened using an upright fluorescence stereomicroscope. The preparations, embedded in 1.5% low-melting-point agarose (Invitrogen, Cat# 15517014), were carefully verified at 5 dpf using an upright fluorescence microscope. The labeling efficiency was evaluated to determine larvae were used for further EM sample preparation. At 6 dpf, larvae were paralyzed with 1 mg/mL α-bungarotoxin (Tocris, Cat# 2133) and dorsally embedded in 1.5% low-melting-point agarose. Morphological fluorescence imaging was performed using an upright confocal laser scanning microscope (FV3000, Olympus, Japan) equipped with a 40× water-immersion objective (NA = 0.8). Imaging parameters included a frame size of 635.9×635.9 μm² (0.621×0.621 μm² per pixel), with two-field stitching along the rostrocaudal axis to cover the SC, Z-axis intervals of 1 – 1.5 μm, and 2 – 3 line averaging. Selected brain regions were further imaged at higher zoom factors to resolve fine spatial patterns of APEX2 signals.

To simultaneously record calcium dynamics in LC-NE, PT-DA, and RN-5HT neurons, we performed *in vivo* calcium imaging on 6-dpf *Ki(th:GAL4-VP16);Ki(tph2:GAL4FF,cmlc2:EGFP);Tg(UAS:GCaMP6s)* larvae, using a confocal microscope (FV3000, Olympus, Japan). Imaging parameters included a frame size of 636.4×636.4 μm² (1.243×1.243 μm² per pixel), with resonant scanning mode, imaging depth of ~200 μm, Z-axis intervals of 10 μm, volumetric imaging rate at ~0.72 Hz, and total acquisition duration of ~30 min per session.

DAB staining

The optimized DAB staining protocol was adapted from published procedures[97]. The 0.1 M Tris-HCl buffer (pH 7.4) was prepared by dissolving 0.6055 g Tris-Base (Millipore, Cat# 648311) in 50 mL of ultrapure water, with pH adjustment using concentrated HCl. A 10 mg tablet of 3,3'-diaminobenzidine tetrahydrochloride (DAB·4HCl, Sigma, Cat# D5905) was subsequently dissolved in 10 mL of this buffer through vigorous vortexing under light-protected conditions. The solution was then filtered through a 0.22-μm filter (Millipore, Millex™-GP 0.22-μm filter) to obtain a 1 mg/mL DAB stock solution, which was stored at 4℃ and protected from light. Samples were sequentially rinsed with 0.1 M phosphate buffer (3 × 10 min), Tris-HCl buffer (2 × 10 min), and DAB solution (2 × 10 min) on a shaker at 4℃ to remove residual aldehydes. For the reaction, 1 μL of 30% H_2O_2 (Sinopharm Chemical Reagent Co., Ltd., Cat# 10011218) was mixed with 1 mL DAB stock to prepare the working solution, and then samples were incubated for 2 h at RT in complete darkness. The reaction was terminated by washing with Tris-HCl buffer (2 × 10 min). The catalysis by APEX2 of DAB polymerization resulted in dark brown precipitates, which were clearly visible under bright-field stereomicroscope. Post-fixation with osmium tetroxide (OsO_4) was conducted immediately following DAB staining to preserve the osmiophilic reaction products.

EM sample preparation

A mixed fixative containing 2% PFA and 2% glutaraldehyde (GA) was used for effective whole-brain fixation. A 1 mL aliquot of 25% GA (Sigma, Cat# G5882, stored at -20℃) was thawed at 4℃ under light protection. Concurrently, 1 g PFA powder was dissolved in 20 mL of ultrapure water within a fume hood, with 1 – 2 drops of 1 M NaOH added. The solution was then sealed, protected from light, and heated at 65℃ with magnetic stirring until fully clarified. After cooling to RT, the solution was filtered through a 0.22 μm filter to yield 5% PFA stock. The 10-mL working fixative was prepared immediately before dissection by mixing 4 mL of 5% PFA, 0.8 mL of 25% GA, 5 mL of 0.1 M phosphate buffer (pH 7.4), and 0.2 mL of ultrapure water. Larvae were embedded in 1.5% low-melting-point agarose dissolved in extracellular solution (in mM: 134 NaCl, 2.9 KCl, 2.1 $CaCl_2 \cdot 2H_2O$, 1.2 $MgCl_2 \cdot 6H_2O$, 10 HEPES, 10 glucose; 290 mOsm, pH 7.8). The agarose covering dorsal epidermis and pectoral fins was removed. Using glass micropipettes, the skin was carefully dissected from the hindbrain ventricle while avoiding damage to tissues. The trunk was rapidly transected anterior to the swim bladder using ophthalmic scissors, after which the head was gently extracted with fine forceps and immediately immersed in fresh fixative. Fixed samples were sealed and stored at 4℃ in darkness overnight.

Subsequent procedures were adapted from the rOTO protocol (*98, 99*). First, a
1:1 (v/v) mixture of 0.2 M phosphate buffer and 2% OsO_4 (TED PELLA, Cat# 18451)
was prepared to obtain an OsO_4-phosphate solution, which was then used to prepare a
reduced OsO_4-phosphate solution containing 0.03 g/mL potassium hexacyanoferrate
trihydrate (Sigma, Cat# P9387). All OsO_4-containing solutions were freshly prepared
within a fume hood, tightly sealed, protected from light at 4℃, and properly collected
for disposal after use. Fixed samples were washed with 0.1 M phosphate buffer at 4℃
(4 × 10 min) and then immersed in the OsO_4-phosphate solution (200 − 500 μL per
1.5 mL tube, 2 h, 4℃). Concurrently, a 0.01 g/mL aqueous thiocarbohydrazide (TCH,
Sigma, Cat# 88535) solution was placed in a 60℃ oven and gently flipped to
facilitate dissolution. The TCH solution was removed from the oven and allowed to
cool slightly above RT before use. The solution was then filtered through a 0.22 μm
filter directly into the sample tube that had been rinsed with ultrapure water (3 × 10
min, 4℃; 1 × 10 min, RT). After a 30-min light-protected TCH treatment at RT,
samples were washed with ultrapure water (4 × 10 min, 4℃). The reduced
OsO_4-phosphate solution was then added (200 − 500 μL per 1.5 mL tube, 30 min, RT).
Afterward, samples were washed with ultrapure water (4 × 10 min, 4℃). Meanwhile,
a 2% aqueous uranyl acetate solution (EMS, Cat# 22400) was centrifuged at 13,000
rpm, and the supernatant was carefully transferred into the sample tube (200 μL per
1.5 mL tube). Samples were then sealed and stored at 4℃ overnight, protected from
light.

On the next day, the uranyl acetate waste solution was collected for proper
disposal, followed by washing samples with ultrapure water (3 × 10 min).
Subsequently, a graded dehydration series was performed on ice using ethanol
solutions (30%, 50%, 70%, 80%, 90%, 95%, and 100% v/v, each for 10 min)
followed by anhydrous acetone (2 × 10 min). On the day of fixation, the resin volume
was calculated, and the premix was prepared by sequentially pipetting 10 mL of
Pon-812 epoxy resin monomer (SPI, Cat# 90529-77-4), 3.36 mL of dodecenyl
succinic anhydride (DDSA, SPI, Cat# 26544-38-7), and 7.04 mL of nadic methyl
anhydride (NMA, SPI, Cat# 25134-21-8) into a pre-dried tube at RT to form a
three-component premix. This was then sealed and mixed on a rotary mixer for at
least 3 hours. Resin infiltration was conducted immediately following dehydration.
400 μL DMP-30 epoxy accelerator (SPI, Cat# 90-72-2) was added to 20 mL premix,
followed by 30-min rotary mixing to prepare the final resin solution. The resin
solution was either used immediately after a 30-min bubble-settling period or kept
mixing to prevent polymerization at RT. Samples underwent graded acetone-resin
infiltration at ratios of 3:1, 1:1, and 1:3 (v/v) for 8 h, 8h, and overnight, respectively.
They were then transferred to resin-filled vials, subject to negative pressure using a 5
mL syringe, and placed on a horizontal shaker for resin infiltration.

This process was conducted over two days with resin changes occurring twice
daily at intervals of 8 – 12 h. All procedures were performed under controlled
environmental conditions (temperatures < 25℃; humidity < 30%). Following the
two-day resin infiltration, samples were transferred to pre-dried 21-well flat molds
and oriented using sterile toothpicks. The molds were then placed in a drying box
containing silica gel beads and polymerized at 37℃ for 8 – 12 h. After verifying and
minimally adjusting sample orientation at night, the temperature was first increased to
45℃ overnight, then to 60℃ the following morning. Fully polymerized blocks were
removed from the oven after 48 h and stored in a dry cabinet.

X-ray tomography
X-ray tomography was conducted to rapidly and non-destructively evaluate the
quality of EM samples using the Xradia 520 Versa system (Zeiss, Germany) at the
Instrumental Analysis Center of Shanghai Jiao Tong University. Samples exhibiting
uniform staining, minimal deformation, and no apparent mechanical damage were
selected for further ultrathin sectioning and SEM imaging. Samples were scanned at
60 kV with 1 μm^3 voxel resolution (~2 h per sample). Data processing was conducted
using Dragonfly (Object Research Systems, Canada). Sequential images were
imported with resampling parameters and voxel resolution specified. Sample
orientation was interactively manipulating reference axes in the display window.
Selected views were exported as projection sequences via the "*Derive new from
current view*" function. 3D animations were created using the *Video marker* tool. The
video denoising was performed using an open-source Real-ESRGAN-General-x4v3
model.

Whole-brain ultrathin section collection
Using an ultramicrotome equipped with an automated tape-collecting system
(ATUMtome, RMC, USA), 33-nm serial sections were obtained[40]. Prior to collection,
trimmed excess blank resin around the larva using a double-edged razor blade. The
front guide slot of the ATUMtome was properly positioned, and parameters including
sectioning speed (0.8 mm/s), tape-advancing speed (0.6 mm/s), and water refilling
interval (5-sec duration every 8 min) were set. The anti-static device and intermittent
water replenishment system were activated to ensure sequential and stable section
collection on hydrophilic, conductive polyimide film strips (KAPTON-PET, DuPont,
USA). If section instability was observed during collection (e.g., incomplete sections),
immediately halted sectioning and either repositioned the knife edge or replaced the
diamond knife (UltraMaxi, Diatome, Switzerland). Throughout the process, ambient
conditions were maintained at 25℃ and 60% humidity. Following collection,
section-bearing strips were segmented and mounted onto 4-inch monocrystalline
silicon wafers using 0.8-mm-wide conductive adhesive tape (SPI, USA), with careful

pressing to eliminate air bubbles. The mounted wafer was placed into a LEICA EM ACE 600 high-vacuum coater for carbon deposition (6 – 10 nm film thickness). The section surfaces were cleaned using nitrogen gas blowing, and subsequently stored in a dry cabinet at 25℃ and 30% humidity. The entire collection process, which spent three days and utilized three diamond knives, yielded an overall damage rate of 6.56%. Sections were numbered #2594 – #25480 in a caudal-to-rostral order. The damaged 1,503 sections with folds, tears, and severe contamination were algorithmically repaired to ensure that intact portions were registered to adjacent sections[100]. The majority of defects did not occur at identical locations across adjacent sections, thus collectively posing no substantial impact on proofreading.

SEM image acquisition

To enhance sample stability and backscattered electron (BSE) signal contrast, wafers were pre-irradiated using a custom-built irradiation system (9 mA, 2 kV, 10 s continuous exposure). Post-irradiation wafers were transferred to an optical microscope (Clippers FBT, China) for the acquisition of navigation images via line-scan imaging with 7.6 μm pixel resolution. Image acquisition was conducted simultaneously across three SEMs (Navigator NeuroSEM100, FBT, China), each equipped with a custom-built collection software designed to maximize efficiency in image output. The pixel resolution was set to 4 nm, the dwell time to 43.5 ns, and the acceleration voltage to 2 kV. Auto focus and astigmatic correction were configured to 80 s per section. To expedite acquisition efficiency and imaging success rate, a region-specific focusing protocol coupled with photography was implemented. Each section was divided into a grid of ~54 tiles (arranged in 9 rows × 6 columns) at 12,000×12,000 pixels per tile. The adjacent tiles were acquired with a 12% overlapping area. 16 tiles (4×4 arrangement) constituted one standard region. Edge regions containing fewer than 16 tiles were processed equivalently. Regions were sequentially numbered and navigated in a serpentine pattern. Prior to high-resolution scanning, a low-resolution focus was conducted to ensure the algorithm efficacy in pinpointing the optimal focal range at the center of the section, which was set at a resolution of 50 nm with a dwell time of 80 ns. The stage was then oriented to the first region's center for high-resolution focusing and astigmatism correction with a resolution of 4 nm and a dwell time of 80 ns, followed by snake-pattern tile acquisition. After capturing all tiles within a region, the stage advanced to the next region (2.5 s/tile for positional adjustment and data transfer). This cycle iterated until full-section coverage was achieved, maintaining serpentine navigation between regions. The process was repeated automatically for subsequent sections until wafer-wide imaging completion.

To enhance the accuracy and robustness of large-scale EM image processing, we

introduced an image quality assessment and tissue validity detection module prior to
algorithmic analysis. A 2D U-Net model was trained to evaluate image sharpness and
assigned a quality score to each section. Low-quality images were then reviewed
using a custom annotation tool, allowing experts to determine whether re-acquisition
or replacement was necessary. Besides, we implemented CNN-based tissue mask
generation. A U-Net model was trained on three manually annotated chunks
(8,600×8,600×31 voxels at 32×32×33 nm^3 per voxel) to classify voxels into brain
tissues, backgrounds, and defect regions (e.g., folds, tears). All predicted masks were
manually reviewed. The quality scores and tissue masks were applied throughout the
processing pipeline for artifact exclusion and region-of-interest selection, supporting
segmentation and agglomeration.

Image stitching and alignment
The 3D volume registration of EM images was accomplished through tile stitching
and section alignment. The stitching workflow proceeded through three computational
stages: 1) feature detection and matching using Scale-Invariant Feature Transform
(SIFT) descriptors[101] extracted from tile overlap areas; 2) pairwise displacement
estimation via coordinate transformation analysis; 3) global position optimization
through linear least squares minimization. Specifically, SIFT-matched feature point
pairs served as the basis for calculating local translation vectors between tiles. To
mitigate error accumulation, a global optimization framework simultaneously
adjusted all tile positions by minimizing the total displacement error across the entire
section.
A multi-scale alignment method was implemented to register serial sections, first
through coarse-scale processing followed by fine-scale refinement. To address error
accumulation, the pipeline employed a sub-volume-division strategy, partitioning the
original dataset into 764 spatially coherent stacks (each containing 31 consecutive
sections). Continuity across stacks was established by selecting the first section of
each stack as a reference section (764 total). These references underwent affine
transformation to correct notable deformations while maintaining spatial invariance
throughout alignment, serving as stable pivots to prevent cumulative error
propagation. Alignment then proceeded within each stack, from coarse to fine scales.
Sections were downsampled 32-fold during coarse-scale alignment. Correspondences
between two adjacent sections were initially extracted using SIFT descriptors and
subsequently filtered using the RANSAC algorithm[102]. Affine transformation was
applied to align adjacent sections. Sections were downsampled 8-fold during
fine-scale alignment. Damaged sections were first aligned to their neighbors using an
expected affine method[91]. Subsequently, an optimized elastic method[103] generated
dense correspondence fields between adjacent sections, guiding the iterative

refinement of deformation meshes that balanced local adaptability with global smoothness. Finally, Thin-Plate Spline (TPS) transformations[104] extrapolated these deformations to original-resolution sections, preserving ultrastructural continuity at synaptic resolution.

Bidirectional CLEM

To match the resolution of the brain template of the Zebrafish Mesoscopic Atlas (1 μm^3 per voxel), the 764 reference sections mentioned above were further downsampled in the xy-plane to generate downsampled whole-brain EM images ($992 \times 992 \times 990$ nm³ per voxel). Multimodal image registration was accomplished through a three-stage approach. First, whole-brain nuclei were extracted from both the Zebrafish Mesoscopic Atlas and the ssEM dataset to create nuclei point clouds, which were then rigidly aligned using the Iterative Closest Point (ICP) algorithm[105]. Manually-annotated nuclei from four 6-dpf *Tg(HuC:H2B-GCaMP6f)* larvae were used to generate the average point cloud for the LM brain template, while nuclei from the ssEM dataset were automatically segmented. Second, the EM images were globally constrained using X-ray tomography data, to correct the distortion along the rostral-caudal axis ("banana effect"). Third, bidirectional deformation fields between LM and EM spaces were computed through 3D template matching and TPS transformations. Soma regions across the entire brain, along with other EM-identifiable regions including the pineal gland, pituitary gland, and bilateral optic tectal neuropils, were manually annotated on downsampled whole-brain EM images and registered with their counterparts in the brain template of the Zebrafish Mesoscopic Atlas, serving as reference landmarks for joint optimization.

Chunk-based 3D reconstruction

Whole-brain 3D reconstruction was performed through an automated pipeline consisting of: (1) image enhancement and denoising, (2) local registration, (3) dense segmentation, (4) intra-chunk agglomeration, and (5) inter-chunk stitching. To efficiently process large-scale computational tasks, we implemented a distributed computing strategy based on spatial block partitioning. Specifically, the original 3D volume was divided into a series of overlapping chunks to ensure boundary continuity and consistency in prediction. These chunks were then preprocessed and spatially realigned, and subsequently treated as independent inference tasks scheduled for parallel execution. The entire computational workflow was deployed on a high-performance distributed parallel computing architecture managed by the Slurm (Simple Linux Utility for Resource Management) workload manager. All sub-tasks were automatically submitted via batch scripts to a Slurm-managed HPC (High-Performance Computing Cluster) consisting of more than 30 nodes. Based on the availability of computational resources, the scheduler dynamically distributed

tasks across multiple physical nodes. Each node was equipped with a heterogeneous configuration of GPUs, including NVIDIA® A100, V100, and GeForce RTX 4090 models, thereby maximizing utilization of available hardware accelerators. Upon completion, the outputs of all sub-tasks were automatically aggregated and post-processed to reconstruct the full prediction results over the entire volume. For tasks including the segmentation of neurons, synapses, and mitochondria, the block and overlap sizes were set to 8,600×8,600×31 and 600×600×1 voxels, with a resolution of 4×4×33 nm^3 per voxel. For tasks including the segmentation of nuclei and vessels, the block and overlap sizes were set to 1,075×1,075×31 and 75×75×1 voxels, with a resolution of 32×32×33 nm^3 per voxel.

Preprocessing and dense segmentation

To address the Poisson and Gaussian noise inherent in EM images and to overcome the challenges of self-supervised denoising with degraded inputs, we implemented Blind2Unblind for implicit lossless self-supervised denoising[106]. This framework leveraged a mask-guided medium to enhance and denoise EM images. It consisted of a globally-aware mask mapper driven by mask-based sampling and a non-blind lossless training strategy based on re-visible loss. Next, to enhance local 3D continuity and improve inference performance, we adopted a spatial registration strategy inspired by the preprocessing pipeline of the Flood-Filling Network (FFN)[107]. For each chunk mentioned above, a 4×4 affine transformation matrix was estimated to align it with a global reference space. Neural network inference, aggregation, and merging were then performed on the aligned chunks. After dense segmentation, inverse transformations were applied to map the results back to the original coordinate system, followed by cropping to restore the original dimensions. All affine matrices were precomputed and cached prior to processing to reduce computational overhead.

After preprocessing, we employed a boundary detection-based dense segmentation approach to process the whole-brain EM dataset. The image segmentation network adopted a U-shaped architecture with residual blocks, and was trained to predict nearest-neighbor affinity maps along all three spatial dimensions. The training dataset consisted of seven randomly selected volumes, each comprising 30 consecutive sections with a spatial size of 2,000×2,000×30 voxels at a resolution of 4×4×33 nm^3. All these volumes were manually annotated by experts. A composite loss function combining weighted cross-entropy and Dice loss was employed to enhance segmentation performance, particularly in the presence of class imbalance. To improve model generalization, we applied a range of data augmentation techniques during training, including basic transformations such as rotation, flipping, and elastic deformation, as well as robustness-oriented strategies such as CutBlur, CutNoise, and Mixup. In addition, we simulated common imaging artifacts including registration

shifts, motion blur, and partial slice dropouts, to enhance the model's capability to

handle real-world imperfections, thereby improving the stability and accuracy of the

segmentation.

Intra-chunk agglomeration

To construct the signature graph, we first performed over-segmentation of the

membrane probability map using a 2D watershed algorithm based on distance

transform. Specifically, small connected membrane boundary components were

removed through thresholding. Given the strong anisotropy of the data (with lateral

resolution of 4 nm per voxel and axial resolution of 33 nm per voxel), the Euclidean

distance transform was applied to each membrane probability map. The maximum

distance value was then used as the seed point to execute the watershed algorithm,

producing 2D watershed superpixels. These superpixels were subsequently merged to

achieve the 3D reconstruction. The image could be represented using watershed

superpixels, where the superpixels served as vertices and the physical connectivity

between them formed the edges of a graph model. The superpixel merging problem

was thus formulated as a vertex partitioning problem on the graph. In this study, the

problem is modeled as the minimum cost multicut problem (MCMP) on the

superpixel adjacency graph[108]. For feature extraction, three types of features were

manually extracted: boundary features between superpixels, edge features within each

superpixel, and inter-layer features derived from the anisotropic dataset. A feature

vector of length 625 was designed for each edge between two superpixels. A random

forest classifier was then trained to predict the probability of merging each pair of

superpixels. Based on maximum likelihood estimation, the predicted probability is

converted into the cost for merging the two superpixels. This resulted in an instance of

the multicut problem. Additionally, previously reconstructed and expert-verified

whole-brain nucleus and synapse segmentation results were introduced as soft

priors[109]. Superpixels belonging to the same nucleus should be merged preferentially,

while superpixels on opposite sides of the same synapse should be merged less likely.

At the same time, the aggregation range was confined to the effective tissue mask area

to prevent the merging of defect regions with other superpixels, thereby avoiding

erroneous propagation. Given that the combinatorial optimization problem defined

here is NP-hard, and exact methods are not polynomial-time solvable, this study

framed the problem as a sequential decision-making task for node aggregation on the

graph. The problem was solved optimally using the heuristic greedy additive edge

contraction (GAEC) framework[110]. Finally, the results were mapped back to the

original over-segmented image obtained from the watershed distance transform.

Inter-chunk stitching

Following the preliminary reconstruction of all individual chunks, we developed an

overlap-aware fragment agglomeration algorithm to identify and merge neuronal
segments across adjacent chunks, thereby enabling the reconstruction of globally
continuous neuronal structures. This algorithm integrated a deep learning-based
candidate identification module with a set of rule-based merging constraints.
Specifically, we trained a 3D-ResNet-based neural network classifier to evaluate
fragment pairs extracted from sliding 3D windows within overlapping regions,
producing a probability score indicating whether the pair originated from the same
cell. Candidate merges were then further filtered based on multiple criteria, including
intersection-over-union (IoU) thresholds, voxel overlap chunk, and the presence of
defect masks, to ensure that the final agglomeration decisions maintained both
structural continuity and biological plausibility. By combining large-scale parallel
processing enabled by the Slurm workload manager mentioned above with this
customized post-processing pipeline, we established an efficient workflow for dense
reconstruction.

Segmentation of nuclei and vessels
We annotated nuclei and vessels in every fifth section of the downsampled ssEM
dataset with a resolution of $32\times32\times165$ nm^3, generating high-quality training sets with
dimensions of $9{,}000\times6{,}000\times764$ voxels. These results were used to train a 2D
segmentation network. We employed nnU-Net[111] for training and inference, with a
patch size set to $1{,}024\times1{,}024\times4$ and a batch size of 2. To maintain training stability,
we chose stochastic gradient descent (SGD) with a momentum of 0.99 as the
optimizer and applied a "poly" learning rate policy, starting with a learning rate of
0.01 and a maximum of 1,000 epochs. Upon completion of inference, we restored the
results to the original space, resulting in a coarse reconstruction of the entire volume.
Next, we implemented a sliding window strategy along the z-axis, systematically
traversing connected instances within each window, to achieve a refined
reconstruction of the whole-brain nuclei and vessels. The window length was set to 30,
with an overlap of 4. When the number of voxels within the receptive field was less
than 1000, the z-axis voxel range was under 2, or the instance did not contact adjacent
windows, we classified it as reconstruction noise and removed it. Approximately 35%
of nuclei and all vessels underwent further manual proofreading, followed by
quantitative analysis and 3D visualization.

Nucleus clustering
We developed a feature extraction pipeline, integrating both traditional handcrafted
morphological features and high-level representations learned through self-supervised
learning, to analyze whole-brain nuclei with diverse morphologies.
Manually-proofread 176,789 nuclei (nuclei with incomplete structures in rostralmost
and caudalmost sections were excluded) were processed using the CloudVolume

framework to compute 3D surface meshes and then extracted 13 traditional morphological features using the Trimesh library. These 13 features collectively characterized nucleus compactness (Φ, κ_A, κ_V), surface irregularity (ΔA, ΔA_b), shape elongation (ρ_{max}, R_{range}), and spatial dispersion (r_{max}, r_{min}). The following notations were adopted:

M: 3D triangular mesh representing a nucleus surface.

V: Volume of the nucleus mesh (μm^3).

A: Surface area of the nucleus mesh (μm^2).

C: Geometric centroid of the nucleus mesh.

M_h: Convex hull mesh of M.

V_h, A_h: Volume and surface area of convex hull, respectively.

S_b: Minimal bounding sphere of M.

V_b, A_b: Volume and surface area of bounding sphere, respectively.

v_i: i-th vertex of mesh M.

$\|\cdot\|$: Euclidean norm in $\mathbb{R}^3$.

Φ: Sphericity defined as $\Phi = \pi^{1/3}(6V)^{2/3}/A$, where $\Phi \in [0, 1]$ and $\Phi = 1$ defines a perfect sphere.

κ_A: Surface convexity defined as $\kappa_A = A/A_h$

κ_V: Volume convexity defined as $\kappa_V = V/V_h$

ΔA: Surface concavity defined as $\Delta A = A - A_h$

ΔV: Volume concavity defined as $\Delta V = V_h - V$

ρ_{max}: Elongation ratio defined as $\rho_{max} = \max_i(\|v_i - C\|)/\min_i(\|v_i - C\|)$

R_{range}: Radial range defined as $R_{range} = \max_i(\|v_i - C\|) - \min_i(\|v_i - C\|)$

r_{max}: Normalized max radius defined as $r_{max} = (\max_i(\|v_i - C\|))^3/V$

r_{min}: Normalized min radius defined as $r_{min} = (\min_i(\|v_i - C\|))^3/V$

σ_A: Spherical area ratio defined as $\sigma_A = A/A_b$

σ_V: Spherical volume ratio defined as $\sigma_V = V/V_b$

ΔA_b: Spherical surface excess defined as $\Delta A_b = A_b - A$

ΔV_b: Spherical volume excess defined as $\Delta V_b = V_b - V$

e：Eccentricity computed based on principal component analysis of the mesh vertices as $e = \sqrt{1 - \frac{\lambda_{min}}{\lambda_{max}}}$, where λ_{max} and λ_{min} are the largest and smallest eigenvalues of the mesh vertex covariance matrix, respectively.

To complement these interpretable handcrafted features, a self-supervised deep learning framework was implemented to extract high-level geometric representations of the nucleus shape. Unlike voxel- or mesh-based formats, point clouds provide a compact and resolution-independent representation that preserved fine morphological details. Each nucleus surface mesh M was uniformly sampled using Poisson disk sampling in Trimesh to obtain a point cloud $P \in \mathbb{R}^{n \times 3}$, where $n = 3{,}072$ represented

the number of sampled vertices. The sampling process ensured uniform spatial
distribution across the mesh surface while preserving geometric details. The point
clouds were processed using a Siamese self-supervised architecture based on
SimSiam[112] to learn transformation-invariant descriptors. During training, each point
cloud P was augmented into two distorted views P_1 and P_2 using random rotations,
Gaussian noise, and jittering. The augmented views were processed through a
three-stage neural network pipeline:

$$f: \mathbb{R}^{n \times 3} \to \mathbb{R}^d \to g: \mathbb{R}^d \to \mathbb{R}^{d'} \to h: \mathbb{R}^{d'} \to \mathbb{R}^{d'}$$

where $f(\cdot)$ (Encoder network, PointNet-based backbone) mapped point clouds to d
$= 1{,}024$-dimensional embeddings, $g(\cdot)$ (Projector network, 2-layer MLP) mapped to d'
$= 256$-dimensional projections, and $h(\cdot)$ (Predictor network, 2-layer MLP) with same
dimensionality. The model was trained to minimize a symmetric negative cosine
similarity loss:

$$\mathcal{L} = -\tfrac{1}{2}[D(h(g(f(P1))), \underline{g(f(P2))}) + D(h(g(f(P2))), \underline{g(f(P1))})]$$

where $D(x, y) = x \cdot y/(\|x\|_2 \|y\|_2)$ denoted the cosine similarity function. The
underlined terms were treated as constants during backpropagation and did not
contribute gradients. This objective encouraged the network to learn rotation- and
noise-invariant representations while preventing representational collapse. Each
nucleus was ultimately represented by a $d = 1{,}024$-dimensional embedding vector $z =$
$f(P)$ extracted from the trained encoder. To reduce redundancy and enable integration
with handcrafted features, we applied Uniform Manifold Approximation and
Projection (UMAP)[113] with the following hyperparameters (Target dimensionality:
$d_{UMAP} = 8$; Number of neighbors: $k = 15$; Minimum distance: $d_{min} = 0.1$; Distance
metric: Euclidean). This yielded an 8-dimensional deep feature representation z_{UMAP}
$\in \mathbb{R}^8$ for each nucleus. The final feature representation for each nucleus was
constructed by concatenating the 8-dimensional deep feature embedding $z_{UMAP} \in$
$\mathbb{R}^8$ with the 13-dimensional handcrafted morphological feature vector $f_{morph} \in \mathbb{R}^{13}$:
$f_{hybrid} = [f_{morph}; z_{UMAP}] \in \mathbb{R}^{21}$, where $[\cdot; \cdot]$ denoted vector concatenation. Prior to
concatenation, all features were standardized using z-score normalization: $f' = (f -$
$\mu)/\sigma$, where μ and σ represented the mean and standard deviation computed across the
entire dataset, respectively. This hybrid descriptor f_{hybrid} leveraged both interpretable
morphological traits (providing biological insight) and high-capacity deep
embeddings (capturing complex geometric patterns), enabling robust and scalable
analysis of nucleus morphology.

Similarly, we adopted the same methodology for morphological analysis of
somata. Since somata could be characterized with the aid of associated structures such
as nuclei and mitochondria, the computation of traditional somatic features
incorporated not only the aforementioned morphological descriptors but also
integrated nucleus- and mitochondria-based features as auxiliary representations.

Segmentation of PSDs, SVCs, and mitochondria

A three-class 3D U-Net network was trained to predict PSDs and SVCs at the resolution of $8\times8\times33$ nm^3 per voxel. We annotated four regions (~$12\times12\times1$ μm^3 per volume), generating training sets encompassing 146M voxels. We optimized the network using the multi-class cross-entropy loss and SGD optimizer. During training, a warm-up learning rate strategy was employed with an initial learning rate of 0.01, a linear warm-up period of 2k steps, and cosine decay over a total of 500k training steps. Data augmentation was performed to improve the network's robustness, including random rotation, random flipping, grayscale transformation, elastic transformation, and blur simulation. To test the network performance, we manually annotated another three regions as test sets (~$8\times8\times1$ μm^3 per volume). The evaluation of these regions yielded an average precision of $85.60 \pm 3.21\%$ and an average recall of $88.37 \pm 1.59\%$.

A 3D U-Net network was trained to detect the mitochondria at the resolution of $8\times8\times33$ nm^3 per voxel. We annotated five regions (~$12\times12\times1$ μm^3 per volume), generating training sets encompassing 467M voxels. To separate adjacent mitochondria, we designed a network to simultaneously predict both a mitochondrial mask (trained with binary cross-entropy loss) and an instance boundary mask (optimized using a combined binary cross-entropy and Dice loss). Training proceeded for 500k iterations with SGD optimizer, applying a learning rate scheduling strategy identical to the synapse detection task. Data augmentation protocols were consistently implemented as in the synapse detection framework. For automated identification of glycinergic neurons, we annotated additional 284 soma-localized mitochondria labeled with APEX2. Neurons were classified as glycinergic if their somata contained ≥ 4 labeled mitochondria.

Proofreading

We conducted extensive manual and semi-automated proofreading of neurons, PSDs, SVs, mitochondria, vessels, and brain regions using a custom software Advanced Connectome Engine (ACE) that integrated the Hadoop- and HBase-based data management, computing modules, and collaborative tools. We used this software to coordinate proofreading tasks, facilitate collaborative proofreading workflows, and log all user operations. Current and immediately preceding proofreading results were compared to evaluate proofreaders' performance. The proofreading team consisted of $30 - 40$ full-time proofreaders from commercial companies and several experts from the laboratories involved. Reconstruction results were represented as graph structures that connected segmentation contours across all serial sections belonging to a single supervoxel. The contours of adjacent layers were compared to identify anomalies and guide proofreaders to target regions requiring correction.

To aid the proofreading workflow, we utilized a series of foundation models for
segmentation[114-116], including the Segment Anything Model (SAM), SAM 2 and
SAEM². Based on these models, we implemented two additional strategies to enhance
ACE's human-computer interaction. First, the SAM ViT-H model checkpoint was
exported to an ONNX (Open Neural Network Exchange) model with additional
quantization and optimization. ONNX Runtime was used to execute these models on
CPU/GPU, which supported cross-platform inference and improved software runtime.
Second, the SAM embeddings generated by encoders were precomputed and saved in
CloudVolume using kempressed format, which demonstrated higher compression
ratios compared to raw and fpzip encodings. For 2D semi-automated segmentation,
we utilized manually traced neuronal skeletons with extra positive/negative points as
input prompts for the SAM model. However, the pretrained SAM failed to accurately
segment densely packed, homogeneous neuronal processes[117]. To address this, we
proposed the finetuned SAEM² with auxiliary tasks to acquire complementary
multi-head information for neuron segmentation in the ssEM dataset. The auxiliary
task involved identifying cell boundaries (plasma membranes) to complement the
original SAM that primarily focused on cell masks. For 3D segmentation, we
employed the SAM 2 augmented with two additional strategies: (1) We trained the
system to predict cross-correlation maps and subsequently computed the shifts
between adjacent 2D EM images. (2) We utilized the distance transformation maps
from the previous cell masks to sample prompt points for generating the next cell
masks.

Micro-mesoscopic morphology comparison
We inferred the neuronal types of APEX2-unlabeled neurons based on morphology
comparison between the ssEM dataset and a zebrafish mesoscopic atlas[12]. A
preprocessed reference file was generated, containing the x-y-z coordinates of all E
and I neurons and their mirrored counterparts from the Zebrafish Mesoscopic Atlas
(n_{Glu} = 12,219, n_{GABA} = 5,371, n_{Gly} = 2,193). We first calculated neuronal skeletons
(stored as swc files) from manually-corrected EM supervoxels. These skeletons were
then transformed into the common physical space of the mesoscopic brain template
using bidirectional CLEM approach mentioned above. Using the soma center of each
EM query neuron as the origin, we retrieved all single-neuron morphological data
within a 25 μm spherical radius based on the preprocessed reference file. For each EM
query neuron, we performed pairwise morphology comparisons with the retrieved
mesoscopic reference neurons. Cross-scale skeleton matching established
morphological correspondences between the same type of neurons across imaging
modalities by quantifying spatial relationships between their reconstructed skeletons.
The matching similarity was constructed by calculating the unidirectional distance (D)

from EM skeletons (S_E) to LM skeletons (S_L) as follows,

$$D = \frac{1}{|S_E|} \sum_{p \in S_E} D_{min}(p, S_L)$$

where D_{min} represents the shortest distance from S_L to the point p in S_E. The

distance for each point in S_E was calculated, and the average was then determined.

The S_L that minimized D was chosen as the most similar skeleton corresponding to S_E.

This distance-based scoring system ranked similarity (lower scores indicated closer

matches), with the top-ranked matches (swc filenames and scores) exported. These

results were manually verified using a custom software Brain Atlas Viewer (BrAVe)[13].

3D visualization

In accordance with open-science principles, we employed the Neuroglancer

precomputed format and CloudVolume interface for data management and 3D

visualization through both Neuroglancer and webKnossos[84]. For data encoding, we

used JPEG compression for ssEM images and compressed_segmentation format for

region silhouettes and segmentation masks. We further implemented the Igneous[118] for

multiprocessed block-wise downsampling, skeletonization and meshing with the

following agglomeration. For skeletonization, we used the Kimimaro implementation

of the automated TEASAR algorithm[119]. We then merged the skeleton splits that

might appear based on nearest Euclidean neighbors, calculated the path length of the

skeleton, and determined the radius of each vertex in the skeleton. For meshing, we

used the automated block-wise Zmesh to generate multi-resolution meshes. These

skeletons and meshes were stored in sharded formats.

We employed mesh representations to visualize brain regions and reconstructions

in Neuroglancer, while utilizing annotation points to demonstrate synaptic locations.

The Neuroglancer *matrix* function served as an affine transformation tool for

manipulating (translate/flip/rotate/zoom) the original image layers. We used the

parallel projection during Neuroglancer rendering to generate 3D scale bars.

Furthermore, we optimized two existing workflows for video creation: (1)

Neuroglancer workflow using Python scripts from the *video_tool* package, and (2) the

open-source computer graphics software Blender, with Python scripts and JSON

descriptions from neuVid. Since we acquired high-resolution images only for the

retina, optic nerve, brain and aSC, we processed a pre-acquired low-resolution dataset

($400 \times 400 \times \sim 1,650$ nm^3 per voxel) of the complete zebrafish head to provide

comprehensive 3D rendering context used in Fig.1b and Supplementary Video 1. We

first verified the section number corresponding to the whole-brain high-resolution

dataset, and then generated uniformly spaced, downsampled images ($992 \times 992 \times 660$

nm^3 per voxel) along the rostral-caudal axis using linear interpolation. These images

were subsequently registered to the high-resolution dataset, ensuring alignment with multiple segmentation and reconstruction results within the same physical space. Finally, grayscale histogram equalization was performed. The ssEM dataset are accessible through the following dedicated links:

https://ng.zebrafish.digital-brain.cn/Fish-X/Brain_EM_images

https://ng.zebrafish.digital-brain.cn/Fish-X/Retina_EM_images

https://ng.zebrafish.digital-brain.cn/Fish-X/Brain_regions

https://ng.zebrafish.digital-brain.cn/Fish-X/Auto_supervoxels

https://ng.zebrafish.digital-brain.cn/Fish-X/Manual_supervoxels

https://ng.zebrafish.digital-brain.cn/Fish-X/Nuclei

https://ng.zebrafish.digital-brain.cn/Fish-X/Vessel

Statistical analysis

No randomization and blinding were used during data collection and analysis. The statistical analysis was performed using GraphPad Prism. Error bars represent mean ± SEM. Kolmogorov-Smirnov test was used for testing normality. Kruskal-Wallis test was used in Fig. 1k (left panel, PSD/SVC/Mito), 1l (left panel), 2j (among LC-NE, PT-DA, and RN-5HT). Unpaired t test was used in Fig. 2j (between LC-NE/PT-DA/RN-5HT and all other types). Two-tailed Mann Whitney U test was used in Figs. 1k (right panel, PSD/SVC/Mito), 1l (right panel), 2f,g, 3d-h, and Extended Data Fig. 10d,h. Two-tailed two-sample Kolmogorov-Smirnov test was used in Figs. 3l,m, 4l. Linear regression analysis was used in Figs. 1j, 2l, 3g,h, and Extended Data Fig. 4h-j. Exponential fitting was performed in Fig. 4d. Chi-square goodness-of-fit test was performed in Fig. 4e. Statistical significance is defined as $P <$ 0.05. The specific statistical tests, P-value ranges, and sample sizes are provided in the corresponding figures and/or figure legends.

Reporting summary

Further information on research design is available in the Nature Port-folio Reporting Summary linked to this article.

Data availability

All raw EM images are openly accessible for visualization at https://ng.zebrafish.digital-brain.cn/Fish-X/Brain_EM_images; additional image data and reconstructed results can also be explored as detailed in the Methods. Further information and requests for materials and additional details will be fulfilled by the corresponding author.

Code availability

All original codes and algorithms are available at https://github.com/MiRA-Han-Lab/Fish-X.

References (Methods)

90. McLean, D. L., Fan, J., Higashijima, S., Hale, M. E. & Fetcho, J. R. A topographic map of recruitment in spinal cord. *Nature* **446**, 71-75 (2007).

91. Quirin, S. et al. Calcium imaging of neural circuits with extended depth-of-field light-sheet microscopy. *Opt. Lett.* **41**, 855-858 (2016).

92. Muto, A. et al. Activation of the hypothalamic feeding centre upon visual prey detection. *Nat. Commun.* **8**, 15029 (2017).

93. Li, J. et al. Intron targeting-mediated and endogenous gene integrity-maintaining knockin in zebrafish using the CRISPR/Cas9 system. *Cell Res.* **25**, 634-637 (2015).

94. Shang, C.-F. et al. Real-time analysis of large-scale neuronal imaging enables closed-loop investigation of neural dynamics. *Nat. Neurosci.* **27**, 1014-1018 (2024).

95. Kawakami, K. Tol2: a versatile gene transfer vector in vertebrates. *Genome Biol.* **8**, S7 (2007).

96. Ariotti, N. et al. Modular Detection of GFP-labeled proteins for rapid screening by electron microscopy in cells and organisms. *Dev. Cell.* **35**, 513-525 (2015).

97. Martell, J. D., Deerinck, T. J., Lam, S. S., Ellisman, M. H. & Ting, A. Y. Ting, Electron microscopy using the genetically encoded APEX2 tag in cultured mammalian cells. *Nat. Protoc.* **12**, 1792-1816 (2017).

98. Willingham, M. C. & Rutherford, A. V. The use of osmium-thiocarbohydrazide-osmium (OTO) and ferrocyanide-reduced osmium methods to enhance membrane contrast and preservation in cultured cells. *J. Histochem. Cytochem.* **32**, 455-460 (1984).

99. Hua, Y.-F., Laserstein, P. & Helmstaedter, M. Large-volume en-bloc staining for electron microscopy-based connectomics. *Nat. Commun.* **6**, 7923 (2015).

100. Xin, T., Shen, L.-J., Li, L.-L., Chen, X. & Han, H. Expected affine: A registration method for damaged section in serial sections electron microscopy. *Front. Neuroinform.* **16**, 944050 (2022).

101. Lowe, D. G. Distinctive image features from scale-invariant keypoints. *Int. J. Comput. Vis.* **60**, 91-110 (2004).

102. Fischler, M. A. & Bolles, R. C. Random sample consensus: a paradigm for model fitting with applications to image analysis and automated cartography. *Commun. ACM.* **24**, 381-395 (1981).

103. Saalfeld, S., Fetter, R., Cardona, A. & Tomancak, P. Elastic volume reconstruction from series of ultra-thin microscopy sections. *Nat. Methods* **9**, 717-720 (2012).

104. Duchon, J. Splines minimizing rotation-invariant semi-norms in Sobolev spaces. In *Constructive Theory of Functions of Several Variables* (eds Schempp, W. & Zeller, K.) **571**, 85-100 (Springer, Berlin, Heidelberg, 1977).

105. Besl, P. J. & McKay, N. D. A method for registration of 3-D shapes. *IEEE Trans on Pattern Analysis and Machine Intelligence* **14**, 239-256 (1992).

106. Wang, Z.-J., Liu, J.-Z., Li, G.-Q. & Han, H. Blind2unblind: Self-supervised image denoising with visible blind spots. In *Proc. IEEE/CVF Conference on Computer Vision and Pattern Recognition (CVPR)* 2027-2036 (2022).

107. Januszewski, M. et al. High-precision automated reconstruction of neurons with flood-filling networks. *Nat. Methods* **15**, 605-610 (2018).

108. Beier, T. et al. Multicut brings automated neurite segmentation closer to human performance. *Nat. Methods* **14**, 101-102 (2017).

109. Hong, B. et al. Graph partitioning algorithms with biological connectivity decisions for neuron reconstruction in electron microscope volumes. *Expert Syst. With Applic.* **222**, 119776 (2023).

110. Keuper, M. et al. Efficient decomposition of image and mesh graphs by lifted multicuts. In *Proc. IEEE/CVF International Conference on Computer Vision (ICCV)* 1751-1759 (2015).

111. Isensee, F., Jaeger, P. F., Kohl, S. A., Petersen, J. & Maier-Hein, K. H. nnU-Net: a self-configuring method for deep learning-based biomedical image segmentation. *Nat. Methods* **18**, 203-211 (2021).

112. Chen, X.-L. & He, K.-M. Exploring simple siamese representation learning. In *Proc. IEEE/CVF Conference on Computer Vision and Pattern Recognition (CVPR)* 15750-15758 (2021).

113. McInnes, L., Healy, J. & Melville, J. UMAP: uniform manifold approximation and projection for dimension reduction. Preprint at https://arxiv.org/abs/1802.03426 (2018).

114. Kirillov, A. et al. Segment anything. In *Proc. IEEE/CVF International Conference on Computer Vision (ICCV)* 4015-4026 (2023).

115. Zhai, H., Guo, J.-Y., Zhang, Y.-C., Liu, J. & Han, H. SAvEM 3: Pretrained and distilled models for general-purpose 3D neuron reconstruction. In *2024 IEEE International Conference on Bioinformatics and Biomedicine (BIBM)* 3972-3977 (2024).

116. Ravi, N. et al. Sam 2: Segment anything in images and videos. In *13th International Conference on Learning Representations (ICLR)* (2025).

117. Januszewski, M. & Jain, V. Next-generation AI for connectomics. *Nat. Methods* **21**, 1398-1399 (2024).

118. Silversmith, W. et al. Igneous: Distributed dense 3D segmentation meshing, neuron skeletonization, and hierarchical downsampling. *Front. Neural Circuits* **16**, 977700 (2022).

119. Sato, M., Bitter, I., Bender, M. A., Kaufman, A. E. & Nakajima, M. TEASAR: tree-structure extraction algorithm for accurate and robust skeletons. In *Proc. 8th Pacific Conference on Computer Graphics and Applications* 281-449 (2000).

Acknowledgements

We are grateful to Drs. Yu-Long Li and Rong-Bo Sun for sharing PKU constructs, Jie He, Misha Ahrens, Koichi Kawakami, and Joe Fetcho for sharing fish lines, and Yun-Feng Hua, Jun Zhang, Xiao-Hong Xu, Ke Zhang, Yu Mu, Dan-Qian Liu, Kai Yang, and Sheng-Xiong Wang for insightful advice. We would like to thank Xu-Long

Wen, and other staffs from Grand View (Beijing) Technology Co., Ltd, and Zhi Zeng,
Le Wang, and other staffs from Chengdu Huizhong Tianzhi Tech. Co., Ltd.
(www.hztzai.com) for proofreading. We would like to thank Kui Wang for
contribution to plasmid construction, Yu Kong, Li-Jun Pan, Xu Wang, Xue Xu from
Electron Microscopy Core Facility (CEBSIT) for assistance in sample preparation,
Ming-Quan Chen and Hong-Li Wan from Facility of Mapping Brain-Wide Mesoscale
Connectome (FMBWMC, CEBSIT) for assistance in utilizing the zebrafish
mesoscopic atlas, Ting-Ting Zhao and Yong-Wei Zhong from FMBWMC for quality
assessment of automated segmentation results, Jia-Wen Huang, Liu-Qin Qian,
Chen-Xi Jin, and other staffs from Big Data of Brain Atlas Facility (CEBSIT) for
establishing a web-sharing platform, Qian Hu from Optical Imaging Core Facility
(CEBSIT) for optical imaging and image processing support, Jia Li for suggestion on
fish line generation and cell type identification, Chun-Feng Shang, Rong-Wei Zhang,
Yu-Fan Wang, Sha Li, Le Sun, Qiu-Sui Deng, Li-Jun Chen for discussions, Ji-Wen Bu
and Ye Hua for lab management, Yan-Hua Zhu from Shanghai Jiao Tong University
for supporting X-ray tomography. We thank our institute (CEBSIT) for providing
access to GeminiSEM 300 (Zeiss, Germany), Talos L120C TEM (Thermo Scientific,
USA), JEM-1230 TEM (JEOL, Japan), FV3000 confocal laser scanning microscope
(Olympus, Japan), and high-performance computing capabilities at the Center for
Data and Computing in Brain Science. We thank the Transdisciplinary Platform of
Functional Connectome and Brain-Inspired Intelligence in Huairou Science City in
Beijing and the Microscopic Technology & Analysis Center at the Institute of
Automation, Chinese Academy of Sciences for providing technical support and device
resources. This work was supported by Brain Science and Brain-like Intelligence
Technology - National Science and Technology Major Project (2021ZD0204500,
2021ZD0204502 and 2021ZD0204503), Creative Research Groups (32321003) and
general grant (32171461) of the National Natural Science Foundation of China,
Shanghai Municipal Science and Technology Major Project (18JC1410100,
2018SHZDZX05, 25511102500), and Key Research Program of Frontier Sciences
(QYZDYSSW-SMC028) and Strategic Priority Research Program (XDB32010200)
of Chinese Academy of Sciences.

1460

Author contributions

Conceptualization and Supervision: JLD, HH, FNL
Methodology - fish line: FNL, CS, RKT, HYH
Methodology - EM sample preparation: FNL, CS
Methodology - sectioning: LLL, XHD
Methodology - imaging: LNZ, HTM, LML
Methodology - registration: YNL, XC, FXZ, TX, BHC, HRC, SC

Methodology - automated reconstruction: JZL, JL, JYG, QWX

Methodology - proofreading: FNL, CS, ZHF, YSW, SRJ

Methodology - software: JBY, LJS, XHZ

Methodology - visualization: HZ

Methodology - mesoscopic atlas: XFD

Methodology - others: YCZ, HYY, JLZ, CYQ

Analysis - LC-NE system: FNL, CS, CMZ, PSX, YXY

Analysis - others: FNL, JZL, CS, JBY, JL, YNL, HZ, YCZ, YQ, YCG

Writing - original draft (methods): JZL, CS, JBY, JL, YNL, YCZ, HYY, HZ, LLL,

LNZ, XHD

Writing - original draft (all sections): FNL, JLD

Writing - review & editing: FNL, JLD

1480

1481 **Competing interests**

1482 The authors declare no conflict of interest.

1483

Figure legends

Fig. 1 | Multiplexed neuromodulatory-type-labeled whole-brain ssEM of larval zebrafish.

a-c, Neuron-type annotation via multiplexed subcellular peroxidase labeling. (**a**) Schematic of APEX2-based multiplexed EM labeling strategy. Assuming three adjacent neurons of distinct types (top) have APEX2 targeted to different subcellular compartments via specific signal peptides and promoters (middle). Following the DAB reaction, APEX2 catalyzes the conversion of DAB into osmiophilic substances and generates localized non-transmembrane electron-dense staining (bottom), which enables the identification of neuron types in the EM dataset. (**b**) Nine distinct labeling patterns were generated by localizing APEX2 (red) to different compartments (six leftmost images), including nucleus, cytosol, plasma membrane, mitochondria, Golgi apparatus, and presynaptic vesicle. Fusion of APEX2 with PKU tags yielded additional fibrous or spherical self-assembled aggregates in the nucleus and cytosol (three rightmost images). Labeling patterns are indicated by red stars or arrows. (**c**) *Gt(vmat2:GAL4FF);Tg(dbh:GAL4FF;UAS:cytosol-APEX2-TagRFP);Tg(hcrt:APEX2-TagRFP-NLS);Tg(glyt2:MITO-APEX2-TagRFP)*, a quadruple transgenic fish line, was established to enable the collection of the Fish-X dataset. Combined with soma location, cytosol, nucleus, and mitochondria labeling patterns were used to identify monoaminergic, Hcrt, and Gly neurons, respectively. The distinct neuron types are listed on the right.

d, Overview of the multiplexed neuromodulatory-type-annotated whole-brain ssEM dataset ("Fish-X") of a 6-dpf zebrafish larva, comprising three partially overlapping volumes: the brain and aSC, the right retina, and the right optic nerve. This panel shows the rendering of the intact head (left; $992 \times 992 \times 660$ nm^3 per voxel) and representative EM images of the aSC (bottom middle; 32×32 nm^2 per pixel), right optic nerve (top right; 8×8 nm^2 per pixel resolution), and right retina (bottom right; 32×32 nm^2 per pixel).

e-i, Multidimensional automated reconstruction or manually proofread results. (**e**) Rendering of 46,536 automatically reconstructed supervoxels in the brain and aSC (dorsal view), with length ranging between $100 - 300$ μm. (**f**) Rendering of randomly color-coded nuclei distributed in a representative horizontal plane at a depth of 108 μm from the dorsal epidermal surface. (**g**) Rendering of the partially reconstructed morphologies for 523 neurons following manual verification. These include 13 NE neurons (orange) in the LC ($n = 10$) and MO ($n = 3$), 10 DA neurons (light green) in the OB ($n = 3$), PT ($n = 4$) and Hc ($n = 3$), 7 5HT neurons (dark blue) in the Pr ($n = 3$) and RN ($n = 4$), 2 Hcrt neurons (magenta) in the hypothalamus (magenta), 24 Gly neurons (sky blue) in the hindbrain, 178 Glu neurons (purple), 38 GABA neurons

(green), 3 RGCs (red), 1 M-cell (dark gray), and 256 type-unidentified neurons (light

gray; Among them, 9 were PNs to a single OB-DA neuron). (**h**) A total of 52 brain

regions and 7 retinal regions were annotated using the CLEM approach, followed by

manual verification. (**i**) Rendering of brain-wide neural tracts (yellow) and vessels

(red). Most vessels remained well-preserved, though some collapse was observed in

the telencephalon. Neural tracts, identified as a bundle of parallel-oriented fibers,

were manually traced. Vessels were automatically reconstructed, followed by manual

proofreading.

j-l, Quantification of nucleus and synapse distribution across brain regions. (**j**) Linear

regression analysis of synapse density versus nucleus density within each of the

forebrain (red), midbrain (yellow), and hindbrain (green) regions. Dashed lines

represent the fitted regression lines, respectively (Forebrain: $Y = -158.1 \times X + 1{,}373$,

$R^2 = 0.6239$, $P < 0.001$; Midbrain: $Y = -221.1 \times X + 2{,}037$, $R^2 = 0.8959$, $P < 0.01$;

Hindbrain: $Y = -165.0 \times X + 1{,}513$, $R^2 = 0.7196$, $P < 0.0001$). Each circle represents a

single region. For analysis, a synapse was defined by the co-occurrence of the PSD

and SVC. The observed excess of putative synapses (23.62 million) over SVCs (15.77

million) arises because multiple PSDs often associate with a single presynaptic SVC.

The normalized density (**k**) and volume in log10 (**l**) of PSD, SVC, and Mito were

analyzed across anatomical and functional groupings. For each structure, the density

in a given region was normalized to the highest density of that structure across regions.

The volume was log10-transformed for analysis. Brain regions were categorized by

anatomy (forebrain, red; midbrain, yellow; hindbrain, green) and function (sensory,

blue; motor, magenta). The dashed line in each radar plot represents the mean value

across all corresponding structures for each group. Panel i in (**k**): P (PSD) = 0.5495, P

(SVC) = 0.6426, P (Mito) = 0.2297, Kruskal-Wallis test; Panel ii in (**k**): P (PSD) =

0.7695, P (SVC) = 0.8547, P (Mito) = 0.5953, Mann Whitney test. Panel i in (**l**): P

(PSD) = 0.1638, P (SVC) < 0.05, P (Mito) = 0.5717, Kruskal-Wallis test with Dunn's

multiple comparison test; Panel ii in (**l**): P (PSD) = 0.1379, P (SVC) < 0.05, P (Mito)

= 0.4151, Mann Whitney test. C, caudal; L, left; V, ventral; LC, locus coeruleus; MO,

medulla oblongata; OB, olfactory bulb; PT, posterior tuberculum; Hc, caudal

hypothalamus; Pr, pretectum; RN, raphe nucleus; NL, neuropil layers (optic tectum);

NE, noradrenergic; DA, dopaminergic; 5HT, serotonergic; Hcrt, hypocretinergic; Gly,

glycinergic; Glu, glutamatergic; GABA, GABAergic; RGC, retinal ganglion cell;

M-cell, Mauthner cell; PSD, postsynaptic density; SVC, synaptic vesicle cloud; Mito,

mitochondria.

Scale bars, 0.5 μm (**b**), 1 μm (**d**, optic nerve), 20 μm (**d**, spinal cord and retina), and

50 μm (**e-i**).

1559

1560 **Fig. 2 | Reconstruction of perisomatic ultrastructures and dendritic synaptic**

inputs of multiple neuromodulatory neurons.

a-e, Representative EM images (4×4 nm^2 per pixel) of seven molecularly identified neuron types with APEX2 labeling (NE, orange; DA, green; 5HT, blue; Hcrt, magenta; Gly, cyan). Individual APEX2-labeled neurons display various degrees of staining intensity and are contoured by color-coded dashed lines. Renderings of all APEX2-labeled nuclei are shown on the left, with brain regions outlined by dashed squares. All types except Gly were manually verified. Gly neurons in hindbrain were identified automatically based on the presence of ≥ 4 labeled mitochondria in the soma. White arrows point to two APEX2-labeled mitochondria in a Gly neuron, with one adjacent to the nuclear envelope infoldings (**e**).

f,g, The mean nucleus/soma volume ratio (**f**) and somatic mitochondrial density (**g**) were analyzed across neuron types. (**f**) LC-NE *vs* PT-DA, $P = 0.1255$; LC-NE *vs* all other types except PT-DA, $P < 0.05$. (**g**) LC-NE *vs* all other types, $P < 0.05$. The Mann Whitney test was used.

h,i, Rendering of EM reconstructions for APEX2-labeled neuromodulatory and Gly neurons in the right hemisphere, shown in dorsal (**h**) and lateral (**i**) views. Gly neurons with APEX2 labeling (cyan), identified as LC-NE presynaptic neurons during reconstruction, are also shown. Presynaptic input sites are marked as small gray dots. The dendritic arbors were fully reconstructed for neuromodulatory types. Neuron and synapse numbers are shown in parentheses. Input mapping was not performed for five LC-NE neurons on the left sides.

j,k, Dendritic input density (**j**) and number (**k**) across neuron types. Dendritic input density (per μm): LC-NE, 0.3493 ± 0.0194; MO-NE, 0.1511 ± 0.0185; OB-DA, 0.0255 ± 0.0094; PT-DA, 0.3420 ± 0.0212; Hc-DA, 0.0415 ± 0.0249; Pr-5HT, 0.0759 ± 0.0249; RN-5HT, 0.3749 ± 0.0413; Hcrt, 0.0747 ± 0.0204; Gly, 0.1228 ± 0.0190. P (LC *vs* MO) < 0.05 (Mann Whitney test); P (among LC, PT, and RN) $= 0.9533$ (Kruskal-Wallis test); P (LC/PT/RN *vs* all other types) < 0.0001 (unpaired t test). Dendritic input number: LC-NE, 430.2 ± 55.48; MO-NE, 56.67 ± 13.35; OB-DA, 13.67 ± 3.180; PT-DA, 321.8 ± 31.89; Hc-DA, 20.67 ± 15.64; Pr-5HT, 44.00 ± 16.52; RN-5HT, 218.0 ± 43.18; Hcrt, 68.00 ± 25.00; Gly, 57.47 ± 13.74. P (among all neuron types) < 0.01 (Kruskal-Wallis test).

l, Linear regression analysis of the total projection length versus the dendritic input number ($Y = 19.82 \times X + 884.4$, $R^2 = 0.6275$, $P < 0.05$). The total projection length was calculated as the mean of three individual neurons from mesoscopic datasets for each neuron type. The dendritic input number were calculated from the same neurons in h to l. Ro, rostral; C, caudal; L, left; Ri, right; V, ventral. Error bars represent mean $\pm$ SEM.

Scale bars, 5 μm (**a-e**), 10 μm (**i**), and 50 μm (**h**).

Fig. 3 | Spatial organization of broad-yet-sparse synaptic inputs of individual LC-NE neurons.

a, Rendering of synaptic input sites on the dendrites of the LC-NE neuron #3, color-coded by sensory/motor modality. Synapses that were traced but not yet reached their soma are marked as gray dots. Dendritic and axonal processes are color-coded as orange and gray, respectively. The number in parentheses represent the number of synapses.

b, 2D spatial distribution of synaptic input sites on the dendrites of LC-NE neuron #3. In zebrafish, LC-NE neurons are pseudounipolar neurons, with both axonal and dendritic processes emerging from a primary branch. The distance between each dot and the soma represents the path length of each input.

c, Rendering of all 374 EM-reconstructed presynaptic neurons (PNs), which form 418 synapses on the LC-NE neuron #3 (orange). All presynapses for the LC-NE neuron #3 were traced and only those successfully traced back to their somata were used for analysis. Based on the soma locations of corresponding PNs, synaptic input sites were empirically classified as sensory-/sensory relay-related (blue; abbreviated as "sensory") or motor-related (magenta). The same color scheme is applied to all related data below. The resulting counts are as follows: Sensory-related, 160 neurons and 182 synapses; Motor-related, 214 neurons and 236 synapses.

d, The mean number of synapses formed by PNs in each of regions. Input number per PN: Sensory, 1.204 ± 0.08268; Motor, 1.102 ± 0.04288. P (sensory vs motor) = 0.5192 (Mann-Whitney test). Region name and PN number are shown on the left. The R1 and R2 rank as the two predominant input sources, accounting for 20.86% (78/374) and 10.16% (38/374) of total PNs, respectively.

e,f, Differential probability distributions of sensory-/motor-related synaptic inputs along dendritic path length, from the soma to distal region. Separate analyses of neuron #3 alone (**e**) and the pooled data from the other four neurons (**f**: #1, #2, #4, and #5; Sensory-related synapse, $n = 104$; Motor-related synapse, $n = 159$) both revealed that motor-related inputs were distributed more proximally than sensory-related inputs. The mean distance to the soma are indicated by dashed lines. ****$P < 0.0001$ (Mann Whitney test).

g,h, Linear regression analyses of dendritic path length and the corresponding synaptic strength quantified by PSD volume (**g**) and SV volume (**h**). Sensory-related (blue): PSD volume vs path length (**g**, left), Y = -1.181e4 × X + 4.927e6, $R^2 = 0.0627$, $P < 0.001$; SV volume vs path length (**h**, left), Y = -3.283e4 × X + 9.356e6, $R^2 = 0.0711$, $P < 0.001$. Motor-related (magenta): PSD volume vs path length (**g**, middle), Y = -2.174e4 × X + 5.7e6, $R^2 = 0.1369$, $P < 0.0001$; SV volume vs path length (**h**, middle), Y = -3.611e4 × X + 8.414e6, $R^2 = 0.0790$, $P < 0.0001$. Right: probability distributions of PSD (**g**) and SV (**h**) volumes for both sensory-related (blue) and

motor-related (magenta) inputs, with dashed lines indicating the mean volumes (Mann Whitney test).

i, Representative examples showing partially EM-reconstructed query neurons (black) with their fully LM-reconstructed top-ranked E/I counterparts (Glu, magenta; GABA, green), revealed by morphology comparison. The index and soma location of the EM query were listed below.

j, E/I composition ratio of PNs innervating the LC-NE neuron #3. The E/I type was determined for about half of sensory-related ($n = 82$) and motor-related ($n = 106$) PNs. Glu (purple, $n = 135$) and GABA (green, $n = 33$) neurons were inferred through morphology comparison, and Gly (blue) neurons were verified via mitochondrial APEX2 labeling ($n = 17$) or inferred through morphology comparison ($n = 3$). PNs with type unknown (UN, $n = 186$) are displayed as gray.

k, Rendering of all 374 PNs of the LC-NE neuron #3, color-coded by sensory/motor modality (LC-NE, orange; Glu, purple; GABA, green; Gly, blue; UN, gray). Inset (right): highlighted somatodendritic area with synaptic input sites.

l,m, Distribution of E (left) and I (right) input sites on the dendritic area of the LC-NE neuron #3, for sensory-related (**l**) and motor-related (**m**) inputs. Orange circles represent the soma. The values are derived from the number of synapses (shown in parentheses) formed by PNs identified as E or I in (**j**) and (**k**). Sensory-related I inputs (**l**, right) and motor-related E inputs (**m**, left) were more clustered in distal and proximal areas, respectively. Left-right axis: P(sensory-E vs sensory-I) < 0.05, P(motor-E vs motor-I) $= 0.1072$, P(sensory-E vs motor-E) < 0.001, P(sensory-I vs motor-I) < 0.05, P(sensory-E vs motor-I) $= 0.6938$, P(sensory-I vs motor-E) < 0.0001. Dorsal-ventral axis, P(sensory-E vs sensory-I) $= 0.2084$, P(motor-E vs motor-I) $= 0.1809$, P(sensory-E vs motor-E) $= 0.0530$, P(sensory-I vs motor-I) $= 0.2053$, P(sensory-E vs motor-I) $= 0.6938$, P(sensory-I vs motor-E) < 0.05. The two-sample Kolmogorov-Smirnov test was used. Ro, rostral; C, caudal; L, left; Ri, right; V, ventral.

Scale bars, 10 μm (**a**), 20 μm (**l,m**), 50 μm (**k**), and 0.5×10^{-7} (**g,h**).

Fig. 4 | Intra- and inter-system common synaptic inputs among monoaminergic systems.

a, Rendering of all 69 EM-reconstructed intra-LC system common presynaptic neurons ("intra-CPNs"), which formed 88 synapses on the LC-NE neuron #3. Sensory-related (blue), 30 neurons and 39 synapses; Motor-related (magenta), 39 neurons and 49 synapses.

b, Regional distribution of 69 intra-CPNs. Neuron numbers are shown at the right of each bar.

c, E/I composition ratios for intra-CPNs with various innervation numbers.

d, Cell ratio of intra-CPNs with various innervation numbers. PN numbers are shown in parentheses. Exponential fitting: $Y = 0.0255 + 28.09 \times e^{-1.816 \times X}$, $R^2 = 0.9973$.

e, Cell ratio of intra-CPNs categorized by innervation combination and target LC-NE identity (inset). Neuron numbers are shown in parentheses. All intra-CPNs analyzed innervated the LC-NE neuron #3. The red symbols represent theoretical cell ratios obtained from 100 iterations of random combinations, using identical cell numbers and matched innervation number distributions. Chi-square goodness-of-fit test was performed (E, $\chi^2(3) = 7.8727$, $P < 0.05$). Error bars represent mean ± SEM.

f, Rendering of all 26 EM-reconstructed inter-monoaminergic system common presynaptic neurons ("inter-CPNs"), which formed 32 synapses on the LC-NE neuron #3. Sensory-related (blue), 12 neurons and 17 synapses; Motor-related (magenta), 14 neurons and 15 synapses.

g, Regional distribution of 26 inter-CPNs. Neuron numbers are shown at the right of each bar.

h, E/I composition ratios for inter-CPNs with different monoaminergic-type combinations.

i,j, Analysis of synchrony in spontaneous activities among three monoaminergic neuron types. Heatmap in (**i**) depicts spontaneous calcium activities (Z-scored) in multiple monoaminergic neurons, and the cross-correlation curves between each pair are shown in (**j**). *Ki(th:GAL4-VP16);Ki(tph2:GAL4FF,cmlc2:EGFP);Tg(UAS:GCaMP6s)* larvae at 6 dpf were used for *in vivo* confocal calcium imaging. Black arrows indicate two events of spontaneous activities synchronized between LC-NE and PT-DA neurons.

k, Rendering of representative DA (green) and 5HT (blue) neurons with their partially reconstructed PNs (gray). Only synapses successfully traced back to their PN somata are shown. PNs of the OB-DA neuron (top-middle) were obtained by dedicated tracing, while PNs of the PT-DA (left) and RN-5HT (right) neurons were identified common inputs (inter-CPNs) discovered during the reconstruction of the five LC-NE neurons' inputs. Notably, some PNs (red arrows) also innervate multiple PT-DA or RN-5HT neurons, serving as intra-CPNs for these systems. The counts of neurons are shown in parentheses.

l, Distribution of non-CPN (left), intra-CPN (middle), and inter-CPN (right) inputs on the dendrites of the LC-NE neuron #3. Orange circles represent the soma. The number in parentheses indicate the number of synapses analyzed. Both intra-CPN and inter-CPN inputs favor proximal distributions, whereas non-CPN inputs display a more dispersive distribution. Left-right axis: P(non-CPN *vs* intra-CPN) < 0.0001, P(non-CPN *vs* inter-CPN) = 0.5716, P(intra-CPN *vs* inter-CPN) < 0.05. Dorsal-ventral axis: P(non-CPN *vs* intra-CPN) < 0.001, P(non-CPN *vs* inter-CPN) < 0.01, P(intra-CPN *vs* inter-CPN) = 0.3284. The two-sample Kolmogorov-Smirnov test

1717   was used.

1718   C, caudal; L, left; V, ventral.

1719   Scale bars, 20 µm (**l**) and 50 µm (**k**).

1720

1721   **Fig. 5 | Multi-level spatial organization of broad-yet-sparse synaptic inputs to**

1722   **individual LC-NE neurons.**

1723   Individual LC-NE neurons receive broad-yet-sparse synaptic inputs (**a**), which are

1724   spatially organized by sensory/motor modality, E/I identity, synaptic strength, and

1725   private/shared types (**b**).

1726

Extended Data Figures and Legends

Extended Data Fig. 1 | Versatile APEX2 genetic toolbox for multiplexed labeling.
a, Schematic of genetic constructs for the nine APEX2 labeling patterns. Red asterisks mark the three transgenic lines employed for the production of larvae used in the Fish-X collection. Yellow boxes indicate corresponding signal peptides or PKU tags.
b, Z-projected whole-brain (left) and expanded (right) fluorescent images of the 6-dpf multiplexed APEX2-labeled larva used for whole-brain ssEM in this study. White arrows indicate cytosol- or nucleus-localized APEX2-conjugated TagRFP signals in LC-NE (i), PT-DA (ii), RN-5HT (iii), and Hcrt (iv) neurons. Mitochondria-localized signals are outlined by white dashed lines in hindbrain Gly neurons (v).
c, Specificity of APEX2 labeling in 6-dpf larvae was validated by immunohistochemical staining (left) and crossing with neuronal type-specific reporter lines (right). Representative fluorescent images show APEX2-TagRFP (top), immunofluorescence/reporter GFP (middle), and merged channels (bottom). *Gt(vmat2:GAL4FF);Tg(dbh:GAL4FF;UAS:cytosol-APEX2-TagRFP)* larvae were immunostained with an anti-tyrosine hydroxylase antibody to validate NE/DA neurons, and were crossed with *Ki(tph2:EGFP)* larvae to validate 5HT neurons. *Tg(hcrt:APEX2-TagRFP-NLS)* larvae were immunostained with an anti-hypocretin antibody to validate Hcrt neurons. *Tg(glyt2:MITO-APEX2-TagRFP)* larvae were crossed with *Tg(glyt2:GFP)* larvae to validate Gly neurons. Ro, rostral; C, caudal; L, left; Ri, right. LC, locus coeruleus; MO, medulla oblongata; OB, olfactory bulb; PT, posterior tuberculum; Hc, caudal hypothalamus; Pr, pretectum; RN, raphe nucleus; NE, noradrenergic; DA, dopaminergic; 5HT, serotonergic; Hcrt, hypocretinergic; Gly, glycinergic.
Scale bars, 10 μm (**b**, right; **c**) and 50 μm (**b**, left).

Extended Data Fig. 2 | Integrated workflow for whole-brain ssEM of larval zebrafish.
This workflow outlines the key steps for generating the Fish-X dataset. We began by using *in vivo* confocal fluorescence imaging to assess the coverage of labeled nuclei and the number of labeled neurons in a larva featuring simultaneous labeling of NE, DA, 5HT, Hcrt, and Gly neurons. The sample was then chemically fixed overnight. The following day, it was treated with a DAB-H_2O_2 solution, and the labeling performance was initially evaluated under a bright-field stereoscope. Subsequently, the sample underwent rOTO staining, dehydration, and resinification. We employed X-ray micro-tomography to rapidly and non-destructively verify the sample's integrity, with the resulting signal serving as a second round of labeling assessment. Using the ATUM system, the sample was sectioned along the rostrocaudal axis to create a library of 33-nm sections, which were collected on 49 wafers. Image acquisition was then carried out over 6 months using three SEMs. Following this, the acquired images were stitched and aligned. The resulting whole-brain volume was partitioned into 34.4×34.4×0.99 μm³ blocks for distributed segmentation, aggregation, and merging, culminating in a fully automated 3D reconstruction (see Methods). The magnified rendering of 1,404 automatically reconstructed optic tectal periventricular cells with

1771   incomplete morphologies is shown (bottom right).

1772   Scale bars, 50 μm.

1773

Extended Data Fig. 3 | Brain-wide heterogeneity of nuclear morphology and
density.

a, UMAP visualization of whole-brain nucleus clustering (176,789 cells) using
combined 8-dimensional deep-learned and 13 traditional features (see Methods). All
nuclei were manually proofread after automated segmentation, resulting in optimal
3D morphological reconstruction. 21 nuclei located at dataset boundaries were
excluded from clustering due to incomplete morphologies. Based on APEX2-labeling,
2,689 Gly neurons were automatically identified, and 309 neurons of
neuromodulatory types were manually annotated by experts. The number of
neuromodulatory-type neurons are shown in parentheses. In addition, 511
vasculature-associated cells, including endothelial cells (ECs, $n = 383$), pericytes (PCs,
$n = 84$), and red blood cells (RBCs, $n = 44$), were manually annotated based on their
location and perisomatic features. Inset (left): vasculature-associated cells tending to
aggregate. Among all identified cells, we classified 70,114 as putative neurons based
on the presence of, at least one synapse in their automatically reconstructed
supervoxels. The remaining cells were presumed to be glial cells or neurons with poor
automated process reconstruction.

b, Rendering of nuclei from diverse cell types. Black arrows indicate shallow grooves
and infoldings on the nuclei of LC-NE and Gly neurons. Two examples of
unclassified neurons with irregular nuclear morphologies are also shown.

c, Distribution of nucleus density distribution across 52 brain regions and 7 retinal
regions. Color code corresponds to brain regions as in Fig. 1i.

Scale bars, 3 μm (**b**).

1797

Extended Data Fig. 4 | Density and volume distributions of PSDs, SVCs, and
Mito.

a, A representative EM image (4×4 nm^2 per pixel) of a synapse, with the PSD (blue),
SVC (orange), and a mitochondrion (green) outlined.

b-d, Summary of spatial colocalization among PSDs (**b**), SVCs (**c**), and Mito (**d**).
Automated 3D reconstructed supervoxels were used to identify whether these
structures were co-localized within the same presynaptic compartments.

e-g, Density distributions of PSDs (**e**), SVCs (**f**), and Mito (**g**) across 52 brain regions.

h-j, Pairwise linear regression of densities among PSDs, SVCs, and Mito. The solid
red lines represent the fitted regression line, and the dashed red lines indicate the 95%
confidence interval of the fit. Each circle represents individual brain regions. SVC *vs*
PSD: $Y = 0.6092 \times X + 12.86$, $R^2 = 0.9398$, $P < 0.0001$ (**h**); Mito *vs* SVC: $Y = 0.7542$
$\times X + 100.9$, $R^2 = 0.8454$, $P < 0.0001$ (**i**); Mito *vs* PSD: $Y = 0.4644 \times X + 107.7$, $R^2 =$
0.8118, $P < 0.0001$ (**j**).

k, Lognormal frequency distributions of volumes among PSDs (top), SVCs (middle),
and Mito (bottom) across brain regions. The volume was calculated using the
automated 3D reconstructed supervoxels, and the data from all 52 regions were

pooled together.

l-n, Lognormal probability density distributions of volumes among PSDs (top), SVCs

(middle), and Mito (bottom) from the forebrain (**l**), midbrain (**m**), and hindbrain (**n**).

All regions exhibit comparable lognormal distributions, with Ep (**l**, brown) and LCa

(**n**, cyan) showing slight differences from the majority, as indicated by arrows.

Regions are randomly color-coded. PSD, postsynaptic density; SVC, synaptic vesicle

cloud; Mito, mitochondria.

Scale bars, 500 nm (**a**).

1823

Extended Data Fig. 5 | Single-cell morphology of diverse neuromodulatory

neurons reconstructed by LM.

a, Workflow for reconstructing single-cell morphologies by LM, adapted from Du et

al[45]. Fluorescent protein-expressing plasmids were injected into fertilized eggs

derived from neuromodulatory-type-specific Gal4 driver lines. This approach

simultaneously enabled sparse labeling for morphology reconstruction and provided

pan-neuronal fluorescence for image registration. At 3 dpf, larvae were screened by

fluorescence microscopy to identify those with single-cell labeling. Selected larvae

were then subjected to *in vivo* confocal imaging at 6 dpf. Neuronal morphologies were

reconstructed through manual skeleton tracing based on the acquired fluorescent

images. In parallel, the confocal images were registered to a mesoscopic brain

template, and the resulting transformation matrices were applied to align the

reconstructed skeletons with the standard reference space.

b-I, Representative mesoscopic single-cell morphologies of diverse neuromodulatory

types, shown in lateral (left), dorsal (middle), and detailed dorsal (right) views.

Examples include LC-NE (**b**), MO-NE (**c**), OB-DA (**d**), PT-DA (**e**), Hc-DA (**f**),

Pr-5HT (**g**), RN-5HT (**h**), and Hcrt (**i**) neurons. Black arrows indicate somata. The

predominantly innervated neuropil (highlighted in light blue) is shown in parentheses.

Ro, rostral; L, left; D, dorsal; L, left.

1843

Extended Data Fig. 6 | Neuron type identification based on soma location and

APEX2 labeling patterns.

a-i, Corresponding 33-nm EM sections (64×64 nm^2 per pixel) containing the

representative APEX2-labeled neurons (marked by asterisks) from Fig. 2a-e. All other

APEX2-labeled neurons within the same sections are also outlined with color-coded

circles (NE, orange; DA, green; 5HT, blue; Hcrt, magenta; Gly, cyan). Section indices

are listed below each image. The sections are arranged sequentially from left to right

and top to bottom, following the rostral-to-caudal axis.

j, The schematic indicates the exact rostral-caudal positions of these sections, with

corresponding section indices and neuronal types listed to the right. V, ventral; L, left.

Scale bars, 50 μm (**a-i**).

1855

Extended Data Fig. 7 | Comparison analysis of perisomatic features between

neuromodulatory neurons and their adjacent unlabeled cells.

a, Rendering of a single LC-NE neuron's soma, containing endoplasmic reticulum

(ER, orange), Golgi apparatus (GA, pink), mitochondria (Mito, blue), nucleus (green),
and nucleolus (red). The red arrow indicates a primary cilia. For clarity, Mito (top
right) and ER (bottom right) are additionally shown in separate renderings.
b, Comparison of perisomatic features between APEX2-labeled LC-NE neurons
(orange, $n = 5$) and their unlabeled neighbors (gray, $n = 69$). Labeled LC-NE neurons
exhibited a lower nucleus-to-soma volume ratio and a higher density of somatic
mitochondria compared to the randomly-selected neighboring cells (left). Based on a
combination of deep-learned and traditional morphological features of the nucleus and
soma, UMAP visualization revealed a consistent structural distinction between the
two groups (right, inset). The blue arrow points to an unlabeled cell that clustered
with LC-NE neurons and displayed similar properties to those of LC-NE neurons,
suggesting it may be an unlabeled member of the same type.
c, Rendering of the five APEX2-labeled (orange) and one unlabeled (blue) LC-NE
neurons shown in **b**. Morphological reconstruction revealed similar axonal projection
patterns between the LC-NE neuron #3 and unlabeled LC-NE neuron.
d-k, Comparisons of perisomatic features between APEX2-labeled neurons and a
10-fold number of adjacent unlabeled cells. Left, nucleus-to-soma volume ratio *vs*
somatic mitochondrial density. Right, UMAP visualization of cell clustering based on
combined perisomatic morphological features. A similar structural divergence was
observed for PT-DA neurons (red outline and asterisk, **f**), when compared to their
neighbors. Cell numbers are shown in parentheses. Ro, rostral; V, ventral.
Scale bars, 2 μm (**a**) and 20 μm (**c**).

1881

Extended Data Fig. 8 | Somatodendritic reconstruction and synaptic input
mapping of a Mauthner cell.
a, Rendering of a right M-cell's soma, lateral dendrites, ventral dendrites, and
proximal axon. The axonal branch is not fully displayed. Small violet dots indicate
synaptic input sites ($n = 1,966$). Both the morphology and synapses were manually
verified based on automated segmentation results.
b,c, Representative EM images (4×4 nm^2 per pixel) of the M-cell's soma, shown in
coronal (**b**) and detailed coronal (**c**) views. The nucleus and cytosol are labeled in red
and yellow, respectively. A typical region of the ventral cell surface was densely
innervated by multiple presynaptic inputs with random color-coding. L, left; Ri, right;
D, dorsal; V, ventral.
Scale bars, 1 μm (**c**), 5 μm (**b**), and 10 μm (**a**).

1894

Extended Data Fig. 9 | Dendritic architecture and synaptic inputs of the other
four APEX2-labeled LC-NE neurons.
a, 2D spatial distribution of dendritic synaptic inputs on LC-NE neurons (#1, #2, #4,
and #5). Each dot represents a single synaptic input site, color-coded by the modality
of PNs (Sensory-related, blue; Motor-related, magenta). Gray dots indicate synaptic
sites untraced or traced but not yet reached soma (UN, undefined). The distance
between each dot and the soma represents the path length of the corresponding
synaptic input. Spatial distribution of five LC-NE neurons is shown via nuclear

positioning in a dorsal view (bottom left).

b,c, Dendritic heterogeneity across individual LC-NE neurons in both branch number (**b**) and maximum branch order (**c**). Values are listed above each bar.

d, Dendritic synaptic input counts for individual LC-NE neurons, with values listed above each bar. Synapses of known modality are proportionally represented within each bar and are color-coded consistently with A. C, caudal; L, left; D, dorsal; V, ventral.

Scale bars, 20 μm (**a**).

Extended Data Fig. 10 | E/I type of reconstructed neurons can be identified via morphology comparison but not synaptic features.

a, The plasmid *UAS:MITO-APEX2-TagRFP* was injected into fertilized eggs derived from *Tg(vglut2a:GAL4FF)* or *TgBAC(gad1b:GAL4-VP16)* lines to specifically label mitochondria in Glu or GABA neurons, respectively.

b,c Representative EM images of APEX2-labeld mitochondria in axon terminals of Glu (**b**) and GABA (**c**) neurons. PSDs, blue; SVCs, orange; Labeled mitochondria, green.

d, Comparisons of ultrastructural features for PSDs between Glu (magenta, $n = 90$) and GABA (green, $n = 102$) synapses. PSD thickness was estimated from the diameter of its maximum inscribed sphere. Dashed lines indicate the average values of PSD thickness (horizontal) and volume (vertical). P(thickness) $= 0.5365$, P(volume) $= 0.0943$ (Mann Whitney test).

e-g, Representative examples showing partially EM-reconstructed query neurons (black) with their fully LM-reconstructed top-ranked E/I type counterparts (Glu, magenta; GABA, green; Gly, blue), revealed by morphology comparison. Some EM queries were matched to LM neurons with similar morphology (**e,f**), whereas others exhibit substantial morphological differences from their top-ranked LM references (**g**). All EM queries used in morphology comparison were PNs of LC-NE neurons. The soma location is shown in parentheses.

h, Comparison of the normalized top penalty scores for all 467 EM queries, with lower penalty scores indicating higher degrees of matching. The penalty score was normalized by skeleton node number of EM queries. All results were manually verified after morphological matching and categorized into E/I-specific match, E/I-ambiguous match, and Un-match. In some brain regions, E and I neurons exhibit similar morphology, causing good matches between EM queries and both E/I neurons. Cell numbers are shown in parentheses. The Mann Whitney test was used, and the error bars represent mean ± SEM.

i,j, EM query neuron #41 (black) exhibited morphological similarity to adjacent LM-reconstructed Glu, GABA, and Gly neurons (**i**), suggesting ambiguous identity based on structure alone. However, it could be unambiguously identified as Gly by its mitochondrial-APEX2 labeling, as indicated by white arrows (**j**).

k-n, Rendering of EM-reconstructed PNs and corresponding LC-NE neuron #1 (**k**), #2 (**l**), #4 (**m**), and #5 (**n**). The counts of PNs with E/I profiles and their corresponding synaptic outputs onto the target LC-NE neuron are shown in parentheses. Notably,

1947   two Hcrt neurons (magenta) with nuclear-localized APEX2 labeling specifically
1948   targeted the LC-NE neuron #4 (**m**), rather than other LC-NE neurons.
1949   Ro, rostral; V, ventral.
1950   Scale bars, 200 nm (**b,c**), 2 μm (**j**), and 50 μm (**k-n**).
1951

 SUPPLEMENTARY INFORMATION

 SUPPLEMENTARY VIDEOS:

1954 **Supplementary Video 1 |** Fish-X: A multidimensional whole-brain EM
1955 reconstruction of a 6-dpf larva characterized by multiplexed APEX2-based
1956 neuromodulatory type labeling, comprising 529 reconstructed neurons of various
1957 types, 176,810 nuclei, neural tracts, and vessels in the brain and anterior spinal cord.
1958 A
1959 *Gt(vmat2:GAL4FF);Tg(dbh:GAL4FF;UAS:cytosol-APEX2-TagRFP);Tg(hcrt:APEX2*
1960 *-TagRFP-NLS);Tg(glyt2:MITO-APEX2-TagRFP)* larva at 6 dpf was used.
1961

1962 **Supplementary Video 2 |** The whole-brain density distribution for nuclei, PSDs,
1963 SVCs, and Mito. The sequence first shows brain regions emerging along the
1964 rostral-to-caudal axis, then displays the corresponding cellular components in
1965 descending order of density.
1966

1967 **Supplementary Video 3 |** Somatodendritic reconstruction of multiple
1968 neuromodulatory and Gly neuronal populations.
1969

1970 **Supplementary Video 4 |** Dendritic input site mapping of multiple monoaminergic
1971 neuronal populations.
1972

1973 **Supplementary Video 5 |** Mauthner cell reconstruction with dendritic input site
1974 mapping.
1975

1976 **Supplementary Video 6 |** Widely distributed inputs from 374 PNs across 32 brain
1977 regions converging onto the LC-NE neuron #3.
1978

1979 **Supplementary Video 7 |** The spatial organization of synaptic inputs to the LC-NE
1980 neuron #3. Inputs are categorized as follows: 1) modalities (sensory-/motor-related), 2)
1981 E/I types (Glu/GABA/Gly), 3) shared inputs within the LC-NE system (intra-CPNs),
1982 and 4) shared inputs across monoaminergic systems (inter-CPNs).

 SUPPLEMENTARY TABLES

 Supplementary Table 1 | Genetic toolbox for subcellularly-targeted APEX2
 labeling

No.	Full name	Simple name	Targeting	Type	Source
1	*Tg(dbh:GAL4FF;5×UAS-hsp:APEX2-TagRFP-NLS)ion143d*	*Tg(dbh:GAL4FF;UAS:APEX2-TagRFP-NLS)*	Nucleus	NE	Line
2	*Tg(dbh:GAL4FF;5×UAS-hsp:cytosol-APEX2-TagRFP)ion142d*	*Tg(dbh:GAL4FF;UAS:cytosol-APEX2-TagRFP)*	Cytosol	NE	Line
3	*Tg(hcrt:APEX2-TagRFP-NLS)ion100d*	*Tg(hcrt:APEX2-TagRFP-NLS)*	Nucleus	Hcrt	Line
4	*Tg(hcrt:MITO-APEX2-TagRFP)ion101d*	*Tg(hcrt:MITO-APEX2-TagRFP)*	Mitochondria	Hcrt	Line
5	*Ki(slc17a6b:GAL4FF,5×UAS-hsp:sfGFP-APEX2-NLS)ion214d*	*Ki(vglut2a:GAL4FF,UAS:sfGFP-APEX2-NLS)*	Nucleus	Glu	Line
6	*TgBAC(slc17a6:cytosol-APEX2-TagRFP)ion140d*	*TgBAC(vglut2a:cytosol-APEX2-TagRFP)*	Cytosol	Glu	Line
7	*TgBAC(gad1b:Hsa.Cox8a-APEX2-P2A-EGFP)ion170d*	*TgBAC(gad1b:MITO-APEX2-P2A-EGFP)*	Mitochondria	GABA	Line
8	*TgBAC(Gad1b:APEX2-EGFP-CAAX)ion141d*	*TgBAC(gad1b:APEX2-EGFP-CAAX)*	Plasma membrane	GABA	Line
9	*Ki(Gad1b:SEPT6-sfGFP-APEX2)ion202d*	*Ki(gad1b:SEPT6-sfGFP-APEX2)*	Fibrous in cytosol	GABA	Line
10	*Tg(slc6a5:APEX2-TagRFP-NLS)ion99d*	*Tg(glyt2:APEX2-TagRFP-NLS)*	Nucleus	Gly	Line
11	*Tg(slc6a5:Hsa.Cox8a-APEX2-TagRFP)ion97d*	*Tg(glyt2:MITO-APEX2-TagRFP)*	Mitochondria	Gly	Line
12	*Tg(5×UAS-hsp:APEX2-TagRFP-NLS)ion195d*	*Tg(UAS:APEX2-TagRFP-NLS)*	Nucleus	-	Line
13	*Tg(5×UAS-hsp:sypb-APEX2-TagRFP)ion144d*	*Tg(UAS:sypb-APEX2-TagRFP)*	Presynaptic vesicle	-	Line
14	-	*yTol2-dbh:GAL4FF,UAS:SEPT6-sfGFP-APEX2-NLS*	Fibrous in cytosol	DBH	Plasmid
15	-	*yTol2-hcrt:Golgi-APEX2-TagRFP*	Golgi apparatus	Hcrt	Plasmid
16	-	*pMD19-T-gad1b-P2A-tetoff-SEPT6-sfGFP-APEX2-NLS KI donor*	Fibrous in cytosol	GABA	Plasmid
17	-	*yTol2-UAS:MITO-APEX2-TagRFP*	Mitochondria	-	Plasmid

| 18 | - | *yTol2-UAS:cytosol-PRDX4m -sfGFP-APEX2* | Spherical in cytosol | - | Plasmid |

1986   Transgenic zebrafish larvae were in *nacre* background. All transgenic lines and
1987   plasmids were first published in this paper.
1988

Supplementary Table 2 | Signal peptide sequences for subcellular targeting and self-assembly of APEX2

No.	Targeting/Morphology (abbr.)	Sequence
1	Nucleus (*NLS*)	ccaaaaaagaagagaaaggtagatccaaaaaagaagagaaaggtagatccaaaaa agaagagaaaggta
2	Cytosol (*cytosol*)	-
3	Plasma membrane	aagctgaaccctcctgatgagagtggccccggctgcatgagctgcaagtgtgtgctc tccttga
4	Mitochondria (*MITO*)	atgtccgtcctgacgccgctgctgctgcgggggcttgacaggctcggcccggcggct cccagtgccgcgcgccaagatccattcgttgggggatctgtccgtcctgacgccgct gctgctgcggggcttgacaggctcggcccggcggctcccagtgccgcgcgccaa gatccattcgttgggggatcc
5	Golgi apparatus (*Golgi*)	atgaggcttcgggagccgctcctgagcggcagcgccgcgatgccaggcgcgtccc tacagcgggcctgccgcctgctcgtggccgtctgcgctctgcaccttggcgtcaccc tcgtttactacctggctggccgcgacctgagccgcctgccccaactggtcggagtct ccacaccgctgcagggcggctcgaacagtgccgccgccatcgggcagtcctccgg ggagctccggaccggaggggccgc
6	Presynaptic vesicle (*sypb*)	atggacgtggtgaatcagctggtggctggggggtcagttccggggtggtcaaggagcc ccttggcttcgtgaaggtgctgcagtgggtctttgccatcttcgcctttgctacgtgcgg cagctacaccggagagcttcggctgagcgtggagtgtgccaacaagacggagagt gccctcaacatcgaagtcgaatttgagtacccattcaggctgcaccaagtgtactttga tgcaccctcctgcgttaaaggggggcactaccaagatcttcctagttggtgactactcct cctcggctgaattctttgtcaccgtggctgtgtttgccttcctctactccatgggggccct ggccacctacatcttcctgcagaacaagtaccgagagaacaacaaagggccaatga tggacttcctggccacagcagtgttcgctttcatgtggctagttagctcatccgcctgg gccaaaggcctgtccgatgtgaagatggccactgacccagagaacattatcaagga gatgcctatgtgccgccagacaggaaacacatgcaaggaactgagggaccctgtg acttcaggactcaacacctcggtggtgtttggcttcctgaacctggtgctctgggttgg caacctatggttcgtgttcaaggagacaggctgggccgcccccattcatgcgcgcacc tccaggcgccccagaaaagcaaccagctcctggcgatgcctacggcgatgcgggc tatgggcagggccccggaggctatgggcccccaggactcctacgggcctcagggtg gttatcaacccgattacgggcagccagccagcggcggtggcggtggctacgggcc tcagggcgactatgggcagcaaggctacggccaacagggtgcgcccacctccttct ccaatcagatg
7	Fibrous (*SEPT6*)	atggcagcgaccgatatagctcgccaggtggggtgaaggttgccgaactgtccccct ggctggacatgtggggtttgacagcttgcctgaccagctggtgaataagtccgtcag ccagggcttctgcttcaacatcctgtgcgtgggagagacaggtttgggcaagtccac cctcatggacaccctgttcaacaccaaattcgaaggggagccagccacccacacac agccgggtgtccagctccagtctaataacctatgacctccaagagagcaacgtgagg ctaaagctcacgatcgttagcacagttggctttggggaccagatcaacaaagaggac agctacaagcctatcgtggaattcatcgatgcacaattcgaggcctacctgcaggaa gagctaaagatccgaagagtgctacacacctaccatgactcccgaatccatgtctgct tgtatttcattgcccccacgggtcattccctgaagtctctggacctagtgactatgaaga

8	Spherical (*PRDX4m*)	agctggacagtaaggtgaacatcatccccatcattgccaaagcagatgccatttcgaa gagtgagctaacaaagttcaaaatcaaaatcaccagcgagcttgtcagcaacggagt ccagatctatcagtttcctacagatgatgagtcggtggcagagatcaatggaaccatg aacgcccacctgccgtttgctgtcattggcagcacagaagaactgaagataggcaac aagatgatgagggcgcggcagtatccttggggcactgtgcaggttgaaaacgaggc ccactgcgactttgtgaagctgcgggagatgctgattcgggtcaacatggaggatct gcgggagcagacccacacccggcactatgagctgtatcgccgctgtaagctggag gagatgggcttcaaggacaccgaccctgacagcaaacccttcagtttacaggagac atatgaggccaaaaggaacgagttcctaggggaactccagaaaaaagaagaggag atgagacagatgttcgtccagcgagtcaaagagaaagaagcggagctcaaagagg cagagaaagagctgcacgagaagtttgaccgtctgaagaaactgcaccaggacga gaagaagaaactggaggataagaagaaatccctggatgatgaagtgaatgctttcaa gcaaagaaagacggcggctgagctgctccagtcccagggctcccaggctggagg ctcacagactctgaagagagacaaagagaagaaaaataatccatggctgtgtactga a atggaggcgctgccgctgctagccgcgacaactccggaccacggccgccaccga aggctgcttctgctgccgctactgctgttcctgctgccggctggagctgtgcagggct gggagacagaggagaggcccccggactcgcgaagaggagtgccacttctacgcgg gtggacaagtgtacccgggagaggcatcccgggtatcggtcgccgaccactccctg cacctaagcaaagcgaagatttccaagccagcgccctactgggaaggaacagctgt gatcgatggagaatttaaggagctgaagttaactgattatcgtgggaaatacttggtttt cttcttctacccacttgatttcacatttgtgtctccaactgaaattatcgcttttggcgaca gacttgaagaattcagatctataaatactgaagtggtagcatgctctgttgattcacagt ttacccatttggcctggattaatacccctcgaagacaaggaggacttgggccaataag gattccacttctttcagatttgacccatcagatctcaaaggactatggtgtatacctaga ggactcaggccacactcttagaggtctcttcattattgatgacaaaggaatcctaagac aaattactctgaatgatcttcctgtgggtagatcagtggatgagacactacgtttggttc aagcattccagtacactgacaaacacggagaagtcagtcctgctggctggaaacctg gtagtgaaacaataatcccagatccagctggaaagctgaagtatttcgataaactgaa t

 Supplementary Table 3 | Standardized nomenclature of brain regions

No.	Name	Abbr.	Volume (µm^3)
1	Olfactory bulb	OB	552,907.80
2	Pallium	P	1,176,906.63
3	Subpallium	S	954,796.37
4	Preoptic region	Po	759,356.45
5	Optic tract and accessory optic system	OTAOS	986,748.95
6	Pretectum	Pr	669,170.64
7	Epiphysis (Pineal gland)	Ep	110,533.43
8	Habenula	Ha	289,854.83
9	Dorsal thalamus	DT	286,673.99
10	Ventral thalamus	VT	336,431.48
11	Posterior tuberculum	PT	377,980.15
12	Migrated posterior tubercular region	M2	658,202.84
13	Rostral hypothalamus	Hr	1,204,135.20
14	Intermediate hypothalamus	Hi	1,874,093.02
15	Diffuse nucleus of the intermediate hypothalamus	DHi	250,549.79
16	Caudal hypothalamus	Hc	176,238.95
17	Pituitary	Pit	83,580.57
18	Neuropil layer (optic tectum)	NL	2,939,375.09
19	Periventricular cell layer (optic tectum)	PL	3,760,502.17
20	Tegmentum	T	536,864.29
21	Nucleus of medial longitudinal fascicle	nMLF	30,611.07
22	Oculomotor nerve nucleus	nIII	77,251.04
23	Torus longitudinalis	TL	115,042.14
24	Torus semicircularis	TS	1,105,957.84
25	Rhombomere 1	R1	1,829,045.90
26	Trochlear nerve motor nucleus	nIV	50,476.46
27	Interpeduncular nucleus	IPN	196,849.62
28	Superior Raphe	SR	122,206.58
29	Corpus cerebelli	CCe	1,345,996.74
30	Valvula cerebelli	Va	282,478.01
31	Eminentia granularis	EG	270,844.81
32	Lobus caudalis cerebelli	LCa	32,741.70
33	Medial octavolateralis nucleus	MON	274,799.18
34	Rhombomere 2	R2	1,039,406.70
35	Dorsal motor nucleus of trigeminal nerve	nVd	36,112.51
36	Rhombomere 3	R3	897,971.87
37	Ventral motor nucleus of trigeminal nerve	nVv	16,845.30
38	Inferior raphe	IR	32,603.36
39	Rhombomere 4	R4	891,197.12
40	Rhombomere 5	R5	1,038,459.75

41	Medial vestibular nucleus	MVN	35,611.76
42	Tangential vestibular nucleus	TVN	30,629.58
43	Rostral facial nerve motor nucleus and octavolateralis efferent neurons	nVIIr/OLer	68,704.18
44	Rostral motor nucleus of abducens nerve	nVIr	6,896.53
45	Rhombomere 6	R6	1,048,429.95
46	Caudal facial nerve motor nucleus and octavolateralis efferent neurons	nVIIc/OLec	22,639.00
47	Caudal motor nucleus of abducens nerve	nVIc	14,460.40
48	Rhombomere 7	R7	930,447.61
49	Rhombomere 8	R8	1,964,286.62
50	Glossopharyngeal/Vagal nerve motor nucleus	nIX/X	188,459.61
51	Vagal lobe	VLo	15,311.87
52	Inferior olive	IO	37,314.70
53	Outer nuclear layer	ONL	1,935,701.58
54	Outer plexiform layer	OPL	218,800.10
55	Inner nuclear layer	INL	1,811,315.73
56	Inner plexiform layer	IPL	642,129.04
57	Retinal ganglion cell layer	RGCL	447,158.97
58	Ciliary marginal zone	CMZ	88,195.60
59	Optic nerve	ON	8,981.38

1993

 Supplementary Table 4 | Nucleus count and density across brain regions

No.	Name (abbr.)	Count	Density (per 10^3 μm^3)
1	OB	2,937	5.31
2	P	6,056	5.15
3	S	5,380	5.63
4	Po	4,733	6.23
5	OTAOS	2,380	2.41
6	Pr	5,021	7.50
7	Ep	461	4.17
8	Ha	2,020	6.97
9	DT	2,347	8.19
10	VT	2,875	8.55
11	PT	2,541	6.72
12	M2	1,738	2.64
13	Hr	4,398	3.65
14	Hi	6,934	3.70
15	DHi	1,589	6.34
16	Hc	1,399	7.94
17	Pit	616	7.37
18	NL	1,489	0.51
19	PL	39,919	10.62
20	T	3,098	5.77
21	nMLF	224	7.32
22	nIII	616	7.97
23	TL	864	7.51
24	TS	6,646	6.01
25	R1	9,401	5.14
26	nIV	503	9.97
27	IPN	524	2.66
28	SR	887	7.26
29	CCe	9,287	6.90
30	Va	2,184	7.73
31	EG	2,233	8.24
32	LCa	314	9.59
33	MON	467	1.70
34	R2	5,307	5.11
35	nVd	241	6.67
36	R3	4,313	4.80
37	nVv	65	3.86
38	IR	152	4.66
39	R4	3,822	4.29
40	R5	5,856	5.64
41	MVN	155	4.35
42	TVN	70	2.29

43	nVIIr/OLer	488	7.10
44	nVIr	25	3.63
45	R6	5,376	5.13
46	nVIIc/OLec	130	5.74
47	nVIc	36	2.49
48	R7	5,031	5.41
49	R8	11,784	6.00
50	nIX/X	1,430	7.59
51	VLo	96	6.27
52	IO	267	7.16
53	ONL	23,576	12.18
54	OPL	1,122	5.13
55	INL	38,063	21.01
56	IPL	89	0.14
57	RGCL	6,988	15.63
58	CMZ	1,051	11.92
59	ON	21	2.34

1995

 Supplementary Table 5 | PSD count and density across brain regions

No.	Name (abbr.)	Count	Density (per 10^3 µm³)
1	OB	168,961	305.59
2	P	619,202	526.13
3	S	570,921	597.95
4	Po	397,593	523.59
5	OTAOS	969,580	982.60
6	Pr	185,228	276.80
7	Ep	5,371	48.59
8	Ha	74,603	257.38
9	DT	51,797	180.68
10	VT	59,776	177.68
11	PT	194,962	515.80
12	M2	890,300	1,352.62
13	Hr	1,248,748	1,037.05
14	Hi	2,269,327	1,210.89
15	DHi	47,967	191.45
16	Hc	29,554	167.69
17	Pit	6,949	83.14
18	NL	6,274,593	2,134.67
19	PL	143,460	38.15
20	T	488,299	909.54
21	nMLF	4,158	135.83
22	nIII	20,325	263.10
23	TL	10,079	87.61
24	TS	997,075	901.55
25	R1	1,817,808	993.86
26	nIV	3,156	62.52
27	IPN	235,891	1,198.33
28	SR	31,382	256.79
29	CCe	543,830	404.04
30	Va	55,456	196.32
31	EG	94,508	348.94
32	LCa	256	7.82
33	MON	387,594	1,410.46
34	R2	1,024,545	985.70
35	nVd	6,723	186.17
36	R3	768,601	855.93
37	nVv	17,612	1,045.51
38	IR	11,246	344.93
39	R4	746,597	837.75
40	R5	660,500	636.04
41	MVN	15,087	423.65
42	TVN	31,957	1,043.34

43	nVIIr/OLer	10,745	156.40
44	nVIr	8,262	1,197.99
45	R6	750,372	715.71
46	nVIIc/OLec	1,141	50.40
47	nVIc	18,099	1,251.63
48	R7	864,232	928.83
49	R8	1,309,422	666.61
50	nIX/X	82,721	438.93
51	VLo	3,171	207.09
52	IO	17,618	472.15

1997

 Supplementary Table 6 | SVC count and density across brain regions

No.	Name (abbr.)	Count	Density (per 10^3 μm^3)
1	OB	157,880	285.54
2	P	440,095	373.94
3	S	418,483	438.30
4	Po	274,409	361.37
5	OTAOS	567,745	575.37
6	Pr	107,340	160.41
7	Ep	6,561	59.36
8	Ha	38,386	132.43
9	DT	28,278	98.64
10	VT	34,647	102.98
11	PT	111,957	296.20
12	M2	578,311	878.62
13	Hr	756,460	628.22
14	Hi	1,493,329	796.83
15	DHi	28,689	114.50
16	Hc	20,016	113.57
17	Pit	11,688	139.84
18	NL	3,864,426	1,314.71
19	PL	81,328	21.63
20	T	325,770	606.80
21	nMLF	1,988	64.94
22	nIII	11,687	151.29
23	TL	6,317	54.91
24	TS	584,917	528.88
25	R1	1,091,715	596.88
26	nIV	1,351	26.76
27	IPN	223,406	1,134.91
28	SR	18,389	150.47
29	CCe	412,867	306.74
30	Va	51,639	182.81
31	EG	45,662	168.59
32	LCa	137	4.18
33	MON	179,624	653.66
34	R2	587,247	564.98
35	nVd	4,383	121.37
36	R3	439,578	489.52
37	nVv	9,836	583.90
38	IR	8,031	246.32
39	R4	447,452	502.08
40	R5	396,779	382.08
41	MVN	9,154	257.05
42	TVN	17,689	577.51

43	nVIIr/OLer	7,489	109.00
44	nVIr	5,165	748.93
45	R6	457,744	436.60
46	nVIIc/OLec	814	35.96
47	nVIc	11,255	778.33
48	R7	516,297	554.89
49	R8	809,742	412.23
50	nIX/X	50,822	269.67
51	VLo	2,377	155.24
52	IO	12,457	333.84

1999

 Supplementary Table 7 | Mito count and density across brain regions

No.	Name (abbr.)	Count	Density (per 10^3 μm^3)
1	OB	146,644	265.22
2	P	365,682	310.71
3	S	364,324	381.57
4	Po	247,663	326.15
5	OTAOS	535,443	542.63
6	Pr	133,626	199.69
7	Ep	11,706	105.90
8	Ha	58,557	202.02
9	DT	45,091	157.29
10	VT	47,103	140.01
11	PT	107,722	284.99
12	M2	411,387	625.02
13	Hr	617,761	513.03
14	Hi	1,115,733	595.35
15	DHi	30,243	120.71
16	Hc	34,030	193.09
17	Pit	17,205	205.85
18	NL	2,582,241	878.50
19	PL	265,164	70.51
20	T	303,911	566.09
21	nMLF	3,942	128.78
22	nIII	18,408	238.29
23	TL	15,109	131.33
24	TS	509,154	460.37
25	R1	905,849	495.26
26	nIV	4,828	95.65
27	IPN	217,883	1,106.85
28	SR	27,805	227.52
29	CCe	399,534	296.83
30	Va	65,467	231.76
31	EG	73,700	272.11
32	LCa	1,190	36.35
33	MON	251,781	916.24
34	R2	490,509	471.91
35	nVd	6,031	167.01
36	R3	402,701	448.46
37	nVv	6,147	364.91
38	IR	13,723	420.91
39	R4	427,100	479.24
40	R5	413,002	397.71
41	MVN	12,173	341.83
42	TVN	25,790	842.00

43	nVIIr/OLer	14,259	207.54
44	nVIr	4,869	706.01
45	R6	497,382	474.41
46	nVIIc/OLec	5,806	256.46
47	nVIc	12,623	872.94
48	R7	487,363	523.79
49	R8	866,776	441.27
50	nIX/X	54,772	290.63
51	VLo	3,139	205.00
52	IO	15,150	406.01

2001

Supplementary Table 8 | Perisomatic features of multiple APEX2-labeled neurons

	LC-NE neurons	MO-NE neurons	OB-DA neurons	PT-DA neurons	Hc-DA neurons	Pr-5HT neurons	RN-5HT neurons	Hcrt neurons	Gly neurons
Soma volume (μm^3)	314 (131)	116 (79.0)	106 (99.5)	303 (111)	92.8 (76.5)	104 (86.5)	139 (103)	137 (103)	82.3 (77.5)
Soma surface area (μm^2)	344 (180)	158 (126)	161 (148)	367 (169)	149 (139)	165 (147)	201 (157)	190 (165)	138 (130)
Soma volume/surface ratio (μm)	0.902 (0.710)	0.732 (0.617)	0.658 (0.675)	0.824 (0.658)	0.614 (0.552)	0.630 (0.591)	0.685 (0.652)	0.720 (0.627)	0.595 (0.598)
Soma sphericity	0.648 (0.687)	0.730 (0.701)	0.674 (0.708)	0.595 (0.669)	0.666 (0.632)	0.652 (0.647)	0.643 (0.679)	0.677 (0.652)	0.663 (0.682)
Soma/convex hull area ratio	1.05 (1.08)	1.01 (1.07)	1.01 (1.04)	1.09 (1.05)	1.04 (0.972)	1.08 (1.08)	1.01 (1.05)	1.12 (1.09)	1.04 (1.04)
Soma/convex hull volume ratio	0.686 (0.737)	0.731 (0.765)	0.694 (0.738)	0.637 (0.687)	0.681 (0.592)	0.693 (0.705)	0.650 (0.720)	0.758 (0.694)	0.712 (0.712)
Soma area minus convex hull area (μm^2)	14.8 (12.7)	1.13 (8.20)	-3.38 (3.28)	32.0 (-15.6)	3.71 (-22.1)	11.6 (9.99)	-0.776 (4.84)	20.0 (11.0)	3.41 (0.745)
Convex hull volume minus soma volume (μm^3)	148 (48.7)	45.1 (25.0)	57.0 (40.8)	171 (84.9)	49.0 (75.6)	48.3 (37.8)	75.9 (45.8)	43.9 (54.2)	35.8 (38.0)
Soma elongation ratio	4.64 (4.42)	3.00 (3.35)	3.92 (3.24)	5.68 (5.38)	3.27 (6.95)	4.15 (4.74)	5.22 (5.07)	3.33 (4.88)	4.65 (4.06)
Soma radial range (μm)	7.88 (4.48)	4.08 (3.25)	4.49 (3.60)	7.41 (5.42)	3.81 (5.66)	4.83 (4.73)	7.04 (4.52)	4.19 (4.16)	4.98 (4.41)
Soma normalized max radius	3.43 (2.25)	2.17 (1.73)	2.47 (1.89)	2.74 (1.85)	1.92 (30.0)	2.70 (3.82)	5.90 (2.80)	1.61 (2.79)	4.33 (4.55)
Soma normalized min radius	0.0337 (0.0486)	0.0723 (0.0598)	0.0354 (0.0548)	0.0140 (0.0575)	0.0544 (0.176)	0.0372 (0.0426)	0.0351 (0.0469)	0.0459 (0.0468)	0.0412 (0.0502)
Soma eccentricity	0.833 (0.742)	0.776 (0.792)	0.820 (0.777)	0.842 (0.757)	0.776 (0.801)	0.774 (0.804)	0.823 (0.787)	0.815 (0.769)	0.845 (0.806)
Nucleus volume (μm^3)	117 (79.5)	78.4 (58.4)	77.9 (71.3)	128 (77.0)	64.8 (55.3)	66.6 (63.0)	78.5 (68.0)	91.4 (71.7)	57.8 (56.7)
Nucleus surface area	126 (96.3)	94.3 (77.7)	96.4 (90.7)	141 (93.7)	84.6 (77.8)	90.1 (85.1)	100 (91.9)	105 (90.8)	81.1 (79.6)

(μm^2)									
Nucleus volume/surface ratio (μm)	0.920 (0.820)	0.831 (0.737)	0.808 (0.782)	0.901 (0.815)	0.762 (0.710)	0.741 (0.741)	0.779 (0.739)	0.866 (0.782)	0.711 (0.710)
Nucleus sphericity	0.913 (0.926)	0.939 (0.923)	0.915 (0.916)	0.867 (0.931)	0.921 (0.904)	0.885 (0.902)	0.881 (0.878)	0.932 (0.917)	0.890 (0.897)
Nucleus/convex hull area ratio	1.01 (1.01)	1.00 (1.00)	1.00 (1.01)	1.01 (1.00)	1.00 (1.01)	1.02 (1.01)	1.01 (1.01)	1.01 (1.00)	1.00 (1.01)
Nucleus/convex hull volume ratio	0.946 (0.961)	0.971 (0.973)	0.965 (0.965)	0.926 (0.966)	0.964 (0.952)	0.937 (0.948)	0.928 (0.928)	0.972 (0.957)	0.946 (0.942)
Nucleus area minus convex hull area (μm^2)	1.55 (0.554)	0.203 (0.279)	0.364 (0.770)	1.79 (0.326)	0.0976 (0.612)	2.06 (0.465)	0.588 (0.956)	0.521 (0.0955)	0.201 (0.621)
Convex hull volume minus nucleus volume (μm^3)	6.72 (3.32)	2.35 (1.41)	2.88 (2.58)	10.4 (2.62)	2.51 (2.91)	4.53 (3.46)	6.00 (5.38)	2.65 (3.33)	3.24 (3.61)
Nucleus elongation ratio	2.05 (1.94)	1.73 (2.02)	2.21 (2.34)	2.46 (2.20)	1.97 (2.15)	2.36 (2.40)	2.57 (3.10)	1.84 (3.37)	2.38 (2.23)
Nucleus radial range (μm)	1.93 (1.58)	1.45 (1.52)	1.98 (1.81)	2.54 (1.58)	1.61 (1.70)	1.88 (1.86)	2.28 (2.12)	1.66 (1.77)	1.90 (1.75)
Nucleus normalized max radius	0.508 (0.540)	0.521 (0.619)	0.676 (0.626)	0.649 (0.608)	0.560 (0.690)	0.606 (0.742)	0.734 (0.855)	0.538 (0.751)	0.726 (0.677)
Nucleus normalized min radius	0.0703 (0.0928)	0.102 (0.0903)	0.0697 (0.0778)	0.0473 (0.0971)	0.0778 (0.0803)	0.0618 (0.0724)	0.0491 (0.0683)	0.0856 (0.0896)	0.0691 (0.0758)
Nucleus eccentricity	0.601 (0.659)	0.697 (0.702)	0.720 (0.718)	0.714 (0.669)	0.738 (0.739)	0.700 (0.701)	0.775 (0.732)	0.671 (0.707)	0.758 (0.716)
Mitochondrial count	165 (13.1)	14.0 (6.48)	9.40 (9.02)	66.3 (5.98)	15.4 (8.06)	5.40 (6.06)	26.2 (9.02)	12.6 (7.38)	6.30 (4.38)
Mitochondrial density per 1 μm^3 soma volume	0.547 (0.0847)	0.120 (0.0803)	0.0838 (0.0912)	0.219 (0.0494)	0.181 (0.106)	0.0528 (0.0671)	0.175 (0.0784)	0.0974 (0.0704)	0.0745 (0.0555)
Mitochondrial density per 1 μm^3 nucleus volume	1.42 (0.147)	0.177 (0.109)	0.117 (0.153)	0.520 (0.0732)	0.243 (0.145)	0.0806 (0.0931)	0.318 (0.122)	0.146 (0.102)	0.107 (0.0758)
Nucleus/soma volume ratio	0.386 (0.640)	0.684 (0.750)	0.741 (0.724)	0.425 (0.712)	0.731 (0.731)	0.653 (0.733)	0.585 (0.683)	0.667 (0.703)	0.710 (0.740)

2004 Data are presented as mean values, with values in parentheses representing unlabeled
2005 cells adjacent to APEX2-labeled neurons. See Extended Data Fig. 7 for cell numbers;
2006 see Methods for calculation details.
2007

Figures

Fig.1

2012   Fig.2

2013
2014

Fig.4

2021   Fig.5

Spatial organization of broad-yet-sparse inputs

a. Broad-yet-sparse synaptic inputs

b. Multi-level spatial organization

→ Single synapse

① Broad-yet-sparse synaptic inputs
② Proximity-dependent synaptic strength scaling
③ Modality and E/I type-based distribution
④ Dual input architecture
⑤ Cross-system shared inputs with proximal bias

2022

2023

2025 Extended Data Fig.1

2026

2027

2028   Extended Data Fig.2

2029
2030

Extended Data Fig.3

Extended Data Fig.4

2038
2039

Extended Data Fig.6

  Extended Data Fig.7

2046   Extended Data Fig.8

2047

2048

 Extended Data Fig.10