1 **Title: Causes and consequences of experimental variation in *Nicotiana benthamiana***
2 **transient expression**
3

4 **Authors:** Sophia N. Tang[1,2,3], Matthew J. Szarzanowicz[1,2,4], Amy Lanctot[1,2,4], Sasilada
5 Sirirungruang[1,2,4], Liam D. Kirkpatrick[1,2,4], Krista Drako[1,2,3], Simón Álamos[1,2,4], Adela Cheng[1,2,7],
6 Lucas M. Waldburger[1,2,5,6], Shuying Liu[1,2], Mitchell G. Thompson[1,2*], Patrick M. Shih[1,2,4,8*]
7

Affilitations:
1 Joint BioEnergy Institute, 5885 Hollis Street, Emeryville, CA 94608, USA.
2 Environmental Genomics and Systems Biology Division, Lawrence Berkeley National
Laboratory, Berkeley, California, USA
3 Department of Molecular and Cell Biology, University of California, Berkeley, CA 94720, USA
4 Department of Plant and Microbial Biology, University of California, Berkeley, CA 94720, USA
5 Department of Bioengineering, University of California, Berkeley, California, USA
6 Biological Systems & Engineering Division, Lawrence Berkeley National Laboratory, Berkeley,
CA 94720, USA
7 Department of Nutritional Sciences and Toxicology, University of California, Berkeley, CA
94720, USA
8 Innovative Genomics Institute, Berkeley, California, USA
20

* Correspondence to Patrick M. Shih (pmshih@berkeley.edu) or Mitchell G. Thompson
(mthompson@lbl.gov)

Abstract

Infiltration of *Agrobacterium tumefaciens* into *Nicotiana benthamiana* has become a
foundational technique in plant biology, enabling efficient delivery of transgenes *in planta* with
technical ease, robust signal, and relatively high throughput. Despite transient expression's
prevalence in disciplines such as synthetic biology, little work has been done to describe and
address the variability inherent in this system, a concern for experiments that rely on highly
quantitative readouts. In a comprehensive analysis of *N. benthamiana* agroinfiltration
experiments, we model sources of variability that affect transient expression. Our findings
emphasize the need to validate normalization methods under the specific conditions of each
study, as distinct normalization schemes do not always reduce variation either within or between
experiments. Using a dataset of 1,813 plants collected over three years, we develop a
comprehensive model of variation in *N. benthamiana* transient expression, using power analysis
to determine the number of individual plants required for a given effect size. Drawing on our
longitudinal data, these findings inform practical guidelines for minimizing variability through
strategic experimental design and power analysis, providing a foundation for more robust and
reproducible use of *N. benthamiana* in quantitative plant biology and synthetic biology
applications.

Introduction

Nicotiana benthamiana is a workhorse of plant molecular biology due to the high efficiency of transient *Agrobacterium*-mediated transformation, enabling researchers to assess gene function and observe phenotypes within days, as opposed to the months it typically takes for stable transformations (1–3). *N. benthamiana*'s capacity for transient expression precipitated its use to study diverse aspects of cell and molecular biology (4), including virus-induced gene silencing (5), subcellular protein localization (6), biosynthetic pathway discovery and engineering (7–11), and gene regulation (12–16). Given the lengthy time required to generate stable transformants and the technical challenges inherent in protoplast transformation (17,18), there are few viable *in planta* alternatives to *N. benthamiana* transient expression, leading to its primacy as the model of plant synthetic biology (19,20).

Despite its widespread use, *N. benthamiana* lacks substantial efforts to assess error and reproducibility, like many other systems routinely leverage in synthetic biology (21). Nor is there a standardized way to design transient assays, and as a result, experimental design varies from publication to publication (13,22–26). These factors obstruct our ability to model and predict variation in this model system, which is crucial for effective experimental design. Limited quantitative understanding of biology has routinely impeded genetic engineers in developing predictable and reliable bioproducts (27), in comparison to the precision associated with traditional engineering disciplines (28). For plant synthetic biology to truly be an engineering discipline, we require robust evaluation of variability and its effects on our design goals.

The variability of transient expression in *N. benthamiana* in highly powered and longitudinal experiments has never been rigorously evaluated. Without robust statistics, researchers risk designing underpowered experiments and overlooking subtle effects. While there have been attempts to determine transgene expression variability in stable plant lines (29,30), it is unclear whether these findings translate to transient expression. Transient expression variability in *N. benthamiana* has been primarily studied using luciferase reporters, but these studies are plagued by small sample size, lack of biological replicates, and the high noise inherent to the luciferase assay (31–33). Most methods used to reduce variability in transient expression experiments have relied on using a second reporter (hereafter referred to as a normalizer) for ratiometric normalization (13,22–24,34,35). In this approach, every tested experimental construct is co-delivered with the same constitutively expressed normalizer, under the assumption that any variation in the experimental construct's expression that tracks with variation in the normalizer's expression must not be due to the sequence of the reporters but rather other variables. Although this approach is widely used, whether normalizing transgene expression actually reduces and controls for variation has not been systematically examined.

Here, we use fluorescence reporter assays to systematically determine the sources of variation of transient expression in *N. benthamiana* leaves. Using a large sample size (>1800 plants) across multiple years of independent experiments, we comprehensively capture transgene expression variation observed between plants and between independent experiments in order to model sources of error. We also complete the first systematic comparison of methods to normalize transgene expression and demonstrate whether and to what extent these methods

reduce variability. Finally, we suggest best practices to mitigate variation through purposeful
experimental design and statistical power analysis.

Results

Plant batch effects are the greatest sources of transient expression variation

To determine sources of variability in *N. benthamiana* transient expression, we analyzed
previously published data wherein the same GFP reporter was agroinfiltrated in 15 independent
experimental replicates using *A. tumefaciens* GV3101::pMP90 (hereafter GV3101) (Fig 1A, Fig.
S1) (36). The reporter strain was infiltrated into the fourth and fifth leaf from the top (leaves T4
and T5) of *N. benthamiana*, and three days post-infiltration, four discs were collected from each
infiltrated leaf. Dates of measurement are unique identifiers for experimental replicates. In this
dataset, we observed as much as a fourfold difference in mean fluorescence between the
lowest (teal, 2022.07.25) and highest (orange, 2023.05.30) replicates (Fig. 1B, 1C). Using a
mixed-effects model, we attributed 41.8%, 13.4%, and 3.2% of the variation to replicate-to-
replicate ("Date"), plant-to-plant ("Plant"), and leaf-to-leaf ("Leaf") differences, respectively (Fig.
1D). The underlying factors of these sources of variation may be batch effects across the fifteen
replicates' plants, non-homogenous growing conditions, and varying leaf cellular ages (Fig. 1E)
(37,38). Though the precise percent contributions of these variables may vary depending on the
transgene(s) introduced, they are nonetheless major sources of variation that should be
accounted for in experimental design.
103
The remaining 41.5% of variation could not be explained using these three variables, so we
expanded our independent parameters to identify other potential sources. In a separate set of
independent experiments, we assessed the effects of 96-well water volume, plant age, and
collection time (both time of day and time elapsed between collection and measurement) and
found that water volume and plant age are negatively correlated with fluorescence, while
collection time had no effect (Fig. S2-S5). Minor changes to water volume (*e.g.*, evaporation,
pipetting error) may have contributed to residual error. All other plants in this study were 4
weeks old at the time of infiltration, however, so no variation can be attributed to differences in
age. With these additional experiments, we have accounted for the majority of variation in our
model and ruled out other potential sources.

Multiple methods of co-delivering a normalizing transgene can decrease expression
variability

Normalizing the expression level of a transgene of interest to that of another independent
transgene is thought to control for any sources of variability independent of the tested
sequences, since they should impact expression of the two transgenes equally. This practice
should theoretically reduce variation both within and between experiments. However, there are
many different conceivable ways the two transgenes could be delivered in *N. benthamiana*, and
indeed normalization schemes differ between publications (13,22–26).
122

We systematically compared normalization schemes for transient expression using fluorescent
protein (FP) reporters, whose gene expression is correlated to an easily measured phenotype,
fluorescence. The normalized readout of the assay is the ratio of eGFP and mCherry
fluorescence measured in a leaf disc. Designating FPs as the reporter or normalizer is arbitrary.
The two expression cassettes were co-delivered in three different ways. First, we co-infiltrated a
mix of two different strains of *A. tumefaciens*, each with a binary vector containing one FP
(schemes 3-7). Second, we infiltrated single *A. tumefaciens* strains carrying binary vectors with
a T-DNA containing both cassettes in every possible orientation, both relative to one another
and to the left and right borders (schemes 8-15). Third, we infiltrated a "BiBi" strain (39), wherein
a single strain of *A. tumefaciens* carries two binary vectors, each with a unique FP gene and ori
(schemes 16-19). All FP genes are driven by the PCM2 promoter and are terminated by the
T_AtUbq3 terminator (Supplementary Table 1, Fig. S1) (14). These 17 methods of co-delivery
were compared to delivering eGFP (scheme 1) or mCherry (scheme 2) alone in two
independent experimental replicates.
137
Co-delivery methods have obvious ramifications for basal transgene expression levels that are
consistent between both experimental replicates (Fig. S6). For both co-infiltrations and BiBis,
the low copy number origin, pSa, consistently produced much lower fluorescence compared to
the higher copy number origins, pVS1 or BBR1 (Fig. 2A) (40). For the stacked, two-cassette T-
DNAs, despite sharing the same pVS1 origin and backbone, these schemes yielded up to eight-
fold difference in fluorescence depending on the cassettes' relative orientations (Fig. 2A, S7).
Convergent, tandem, and divergent cassettes produced the strongest, intermediate, and
weakest expression, respectively (Fig. S7). The expression of one gene is affected by the
orientation of the other gene, even when its own position and orientation is maintained, as with
schemes 8 and 10 for eGFP or schemes 9 and 11 for mCherry (Fig. S7). These results suggest
that gene position relative to the left and right borders and, most importantly, relative orientation
strongly affect expression in multi-gene T-DNAs.
150
Variability, as measured by coefficient of variation (CV) from all leaf discs, decreases for all
normalization schemes (Fig. S8). Furthermore, all schemes have a lower CV for
reporter/normalizer than for reporter (Fig. S9). The more appropriate comparison, however, is
whether a scheme's reporter/normalizer CV is lower than the corresponding unnormalized
control's reporter CV. Few schemes produce an eGFP/mCherry CV that is significantly less than
scheme 1's eGFP CV following a Bonferroni correction (Fig. 2B), and only scheme 3 (co-
infiltration of two pVS1 binary vectors) meets this condition when mCherry is treated as the
reporter (Fig. 2C). Such a test is conservative, and it should be noted that there is at least one
scheme per co-delivery method that reduces CV by >50% compared to the unnormalized
control, whether CV is calculated per plant or pooling all discs together (Fig. 2B, 2C, S8). All co-
delivery methods can offer reductions in variability, thereby increasing the statistical power of
transient expression-based experiments.

Inoculum densities of co-infiltrated strains affect signal strength but not variability

Because the decrease in CV was the greatest and most statistically significant for co-infiltration
of two pVS1 binary vectors (scheme 3), we used this scheme to explore other possible

experimental variables for further reductions in variability. Additionally, co-infiltration is highly
modular, and multiple T-DNAs can efficiently be delivered to each plant cell (39,41). Since *A.*
tumefaciens strains in *N. benthamiana* leaves can antagonize or synergize with one another
depending on their densities (39), we next determined whether the OD_{600} of each strain in the
co-infiltration affects the resulting CV. Each strain was added to the infiltration mix at an OD_{600}
of 0, 0.01, 0.1, 0.5, or 1 and combined with the other strain, also added at the same densities for
a total of 24 possible combinations.
173
Total OD_{600} of the co-infiltrated mix and the OD_{600} of the individual strains do not greatly nor
predictably affect variability, but inoculum density of a strain dictates its total transgene
expression levels. There is no discernible pattern in eGFP CV with respect to the OD_{600} of the
co-infiltrated strains (Fig. 3A), and while all CVs for the ratio of eGFP/mCherry are
comparatively much lower (Fig. 3B), there is no one combination of ODs that is significantly
superior to all others. The same trends are apparent when mCherry is treated as the reporter
and eGFP as the normalizer (Fig. S10). For raw fluorescence, signal begins to saturate beyond
an OD_{600} of 0.5, and when one strain is held at a given OD_{600} while the OD_{600} of another
competing strain increases, fluorescence from the fixed strain decreases (Fig. 3C, 3D). The
primary concern when selecting OD_{600}, then, should be the desired signal strength. Of the
densities tested, 0.1 results in strong, measurable signal and, in agreement with previous
findings, does not saturate the transgene expression capacity of the plant cell (39,41).

Promoter choice determines the utility of normalization

Given that the choice of promoter(s) in transient transformations affects not only signal strength
but also normalization outcomes (14), we more thoroughly explored additional promoter
combinations. All transgenes in this study have hitherto been driven by a medium-strength
promoter, PCM2. From a library of constitutive promoters characterized by Zhou et al., we
selected low- and high-strength promoters, PCL2 and PCH5, to test alongside PCM2. All three
promoters were used to drive either eGFP and mCherry expression to make a total of six binary
vectors. The eGFP binary vectors were co-infiltrated with the mCherry binary vectors in every
possible combination, and all binary vectors were also infiltrated alone for a total of 15 unique
infiltrations.
196
There is no discernible pattern in eGFP CV for any of the infiltrations (Fig. 4A), but for
eGFP/mCherry, identical promoters driving the two FP genes yield the lowest CVs (Fig. 4B).
Outside of these pairs, normalization does not provide as large reductions in variations, if at all.
In fact, for three nonidentical promoter pairs, their eGFP/mCherry CV is greater than their own
eGFP CV and, for one pair, also greater than the eGFP CV of its corresponding unnormalized
control (Fig. 4A, 4B). Similar observations are true when treating mCherry as the reporter and
eGFP as the normalizer (Fig. S11). Additionally, fluorescence from a given strain diminishes the
stronger a competing strain's FP promoter is, though this trend is weaker for mCherry than it is
for GFP (Fig. 4C, 4D).
206
We then sought to clarify whether the large decrease in CV when using identical promoters is
due to the similar promoter strengths or to other variables, such as shared trans factors

affecting mRNA abundance. To do so, we generated another three binary vectors with mCherry
driven by PCL1, PCM1, and PCH4, which are the promoters in the library closest in expression
to the three already tested. As with the first promoter combination experiment, the existing
eGFP binary vectors were co-infiltrated with the new mCherry binary vectors in every possible
combination, and all six binary vectors were also infiltrated alone.

With these promoters, both the CVs of eGFP and of eGFP/mCherry follow no apparent pattern
(Fig. 4E, 4F), but the negative trend between fluorescence signals of the two competing strains
holds true, although PCH4 fits poorly in the trend (Fig. 4G, 4H). Several promoter pairs have
higher eGFP/mCherry CVs than their own eGFP CVs and/or the eGFP CV of the unnormalized
control (Fig. 4E, 4F), and vice versa when treating mCherry as the reporter (Fig. S11). Similarity
in promoter strength, then, does not guarantee lower CVs upon normalization. Depending on
the promoters used to normalize, normalization may even worsen the assay's statistical power.
Since the co-delivery comparison and the OD optimization experiments were conducted using
PCM2 for every expression cassette, we had fortuitously picked the best case scenario,
identical promoters. Had we chosen nonidentical promoters, all methods of co-delivery might
not have decreased CV compared to the unnormalized controls. When possible, identical
promoters should be used.

Normalization does not yield reproducibility of absolute quantification

Normalization can serve two distinct purposes: (1) decreasing within-experiment variation to
improve detection of small effect sizes, or (2) decreasing between-experiment variation to allow
for comparison across independent experiments. Mitigating within-experiment variation does not
necessarily have an effect on between-experiment variation. To explore whether normalization
makes comparisons between replicates or to historical data more valid, we co-infiltrated
PCL2:GFP alone and with all six mCherry binary vectors thus far generated in six independent
replicates.

Normalization makes replicates more similar to one another for most, but not all, promoter pairs
(Fig. 5A, 5B, S12). We performed one-sample Kolmogorov-Smirnov tests for every condition to
determine the likelihood that a sample distribution (an individual replicate) is drawn from a
reference distribution (all replicates, pooled). The cumulative density functions (CDFs) show that
sample distributions nearly all become more comparable to the reference distribution with
normalization, *i.e.*, the maximum vertical distance between the distributions decreases (Fig. 5A,
5B, S12). Replicate-to-replicate, the PCL2/PCM2 promoter pair produces the most consistent
distribution of GFP/mCherry values, far more reproducible than its raw eGFP fluorescences
(Fig. 5A, 5B). The sole exception is the PCL2/PCM1 promoter pair, whose GFP/mCherry CDFs
are actually less comparable to the reference distribution than its GFP CDFs are (Fig. 5A, 5B);
there is no singular GFP/mCherry value that normalization causes the data to reliably converge
upon. However, this pair and PCL2/PCL2 were the only promoter pairs for which replicate CVs,
representing within-experiment variation, were regularly lower than the unnormalized control's
(Fig. 5C). Additionally, PCL2/PCL2 produces the most consistent replicate CVs–it alone has an
interquartile range of replicate CVs narrower than the unnormalized control's (Fig. 5C).
Decreasing within-experiment variation, then, is not equivalent to decreasing between-

experiment variation. Normalization schemes must be designed with the desired outcome in
mind (minimizing within- and/or between-experiment variation) and then validated.

Developing a generalizable power analysis framework of N. benthamiana *transient expression*

To evaluate whether normalization meaningfully reduces the number of plants required to detect
a given effect size, we developed a model informed by a large dataset of *N. benthamiana*
transient expression experiments. This dataset comprises 1,813 plants infiltrated with a GFP
reporter over nearly three years, spanning experiments conducted by multiple researchers and
diverse conditions, including different inoculum densities and *Agrobacterium* strains (see
Supplementary Information for details). We calculated the CV for each plant infiltrated with one
of two *Agrobacterium* strains: GV3101 and the hypervirulent EHA105, which displays both
elevated transgene expression and variance compared to GV3101 (Fig. 6A). For batches of
plants for which at least 30 plants were used, we visualized the plant CVs in a CDF (Fig. 6B
solid gray line). With these per-plant and per-experiment CVs, we developed a Monte Carlo
simulation of assay variability, which closely recapitulates the observed distribution of real plant
CVs across the training dataset (Fig. 6B dotted black line; Wasserstein distance = 0.023, mean
ECDF difference = 0.05).

We then simulated how variability propagates through an experiment on the per-plant and per-
experiment levels. First, a batch CV is drawn from the empirical distribution shown in Fig. 6A by
randomly generating a percentile and selecting the corresponding CV, which represents the
average CV of plants from a hypothetical replicate. Around this batch-level value, we simulate
CVs of individual plants by incorporating additional noise that approximates within-experiment
heterogeneity between plants. For each simulated plant, eGFP fluorescence data for eight leaf
discs is generated based on the plant CV and a fixed, construct-specific mean expression. Two
conditions are compared with an independent, two-tailed Student's *t*-test at varying effect size
differences and sample sizes (Fig. 6C). For each comparison, 1000 independent replicates are
simulated and used to calculate the probability of a successful comparison.

This approach enables us to estimate the minimum number of plants needed to reliably (>95%
of simulations) and significantly ($p < 0.05$) distinguish between two constructs. The smallest
detectable effect size using 50 plants for unnormalized EHA105 was 15.9%, compared to 10.0%
for unnormalized GV3101 and 8.3% for GV3101 under optimal normalization (Fig. 6D). While
larger sample sizes can resolve even smaller differences, practical constraints limit the number
of plants that can be included in an experiment, but relatively small effect sizes can be
accurately distinguished with a small number of plants. For instance, with $n = 3$, the minimum
reliably detectable effect size for unnormalized GV3101 is ~40%. Normalization is most
beneficial in experiments where small effect sizes must be resolved using limited sample sizes.
However, since normalization may also reduce statistical power in some cases (Fig. S13), any
normalization strategy should be validated empirically prior to implementation.

Discussion

Here, we systematically measure expression variability in *N. benthamiana* agroinfiltration. The inherent noise in the transient expression of FPs is considerable (as much as fourfold difference in mean expression across replicates) but mitigable through proper experimental design (Fig. 1B). Using a mixed-effects model, we attribute 41.8% of the observed variation to batch effects among the plants (Fig. 1D). We find that multiple co-delivery methods for dual reporters can decrease variation via ratiometric normalization, and of the examined methods, co-infiltration best reduces variation, approximately halving the CV compared to no normalization (Fig. 2). Neither inoculum density nor promoter strength in co-infiltrations predictably affect expression variation, but normalization efficacy is highly sensitive to promoter choice (Fig. 3, Fig. 4). Some promoter pairs may in fact increase variation or make independent experiments less comparable, and using identical promoters is the surest and most effective way to reduce CV within experiments (Fig. 4, Fig. 5). Normalization should not be done arbitrarily but rather validated on a case-by-case basis for the co-delivery method and experimental design used, as this is absolutely necessary for results that are reproducible and robust to potential sources of variability.

Though unambiguously beneficial, the use of constitutive, identical promoters is also particularly limiting given the popularity of *N. benthamiana* to characterize the effects of promoter identity and architecture on transcription. This problem is undoubtedly compounded in more complex experimental designs. To increase predictability and standardization of normalization methods, multi-lab efforts to quantify variation using diverse methods will be necessary. More robust transcriptomic datasets of *N. benthamiana* expressing diverse transgenes in distinct host strains could discourage reliance on a single, unvalidated normalizer. Additionally, multi-factor analysis of binary vector construction and expression cassette positioning could elucidate the extent to which these variables affect both expression and variability.

There has yet to be a published analysis, to our knowledge, of how the design of multi-gene T-DNA structure affects the transient expression of those genes. We find that convergent cassettes are far more highly expressed than tandem or divergent cassettes. Since *Agrobacteria*'s VirD2 virulence protein covalently binds the right border (46), it is thought that transgenes proximal to the right border would be protected from exonucleolytic degradation and thus more likely to remain intact inside the plant cell, but this hypothesis is not fully supported by our data (Fig. 2A, Fig. S7). Clearly, T-DNA architecture affects transgene expression strength and variation, and a high-throughput library approach would be ideal to address the multiplicity of orientation and position combinations in multi-gene binary vectors. Determining these rules of expression and assessing them in both stable and transient systems requires further, data-intensive work.

Given the lack of large, curated datasets of transient gene expression in *N. benthamiana*, it is likely that many published *N. benthamiana* experiments chose sample sizes arbitrarily based on unvalidated assumptions. Our model of the statistical power required to determine different effect sizes with high confidence shows that in cases where the expected effect size is large, it is easy to achieve the necessary sample size. For example, for effect sizes >50%, fewer than

three plants are required using the GV3101 strain. For effect sizes <20%, many plants are
required, to the extent that the number of plants needed may be infeasible for single
experiments (*e.g.* 50 plants are needed to show a 15.9% effect using EHA105). The majority of
synthetic biology applications for transient expression, such as bioproduction or circuit
reconstruction, require strong effects, which call for small numbers of plants. However, these
results could have strong implications for subtle phenotypes, like the transformation efficiency of
various *Agrobacterium* strains or mutants (36,42), or inducible gene expression systems (43–
45).
344
While the data in this work required many hundreds of plants and multiple years to collect, we
were nonetheless only able to capture variation in *N. benthamiana* transient expression from a
rather narrow perspective. All of our data are from fluorescent reporters from 4-week-old plants,
with data collected three days after infiltration. We demonstrated that *Agrobacterium* strain,
promoter choice, and T-DNA design all impact variation, but many more strains, plasmids, and
expression cassette designs remain untested. There are other variables affecting
Agrobacterium-mediated transformation that we did not address, such as binary vector origin of
replication or T-DNA length. Furthermore, as novel *Agrobacterium*-independent technologies to
introduce nucleic acid and proteins into plant cells mature (47,48), their variability should also be
estimated. It is critically important to establish whether our values of variance are broadly
applicable, and future work should prioritize cross-lab validation.
356
Even in this simple context, variation in *N. benthamiana* transient expression depends strongly
on a host of inputs, and attempts to normalize and mitigate this variation must be individually
validated. As complexity increases in synthetic biology experimental designs, this
unpredictability in variation and in confidence will likely be compounded. Our results show that
the choice and validation of normalization methods and data collection are critical for *N.*
benthamiana to be a reliable platform for applications such as reconstituting complex metabolic
pathways and designing large genetic circuits. As we have thoroughly demonstrated, careful,
purposeful experimental design and interpretation are paramount to ensure robust and
reproducible results when using this essential technique for synthetic plant biology.

Methods

Media, chemicals, and culture conditions

Routine bacterial cultures were grown in Luria-Bertani (LB) Miller medium (BD Biosciences,
USA). *E. coli* was grown at 37 °C, while *A. tumefaciens* was grown at 30 °C unless otherwise
noted. Cultures were supplemented with kanamycin (50 mg/L, Sigma Aldrich, USA),
gentamicin (30 mg/L, Fisher Scientific, USA), or spectinomycin (100mg/L, Sigma Aldrich,
USA), when indicated. All other compounds unless otherwise specified were purchased through
Sigma Aldrich.

Bacterial strain and plasmid construction

All bacterial strains and plasmids used in this work are listed in Supplementary Table 1. All strains and plasmids created in this work are viewable through the public instance of the Joint BioEnergy Institute (JBEI) registry. All strains and plasmids created in this work can be requested from the strain archivist at JBEI with a signed material transfer agreement. Plasmids were assembled by Gibson assembly using standard protocols. Plasmids were routinely isolated using the QIAprep spin miniprep kit (Qiagen) and all primers were purchased from Integrated DNA Technologies (IDT). Plasmid sequences were verified using whole-plasmid sequencing (Primordium Labs). *Agrobacterium* was routinely transformed by electroporation as described previously using a 1-mm cuvette and a 2.4-kV, 25-µF, 200-Ω pulse (49).

N. benthamiana growth conditions

Wild type *N. benthamiana* (LAB accession) plants were obtained from the in-house seed bank at the Joint BioEnergy institute. All seedlings and plants were grown at 25 °C in 60% humidity under long-day conditions (16 h of light, 8 h of darkness) of 150 µmol m^{-2} s^{-1} photosynthetically active radiation (PAR; wavelength: 400–700 nm).

As many pots as needed of Sungro Sunshine mix #4 (aggregate plus) were wetted by running excess tap water from above and allowing it to drain through. The soil was then topped with a layer of topsoil that was thoroughly wetted with a spray bottle. Using a spatula, a pinch of seeds were sprinkled evenly over the soil. Ideally, 50-100 good seedlings per pot germinate. If seeds are sown too densely, the seedlings are smaller, and having more sparse, healthy seedlings results in superior plant quality at later stages. All seedling pots were placed in a flat with 1L tapwater and covered with a hood, vents closed, in the growth room. If algal or cyanobacterial contamination grows in the water, reduce the volume of water added to the tray.

After one week, flats (as many as needed) were filled with 18 NxN pots. These pots were filled to the top with Sungro Sunshine mix #4 (aggregate plus), supplemented with 1.5 Tbsp Osmocote (14-14-14) pellets per 4L soil, and then 3L of tap water was added to the flat. The soil was allowed to soak up the water for 2-3 hours, and then any excess was drained off and the pots broken apart. Small holes were made in each pot's soil, and the germinated seedlings were transplanted into individual pots. The root of the seedling must not be damaged during transplantation. The block of soil from a seedling pot can be removed and placed on its side to facilitate removing seedlings from soil without causing damage. Flats were covered with a hood, vents closed, in the growth room. After one week, all the vents were opened to allow the hood to slowly equilibrate to the ambient conditions in the growth room. Five days later, the hoods were removed. Two days later, each flat was watered with tap water supplemented with Peter's Professional (20-20-20) at 1 tsp per 4L water. After five days, the pots were rearranged into a checkerboard pattern within the flats to maximize the amount of space per plant, decreasing the density from 18 plants per flat to 9 plants per flat. Each flat was also watered with 1L water. Three days later, plants were infiltrated. A highly detailed protocol of *N. benthamiana* growth and care is provided in Supplemental Methods 1.

Agroinfiltration of N. benthamiana

Generated binary vectors were transformed into *A. tumefaciens* strain GV3101 via electroporation (49). BiBi strains were generated by growing a liquid culture of single transformants to saturation, pelleting and washing thrice with ice cold 10% glycerol, resuspending in 100ul water, and electroporating the second binary vector into the cells. Selected transformants were inoculated in liquid media with appropriate selection the night before the experiment. *A. tumefaciens* strains were grown until OD_{600} between 0.8-1.2, centrifuged for 10 min at 4000 g, and resuspended in infiltration buffer (10 mM MgCl2, 10 mM MES, and 200 µM acetosyringone, pH 5.6) to achieve the desired OD_{600}. Cultures were induced for 1 h at room temperature on a rocking shaker. Leaves T4 and T5 of 4-week-old *N. benthamiana* plants were syringe infiltrated with the *A. tumefaciens* suspensions. After infiltration, *N. benthamiana* plants were maintained in the same growth conditions as described above with 1L tap water per tray. Three days post infiltration, 4 leaf discs per infiltrated leaf were hole punched. The leaf discs were placed on 350 µL of water in 96 well microtiter plates and GFP (Ex.λ = 488 nm, Em.λ = 520 nm) and mCherry (Ex.λ = 587 nm, Em.λ = 615 nm) fluorescence were measured using a Synergy 4 microplate reader (Bio-tek).

Statistical analysis

Mixed effects model and Monte Carlo simulations were run using R (v4.2.0) and the following packages: tidyverse (v2.0.0) (50), transport (v0.14.6), car (v3.1.3), lme4 (v1.1.31) (51), and performance (v0.13.0) (52). All other analyses and figures were generated with Python (v3.11.4) and the following packages: jupyterlab (v4.0.3) (53), pandas (v2.0.3), numpy (v1.25.1) (54), seaborn (v0.12.2), matplotlib (v3.7.2), scipy (v1.11.3) (55), and statsmodels (v0.14.2).

Data availability

All code related to this study is publicly available on GitHub (github.com/shih-lab/benthi_variation).

References

1.   Yang S-J, Carter SA, Cole AB, Cheng N-H, Nelson RS. A natural variant of a host RNA-dependent RNA polymerase is associated with increased susceptibility to viruses by Nicotiana benthamiana. Proc Natl Acad Sci USA. 2004 Apr 20;101(16):6297–302.

2.   Kapila J, De Rycke R, Van Montagu M, Angenon G. An Agrobacterium-mediated transient gene expression system for intact leaves. Plant Sci. 1997 Jan;122(1):101–8.

3.   Grimsley N, Hohn B, Hohn T, Walden R. "Agroinfection," an alternative route for viral infection of plants by using the Ti plasmid. Proc Natl Acad Sci USA. 1986 May;83(10):3282–6.

4.   Schöb H, Kunz C, Meins F. Silencing of transgenes introduced into leaves by agroinfiltration: a simple, rapid method for investigating sequence requirements for gene

silencing. Mol Gen Genet. 1997 Nov;256(5):581–5.

5.   Ruiz MT, Voinnet O, Baulcombe DC. Initiation and maintenance of virus-induced gene
silencing. Plant Cell. 1998 Jun;10(6):937–46.

6.   Chakrabarty R, Banerjee R, Chung S-M, Farman M, Citovsky V, Hogenhout SA, et al.
PSITE vectors for stable integration or transient expression of autofluorescent protein
fusions in plants: probing Nicotiana benthamiana-virus interactions. Mol Plant Microbe
Interact. 2007 Jul;20(7):740–50.

7.   Irmisch S, Jo S, Roach CR, Jancsik S, Man Saint Yuen M, Madilao LL, et al. Discovery of
UDP-Glycosyltransferases and BAHD-Acyltransferases Involved in the Biosynthesis of the
Antidiabetic Plant Metabolite Montbretin A. Plant Cell. 2018 Aug;30(8):1864–86.

8.   Wood CC, Petrie JR, Shrestha P, Mansour MP, Nichols PD, Green AG, et al. A leaf-
based assay using interchangeable design principles to rapidly assemble multistep
recombinant pathways. Plant Biotechnol J. 2009 Dec;7(9):914–24.

9.   Martin LBB, Kikuchi S, Rejzek M, Owen C, Reed J, Orme A, et al. Complete biosynthesis
of the potent vaccine adjuvant QS-21. Nat Chem Biol. 2024 Apr;20(4):493–502.

10. Calgaro-Kozina A, Vuu KM, Keasling JD, Loqué D, Sattely ES, Shih PM. Engineering
plant synthetic pathways for the biosynthesis of novel antifungals. ACS Cent Sci. 2020
Aug 26;6(8):1394–400.

11. Nett RS, Lau W, Sattely ES. Discovery and engineering of colchicine alkaloid
biosynthesis. Nature. 2020 Aug;584(7819):148–53.

12. Jores T, Tonnies J, Wrightsman T, Buckler ES, Cuperus JT, Fields S, et al. Synthetic
promoter designs enabled by a comprehensive analysis of plant core promoters. Nat
Plants. 2021 Jun 3;7(6):842–55.

13. Hummel NFC, Zhou A, Li B, Markel K, Ornelas IJ, Shih PM. The trans-regulatory
landscape of gene networks in plants. Cell Syst. 2023 Jun 21;14(6):501-511.e4.

14. Zhou A, Kirkpatrick LD, Ornelas IJ, Washington LJ, Hummel NFC, Gee CW, et al. A suite
of constitutive promoters for tuning gene expression in plants. ACS Synth Biol. 2023 May
19;12(5):1533–45.

15. Brophy JAN, Magallon KJ, Duan L, Zhong V, Ramachandran P, Kniazev K, et al.
Synthetic genetic circuits as a means of reprogramming plant roots. Science. 2022 Aug
12;377(6607):747–51.

16. Belcher MS, Vuu KM, Zhou A, Mansoori N, Agosto Ramos A, Thompson MG, et al.
Design of orthogonal regulatory systems for modulating gene expression in plants. Nat
Chem Biol. 2020 Aug;16(8):857–65.

17. Wu F-H, Shen S-C, Lee L-Y, Lee S-H, Chan M-T, Lin C-S. Tape-Arabidopsis Sandwich -
a simpler Arabidopsis protoplast isolation method. Plant Methods. 2009 Nov 24;5:16.

18. Yoo S-D, Cho Y-H, Sheen J. Arabidopsis mesophyll protoplasts: a versatile cell system
for transient gene expression analysis. Nat Protoc. 2007;2(7):1565–72.

19. Morey K, Khakhar A. Exploring the frontier of rapid prototyping technologies for plant

synthetic biology and what could lie beyond. New Phytol. 2024 May;242(3):903–8.

20. Bally J, Jung H, Mortimer C, Naim F, Philips JG, Hellens R, et al. The Rise and Rise of
Nicotiana benthamiana: A Plant for All Reasons. Annu Rev Phytopathol. 2018 Aug
25;56:405–26.

21. Lux MW, Strychalski EA, Vora GJ. Advancing reproducibility can ease the "hard truths"
of synthetic biology. Synth Biol (Oxf). 2023 Oct 28;8(1):ysad014.

22. Zhou X, Mehta S, Zhang J. Genetically encodable fluorescent and bioluminescent
biosensors light up signaling networks. Trends Biochem Sci. 2020 Oct;45(10):889–905.

23. Khosla A, Rodriguez-Furlan C, Kapoor S, Van Norman JM, Nelson DC. A series of dual-
reporter vectors for ratiometric analysis of protein abundance in plants. Plant Direct. 2020
Jun 21;4(6):e00231.

24. Rattan R, Alamos S, Szarzanowicz M, Markel K, Shih PM. Predictable modulation of plant
root development using engineered cytokinin regulators. BioRxiv. 2024 Dec 6;

25. Morffy N, Van den Broeck L, Miller C, Emenecker RJ, Bryant JA, Lee TM, et al.
Identification of plant transcriptional activation domains. Nature. 2024
Aug;632(8023):166–73.

26. Xu P, Lai S, Yin B, Yao C, Gu X, Huang J, et al. Interplay between Agrobacterium T-DNA
and backbone DNA in transgenic plant cells. Transgenic Res. 2024 Dec 30;34(1):1.

27. Lopatkin AJ, Collins JJ. Predictive biology: modelling, understanding and harnessing
microbial complexity. Nat Rev Microbiol. 2020 Sep;18(9):507–20.

28. Covert MW, Gillies TE, Kudo T, Agmon E. A forecast for large-scale, predictive biology:
Lessons from meteorology. Cell Syst. 2021 Jun 16;12(6):488–96.

29. Breyne P, Gheysen G, Jacobs A, Van Montagu M, Depicker A. Effect of T-DNA
configuration on transgene expression. Mol Gen Genet. 1992 Nov;235(2–3):389–96.

30. Bhattacharyya MK, Stermer BA, Dixon RA. Reduced variation in transgene expression
from a binary vector with selectable markers at the right and left T-DNA borders. The
Plant Journal [Internet]. 1994 Dec [cited 2025 Jan 26]; Available from:
https://onlinelibrary.wiley.com/doi/10.1046/j.1365-313X.1994.6060957.x

31. Bashandy H, Jalkanen S, Teeri TH. Within leaf variation is the largest source of variation
in agroinfiltration of Nicotiana benthamiana. Plant Methods. 2015 Oct 14;11:47.

32. Vazquez-Vilar M, Quijano-Rubio A, Fernandez-Del-Carmen A, Sarrion-Perdigones A,
Ochoa-Fernandez R, Ziarsolo P, et al. GB3.0: a platform for plant bio-design that
connects functional DNA elements with associated biological data. Nucleic Acids Res.
2017 Feb 28;45(4):2196–209.

33. Moyle RL, Carvalhais LC, Pretorius L-S, Nowak E, Subramaniam G, Dalton-Morgan J, et
al. An optimized transient dual luciferase assay for quantifying microrna directed
repression of targeted sequences. Front Plant Sci. 2017 Sep 20;8:1631.

34. Cazzonelli CI, Velten J. Analysis of RNA-mediated gene silencing using a new vector
(pKNOCKOUT) and an in plantaAgrobacterium transient expression system. Plant Mol

Biol Rep. 2004 Dec;22(4):347–59.

35.   Hellens RP, Allan AC, Friel EN, Bolitho K, Grafton K, Templeton MD, et al. Transient
expression vectors for functional genomics, quantification of promoter activity and RNA
silencing in plants. Plant Methods. 2005 Dec 18;1:13.

36.   Thompson M, Kirkpatrick LD, Geiselman GM, Waldburger LM, Pearson AN,
Szarzanowicz M, et al. Genetically refactored Agrobacterium-mediated transformation.
BioRxiv. 2023 Oct 13;

37.   Huang YW, Hu CC, Tsai CH, Lin NS, Hsu YH. Chloroplast Hsp70 Isoform Is Required for
Age-Dependent Tissue Preference of Bamboo mosaic virus in Mature Nicotiana
benthamiana Leaves. Mol Plant Microbe Interact. 2017 Aug;30(8):631–45.

38.   Sheludko YV, Sindarovska YR, Gerasymenko IM, Bannikova MA, Kuchuk NV.
Comparison of several Nicotiana species as hosts for high-scale Agrobacterium-mediated
transient expression. Biotechnol Bioeng. 2007 Feb 15;96(3):608–14.

39.   Alamos S, Szarzanowicz MJ, Thompson MG, Stevens DM, Kirkpatrick LD, Dee A, et al.
Quantitative dissection of Agrobacterium T-DNA expression in single plant cells reveals
density-dependent synergy and antagonism. Nat Plants. 2025 May 12;

40.   Szarzanowicz MJ, Waldburger LM, Busche M, Geiselman GM, Kirkpatrick LD, Kehl AJ, et
al. Binary vector copy number engineering improves Agrobacterium-mediated
transformation. Nat Biotechnol. 2024 Nov 4;

41.   Carlson ED, Rajniak J, Sattely ES. Multiplicity of the Agrobacterium Infection of Nicotiana
benthamiana for Transient DNA Delivery. ACS Synth Biol. 2023 Aug 18;12(8):2329–38.

42.   Lopez-Agudelo JC, Goh F-J, Tchabashvili S, Huang Y-S, Huang C-Y, Lee K-T, et al.
Rhizobium rhizogenes A4-derived strains mediate hyper-efficient transient gene
expression in Nicotiana⬚benthamiana and other solanaceous plants. Plant Biotechnol J.
2025 Apr 9;

43.   Garcia-Perez E, Diego-Martin B, Quijano-Rubio A, Moreno-Giménez E, Selma S, Orzaez
D, et al. A copper switch for inducing CRISPR/Cas9-based transcriptional activation tightly
regulates gene expression in Nicotiana benthamiana. BMC Biotechnol. 2022 Mar
24;22(1):12.

44.   Kallam K, Moreno-Giménez E, Mateos-Fernández R, Tansley C, Gianoglio S, Orzaez D,
et al. Tunable control of insect pheromone biosynthesis in Nicotiana benthamiana. Plant
Biotechnol J. 2023 Jul;21(7):1440–53.

45.   Larsen B, Hofmann R, Camacho IS, Clarke RW, Lagarias JC, Jones AR, et al.
Highlighter: An optogenetic system for high-resolution gene expression control in plants.
PLoS Biol. 2023 Sep 21;21(9):e3002303.

46.   Dürrenberger F, Crameri A, Hohn B, Koukolíková-Nicola Z. Covalently bound VirD2
protein of Agrobacterium tumefaciens protects the T-DNA from exonucleolytic
degradation. Proc Natl Acad Sci USA. 1989 Dec;86(23):9154–8.

47.   Demirer GS, Zhang H, Matos JL, Goh NS, Cunningham FJ, Sung Y, et al. High aspect

ratio nanomaterials enable delivery of functional genetic material without DNA integration
in mature plants. Nat Nanotechnol. 2019 May;14(5):456–64.

48.   Demirer GS, Zhang H, Goh NS, González-Grandío E, Landry MP. Carbon nanotube-
mediated DNA delivery without transgene integration in intact plants. Nat Protoc. 2019
Oct;14(10):2954–71.

49.   Kámán-Tóth E, Pogány M, Dankó T, Szatmári Á, Bozsó Z. A simplified and efficient
Agrobacterium tumefaciens electroporation method. 3 Biotech. 2018 Mar;8(3):148.

50.   Wickham H, Averick M, Bryan J, Chang W, McGowan L, François R, et al. Welcome to
the tidyverse. JOSS. 2019 Nov 21;4(43):1686.

51.   Bates D, Mächler M, Bolker B, Walker S. Fitting linear mixed-effects models using lme4. J
Stat Softw. 2015;67(1):1–48.

52.   Lüdecke D, Ben-Shachar M, Patil I, Waggoner P, Makowski D. performance: An R
Package for Assessment, Comparison and Testing of Statistical Models. JOSS. 2021 Apr
21;6(60):3139.

53.   Perez F, Granger BE. IPython: A System for Interactive Scientific Computing. Comput Sci
Eng. 2007;9(3):21–9.

54.   van der Walt S, Colbert SC, Varoquaux G. The NumPy Array: A Structure for Efficient
Numerical Computation. Comput Sci Eng. 2011 Mar;13(2):22–30.

55.   Virtanen P, Gommers R, Oliphant TE, Haberland M, Reddy T, Cournapeau D, et al. SciPy
1.0: fundamental algorithms for scientific computing in Python. Nat Methods. 2020
Mar;17(3):261–72.

Acknowledgements

593 We would like to thank Dr. Tyler Backman and Dr. Hector Garcia-Martin for their invaluable
594 advice on statistics, and to all members of the Shih lab for their helpful discussions. We also
595 thank Konami for facilitating discussion of expression cassette orientation via Dance Dance
596 Revolution artwork.
597
BioRender was used to generate figures 1 (https://BioRender.com/c5zadct), 2
(https://BioRender.com/2vhubua), S1 (https://biorender.com/82xc4es), S6
(https://biorender.com/4ila5ts), and S7 (https://biorender.com/dcf88wn). This material is based
upon work supported by the National Science Foundation Graduate Research Fellowship under
Grant No. DGE 2146752. This work was part of the DOE Joint BioEnergy Institute
(https://www.jbei.org) supported by the U. S. Department of Energy, Office of Science, Office of
Biological and Environmental Research, supported by the U.S. Department of Energy, Energy
Efficiency and Renewable Energy, Bioenergy Technologies Office, through contract DE-AC02-
05CH11231 between Lawrence Berkeley National Laboratory and the U.S. Department of
Energy. The funders had no role in manuscript preparation or the decision to publish. The views
and opinions of the authors expressed herein do not necessarily state or reflect those of the

Author information

Authors and affiliations

Joint BioEnergy Institute, 5885 Hollis Street, Emeryville, CA 94608, USA.
Sophia N. Tang, Matthew J. Szarzanowicz, Sasilada Sirirungruang, Amy Lanctot, Liam D. Kirkpatrick, Krista Drako, Simón Álamos, Lucas M. Waldburger, Adela Cheng, Shuying Liu, Mitchell G. Thompson, Patrick M. Shih

Environmental Genomics and Systems Biology Division, Lawrence Berkeley National Laboratory, Berkeley, California, USA
Sophia N. Tang, Matthew J. Szarzanowicz, Sasilada Sirirungruang, Amy Lanctot, Liam D. Kirkpatrick, Krista Drako, Simón Álamos, Lucas M. Waldburger, Shuying Liu, Mitchell G. Thompson, Patrick M. Shih

Department of Molecular and Cell Biology, University of California, Berkeley, CA 94720, USA
Sophia N. Tang, Krista Drako

Department of Plant and Microbial Biology, University of California, Berkeley, CA 94720, USA
Matthew J. Szarzanowicz, Sasilada Sirirungruang, Amy Lanctot, Liam D. Kirkpatrick, Simón Álamos, Patrick M. Shih

Department of Bioengineering, University of California, Berkeley, California, USA
Lucas M. Waldburger

Biological Systems & Engineering Division, Lawrence Berkeley National Laboratory, Berkeley, CA 94720, USA
Lucas M. Waldburger

650 **Department of Nutritional Sciences and Toxicology, University of California, Berkeley, CA**
651 **94720, USA**
652 Adela Cheng
653
Innovative Genomics Institute, Berkeley, California, USA
Patrick M. Shih

Contributions

P.M.S. and M.G.T. conceptualized the initial study. M.G.T., S.A., L.M.W., and S.N.T. designed
experiments. S.N.T., K.D., and A.C. generated plasmids and strains. S.N.T, S.S., L.D.K., S.L.,
M.G.T., M.J.S., K.D., and A.L. performed experiments. M.J.S. performed mixed-effect modeling
and Monte Carlo simulations, and S.N.T. completed all other data analysis. S.N.T. wrote the
draft manuscript. All authors discussed the results, reviewed the article, and approved the final
article.

Corresponding authors

Correspondence to Patrick M. Shih or Mitchell G. Thompson.

Ethics declarations

M.G.T., M.J.S., and P.M.S. have financial interest in BasidioBio.
667

Figure 1. Longitudinal assessment of transient expression identifies sources of
variability in *N. benthamiana* fluorescence assays. **A)** Schematic of the transient
fluorescence assay. *N. benthamiana* is agroinfiltrated with an *A. tumefaciens* strain at an OD of
0.5. The binary vector contains, from left to right border, CaMV35S2:nptII and
PCM2:eGFP:At_Ubq3 oriented divergently in the T-DNA. Discs are collected from leaves 3 days
post-infiltration and measured in a plate reader. **B)** Distributions of GFP fluorescence for 15
experimental replicates, n=64 discs, except for 2022.06.06 and 2022.06.13 which are n=40.
Dates of sample collection and measurement are written YYYY.MM.DD. **C)** Boxplots for each
plant on the dates with the lowest (2022.07.25) and highest (2023.05.30) mean raw
fluorescence show the median and interquartile range (IQR), and whiskers show the range of
data within 1.5 IQR. Scatterplots show T4 discs in white and T5 discs in black. **D)** Percent
contributions to observed variability as calculated from a mixed-effects model. **E)** Illustrations of
sources of variability.
682

Figure 2. Comparison of three discrete methods to deliver normalizing reporters. **A)** Left, categories of delivery methods: unnormalized (green or magenta), co-infiltration (yellow), same T-DNA (pink), and BiBi (blue). Center, distributions of eGFP and mCherry fluorescence, n=96 leaf discs per row. Right, cartoons showing the binary vector origin of replication, resistance marker, and orientations of FP expression cassettes in the T-DNA. All binary vector cartoons are read from left to right: ori, left border, T-DNA, right border. Origins of replication are pVS1 (diamond, black), BBR1 (circle, teal), and pSa (square, gray). Circles enclosing an X represent tOcs, a 722bp spacer in between the two expression cassettes. **B)** Plant coefficients of variation (CV), as calculated from the 8 discs per plant, when eGFP is treated as the reporter. All values are eGFP/mCherry CV except for scheme 1, which is GFP CV. **C)** Plant CVs when mCherry is treated as the reporter. All values are mCherry/eGFP CV except for scheme 2, which is

mCherry CV. Normalization scheme IDs match across all subpanels. For **B)** and **C)**, an independent, one-tailed Welch's t-test and a Bonferroni correction were conducted to determine whether the reporter/normalizer CV of a scheme is significantly less than the reporter CV for the corresponding unnormalized scheme. Asterisks indicate p-values: * < 0.05, ** < 0.01. Total OD infiltrated in all schemes is 0.5. ODs of co-infiltrated strains are 0.25 each. In the legend, "&" indicates co-infiltration, arrows indicate the direction of an expression cassette, "+" indicates BiBi, and GFP is abbreviated to "G" and mCherry to "R". Leaves T4 and T5 of six four-week-old *N. benthamiana* plants were infiltrated in each experimental replicate, and four discs were collected from each infiltrated leaf.

Figure 3. OD$_{600}$ of two co-infiltrated strains affect transgene expression strength but not variability. Raw fluorescence signal from **A)** eGFP and **B)** mCherry for all OD$_{600}$ combinations. Darker hues indicate increasing OD$_{600}$ of the competing strain, which carries the other FP. Error bars show the standard error. Matrix of all OD$_{600}$ combinations' CV of **C)** eGFP fluorescence and **D)** ratio of eGFP/mCherry. Each condition was infiltrated into leaves T4 and T5 of six plants. Four discs were collected from each leaf for a total of n=48 leaf discs.

Figure 4. Promoters of co-infiltrated transgenes determine the efficacy of normalization.
eGFP and mCherry were driven by PCL2, PCM2, or PCH5 for a total of six unique binary
vectors. Log$_2$ raw fluorescence from **A)** eGFP and **B)** mCherry for all promoter combinations.

Darker hues indicate stronger promoter of the competing strain. Error bars show the standard
error. Matrix of all promoter combinations' CV of **C)** eGFP fluorescence and **D)** ratio of
eGFP/mCherry. Log$_2$ raw fluorescence signal from **E)** eGFP and **F)** mCherry for all promoter
combinations when instead the mCherry binary vectors are driven by PCL1, PCM1, or PCH4.
Darker hues indicate stronger promoter of the competing strain. Error bars show the standard
error. Matrix of all promoter combinations' CV of **G)** eGFP fluorescence and **H)** ratio of
eGFP/mCherry. Conditions from the "Same Promoters" and "Different Promoters" experiments
were infiltrated into leaves T4 and T5 of six plants, and four discs were collected from each leaf
for a total of n=48 leaf discs. The experiments were performed independently on separate days.
726

727
Figure 5. Promoter choice affects variation between experimental replicates. A)
Cumulative density functions of PCL2:eGFP alone or normalized by mCherry driven by PCL2,
PCM1, or PCM2. Left: eGFP, right: eGFP/mCherry. Each experimental replicate is a unique
color. The black line is the CDF for the pooled data of all six replicates. The p-values of one-
sample Kolmogorov-Smirnov tests appear to the right of each CDF, colored by replicate.
Asterisks indicate p-values: * < 0.05, ** < 0.01, *** < 0.001, and ns = not significant. **B)** Absolute
value of every experimental replicate's D, the greatest vertical distance between the CDF a
given replicate (color) and the pooled CDF (black) from **A)**. White boxplots indicate D values for

eGFP CDFs and gray boxplots for eGFP/mCherry CDFs. **C)** CVs of the six experimental
replicates for each condition. As in **B)**, white boxplots indicate eGFP and gray eGFP/mCherry.
Every condition was infiltrated into leaves T4 and T5 of four plants, and four discs were
collected from each leaf in all six experimental replicates for a total of n=192 leaf discs.
740

Figure 6. Modeling variation of transgene expression in 4-week-old *N. benthamiana* plants. A) Per-plant CV of unnormalized GFP calculated from data compiled across many years and independent experiments. CV for a plant is calculated using all discs derived from that plant, regardless of leaf. The number of plants included appears above the violin plots. **B)**

Cumulative density function of per-plant, unnormalized eGFP CVs. Solid gray, empirical data
from 32 batches of at least 30 plants. Dotted black, Monte Carlo simulated data. **C)** Minimum
number of simulated plants needed to detect a given effect size with the CV of unnormalized
EHA105, unnormalized GV3101, or optimally normalized GV3101 assuming 95% accuracy and
statistical significance ($p<0.05$). **D)** Exponential regression fit to the simulated data in **C)**.
Dashed horizontal line indicates the arbitrary 50 plant cap, and dotted vertical lines indicate the
smallest detectable effect size with 50 plants. Orange, EHA105. Gray, unnormalized GV3101.
Teal, optimally normalized GV3101.