Title: Causes and consequences of experimental variation in *Nicotiana benthamiana*
transient expression

Authors: Sophia N. Tang[1,2,3], Matthew J. Szarzanowicz[1,2,4], Amy Lanctot[1,2,4], Sasilada
Sirirungruang[1,2,4], Liam D. Kirkpatrick[1,2,4], Krista Drako[1,2,3], Simón Álamos[1,2,4], Lyurui Cheng[1,2,7],
Lucas M. Waldburger[1,2,5,6], Shuying Liu[1,2], Lena Huang[1,2], Sami Kazaz[1,2], Emine Akyuz
Turumtay[1], Edward Baidoo[1], Aymerick Eudes[1,2], Mitchell G. Thompson[1,2]*, Patrick M. Shih[1,2,4,8]*

Affiliations:
1 Joint BioEnergy Institute, 5885 Hollis Street, Emeryville, CA 94608, USA.
2 Environmental Genomics and Systems Biology Division, Lawrence Berkeley National
Laboratory, Berkeley, California, USA
3 Department of Molecular and Cell Biology, University of California, Berkeley, CA 94720, USA
4 Department of Plant and Microbial Biology, University of California, Berkeley, CA 94720, USA
5 Department of Bioengineering, University of California, Berkeley, California, USA
6 Biological Systems & Engineering Division, Lawrence Berkeley National Laboratory, Berkeley,
CA 94720, USA
7 Department of Nutritional Sciences and Toxicology, University of California, Berkeley, CA
94720, USA
8 Innovative Genomics Institute, Berkeley, California, USA

* Correspondence to Patrick M. Shih (pmshih@berkeley.edu) or Mitchell G. Thompson
(mthompson@lbl.gov)

Abstract

Infiltration of *Agrobacterium tumefaciens* into *Nicotiana benthamiana* has become a
foundational technique in plant biology, enabling efficient delivery of transgenes *in planta* with
technical ease, robust signal, and relatively high throughput. Despite transient expression's
prevalence in disciplines such as synthetic biology, little work has been done to describe and
address the variability inherent in this system, a concern for experiments that rely on highly
quantitative readouts. In a comprehensive analysis of *N. benthamiana* agroinfiltration
experiments, we model sources of variability that affect transient expression. Our findings
emphasize the need to validate normalization methods under the specific conditions of each
study, as distinct normalization schemes do not always reduce variation either within or between
experiments. Using a dataset of 1,915 plants collected over three years, we develop a model of
variation in *N. benthamiana* transient expression, using power analysis to determine the number
of individual plants required for a given effect size. Drawing on our longitudinal data, these
findings inform practical guidelines for minimizing variability through strategic experimental
design and power analysis, providing a foundation for more robust and reproducible use of *N.*
benthamiana in quantitative plant biology and synthetic biology applications.

Introduction

Nicotiana benthamiana is a workhorse of plant molecular biology due to the high efficiency of transient *Agrobacterium*-mediated transformation, enabling researchers to assess gene function and observe phenotypes within days, as opposed to the months it typically takes for stable transformations [1–3]. *N. benthamiana*'s capacity for transient expression precipitated its use to study diverse aspects of cell and molecular biology [4], including virus-induced gene silencing [5], subcellular protein localization [6], biosynthetic pathway discovery and engineering [7–11], and gene regulation [12–16]. Given the lengthy time required to generate stable transformants and the technical challenges inherent in protoplast transformation [17,18], there are few viable *in planta* alternatives to *N. benthamiana* transient expression, leading to its primacy as the model of plant synthetic biology [19,20].

Despite its widespread use, *N. benthamiana* lacks substantial efforts to assess error and reproducibility, like many other systems routinely leverage in synthetic biology [21]. Nor is there a standardized way to design transient assays, and as a result, experimental design varies from publication to publication [13,22–26]. These factors obstruct our ability to model and predict variation in this model system, which is crucial for effective experimental design as has been demonstrated in the closely related *Nicotiana tabacum* [27]. Limited quantitative understanding of biology has routinely impeded genetic engineers in developing predictable and reliable bioproducts [28], in comparison to the precision associated with traditional engineering disciplines [29]. For plant synthetic biology to truly be an engineering discipline, we require robust evaluation of variability and its effects on our design goals.

The variability of transient expression in *N. benthamiana* in highly powered and longitudinal experiments has never been rigorously evaluated. Without robust statistics, researchers risk designing underpowered experiments and overlooking subtle effects. While there have been attempts to determine transgene expression variability in stable plant lines [30,31], it is unclear whether these findings translate to transient expression. Transient expression variability in *N. benthamiana* has been primarily studied using luciferase reporters, but these studies are plagued by small sample size, lack of biological replicates, and the high noise inherent to the luciferase assay [32–34]. Most methods used to reduce variability in transient expression experiments have relied on using a second reporter (hereafter referred to as a normalizer) for ratiometric normalization [13,22–24,35,36]. In this approach, every tested experimental construct is co-delivered with the same constitutively expressed normalizer, under the assumption that any variation in the experimental construct's expression that tracks with variation in the normalizer's expression must not be due to the sequence of the reporters but rather other variables. Although this approach is widely used, whether normalizing transgene expression actually reduces and controls for variation has not been systematically examined.

Here, we use fluorescence reporter assays to systematically determine the sources of variation of transient expression in *N. benthamiana* leaves. Using a large sample size (>1900 plants) across multiple years of independent experiments, we comprehensively capture transgene expression variation observed between plants and between independent experiments in order to model sources of error. We also complete the first systematic comparison of methods to

normalize transgene expression and demonstrate whether and to what extent these methods
reduce variability. Finally, we suggest best practices to mitigate variation through purposeful
experimental design and statistical power analysis.

Results

Categorizing the sources of transient expression variation

To determine sources of variability in *N. benthamiana* transient expression, we analyzed
previously published data wherein the same GFP reporter (vector 1 of Supplementary Table 1)
was agroinfiltrated in 15 independent experimental replicates using *A. tumefaciens*
GV3101::pMP90 (hereafter GV3101) (Fig 1A, Fig. S1) [37]. The reporter strain was infiltrated
distal to the petiole of the fourth and fifth leaf from the top (leaves T4 and T5) of *N.*
benthamiana, and three days post-infiltration, four discs were collected from each infiltrated leaf.
All plants germinated at the same time and which experienced identical growing conditions and
care are referred to as belonging to the same batch. In this dataset, we observed as much as a
fourfold difference between experimental replicates with the lowest (teal, 2022.07.25) and
highest (orange, 2023.05.30) mean fluorescence (Fig. 1B, 1C). Using a mixed-effects model, we
attributed nearly all of the variation to fluctuations in batch-level mean GFP (23.8%) and GFP
standard deviation (15.6%); plant-level mean GFP (9.9%) and GFP standard deviation (28.2%)
within a given batch; and disc-level GFP standard deviation (22.3%) within a given plant (Fig.
1D). Though the precise percent contributions of these components varies between
experimental replicates (Fig. S2), they are nonetheless all major sources of variation that should
be accounted for in experimental design. The underlying factors of these sources of variation
may be batch effects across the fifteen experimental replicates' plants, non-homogenous
growing conditions, and varying leaf cellular ages (Fig. 1E) [38,39].

107
Since the experimental conditions across these 15 replicates were identical, we separately
investigated other potential contributors to variation that were controlled for in that experiment.
In other independent experiments, we assessed the effects of 96-well water volume, plant age,
leaf infiltration site, and collection time (both time of day and time elapsed between collection
and measurement). We found that water volume and plant age are negatively correlated with
fluorescence (Fig. S3, S4), infiltration site significantly affects transgene expression in a leaf-
and strain-dependent manner (Fig. S5), and collection time has no effect (Fig. S6, S7). Minor
changes to water volume (*e.g.*, evaporation, pipetting error) may have contributed to residual
error in the data from Fig. 1B. However, all plants in this study were 4 weeks old at the time of
infiltration and infiltrated distal to the petiole, so no variation can be attributed to differences in
age or infiltration site. With these additional experiments, we have both accounted for the
majority of variation in our model and now ruled out other potential sources.

120
To demonstrate that these findings are generalizable beyond fluorescent proteins, we also
quantified variation for two different metabolic pathways to produce betalain (3 experimental
replicates) and 2-pyrone-4,6-dicarboxylic acid (PDC) (4 experimental replicates) [40,41]. The three
enzymes of the betalain pathway were co-infiltrated on three separate T-DNAs or infiltrated as

one T-DNA separated by self-cleaving T2A peptides (RUBY reporter) [42]. The absorbance of one
of the three betalain experimental replicates was significantly different from the other two for
both methods of delivery, and the greatest fold change between two replicates was about
twofold for co-infiltration and about fourfold for T2A, on par with the maximal difference
observed for the GFP reporter (Fig. S8, Fig. 1B-D). For the PDC experimental replicates, yields
were not significantly different after Bonferroni correction, but there was a ~20% difference
between the highest and lowest yields (Fig. S8). Multi-step metabolic pathway yields are subject
to enzyme subcellular localization, substrate diffusion, and the pathway's underlying dynamics,
which may mask or exacerbate the effects of variable transgene expression. There are
considerable variations in yields, specific to each pathway, and the trends observed in FP
fluorescence likely translate to the variability of each component transgene in a metabolic
pathway.

Multiple methods of co-delivering a normalizing transgene can decrease expression variability

Normalizing the expression level of a transgene of interest to that of another independent
transgene is thought to control for any sources of variability independent of the tested
sequences, since they should impact expression of the two transgenes equally. This practice
should theoretically reduce variation both within and between experiments. However, there are
many different conceivable ways the two transgenes could be delivered in *N. benthamiana*, and
indeed normalization schemes differ between publications [13,22–26].

We systematically compared normalization schemes for transient expression using fluorescent
protein (FP) reporters, whose gene expression is correlated to an easily measured phenotype,
fluorescence. The normalized readout of the assay is the ratio of eGFP and mCherry
fluorescence measured in a leaf disc. Designating FPs as the reporter or normalizer is arbitrary.
The two expression cassettes were co-delivered in three different ways. First, we co-infiltrated a
mix of two different strains of *A. tumefaciens*, each with a binary vector containing one FP
(schemes 3-7). Second, we infiltrated single *A. tumefaciens* strains carrying binary vectors with
a T-DNA containing both cassettes in every possible orientation, both relative to one another
and to the left and right borders (schemes 8-15). Third, we infiltrated a "BiBi" strain [43], wherein a
single strain of *A. tumefaciens* carries two binary vectors, each with a unique FP gene and ori
(schemes 16-19). All FP genes are driven by the medium strength PCM2 constitutive promoter
derived from the *A. thaliana* HTR5 histone gene (AT4G40040) and are terminated by the *A.
thaliana* Ubq3 terminator (Supplementary Table 1, Fig. S1) [14]. These 17 methods of co-delivery
were compared to delivering eGFP (scheme 1) or mCherry (scheme 2) alone in two
independent experimental replicates.

Co-delivery methods have obvious ramifications for basal transgene expression levels that are
consistent between both experimental replicates (Fig. S9). For both co-infiltrations and BiBis,
the low copy number origin, pSa, consistently produced much lower fluorescence compared to
the higher copy number origins, pVS1 or BBR1 (Fig. 2A) [44]. For the stacked, two-cassette T-
DNAs, despite sharing the same pVS1 origin and backbone, these schemes yielded up to eight-
fold difference in fluorescence depending on the cassettes' relative orientations (Fig. 2A, S10).

Convergent, tandem, and divergent cassettes produced the strongest, intermediate, and
weakest expression, respectively (Fig. S10). The expression of one gene is affected by the
orientation of the other gene, even when its own position and orientation is maintained, as with
schemes 8 and 10 for eGFP or schemes 9 and 11 for mCherry (Fig. S10). The formation of a
double terminator with the tOcs spacer in some cassettes may be partly responsible for
increased expression, as has been shown with other binary vectors [45], but reversing tOcs does
not consistently increase fluorescence from a doubly terminated cassette over the
corresponding singly terminated cassette (Fig. S10). These results suggest that gene position
relative to the left and right borders and, most importantly, relative orientation strongly affect
expression in multi-gene T-DNAs.
178
Variability, as measured by coefficient of variation (CV) from all leaf discs, decreases for all
normalization schemes (Fig. S11). Furthermore, all schemes have a lower CV for
reporter/normalizer than for reporter (Fig. S12). The more appropriate comparison, however, is
whether a scheme's reporter/normalizer CV is lower than the corresponding unnormalized
control's reporter CV. Most schemes produce an eGFP/mCherry CV that is significantly less
than scheme 1's eGFP CV following a Bonferroni correction (Fig. 2B), but only scheme 3 (co-
infiltration of two pVS1 binary vectors) meets this condition when mCherry is treated as the
reporter (Fig. 2C). Such a test is conservative, and it should be noted that there is at least one
scheme per co-delivery method that reduces CV by >50% compared to the unnormalized
control, whether CV is calculated per plant or pooling all discs together (Fig. 2B, 2C, S11). All
co-delivery methods can offer reductions in variability, thereby increasing the statistical power of
transient expression-based experiments.

Inoculum densities of co-infiltrated strains affect signal strength but not variability

Because the decrease in CV was the greatest and most statistically significant for co-infiltration
of two pVS1 binary vectors (scheme 3), we used this scheme to explore other possible
experimental variables for further reductions in variability. Additionally, co-infiltration is highly
modular, and multiple T-DNAs can efficiently be delivered to each plant cell [43,46]. Since *A.*
tumefaciens strains in *N. benthamiana* leaves can antagonize or synergize with one another
depending on their densities [43], we next determined whether the OD_{600} of each strain in the co-
infiltration affects the resulting CV. Each strain was added to the infiltration mix at an OD_{600} of 0,
0.01, 0.1, 0.5, or 1 and combined with the other strain, also added at the same densities for a
total of 24 possible combinations.
201
Total OD_{600} of the co-infiltrated mix and the OD_{600} of the individual strains do not greatly nor
predictably affect variability, but inoculum density of a strain dictates its total transgene
expression levels. There is no discernible pattern in eGFP CV with respect to the OD_{600} of the
co-infiltrated strains (Fig. 3A), and while all CVs for the ratio of eGFP/mCherry are
comparatively much lower (Fig. 3B), there is no one combination of ODs that is significantly
superior to all others. The same trends are apparent when mCherry is treated as the reporter
and eGFP as the normalizer (Fig. S13). For raw fluorescence, signal begins to saturate beyond
an OD_{600} of 0.5, and when one strain is held at a given OD_{600} while the OD_{600} of another
competing strain increases, fluorescence from the fixed strain decreases (Fig. 3C, 3D). The

primary concern when selecting OD_{600}, then, should be the desired signal strength. Of the
densities tested, 0.1 results in strong, measurable signal and, in agreement with previous
findings, does not saturate the transgene expression capacity of the plant cell [43,46].

Promoter choice determines the utility of normalization

Given that the choice of promoter(s) in transient transformations affects not only signal strength
but also normalization outcomes [14], we more thoroughly explored additional promoter
combinations. All transgenes in this study have hitherto been driven by a medium-strength
promoter, PCM2. From a library of low, medium, and high strength constitutive promoters
characterized by Zhou et al. (annotated as "PCL", "PCM", and "PCH," respectively), we selected
low- and high-strength promoters, PCL2 and PCH5, to test alongside PCM2. eGFP and
mCherry were driven by this same set of three promoters ("same promoter set"). The eGFP
binary vectors were co-infiltrated with the mCherry binary vectors in every possible combination,
and all binary vectors were also infiltrated alone for a total of 15 unique infiltrations.

There is no discernible pattern in eGFP CV for any of the infiltrations (Fig. 4A), but for
eGFP/mCherry, identical promoters driving the two FP genes yield the lowest CVs (Fig. 4B).
Outside of these pairs, normalization does not provide as large reductions in variations, if at all.
In fact, for three nonidentical promoter pairs, their eGFP/mCherry CV is greater than their own
eGFP CV and, for one pair, also greater than the eGFP CV of its corresponding unnormalized
control (Fig. 4A, 4B). Similar observations are true when treating mCherry as the reporter and
eGFP as the normalizer (Fig. S14). Additionally, fluorescence from a given strain diminishes the
stronger a competing strain's FP promoter is, though this trend is weaker for mCherry than it is
for GFP (Fig. 4C, 4D).

We then sought to clarify whether the large decrease in CV when using identical promoters is
due to the similar promoter strengths or to other variables, such as shared trans factors
affecting mRNA abundance. To do so, we generated another set of three binary vectors with
mCherry driven by PCL1, PCM1, and PCH4 ("different promoter set"), which are the promoters
in the library closest in expression to the three already tested. As with the first promoter
combination experiment, the existing eGFP binary vectors were co-infiltrated with the new
mCherry binary vectors in every possible combination, and all six binary vectors were also
infiltrated alone.

With these promoters, both the CVs of eGFP and of eGFP/mCherry follow no apparent pattern
(Fig. 4E, 4F), but the negative trend between fluorescence signals of the two competing strains
holds true, although PCH4 fits poorly in the trend (Fig. 4G, 4H). Several promoter pairs have
higher eGFP/mCherry CVs than their own eGFP CVs and/or the eGFP CV of the unnormalized
control (Fig. 4E, 4F), and vice versa when treating mCherry as the reporter (Fig. S14). Similarity
in promoter strength, then, does not guarantee lower CVs upon normalization. Depending on
the promoters used to normalize, normalization may even worsen the assay's statistical power.
Since the co-delivery comparison and the OD optimization experiments were conducted using
PCM2 for every expression cassette, we had fortuitously picked the best case scenario,
identical promoters. Had we chosen nonidentical promoters, all methods of co-delivery might

not have decreased CV compared to the unnormalized controls. When possible, identical
promoters should be used.

Normalization does not yield reproducibility of absolute quantification

Normalization can serve two distinct purposes: (1) decreasing within-experiment variation to
improve detection of small effect sizes, or (2) decreasing between-experiment variation to allow
for comparison across independent experiments. Mitigating within-experiment variation does not
necessarily have an effect on between-experiment variation. To explore whether normalization
makes comparisons between experimental replicates or to historical data more valid, we co-
infiltrated GFP driven by PCL2 (which yielded the lowest average normalized CVs when driving
both FPs (Fig. 4, S11)) alone and with all six mCherry binary vectors thus far generated in six
independent experimental replicates.

Normalization makes experimental replicates more similar to one another for most, but not all,
promoter pairs (Fig. 5A, 5B, S15). We performed one-sample Kolmogorov-Smirnov tests for
every condition to determine the likelihood that a sample distribution (an individual experimental
replicate) is drawn from a reference distribution (all experimental replicates, pooled). The
cumulative density functions (CDFs) show that sample distributions nearly all become more
comparable to the reference distribution with normalization, *i.e.*, the maximum vertical distance
between the distributions decreases (Fig. 5A, 5B, S15). Replicate-to-replicate, the PCL2/PCM2
promoter pair produces the most consistent distribution of GFP/mCherry values, far more
reproducible than its raw eGFP fluorescences (Fig. 5A, 5B). The sole exception is the
PCL2/PCM1 promoter pair, whose GFP/mCherry CDFs are actually less comparable to the
reference distribution than its GFP CDFs are (Fig. 5A, 5B); there is no singular GFP/mCherry
value that normalization with this promoter pair causes the data to reliably converge upon.
However, this pair and PCL2/PCL2 were the only promoter pairs for which experimental
replicate CVs, representing within-experiment variation, were regularly lower than the
unnormalized control's (Fig. 5C). Additionally, PCL2/PCL2 produces the most consistent
experimental replicate CVs–it alone has an interquartile range of experimental replicate CVs
narrower than the unnormalized control's (Fig. 5C). Decreasing within-experiment variation,
then, is not equivalent to decreasing between-experiment variation. Normalization schemes
must be designed with the desired outcome in mind (minimizing within- and/or between-
experiment variation) and then validated.

Developing a generalizable power analysis framework of N. benthamiana *transient expression*

To evaluate whether normalization meaningfully reduces the number of plants required to detect
a given effect size, we developed a model informed by a large dataset of *N. benthamiana*
transient expression experiments. This dataset comprises 1,813 plants infiltrated with a GFP
reporter over nearly three years, spanning experiments conducted by multiple researchers and
diverse conditions, including different inoculum densities and *Agrobacterium* strains (see
Supplementary Methods 2 for details). Because this dataset accounts for historical variation
across diverse experiments, the simulations trained with it will give conservative estimates of

variance. Individual, well-controlled experiments may yield lower variations, but since the
precise variabilitycannot be known beforehand, erring on the side of excessive rather than
insufficient statistical power is preferable. We calculated the CV for each plant infiltrated with
one of two *Agrobacterium* strains: GV3101 and the hypervirulent EHA105, which displays both
elevated transgene expression and variance compared to GV3101 (Fig. 6A). For batches of
plants for which at least 30 plants were used, we visualized the plant CVs in a CDF (Fig. 6B
solid gray line). With these per-plant and per-experiment CVs, we developed a Monte Carlo
simulation of assay variability, which closely recapitulates the observed distribution of real plant
CVs across the training dataset (Fig. 6B dotted black line; Wasserstein distance = 0.023, mean
ECDF difference = 0.05).
305
We then simulated how variability propagates through an experiment on the per-plant and per-
experiment levels. First, a batch CV is drawn from the empirical distribution shown in Fig. 6A by
randomly generating a percentile and selecting the corresponding CV, which represents the
average CV of plants from a hypothetical experimental replicate. Around this batch-level value,
we simulate CVs of individual plants by incorporating additional noise that approximates within-
experiment heterogeneity between plants. For each simulated plant, eGFP fluorescence data
for eight leaf discs is generated based on the plant CV and a fixed, construct-specific mean
expression. Two conditions are compared with an independent, two-tailed Student's *t*-test at
varying effect size differences and sample sizes (Fig. 6C). For each comparison, 1000
independent experimental replicates are simulated and used to calculate the probability of a
successful comparison.
317
This approach enables us to estimate the minimum number of plants needed to reliably (>95%
of simulations) and significantly ($p < 0.05$) distinguish between two constructs. From these
calculations of plants needed for given effect sizes, we fit an exponential decay curve assuming
fixed CV for three situations: unnormalized EHA105, unnormalized GV3101, and optimally
normalized GV3101 (lowest CV achieved in this publication) (Fig. 6D). The smallest detectable
effect size using 50 plants for unnormalized EHA105 was 13.7%, compared to 10.7% for
unnormalized GV3101 and 10.1% for GV3101 under optimal normalization (Fig. 6D). While
larger sample sizes can resolve even smaller differences, practical constraints limit the number
of plants that can be included in an experiment, but relatively small effect sizes can be
accurately distinguished with a small number of plants. Based on the Monte Carlo simulations,
with $n = 3$, the minimum reliably detectable effect size for unnormalized GV3101 is ~40%.
Normalization is most beneficial in experiments where small effect sizes must be resolved using
limited sample sizes. However, since normalization may also reduce statistical power in some
cases (Fig. S16), any normalization strategy should be validated empirically prior to
implementation.

Discussion

Here, we systematically measure expression variability in *N. benthamiana* agroinfiltration. The
inherent noise in the transient expression of FPs is considerable (as much as fourfold difference
in mean expression across experimental replicates) but mitigable through proper experimental

design (Fig. 1B). Using a mixed-effects model, we categorize nearly all observed variation in our
dataset (Fig. 1D). We find that multiple co-delivery methods for dual reporters can decrease
variation via ratiometric normalization, and of the examined methods, co-infiltration best reduces
variation, approximately halving the CV compared to no normalization (Fig. 2). Neither inoculum
density nor promoter strength in co-infiltrations predictably affect expression variation, but
normalization efficacy is highly sensitive to promoter choice (Fig. 3, Fig. 4). Some promoter
pairs may in fact increase variation or make experimental replicates less comparable, and using
identical promoters is the surest and most effective way to reduce CV within experiments (Fig.
4, Fig. 5). Normalization should not be done arbitrarily but rather validated on a case-by-case
basis for the co-delivery method and experimental design used, as this is absolutely necessary
for results that are reproducible and robust to potential sources of variability.
348
Though unambiguously beneficial, the use of constitutive, identical promoters is also particularly
limiting given the popularity of *N. benthamiana* to characterize the effects of promoter identity
and architecture on transcription. This problem is undoubtedly compounded in more complex
experimental designs. To increase predictability and standardization of normalization methods,
multi-lab efforts to quantify variation using diverse methods will be necessary. More robust
transcriptomic datasets of *N. benthamiana* expressing diverse transgenes in distinct host strains
could discourage reliance on a single, unvalidated normalizer. Additionally, multi-factor analysis
of binary vector construction and expression cassette positioning could elucidate the extent to
which these variables affect both expression and variability.
358
There is little published analysis of how the design of multi-gene T-DNA structure affects the
transient expression of those genes [47]. We find that convergent cassettes are far more highly
expressed than tandem or divergent cassettes. Since *Agrobacteria*'s VirD2 virulence protein
covalently binds the right border [48], it is thought that transgenes proximal to the right border
would be protected from exonucleolytic degradation and thus more likely to remain intact inside
the plant cell, but this hypothesis is not fully supported by our data (Fig. 2A, Fig. S10). Clearly,
T-DNA architecture affects transgene expression strength and variation, and a high-throughput
library approach would be ideal to address the multiplicity of orientation and position
combinations in multi-gene binary vectors. Determining these rules of expression and assessing
them in both stable and transient systems requires further, data-intensive work.
369
Given the lack of large, curated datasets of transient gene expression in *N. benthamiana*, it is
likely that many published *N. benthamiana* experiments chose sample sizes arbitrarily based on
unvalidated assumptions. Our model of the statistical power required to determine different
effect sizes with high confidence shows that in cases where the expected effect size is large, it
is easy to achieve the necessary sample size. For example, for effect sizes >50%, fewer than
three plants are required using the GV3101 strain. For effect sizes <20%, many plants are
required, to the extent that the number of plants needed may be infeasible for single
experiments (*e.g.* 50 plants are needed to show a 13.7% effect using EHA105). The majority of
synthetic biology applications for transient expression, such as bioproduction or circuit
reconstruction, require strong effects, which call for small numbers of plants. However, these

results could have strong implications for subtle phenotypes, like the transformation efficiency of
various *Agrobacterium* strains or mutants [37,49], or inducible gene expression systems [47,50,51].
382
While the data in this work required many hundreds of plants and multiple years to collect, we
were nonetheless only able to capture variation in *N. benthamiana* transient expression from a
rather narrow perspective. All of our data are from fluorescent reporters from 4-week-old plants,
with data collected three days after infiltration. We demonstrated that *Agrobacterium* strain,
promoter choice, and T-DNA design all impact variation, but many more strains, plasmids, and
expression cassette designs remain untested. There are other variables affecting
Agrobacterium-mediated transformation that we did not address, such as binary vector origin of
replication or T-DNA length. Furthermore, as novel *Agrobacterium*-independent technologies to
introduce nucleic acid and proteins into plant cells mature [52,53], their variability should also be
estimated. It is critically important to establish whether our values of variance are broadly
applicable, and future work should prioritize cross-lab validation.
394
Even in this simple context, variation in *N. benthamiana* transient expression depends strongly
on a host of inputs, and attempts to normalize and mitigate this variation must be individually
validated. As complexity increases in synthetic biology experimental designs, this
unpredictability in variation and in confidence will likely be compounded. Our results show that
the choice and validation of normalization methods and data collection are critical for *N.*
benthamiana to be a reliable platform for applications such as reconstituting complex metabolic
pathways and designing large genetic circuits. As we have thoroughly demonstrated, careful,
purposeful experimental design and interpretation are paramount to ensure robust and
reproducible results when using this essential technique for synthetic plant biology.

Methods

Media, chemicals, and culture conditions

Routine bacterial cultures were grown in Luria-Bertani (LB) Miller medium (BD Biosciences,
USA). *E. coli* was grown at 37 °C, while *A. tumefaciens* was grown at 30 °C unless otherwise
noted. Cultures were supplemented with rifampicin (100 µg/mL), kanamycin (50 µg/L, Sigma
Aldrich, USA), gentamicin (30 µg/L, Fisher Scientific, USA), or spectinomycin (100µg/L, Sigma
Aldrich, USA), when indicated.

Bacterial strain and plasmid construction

All bacterial strains and plasmids used in this work are listed in Supplementary Table 1. All
strains and plasmids created in this work are viewable through the public instance of the Joint
BioEnergy Institute (JBEI) registry: https://public-registry.jbei.org/folders/929. All strains and
plasmids created in this work can be requested from the strain archivist at JBEI with a signed
material transfer agreement. Plasmids were assembled by Gibson assembly using standard
protocols. Plasmids were routinely isolated using the QIAprep spin miniprep kit (Qiagen) and all
primers were purchased from Integrated DNA Technologies (IDT). Plasmid sequences were
verified using whole-plasmid sequencing (Primordium Labs). *Agrobacterium* was routinely

transformed by electroporation as described previously using a 1-mm cuvette and a 2.4-kV, 25-
µF, 200-Ω pulse [54].

N. benthamiana growth conditions

Wild type *N. benthamiana* (LAB accession) plants were obtained from the in-house seed bank
at the Joint BioEnergy institute. All seedlings and plants were grown at 25 °C in 60% humidity
under long-day conditions (16 h of light, 8 h of darkness) of 150 µmol m^{-2} s^{-1}
photosynthetically active radiation (PAR; wavelength: 400–700 nm).
427
As many pots as needed of Sungro Sunshine mix #4 (aggregate plus) were wetted by running
excess tap water from above and allowing it to drain through. The soil was then topped with a
layer of topsoil that was thoroughly wetted with a spray bottle. Using a spatula, a pinch of seeds
were sprinkled evenly over the soil. Ideally, 50-100 good seedlings per pot germinate. If seeds
are sown too densely, the seedlings are smaller, and having more sparse, healthy seedlings
results in superior plant quality at later stages. All seedling pots were placed in a flat with 1L
tapwater and covered with a hood, vents closed, in the growth room. If algal or cyanobacterial
contamination grows in the water, reduce the volume of water added to the tray.
436
After one week, flats (as many as needed) were filled with 18 3.11" x 3.11" x 2.25" Traditional
Inserts pots (Greenhouse Megastore). These pots were filled to the top with Sungro Sunshine
mix #4 (aggregate plus), supplemented with 1.5 Tbsp Osmocote (14-14-14) pellets per 4L soil,
and then 3L of tap water was added to the flat. The soil was allowed to soak up the water for 2-3
hours, and then any excess was drained off and the pots broken apart. Small holes were made
in each pot's soil, and the germinated seedlings were transplanted into individual pots. The root
of the seedling must not be damaged during transplantation. The block of soil from a seedling
pot can be removed and placed on its side to facilitate removing seedlings from soil without
causing damage. Flats were covered with a hood, vents closed, in the growth room. After one
week, all the vents were opened to allow the hood to slowly equilibrate to the ambient conditions
in the growth room. Five days later, the hoods were removed. Two days later, each flat was
watered with tap water supplemented with Peter's Professional (20-20-20) at 1 tsp per 4L water.
After five days, the pots were rearranged into a checkerboard pattern within the flats to
maximize the amount of space per plant, decreasing the density from 18 plants per flat to 9
plants per flat. Each flat was also watered with 1L water. Three days later, plants were
infiltrated. A highly detailed protocol of *N. benthamiana* growth and care is provided in
Supplemental Methods 1.

Agroinfiltration of N. benthamiana

Generated binary vectors were transformed into *A. tumefaciens* strain GV3101 via
electroporation [54]. BiBi strains were generated by growing a liquid culture of single
transformants to saturation, pelleting and washing thrice with ice cold 10% glycerol,
resuspending in 100ul water, and electroporating the second binary vector into the cells.
Selected transformants were inoculated in liquid media with appropriate selection the night

before the experiment. *A. tumefaciens* strains were grown until OD_{600} between 0.8-1.2,
centrifuged for 10 min at 4000 g, and resuspended in infiltration buffer (10 mM MgCl2, 10 mM
MES, and 200 µM acetosyringone, pH 5.6) to achieve the desired OD_{600}. Cultures were induced
for 1 h at room temperature on a rocking shaker. Leaves T4 and T5 of 4-week-old *N.*
benthamiana plants were syringe infiltrated with the *A. tumefaciens* suspensions. After
infiltration, *N. benthamiana* plants were maintained in the same growth conditions as described
above with 1L tap water per tray. Three days post infiltration, four 6-mm leaf discs per infiltrated
leaf were hole punched. The leaf discs were placed abaxial side up on 350 µL of water in black,
360-µL 96 well Costar Assay Plates with clear flat bottoms (Corning). eGFP (Ex.λ = 488 nm,
Em.λ = 520 nm) and mCherry (Ex.λ = 587 nm, Em.λ = 615 nm) fluorescences were measured
using a Synergy 4 microplate reader (Bio-tek). Gain was set at 100 and read height at 10.5 mm.
471

RUBY extraction and quantification
Leaves T4 and T5 of 4-week *N. benthamiana* were co-infiltrated with equal amounts of three
strains (carrying vectors 2-4 from Supplementary Table 1) on side of the leaf and infiltrated with
one strain (carrying vector 5 from Supplementary Table 5) on the other. Total OD_{600} for both
delivery methods was 0.5, and the infiltrated spots were not touching. Leaf tissues infiltrated
with RUBY constructs were excised with a razor blade from the leaf 5 dpi, frozen in liquid
nitrogen, and lyophilized for 2 d. Tissues were then weighed to record dry weights and
homogenized with metal beads in a PowerLyzer at 1000 g for 2 min. 80 µL of 20% methanol per
mg of dry weight was added to each sample. Samples were then centrifuged at 15,000 rcf for 5
min, and 100 µl of the supernatant and diluted with an additional 500 µl of water. Absorbances
at λ = 538 nm of 300 µl of the diluted extracts in black, 360-µL 96 well Costar Assay Plates with
clear flat bottoms (Corning) were measured in a Syngery 4 microplate reader (Bio-tek).
484

2-pyrone-4,6-dicarboxylic acid (PDC) extraction and quantification
The entirety of leaves T4 and T5 of 4-week old *N. benthamiana* were infiltrated with 0.2 OD_{600} of
each of the five enzymes in the PDC pathway (total OD_{600} = 1). At 5 dpi, both leaves were
frozen in liquid nitrogen together and lyophilized for 2 d. Leaves were then ball-milled and
extracted with 80% (v/v) methanol-water as solvent as previously described [40]. Metabolites were
analyzed using an HPLC-ESI-TOF-MS as previously described and quantified with a 6-point
calibration curve of PDC standard [40,41].
492

Statistical analysis
Mixed effects model and Monte Carlo simulations were run using R (v4.2.0) and the following
packages: tidyverse (v2.0.0) [55], transport (v0.14.6), car (v3.1.3), lme4 (v1.1.31) [56], and
performance (v0.13.0) [57]. All other analyses and figures were generated with Python (v3.11.4)
and the following packages: jupyterlab (v4.0.3) [58], pandas (v2.3.3), numpy (v2.3.3) [59], seaborn
(v0.13.2), matplotlib (v3.10.6), scipy (v1.16.1) [60], and statsmodels (v0.14.5).

Data availability

All code related to this study is publicly available on GitHub (github.com/shih-lab/benthi_variation).

References

1. Yang, S.-J., Carter, S. A., Cole, A. B., Cheng, N.-H. & Nelson, R. S. A natural variant of a host RNA-dependent RNA polymerase is associated with increased susceptibility to viruses by Nicotiana benthamiana. *Proc Natl Acad Sci USA* **101**, 6297–6302 (2004).

2. Kapila, J., De Rycke, R., Van Montagu, M. & Angenon, G. An Agrobacterium-mediated transient gene expression system for intact leaves. *Plant Sci.* **122**, 101–108 (1997).

3. Grimsley, N., Hohn, B., Hohn, T. & Walden, R. "Agroinfection," an alternative route for viral infection of plants by using the Ti plasmid. *Proc Natl Acad Sci USA* **83**, 3282–3286 (1986).

4. Schöb, H., Kunz, C. & Meins, F. Silencing of transgenes introduced into leaves by agroinfiltration: a simple, rapid method for investigating sequence requirements for gene silencing. *Mol. Gen. Genet.* **256**, 581–585 (1997).

5. Ruiz, M. T., Voinnet, O. & Baulcombe, D. C. Initiation and maintenance of virus-induced gene silencing. *Plant Cell* **10**, 937–946 (1998).

6. Chakrabarty, R. *et al.* PSITE vectors for stable integration or transient expression of autofluorescent protein fusions in plants: probing Nicotiana benthamiana-virus interactions. *Mol. Plant Microbe Interact.* **20**, 740–750 (2007).

7. Irmisch, S. *et al.* Discovery of UDP-Glycosyltransferases and BAHD-Acyltransferases Involved in the Biosynthesis of the Antidiabetic Plant Metabolite Montbretin A. *Plant Cell* **30**, 1864–1886 (2018).

8. Wood, C. C. *et al.* A leaf-based assay using interchangeable design principles to rapidly assemble multistep recombinant pathways. *Plant Biotechnol. J.* **7**, 914–924 (2009).

9. Martin, L. B. B. *et al.* Complete biosynthesis of the potent vaccine adjuvant QS-21. *Nat. Chem. Biol.* **20**, 493–502 (2024).

10. Calgaro-Kozina, A. *et al.* Engineering plant synthetic pathways for the biosynthesis of novel

antifungals. *ACS Cent. Sci.* **6**, 1394–1400 (2020).

11. Nett, R. S., Lau, W. & Sattely, E. S. Discovery and engineering of colchicine alkaloid

biosynthesis. *Nature* **584**, 148–153 (2020).

12. Jores, T. *et al.* Synthetic promoter designs enabled by a comprehensive analysis of plant

core promoters. *Nat. Plants* **7**, 842–855 (2021).

13. Hummel, N. F. C. *et al.* The trans-regulatory landscape of gene networks in plants. *Cell

Syst.* **14**, 501-511.e4 (2023).

14. Zhou, A. *et al.* A suite of constitutive promoters for tuning gene expression in plants. *ACS

Synth. Biol.* **12**, 1533–1545 (2023).

15. Brophy, J. A. N. *et al.* Synthetic genetic circuits as a means of reprogramming plant roots.

Science **377**, 747–751 (2022).

16. Belcher, M. S. *et al.* Design of orthogonal regulatory systems for modulating gene

expression in plants. *Nat. Chem. Biol.* **16**, 857–865 (2020).

17. Wu, F.-H. *et al.* Tape-Arabidopsis Sandwich - a simpler Arabidopsis protoplast isolation

method. *Plant Methods* **5**, 16 (2009).

18. Yoo, S.-D., Cho, Y.-H. & Sheen, J. Arabidopsis mesophyll protoplasts: a versatile cell

system for transient gene expression analysis. *Nat. Protoc.* **2**, 1565–1572 (2007).

19. Morey, K. & Khakhar, A. Exploring the frontier of rapid prototyping technologies for plant

synthetic biology and what could lie beyond. *New Phytol.* **242**, 903–908 (2024).

20. Bally, J. *et al.* The Rise and Rise of Nicotiana benthamiana: A Plant for All Reasons. *Annu.

Rev. Phytopathol.* **56**, 405–426 (2018).

21. Lux, M. W., Strychalski, E. A. & Vora, G. J. Advancing reproducibility can ease the "hard

truths" of synthetic biology. *Synth Biol (Oxf)* **8**, ysad014 (2023).

22. Zhou, X., Mehta, S. & Zhang, J. Genetically encodable fluorescent and bioluminescent

biosensors light up signaling networks. *Trends Biochem. Sci.* **45**, 889–905 (2020).

23. Khosla, A., Rodriguez-Furlan, C., Kapoor, S., Van Norman, J. M. & Nelson, D. C. A series
of dual-reporter vectors for ratiometric analysis of protein abundance in plants. *Plant Direct*
4, e00231 (2020).

24. Rattan, R., Alamos, S., Szarzanowicz, M., Markel, K. & Shih, P. M. Predictable modulation
of plant root development using engineered cytokinin regulators. *BioRxiv* (2024)
doi:10.1101/2024.12.06.627221.

25. Morffy, N. *et al.* Identification of plant transcriptional activation domains. *Nature* **632**, 166–
173 (2024).

26. Xu, P. *et al.* Interplay between Agrobacterium T-DNA and backbone DNA in transgenic
plant cells. *Transgenic Res.* **34**, 1 (2024).

27. Buyel, J. F. & Fischer, R. Predictive models for transient protein expression in tobacco
(Nicotiana tabacum L.) can optimize process time, yield, and downstream costs.
Biotechnol. Bioeng. **109**, 2575–2588 (2012).

28. Lopatkin, A. J. & Collins, J. J. Predictive biology: modelling, understanding and harnessing
microbial complexity. *Nat. Rev. Microbiol.* **18**, 507–520 (2020).

29. Covert, M. W., Gillies, T. E., Kudo, T. & Agmon, E. A forecast for large-scale, predictive
biology: Lessons from meteorology. *Cell Syst.* **12**, 488–496 (2021).

30. Breyne, P., Gheysen, G., Jacobs, A., Van Montagu, M. & Depicker, A. Effect of T-DNA
configuration on transgene expression. *Mol. Gen. Genet.* **235**, 389–396 (1992).

31. Bhattacharyya, M. K., Stermer, B. A. & Dixon, R. A. Reduced variation in transgene
expression from a binary vector with selectable markers at the right and left T-DNA borders.
The Plant Journal (1994).

32. Bashandy, H., Jalkanen, S. & Teeri, T. H. Within leaf variation is the largest source of
variation in agroinfiltration of Nicotiana benthamiana. *Plant Methods* **11**, 47 (2015).

33. Vazquez-Vilar, M. *et al.* GB3.0: a platform for plant bio-design that connects functional DNA
elements with associated biological data. *Nucleic Acids Res.* **45**, 2196–2209 (2017).

34. Moyle, R. L. *et al.* An optimized transient dual luciferase assay for quantifying microrna

directed repression of targeted sequences. *Front. Plant Sci.* **8**, 1631 (2017).

35. Cazzonelli, C. I. & Velten, J. Analysis of RNA-mediated gene silencing using a new vector

(pKNOCKOUT) and an in plantaAgrobacterium transient expression system. *Plant Mol.*

Biol. Rep. **22**, 347–359 (2004).

36. Hellens, R. P. *et al.* Transient expression vectors for functional genomics, quantification of

promoter activity and RNA silencing in plants. *Plant Methods* **1**, 13 (2005).

37. Thompson, M. *et al.* Genetically refactored Agrobacterium-mediated transformation.

BioRxiv (2023) doi:10.1101/2023.10.13.561914.

38. Huang, Y. W., Hu, C. C., Tsai, C. H., Lin, N. S. & Hsu, Y. H. Chloroplast Hsp70 Isoform Is

Required for Age-Dependent Tissue Preference of Bamboo mosaic virus in Mature

Nicotiana benthamiana Leaves. *Mol. Plant Microbe Interact.* **30**, 631–645 (2017).

39. Sheludko, Y. V., Sindarovska, Y. R., Gerasymenko, I. M., Bannikova, M. A. & Kuchuk, N. V.

Comparison of several Nicotiana species as hosts for high-scale Agrobacterium-mediated

transient expression. *Biotechnol. Bioeng.* **96**, 608–614 (2007).

40. Lin, C.-Y. *et al.* In-planta production of the biodegradable polyester precursor 2-pyrone-4,6-

dicarboxylic acid (PDC): Stacking reduced biomass recalcitrance with value-added co-

product. *Metab. Eng.* **66**, 148–156 (2021).

41. Eudes, A. *et al.* Production of hydroxycinnamoyl anthranilates from glucose in Escherichia

coli. *Microb. Cell Fact.* **12**, 62 (2013).

42. He, Y., Zhang, T., Sun, H., Zhan, H. & Zhao, Y. A reporter for noninvasively monitoring

gene expression and plant transformation. *Hortic. Res.* **7**, 152 (2020).

43. Alamos, S. *et al.* Quantitative dissection of Agrobacterium T-DNA expression in single plant

cells reveals density-dependent synergy and antagonism. *Nat. Plants* (2025)

doi:10.1038/s41477-025-01996-w.

44. Szarzanowicz, M. J. *et al.* Binary vector copy number engineering improves Agrobacterium-

mediated transformation. *Nat. Biotechnol.* **43**, 1708–1716 (2025).

45. Diamos, A. G. & Mason, H. S. Chimeric 3' flanking regions strongly enhance gene

expression in plants. *Plant Biotechnol. J.* **16**, 1971–1982 (2018).

46. Carlson, E. D., Rajniak, J. & Sattely, E. S. Multiplicity of the Agrobacterium Infection of

Nicotiana benthamiana for Transient DNA Delivery. *ACS Synth. Biol.* **12**, 2329–2338

(2023).

47. Kallam, K. *et al.* Tunable control of insect pheromone biosynthesis in Nicotiana

benthamiana. *Plant Biotechnol. J.* **21**, 1440–1453 (2023).

48. Dürrenberger, F., Crameri, A., Hohn, B. & Koukolíková-Nicola, Z. Covalently bound VirD2

protein of Agrobacterium tumefaciens protects the T-DNA from exonucleolytic degradation.

Proc Natl Acad Sci USA **86**, 9154–9158 (1989).

49. Lopez-Agudelo, J. C. *et al.* Rhizobium rhizogenes A4-derived strains mediate hyper-

efficient transient gene expression in Nicotiana benthamiana and other solanaceous

plants. *Plant Biotechnol. J.* (2025) doi:10.1111/pbi.70083.

50. Garcia-Perez, E. *et al.* A copper switch for inducing CRISPR/Cas9-based transcriptional

activation tightly regulates gene expression in Nicotiana benthamiana. *BMC Biotechnol.* **22**,

12 (2022).

51. Larsen, B. *et al.* Highlighter: An optogenetic system for high-resolution gene expression

control in plants. *PLoS Biol.* **21**, e3002303 (2023).

52. Demirer, G. S. *et al.* High aspect ratio nanomaterials enable delivery of functional genetic

material without DNA integration in mature plants. *Nat. Nanotechnol.* **14**, 456–464 (2019).

53. Demirer, G. S., Zhang, H., Goh, N. S., González-Grandío, E. & Landry, M. P. Carbon

nanotube-mediated DNA delivery without transgene integration in intact plants. *Nat. Protoc.*

14, 2954–2971 (2019).

54. Kámán-Tóth, E., Pogány, M., Dankó, T., Szatmári, Á. & Bozsó, Z. A simplified and efficient

Agrobacterium tumefaciens electroporation method. *3 Biotech* **8**, 148 (2018).

55. Wickham, H. *et al.* Welcome to the tidyverse. *JOSS* **4**, 1686 (2019).

56. Bates, D., Mächler, M., Bolker, B. & Walker, S. Fitting linear mixed-effects models using lme4. *J. Stat. Softw.* **67**, 1–48 (2015).

57. Lüdecke, D., Ben-Shachar, M., Patil, I., Waggoner, P. & Makowski, D. performance: An R Package for Assessment, Comparison and Testing of Statistical Models. *JOSS* **6**, 3139 (2021).

58. Perez, F. & Granger, B. E. IPython: A System for Interactive Scientific Computing. *Comput. Sci. Eng.* **9**, 21–29 (2007).

59. van der Walt, S., Colbert, S. C. & Varoquaux, G. The NumPy Array: A Structure for Efficient Numerical Computation. *Comput. Sci. Eng.* **13**, 22–30 (2011).

60. Virtanen, P. *et al.* SciPy 1.0: fundamental algorithms for scientific computing in Python. *Nat. Methods* **17**, 261–272 (2020).

Acknowledgements

We would like to thank Dr. Tyler Backman and Dr. Hector Garcia-Martin for their invaluable advice on statistics, and to all members of the Shih lab for their helpful discussions. We also thank Konami for facilitating discussion of expression cassette orientation via Dance Dance Revolution artwork.

BioRender was used to generate figures 1 (https://biorender.com/kh6zx0s), 2 (https://BioRender.com/2vhubua), S1 (https://biorender.com/82xc4es), S8 (https://biorender.com/4ila5ts), and S9 (https://biorender.com/w0f87at). This material is based upon work supported by the National Science Foundation Graduate Research Fellowship under Grant No. DGE 2146752. This work was part of the DOE Joint BioEnergy Institute (https://www.jbei.org) supported by the U. S. Department of Energy, Office of Science, Office of Biological and Environmental Research, supported by the U.S. Department of Energy, Energy Efficiency and Renewable Energy, Bioenergy Technologies Office, through contract DE-AC02-05CH11231 between Lawrence Berkeley National Laboratory and the U.S. Department of Energy. The funders had no role in manuscript preparation or the decision to publish. The views and opinions of the authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, expressed or implied, or

Author information

Authors and affiliations

Joint BioEnergy Institute, 5885 Hollis Street, Emeryville, CA 94608, USA.
Sophia N. Tang, Matthew J. Szarzanowicz, Sasilada Sirirungruang, Amy Lanctot, Liam D. Kirkpatrick, Krista Drako, Simón Álamos, Lucas M. Waldburger, Lyurui Cheng, Shuying Liu, Lena Huang, Sami Kazaz, Aymerick Eudes, Edward Baidoo, Mitchell G. Thompson, Patrick M. Shih

Environmental Genomics and Systems Biology Division, Lawrence Berkeley National Laboratory, Berkeley, California, USA
Sophia N. Tang, Matthew J. Szarzanowicz, Sasilada Sirirungruang, Amy Lanctot, Liam D. Kirkpatrick, Krista Drako, Simón Álamos, Lucas M. Waldburger, Shuying Liu, Lena Huang, Sami Kazaz, Aymerick Eudes, Edward Baidoo, Mitchell G. Thompson, Patrick M. Shih

Department of Molecular and Cell Biology, University of California, Berkeley, CA 94720, USA
Sophia N. Tang, Krista Drako

Department of Plant and Microbial Biology, University of California, Berkeley, CA 94720, USA
Matthew J. Szarzanowicz, Sasilada Sirirungruang, Amy Lanctot, Liam D. Kirkpatrick, Simón Álamos, Patrick M. Shih

Department of Bioengineering, University of California, Berkeley, California, USA
Lucas M. Waldburger

Biological Systems & Engineering Division, Lawrence Berkeley National Laboratory, Berkeley, CA 94720, USA
Lucas M. Waldburger

704 **Department of Nutritional Sciences and Toxicology, University of California, Berkeley, CA**
705 **94720, USA**
706 Lyurui Cheng
707
Innovative Genomics Institute, Berkeley, California, USA
Patrick M. Shih

Contributions

P.M.S. and M.G.T. conceptualized the initial study. M.G.T., S.A., L.M.W., and S.N.T. designed
experiments. S.N.T., K.D., and A.C. generated plasmids and strains. S.N.T, S.S., L.D.K., S.L.,
M.G.T., M.J.S., K.D., L.H., S.K., E.A.T, A.E., E.E., and A.L. performed experiments. M.J.S.
performed mixed-effect modeling and Monte Carlo simulations, and S.N.T. completed all other
data analysis. S.N.T. wrote the draft manuscript. All authors discussed the results, reviewed the
article, and approved the final article.

Corresponding authors

Correspondence to Patrick M. Shih or Mitchell G. Thompson.

Ethics declarations

M.G.T., M.J.S., and P.M.S. have financial interest in BasidioBio. PMS also has financial interest
in Totality Biosciences.
722

723
Figure 1. Longitudinal assessment of transient expression identifies sources of
variability in *N. benthamiana* fluorescence assays. **A)** Schematic of the transient

fluorescence assay. *N. benthamiana* is agroinfiltrated with an *A. tumefaciens* strain at an OD of
0.5. The binary vector contains, from left to right border, CaMV35S2:nptII and
PCM2:eGFP:At_Ubq3 oriented divergently in the T-DNA. Discs are collected from leaves 3 days
post-infiltration and measured in a plate reader. **B)** Distributions of GFP fluorescence for 15
experimental replicates, n=64 discs from 8 separate plants, except for 2022.06.06 and
2022.06.13 which are n=40. Dates of sample collection and measurement are written
YYYY.MM.DD. **C)** Boxplots for each plant in experimental replicates with the lowest
(2022.07.25) and highest (2023.05.30) mean raw fluorescence show the median and
interquartile range (IQR), and whiskers show the range of data within 1.5 IQR. Scatterplots
show T4 discs in white and T5 discs in black. **D)** Percent contributions to observed variability as
calculated from a mixed-effects model and illustrations of sources of variability.
737

Figure 2. Comparison of three discrete methods to deliver normalizing reporters. **A)** Left,
categories of delivery methods: unnormalized (green or magenta), co-infiltration (yellow), same
T-DNA (pink), and BiBi (blue). Center, distributions of eGFP and mCherry fluorescence, n=96
leaf discs per row. Right, cartoons showing the binary vector origin of replication, resistance
marker, and orientations of FP expression cassettes in the T-DNA. All binary vector cartoons
are read from left to right: ori, left border, T-DNA, right border. Origins of replication are pVS1
(diamond, black), BBR1 (circle, teal), and pSa (square, gray). Circles enclosing an X represent
tOcs, a 722bp spacer in between the two expression cassettes. **B)** Plant coefficients of variation
(CV), as calculated from the 8 discs per plant, when eGFP is treated as the reporter. All values
are eGFP/mCherry CV except for scheme 1, which is GFP CV. **C)** Plant CVs when mCherry is
treated as the reporter. All values are mCherry/eGFP CV except for scheme 2, which is

mCherry CV. Normalization scheme IDs match across all subpanels. For **B)** and **C)**, an
independent, one-tailed Welch's t-test and a Bonferroni correction were conducted to determine
whether the reporter/normalizer CV of a scheme is significantly less than the reporter CV for the
corresponding unnormalized scheme. Asterisks indicate p-values: * < 0.05, ** < 0.01. Total OD
infiltrated in all schemes is 0.5. ODs of co-infiltrated strains are 0.25 each. In the legend, "&"
indicates co-infiltration, arrows indicate the direction of an expression cassette, "+" indicates
BiBi, and GFP is abbreviated to "G" and mCherry to "R". Leaves T4 and T5 of six four-week-old
N. benthamiana plants were infiltrated in each experimental replicate, and four discs were
collected from each infiltrated leaf.

**Figure 3. OD$_{600}$ of two co-infiltrated strains affect transgene expression strength but not
variability**. Raw fluorescence signal from **A)** eGFP and **B)** mCherry for all OD$_{600}$ combinations.
Darker hues indicate increasing OD$_{600}$ of the competing strain, which carries the other FP. Error
bars show the standard error. Matrix of all OD$_{600}$ combinations' CV of **C)** eGFP fluorescence
and **D)** ratio of eGFP/mCherry. Each condition was infiltrated into leaves T4 and T5 of six
plants. Four discs were collected from each leaf for a total of n=48 leaf discs.

Figure 4. Promoters of co-infiltrated transgenes determine the efficacy of normalization.
eGFP and mCherry were driven by PCL2, PCM2, or PCH5 for a total of six unique binary

vectors. Log_2 raw fluorescence from **A)** eGFP and **B)** mCherry for all promoter combinations. Darker hues indicate stronger promoter of the competing strain. Error bars show the standard error. Matrix of all promoter combinations' CV of **C)** eGFP fluorescence and **D)** ratio of eGFP/mCherry. Log_2 raw fluorescence signal from **E)** eGFP and **F)** mCherry for all promoter combinations when instead the mCherry binary vectors are driven by PCL1, PCM1, or PCH4. Darker hues indicate stronger promoter of the competing strain. Error bars show the standard error. Matrix of all promoter combinations' CV of **G)** eGFP fluorescence and **H)** ratio of eGFP/mCherry. Conditions from the "Same Promoter Set" and "Different Promoter Set" experiments were infiltrated into leaves T4 and T5 of six plants, and four discs were collected from each leaf for a total of n=48 leaf discs. The experiments were performed independently on separate days.

Figure 5. Promoter choice affects variation between experimental replicates. A) Cumulative density functions of PCL2:eGFP alone or normalized by mCherry driven by PCL2, PCM1, or PCM2. Left: eGFP, right: eGFP/mCherry. Each experimental replicate is a unique color. The black line is the CDF for the pooled data of all six experimental replicates. The p-values of one-sample Kolmogorov-Smirnov tests appear to the right of each CDF, colored by experimental replicate. Asterisks indicate p-values: * < 0.05, ** < 0.01, *** < 0.001, and ns = not

significant. **B)** Absolute value of every experimental replicate's D, the greatest vertical distance
between the CDF a given experimental replicate (color) and the pooled CDF (black) from **A)**.
White boxplots indicate D values for eGFP CDFs and gray boxplots for eGFP/mCherry CDFs.
C) CVs of the six experimental replicates for each condition. As in **B)**, white boxplots indicate
eGFP and gray eGFP/mCherry. Every condition was infiltrated into leaves T4 and T5 of four
plants, and four discs were collected from each leaf in all six experimental replicates for a total
of n=192 leaf discs.
798

Figure 6. Modeling variation of transgene expression in 4-week-old *N. benthamiana* plants. A) Per-plant CV of unnormalized GFP calculated from data compiled across many years and independent experiments. CV for a plant is calculated using all discs derived from that plant, regardless of leaf. The number of plants included appears above the violin plots. **B)** Cumulative density function of per-plant, unnormalized eGFP CVs. Solid gray, empirical data from 32 batches of at least 30 plants. Dotted black, Monte Carlo simulated data. **C)** Minimum number of simulated plants needed to detect a given effect size with the CV of unnormalized EHA105, unnormalized GV3101, or optimally normalized GV3101 assuming 95% accuracy and statistical significance (p<0.05). **D)** Exponential regression fit to the simulated data in **C)**. Dashed horizontal line indicates the arbitrary 50 plant cap, and dotted vertical lines indicate the smallest detectable effect size with 50 plants. Points represent predictions from the Monte Carlo simulations in **C**. Orange, EHA105. Gray, unnormalized GV3101. Teal, optimally normalized GV3101.