Genomic adaptations of novel halotolerant bacteria from extreme North Greenland

Miguel Ángel Salinas García[1,2,3], *Anders Priemé[2,1,3]

[1] Centre for Exolife Sciences (CELS), Niels Bohr Institute, University of Copenhagen, Jagtvej 155A, DK-2200, Copenhagen, Denmark

[2] Center for Volatile Interactions (VOLT), Department of Biology, University of Copenhagen, Universitetsparken 15, DK-2100 Copenhagen, Denmark

[3] Section of Microbiology, Department of Biology University of Copenhagen, Universitetsparken 15, DK-2100 Copenhagen, Denmark

***Corresponding author**: Anders Priemé (aprieme@bio.ku.dk)

Keywords: extremophile, Greenland, halophile, halotolerant, genome, adaptations

ABSTRACT

Peary Land, northern-most Greenland, is a cold desert region whose soil microbiome remains underexplored. This is the first study to explore the halotolerant soil microbiome of this region with culture-dependent methods. Forty-nine taxonomically diverse bacterial isolates were obtained from dry biological soil crust, surface soil and permafrost soil. Nine isolates were selected for whole genome sequencing; these genomes showed adaptations to their cold, saline native environment, such as cold shock proteins and a large number of osmoprotectant transporters. The genomes also contained several biosynthetic gene clusters, including a large variety of clusters involved in the production of antimicrobial compounds. Furthermore, most of the sequenced genomes showed low digital DNA-DNA hybridisation to their closest relatives, suggesting that they represent novel species. Overall, this study demonstrates the unexplored potential of High Arctic deserts as a source of novel halotolerant bacteria and their secondary metabolites.

INTRODUCTION

Peary Land in extreme North Greenland is the northern-most landmass of our planet. The interior of this remote and uninhabited peninsula is characterized by very sparse vegetation, very low precipitation and very low temperatures (Mølgaard 1982). Due to the low precipitation, most of Peary Land is ice-free and vegetation is mainly found along the moist northern coast bordering the icy Arctic Ocean and in small depressions in the landscape fed by water from melting snowpack. In addition, Peary Land is home to one of the oldest permafrost formations in the world, located in Kap København and around 2 million years old (Fernandez-Guerra et al. 2023). Other notable locations include Cape Morris Jessup, the most northern point of mainland Greenland, and Citronen Fjord, which contains rich deposits of zinc and lead and is being considered for exploitation (Ironbark Zinc ltd.).

Due to the low precipitation and sparse vegetation, the soils in Peary Land are generally thin, alkaline and salt-rich, and poor in carbon, nitrogen and phosphorus (Stimmler et al. 2022). Surface soils of the High Arctic desert of Peary Land tend to harden and crack due to lack of water and evaporation, creating the typical patterns found in deserts (Charlier 1969). In contrast to warm or hot deserts, UV radiation is relatively low, with noon UV index averages of 2-3 in July (Acker and Leptoukh 2007; Hovila et al. 2009).

Biological soil crusts are common in hot and cold deserts and have been defined as communities of organisms dwelling on the soil surface. These communities create an almost living 'skin' of intimate associations between soil particles and photoautotrophic (cyanobacteria, algae, lichens, bryophytes) and heterotrophic (bacteria, fungi, archaea) organisms (Weber et al. 2022). Biological soil crusts regularly experience low water activity and are hence expected to harbour halophilic, salt-loving, and halotolerant taxa. These biological crusts prevent soil erosion and retain moisture, properties relevant in desert regions where bioremediation of biological soil crusts could be particularly effective (Guida et al. 2023).

Permafrost soil remains frozen for at least two consecutive years and is covered by the active layer, i.e., soil that thaws during summer. Despite the frozen conditions, permafrost soils are home to a large diversity of microorganisms (Jansson and Taş 2014). In permafrost and other frozen environments, liquid brine inclusions (cryopegs) form due to salt fractionation (Iwahana et al. 2021). In contrast to the bulk permafrost, nutrients and microorganisms can move in the brine, enabling microbial activity. These brines can reach salt concentrations near saturation, but nevertheless provide a liquid environment for microorganisms to inhabit (Steven et al. 2006). Permafrost soil, and especially the cryopegs within, has been a source of many

halophilic microorganisms described in the literature (Niederberger et al. 2009; Shcherbakova et
al. 2009; Spirina et al. 2017). Microorganisms isolated from these environments can serve as
models in astrobiology (Nicholson et al. 2013), and their enzymes and metabolites have
biotechnological and industrial applications (Flores-Gallegos et al. 2019; Parvizpour et al. 2021).

Microbial life in Peary Land soils require special adaptations to the low
temperatures, low water activity and low nutrient conditions; including modifications to cellular
membranes, proteins, genetic material and metabolism (Sarma et al. 2023). The extremophiles
that thrive under these conditions are known as psychrophiles or psychrotolerant (organisms
that grow below 10ºC) and xerotolerant (organisms that can grow at water activity below 0.8,
where 1 is pure water) (Lebre et al. 2017). Halophiles are a group of xerotolerants that have
evolved to grow under high salt conditions with low water activity (Stevenson et al. 2015).
Halophilic microorganisms have previously been isolated from non-Polar deserts such as the
Atacama Desert (Ruginescu et al. 2019), but very little is known about halophilic and halotolerant
microorganisms in Polar deserts.

Previous studies have shown that the Greenland Ice Sheet is a promising source of
biosynthetic gene clusters (Jaarsma et al. 2023). With this study, we aim to extend this potential
to different Peary Land soils, whose microbiome remains underexplored and may harbour novel
bacterial isolates with potential applications in biotechnology. We hypothesize that i) permafrost
soil, biological soil crust and surface soil in Peary Land are inhabited by novel culturable
halophilic and halotolerant bacteria, ii) the genomes of these bacteria reveal adaptations to life
at low water activity and iii) Northern Greenland represents a potential source of secondary
bacterial metabolites.

MATERIALS AND METHODS

Sampling sites

The Kap København formation (sampled at N82.341460, W21.256961, 160 meters
above sea level, m.a.s.l.) (**Fig. 1a**), consists of a 100-m thick succession of shallow marine
sediments which were deposited during an interglacial period 2 million years ago. The deposits
were later uplifted and exposed subaerially about 1 million years ago, at which time the deposits
became permafrost (Fernandez-Guerra et al. 2023). Samples were extracted using a sterilized
shovel from the permafrost ca. 4 meter below the surface accessed from the side of the
formation, which was exposed due to recent erosion. Within minutes of extraction, the extracted
permafrost material was opened using a sterile chisel and subsamples from the freshly exposed

material were immediately transferred to sterile plastic tubes where they were kept until
initiation of the isolation work at our laboratories.

At Citronen Fjord (**Fig. 1b**), conditions are desert-like with very sparse vegetation
consisting of scattered specimens of Arctic Willow (*Salix arctica*) and Entire-leaved Mountain
Avens (*Dryas integrifolia*). Here, samples of biological soil crust were obtained from a site devoid
of vegetation (N83.085782, W28.2496755, 28 m.a.s.l.). Biological soil crust was scooped from
the surface using a sterile spoon and transferred to sterile plastic tubes.

Fig. 1 Sampling sites at Kap København (a) where permafrost soil samples were obtained at the erosion front left of the
snowpack, Citronen Fjord (b) and Cape Morris Jessup (c). d: Location of Peary Land within Greenland and sampling
sites: Cape Morris Jessup (magenta diamond), Citronen Fjord (red triangle) and Kap København (dark green circle).
Greenland image: NASA. Peary Land map: Blue Green Atlas, licensed under CC BY 4.0

Due to its proximity to the Arctic Ocean, Cape Morris Jessup (**Fig. 1c**) harbours
more dense vegetation (5-15 percent ground cover) compared to Citronen Fjord. The only
weather station in Peary Land, WMO 04301 in Cape Morris Jessup, reported a mean temperature
of -14ºC, mean annual precipitation of 234 mm, and temperature maxima of around 12ºC in the
summer for the period 2020-2023 (Meteostat). The vegetation is dominated by Purple Saxifrage
(*Saxifraga oppositifolia*) and Arctic Willow. At N83.657236, W33.368495, 5 m.a.s.l., we sampled
biological soil crust (as described above) and soil from 0-4 cm depth by hammering sterile steel
cores (4.5 cm diameter) into the soil. A subsample from each soil core was transferred to sterile
plastic tubes.

Strain isolation

Samples were stored at 4ºC for a period between one and three months, which included the travel time from the sample sites in Peary Land to Copenhagen, Denmark, and pre-incubation of the samples, known as cold enrichment, which has been shown to increase the number of isolated bacteria from permafrost (Vishnivetskaya et al. 2000).

After pre-incubation, the samples were plated on agar plates. Under sterile conditions, approximately 1 gram of soil was suspended in 1 mL of phosphate-buffered saline (PBS) solution supplemented with 10% NaCl. This buffer was chosen to initially select for bacteria that were resistant to shock from high osmolarity. After gentle mixing for 5 minutes, the samples were briefly centrifuged to remove solid particles from suspension. 100µL of the supernatant were plated on agar plates (see below) and left to grow. For the permafrost samples, the supernatant was diluted 100 and 1000 times prior to plating. The plates were incubated at either 4ºC, 10ºC or 25ºC, and checked weekly and daily, respectively. New colonies were randomly selected based on unique morphologies, then plated into fresh agar plates and incubated again. After incubation, a single colony was picked, plated and incubated. This process was repeated several times until only a single type of colony was observed, at which point we considered that the culture was axenic. Single colonies from the axenic plates were transferred into liquid medium and incubated overnight. 1mL of these cultures was mixed with a glycerol stock (20% v/v final concentration) and frozen at -70ºC for long-term storage.

For isolation of halotolerant and halophilic bacteria, three saline media were used. The first was a modified HM medium (Ventosa et al. 1982), referred to in this manuscript HM10 for its 10% w/v content of NaCl, and containing, in g L^{-1}: NaCl (100) $MgSO_4 \cdot 7H_2O$ (1), $CaCl_2 \cdot 2H_2O$ (0,36), KCl (2), KBr (0,27), $FeCl_3 \cdot 6H_2O$ (0,002), $NaHCO_3$ (0,06), Proteose-peptone no. 3 (5), yeast extract (10), glucose (1) and Bacto agar (20, when applicable) adjusted to pH 7,8-8. The second was a medium devised for this study named OHAM (Oligotrophic Halophilic Actinobacteria Medium) containing, in g L^{-1}: soluble starch (20), KNO_3 (0.2), NaCl (100), $MgSO_4 \cdot 7H_2O$ (0.5), K_2HPO_4 (0.5), $FeSO_4 \cdot 7H_2O$ (0.01), sodium pyruvate (0.05) and Bacto agar (15, when applicable), and a final pH of 7.2. The third was R2A medium (Sigma Aldritch, Germany) supplemented with 10% NaCl.

In addition, a non-saline medium, Gauze Medium 1 (GM1), was also used, containing in g/L: Soluble starch (20), KNO_3 (1), NaCl (0.5), $MgSO_4 \cdot 7H_2O$ (0.5), K_2HPO_4 (0.5), $FeSO_4 \cdot 7H_2O$ (0.01) and bacto agar (15, when applicable). Final pH was adjusted to 7.2-7.4. This

147 medium has been used in previous works to isolate bacterial strains from permafrost (Wang et
148 al. 2018).

149   Premade marine agar (MarA, Sigma Aldritch, Germany) and Tryptic Soy Broth (TSA,
150 Sigma Aldritch, Germany) were used for routine culturing of the isolates after axenic cultures
151 were obtained.

Determination of the effect of NaCl on growth

To test the NaCl tolerance of the selected isolates, 96-well plates were prepared
with HM in different concentrations of NaCl, from 0% (w/v) to 20% in increments of 2.5%. The
plates were inoculated with overnight cultures to a constant initial OD_{600}, blanks were made for
each different NaCl concentration. The plates were then incubated at 25ºC in a BioTek Synergy
H1 Multimode Reader (Agilent, USA) for 5 days with constant rotation and OD measurements
every 30 minutes. Range of growth was determined based on the observed growth in the wells at
the end of the experiment. To calculate the effect of NaCl on growth, blanks were subtracted and
each replicate was fit to a Gompertz model using the R package *growthrates* (Petzoldt 2016)
version 0.8.4. The resulting parameters were used to calculate the area under the curve using the
R package *pracma* (Borchers 2011) version 2.4.4.

16S rRNA gene sequencing and analysis

Total DNA was isolated from overnight cultures using the DNeasy PowerSoil Pro Kit
(Qiagen, Germany). The 16S rRNA gene was then amplified with the PCRBio kit (PCR Biosystems,
UK) using the universal primers 27F (5'-AGA GTT TGA TCM TGG CTC AG-3') and 1492R (5'-ACG
GYT ACC TTG TTA CGA CTT-3') purchased from TAG Copenhagen (www.tagc. com). After the size
and quality of the amplicons was assessed with electrophoresis on agarose gel and ethidium
bromide staining, the amplified product was sent for Sanger sequencing to Eurofins (Germany)
with the same primers, to obtain both forward and reverse reads.

The forward and reverse reads were aligned in MEGA11 (Tamura et al. 2021) using
the MUSCLE algorithm (Edgar 2004) to produce the near full-length 16S rRNA gene sequences.
Then, BlastN (Sayers et al. 2022) was used to identify the strains to the genus level; this search
was restricted to type strain sequences. To study phylogenetic relationships, an alignment file
was produced in MEGA11 using the MUSCLE algorithm with the near full-length 16S rRNA gene
sequences and the full-length 16S rRNA gene sequences of related type strains. A phylogenetic
tree was constructed with this alignment file using IQ-TREE 2 (Minh et al. 2020) with ModelFinder
(Kalyaanamoorthy et al. 2017) and ultrafast bootstrap (Hoang et al. 2018). Lastly, the tree was

drawn with FigTree v1.4.4 (http://tree.bio.ed.ac.uk/software/figtree/) and edited using Inkscape
v 1.3.2 (https://inkscape.org/).

Genome sequencing and assembly

Total genomic DNA was isolated from overnight cultures using the DNeasy
PowerSoil Pro Kit (Qiagen, Germany). Short read libraries were prepared using the DNA-Seq Enz
kit for MagicPrep (Tecan, USA). The quality of the libraries was checked using a 5200 Fragment
Analyzer System (Agilent, USA) before sequencing of the DNA fragments using Miseq2 (Illumina,
Germany). The quality of the reads was examined with FastQC v.0.12.0
(https://www.bioinformatics.babraham.ac.uk/projects/fastqc/) and Trimmomatic v0.5.1 (Bolger
et al. 2014) was used to remove adapter sequences and low quality reads. Unicycler v0.5.1 (Wick
et al. 2017) was used with SPAdes v4.0.0 (Prjibelski et al. 2020) to assemble the genomes. The
basic statistics and quality of the assemblies were determined with QUAST v5.2 (Mikheenko et
al. 2018), checkM v1.2.3 (Parks et al. 2015) and BUSCO v5.7.1 (Manni et al. 2021). To sequence
the genome of *Arthrobacter halodurans* DSM 21081[T], genomic DNA was directly ordered from the
German Collection of Microorganisms and Cell Cultures (DSMZ) and sequenced as described
above.

Genome analysis

The phylogeny of the assembled genomes was determined with the Type (Strain)
Genome Server (TYGS) at DSMZ (tygs.dsmz.de), with nomenclature provided by the List of
Prokaryotic Names with Standing in Nomenclature (LPSN) (Meier-Kolthoff and Göker 2019;
Meier-Kolthoff et al. 2022). A threshold of 70% digital DNA-DNA-Hybridation (dDDH), as
determined by the Genome-to-Genome Distance Calculator d_4 formula (GGDC formula 2), was
used to determine the species of the isolates. The genomes were annotated with Bakta
(Schwengers et al. 2021) and their metabolic pathways were explored by obtaining KEGG
Orthology (KO) numbers using the BlastKOALA and Reconstruct tools of the Kyoto Encyclopedia
of Genes and Genomes (KEGG) (Kanehisa et al. 2016). AntiSMASH (Blin et al. 2023) in relaxed
mode was used to find biosynthetic gene clusters (BGCs) in the assemblies.

RESULTS AND DISCUSSION

Bacterial isolates

A total of 49 bacterial isolates were obtained from the samples (**Table 1**). The soil
samples from Citronen Fjord (CF) and Kap København (KK) were the most prolific, while a lower
number of isolates was obtained from Cape Morris Jessup soil (CMS) and biological crust (CMC).

In addition to bacterial colonies, fungal colonies were observed in plates from all samples. We
focused on the bacterial isolates and fungal isolates were not further investigated.

The 17 strains isolated with saline media were capable of growth under a wide
range of salt concentrations and showed different responses to salt (**Fig. 2**). Three of these
isolates, KK5.5, CF3.5 and CF4.18 showed the highest growth at 0% w/v NaCl, while other
isolates such as CF4.1, CF4.2 or CMS1.3 showed almost or no growth in medium without NaCl.
The growth of all isolates was impaired at high salt concentrations, but eight isolates showed
some level of growth at the maximum concentration tested, 20% w/v NaCl. Interestingly, some
isolates belonging to the same genus, e.g., *Nesterenkonia*, showed similar responses to NaCl,
while both *Arthrobacter* isolates, KK5.5 and KK5.7, showed different growth patterns as NaCl
concentrations increased. Halophiles are commonly classified as such if their growth optimum
is above 2% w/v NaCl, while halotolerant organisms grow optimally below 2% and can tolerate a
wide range of salinities (Ollivier et al. 1994). Based on this classification, the tested isolates
consisted of a mixture of halotolerant (CF3.5, CF4.18 and KK5.5) and halophiles (the rest of the
strains).

Fig. 2 *Growth of strains isolated in saline media at 0 to 20% w/v NaCl. Y axis is Area Under the Curve of a Gompertz*

228 *model fitted to the data to indicate total growth during the 5-days incubation period. Black dots represent datapoints.*

229 *Lines indicate LOESS fit to data with a span of 0.75. Shaded area is the 95% confidence interval. Colour of lines*

230 *indicates genus. Isolates whose genomes were sequenced are highlighted in bold and italics*

 Table 1 *Number of bacterial isolates obtained from each site and medium. The total number of bacterial isolates was*
237   *49. Saline media are marked with an asterisk (*)*

Medium	Citronen Fjord (biological crust)	Cape Morris Jessup (soil)	Cape Morris Jessup (biological crust)	Kap København (permafrost soil)
GM1	10	4	2	16
HM10*	7	0	0	0
R2A+10% NaCl*	4	2	0	0
OHAM*	2	0	0	2
Total	23	6	2	18

16S rRNA gene sequencing

Based on their 16S rRNA gene sequence, the affiliation of the isolates with known
type strains was visualized (**Fig. 3**). Full details of each isolate and accession numbers are
available in Table S1. The isolates mainly belonged to three phyla: Actinomycetota, Bacillota and
Pseudomonadota, while Bacteroidota was only represented by one isolate, *Acidovorax* sp.
CMS1.1 (Fig. 3B). All isolates demonstrated the ability to grow at 4ºC. Many of the strains were
isolated in saline media, thus demonstrating their halotolerance. These included all the
Bacillota, as well as isolates belonging to the genus *Nesterenkonia*. Some strains isolated in non-
saline media also proved to be halotolerant, as they grew on MarA (containing 1.945% w/v NaCl),
including most members of *Arthrobacter* and *Pseudomonas* sp. KK6.1. However, other isolates,
like *Massilia* sp. CMS3.1 and *Flavobacterium* sp. CMS1.2 were unable to grow in medium
supplemented with >1% NaCl (data not shown).

The most abundant genera were *Arthrobacter*, predominantly isolated from
permafrost soil, and *Streptomyces* and *Nesterenkonia*, obtained from biological crust. Previous
studies have shown that these genera, as well as other genera isolated in this study, are
commonly found in polar environments. For example, Singh and colleagues (Singh et al. 2017)
reported that *Arthrobacter* and *Pseudomonas* were the most commonly isolated genera in
permafrost soil from Svalbard, Norway, where they also isolated strains belonging to
Nesterenkonia, Bacillus, and others. Interestingly, the authors also obtained isolates belonging
to *Streptomyces*, while none of our isolates in this genus were obtained from permafrost soil.
Streptomyces and other Actinomycetota are commonly found in arid and desert soils (Bull and
Asenjo 2013) as well as polar environments (Abirami et al. 2021), and extremophilic and
extremotolerant *Streptomyces* have drawn attention as potential producers of novel
antibacterial compounds and other compounds of interest (Sivalingam et al. 2019). In addition,

Nesterenkonia has been isolated from Arctic permafrost soil (Singh et al. 2021), as well as other

cold environments, like soils of Antarctica and the Tibetan plateau (Finore et al. 2016; Dai et al.

2022).

Arthrobacter is commonly found in permafrost soil with both culture-dependent

(Hinsa-Leasure et al. 2010; Singh et al. 2017) and culture-independent methods (Shen et al.

2021), the latter of which have shown that *Arthrobacter spp.* can constitute up to 4% of the total

bacterial community (Dsouza et al. 2015). This suggests that either *Arthrobacter spp.* are

especially adapted to survive in permafrost soil, or that they can quickly adapt to newly forming

permafrost. The *Arthrobacter* pangenome (Gushgari-Doyle et al. 2022) revealed great potential

adaptability, which could explain its prevalence in permafrost soil. The same study showed the

halotolerance of several *Arthrobacter* strains, a similar observation to our *Arthrobacter* isolates,

as many of them could grow on MarA or were directly isolated in saline medium. Based on its

prevalence in a variety of environments, some authors have used *Arthrobacter* as a model to

study bacterial adaptations in soil (Chauhan et al. 2018; Shen et al. 2021). In our study, the

isolated *Arthrobacter* strains were initially isolated in either GM1 or OHAM, media with no

nitrogen source other than nitrate, and with glycogen as the only carbon source (along with

pyruvate in OHAM). Some members of this genus are known to be able to assimilate nitrate and

nitrite for growth (He et al. 2020; Liu et al. 2023). Based on our results, GM1 may be an effective

selective medium to isolate *Arthrobacter* strains from permafrost soil.

The diversity of isolates obtained from Kap København, Cape Morris Jessup and

Citronen Fjord, and the displayed halotolerance of the isolates, show that the sampled sites are

inhabited by a wide range of halotolerant bacteria, thus partially failing to reject our first

hypothesis. In the next section, we discuss the novelty of the isolates.

Fig. 3 [next page] *Phylogenetic affiliation of isolates. a: Consensus cladogram based on the 16S rRNA gene sequences*

of the isolates and selected type strains calculated with TIM2+F+I+G4. Bootstrap values (as percentage of 10000

repetitions) are shown at the branch points. Phyla are highlighted with colours as indicated in panel b. Within

Actinomycetota, genera are further highlighted with colours as indicated in panel b. The markers indicate the source

of isolation: Citronen Fjord (black circle), Kap København (white circle), and Cape Morris Jessup soil (semicircle), and

crust (dots). Type strains have no markers. Aquifex pyrophilus DSM 6858 was used as the outgroup. ***b:*** *Number of*

isolates of each detected genus, and colours used to highlight groups in panel a

a

b

Genus	Phylum	Isolates (saline media)	Isolates (non-saline media)
● Aeromicrobium	● Actinomycetota	2	-
● Arthrobacter	● Actinomycetota	2	13
● Micrococcus	● Actinomycetota	-	1
● Nesterenkonia	● Actinomycetota	6	-
● Paenarthrobacter	● Actinomycetota	-	1
● Pseudarthrobacter	● Actinomycetota	1	2
● Streptomyces	● Actinomycetota	-	8
● Jeotgalibacillus	● Bacillota	2	-
● Oceanobacillus	● Bacillota	3	-
● Salipaludibacillus	● Bacillota	2	-
● Flavobacterium	● Bacteroidota	-	1
● Acidovorax	● Pseudomonadota	-	1
● Massilia	● Pseudomonadota	-	1
● Pseudomonas	● Pseudomonadota	-	3

Genomes and functional analyses of selected isolates

297
From the 16S rRNA gene-based phylogeny, nine isolates were selected for whole
genome analysis (**Table 2**), of which eight were halotolerant. *Massilia* sp. CMS3.1 was
determined to not be halotolerant, as it was only able to grow in GM1 with <1% NaCl, but was
chosen for genome sequencing (a) as a Gram-negative and non-halotolerant representative and
(b) to investigate its adaptations to its natural environment as a non-halotolerant bacterium. In
the case of *Arthrobacter* sp. KK5.5, its closest strain was *A. halodurans* DSM 21081^T, whose
genome was not available in any database. To confirm the phylogeny of our isolate, we
sequenced the genome of *A. halodurans* DSM 21081^T. All genomes showed completeness above
98% and contamination below 2%, which was determined sufficient to proceed to the next part
of the analysis. Six of the nine sequenced isolates showed dDDH similarities to their closest
relatives below 30%, well below the established threshold of 70% to distinguish between
different species (**Table 2**). Following the phylogenetic analysis, we conducted a functional
analysis of the genomes using the KEGG database and KEGG Orthology numbers.

316 **Table 2** *Isolates selected for genome sequencing, information on the assemblies, and similarity to their closest relatives bases on digital DNA-DNA hybridization (dDDH).*

317 *Completeness and contamination were calculated with CheckM. Completeness calculated by BUSCO is available in Table S2. GGDC: Genome-to-Genome Distance Calculator.*

318 **[DISCLAIMER: We have received permission from the Greenlandic authorities to deposit our genome-sequenced strains at DMSZ and are currently in the process of doing so. DSMZ*

319 *numbers will be added to this table.]*

Isolate	NaCl range (% w/v) (optimum)	Genome size (Mbp)	No. of contigs	N50	GC content (%)	Completeness (%)	Contamination (%)	Number of CDS	Closest match (dDDH)	Similarity to closest dDDH match (GGDC formula 2, %)	DSMZ strain number*	Genome accession number
CF4.18	0-17.5 (0)	5.27	159	76384	37.32	99.34	1.77	5087	*Salipaludibacillus neizhouensis* DSM 19794	74.7	TBD	JBIMEV000000000
CF4.6	0-15 (7.5)	4.07	39	241120	37.48	99.33	1.22	4050	*Oceanobacillus rekensis* PT-11[T]	27.1	TBD	JBIMEX000000000
CMS3.1	0-1 (0)	5.37	85	300246	62.92	100	0.95	4727	*Massilia litorea* LPB0304[T]	29.4	TBD	JBIMES000000000
CF4.12	0-20 (5)	2.94	13	734929	66.33	99.66	0.43	2687	*Nesterenkonia halotolerans* DSM 15474[T]	81.8	TBD	JBIMEW000000000
CF4.4	0-20 (5)	3.31	19	389302	66.90	99.24	0.61	3018	*Nesterenkonia aurantiaca* DSM 27373[T]	31.8	TBD	JBIMEY000000000
CMS1.6	0-20 (5)	3.05	37	330592	67.29	99.11	0.46	2775	*Nesterenkonia aurantiaca* DSM 27373[T]	82.6	TBD	JBIMET000000000
KK5.5	0-20 (0)	3.34	40	190007	68.49	99.77	0	3053	*Arthrobacter halodurans* DSM 21081[T]	22.6	TBD	JBIMER000000000
CF3.5	0-10 (2.5)	3.40	29	432807	68.19	99.74	0	3303	*Aeromicrobium fastidiosum* NRBC 14897[T]	20.3	TBD	JBIMEZ000000000
CF4.19	0-15 (5)	3.49	21	441989	70.42	98.45	0.52	3442	*Aeromicrobium halocynthiae* JCM15749[T]	21.2	TBD	JBIMEU000000000

Adaptations to high salinity and low water activity

All sequenced isolates, except for *Massilia* sp. CMS3.1, possessed orthologs of

genes encoding importers of osmoprotectants like proline, ectoine and betaine choline (**Fig. 4**

and **Table S3**). Osmoprotectants are small, neutral organic solutes that do not interfere with

cellular processes, and can thus be accumulated to mitigate osmotic and low-temperature

stress (Pathania et al. 2022). The ability to import osmoprotectants from the environment offers

flexibility, allowing bacteria to adapt to fluctuating salinity. This strategy reduces the energy cost

associated with de novo synthesis, providing a competitive advantage. Interestingly, the non-

halotolerant *Massilia* sp. CMS3.1 possessed only one ortholog of osmoprotectant transporters

(*yaaJ*), whereas its close relative, *Telluria aromaticivorans*, has 16 transporters and 9 synthesis-

related genes, despite its low salt tolerance (Son et al. 2021). Conversely, *Salipaludibacillus*

neizhouensis CF4.18 had 33 orthologues related to osmoprotectant transport, almost twice that

of the type strain of its species, revealing large intra-species variation. These findings suggests

that halotolerance may not be directly corelated with the copy number of genes related to

osmoprotectant synthesis and import.

Other abundant groups were orthologues related to ion transport (including *trkA*,

kdpABC, *tlyC*, and *nhaAC*). Ion transporters are very common in bacteria and are not necessarily

indicative of halotolerance, as evidenced by similar numbers of ion transporter orthologues in

non-halotolerant strains (**Fig. 4**). However, ion transporters may have specific adaptations that

aid with halotolerance. For example, the cyanobacterium *Aphanothece halophytica* has ion

transporters with different ion specificities that allow it to tolerate high salinity and pH

(Wutipraditkul et al. 2005). In the halophilic bacterium *Halomonas beimenensis*, the genes *nqrA*,

trkA, *atpC*, *nadA*, and *gdhB* — related to ion transport — along with *spoT*, *prkA*, *mtnN*, *rsbV*, *lon*,

smpB, *rfbC*, *rfbP*, *tatB*, *acrR1*, and *lacA* — related to cellular signalling, quorum sensing,

transcription/translation, and cell motility — are critical for halotolerance, and their deletion

impairs growth at high salinity (Chen et al. 2017). Similarly, the genes *trkA2*, *smpB*, *nadA*, *mtnN2*,

rfbP, *lon*, and *atpC* are upregulated in high salinity in *Virgibacillus chiguensis* (Chen et al. 2018).

Orthologues of many of these genes were found in varying numbers in the sequenced isolates

(**Table S3**), suggesting their potential role in the observed halotolerance across the bacteria

isolated in this study.

Cold temperatures can lead to similar stress variables than high salinity. For

example, water activity is reduced as ice crystals form, leading to osmotic stress, and brines can

form as salt ions are excluded from ice crystals (Iwahana et al. 2021). Thus, the halotolerance-

related genes discussed may also aid the isolates against the low temperatures in their native

Northern Greenland. In addition, the genomes also contained several orthologues of cold shock

proteins (*cspA*). These proteins have a protective effect against the stresses associated with low

temperatures by binding to single stranded nucleic acids and regulating transcription and

translation (Horn et al. 2007).

Fig. 4 KEGG Orthology count numbers of groups of genes that have been described to be important for halotolerance in the sequenced genomes and in genomes of close relatives.

The complete dataset is available in Table S3. The close relatives are as follows, from left to right: *Aeromicrobium halocynthiae* JCM 15749[T], *Arthrobacter halodurans* DSM 21081[T], *Nesterenkonia halotolerans* DSM 15474[T], *Nesterenkonia aurantiaca* DSM 27373[T], *Oceanobacillus rekensis* PT11[T], *Salipaludibacillus neizhouensis* DSM 19794[T], and *Telluria aromaticivorans* ML15P13[T]

Other metabolic pathways

Some of the sequenced genomes contained genes encoding enzymes related to nitrogen and sulphur cycling. *S. neizhouensis CF*4.18 and *Oceanobacillus* sp. CF4.6 had a complete assimilatory sulfate reduction pathway, while *N. aurantiaca* CMS1.6 and *N. halotolerans* CF4.12 lacked this pathway completely. The rest of the sequenced isolates had an incomplete pathway, lacking the enzymes to reduce sulphate to sulphite. However, all genomes contained transporters for alkanesulfonates, and most genomes contained genes encoding for alkanesulfonate monooxigenases. This alternative source of sulphur may aid against sulphur starvation (Ellis 2011), as the low amount of organic carbon in Peary Land soils could limit the amount of sulphur available from organic matter (Stimmler et al. 2022). However, sulphur may not be a limiting nutrient in the permafrost at Kap København, as it consists mostly of marine deposits, which are typically enriched in sulphur (Funder et al. 1984; Francois 1987).

Nesterenkonia sp. CF4.4, *Aeromicrobium* sp. CF3.5, *Arthrobacter* sp. KK5.5, *S. neizhouensis* CF4.18 and *Massilia* sp. CMS3.1 had genes encoding for both nitrate and nitrate reductases, while these were not found in the other isolates. Interestingly, *Oceanobacillus* sp. CF4.6 was the only isolate whose genome contained *nirK*, which encodes for a nitric oxide-producing nitrite reductase. The reduction of nitrite to nitric oxide (NO) is the second step in denitrification after the reduction of nitrate to nitrite. NO is then reduced to nitrous oxide (N_2O), which is finally reduced to N_2. NO and N_2O have significant impact on atmospheric chemistry, as they can induce the formation of tropospheric ozone and form NO_x; furthermore, N_2O is a potent greenhouse gas. In arid areas, especially in the polar regions, nitrification and denitrification remain poorly understood (Ramond et al. 2022). There are no available studies on the nitrification or denitrification rates, or on the emission of NO and N_2O, from Northern Greenland, but ecosystem N losses from Arctic soils through N_2O emission are generally small, which may be linked to the N limitation of these soils (Tamm 1991; Elser et al. 2007). *Oceanobacillus* sp. CF4.6 may play a role in denitrification and ecosystem loss of nitrogen in its native environment, although environmental studies are needed to characterise the soil nitrogen cycle in the High Arctic desert of Northern Greenland.

All the sequenced genomes encoded pathways for the synthesis of terpenoid backbones. Apart from *Oceanobacillus* sp. CF4.6, whose genome encoded for the mevalonate pathway, the sequenced genomes encoded for the MEP/DOXP pathway for isoprenoid backbone synthesis. Interestingly, the pathway in *N. halotolerans* CF4.12 is incomplete, as it is missing 2-C-methyl-D-erythritol 4-phosphate cytidylyltransferase (EC 2.7.7.60) encoded by *IspD*, and 2-C-methyl-D-erythritol 2,4-cyclodiphosphate synthase (EC 4.6.1.12), encoded by *IspF* (Phillips et al.

2008). *N. halotolerants* DSM 15474[T] shows the same missing enzymes in its draft genome. These

findings suggest that *N. halotolerans* may contain undescribed isoforms of these enzymes. The

synthesis of terpenoid backbones is an important pathway to produce essential and secondary

metabolites like undecaprenol phosphate and carotenoids. In fact, all isolates apart from

Oceanobacillus sp. CF4.6 and the two *Aeromicrobium* sp. had BGCs encoding for terpene-

related products, with many of them encoding for carotenoids (see next section). This pathway

may be of particular relevance in *Nesterenkonia*, as all isolates belonging to this genus had a

bright red or orange colour, suggesting a significant content of carotenoids.

The adaptations discussed in this section allow the isolates to sustain activity in

their cold, nutrient-poor, and water-scarce environment. Furthermore, most of the sequenced

genomes showed very low dDDH percentages towards their closest known relative, suggesting

that they represent previously undescribed species. Thus, we fail to reject hypotheses i and ii.

Biosynthetic gene clusters in the genomes of the sequenced isolates

The sequenced genomes were found to contain several BGCs (**Fig. 5** and **Tables**

S4-S12), including several with no hits to known BGCs. 32% of the identified BGCs putatively

encoded for antimicrobial compounds, such as the lassopeptide paeninodin found in *S.*

neizhouensis CF4.18 and *Oceanobacillus* sp. CF4.6, ε-Poly-L-lysine synthesis clusters found in

N. halotolerans CF4.12 and *Nesterenkonia* sp. CF4.4, and other BGCs encoding for lactones

found in several isolates. Most of these putatively antimicrobial clusters encoded for polyketide

synthases (PKS) and non-ribosomal peptide synthases (NRPS). Antimicrobial compounds and

antimicrobial resistance are widely distributed in soils, where competition for available space

and resources exerts significant selective pressure (Mullis et al. 2019). The production of

antimicrobial compounds typically incurs a high metabolic cost; for example, Schlatter and

Kinkel (Schlatter and Kinkel 2015) found trade-offs between efficient growth, antimicrobial

resistance and antimicrobial production in *Streptomyces spp*., and suggested that a "super-

killer" phenotype results in a reduction in niche width. The most common strategy in

Streptomyces spp. was the simultaneous production of different kinds of antimicrobial

compounds to increase their synergistic effect and slow down the evolution of resistance.

However, in carbon-poor soils, as is the case in Peary Land, nutrient limitation may present an

obstacle for this strategy. On the other hand, nutrient scarcity may increase evolutionary

pressure to inhibit potential competitors. Many studies have demonstrated the production of

antibacterial and antifungal compounds by bacteria isolated from arid soils (e.g., Saadoun et al.

2008; Nasfi et al. 2018), and even the hyper-arid Atacama desert is a potential source of novel

antimicrobials (Bull and Asenjo 2013). Polar desert soils, although not as explored, have also demonstrated this potential (van Dorst et al. 2017). In permafrost, where biomass can be higher and environmental conditions are stable, antibiotic resistance is widespread, even in ancient permafrost (Mindlin and Petrova 2017; Gomathinayagam and Muthukaliannan 2024). Although measuring the production of antimicrobial compounds of our isolates was beyond the scope of this study, the diversity of BGCs encoding for antimicrobial compounds found in the sequenced genomes suggest a competitive environment where it may be advantageous to have different antimicrobial strategies available.

Desert soils often have higher pH compared to soils from forests and other biomes. The high pH is caused by accumulation of minerals and salts that are not washed away by rain. Peary Land soil is no exception, with soils at Cape Morris Jesup and Citronen Fjord having pH of 7.8 and 9.2, respectively. At these pH values, the bioavailability of many metals is lowered. Thus, bacteria in desert soils need effective metal-acquiring systems such as siderophores. Siderophores allow for the solubilisation of iron essential for the production of, e.g., heme proteins. Both *Aeromicrobium* isolates, as well as *Arthrobacter* sp. KK5.5 had BGCs encoding for NRPS-independent siderophores; their genomes also contained orthologs of *fepBDG*, which encode for Fe-siderophore complex transporters, and both *Aeromicrobium* isolates additionally had *fepC*. Notably, *N. halotolerans* CF4.12 also had a complete set of *fepBCDG*, while the other two *Nesterenkonia* isolates had orthologues of *fepCG*. This absence of siderophore synthesis genes, while transport genes are present, may be due to a variety of causes. One possibility is that these bacteria rely on siderophores produced by other bacteria for iron uptake (Kramer et al. 2020).

Lastly, *Massilia* sp. CMS3.1 had a BGC whose putative product is a homoserine lactone. While it had no hits to known clusters, it contained, among others, a protein with an acyltransferase domain; this suggests that this BGC may encode for an N-Acyl homoserine lactone, a group of signalling molecules involved in quorum sensing and biofilm formation. This aligns with the possible ability of *Massilia* sp. CMS3.1 to form biofilms. Indeed, the genome of this isolate contained several genes involved in quorum sensing and biofilm formation, such as orthologues of *lasIR* and *wspBEFR*, as well as *pilGHIJ*, which are involved in type IV pilus formation. Young (<24h) colonies of *Massilia* sp. CMS3.1 were orange, soft and easily pickable, but older (especially >48h) colonies became darker, hard, and difficult to disaggregate, which suggests the formation of a hard extracellular matrix that may be mediated by these genes. Biofilm formation has been shown to be a successful strategy to improve survivability against extreme environmental factors like temperature and pH (high and low), aridity and high salinity

465   (Khan and Rao 2023). For example, biofilms are employed by epiphytic bacteria to survive periods

466   of extreme cold and freeze-thaw cycles (Wu et al. 2012).

467   Other types of BGCs found in the genomes included an ectoine synthesis cluster

468   and terpenoid synthesis clusters, which were discussed in the previous section; the terpenoid

469   clusters encoded mainly for carotenoids.

470   It is possible that some BGCs were missed by antiSMASH. Indeed, BGC detection

on contig-level assemblies can be challenging, as BGCs may be split into different contigs

(Meleshko et al. 2019). This is evidenced by a few BGCs that were found at the beginning or the

end of a contig, suggesting that these BGCs are incomplete. This, combined with the significant

amount of BGCs that had no hits with known BGCs, suggest that Peary Land is indeed a source

of potentially novel secondary metabolites, as per hypothesis iii; however, further investigation

in vivo will be necessary to fully characterise these secondary metabolites.

Fig. 5 *Biosynthetic Gene Clusters (BGCs) found in the assembled genomes. The dotted pattern represents BGCs with no hits to known BGCs. NRP: Non-Ribosomal Peptide. RiPP: Ribosomally synthesized and Post-translationally modified Peptide*

CONCLUSIONS

We isolated 49 bacteria from three different sites in Peary Land, northern Greenland, representing some of the most northernly bacterial isolates, and the first strains isolated from these areas. We showed the diversity of culturable halotolerant bacteria at the three sites and found potentially novel BGCs that suggest a competitive environment with many antimicrobial compounds present. The isolates showed adaptations to their cold and saline native environments, including a variety of ion transporters and genes related to osmoprotectant synthesis and transport. These findings underscore the importance of osmoprotectants in cold

and saline environments and expand our understanding of microbial life in extreme environments. Overall, this work not only provides the first insights into the diversity of culturable halotolerant bacteria of northern-most Greenland, but also suggests that the region harbours untapped microbial resources that may hold significant biotechnological value.

DATA AVAILABILITY

All sequences obtained and used in this study, including near-complete 16S rRNA gene sequences, genome assemblies, and short-read libraries, are available in the National Center for Biotechnology Information (NCBI) database under the BioProject accession number PRJNA1156031.

ACKNOWLEDGEMENTS

We would like to thank Lorrie Maccario for her help in the library preparation and sequencing of the genomes, and Anette Hørdum Løth and Ayoe Lüchau for their indispensable assistance in the lab. We would also like to thank Magnus Per Damsø Jeppesen for his help in the collection of some of the salt response data. We are grateful to the Greenlandic authorities for granting permission to sample in the Northeast Greenland National Park (Expedition Permit C-21-549) and to export the soil samples to Denmark for scientific work (Non-exclusive license number G21-031 for utilization of Greenlandic genetic resources).

FUNDING

This project was funded by the Novo Nordisk Foundation under the NNF Interdisciplinary Synergy Program (grant number NNF19OC0057374): Effects of bacteria on atmospheres of Earth, Mars, and exoplanets -- adapting and identifying life in extraterrestrial environments. This project was also supported by The Danish National Research Foundation within the Center for Volatile Interactions (VOLT, DNRF168). The bacteria in this study were isolated from samples collected during the Leister Around North Greenland 2021 Expedition financed by the Leister Stiftung, Switzerland.

CONFLICTS OF INTEREST

The authors declare no conflicts of interest.

REFERENCES

Abirami B, Manigundan K, Radhakrishnan M, et al (2021) Diversity and Bioprospecting Potentials of Antarctic (Polar) Microbes. In: Microbiome-Host Interactions. CRC Press

Acker JG, Leptoukh G (2007) Online analysis enhances use of NASA Earth science data. EoS Transactions 88:14–17. https://doi.org/10.1029/2007eo020003

Blin K, Shaw S, Augustijn HE, et al (2023) antiSMASH 7.0: new and improved predictions for detection, regulation, chemical structures and visualisation. Nucleic Acids Research 51:W46–W50. https://doi.org/10.1093/nar/gkad344

Bolger AM, Lohse M, Usadel B (2014) Trimmomatic: a flexible trimmer for Illumina sequence data. Bioinformatics 30:2114–2120. https://doi.org/10.1093/bioinformatics/btu170

Borchers HW (2011) pracma: Practical Numerical Math Functions. 2.4.4

Bull AT, Asenjo JA (2013) Microbiology of hyper-arid environments: recent insights from the Atacama Desert, Chile. Antonie van Leeuwenhoek 103:1173–1179. https://doi.org/10.1007/s10482-013-9911-7

Charlier RH (1969) The Geographic Distribution of Polar Desert Soils in the Northern Hemisphere. Geol Soc America Bull 80:1985. https://doi.org/10.1130/0016-7606(1969)80[1985:TGDOPD]2.0.CO;2

Chauhan A, Pathak A, Jaswal R, et al (2018) Physiological and Comparative Genomic Analysis of Arthrobacter sp. SRS-W-1-2016 Provides Insights on Niche Adaptation for Survival in Uraniferous Soils. Genes 9:31. https://doi.org/10.3390/genes9010031

Chen Y-H, Lu C-W, Shyu Y-T, Lin S-S (2017) Revealing the Saline Adaptation Strategies of the Halophilic Bacterium Halomonas beimenensis through High-throughput Omics and Transposon Mutagenesis Approaches. Sci Rep 7:13037. https://doi.org/10.1038/s41598-017-13450-9

Chen Y-H, Shyu Y-T, Lin S-S (2018) Characterization of candidate genes involved in halotolerance using high-throughput omics in the halotolerant bacterium Virgibacillus chiguensis. PLOS ONE 13:e0201346. https://doi.org/10.1371/journal.pone.0201346

Dai D, Lu H, Xing P, Wu Q (2022) Comparative Genomic Analyses of the Genus Nesterenkonia Unravels the Genomic Adaptation to Polar Extreme Environments. Microorganisms 10:233. https://doi.org/10.3390/microorganisms10020233

Dsouza M, Taylor MW, Turner SJ, Aislabie J (2015) Genomic and phenotypic insights into the ecology of Arthrobacter from Antarctic soils. BMC Genomics 16:36. https://doi.org/10.1186/s12864-015-1220-2

Edgar RC (2004) MUSCLE: multiple sequence alignment with high accuracy and high throughput. Nucleic Acids Res 32:1792–1797. https://doi.org/10.1093/nar/gkh340

Ellis HR (2011) Mechanism for sulfur acquisition by the alkanesulfonate monooxygenase system. Bioorganic Chemistry 39:178–184. https://doi.org/10.1016/j.bioorg.2011.08.001

Elser JJ, Bracken MES, Cleland EE, et al (2007) Global analysis of nitrogen and phosphorus
limitation of primary producers in freshwater, marine and terrestrial ecosystems. Ecology
Letters 10:1135–1142. https://doi.org/10.1111/j.1461-0248.2007.01113.x

Fernandez-Guerra A, Borrel G, Delmont TO, et al (2023) A 2-million-year-old microbial and viral
communities from the Kap København Formation in North Greenland

Finore I, Orlando P, Di Donato P, et al (2016) Nesterenkonia aurantiaca sp. nov., an alkaliphilic
actinobacterium isolated from Antarctica. International Journal of Systematic and
Evolutionary Microbiology 66:1554–1560. https://doi.org/10.1099/ijsem.0.000917

Flores-Gallegos AC, Delgado-García M, Ascacio-Valdés JA, et al (2019) Chapter 13 - Hydrolases
of Halophilic Origin With Importance for the Food Industry. In: Kuddus M (ed) Enzymes in
Food Biotechnology. Academic Press, pp 197–219

Francois R (1987) A study of sulphur enrichment in the humic fraction of marine sediments during
early diagenesis. Geochimica et Cosmochimica Acta 51:17–27.
https://doi.org/10.1016/0016-7037(87)90003-2

Funder S, Bennike E, Mogensen GS, et al (1984) The Kap København Formation, a late Cainozoic
sedimentary sequence in North Greenland. Rapport Grønlands Geologiske
Undersøgelse 120:9–18. https://doi.org/10.34194/rapggu.v120.7851

Gomathinayagam S, Muthukaliannan GK (2024) Profiling Antibiotic Resistance Determinants in
Ancient Permafrost Microbiomes. 2024.05.20.595077

Guida G, Nicosia A, Settanni L, Ferro V (2023) A review on effects of biological soil crusts on
hydrological processes. Earth-Science Reviews 243:104516.
https://doi.org/10.1016/j.earscirev.2023.104516

Gushgari-Doyle S, Lui LM, Nielsen TN, et al (2022) Genotype to ecotype in niche environments:
adaptation of Arthrobacter to carbon availability and environmental conditions. ISME
Communications 2:32. https://doi.org/10.1038/s43705-022-00113-8

He T, Xie D, Ni J, et al (2020) Characteristics of nitrogen transformation and intracellular nitrite
accumulation by the hypothermia bacterium *Arthrobacter arilaitensis*. Science of The
Total Environment 701:134730. https://doi.org/10.1016/j.scitotenv.2019.134730

Hinsa-Leasure SM, Bhavaraju L, Rodrigues JLM, et al (2010) Characterization of a bacterial
community from a Northeast Siberian seacoast permafrost sample. FEMS Microbiology
Ecology 74:103–113. https://doi.org/10.1111/j.1574-6941.2010.00945.x

Hoang DT, Chernomor O, von Haeseler A, et al (2018) UFBoot2: Improving the Ultrafast Bootstrap
Approximation. Molecular Biology and Evolution 35:518–522.
https://doi.org/10.1093/molbev/msx281

Horn G, Hofweber R, Kremer W, Kalbitzer HR (2007) Structure and function of bacterial cold
shock proteins. Cell Mol Life Sci 64:1457. https://doi.org/10.1007/s00018-007-6388-4

Hovila J, Arola A, Tamminen J (2009) OMI/Aura Surface UVB Irradiance and Erythemal Dose Daily
L3 Global 1.0x1.0 deg Grid

Iwahana G, Cooper ZS, Carpenter SD, et al (2021) Intra-ice and intra-sediment cryopeg brine occurrence in permafrost near Utqiaġvik (Barrow). Permafrost and Periglacial Processes 32:427–446. https://doi.org/10.1002/ppp.2101

Jaarsma AH, Zervas A, Sipes K, et al (2023) The undiscovered biosynthetic potential of the Greenland Ice Sheet microbiome. Front Microbiol 14:. https://doi.org/10.3389/fmicb.2023.1285791

Jansson JK, Taş N (2014) The microbial ecology of permafrost. Nat Rev Microbiol 12:414–425. https://doi.org/10.1038/nrmicro3262

Kalyaanamoorthy S, Minh BQ, Wong TKF, et al (2017) ModelFinder: fast model selection for accurate phylogenetic estimates. Nat Methods 14:587–589. https://doi.org/10.1038/nmeth.4285

Kanehisa M, Sato Y, Morishima K (2016) BlastKOALA and GhostKOALA: KEGG Tools for Functional Characterization of Genome and Metagenome Sequences. Journal of Molecular Biology 428:726–731. https://doi.org/10.1016/j.jmb.2015.11.006

Khan A, Rao TS (2023) Chapter 13 - Natural biofilms: Structure, development, and habitats. In: Das S, Kungwani NA (eds) Understanding Microbial Biofilms. Academic Press, pp 187–206

Kramer J, Özkaya Ö, Kümmerli R (2020) Bacterial siderophores in community and host interactions. Nat Rev Microbiol 18:152–163. https://doi.org/10.1038/s41579-019-0284-4

Lebre PH, De Maayer P, Cowan DA (2017) Xerotolerant bacteria: surviving through a dry spell. Nat Rev Microbiol 15:285–296. https://doi.org/10.1038/nrmicro.2017.16

Liu Y, Zhang Y, Huang Y, et al (2023) Spatial and temporal conversion of nitrogen using Arthrobacter sp. 24S4–2, a strain obtained from Antarctica. Front Microbiol 14:. https://doi.org/10.3389/fmicb.2023.1040201

Manni M, Berkeley MR, Seppey M, et al (2021) BUSCO Update: Novel and Streamlined Workflows along with Broader and Deeper Phylogenetic Coverage for Scoring of Eukaryotic, Prokaryotic, and Viral Genomes. Molecular Biology and Evolution 38:4647–4654. https://doi.org/10.1093/molbev/msab199

Meier-Kolthoff JP, Carbasse JS, Peinado-Olarte RL, Göker M (2022) TYGS and LPSN: a database tandem for fast and reliable genome-based classification and nomenclature of prokaryotes. Nucleic Acids Research 50:D801–D807. https://doi.org/10.1093/nar/gkab902

Meier-Kolthoff JP, Göker M (2019) TYGS is an automated high-throughput platform for state-of-the-art genome-based taxonomy. Nat Commun 10:2182. https://doi.org/10.1038/s41467-019-10210-3

Meleshko D, Mohimani H, Tracanna V, et al (2019) BiosyntheticSPAdes: reconstructing biosynthetic gene clusters from assembly graphs. Genome Res 29:1352–1362. https://doi.org/10.1101/gr.243477.118

Meteostat Kap Morris Jesup | Weather History & Climate. In: Meteostat.
https://meteostat.net/en/station/04301?t=2020-01-01/2023-12-31. Accessed 26 Sep
2024

Mikheenko A, Prjibelski A, Saveliev V, et al (2018) Versatile genome assembly evaluation with
QUAST-LG. Bioinformatics 34:i142–i150. https://doi.org/10.1093/bioinformatics/bty266

Mindlin SZ, Petrova MA (2017) On the Origin and Distribution of Antibiotic Resistance: Permafrost
Bacteria   Studies.   Mol   Genet   Microbiol   Virol   32:169–179.
https://doi.org/10.3103/S0891416817040048

Minh BQ, Schmidt HA, Chernomor O, et al (2020) IQ-TREE 2: New Models and Efficient Methods
for Phylogenetic Inference in the Genomic Era. Molecular Biology and Evolution 37:1530–
1534. https://doi.org/10.1093/molbev/msaa015

Mølgaard P (1982) Temperature Observations in High Arctic Plants in Relation to Microclimate in
the Vegetation of Peary Land, North Greenland. Arctic and Alpine Research 14:105–115.
https://doi.org/10.1080/00040851.1982.12004287

Mullis MM, Rambo IM, Baker BJ, Reese BK (2019) Diversity, Ecology, and Prevalence of
Antimicrobials in Nature. Front Microbiol 10:. https://doi.org/10.3389/fmicb.2019.02518

Nasfi Z, Busch H, Kehraus S, et al (2018) Soil Bacteria Isolated From Tunisian Arid Areas Show
Promising Antimicrobial Activities Against Gram-Negatives. Front Microbiol 9:.
https://doi.org/10.3389/fmicb.2018.02742

Nicholson WL, Krivushin K, Gilichinsky D, Schuerger AC (2013) Growth of Carnobacterium spp.
from permafrost under low pressure, temperature, and anoxic atmosphere has
implications   for   Earth   microbes   on   Mars.   PNAS   110:666–671.
https://doi.org/10.1073/pnas.1209793110

Niederberger TD, Steven B, Charvet S, et al (2009) Virgibacillus arcticus sp. nov., a moderately
halophilic, endospore-forming bacterium from permafrost in the Canadian high Arctic.
International Journal of Systematic and Evolutionary Microbiology 59:2219–2225.
https://doi.org/10.1099/ijs.0.002931-0

Ollivier B, Caumette P, Garcia JL, Mah RA (1994) Anaerobic bacteria from hypersaline
environments. Microbiological Reviews 58:27. https://doi.org/10.1128/mr.58.1.27-
38.1994

Parks DH, Imelfort M, Skennerton CT, et al (2015) CheckM: assessing the quality of microbial
genomes recovered from isolates, single cells, and metagenomes. Genome Res
25:1043–1055. https://doi.org/10.1101/gr.186072.114

Parvizpour S, Hussin N, Shamsir MS, Razmara J (2021) Psychrophilic enzymes: structural
adaptation, pharmaceutical and industrial applications. Appl Microbiol Biotechnol
105:899–907. https://doi.org/10.1007/s00253-020-11074-0

Pathania S, Solanki P, Putatunda C, et al (2022) Adaptation to Cold Environment: The Survival
Strategy of Psychrophiles. In: Goel R, Soni R, Suyal DC, Khan M (eds) Survival Strategies
in Cold-adapted Microorganisms. Springer, Singapore, pp 87–111

Petzoldt T (2016) growthrates: Estimate Growth Rates from Experimental Data. 0.8.4

Phillips MA, León P, Boronat A, Rodríguez-Concepción M (2008) The plastidial MEP pathway: unified nomenclature and resources. Trends in Plant Science 13:619–623. https://doi.org/10.1016/j.tplants.2008.09.003

Prjibelski A, Antipov D, Meleshko D, et al (2020) Using SPAdes De Novo Assembler. Current Protocols in Bioinformatics 70:e102. https://doi.org/10.1002/cpbi.102

Ramond J-B, Jordaan K, Díez B, et al (2022) Microbial Biogeochemical Cycling of Nitrogen in Arid Ecosystems. Microbiol Mol Biol Rev 86:. https://doi.org/10.1128/mmbr.00109-21

Ruginescu R, Purcărea C, Dorador C, et al (2019) Exploring the hydrolytic potential of cultured halophilic bacteria isolated from the Atacama Desert. FEMS Microbiology Letters 366:fnz224. https://doi.org/10.1093/femsle/fnz224

Saadoun I, Wahiby L, Ababneh Q, et al (2008) Recovery of soil streptomycetes from arid habitats in Jordan and their potential to inhibit multi-drug resistant Pseudomonas aeruginosa pathogens. World J Microbiol Biotechnol 24:157–162. https://doi.org/10.1007/s11274-007-9451-2

Sarma J, Sengupta A, Laskar MK, et al (2023) Microbial adaptations in extreme environmental conditions. In: Bacterial Survival in the Hostile Environment. Elsevier, pp 193–206

Sayers EW, Bolton EE, Brister JR, et al (2022) Database resources of the national center for biotechnology information. Nucleic Acids Res 50:D20–D26. https://doi.org/10.1093/nar/gkab1112

Schlatter DC, Kinkel LL (2015) Do tradeoffs structure antibiotic inhibition, resistance, and resource use among soil-borne Streptomyces? BMC Evolutionary Biology 15:186. https://doi.org/10.1186/s12862-015-0470-6

Schwengers O, Jelonek L, Dieckmann MA, et al (2021) Bakta: rapid and standardized annotation of bacterial genomes via alignment-free sequence identification. Microbial Genomics 7:000685. https://doi.org/10.1099/mgen.0.000685

Shcherbakova VA, Chuvil'skaya NA, Rivkina EM, et al (2009) Novel halotolerant bacterium from cryopeg in permafrost: Description of Psychrobacter muriicola sp. nov. Microbiology 78:84–91. https://doi.org/10.1134/S0026261709010111

Shen L, Liu Y, Allen MA, et al (2021) Linking genomic and physiological characteristics of psychrophilic Arthrobacter to metagenomic data to explain global environmental distribution. Microbiome 9:136. https://doi.org/10.1186/s40168-021-01084-z

Singh P, Kapse N, Gowdaman V, et al (2021) Comparative Genomic Analysis of Arctic Permafrost Bacterium Nesterenkonia sp. PF2B19 to Gain Insights into Its Cold Adaptation Tactic and Diverse Biotechnological Potential. Sustainability 13:4590. https://doi.org/10.3390/su13084590

Singh P, Singh SM, Singh RN, et al (2017) Bacterial communities in ancient permafrost profiles of Svalbard, Arctic. Journal of Basic Microbiology 57:1018–1036. https://doi.org/10.1002/jobm.201700061

708   Sivalingam P, Hong K, Pote J, Prabakar K (2019) Extreme Environment *Streptomyces* : Potential
709   Sources for New Antibacterial and Anticancer Drug Leads? International Journal of
710   Microbiology 2019:1–20. https://doi.org/10.1155/2019/5283948

711   Son J, Lee H, Kim M, et al (2021) Massilia aromaticivorans sp. nov., a BTEX Degrading Bacterium
712   Isolated from Arctic Soil. Curr Microbiol 78:2143–2150. https://doi.org/10.1007/s00284-
713   021-02379-y

714   Spirina EV, Durdenko EV, Demidov NE, et al (2017) Halophilic-psychrotrophic bacteria of an
715   Alaskan cryopeg—a model for astrobiology. Paleontol J 51:1440–1452.
716   https://doi.org/10.1134/S0031030117120036

717   Steven B, Léveillé R, Pollard WH, Whyte LG (2006) Microbial ecology and biodiversity in
718   permafrost. Extremophiles 10:259–267. https://doi.org/10.1007/s00792-006-0506-3

719   Stevenson A, Cray JA, Williams JP, et al (2015) Is there a common water-activity limit for the three
720   domains of life? ISME J 9:1333–1351. https://doi.org/10.1038/ismej.2014.219

721   Stimmler P, Priemé A, Elberling B, et al (2022) Arctic soil respiration and microbial community
722   structure driven by silicon and calcium. Science of The Total Environment 838:156152.
723   https://doi.org/10.1016/j.scitotenv.2022.156152

724   Tamm CO (1991) Nitrogen-Limited and Nitrogen-Depleted Terrestrial Ecosystems: Ecological
725   Characteristics. In: Nitrogen in Terrestrial Ecosystems. Springer Berlin Heidelberg, Berlin,
726   Heidelberg, pp 34–49

727   Tamura K, Stecher G, Kumar S (2021) MEGA11: Molecular Evolutionary Genetics Analysis Version
728   11. Molecular Biology and Evolution 38:3022–3027.
729   https://doi.org/10.1093/molbev/msab120

730   van Dorst J, Benaud N, Ferrari B (2017) New Insights into the Microbial Diversity of Polar Desert
731   Soils: A Biotechnological Perspective. In: Chénard C, Lauro FM (eds) Microbial Ecology of
732   Extreme Environments. Springer International Publishing, Cham, pp 169–183

733   Ventosa A, Quesada E, Rodriguez-Valera F, et al (1982) Numerical Taxonomy of Moderately
734   Halophilic Gram-negative Rods. Microbiology 128:1959–1968.
735   https://doi.org/10.1099/00221287-128-9-1959

736   Vishnivetskaya T, Kathariou S, McGrath J, et al (2000) Low-temperature recovery strategies for
737   the isolation of bacteria from ancient permafrost sediments. Extremophiles 4:165–173.
738   https://doi.org/10.1007/s007920070031

739   Wang H, Zhang X, Wang S, et al (2018) Massilia violaceinigra sp. nov., a novel purple-pigmented
740   bacterium isolated from glacier permafrost. International Journal of Systematic and
741   Evolutionary Microbiology 68:2271–2278. https://doi.org/10.1099/ijsem.0.002826

742   Wick RR, Judd LM, Gorrie CL, Holt KE (2017) Unicycler: Resolving bacterial genome assemblies
743   from short and long sequencing reads. PLOS Computational Biology 13:e1005595.
744   https://doi.org/10.1371/journal.pcbi.1005595

745   Wu Z, Kan FWK, She Y-M, Walker VK (2012) Biofilm, ice recrystallization inhibition and freeze-
746   thaw protection in an epiphyte community. Appl Biochem Microbiol 48:363–370.
747   https://doi.org/10.1134/S0003683812040138

748 Wutipraditkul N, Waditee R, Incharoensakdi A, et al (2005) Halotolerant Cyanobacterium
749   Aphanothece halophytica Contains NapA-Type Na+/H+ Antiporters with Novel Ion
750   Specificity That Are Involved in Salt Tolerance at Alkaline pH. Applied and Environmental
751   Microbiology 71:4176–4184. https://doi.org/10.1128/AEM.71.8.4176-4184.2005

752