Continuous flashing suppression of neural responses and population orientation coding in

macaque V1

[1]Cai-Xia Chen[1#], Xin Wang[1#], Dan-Qing Jiang[1], Shi-Ming Tang[2,3*], & Cong Yu[4*]   Mar. 03, 2026

[1] School of Psychological and Cognitive Sciences, [2] School of Life Sciences, and [3] IDG-McGovern

Institute for Brain Research, Peking University, Beijing, China, 100871; [4] Department of Psychology and

Behavioral Sciences, Zhejiang University, Hangzhou, China

Equal-contribution first authors

* Correspondence authors: Shi-Ming Tang (tangshm@pku.edu.cn) & Cong Yu (yu_cong@zju.edu.cn)

Abstract

Continuous flash suppression (CFS), a dynamic masker presented to one eye

suppressing the conscious perception of a stimulus shown to the other eye, has been

extensively used to study visual consciousness. Various studies reported high-level visual and

cognitive functions under CFS, which, however, has more recently been questioned and at

least partially attributed to low-level stimulus properties. A key but unsettled issue is the

extent to which the responses of V1 neurons, where inputs from two eyes first merge, are

affected, as severely suppressed V1 responses would not sustain high-level processing. Here,

we used two-photon calcium imaging to record the responses of large samples of V1 neurons

to a grating stimulus under CFS in awake, fixating macaques. The results revealed that CFS

substantially suppressed V1 orientation responses. Ocularity-wise, it nearly completely

eliminated the orientation responses of V1 neurons preferring the masker eye or both eyes,

while also significantly suppressing the responses of those preferring the grating eye.

Modeling analyses suggest that, under CFS, the brain retains the ability of classifying coarse

orientations, but may become less capable of reconstructing the grating stimulus.

Consequently, while CFS-suppressed orientation information still supports low-level

orientation discrimination, it may not suffice for high-level visual and cognitive processing.

Introduction

When a target stimulus is presented to one eye and a flickering Mondrian-like masker

to the other eye, the target can be rendered invisible for an extended period (Tsuchiya & Koch,

2005). This paradigm, known as continuous flash suppression (CFS), has been widely used to

investigate subconscious visual processing (Yang, Brascamp, Kang, & Blake, 2014; Moors,

Hesselmann, Wagemans, & van Ee, 2017; Pournaghdali & Schwartz, 2020). Among the most

intriguing findings are the subconscious high-level visual and cognitive functions under the

influence of CFS (e.g., Fang & He, 2005; Almeida, Mahon, Nakayama, & Caramazza, 2008;

Adams, Gray, Garner, & Graf, 2010; Mudrik, Breska, Lamy, & Deouell, 2011; Sklar et al., 2012;

Zabelina et al., 2013; Tettamanti, Conca, Falini, & Perani, 2017). For example, as reported,

priming effects are evident when the target and the invisible primer are categorically (Almeida

et al., 2008) or semantically (Zabelina et al., 2013) consistent. However, many of these

observations have been questioned by more recent studies, with at least some of the high-

level effects being attributed to low-level feature processing (Guido Hesselmann & Malach,

2011; Sakuraba, Sakai, Yamanaka, Yokosawa, & Hirayama, 2012; Gray, Adams, Hedger, Newton,

& Garner, 2013; Moors, Boelens, van Overwalle, & Wagemans, 2016; Moors et al., 2017;

Moors & Hesselmann, 2018; Pournaghdali & Schwartz, 2020; Stuit, Paffen, & Van der Stigchel,

2023).

A critical issue in this debate is the impact of CFS on V1 neuronal activity. CFS has been

hypothesized to arise from mechanisms similar to those in binocular rivalry (Tsuchiya & Koch,

2005; Yang et al., 2014; Moors et al., 2017), which likely suppress V1 responses through

interocular inhibition. Only the surviving stimulus information would then be relayed to

downstream areas for potential subconscious higher-level visual and cognitive processing

(Jiang, Costello, & He, 2007; Adams et al., 2010; Almeida, Mahon, & Caramazza, 2010).

Importantly, if V1 activity is suppressed to a sufficient degree, the low-level stimulus

information carried by the remaining V1 responses may not suffice to sustain high-level

processing of more complex stimuli defined by those low-level features.

Two prominent fMRI studies have examined the impact of CFS on V1 activity

(Watanabe et al., 2011; Yuval-Greenberg & Heeger, 2013). Watanabe et al. (2011) compared

monocular CFS masking (stimulus visible) and dichoptic CFS masking (stimulus invisible), and

reported that V1 BOLD responses were largely insensitive to stimulus visibility when attention

was carefully controlled. However, using similar experimental design, Yuval-Greenberg and

Heeger (2013) observed reduced BOLD responses in V1 under dichoptic masking, suggesting

that V1 activity changed with stimulus visibility. They attributed the difference of results

between two studies mainly to differences in number of trials and thus the statistical power

(~250 trials per condition vs. ~90 trials per condition). Nevertheless, these studies contrasted

monocular and dichoptic masking conditions to equate stimulus input while manipulating

perceptual visibility, which were not designed to quantify the pure effect of CFS on stimulus-

evoked V1 responses, that is, the difference of BOLD signals between binocular masking and

stimulus alone conditions. In contrast, original psychophysical studies (Tsuchiya & Koch, 2005;

Tsuchiya, Koch, Gilroy, & Blake, 2006) demonstrated CFS masking by contrasting the visibility of

the target stimulus with and without the presence of dichoptic mask. Consequently, the

impact of CFS on V1 activity should be larger than what has been reported by Yuval-Greenberg

and Heeger (2013).

Neurons in V1 exhibit various degrees of ocular dominance (Hubel & Wiesel, 1962),

which influences each neuron's binocular combination of monocular visual inputs from two

eyes (Kato, Bishop, & Orban, 1981; Mitchell, Carlson, Westerberg, Cox, & Maier, 2023; Zhang,

Zhao, Jiang, Tang, & Yu, 2024). In the present study, we used a with-or-without-dichoptic-

masker design similar to those used in original psychophysical studies, and examined the

extent to which V1 neuronal responses were affected by CFS and how neurons preferring the

target eye, masker eye, or both eyes were differently impacted. Using a customized two-

photon imaging setup for awake macaques (Li, Liu, Jiang, Lee, & Tang, 2017), we sampled large

neural populations at cellular resolution and measured ocular dominance for each individual

neuron. This approach enabled us to investigate the potentially differential impacts of CFS on

the responses of V1 neurons with varying ocular preferences, as well as apply machine

learning tools to understand the impacts of CFS on V1 orientation coding at the population

level.

Results

We used two-photon calcium imaging to record responses of V1 superficial neurons

from two awake, fixating macaques, each with two response fields of view (FOVs, 850 x 850

μm^2) (Fig. 1A). During the initial recording, the stimulus was a binocular 0.45-contrast square-

wave grating varying at twelve orientations and two spatial frequencies (3 & 6 cpd) (Fig. 1B). A

total of 3,564 neurons were identified through image processing, including 3,004 (84.29%)

orientation-tuned neurons that were included in following data analyses.

The same grating stimulus was then presented monocularly (Fig. 1B) to each eye to

characterize individual neurons' eye preferences. Each neuron's ocular dominance index (ODI)

was calculated as ODI = $(R_i - R_c)/(R_i + R_c)$, where R_i and R_c were the neuron's peak responses to

ipsilateral and contralateral stimulations, respectively. Neurons with an ODI at –1 or +1 would

exclusively prefer the contralateral or ipsilateral eye, while neurons with an ODI at 0 would

prefer both eyes equally. Consistent with previous findings (Hubel & Wiesel, 1962; Horton &

Hocking, 1996; Livingstone, 1996; Zhang et al., 2024), neurons with similar eye preferences

clustered together (Fig. 1C), indicating ocular dominance columns. The ODI followed unimodal

distributions (Fig. 1D), in which the majority of neurons were binocular, showing comparable

preferences for either eye. Only a small portion of neurons were monocular, being more

responsive to the ipsilateral or contralateral eye.

Figure 1. Two-photon imaging and ocular dominance mapping. **A**. Optical windows for imaging of

two macaques. Green crosses indicate the regions for viral vector injections, and yellow boxes

indicate the FOVs chosen for imaging. **B**. Stimuli used for OD mapping. A circular-windowed

square-wave grating was presented monocularly to each eye, respectively, to probe each neuron's

ODI. **C**. Ocular dominance functional maps of each FOV at single-neuron resolution showing OD

clusters. **D**. Frequency distributions of individual neurons' ODIs in each FOV.

In a third and last step, the grating stimulus and the flashing noise masker were

presented dichoptically to evaluate the impact of CFS on neurons' orientation responses (Fig.

2A). The results are summarized as population orientation tuning functions under the baseline

no-CFS condition and the CFS condition following the procedure in Busse, Wade, and Carandini

(2009). Specifically, neurons with similar orientation preferences were binned (bin width = 15°)

relative to the target orientation for a total of 12 bins, and the resultant population orientation

tuning functions based on the mean responses of these bins (Fig. 2C) were fitted with a

Gaussian function. Compared to the baseline population orientation tuning functions, those

under the influence of CFS displayed profound reductions in orientation response. The

amplitude decreased by 84.18% in Monkey A and 60.78% in Monkey B on the basis of

Gaussian fitting, while the slope decreased by 91.31% in Monkey A and 71.50% in Monkey B

(Fig. 2B). These results support the conclusion of Yuval-Greenberg and Heeger (2013) that V1

activity is degraded by CFS, 'akin' to a loss in the contrast-to-noise ratio of neural activity.

Furthermore, neurons were divided into three groups according to their ODIs, and the

impacts of CFS on their respective orientation responses were examined: neurons preferring

the grating eye (ODI > 0.2 or < -0.2, depending on whether the grating stimulation was

ipsilateral or contralateral), binocular neurons (-0.2 <= ODI <= 0.2), and neurons preferring the

masker eye (ODI < -0.2 or > 0.2 relative to the grating eye). Compared to the baseline

condition, the orientation tuning of neurons preferring the masker eye was completely wiped

out by CFS (Fig. 2B, D left), leading to flattened tuning curves with unmeasurable amplitudes

or bandwidths. The orientation tuning of binocular neurons was either nearly completely

wiped out (Monkey A) or substantially abolished (Monkey B) (Fig. 2B, D middle). There were

85.68% and 68.32% decreases in amplitude, and 92.64% and 77.07% decreases in slope, for

Monkeys A and B, respectively. The orientation tuning of neurons preferring the grating eye

was the least but still substantially affected (Fig. 2B, D right), with respective 77.78% and

41.75% decreases in amplitude and 85.23% and 57.56% decreases in slope for two monkeys.

To quantify the loss of V1 population orientation encoding due to continuous flash

suppression (CFS), we compared the Fisher information (Averbeck & Lee, 2006) under both

baseline and CFS conditions. Here, Fisher information serves as a statistical measure that

reflects how much information the responses of neurons can provide about the grating

orientation. Specifically, it indicates the sensitivity of neural responses to small changes in

orientation, in that higher values signify greater precision in encoding orientation information.

As illustrated in Fig. 2E, Fisher information was reduced by CFS primarily for orientations

deviated by less than 15° from the neurons' preferred orientations. The average Fisher

information for stimuli within this 15° range decreased to 29.1% and 43.4% of the baseline

values in two macaques, respectively (Fig. 2F), indicating the detrimental impact of CFS on the

ability of V1 populations to accurately encode and represent orientation information,

especially for orientations closely aligned with neuronal preferences.

149
150
Figure 2. The impacts of CFS on population orientation tuning in two macaques. **A**. Stimuli used in

the CFS experiment for one macaque. The grating target was presented to one eye, which was

dichoptically masked by a circular flashing masker presented to the other eye. The white dot was

the fixation point. **B**. Exemplar baseline and CFS orientation tuning functions for neurons with

different eye preferences based on Gaussian fitting. **C**. Population orientation tuning functions of all

neurons without CFS as the baseline and with CFS based on Gaussian fitting. Data from two FOVs

of each monkey were pooled due to highly consistent results. **D**. Population orientation tuning

functions of sub-groups of neurons with different eye preferences without and with CFS. Data from

two FOVs of each monkey were pooled. Curves are fitting results using an ocular dominance-

dependent gain control model. Error bars represent ±1 SE. **E**. The impacts of CFS on Fisher

information. Fisher information is plotted as a function of relative orientation (to the neuron's

preferred orientation) without and with CFS. Shaded areas denote ±1 SE. **F**. The ratio of

baseline/CFS Fisher information within 15° of neurons' preferred orientations. Data from two FOVs

of each monkey were pooled due to highly consistent results.

An ocular-dominance-dependent gain control model

We developed an ocular dominance-dependent gain control model to account for the

impact of CFS on V1 population orientation tuning. The model development followed two

steps.

Step I. Population orientation tuning functions before CFS

The population orientation tuning functions due to monocular stimulation exhibited

different amplitudes among OD groups (Fig. 2D, red curves), which could be simulated with

Equation 1, an OD-weighted Gaussian basis function:

$$\boldsymbol{R} = \boldsymbol{w}^{S} \times \left[A \times exp^{-\left(\frac{\theta}{\sigma}\right)^{2}} + B \right] \qquad\qquad \text{(Eq. 1)}$$

where parameters *A*, *σ*, and *B* corresponded to the amplitude, standard deviation, and

minimal response of the Gaussian basis function, respectively, and θ represented the

preferred orientation of a bin of neurons relative to the actual orientation of the grating

stimulus. The weight parameter *w* was the mean of linearly transformed ODIs of neurons in a

neuronal group, which equated to *(ODI +1)/2* or *1 - (ODI + 1)/2*, depending on contralateral or

ipsilateral eye grating stimulation, and ranged from 0-1. Thus, a smaller w would indicate a

higher preference for the eye seeing the grating, and a larger w would indicate a higher

preference for the unstimulated eye (or the eye seeing the flashing masker under CFS). The w

equated to 0.33, 0.50, and 0.67 in Monkey A, and 0.32, 0.5, and 0.68 in Monkey B, for the

grating eye-preferring group, binocular group, and the masker eye-preferring group,

respectively. The exponent s represented a nonlinear transformation.

Equation 1 fitted the baseline data well (Fig. 2D, red curves), resulting in goodness-of-

fit (R^2) values at 0.94 and 0.95 for the two monkeys, respectively. This indicated that the

equation captured the OD-dependent population orientation tuning characteristics of V1

neurons with monocular stimulation before CFS.

Step II. The impacts of CFS

In step II, the model introduced several binocular combination factors to account for

population orientation tuning functions under CFS.

To account for the OD-dependent changes of orientation tuning bandwidths under CFS,

a w-dependent inhibition factor w^t was introduced, which scaled the σ of the tuning functions,

changing the monocular tunings R into R':

$$\mathbf{R'} = \boldsymbol{w}^s \times \left[A \times exp^{-\left(\frac{\theta}{\sigma \times w^t}\right)^2} + B \right] \qquad \text{(Eq. 2)}$$

This allowed different groups of neurons to exhibit various degrees of orientation

tuning function broadening, capturing the pattern in which neurons preferring the eye seeing

the grating displayed a sharper population orientation tuning curve under CFS than those

preferring the eye seeing the masker.

Previous studies have shown that binocular neuronal responses can be modeled by

incorporating interocular suppression and summation processes (Kato et al., 1981; Dougherty,

Cox, Westerberg, & Maier, 2019; Zhang et al., 2024). Therefore, R' was further normalized by

the neural response to the flashing masker to simulate interocular suppression, which was the

first component of Equation 3. Additionally, the neural response to the flashing masker was

summed to simulate binocular summation, which was the second component of Equation 3.

These two components when summed, determining the final neural responses under CFS:

$$\mathbf{R}_{CFS} = \frac{R'}{aN^k} + bN^m \qquad \text{(Eq. 3)}$$

where N was the empirical neural response to the monocularly presented flashing masker

stimulation, a and b were scaling parameters, and k and m were nonlinearity parameters. The

interocular normalization by masker response led to amplitude reduction of population

orientation tuning functions for different groups of neurons, while the binocular summation

with masker response elevated the minimal responses of tuning functions to their

corresponding heights.

During the step II model fitting, the parameters A, σ, and s were inherited from the

monocular tuning fits derived from Equation 1 and served as inputs, while the parameters a, k,

b, m, and t were optimized. The model captured the CFS modulation on population orientation

tuning curves well, with R^2 = 0.99 and 0.98 for Monkeys A and B, respectively (Fig. 2D, red

curves).

Orientation classification and reconstruction under CFS

What are the impact of CFS-induced suppression on V1 orientation decoding? To

answer this question, which is crucial for understanding subconscious processing under CFS,

we trained linear decoders to classify neighboring stimulus orientations (15°) in our

experiments, as well as transformer models to reconstruct the stimulus images. Here,

orientation classification was parallel to coarse orientation discrimination, and image

reconstruction was parallel to orientation recognition, both suggesting the upper bounds of

performance assuming an ideal observer.

For orientation classification, we trained an all-pair multiclass support vector machine

(SVM) classifier to discriminate 12 orientations based on trial-by-trial population neural

responses from all trials (Allwein, Schapire, & Singer, 2000). Decoders for different FOVs,

ipsilateral/contralateral target presentations, and baseline vs. CFS conditions were trained

separately. Under the baseline condition, the decoders achieved mean classification accuracies

of 89.5 ± 2.0% and 91.5 ± 2.1% across ipsilateral and contralateral eye conditions in Monkeys A

and B, respectively, in contrast to a chance level of 8.3% (1 out of 12). Under CFS, decoding

accuracy slightly decreased in Monkey A (81.7 ± 1.9%) but remained stable in Monkey B (90.4

± 2.1%, Fig. 3A). These results suggest that under CFS, there is still sufficient information for

coarse orientation discrimination, even for Monkey A whose V1 neuronal responses were

substantially suppressed.

Next, we trained transformer models to reconstruct the grating images on the basis of

corresponding neuronal responses under baseline and CFS conditions. The motivation for this

part of the modeling work was the assumption that high-level tasks would be difficult to carry

out if the basic stimulus features forming more complex patterns were not intact. Our

transformer model contained an architecture that integrated embedding, self-attention, and

unembedding modules, as well as a fully connected feedforward layer (Fig. 3B). The model

inputs were the responses of all neurons within a FOV to the grating stimulus (ipsilateral and

contralateral presentations of the same stimulus were modeled separately), and the model

output was the reconstructed grating image. During the training process, the model typically

reached two successive learning plateaus, where the validation loss temporarily stagnated

(Fig. 3C). Moreover, the validation loss decreased more rapidly when training on the baseline

neural response data compared to the CFS data. To compare the differences, we identified the

epoch at which the validation loss of the baseline model reached 75% of its total decrease

between the two plateaus using a sigmoid fit, and then we retrained both the baseline and

CFS models up to this epoch.

The retained baseline models reconstructed the grating stimuli significantly better than

the CFS models in Monkey A, but this discrepancy was less pronounced in Monkey B (Fig. 3D),

consistent with the neuronal data that Monkey A exhibited substantially more CFS suppression

than Monkey B in terms of population orientation tuning and Fisher information (Fig. 2). We

used a structural similarity index (SSIM) (Brunet, Vrscay, & Wang, 2012) to quantify the

reconstruction performances. Across the grating-presenting ipsilateral and contralateral eyes,

the baseline models reconstructed the grating with median SSIMs of 0.52 and 0.61 for the two

FOVs of Monkey A, and 0.57 and 0.63 for the two FOVs of Monkey B, respectively, while the

corresponding SSIMs for the CFS models were 0.16 and 0.19 for Monkey A, and 0.55 and 0.53

for Monkey B (Fig. 3E).

Figure 3. Decoding consequences of CFS revealed by machine learning. **A**. Multiway orientation

classification accuracies under CFS vs. baseline conditions obtained using SVM decoders. Each

datum represents results from a contralateral or ipsilateral grating condition with a specific FOV

averaged across 10-fold cross-validations. Error bars denote 95% confidence intervals. **B**. A

diagram of the transformer model for stimulus image reconstruction. **C**. Exemplar learning curves

of transformer models under baseline and CFS conditions from two FOVs. The vertical dashed line

indicates the epoch at which the baseline model reaches 75% of its total loss decrease between

the two learning plateaus estimated using a sigmoid fit. **D**. Illustrations of corresponding

reconstructed stimulus images on the basis of learning curves in C. **E**. Box plots of SSIM scores

between the original and reconstructed images with baseline and CFS transformers. Within a FOV,

results from contralateral eye and ipsilateral eye conditions are combined.

To estimate the impact of CFS-induced V1 suppression on downstream processing, we

also recorded neuronal responses from two V2 FOVs in Monkey A (FOVs 3 & 4). As anticipated,

V2 neurons were binocular, with over 90% of them showing ODIs within the range of -0.2 to

0.2 (Fig. 4A). Similar to V1 results from the same monkey, CFS on average reduced the

amplitudes of the population orientation tuning functions by 80.05% and the slopes by

89.44% (Fig. 4B). It also reduced the Fisher information to 33.1% of the baseline value (Fig.

4C). Furthermore, we applied the same orientation classification and image reconstruction

procedures to the V2 data. For orientation classification, the SVM decoders achieved near-

perfect performance in classifying 12 orientations under both baseline and CFS conditions,

with classification accuracies exceeding 94% across all cases (Fig. 4D). In the image

reconstruction task, the baseline model outperformed the CFS model. Specifically, the baseline

transformer models reconstructed the stimulus images with the median SSIM values of 0.61

and 0.53 for the two V2 FOVs, respectively, which dropped to 0.42 and 0.18 in the CFS models

(Fig. 4E), implying poorer or failed reconstruction of stimulus images.

Figure 4. Effects of CFS on V2 orientation responses. **A**. OD maps of the two V2 FOVs of Monkey

A (MA3 & MA4). **B**. Population orientation tuning functions for all orientation-tuned neurons with

baseline and CFS conditions. Curves represent the results of Gaussian fittings. Error bars

represent ±1 SE. **C**. Fisher information as a function of the relative orientation (to the neuron's

preferred orientation) with baseline and CFS conditions. Shaded areas denote ±1 SE. Fisher

information was lower in MA4 due to higher variations in the data. **D**. Multiway orientation

classification accuracies under CFS vs. baseline conditions using SVM decoders. Each datum

represents results from a contralateral or ipsilateral grating condition with one FOV, averaged

across 5-fold cross-validations. Error bars denote 95% confidence intervals. **E**. Box plots of SSIM

scores between the original and reconstructed images with baseline and CFS transformers. Within

a FOV, results from contralateral eye and ipsilateral eye conditions are combined.

Discussion

Our study demonstrates that CFS severely compromises orientation information in V1

neurons in an ocular dominance-dependent manner. Orientation information carried by

neurons preferring the masker eye or both eyes is completely or nearly completely wiped out,

while information carried by those preferring the grating eye is partially retained.

Downstream, orientation information in V2 neurons is also substantially weakened. Linear

decoding and transformer models suggest that CFS-compromised orientation information may

still allow coarse orientation discrimination, but will most likely impair orientation recognition

when the suppression is sufficiently strong as in Monkey A. Similarly strong suppression is also

possible in Monkey B if the current grating contrast (0.45) is lower to be 0.1-0.3 as in many CFS

experiments (Watanabe et al., 2011; Yuval-Greenberg & Heeger, 2013; Lunghi & Pooresmaeili,

2023; Alais, Coorey, Blake, & Davidson, 2024).

CFS-compromised V1 orientation information transmits for downstream visual

processing, which may explain the unconscious orientation processing observed in human CFS

studies. The "invisible" orientation information can be processed, as demonstrated by

adaptation (Kanai, Tsuchiya, & Verstraten, 2006; Bahrami, Carmel, Walsh, Rees, & Lavie, 2008)

and priming (Koivisto & Grassini, 2018) studies. The adaptation aftereffect is reduced

compared to the visible condition but not entirely abolished (Kanai et al., 2006; Bahrami et al.,

2008), likely a result of the degraded orientation information surviving CFS. For the same

reason, the priming effect also decreases during trials in which the stimulus is rendered

invisible by CFS, compared to those in which the stimulus is visible or partially visible (Koivisto

& Grassini, 2018), as the degraded stimulus information provides insufficient evidence for

decision-making, resulting in a diminished priming effect (Dehaene, 2011; Gomez, Perea, &

Ratcliff, 2013).

Furthermore, our linear decoding and transformer results can help elucidate the

debate on whether visual processing still functions at the categorization level under the

influence of CFS. Previous studies have provided evidence for the preserved category

information of the target, as demonstrated by tool-specific priming effects (Almeida et al.,

2008; Almeida et al., 2010) and differential BOLD response patterns between tools and other

object categories under CFS (Guido Hesselmann, Hebart, & Malach, 2011; Tettamanti et al.,

2017). However, an intriguing question is: Do these results rather reflect low-level feature

differences between tools and other object categories? It has been reported that elongated

objects, irrespective of their categorical affiliation, elicit similar priming effects (Sakuraba et

al., 2012). Consistent with this, when tools are categorized by their shape (elongated vs. non-

elongated), only the neural response patterns elicited by elongated tools can be discriminated

from other object categories under CFS (Fogelson, Kohler, Miller, Granger, & Tse, 2014; Ludwig,

Kathmann, Sterzer, & Hesselmann, 2015). In line with this interpretation, Guido Hesselmann,

Darcy, Rothkirch, and Sterzer (2018) reported that tool-specific priming under CFS does not

reliably emerge under conditions designed to produce strong interocular suppression,

suggesting that previously observed category effects may reflect access to low-level shape

features rather than preserved category representations. Moreover, a recent study measuring

the contrast thresholds required to both break from and suppress CFS found that stimuli

exhibited similar suppression strengths across various categories (Alais et al., 2024). According

to our results, when suppression is too strong to allow for stimulus reconstruction, as in the

case of Monkey A (Fig. 3C), the orientation information under CFS may not accumulate to a

level sufficient for resolving semantic category boundaries. The latter might require somewhat

intact stimulus information, even if subconsciously. However, it could potentially assist in

category discrimination when categorical differences lie in certain low-level shape dimensions

like orientation, as coarse orientation discrimination appears unaffected by CFS suppression

(Fig. 3A).

A related issue is the dorsal-ventral CFS hypothesis, which proposes that CFS

suppression may disproportionately affect ventral visual processing while relatively preserving

dorsal pathways involved in visuomotor functions, potentially allowing category- or action-

related information to remain accessible under suppression (Fang & He, 2005). However,

subsequent fMRI studies have failed to provide consistent support for this dissociation,

reporting either stream-invariant awareness effects (Guido Hesselmann & Malach, 2011;

Ludwig et al., 2015; Tettamanti et al., 2017), residual signal in ventral rather than dorsal

regions (Guido Hesselmann et al., 2011; Fogelson et al., 2014), or residual low-level feature

information/partial visibility rather than preserved dorsal processing (Ludwig et al., 2015).

Although our study does not directly test dorsal-ventral dissociations, our V1 results provide a

constraint on what information downstream visual pathways could access under suppression.

When CFS- induced interocular suppression was strong enough and stimuli reconstruction was

markedly reduced, as in the case of Monkey A, the information required for category-level or

action-related processing may not be sufficient for high-level cortical representation.

Interocular suppression under CFS is known to vary substantially across individuals

(Yamashiro et al., 2013; Gayet & Stein, 2017; Blake, Goodman, Tomarken, & Kim, 2019). This

inter-individual variability may contribute to the heterogeneity observed in the CFS literature.

We also found that the strength of V1 response suppression during CFS differed between two

monkeys, as reflected by population orientation tuning functions (Fig. 2C), Fisher information

(Fig. 2F), and reconstruction performance by the transformer (Fig. 3E). Several experimental

factors may have contributed to the relatively weaker suppression observed in Monkey B.

Because monkeys viewed the stimuli passively, we could not determine the dominant eye for

each monkey (instead we switched the eyes and averaged the results), and the target was

presented at relatively high contrast. Both factors are known to reduce the effectiveness of

CFS suppression (Yang, Blake, & McDonald, 2010; Yuval-Greenberg & Heeger, 2013). In

addition, the random-noise masker we used might not be as effective as Mondrian patterns

(G. Hesselmann, Darcy, Ludwig, & Sterzer, 2016). If reduced stimulus contrast and a Mondrian

masker were used, we predict that CFS suppression in Monkey B would strengthen, potentially

approaching the level observed in Monkey A. Nevertheless, it is worth emphasizing that our

main conclusions are primarily based on data from Monkey A, who exhibited much stronger

CFS suppression.

Materials and Methods

Monkey preparation

390   Monkey preparation was identical to procedures reported in previous studies (Ju,

Guan, Tao, Tang, & Yu, 2020; Guan, Ju, Tao, Tang, & Yu, 2021; Zhang et al., 2024). Two rhesus

monkeys (*Macaca mulatta*, aged 5 and 6, respectively) underwent two sequential surgeries

under general anesthesia and strictly sterile conditions. During the first surgery, a 20-mm

diameter craniotomy was performed on the skull over V1. The dura was opened and multiple

tracks of 100-150 nil AAV1.hSynap.GCaMP5G.WPRE.SV40 (AV-1-PV2478, titer 2.37e13 (GC/ml),

Penn Vector Core) were pressure-injected at a depth of ~350 μm at multiple locations. The

dura was then sutured, the skull cap was re-attached with three titanium lugs and six screws,

and the scalp was sutured. After the surgery, the animal was returned to the cage and treated

with injectable antibiotics (Ceftriaxone sodium, Youcare Pharmaceutical Group, China) for one

week. Postoperative analgesia was also administered. The second surgery was performed 45

days later. A T-shaped steel frame was installed for head stabilization, and an optical window

was inserted onto the cortical surface. Data collection could start as early as one week later.

More details about the preparation and surgical procedures can be found in Li et al. (2017).

The procedures were approved by the Institutional Animal Care and Use Committee, Peking

University.

Behavioral task

After a ten-day recovery period following the second surgery, monkeys were placed in

a primate chair with head restraint. They were trained to hold fixation on a small white spot

(0.2°) with eye positions monitored by an Eyelink-1000 eye tracker (SR Research) at a 1000-Hz

sampling rate. During the experiment, trials with the eye position deviated 1.5° or more from

the fixation before stimulus offset were discarded as ones with saccades and repeated.

Visual stimuli and experimental procedures

Visual stimuli were generated with a Matlab-based Psychtoolbox-3 software (Pelli &

Zhang, 1991) and presented on a ROG Swift PG278QR monitor (refresh rate = 120 Hz,

resolution = 2560 × 1440 pixel, pixel size = 0.23 mm × 0.23 mm). The screen luminance was

linearized by an 8-bit look-up table, and the mean luminance was 47 cd/m^2. The viewing

distance was 60 cm.

A drifting square-wave grating (spatial frequency = 4 cpd, contrast = full, speed = 3

cycles/sec, starting phase = 0°, size = 0.4° in diameter) was first used to determine the

population receptive field (pRF) location, shape, and approximate size associated with a

specific FOV. The same stimulus was also monocularly presented to confirm the V1 location as

ocular dominance columns would appear. This fast process used a 4 × objective lens mounted

on the two-photon microscope and did not provide cell-specific information. The recorded V1

pRFs were centered at ~0.90° eccentricity in Monkey A and ~1.93° in Monkey B. V2 pRFs were

centered at ~0.67° in Monkey A. All pRFs were approximately circular with a diameter of 0.9°.

428   The target stimulus used in the experiments was a 0.45-contrast circular-windowed

square-wave grating. It drifted at 4 cycles per second in opposite directions perpendicular to

the orientation with a starting phase of 0°, and varied at 12 orientations (0° to 165° in 15°

increments) and two spatial frequencies (3 & 6 cpd) trial by trial. The circular envelope had a

diameter of 1°, which approximated the size of pRFs for recorded FOVs, with the edge blurred

by a linear ramp starting at a radius of 0.38°. The flashing masker was a circular white noise

pattern with a diameter of 1.89°, a contrast of 0.5, and a flickering rate of 10 Hz. The white

noise consisted of randomly generated black and white blocks (0.07° × 0.07° each). The target

grating and the flashing masker were presented through a pair of NVIDIA 3D Vision 2 active

shutter glasses. To mitigate the ghost image, a low contrast (RMS contrast = 0.08) white noise

was added to the grating. The width of the noise element was half of the bar width of the

square grating, and the white noise was regenerated every frame.

440   Each block of trials consisted of four groups of stimuli: binocular, monocular, CFS, and

flashing masker-only. In the binocular group, the grating was presented to both eyes

simultaneously. The relevant data were only used to help identify ROIs and orientation-tuned

neurons along with data from other stimulus conditions. In the monocular group, the grating

was monocularly presented to the contralateral or ipsilateral eye, which served as the baseline

conditions without the influences of CFS. In the CFS group, the grating and flashing masker

were presented dichoptically. In the flashing masker-only group, the flashing masker was

presented monocularly to either eye. Each stimulus condition was repeated for 10-12 trials.

For conditions involving the grating, the trials were split for two opposite drifting directions. A

block of trial contained 242 trials, two trials for each stimulus condition, with the order of

stimulus conditions arranged in a pseudorandom manner. There were 5 to 6 blocks of trials

with each FOV.

Each stimulus was presented for 1000 ms, followed by an inter-stimulus interval of

1500 ms, allowing sufficient time for the calcium signals to return to the baseline level (Guan,

Zhang, Zhang, Tang, & Yu, 2020). For each FOV, the recording was completed in a single

session with 5-6 experiment blocks and lasted for 2-3 hours.

Two-photon imaging

Two-photon imaging was performed using a FENTOSmart two-photon microscope

(Femtonics), along with a Ti:sapphire laser (Mai Tai eHP, Spectra Physics). GCaMP5 was chosen

as the indicator of calcium signals because the fluorescence activities it expresses are linearly

proportional to neuronal spike activities within a wide range of firing rates from 10-150 Hz (Li

et al., 2017). During imaging, a 16× objective lens (0.8 N.A., Nikon) with a resolution of 1.6

μm/pixel was used, along with a 1000 nm femtosecond laser. A fast resonant scanning mode

(32 fps) was chosen to obtain continuous images of neuronal activity (8 frames per second

after averaging every 4 frames). The strength of fluorescent signals (mean luminance of a

small area) was monitored and adjusted if necessary for the drift of fluorescent signals. Two

response fields of view (FOVs) measuring 850 × 850 μm^2 in V1 were selected in both

macaques, and two FOVs of the same size in V2 were selected in Macaque A.

Imaging data analysis: Initial screening of ROIs

Data were analyzed with customized MATLAB codes. A normalized cross-correlation

based translation algorithm was used to reduce motion artifacts (Li et al., 2017). Then the

fluorescence changes were associated with corresponding visual stimuli through the time

sequence information recorded by Neural Signal Processor (Cerebus system, Blackrock

Microsystem). By subtracting the mean of the 4 frames before stimuli onset ($F0$) from the

average of the 6th-9th frames after stimuli onset (F) across 5 or 6 repeated trials for the same

stimulus condition (same orientation, spatial frequency, size, and drifting direction), the

differential image ($\Delta F = F - F0$) was obtained.

For a specific FOV, the regions of interest (ROIs) or possible cell bodies were decided

through sequential analysis of 242 differential images in the order of CFS, monocular,

binocular, and flashing masker-only conditions. CFS conditions consisted of 96 ($2\times2\times12\times2 =$

96) differential images, with the grating presented to either eye (2), at two spatial frequencies

(2), twelve orientations (12), and two motion directions (2). Monocular conditions were

identical to the CFS conditions except that the flashing masker was absent. In the binocular

conditions, gratings at two spatial frequencies (2), twelve orientations (12), and two motion

directions (2) were binocularly presented, resulting in 48 differential images. The flashing

masker-only conditions consisted of the flashing masker presented to either eye, resulting in 2

differential images.

The first differential image was filtered with a band-pass Gaussian filter (size = 2–10

pixels), and connected subsets of pixels (>25 pixels, which would exclude smaller vertical

neuropils) with average pixel value >3 standard deviations of the mean brightness were

selected as ROIs. Then the areas of these ROIs were set to mean brightness in the next

differential image before the bandpass filtering and thresholding were performed. This

measure gradually reduced the standard deviations of differential images and facilitated the

detection of neurons with relatively low fluorescence responses. If a new ROI and an existing

ROI from the previous differential image overlapped, the new ROI would be on its own if the

overlapping area OA < 1/4 ROI_{new}, discarded if 1/4 ROI_{new} < OA < 3/4 ROI_{new}, and merged with

the existing ROI if OA > 3/4 ROI_{new}. The merges would help smoothen the contours of the final

ROIs. This process went on through all differential images twice to select ROIs. Finally, the

roundness for each ROI was calculated as:

$$Roundness = \frac{\sqrt{4\pi \times A}}{P}$$

where A was the ROI's area, and P was the perimeter. Only ROIs with roundness larger than

0.9, which would exclude horizontal neuropils, were selected for further analysis.

Imaging data analysis: Orientation tuning and ocular dominance

The ratio of fluorescence change (*ΔF/F0*) was calculated as a neuron's response to a

specific stimulus condition. For a specific neuron's response to a specific stimulus condition,

the *F0n* of the n-th trial was the average of 4 frames before stimulus onset (-500 - 0 ms), and

Fn was the average of the *5th-8th, 6th-9th, or 7th-10th* frames after stimulus onset, whichever

was the greatest. *F0n* was then averaged across 10 or 12 repeated trials to obtain the baseline

F0 for all trials (to reduce noise in the calculation of responses), and *ΔFn/F0 = (Fn- F0)/F0* was

taken as the neuron's response to this stimulus at the n-th trial.

Several steps were taken to determine whether a neuron was orientation-selective. For

each monocular or binocular condition, the orientation and SF eliciting the maximal response

were designated as the neuron's preferred SF and orientation. We then compared responses

across all 12 orientations at the preferred SF by performing a non-parametric Friedman test to

determine whether the neuron's responses at various orientations were significantly different

from each other. To reduce Type I errors, the significance level was set at α = 0.01. Neurons

that passed the Friedman test at least under one viewing condition were selected as

orientation-tuned neurons.

The ocular dominance index (ODI) was calculated to characterize each neuron's eye

preference: ODI = $(R_i - R_c)/(R_i + R_c)$, where R_i and R_c were the neuron's peak responses at the

best orientation and SF to ipsilateral and contralateral monocular grating conditions,

respectively. Neurons with an ODI of -1 or 1 would be completely contralateral or ipsilateral

eye dominant, and of 0 would be equally dominant by both eyes.

Population orientation tuning

For each neuron, neural responses at the preferred SF were selected for tuning

analysis. To derive population orientation tuning curves under a specific condition, we

categorized neurons into twelve orientation preference bins according to their preferred

orientations (bin width = 15°). For each orientation presented, the responses of all orientation

preference bins were reorganized according to the relative orientation preference.

Subsequently, neuronal responses of the same relative orientation preference were averaged

to generate the final population orientation tuning function. For CFS conditions, the selected

SF and binning procedures were the same as their corresponding monocular conditions.

The population orientation tuning function was fitted with a Gaussian model with

MATLAB's nonlinear least-squares function 'lsqnonlin':

538
$$R(\theta) = a \times 2^{-\left(\frac{\theta - \theta_0}{\sigma}\right)^2} + b$$

where R(θ) was the response at orientation θ, free parameters a, θ_0, σ, and b were the

amplitude, peak orientation, standard deviation of the Gaussian function (equal to half width

at half height), and minimal response, respectively.

The population orientation tuning curves for different eye preference groups were

derived using the same procedure, with additional binning of neurons according to their ODI.

To obtain the tuning curve of the neurons preferring the eye seeing the grating, responses of

neurons with an ODI < -0.2 (preferentially responding to the contralateral eye) under

contralateral eye grating presentation and those with an ODI > 0.2 (preferentially responding

to the ipsilateral eye) under ipsilateral eye grating presentation were combined. Similarly, for

neurons preferring the eye seeing the masker, responses of neurons with an ODI < -0.2 under

ipsilateral eye grating presentation and those with an ODI > 0.2 under contralateral eye grating

presentation were combined. For binocular neurons (-0.2 < ODI < 0.2), responses under both

grating presentation conditions were combined.

Fisher information

The Fisher information assesses the amount of information contained in a neuron

population using an optimal decoder (Pouget, Deneve, Ducom, & Latham, 1999). Assume

independent Gaussian noise distributions, the Fisher information for a population of N

neurons was given as

$$I = \sum_{i=1}^{N} \frac{f_i'(\theta)^2}{\sigma_i(\theta)^2}$$

where $f_i(\theta)$ was the mean activity of neuron i in response to the presentation angle, θ, and

$f_i'(\theta)$ was its derivative with respect to θ. We fitted each neuron's response tuning $f_i(\theta)$

and variance tuning $\sigma_i(\theta)$ with Gaussian functions and calculated the averaged Fisher

information across neurons at each orientation.

SVM-based orientation classification

In the orientation classification task (Fig. 3A), we trained a support vector machine

(SVM) with a one-vs.-one coding scheme to classify orientations from standardized population

neural activity. The SVM decoder was implemented using MATLAB's '*fitcsvm*' function with a

linear kernel. To prevent overfitting and evaluate the generalization ability of the model, we

employed a 5-fold cross-validation procedure, and the model performance on the validation

dataset was reported.

Decoders were trained independently for each experiment condition, resulting in 4

models per FOV (contralateral/ipsilateral × baseline/CFS). Neural response data from two

spatial frequencies were used as input, with each neuron treated as a feature. In this way,

each model were trained and tested on 288 or 240 samples (2 SFs × 12 orientations × 12/10

repeats).

The transformer model

Model input, output, and training procedure

We implemented a transformer-based model to reconstruct grating stimuli from

population neuronal responses recorded under different experiment conditions. The model

input was a vector of neuronal responses, each corresponding to an individual neuron, and the

output was the reconstructed grating image of size 70 × 70 pixels.

The transfromer was trained independently for each experimental condition, resulting

in four models per FOV (contralateral/ipsilateral × baseline/CFS). Pilot experiments revealed

that our original dataset was insufficient for the model to converge. To address this, we

augmented the dataset to four times its original size before training. Augmentation was

performed by sampling from a normal distribution centered at each neuron's response mean,

with a standard deviation equal to its original standard deviation. Within the augmented

dataset, 6% was reserved for validation. Responses were normalized to [0, 1] before fed into

the model.

We implemented a two-phase training procedure to assess the reconstruction ability of

models trained on different neural data. During the training process, the model typically

reached two learning plateaus, where the validation loss temporarily stagnated (Fig. 3C). In

the first training session, we analyzed the learning curve to determine the epoch at which the

baseline model's validation loss had completed 75% of its total decrease between the two

plateaus. This was estimated using a modified sigmoid fit:

597
$$\text{loss(t)} = A - \frac{C}{1 + e^{k(b-t)}}$$

598   where A and C defined the function range, b was the symmetry point, k was the steepness

599   parameter, and t represented the epoch number, counted from epoch 500 (initial epochs were

600   discarded due to a drastic drop in validation loss across all training runs, see Fig. 3C). The 75%

601   decrease point was computed as:

602
$$\text{Stop point } = \text{ b} + \frac{\ln(3)}{k}$$

In the second training session, we retrained both models up to the identified epoch

and evaluated their performances.

The model was trained to minimize the mean squared error (MSE) between the

reconstructed and actual stimuli. Optimization was performed using RMSprop with a learning

rate of 0.00005 and a smoothing factor $\rho = 0.85$.

Model structure

Each neuron's response was embedded into a higher-dimensional space using a

learned weight vector as follows:

611
$$R^1 = W^{emb} \odot R^0$$

where the R^0 (n×1) represented the original response vector from n neurons, and W^{emb} (n×

d_{model}) was the embedding weight matrix, with each row corresponding to a neuron-specific

weight vector. Here we used d_{model} = 2. The symbol ⊙ denoted row-wise multiplication, such

that the ith response r_i^0 was multiplied by both elements in its embedding weight vector

w_i^{emb}. The resulting embedding matrix R^1 (n×d_{model}) contained the high-dimensional

representations of the neuronal responses.

The enriched embedding matrix was then passed through a self-attention module. In

this module, R^1 was first projected into queries (Q), keys (K), and values (V) through

independent learnable weight matrices, respectively. Then the attention map was computed

as:

622
$$AttentionMap = SoftMax(\frac{QK^T}{\sqrt{d_k}})$$

623   where d_k represented the dimensionality of the key vectors, which scaled the dot-product to

624   control the variance of the attention scores.

625   The output of self-attention was calculated as:

626
$$R^2 = AttentionMap \times V$$

The output from self-attention was unembedded by projecting each neuron's high-

dimensional representation back to one-dimensional. A feedforward layer transformed the

unembedded vector into a stimulus vector, which was then reshaped into the final 70 × 70

image.

Model evaluation

The original, non-augmented data was used for analysis, which had been seen during

training in both the training and validation sets. We used a structural similarity index (SSIM)

(Brunet et al., 2012) to quantify the reconstruction performances.

The SSIM (Brunet et al., 2011) between two images x and y (both 70×70) is defined as:

$$SSIM(x, y) = \frac{(2\mu_x\mu_y + c_1)(2\sigma_{xy} + c_2)}{(\mu_x^2 + \mu_y^2 + c_1)(\sigma_x^2 + \sigma_y^2 + c_2)}$$

where μ_x and μ_y are the mean intensities, σ_x^2 and σ_y^2 are the variances, σ_{xy} is the covariance,

and c_1 and c_2 are constants for numerical stability.

Acknowledgments

This study was supported by a Natural Science Foundation of China STI2030-Major

Projects grant (2022ZD0204600) to SMT and CY.

Competing interests

None.

References

Adams, W. J., Gray, K. L. H., Garner, M., & Graf, E. W. (2010). High-Level Face Adaptation
Without Awareness. *Psychological Science, 21*, 205-210.
Alais, D., Coorey, J., Blake, R., & Davidson, M. J. (2024). A new 'CFS tracking' paradigm
reveals uniform suppression depth regardless of target complexity or salience. *eLife,*
12, RP91019.
Allwein, E. L., Schapire, R. E., & Singer, Y. (2000). Reducing multiclass to binary: A unifying
approach for margin classifiers. *Journal of machine learning research, 1*, 113-141.
Almeida, J., Mahon, B. Z., & Caramazza, A. (2010). The Role of the Dorsal Visual Processing
Stream in Tool Identification. *Psychological Science, 21*, 772-778.
Almeida, J., Mahon, B. Z., Nakayama, K., & Caramazza, A. (2008). Unconscious processing
dissociates along categorical lines. *Proceedings of the National Academy of Sciences*
of the United States of America, 105, 15214-15218.
Averbeck, B. B., & Lee, D. (2006). Effects of Noise Correlations on Information Encoding and
Decoding. *Journal of Neurophysiology, 95*, 3633-3644.
Bahrami, B., Carmel, D., Walsh, V., Rees, G., & Lavie, N. (2008). Unconscious orientation
processing depends on perceptual load. *Journal of Vision, 8*, 12-12.
Blake, R., Goodman, R., Tomarken, A., & Kim, H.-W. (2019). Individual differences in
continuous flash suppression: Potency and linkages to binocular rivalry dynamics.

Vision Research, 160, 10-23.

Brunet, D., Vrscay, E. R., & Wang, Z. (2012). On the Mathematical Properties of the Structural
Similarity Index. *IEEE Transactions on Image Processing, 21*, 1488-1499.

Busse, L., Wade, A. R., & Carandini, M. (2009). Representation of concurrent stimuli by
population activity in visual cortex. *Neuron, 64*, 931-942.

Dehaene, S. (2011). Conscious and Nonconscious Processes:Distinct Forms of Evidence
Accumulation? In Vincent Rivasseau (Ed.), *Biological Physics: Poincaré Seminar 2009*
(pp. 141-168). Basel: Springer Basel.

Dougherty, K., Cox, M. A., Westerberg, J. A., & Maier, A. (2019). Binocular Modulation of
Monocular V1 Neurons. *Current Biology, 29*, 381-391.e384.

Fang, F., & He, S. (2005). Cortical responses to invisible objects in the human dorsal and
ventral pathways. *Nature Neuroscience, 8*, 1380-1385.

Fogelson, S. V., Kohler, P. J., Miller, K. J., Granger, R., & Tse, P. U. (2014). Unconscious neural
processing differs with method used to render stimuli invisible. *5*.

Gayet, S., & Stein, T. (2017). Between-Subject Variability in the Breaking Continuous Flash
Suppression Paradigm: Potential Causes, Consequences, and Solutions. *Frontiers in*
Psychology, Volume 8 - 2017.

Gomez, P., Perea, M., & Ratcliff, R. (2013). A diffusion model account of masked versus
unmasked priming: Are they qualitatively different? *Journal of Experimental*
Psychology: Human Perception and Performance, 39, 1731-1740.

Gray, K. L., Adams, W. J., Hedger, N., Newton, K. E., & Garner, M. (2013). Faces and
awareness: low-level, not emotional factors determine perceptual dominance.
Emotion, 13, 537-544.

Guan, S.-C., Ju, N.-S., Tao, L., Tang, S.-M., & Yu, C. (2021). Functional organization of spatial
frequency tuning in macaque V1 revealed with two-photon calcium imaging.
Progress in Neurobiology, 205, 102-120.

Guan, S.-C., Zhang, S.-H., Zhang, Y.-C., Tang, S.-M., & Yu, C. (2020). Plaid detectors in
macaque V1 revealed by two-photon calcium imaging. *Current Biology, 30*, 934-940.

Hesselmann, G., Darcy, N., Ludwig, K., & Sterzer, P. (2016). Priming in a shape task but not in
a category task under continuous flash suppression. *J Vis, 16*, 17.

Hesselmann, G., Darcy, N., Rothkirch, M., & Sterzer, P. (2018). Investigating masked priming
along the "vision-for-perception" and "vision-for-action" dimensions of unconscious
processing. *Journal of Experimental Psychology: General, 147*, 1641.

Hesselmann, G., Hebart, M., & Malach, R. (2011). Differential BOLD activity associated with
subjective and objective reports during "blindsight" in normal observers. *Journal of*
Neuroscience, 31, 12936-12944.

Hesselmann, G., & Malach, R. (2011). The link between fMRI-BOLD activation and perceptual
awareness is stream-invariant in the human visual system. *Cerebral Cortex, 21*, 2829-
2837.

Horton, J. C., & Hocking, D. R. (1996). Intrinsic Variability of Ocular Dominance Column
Periodicity in Normal Macaque Monkeys. *The Journal of Neuroscience, 16*, 7228.

Hubel, D. H., & Wiesel, T. N. (1962). Receptive fields, binocular interaction and functional
architecture in the cat's visual cortex. *The Journal of Physiology, 160*, 106-154.

Jiang, Y., Costello, P., & He, S. (2007). Processing of Invisible Stimuli: Advantage of Upright
Faces and Recognizable Words in Overcoming Interocular Suppression. *Psychological*
Science, 18, 349-355.

Ju, N.-S., Guan, S.-C., Tao, L., Tang, S.-M., & Yu, C. (2020). Orientation Tuning and End-

stopping in Macaque V1 Studied with Two-photon Calcium Imaging. *Cerebral Cortex,*
31, 2085-2097.
Kanai, R., Tsuchiya, N., & Verstraten, F. A. J. (2006). The Scope and Limits of Top-Down
Attention in Unconscious Visual Processing. *Current Biology, 16*, 2332-2336.
Kato, H., Bishop, P. O., & Orban, G. A. (1981). Binocular interaction on monocularly
discharged lateral geniculate and striate neurons in the cat. *J Neurophysiol, 46*, 932-
951.
Koivisto, M., & Grassini, S. (2018). Unconscious response priming during continuous flash
suppression. *PLOS ONE, 13*, e0192201.
Li, M., Liu, F., Jiang, H., Lee, T. S., & Tang, S. (2017). Long-term two-photon imaging in awake
macaque monkey. *Neuron, 93*, 1049-1057.
Livingstone, M. S. (1996). Ocular dominance columns in New World monkeys. *The Journal of*
Neuroscience, 16, 2086.
Ludwig, K., Kathmann, N., Sterzer, P., & Hesselmann, G. (2015). Investigating category- and
shape-selective neural processing in ventral and dorsal visual stream under
interocular suppression. *Human Brain Mapping, 36*, 137-149.
Lunghi, C., & Pooresmaeili, A. (2023). Learned value modulates the access to visual
awareness during continuous flash suppression. *Scientific Reports, 13*, 756.
Mitchell, B. A., Carlson, B. M., Westerberg, J. A., Cox, M. A., & Maier, A. (2023). A role for
ocular dominance in binocular integration. *Current Biology, 33*, 3884-3895.e3885.
Moors, P., Boelens, D., van Overwalle, J., & Wagemans, J. (2016). Scene Integration Without
Awareness: No Conclusive Evidence for Processing Scene Congruency During
Continuous Flash Suppression. *Psychol Sci, 27*, 945-956.
Moors, P., & Hesselmann, G. (2018). A critical reexamination of doing arithmetic
nonconsciously. *Psychon Bull Rev, 25*, 472-481.
Moors, P., Hesselmann, G., Wagemans, J., & van Ee, R. (2017). Continuous Flash Suppression:
Stimulus Fractionation rather than Integration. *Trends Cogn Sci, 21*, 719-721.
Mudrik, L., Breska, A., Lamy, D., & Deouell, L. Y. (2011). Integration without awareness:
expanding the limits of unconscious processing. *Psychol Sci, 22*, 764-770.
Pelli, D. G., & Zhang, L. (1991). Accurate control of contrast on microcomputer displays.
Vision Research, 31, 1337-1350.
Pouget, A., Deneve, S., Ducom, J.-C., & Latham, P. E. (1999). Narrow Versus Wide Tuning
Curves: What's Best for a Population Code? *Neural Computation, 11*, 85-90.
Pournaghdali, A., & Schwartz, B. L. (2020). Continuous flash suppression: Known and
unknowns. *Psychonomic Bulletin & Review, 27*, 1071-1103.
Sakuraba, S., Sakai, S., Yamanaka, M., Yokosawa, K., & Hirayama, K. (2012). Does the human
dorsal stream really process a category for tools? *Journal of Neuroscience, 32*, 3949-
3953.
Sklar, A. Y., Levy, N., Goldstein, A., Mandel, R., Maril, A., & Hassin, R. R. (2012). Reading and
doing arithmetic nonconsciously. *Proc Natl Acad Sci U S A, 109*, 19614-19619.
Stuit, S. M., Paffen, C. L. E., & Van der Stigchel, S. (2023). Prioritization of emotional faces is
not driven by emotional content. *Sci Rep, 13*, 549.
Tettamanti, M., Conca, F., Falini, A., & Perani, D. (2017). Unaware processing of tools in the
neural system for object-directed action representation. *Journal of Neuroscience, 37*,
10712-10724.
Tsuchiya, N., & Koch, C. (2005). Continuous flash suppression reduces negative afterimages.
Nature Neuroscience, 8, 1096-1101.

Tsuchiya, N., Koch, C., Gilroy, L. A., & Blake, R. (2006). Depth of interocular suppression
associated with continuous flash suppression, flash suppression, and binocular
rivalry. *Journal of Vision, 6*, 6-6.

Watanabe, M., Cheng, K., Murayama, Y., Ueno, K., Asamizuya, T., Tanaka, K., & Logothetis, N.
(2011). Attention But Not Awareness Modulates the BOLD Signal in the Human V1
During Binocular Suppression. *Science, 334*, 829-831.

Yamashiro, H., Yamamoto, H., Mano, H., Umeda, M., Higuchi, T., & Saiki, J. (2013). Activity in
early visual areas predicts interindividual differences in binocular rivalry dynamics.
Journal of Neurophysiology, 111, 1190-1202.

Yang, E., Blake, R., & McDonald, J. E., II. (2010). A New Interocular Suppression Technique for
Measuring Sensory Eye Dominance. *Investigative Ophthalmology & Visual Science,*
51, 588-593.

Yang, E., Brascamp, J., Kang, M.-S., & Blake, R. (2014). On the use of continuous flash
suppression for the study of visual processing outside of awareness. *Frontiers in*
Psychology, 5.

Yuval-Greenberg, S., & Heeger, D. J. (2013). Continuous Flash Suppression Modulates
Cortical Activity in Early Visual Cortex. *The Journal of Neuroscience, 33*, 9635.

Zabelina, D. L., Guzman-Martinez, E., Ortega, L., Grabowecky, M., Suzuki, S., & Beeman, M.
(2013). Suppressed semantic information accelerates analytic problem solving.
Psychon Bull Rev, 20, 581-585.

Zhang, S.-H., Zhao, X.-N., Jiang, D.-Q., Tang, S.-M., & Yu, C. (2024). Ocular dominance-
dependent binocular combination of monocular neuronal responses in macaque V1.
eLife, 13, RP92839.

783