Behavioral algorithms of ontogenetic switching in larval and juvenile zebrafish phototaxis

Authors

Maxim Q. Capelle[1,2,3,4] *, Katja Slangewal[1,2,4], Armin Bahl[1,2,3,4] *

Affiliations

[1] Department of Biology, University of Konstanz

[2] Centre for the Advanced Study of Collective Behaviour, University of Konstanz

[3] Zukunftskolleg, University of Konstanz

[4] IMPRS, Max Planck Institute of Animal Behavior

* Correspondence to maxim.capelle@uni-konstanz.de; armin.bahl@uni-konstanz.de

Abstract

Animals undergo major behavioral adjustments during ontogeny, but the cognitive and neural computations controlling these processes remain elusive. Here, we describe that zebrafish transition from light-seeking to dark-seeking, as they grow from larval to juvenile stage, within the first few weeks of their life. We apply a combination of complementary phototaxis assays in virtual reality and modeling to dissect the algorithmic basis of this transition. We identify three parallel pathways, one analyzing whole-field luminance levels, one comparing spatial light levels across eyes, and one computing eye-specific temporal derivatives. Larvae mostly use the latter two spatio-temporal computations for navigation, while juveniles largely employ the first one. We build a library of agent-based models to predict animal behavior across stimulation conditions and in more complex environments. Model-based extraction of latent cognitive variables points towards potential neural correlates of the observed behavioral inversion and illustrates a novel way to explore the mechanisms of vertebrate ontogeny. We suggest that zebrafish phototaxis is regulated via parallel processing streams, which could be a universal implementation to change strategies depending on developmental stage, context, or internal state, making behavior flexible and goal-oriented.

Keywords

Zebrafish (*Danio rerio*); behavior; phototaxis; ontogeny; gradient navigation; agent-based simulations

Introduction

Animal behavior changes during development and in response to different environmental constraints. For example, growing up from infancy to adulthood, humans undergo a long phase of immaturity in which they crucially depend on others for food and protection (*1*). Rodent reflexive responses, anxiety behavior, and memory formation change with development (*2–4*). Locusts display increased threat-induced hiding behavior over early-life stages (*5*). While *Drosophila* larvae avoid light (*6, 7*), adult *Drosophila* are attracted by light (*8, 9*). It remains unknown whether such ontogenetic behavioral transitions involve remodeling circuit mechanisms or if they can be explained through adjustments of the algorithmic strategies underlying sensory-motor decision-making.

An excellent system for the study of ontogeny is the vertebrate zebrafish (*10*). This sweetwater fish shows rapid behavioral adjustments in early development, while continuously interacting with the environment and without the need for parental care. These features provide unique advantages compared to insects and mammalian systems. The majority of insects undergo fundamental structural changes during metamorphosis. While some neural structures remain intact, most of the insect nervous system and body plan get completely remodeled (*11*), making it challenging to interpret insect ontogeny. Mammalian development, in general, requires complex social interactions with the parental generation (*12*), largely complicating experimentation.

The visual system and motor control elements of zebrafish become fully functional a few days after hatching (*13–16*), allowing animals to perform a large range of visually driven behaviors (*17*). Around three weeks of age, zebrafish start to engage in social interactions (*18–23*) and display robust learning and memory abilities (*24–26*). In addition to these behavioral transitions, it has been shown that even certain innately present sensory-motor responses, like the optomotor response and phototaxis, may invert while the animals grow from larval to juvenile to adult stage (*27–30*). However, quantitative behavioral analyses of such inversions are largely absent.

In particular, phototaxis—the tendency to alter behavioral responses to light—is well-suited for the study of ontogeny (*10*). Luminance stimuli are easy to control, allow for rich variations of spatial and temporal features, and can be used to probe complex gradient tracking strategies and search behavior. Larval zebrafish display positive phototaxis by approaching brighter regions (*31, 32*). Many studies have explored potential behavioral algorithms and neural implementations of this behavior. When presented with spatial contrast stimuli locked to the position and orientation of the fish, larvae swim towards the brighter side (*33, 34*), which suggests that spatial contrast across the eyes plays a major role in phototaxis. Temporal luminance dynamics have been shown to provide additional cues for gradient navigation. In response to sudden decreases in whole-field brightness, animals swim more vigorously, and with larger and more frequent turns (*32, 35–38*). These behaviors can be modulated by adapting animals to different brightness levels (*35, 39, 40*), indicating that phototaxis flexibly depends on the recent environmental context. With advances in virtual reality for high-throughput zebrafish experimentation (*21*), it is now becoming possible to extend behavioral assays towards more complex stimulus arrangements and beyond the larval stage.

Quantitative behavioral experiments and computational modeling allow for the extraction and validation of the algorithmic processes underlying ontogeny. In addition, they assist in the mapping of how corresponding cognitive parameters change over time. Such analyses provide a theoretical basis to interpret potential ecologically relevant reasons for the observed changes and guide experiments to assess their mechanistic implementation (*41, 42*). Several models have been suggested to explain phototaxis in larval zebrafish: Fish may turn away from the eye that experiences the greater reduction in light intensity (*39*). Whole-field brightness change magnitudes may determine whether animals should swim forward or make a turn (*34*). Brightness levels in each eye may also be compared with a dynamically evolving eye-specific luminance set point to determine turn direction (*40*). Focusing on purely temporal aspects of the behavior, navigational strategies could also employ the tendency of

animals to repeat the same turn direction upon sudden decreases in brightness, thus enabling the
return to the original location (*37*). However, none of these previous works have systematically
validated their models over multiple stimulation configurations, in more complex environments, and
across the ontogenic timeline.

In this study, we use phototaxis in zebrafish transitioning from the larval to the juvenile stage. We
find a fundamental shift in behavior as animals mature: larvae spend more time in bright areas,
whereas juveniles spend more time in the dark. Using specifically designed visual stimuli, we probe
the influence of luminance on swim statistics. Separating temporal and spatial features enables us to
disentangle three parallel sensory-motor pathways and build a universal computational model of
phototaxis, which allows generating testable predictions in new brightness environments. Our
proposed model indicates that animals transition from a spatial contrast comparison to a whole-field
luminance averaging strategy. Combining high-throughput tracking experiments with computational
modeling in the context of zebrafish phototaxis, our study provides a general framework to describe
how behavioral strategies evolve during ontogeny.

Results

Inversion of phototactic behavior during zebrafish development

To establish how zebrafish phototaxis changes across two developmental time points, we
characterized the behavior of larval (5 days-post-fertilization; dpf) and juvenile zebrafish (26–27 dpf)
in different brightness environments. We placed fish in opaque watch glasses, where they could swim
freely while their body position and orientation were continuously tracked at high framerates (**Fig. 1A**).
Our setups allow for the presentation of well-controlled visual stimuli—either locked in closed-loop
relative to the fish or in a static open-loop configuration. Carefully designed experiments using such
stimuli enable us to build, constrain, and validate behavioral algorithmic models of phototaxis across
development (**Fig. 1B**). To probe behavioral responses to light, we used four static luminance stimuli
with different spatial characteristics (**Fig. 1C**): two half-circles with distinct brightness levels on each
side (dark vs. bright); a circular gradient that decreased from bright to dark from 0 to 180 degrees and
then increased back to bright from 180 to 360 degrees; an inward gradient where brightness increased
from the rim towards the center; and an outward gradient where brightness increased from the center
to the rim. We tracked individual fish over time (**Fig. 1D**), allowing us to compute the fraction of time
animals spend at different locations across the arena (**Fig. 1E**). We also performed experiments in a
uniform luminance control stimulus, showing largely homogeneous position distributions for larvae
and juveniles (fig. S1A). This experiment allowed us to correct for the geometry of the dish by
computing values relative to control conditions (**Fig. 1F**).

Across our four tested stimuli, we observed that larvae spent more time in brighter areas, whereas
juveniles were more likely to be found in darker regions (**Fig. 1, E** and **F**). We further computed
occupancies along stimulus-specific axes, allowing for individual fish analyses (**Fig. 1, G** and **H**),
confirming the switch in phototactic behavior from larval to juvenile stage. Wall interactions added
some attraction toward the rim (fig. S1B), slightly confounding results for juveniles for the two tested
radial gradients. We did not observe body size-dependent effects within each age group, supporting
the idea that the ontogeny of the behavior depends on neural rather than bodily development (fig.
S1C).

Thus, we find a robust behavioral switch in phototaxis behavior across a range of visual stimuli as
animals mature from the larval to the juvenile stage. These results hint towards specific adjustments
in the behavioral algorithms that animals employ during development. A possible strategy may be
based on swimming modulation as a function of whole-field luminance. To probe this idea, we next

developed a stimulus that allowed us to measure behavioral responses to varying but spatially uniform
brightness levels.
129

Fig. 1. Zebrafish invert phototactic behavior while growing from larval to juvenile stage. (**A**) Experimental
setup: The position and orientation of freely swimming fish are tracked in real-time using infrared illumination
and a camera. Brightness stimuli are projected from below onto shallow watch glasses (12 cm diameter) using
a video projector. (**B**) Our conceptual framework for model-based dissection of ontogenic switching of visual
behaviors. (**C**) Four static test stimuli: half-circles (10 vs. 300 lux), circular gradient (300 to 10 lux, and back),
inward gradient (10 to 300 lux), and outward gradient (10 to 600 lux). (**D**) Example trajectories for a single larva
and juvenile (same two animals across stimuli). (**E**) Two-dimensional distribution of the fraction of time spent
(%) across arena locations (calculated per fish, then averaged). (**F**) Same as in (E) but relative to the control
stimulus (uniform 300 lux across arena, fig. S1A). (**G**) Spatial distributions as fraction of time (%) spent at arena
locations, binned using stimulus-specific metrics (see axis labels in C), relative to the control stimulus. Mean ±
SEM over fish. (**H**) Stimulus-specific metrics for individual fish (circles). For the circular gradient, angular
positions are cosine-weighted. N = 70 larvae (blue) and 73 juveniles (orange). Each fish was presented with all
stimuli. Black circles in (H) indicate control stimulus (larvae, left; juveniles, right). Annotations indicate statistical
significance (p*** < 0.001; p** < 0.01; p* < 0.05; n.s. >= 0.05; Mann-Whitney U-test comparing test stimuli for
larvae vs. juveniles as well as the test stimuli vs. control within each age group).

Modulation of swimming behavior by whole-field brightness

148 Phototactic behavior may be controlled by whole-field luminance levels, temporal luminance changes,
149 spatial cues across the eyes, or combinations of these features. The presented static stimuli (**Fig. 1C**)
150 revealed a fundamental switch in behavior across development. However, such experiments do not
151 allow disentangling which of these components are most relevant for the observed effects because
152 spatial and temporal cues vary simultaneously as the fish swims through the arena. To address this

problem, we developed a set of well-controlled stimulus conditions to precisely probe temporal and spatial features independently from one another.

We first sought to explore the influence of whole-field luminance levels on behavior. To this end, we developed a virtual phototaxis paradigm with spatially uniform brightness. We picked the brightness level in real-time based on the position of the fish, using the circular gradient stimulus as a lookup function (**Fig. 2A**). This stimulus mimics the overall time scale of fluctuations when animals move through the arena, while not displaying any spatial contrasts. If animals rely only on spatial luminance cues to guide sensory-motor decision-making, then phototactic behavior should disappear in this configuration, because animals never experience any luminance contrast across the two eyes. On the contrary, if they use an averaging strategy of luminance across the visual field, their phototactic behavior should remain intact. Using the same stimulus-specific metrics as before (**Fig. 1G**), we find that larval ability to stay in brighter areas drops to almost chance levels, while juvenile performance improves (**Fig. 2B**).

To further explore the influence of whole-field luminance on the statistics of swimming, we analyzed behavior from the perspective of individual swims. Both larvae and juveniles exhibit burst-and-glide swimming, allowing us to characterize each event by orientation change, inter-swim interval, and displacement (**Fig. 2C**). Larvae and juveniles display symmetric orientation change distributions (**Fig. 2D**), as expected for uniform environments. Whereas larvae follow a trimodal distribution, with an emphasis on forward swims, juveniles predominantly exhibit turns, resulting in a bimodal distribution. These distributions were largely independent of radial position in the arena, and thus their shapes cannot be explained by wall-following behavior (fig. S2, A and B). We classified each swim as a turn when the absolute orientation change exceeded 10 degrees, a threshold corresponding to two standard deviations of the normal peak for larval orientation changes. Otherwise, swims were labeled as straight swims. We used these classifications to compute the percentage of turns relative to all swims and the median absolute angle of all turns.

The extracted swimming metrics allowed us to quantify the effect of uniform brightness levels on behavior (**Fig. 2E**) and develop a computational model (**Fig. 2F**). This model uses an Averaging pathway (A) in which the luminance average across the eyes is transformed through a logarithmic nonlinearity into behavioral properties. Fitting this model to our experiments allowed us to quantify brightness-dependent modulation across individual fish (**Fig. 2G**). For juveniles, we observed that with brightness, the absolute angle of turns decreases, the inter-swim interval increases, and the displacement per swim increases. Overall, these modulations should result in straighter trajectories using fewer but stronger swims to leave brighter areas more efficiently. In contrast, larvae did not significantly adjust any of the measured swim properties based on overall brightness. These results are in line with our observations of animal performance in the virtual circular gradient stimulus, where larvae failed to follow the luminance gradient but juveniles performed well (**Fig. 2B**). Thus, our results suggest that larvae do not navigate brightness environments based on a luminance averaging strategy, while juveniles do.

Notably, we observe an enhancement of phototaxis for juveniles in the virtual setting compared to the real static stimulus (**Figs. 2B** and **1G**), which hints towards a specific field-of-view for the luminance averaging process. To illustrate this logic, one may assume an extreme case in which juveniles average over large parts of the arena. In such a case, one would obtain similar brightness perceptions independent of their position. If this were true, we would expect uniform distributions of juveniles across the arena for the static stimuli. However, this is not the case (**Fig. 1G**). In another extreme, animals may sample only directly under them. In this case, the perceived brightness levels in the virtual and real configurations would be identical, leading to the same performance in the two cases. However, juveniles perform better in the virtual configuration (**Fig. 2B**). Therefore, a highly localized averaging strategy is not present either. We conclude that juveniles sample luminance within

an intermediate range around their body. We experimentally quantify the spatial extent of this visual
range below as part of our agent model design (**Fig. 4** and fig. S4).

Fig. 2. Whole-field brightness modulates swimming statistics in juveniles, but not in larvae. (**A**) Fish are
shown spatially uniform brightness levels that are dynamically updated in real-time based on the position of the
animal in a virtual circular gradient stimulus. (**B**) Same as in **Fig. 1G**, but for the virtual circular gradient stimulus.
Compared to the real circular gradient stimulus (thin lines), luminance navigation performance is largely
impaired in larvae, while it improves for juveniles. Mean ± SEM over fish. (**C**) Swims are detected based on
sudden changes in heading orientation and quantified by three metrics (orientation change, inter-swim interval,
and displacement). (**D**) Normalized distribution of orientation change for larvae and juveniles, and definitions of
turns and straight swims. The black arrowhead indicates the absolute angle of turns. Dashed black lines are
multimodal distribution fits (Methods). (**E**) Swimming behavior as a function of uniform brightness: Percentage
of turns relative to all swims, absolute angle of turns, inter-swim interval, and displacement. Mean ± SEM over
fish. Thin colored lines are medians within individual fish, and the black-dashed line is the average over
logarithmic fits of individual fish. (**F**) Averaging pathway (A) model (one separate model for each swim metric).
(**G**) Fitted logarithmic slopes for individual fish (circles) of the same data as in E. The black cross is the mean
over individual slopes. N = 46 larvae (blue) and 30 juveniles (orange). Annotations indicate bootstrapped
significance from zero (p*** < 0.005; p* < 0.05; n.s. >= 0.05; Methods).

As animals tend to repeat the same turning direction with consecutive swims (*43*), we argued that
motor history may perhaps also influence phototactic behavior (*37*). We found such effects to be weak
in our assays for both larvae and juveniles (fig. S2, C and D), indicating that motor history-dependent
dynamics are likely not the main drivers of phototaxis.

After having explored the influence of whole-field luminance levels on behavior, we next asked to
what degree whole-field luminance temporal changes may modulate swimming. To this end, we
tested a stimulus in which the complete arena linearly darkened or brightened at different speeds (fig.
S3A). We chose a broad range of speeds to also cover extreme scenarios. For both larvae and
juveniles, we find modulations of swimming behavior only for large brightness changes (fig. S3B). For
example, larvae increased their turning angle for strong brightness decrements, reminiscent of a dark-
flash response (*44*). Juveniles did the same, but for strong brightness increments. Yet, within the
intermediate luminance change regimes, modulation of swimming was rather small. We therefore
argue that luminance changes are likely less relevant to guide phototaxis under realistic conditions.

Based on the observed behavior, we developed a model using an Averaging-Derivative pathway (AD)
(fig. S3C), in which swim statistics are determined by changes in whole-field brightness levels with
different linear transformations for luminance decreases and increases. We fit this model to our
experimentally observed swim modulations (fig. S3B) to obtain negative and positive slopes for
individual fish (fig. S3, D and E). Using agent simulations, we will later further quantify the performance
of this model (**Fig. 4** and fig. S5G).

In summary, our experimental dissections of the influence of whole-field luminance dynamics on
behavior suggest that juveniles navigate brightness gradients using an Averaging strategy based on
instantaneously measured luminance levels within an intermediate spatial range. Larval behavior,
however, does not seem to be modulated by such cues, suggesting that phototaxis at that stage of
development likely involves other pathways, such as spatial computations across the two eyes. We
next developed a range of well-controlled spatio-temporal luminance configurations to explore this
possibility.

Modulation of turns by lateral brightness stimuli
We have shown that larval phototactic behavior is largely absent in our virtual critical gradient stimulus
(**Fig. 2B**) and that behavior is not modulated by whole-field brightness (**Fig. 2, E** and **G**). We thus
hypothesized that larvae use spatial contrast cues across the eyes to perform phototaxis. For
juveniles, we found that they can perform phototaxis using only whole-field luminance cues (**Fig. 2B**).
The average brightness modulates their swimming statistics (**Fig. 2, E** and **G**), which may be the basis
for the observed behavior. However, this result does not rule out that juveniles also employ spatial
information in more complex environments. To explore how these spatial aspects influence directed
turns, we implemented a range of visual stimuli in which luminance cues on either lateral side of the
fish could be independently controlled. To achieve this, while allowing animals to freely move through
the arena, we operated our tracking setup in a real-time closed-loop configuration, such that stimuli
are continuously locked to the position and orientation of the fish (**Fig. 3A**).

Specifically, we designed a lateral brightness stimulus consisting of periods of homogeneous
brightness, homogeneous darkness, and eye-specific luminance contrast cues locked to the midline
axis of the fish (**Fig. 3, A** and **B**). We characterized the temporal dynamics of the probability that
animals turn left, revealing three major components in the behavioral responses: First, fish tended to
transiently turn away from the side that underwent decreasing or increasing changes in brightness
(see **Fig. 3B** at 30 s and 150 s). Notably, this behavior also occurred when transitioning to a uniform
stimulus, even though spatial contrasts are absent during this period (see 90 s and 210 s). These
transient responses were present in both larvae and juveniles. Second, larval turning tendencies
converged to swimming towards the brighter side when spatial contrast is present (see 50 s, 80 s,
170 s, and 200 s). For juveniles, we found this contrast-dependent behavior to be much weaker and
close to chance levels. Third, when luminance contrast abruptly switched sides (see 60 s and 180 s),
we found larvae to show a rapid transition between contrast-dependent turning directions while
transient turning tendencies were absent. Juveniles, however, transiently swam towards the side that
decreased in brightness.

To quantify latent algorithmic parameters and compare values across larvae and juveniles, we
developed a unified model architecture to be fitted to each of the two age groups (**Fig. 3C**, Methods).
This model is a superposition of two parallel pathways: an eye-specific Derivative pathway (D) and a
spatial Contrast pathway (C). This model takes two inputs over time: the brightness on the left and
the right side of the fish. For each side, we used a high-pass filter acting like a derivative to detect
temporal changes in brightness. We implemented the high-pass filter as a mathematical operation
that compares an input signal with its low-pass filtered version (**Fig. 3C**, right). We then applied

different weight factors, depending on whether brightness increased or decreased. This resulted in a pathway in which luminance changes on the right side induced swims to the left, and the other way around for luminance changes on the left side. The different weight factors can lead to phototactic behavior, favoring increases over decreases or the other way around. We further computed a signed spatial contrast across the eyes by subtracting the brightness on the right side from the brightness on the left side and applied an additional weight factor to the resulting value. Combining these pathways results in a bias in turning direction. The model has only four parameters: the time constant (τ) and two asymmetric weights for the Derivative pathway, as well as the weight of the Contrast pathway.

We then fitted this model to our experimental data, allowing us to extract the latent cognitive variables underlying behavior (**Fig. 3D**, Methods). Comparison of the resulting values across the larval and juvenile stages then hints toward possible mechanisms of the computational adjustments during development. In the eye-specific Derivative pathway, the time constant was more than twice as high for larvae as for juveniles, suggesting faster temporal processing of visual information in older animals. For both larvae and juveniles, we found the two weight factors for brightness increments and decrements to be of opposing signs. This means that this pathway generally biases turns to go away from localized absolute brightness change for the two age groups. Interestingly, for juveniles, the magnitude of these weight factors was much lower for decrements than for increments, pointing towards asymmetric contrast change computations favoring darkness at this developmental stage (see **Fig. 3B**, 60 s and 180 s). For the Contrast pathway, larvae showed a strong positive weight, while the weight for juveniles was almost zero. This result highlights that spatial luminance cues contribute differently in larval and juvenile brightness navigation.

Our model design and its parameters are based on a stimulus that changes on fixed 30-second intervals (**Fig. 3B**). We next sought to challenge our model and test its predictive power with new, untested stimuli. To this end, we developed a paradigm in which uniform brightness levels and spatial contrast cues are temporally separated from one another with varying delay intervals (**Fig. 3E**). Specifically, we started with a dark uniform environment and then changed the luminance to bright for variable interval lengths. We then switched on the luminance contrast cue, with the bright stimulus on the right side, and probed the initial response during this period. For short intervals, our model predicted that during the luminance contrast phase, turns should be more likely to go towards the dark side, because the model is still adapted to the initial period of darkness. For longer intervals, the model should get increasingly adapted to the bright environment and thus predict more turns towards the bright side. To test these predictions, we performed experiments with fish in the same stimulus configuration. Animal behavior closely matched our modeling results for both larvae and juveniles. As an additional validation step, we repeated analyses on a luminance-inverted variant of the stimulus, where model and experiment were also in close agreement.

We conclude that two more luminance processing pathways are present in zebrafish, in addition to the Averaging pathway discussed in the previous section (**Fig. 2**). The Derivative pathway lets larval and juvenile zebrafish transiently swim away from changing brightness levels. The Contrast pathway controls responses to brightness differences across the eyes. Here, when presented with spatial luminance contrasts, larvae swim towards the brighter side, while juveniles remain largely undirected. We have now successfully described behavior in response to well-controlled whole-field (**Fig. 2)** and eye-specific spatial-temporal (**Fig. 3)** brightness environments. These results enable us to explore whether our fitted models, without further changing their structure and parameters, may predict phototactic behavior when fish dynamically interact with more complex brightness patterns.

Fig. 3. Lateral brightness stimuli modulate turn direction. (**A**) Visual stimuli are updated based on fish position in real-time, allowing independent control of brightness on the left and right sides of the fish (closed-loop configuration). (**B**) Left turns (%) over time in response to changing lateral brightness levels as depicted in the stimulus bar (top). Black dashed line: mean over individual model fits (see C, D). (**C**) Spatial-temporal model: A bias for left turns is determined by a Derivative (D) and Contrast (C) pathway. Increasing and decreasing brightness levels are detected with a high-pass filter. (**D**) Fitted model parameters for individual bootstrapped sets of fish (Methods). The black cross shows the mean over all fits. (**E**) Model prediction and experimental validation. Fish are adapted to one uniform brightness, then exposed to a new brightness for a variable interval (left: first 10 lux then 300 lux; right: first 300 lux then 10 lux), and subsequently presented with a dark-bright choice. Black dashed lines indicate model predictions. Quantification is based on left turn peak heights after the transition to the spatial contrast. Experimental data show mean ± SEM over fish. B, D: N = 45 larvae (blue) and 52 juveniles (orange), bootstrapped only for D. E: N = 63 larvae and 49 juveniles. Annotations indicate statistical significance (p*** < 0.001; p** < 0.01; p* < 0.05; Mann-Whitney U-test comparing larvae vs juveniles).

Model simulations of phototactic behavior in complex brightness environments

We have developed algorithmic models for larval and juvenile luminance-based behavioral modulations, whose parameters we fitted to simplified experimental configurations (**Figs. 2** and **3**). With these models, we can now test more complex stimuli where spatial and temporal stimulus features are intertwined and where the luminance cues perceived by the animal dynamically depend on its behavior. Moreover, such model simulations allow us to evaluate the relevance of each of the identified pathways for phototactic behavior and how pathway contributions may change across development.

To test our models, we performed agent simulations (**Fig. 4A**), in which virtual fish navigate stimulus landscapes identical to those presented to real animals (**Fig. 1C**). We projected the visual scene onto both eyes of the agent based on its position and orientation in the arena. To identify the spatial extent of these projections, we performed experiments in which we varied luminance cues as a function of distance to the animal (fig. S4). We found that swim modulations saturated beyond 2 cm and therefore set the field of view of our agent to a radius of 2 cm. To obtain eye-specific luminance levels, we averaged retinal images. We fed these time-evolving signals into our models to generate agent movement decisions. As the agents move through the arena, they perceive updated visual

inputs. We ran simulations for the same amount of time as for our real fish and generated the same
data structure, largely facilitating comparisons between real and virtual animals.

We set the parameters of our agent models solely based on the fitted model results from our
separate experiments (**Figs. 2** and **3**). This approach allowed us to validate models without fitting
them to test stimuli. To compare across different models (**Fig. 4B**), we computed a performance score
based on the mean squared error between the model and the experimental spatial distributions across
the four test stimuli, normalized by the error of a 'blind' control model that is fed with constant
luminance cues (Methods). A positive performance score indicates that the model simulations show
the same qualitative phototaxis tendency as real animals, while a negative value points toward
opposing behaviors between the model and experiment.

We first tested the Averaging pathway model (**Fig. 2F**). Juveniles show modulations of swimming
as a function of whole-field brightness, such as an increased displacement in brighter environments
(**Fig. 2, E** and **G**). Such a modulation should allow animals to more rapidly leave brighter areas. As
larvae did not show such modulations, we expect our larval agent model not to be able to perform the
behavior. Our simulations were in agreement with such reasoning (**Fig. 4B** and fig. S5A).

Next, we tested models composed of the Contrast and Derivative pathways (**Fig. 3C**). We first
probed each pathway in isolation. The Contrast pathway induces a tendency to swim towards the
brighter side for both age groups, with strong effects in larvae but weak effects in juveniles (**Fig. 3D**).
We expected that such dynamics would be reflected in our agent simulations, which was indeed the
case (fig. S5B). While our predictions were well aligned with the behavior of real larvae, they were the
opposite of what real juveniles do, thus leading to positive and negative model performance scores
for larvae and juveniles, respectively. For the Derivative pathway model, our agent simulation
qualitatively captured both larval and juvenile behavior (fig. S5C), resulting in positive performance
scores for both developmental stages. The combination of the Derivative and Contrast pathways
(D+C, same model as in **Fig. 3C**) further improves the spatial distributions across the four test stimuli
for larvae but not for juveniles (fig. S5D), reflected by the corresponding positive performance scores.

While these separate pathways in isolation can qualitatively explain individual elements of
phototaxis, we require their combination to fully capture the different configurations presented in **Figs.**
1 to **3**. We therefore propose a model that combines three parallel processing pathways (**Fig. 4C**):
The Averaging pathway modulates the probability that a swim is a turn, the absolute turn angle, the
inter-swim interval, and the displacement per swim. The Derivative and the Contrast pathways then
regulate the direction of turns. Not only is this proposed model able to capture the behavioral
responses discussed in the previous sections (**Figs. 1** to **3**), but it also shows the highest performance
score when comparing simulations with the experiment (**Fig. 4B**).

To facilitate easy comparison between our proposed model results and experimental data, we
plotted simulated data in the same style as displayed before: Our virtual agent simulations mimic the
trajectories of real fish (compare **Figs. 4D** and **1D**). Looking at the two-dimensional spatial
distributions, we observed that simulated larvae spend more time in bright areas, while juveniles go
to darker areas (compare **Figs. 4, E** and **F**, and **1**, **E** and **F**). Moreover, our agents largely capture the
same stimulus-specific distributions (compare **Figs. 4G** and **1G**). We attribute differences in the radial
configurations to wall interactions in the experimental data, which we did not explicitly include in our
models. Quantifying stimulus-specific metrics for individual fish showed significant differences
between larval and juvenile agents (**Fig. 4H**), as found in the experiments (**Fig. 1H**).

Notably, we obtained these results without tuning any of the model parameters. For juveniles,
simulation results and experimental data were in close agreement (fig. S5E). In larvae, however, we
only obtained qualitative matching of our stimulus-specific metrics (fig. S5E). Together, these results
indicate that our proposed model likely represents major important pathways in visual processing. To

show that our larval simulations would be capable of matching data quantitatively, we next manually tuned one of the parameters in our proposed model. Specifically, we increased the weight of the Contrast pathway in the larval model by a factor of 5. A model with such a configuration mimicked the experimentally observed distributions well (fig. S5F).

Fig. 4. Predicting brightness preferences in complex brightness environments. (**A**) Agents are simulated in virtual arenas of the same dimensions and seeing identical visual stimuli as in **Fig. 1C**. (**B**) Performance scores comparing agent spatial distributions to real larval and juvenile data, normalized to blind control agent. (**C**) Proposed model architecture: percentage of turns relative to all swims, absolute turn angle, inter-swim interval, and displacement are determined by the Averaging pathway (A, **Fig. 2F**), with turn direction controlled by the Derivative (D) and Contrast (C) pathways (**Fig. 3C**). (**D, E, F, G, H**) Same as in **Fig. 1** but for agent data. N = 96 larva-based agents (blue) and 96 juvenile-based agents (red). Annotations indicate statistical significance (p*** < 0.001; p** < 0.01; p* < 0.05; n.s. >= 0.05; Mann-Whitney U-test comparing test stimuli for larvae vs. juveniles as well as the test stimuli vs. control within each age group).

Finally, we wanted to know if more complex models, capturing additional behavioral features, would perform better than the proposed model. Further analyzing the behavioral response dynamics to the lateral brightness stimulus (**Fig. 3B**), we found that the other swim statistics, besides the fraction of left turns, were also affected (fig. S6A). We therefore designed a five-pathway model that independently modulates each of the five swim statistics (fig. S6B): In addition to the Averaging (A), Derivative (D), and Contrast (C) pathways, this extended model also uses an Averaging-Derivative (AD) (fig. S3C) and a Derivative-Averaging (DA) pathway. In the latter, we use the eye-specific Derivative pathways and then average signals across eyes. We simultaneously fit our model to the experimental data obtained from both the lateral brightness stimulus (**Fig. 3B**) as well as the virtual

circular gradient stimulus (**Fig. 2**), obtaining good matches between model and experiment (fig. S6,
A and C, Methods). We then probed the performance of this extended model in our agent simulations
(fig. S5G). Although this five-pathway model shows an improvement in the performance score
compared to the proposed model, it does not capture the qualitative aspects of the spatial densities
better than the proposed model. For completeness, we also tested two models composed only of the
Averaging-Derivative pathway (fig. S3C) or the Derivative-Averaging pathway, which did not perform
better than our proposed model (fig. S5, H and I).

In summary, through systematic testing of a variety of different models for phototaxis, we propose
a three-pathway configuration regulating swimming behavior. The Averaging pathway controls
undirected swimming statistics, while a superposition of a Derivative and a Contrast pathway
modulates turn direction tendencies. Even though our proposed model is relatively simple in structure
and parameter numbers, it could qualitatively capture behavior across a variety of experimental
configurations without parameter tuning. Manual tuning of a single model parameter allowed us to
obtain quantitatively matching results. The five-pathway model that modulates each swim statistic
separately also performed well, however, with the cost of a largely extended parameter space.
Comparing performance score and number of parameters across all tested model arrangements (fig.
S7A), we conclude that our proposed model provides an adequate intermediate level of model
complexity and biological realism.

Discussion

Several studies have explored larval zebrafish phototaxis across a range of stimulus configurations,
e.g. (*31–34*, *37*, *40*). Yet, there have been no attempts to obtain a comprehensive generalized model
that can explain behavior across conditions and over development. Here, we developed a systematic
strategy to unravel the algorithmic principles of zebrafish phototaxis, proposing that animals employ
three parallel pathways for luminance-based navigation. Our modelling approach allowed us to
dissect how computations change as animals become older. It has been reported that juvenile
zebrafish tend to swim into darkness, similarly to *Danionella cerebrum* adults (*30*). This behavior of
juvenile zebrafish is in stark contrast to what larval zebrafish do, which swim into the light, making this
transition a suitable example of vertebrate ontogeny.

Although little is known about zebrafish behavior in their natural environment (*45*), we hypothesize
that eggs are laid in shallow, brighter waters, where the semi-transparent larvae hatch and spend
their first days, likely staying hidden from predators or strong currents. As the fish grow and become
more visible, they may move to deeper or darker waters for less visual exposure or to start hunting.
Another example is the shell-dwelling cyclid, in which the leaving of the nest coincides with an
inversion from dark-seeking to light-seeking (*46*).

We found that juvenile zebrafish, like larvae, swim in a burst-and-glide pattern. Noteworthy, the
two age groups differed in their distribution of turn angles: whereas about half of larval swims are
directed straight, juveniles turned more frequently. This increase in turn frequency may reflect a shift
in their navigational strategy. With laterally positioned eyes, stronger relative sideways motion could
support a better scan of the environment ahead.

While larval swim characteristics showed little sensitivity to uniform brightness levels, juvenile
responses to whole-field brightness followed a natural logarithmic relationship, consistent with the
Weber–Fechner law (*47*). This developmental change may relate to earlier findings that retinal cells
in zebrafish keep developing until the juvenile stage (*16*). We conclude that whole-field brightness-
based behavioral modulations, which can be described via an Averaging pathway, play a major role
in the navigational strategy of juvenile zebrafish.

Using closed-loop stimulus delivery methods, previous studies have shown that larvae tend to
swim toward brighter regions (*33*, *34*) and away from sides where luminance changes occur (*37*, *40*).
We developed this paradigm further to capture both of these behavioral responses within a single
experiment. We observed that larval zebrafish behavior is predominantly guided by a Contrast
pathway that steers swimming toward the brighter side. Juveniles, however, show very little directional
swimming as a function of spatial contrast. Moreover, we find that animals at both developmental
stages show a strong tendency to swim away from luminance change, which we describe through a
Derivative pathway. This Derivative pathway may be linked to retinal ON- and OFF-pathways, which
have been attributed to guide zebrafish larvae along luminance gradients (*39*). Yet, as we find that
our Derivative pathway induces turns in response to both increments and decrements, it is unlikely to
be the only pathway involved in luminance gradient navigation. Instead, it may act as a conservative
mechanism to stabilize movement, which becomes particularly notable in high-contrast regions, such
as near the midline edge of the half-circle stimulus.

Based on our experiments, we propose a three-pathway model for zebrafish phototaxis. Our
model includes an Averaging pathway to control the percentage of turns relative to all swims, the
absolute turn angle, the inter-swim interval, and the displacement per swim. Derivative and Contrast
pathways then bias the turning direction.

Agent-based simulations are powerful tools for developing and validating computational models
of behavior and for probing their generalizability (*48*). We implemented our proposed model as an
agent navigating the same environments as our real fish. Importantly, even though this model was
not fitted to the test data, we found that model predictions and experimental results qualitatively
match. Comparing this model to a library of alternative model implementations, we further confirm
that our proposed model achieves the best predictive performance using only a moderate number of
parameters.

Our agent model describes both larval and juvenile behavior using the same algorithmic
architecture. We therefore hypothesize that the underlying neuronal connectivity and circuit
mechanisms underlying phototaxis are largely conserved during development. The observed switch
from light-seeking to dark-seeking may then be implemented through adjustments in synaptic
strengths and potentially rebalancing of pathways. This configuration allows behavior to flexibly adapt
to context, internal state, and changing environmental goals, without the need for an energy-costly
restructuring of the brain.

Despite these advances, our study has limitations. In **Fig. 3**, we applied bootstrapping over
individuals, due to the limited number of swim events per fish per time bin, which prevented fitting
models to individual fish. Shorter stimulation periods and longer experimental runs may help to
mitigate this problem in future experimental designs. Moreover, after manually adjusting the weight of
the Contrast pathway, our proposed model could not only qualitatively, but also quantitatively,
reproduce larval spatial distributions. Future experiments with a broader range of contrasts may
contribute to retrieving a more optimal value of the Contrast pathway weight. Experiments shown in
Figs. 1 to **3** were not conducted on the same batches of fish. One could also test the same individuals
across all conditions, allowing for single-fish comparisons and individualized model fitting.
Furthermore, our tested stimuli are only abstractions of naturalistic visual environments, which may
include dynamic motion, depth, and color variation (*49*, *50*). Future work could test whether our
findings hold under more ecologically realistic conditions, potentially even in the wild. Our study further
assumes that each fish behaves consistently throughout each experiment, which may not be true
(*51*). Exploring intra-individual variability over time could provide further insights into behavioral
flexibility and adaptation.

Further studies could explore behavior along more developmental time points to better understand if
transitions happen gradually or at specific maturation events, for example, with the onset of active

feeding. Another promising direction is to study phototactic behavior in animal groups, where social
cues may benefit collective decision-making (*52*). Here, it will be interesting to combine our
experimentally constrained model of phototaxis in juveniles with additional inter-animal attraction rules
and probe joint navigational performance. Our modeling framework can be used to make testable
predictions for experiments dissecting the structure and function of the nervous system within and
across species. For example, it will be interesting to repeat our behavioral analyses in *Danionella*
cerebrum, as it has been shown that these fish swim towards the dark already at the larval stage (*53*).
We have previously used a modeling approach to link specific disease-related mutations to social
behavior (*21*). Similarly, using recently developed mutant zebrafish with defects in brightness and
contrast perception (*54*), it will now be possible to test how such manipulations impact our proposed
visual processing pathways.

Our work takes advantage of the fact that development and behavior are in an interdependent
relationship. We explore two different developmental stages in zebrafish to constrain a universal
three-pathway model for phototaxis. This approach allows us to study how decision-making
algorithms unfold with development, in organisms that must solve context-specific problems at an
earlier stage, while still supporting the development of their behavioral repertoire. In general, our
findings provide important new insights into the computational mechanisms that guide vertebrate
ontogeny and exemplify the power of model-driven behavioral research.

Materials and Methods

Zebrafish handling and rearing

Wild-type *Danio rerio* (Konstanz strain) larvae, at 5 days post-fertilization (dpf), and juveniles, at 26
and 27 dpf, were used. Larvae were reared in Petri dishes containing E3 solution under a 14/10 hr.
light/dark cycle at 28°C. Juveniles were housed in ZebTec tanks under a 12/12 hr light-dark cycle and
at 28°C. Juveniles were fed dry Zebrafeed by Sparos 3 times a day. In addition, they were fed live
high-hufa artemia, according to the following schedule: 5–10 dpf: None; 11–15 dpf: 4x a day 1 drop
per feeding; 16–20 dpf: 4x a day 3 drops per feeding; 21–30 dpf: 4x a day 5 drops per feeding. This
protocol minimized variation in body size. Care was taken to always include fish from multiple
clutches. All experiments were conducted during daytime hours (9 AM to 5 PM). Experiments
involving juveniles were approved by the Regierungspräsidium Freiburg under permit number
G21/153. To reduce the number of animals used, only zebrafish that were surplus-bred for other
projects and could not be used in those projects were included in this study. Sex could not be
determined at this age.

Experimental setup

Experiments were conducted in diffusive watch-glass arenas, each filled with 75 mL of E3 solution,
resulting in a water volume having a diameter of 12 cm and a maximum depth of 5 mm. The water
temperature was stably maintained at 26°C. Before trials began, fish were given 15 minutes to
habituate to the experimental arenas. The visual stimuli were created using Panda3D. Visual stimuli
were projected from below using an AAXA P300 Pico Projector. We ensured control over the
projected brightness levels by measuring intensities for a fixed interval of projector input values using
an LX1330B light meter (Hongkong Thousandshores Ltd., Central Hong Kong). Subsequently, a
sigmoid function was fitted to the measured values for each projector and inverted to retrieve the
appropriate projector input value for each desired brightness level. Luminance levels varied between
10 lux (dark) and 300 lux (bright) for all stimuli except the outward gradient (**Figs. 1C** and **4C**), where
brightness increased to 600 lux. For the circular and radial gradients, the brightness scales linearly
between this brightness range along the azimuth and radius, respectively. The stimulus position was
calibrated to match the camera position, and a computational warping grid was applied to allow

orthogonal projection onto the curved watch-glass arenas. Stimuli were shown in random order, and
the same fish were exposed to stimulus-corresponding controls. Stimuli were shown either as static
or locked to the fish's position in a closed-loop setup configuration (full closed-loop delay: 60 ms). All
stimuli were also always shown in their mirrored version to control for potential asymmetries in
behavior and arena shape: around the arena's y-axis for static stimuli (**Figs. 1**, **2**, and **4**) or around
the fish's longitudinal body axis for fish-locked stimuli (**Fig. 3**).

Infrared illumination was provided by an LED array (940 nm wavelength), and behavioral recordings
were made using an IR-sensitive camera (Basler acA2040-90um-NIR) equipped with an adjustable
macro lens (Navitar ZOOM 7000) and an IR longpass filter (Linghuang Zomei IR 850 nm 52 mm).
The camera operated at 90 Hz. To increase experimental throughput, we used two computers, each
connected to eight cameras and four projectors, allowing for independent stimulation and closed-loop
tracking of 16 fish simultaneously. Fish position and orientation were continuously measured using
background subtraction-based tracking, implemented in custom Python software as described
previously (*55*). To compute body size (fig. S1C), we implemented a real-time posture-based tracking
to detect the midline. Experiments, simulations, and data analysis were performed using custom-
written Python code (Python 3.11).

Agent simulations

Each simulation ran at 60 Hz and included 96 model agents. The simulations were run in parallel at
the agent level, in which each agent was assigned a unique random number generator seed. The
simulated arena matched the real experimental arena (6 cm radius). Simulated agents received
identical visual stimuli as the real fish, with the same trial structure, trial duration, and number of trials.
Each agent had a left and right eye, each with a 2 cm half-disk field of view, where the half-circle
region on the side of each eye was considered for visual input. Pixels outside the dish but within the
field of view were assigned not-a-number values and ignored in further processing. The stimulus was
rendered on a grid of 256 by 256 pixels. At the start of each trial, each agent was assigned a random
orientation and position within the arena.

The swim events of agents were determined based on the brightness input to the corresponding
models. All models had parameters obtained from real fish data (**Figs. 2** and **3,** and fig. S3), fitted for
each age group and swim statistical property. The Averaging pathway (A) models (**Fig. 2D**) took the
average brightness (B) across the two eyes

$$B_{\mathrm{A}} = \frac{B_{\mathrm{right}} + B_{\mathrm{left}}}{2}$$

and the contribution to the swim characteristics by evaluating the natural logarithm

$y_{\mathrm{A}} = b_{\mathrm{A}} \log B_{\mathrm{A}}$.

The Contrast pathway models (**Fig. 3C**) compute the contrast across the eyes following

$B_{\mathrm{C}} = B_{\mathrm{right}} - B_{\mathrm{left}}$

and its contribution to swim characteristics was obtained by evaluating the linear function

$y_{\mathrm{C}} = b_{\mathrm{C}} B_{\mathrm{contrast}}$.

The Derivative (D, **Fig. 3C**), Averaging-Derivative (AD), and Derivative-Averaging pathway (DA)
involve a high-pass filter to detect luminance changes. This filter is implemented via a low-pass filter
that is slowly updated to match the new brightness value. The low-pass filtered value is updated at
each timestep n following

$B_{\mathrm{LPF},\, n} = \alpha * B_n + (1 - \alpha) * B_{\mathrm{LPF},\, n-1}$

where the low-pass filter parameter α is based on the time constant τ and time since last processed
frame Δt:

$\alpha = \Delta t / (\tau + \Delta t)$.

Subsequently, the high-pass filter value is obtained by computing the difference between the current
perceived brightness B_n and the low-pass filtered $B_{\mathrm{LPF},\,n}$ by

$$\Delta B = B_{\mathrm{LPF},\,n} - B_n$$.

The resulting brightness change value is subsequently multiplied by sign-dependent weights and
linearly transformed to obtain the swim characteristics according to

$$y = b \begin{cases} w^+ \Delta B & \text{if } \Delta B > 0 \\ w^- \Delta B & \text{if } \Delta B \leq 0 \\ 0 & \text{otherwise} \end{cases}$$

Where b, w^+, and w^- varied for the D, AD, and DA pathways. The contributions of each pathway are
summed, and an offset a is added. For the extended five-pathway model, this results in

$$Y = y_A + y_C + y_D + y_{AD} + y_{DA} + a.$$

In the proposed model, the percentage of turns relative to all swims, turn angle, inter-swim interval,
and displacement are determined by the Averaging pathway

$$Y = y_A + a$$

and the fraction of left turns by a superposition of the Derivative (D) and Contrast (C) pathways

$$Y = y_D + y_C + a.$$

Upon reaching the arena wall, agents maintained their general swimming direction but were assigned
a new heading angle relative to the wall, drawn from a normal distribution centered around 115
degrees and with a 10-degree standard deviation. The angle drawing process was repeated until
agents moved away from the wall again.

Data analysis

<u>Edge interactions, pre-screening, and data exclusion:</u> To exclude edge effects, only data from fish
located at least 1 cm away from the edge were included in the analysis. As part of the habituation to
the experimental arena, larvae underwent a 10-minute pre-screening period to exclude unhealthy
individuals. Larvae that did not adhere to the following behavioral criteria were excluded: Leftward
and rightward gratings: at least 50% of turns aligned with the grating direction; Homogeneous gray
background: maximum of 20% of turns in a single direction. For juveniles, the arena was left gray
during the habituation phase without performing a pre-screening protocol. To exclude potential
tracking errors during the pre-screening and the experiment, swim events were discarded if they
exceeded predefined thresholds. Generally, less than 1% of all data was discarded. Larvae: maximum
contour area = 12,000 px, maximum average speed per swim = 1.5 cm/s. Juveniles: max. contour
area = 16,000 px, maximum average speed per swim = 3 cm/s. All fish: maximum inter-swim interval
= 10 s, maximum absolute orientation change = 150°. A complete trial was discarded if more than 5%
of swim events in that trial exceeded any of the above thresholds.

<u>Swim event detection:</u> Fish position and orientation were extracted after background subtraction, with
the background being updated every 120 s. Swim events were identified based on windowed variance
thresholds applied to movement data, using a variance window of 0.050 s, a minimum duration above
the start threshold of 0.020 s, a minimum duration below the end threshold of 0.050 s, and a maximal
event length of 4 s. The start and end variance thresholds differed between larvae (start threshold =
5, end threshold = 1.5) and juveniles (start threshold = 7.5, end threshold = 5).

<u>Figures 1 and 4:</u> The tracking data described the x and y positions for every individual, stimulus, and
trial, with each row denoting a frame. Spatial binning was applied using 10 bins. Bins were defined
for x-position, radius, and azimuthal angle, with values ranging from –5 to 5 cm for x, 0 to 5 cm for
radius, and 0 to 360 degrees for the angle. The spatial distributions were first averaged within
individuals before calculating the mean and standard error of the mean over each population. The
average cosine-weighted angle was used to capture the average position of individuals for the circular

gradient test stimulus. This was computed by multiplying the number of swims within each azimuthal
bin $N(\alpha_i)$ with the cosine-weighted value $\cos(\alpha_i)$ for that bin and subsequently summing over all bins

658
$$100 \cdot \sum_{\alpha_i \in \text{angles}} N(\alpha_i)\cos(\alpha_i)$$
.

659 Simulations produced the same format as the experimental data, allowing us to perform the exact
660 same analysis for both figures. The agents' performance score indicates how well an agent captures
661 the spatial distribution of the real fish. We calculate the average mean-squared error for each of the
662 four test stimuli in their stimulus-specific spatial bins, comparing the spatial distributions of simulated
663 agents (p_{agent}) to real fish (p_{ref})

664
$$\text{MSE} = \frac{1}{N_{\text{stim}}} \sum_{i \in \text{stimuli}} \frac{1}{N_{\text{bins}}} \sum_{j \in \text{bins}} \sqrt{\left(p_{\text{ref},i,j} - p_{\text{agent},i,j}\right)^2}$$
.

665 Subsequently, the performance was calculated as the percentage improvement relative to a blind
666 control model (perceiving constant luminance levels),

667
$$\frac{\text{MSE}_{\text{blind}} - \text{MSE}}{\text{MSE}_{\text{blind}}} \cdot 100\%$$
.

<u>Figure 2:</u> The swim-event characteristics (orientation change, inter-swim interval, and displacement)
are summarized as median values over trials, within individual fish and stimuli, as the underlying
distributions are not necessarily symmetric. Orientation change can be best described by a mixture
of a Normal distribution for forward swims and two Maxwell-Boltzmann distributions for left and right
turns. The proportions of left and right turns can be captured using weighting factors in the mixture
model. Displacement and inter-swim intervals both follow a Gamma distribution, which is known to be
well-suited for representing temporal and spatial intervals.

In the virtual circular gradient paradigm, the arena brightness was set based on the azimuthal position
of the fish. Swims were grouped into brightness bins, based on the brightness at the start of the swim,
and medians were computed within each brightness bin for individual fish. A logarithmic function was
then fitted to the brightness data (see below), allowing for an estimation of the relationship between
brightness and swim characteristics. To assess whether the fitted slope parameters differed
significantly from zero, we performed bootstrapping over fish with 10,000 iterations to compute two-
tailed p-values and 95% confidence intervals. This approach provided robust statistical inference for
evaluating the modulation of swim parameters by brightness. The Averaging pathway (A) model (**Fig.
2D**) is described above in the subsection called 'Agent simulations'.

<u>Figure 3:</u> To ensure sufficient data points per temporal bin for model fitting, we generated
bootstrapped sets of fish. For each age group, the total number of bootstrapped fish matched the
number of original individuals. Each bootstrapped fish was constructed by randomly sampling, with
replacement, two-thirds of the available real fish. For time-series analysis, data were resampled by
calculating the median value within 1-second time windows for each individual. The Contrast (C) and
Derivative (D) pathways (**Fig. 3C**) are described above in the 'Agent simulations' subsection. Models
were fitted to individual bootstrapped fish (see below), and mean model parameters were retrieved
by averaging over parameters of the individual fits.

To detect peaks in the percentage of left turns (**Fig. 3E**), values before stimulus onset were excluded,
and the peak was determined by maximizing the absolute deviation from the 50% chance level.

<u>Statistical analysis and model fits:</u> All statistical tests are specified in the figure legends. To compare
distributions against zero, we used bootstrapping to compute a two-tailed p-value. Unless specified
otherwise, resampling was performed 10,000 times, each time selecting 2/3 of the individuals to
create a bootstrapped dataset matching the original sample size. For comparisons between
distributions, we applied the Mann-Whitney U-test.

Model parameters were obtained by fitting the analytic model equations to experimental data
using the optimization and root-finding functions of the Python package SciPy. Fitting to separate
experimental datasets–virtual circular gradient stimulus (**Fig. 2**), lateral brightness stimulus (**Fig. 3**),
and whole-field luminance changes stimulus (fig. S3)–was performed using the non-linear least
squares fit function *curve_fit*. The extended five-pathway model was fitted to combined experimental
data from the lateral brightness stimulus and virtual circular gradient stimulus by minimizing the mean-
squared error to both datasets using the *minimize* function.

Data and materials availability

The code for simulations and data analysis is available on GitHub ([github.com/bahl-lab-](github.com/bahl-lab-konstanz/phototaxis_ontogeny)
[konstanz/phototaxis_ontogeny](github.com/bahl-lab-konstanz/phototaxis_ontogeny)). Data will be available upon request. GitHub Copilot was used for low-
level coding suggestions.
710

References

1. M. Tomasello, *Becoming Human: A Theory of Ontogeny* (Harvard University Press, Cambridge, Massachusetts, 2019).

2. W. M. Fox, Reflex-ontogeny and behavioural development of the mouse. *Anim. Behav.* **13**, 234-IN5 (1965).

3. D. A. Lynn, G. R. Brown, The ontogeny of anxiety-like behavior in rats from adolescence to adulthood. *Dev. Psychobiol.* **52**, 731–739 (2010).

4. F. Donato, C. M. Alberini, D. Amso, G. Dragoi, A. Dranovsky, N. S. Newcombe, The ontogeny of hippocampus-dependent memories. *J. Neurosci.* **41**, 920–926 (2021).

5. J. Sztarker, F. C. Rind, A look into the cockpit of the developing locust: Looming detectors and predator avoidance. *Dev. Neurobiol.* **74**, 1078–1095 (2014).

6. E. P. Sawin-McCormack, M. B. Sokolowski, A. R. Campos, Characterization and genetic analysis of *Drosophila melanogaster* photobehavior during larval development. *J. Neurogenet.* **10**, 119–135 (1995).

7. M. L. Zhu, K. J. Herrera, K. Vogt, A. Bahl, Navigational strategies underlying temporal phototaxis in *Drosophila* larvae. *J. Exp. Biol.* **224**, jeb242428 (2021).

8. S. Benzer, Behavioral mutants of *Drosophila* isolated by countercurrent distribution. *Proc. Natl. Acad. Sci.* **58**, 1112–1119 (1967).

9. J. S. Kain, C. Stokes, B. L. de Bivort, Phototactic personality in fruit flies and its suppression by serotonin and *white*. *Proc. Natl. Acad. Sci.* **109**, 19834–19839 (2012).

10. H. Baier, E. K. Scott, The visual systems of zebrafish. *Annu. Rev. Neurosci.* **47**, 255–276 (2024).

11. K. Lee, C. Q. Doe, A locomotor neural circuit persists and functions similarly in larvae and adult *Drosophila*. *eLife* **10**, e69767 (2021).

12. J. Kohl, B. M. Babayan, N. D. Rubinstein, A. E. Autry, B. Marin-Rodriguez, V. Kapoor, K. Miyamishi, L. S. Zweifel, L. Luo, N. Uchida, C. Dulac, Functional circuit architecture underlying parental behaviour. *Nature* **556**, 326–331 (2018).

13. S. S. Easter, G. N. Nicola, The development of vision in the zebrafish (*Danio rerio*). *Dev. Biol.* **180**, 646–663 (1996).

14. J. C. Beck, E. Gilland, D. W. Tank, R. Baker, Quantifying the ontogeny of optokinetic and vestibuloocular behaviors in zebrafish, medaka, and goldfish. *J. Neurophysiol.* **92**, 3546–3561 (2004).

15. H. K. Koning, A. Ahemaiti, H. Boije, A deep-dive into fictive locomotion – a strategy to probe cellular activity during speed transitions in fictively swimming zebrafish larvae. *Biol. Open* **11**, bio059167 (2022).

16. T. Branchek, R. Bremiller, The development of photoreceptors in the zebrafish, *Brachydanio rerio*. I. Structure. *J. Comp. Neurol.* **224**, 107–115 (1984).

17. R. Portugues, F. Engert, The neural basis of visual behaviors in the larval zebrafish. *Curr. Opin. Neurobiol.* **19**, 644–647 (2009).

18. S. J. Stednitz, P. Washbourne, Rapid progressive social development of zebrafish. *Zebrafish* **17**, 11–17 (2020).

19. E. Dreosti, G. Lopes, A. R. Kampff, S. W. Wilson, Development of social behavior in young zebrafish. *Front. Neural Circuits* **9**, 39 (2015).

20. R. Harpaz, M. N. Nguyen, A. Bahl, F. Engert, Precise visuomotor transformations underlying
collective behavior in larval zebrafish. *Nat. Commun.* **12**, 6578 (2021).

21. R. Harpaz, A. C. Aspiras, S. Chambule, S. Tseng, M.-A. Bind, F. Engert, M. C. Fishman, A.
Bahl, Collective behavior emerges from genetically controlled simple behavioral motifs in
zebrafish. *Sci. Adv.* **7**, eabi7460 (2021).

22. J. Larsch, H. Baier, Biological motion as an innate perceptual mechanism driving social
affiliation. *Curr. Biol.* **28**, 3523-3532.e4 (2018).

23. R. C. Hinz, G. G. de Polavieja, Ontogeny of collective behavior reveals a simple attraction rule.
Proc. Natl. Acad. Sci. **114**, 2295–2300 (2017).

24. A. Valente, K.-H. Huang, R. Portugues, F. Engert, Ontogeny of classical and operant learning
behaviors in zebrafish. *Learn. Mem.* **19**, 170–177 (2012).

25. K. Yashina, Á. Tejero-Cantero, A. Herz, H. Baier, Zebrafish exploit visual cues and geometric
relationships to form a spatial memory. *iScience* **19**, 119–134 (2019).

26. F. Palumbo, B. Serneels, R. Pelgrims, E. Yaksi, The zebrafish dorsolateral habenula is
required for updating learned behaviors. *Cell Rep.* **32**, 108054 (2020).

27. J. Bak-Coleman, D. Smith, S. Coombs, Going with, then against the flow: evidence against the
optomotor hypothesis of fish rheotaxis. *Anim. Behav.* **107**, 7–17 (2015).

28. S. Coombs, J. Bak-Coleman, J. Montgomery, Rheotaxis revisited: a multi-behavioral and
multisensory perspective on how fish orient to flow. *J. Exp. Biol.* **223**, jeb223008 (2020).

29. E. L. Serra, C. C. Medalha, R. Mattioli, Natural preference of zebrafish (*Danio rerio*) for a dark
environment. *Braz. J. Med. Biol. Res.* **32**, 1551–1553 (1999).

30. T. J. Lee, K. L. Briggman, Visually guided and context-dependent spatial navigation in the
translucent fish *Danionella cerebrum*. *Curr. Biol.* **33**, 5467-5477.e4 (2023).

31. S. E. Brockerhoff, J. B. Hurley, U. Janssen-Bienhold, S. C. Neuhauss, W. Driever, J. E.
Dowling, A behavioral screen for isolating zebrafish mutants with visual system defects. *Proc.
Natl. Acad. Sci. U. S. A.* **92**, 10545–10549 (1995).

32. H. A. Burgess, M. Granato, Modulation of locomotor activity in larval zebrafish during light
adaptation. *J. Exp. Biol.* **210**, 2526–2539 (2007).

33. K.-H. Huang, M. B. Ahrens, T. W. Dunn, F. Engert, Spinal projection neurons control turning
behaviors in zebrafish. *Curr. Biol.* **23**, 1566–1573 (2013).

34. S. Karpenko, S. Wolf, J. Lafaye, G. Le Goc, T. Panier, V. Bormuth, R. Candelier, G.
Debrégeas, From behavior to circuit modeling of light-seeking navigation in zebrafish larvae.
eLife **9**, e52882 (2020).

35. M. B. Orger, H. Baier, Channeling of red and green cone inputs to the zebrafish optomotor
response. *Vis. Neurosci.* **22**, 275–281 (2005).

36. A. M. Fernandes, K. Fero, A. B. Arrenberg, S. A. Bergeron, W. Driever, H. A. Burgess, Deep
brain photoreceptors control light-seeking behavior in zebrafish larvae. *Curr. Biol.* **22**, 2042–
2047 (2012).

37. X. Chen, F. Engert, Navigational strategies underlying phototaxis in larval zebrafish. *Front.
Syst. Neurosci.* **8**, 39 (2014).

38. E. J. Horstick, Y. Bayleyen, J. L. Sinclair, H. A. Burgess, Search strategy is regulated by
somatostatin signaling and deep brain photoreceptors in zebrafish. *BMC Biol.* **15**, 4 (2017).

39. H. A. Burgess, H. Schoch, M. Granato, Distinct retinal pathways drive spatial orientation
behaviors in zebrafish navigation. *Curr. Biol.* **20**, 381–386 (2010).

40. A. B. Chen, D. Deb, A. Bahl, F. Engert, Algorithms underlying flexible phototaxis in larval
zebrafish. *J. Exp. Biol.* **224**, jeb238386 (2021).

41. D. Marr, *Vision: A Computational Investigation into the Human Representation and Processing
of Visual Information* (MIT Press, Cambridge, Mass, 1982).

42. V. Braitenberg, *Vehicles, Experiments in Synthetic Psychology* (MIT Press, Cambridge, Mass,
1984).

43. T. W. Dunn, Y. Mu, S. Narayan, O. Randlett, E. A. Naumann, C.-T. Yang, A. F. Schier, J.
Freeman, F. Engert, M. B. Ahrens, Brain-wide mapping of neural activity controlling zebrafish
exploratory locomotion. *eLife* **5**, e12741 (2016).

44. L. A. Lamiré, M. Haesemeyer, F. Engert, M. Granato, O. Randlett, Functional and
pharmacological analyses of visual habituation learning in larval zebrafish. *eLife* **12**, RP84926
(2023).

45. J. Sundin, R. Morgan, M. H. Finnøen, A. Dey, K. Sarkar, F. Jutfelt, On the observation of wild
zebrafish (*Danio rerio*) in India. *Zebrafish* **16**, 546–553 (2019).

46. A. V. Parker, M. Stemmer, S. Grätsch, A. Dorigo, O. Rodriguez Ramirez, A. Adel, A. Jordan,
H. Baier, Intrinsic timing of brood care in shell-dwelling cichlids. *Curr. Biol.* **35**, 672-680.e4
(2025).

47. S. Dehaene, The neural basis of the Weber-Fechner law: a logarithmic mental number line.
Trends Cogn Sci **7**, 145–147 (2003).

48. S. H. Singh, F. Van Breugel, R. P. N. Rao, B. W. Brunton, Emergent behaviour and neural
dynamics in artificial agents tracking odour plumes. *Nat. Mach. Intell.* **5**, 58–70 (2023).

49. L. T. Cai, V. S. Krishna, T. C. Hladnik, N. C. Guilbeault, C. Vijayakumar, M. Arunachalam, S.
A. Juntti, A. B. Arrenberg, T. R. Thiele, E. A. Cooper, Spatiotemporal visual statistics of aquatic
environments in the natural habitats of zebrafish. *Sci. Rep.* **13**, 12028 (2023).

50. T. Baden, Circuit mechanisms for colour vision in zebrafish. *Curr. Biol.* **31**, R807–R820 (2021).

51. K. Krishnan, A. Muthukumar, S. Sterrett, P. Pflitsch, A. Fairhall, M. Fishman, A. Bahl, H.
Zwaka, F. Engert, Attentional switching in larval zebrafish: The attentive leaky integrator.
Research Square [Preprint] (2023).

52. A. Berdahl, C. J. Torney, C. C. Ioannou, J. J. Faria, I. D. Couzin, Emergent sensing of complex
environments by mobile animal groups. *Science* **339**, 574–576 (2013).

53. N. Lindemann, L. Kalix, J. Possiel, R. Stasch, T. Kusian, R. W. Köster, J. W. von Trotha, A
comparative analysis of *Danionella cerebrum* and zebrafish (*Danio rerio*) larval locomotor
activity in a light-dark test. *Front. Behav. Neurosci.* **16** (2022).

54. E. Putti, G. Faini, J. T.-M. Dang, F. Eggeler, J. H. Savaliya, K. Duroure, J. Vougny, A. Bahl, J.
W. Lichtman, F. Engert, J. Boulanger-Weill, F. Del Bene, S. Albadri, Lrrns define a visual
circuit underlying brightness and contrast perception. bioRxiv [Preprint] (2025).

55. A. Bahl, F. Engert, Neural circuits for evidence accumulation and decision making in larval
zebrafish. *Nat. Neurosci.* **23**, 94–102 (2020).

836
837

Acknowledgments

We thank the staff of the Animal Facility of the Couzin Lab, and in particular Dominique Leo and Jayme Weglarski, for their generous supply of zebrafish eggs and juvenile fish. In addition, we thank Heike Naumann and Ulrike Bonitz for their vital work in keeping the lab running. We are thankful to the staff support of our central machine shop, especially Ralf Honz, for the invaluable assistance with building our tracking setups. We are grateful for the inspiring and stimulating discussions with Daan Brinks, Iain Couzin, and Osama M. Ahmed. Finally, we would like to thank Daan Brinks, Fabienne Roth, Lukas Pijnacker-Hordijk, Margherita Zaupa, Panagiotis Eleftheriadis, Osama M. Ahmed, and Sophie Aimon for proofreading and providing constructive feedback on the manuscript.

Funding

This work was funded by the Emmy Noether Program (BA 5923/1-1), an ERC Starting Grant (101075541 – CollectiveDecisions), and the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under Germany's Excellence Strategy (EXC 2117 – 422037984). In addition, the Zukunftskolleg Konstanz supported A.B. and M.Q.C. The Max Planck Institute for Animal Behavior provided bridge funding for M.Q.C. K.S. was supported by a Boehringer Ingelheim Fonds graduate fellowship.

Author contributions

Conceptualization: M.Q.C., K.S., and A.B.; Data curations: M.Q.C., K.S., and A.B.; Formal analysis: M.Q.C.; Funding acquisition: M.Q.C., A.B.; Investigations: M.Q.C.; Methodology: M.Q.C., K.S.; Project administration: M.Q.C.; Resources: M.Q.C., A.B.; Software: M.Q.C., K.S., A.B.; Supervision: A.B.; Validation: M.Q.C., K.S., A.B.; Visualization: M.Q.C.; Writing - original draft: M.Q.C., A.B.; Writing - review & editing: M.Q.C., K.S., A.B.

Competing interests

The authors declare no competing interests.

866 **Supplementary Materials**

867 Figs. S1 to S7

869
Fig. S1. Controls for static luminance experiments. (**A**) Larval and juvenile distributed homogeneously in a
control stimulus (uniform 300 lux across the arena). Fraction of time (%) spent at arena locations, binned using
the same metrics as in **Fig. 1G**, but not normalized to control stimulus. Mean ± SEM over fish. (**B**) Wall
interactions add a small additional attraction towards the rim for all tested stimuli. (**C**) The relation between body
size (as measured by the length of the midline) and phototactic behavior for individual fish. Body size within
each age group does not influence the position of the fish in the arena. N = 70 larvae (blue) and 73 juveniles
(orange), thin black lines indicate mathematical chance levels, and black dashed lines indicate linear fits to age
populations.

879

880 **Fig. S2. Directed turns in the virtual circular gradient stimulus.** (**A**) Area-normalized distributions of
881 orientation changes at different radial bins in the arena. (**B**) Fraction of left-directed turns as a function of whole-
882 field brightness. Annotations indicate bootstrapped significance from zero (n.s. >= 0.05). (**C**) Following the
883 analysis performed in (*43*), we only observe a weak, not significant effect in streak length probabilities when
884 compared to randomly generated turns. P-values indicate statistical significance (Mann-Whitney U-test
885 comparing randomly generated turns to real data). (**D**) Turn pair counting for individual fish does not show a
886 significant difference from shuffled data either. N = 46 larvae (blue) and 30 juveniles (orange). Black vertical
887 bars indicate 95% bootstrapped confidence intervals.

888

Fig. S3. Whole-field luminance changes. (**A**) We present whole-field luminance changes by linearly
increasing or decreasing the arena brightness with either 30, 60, 120, or 300 lux/s. (**B**) Swimming behavior as
a function of temporally changing whole-field brightness levels: Percentage of turns relative to all swims,
absolute angle of turns, inter-swim interval, and displacement. Mean ± SEM over fish. Thin colored lines are
median values within individual fish, and the black-dashed line is the average over fits of individual fish. (**C**)
Averaging-Derivative pathway (AD) model. (**D**, **E**) Fitted slopes for individual fish (circles) of the same data as
in B. N = 54 larvae (blue) and 43 juveniles (orange). The black cross is the mean over individual slopes.
Annotations indicate bootstrapped significance from zero (p*** < 0.005; p* < 0.05; n.s. >= 0.05; Methods).

899
Fig. S4. Field of view measurements reveal spatial extent of luminance calculations. To gain an
impression of the field of view of larval and juvenile zebrafish for luminance calculations, we quantified swim
characteristics for three different fish-locked stimuli. (**A**) A spatial-contrast stimulus where the lateral offset
position of the contrast edge is varied relative to the midline of the fish. (**B**) A dark disk centered at the position
of the fish in a bright environment. (**C**) A bright disk in a dark environment. For each configuration, we quantified
how behavior is modulated by the offset or radius of the stimulus. Mean ± SEM over individual fish, thin colored
lines are medians within individual fish. A: N = 47 larvae (blue) and 20 juveniles (orange), B and C: N = 10
larvae and 16 juveniles.
908

Fig. S5. Detailed simulation results for all agent model variants. (**A**–**I**) First four columns: Fraction of time (%) spent at arena locations, binned using stimulus-specific metrics (see axis labels in **Fig. 1C**), relative to control stimulus. Mean ± SEM over agents. Fifth column: model illustration. Sixth column: Performance score. A to I: Averaging pathway model (A), Contrast pathway model (C), Derivative pathway model (D), Derivative and Contrast pathway model (D+C), the proposed model (Proposed), the proposed model (Proposed*) with manually tuned Contrast weight (larval agent only), Extended model (Five-pathways), Averaging-Derivative pathway model (AD), and Derivative-Averaging pathway model (DA). N = 96 agents for each model simulation and age group, larval-agents (hollow light blue), juvenile-agents (hollow red), 70 real larvae (solid blue), and 72 real juveniles (solid orange).

920
Fig. S6. Extended five-pathway model for lateral brightness and virtual circular gradient stimuli. (**A**)
Swimming behavior in response to lateral brightness stimulus (**Fig. 3B**); percentage of turns relative to all swims
of all swims, fraction of left turns of all turns, absolute angle of turns, inter-swim interval, and displacement over
time. (**B**) Extended model including all pathways: Averaging (A), Averaging-Derivative (AD), Derivative (D),
Contrast (C), and Derivative-Averaging (DA) pathway. (**C**) Swimming behavior as a function of uniform
brightness level (**Fig. 2A**). A: N = 45 larvae (blue) and 52 juveniles (orange), C: N = 46 larvae and 30 juveniles.
Thin, solid, colored lines are individual fish. Black dashed line: mean over individual model fits.

Fig. S7. Simulated model performance across all tested agent configurations. (**A**) Mean performance
scores over larval and juvenile-based models vs model parameters. Blind: Constant luminance input, C:
Contrast pathway, D: Derivative pathway, D+C: Derivative and Contrast pathway, Proposed: proposed model,
Proposed*: proposed model with manually tuned Contrast weight (larval agent only), DA: Derivative-Averaging
pathway, AD: Averaging-Derivative pathway, Five-pathways: extended model.