Sequence-based prediction of condensate composition reveals that specificity can emerge from multivalent interactions among disordered regions

Jonas Wessén[,1,4], Nancy De La Cruz[2,4], Heankel Lyons[2], Hue Sun Chan[1,*], Benjamin R. Sabari[2,3,*]

[1] Department of Biochemistry, Temerty Faculty of Medicine, University of Toronto, Toronto, Ontario, Canada

[2] Laboratory of Nuclear Organization, Cecil H. and Ida Green Center for Reproductive Biology Sciences, University of Texas Southwestern Medical Center, Dallas, TX, USA

[3] Department of Molecular Biology, Hamon Center for Regenerative Science and Medicine, University of Texas Southwestern Medical Center, Dallas, TX, USA

[4] These authors contributed equally to this work

*Correspondence: H.S.C. huesun.chan@utoronto.ca; B.R.S. benjamin.sabari@utsouthwestern.edu

Abstract

While specificity of biomolecular interactions is typically understood to require interactions involving ordered structures, several biomolecular condensates exhibit specificity in the absence of apparent structural order. We have previously shown that condensates composed of the disordered region of MED1 partition specific proteins, mediated by sequence patterns of charged amino acids on the disordered regions of both MED1 and the interacting partner. Whether this specificity is due to an unknown ordered-structure-mediated interaction or from the dynamic multivalent interactions between the patterned charged amino acids in the disordered regions was unresolved. Here we show that a polymer physics-based model that only accounts for multivalent interactions among polymers in a statistical manner can largely explain published data on selective partitioning and make predictions that are subsequently experimentally validated. These results suggest that the specificity of condensate composition is underpinned to a substantial extent by multivalent interactions in the context of conformational disorder.

Introduction

Specificity of biomolecular interactions is often conflated with folded or otherwise ordered structures. It has long been the goal of molecular biology to catalog the various ordered structures that biomolecules adopt and to characterize how their interfaces engage with one another. The language and criteria used to define specificity of biomolecular interactions is informed by and assumes ordered-structure-mediated interactions or "site-specific interactions", yet there is growing evidence that dynamic multivalent interactions involving intrinsically disordered regions (IDRs) can also exhibit specificity with functional outcomes as soluble complexes[1-9] and in the context of meso-scale assemblies referred to as biomolecular condensates[10-14]. Theoretical examples of such a statistical mechanism of "fuzzy" molecular recognition afforded by IDRs include the sequence-dependent co-mixing and de-mixing of condensed polyampholyte species[13,15,16] and other model IDRs[14,17].

We recently demonstrated experimentally that condensates composed of a large IDR of MED1 (MED1[IDR]), the largest subunit of the transcriptional coactivator complex Mediator, selectively partitioned RNA Polymerase II (RNA Pol II) together with positive regulators of Pol II (e.g. SPT6) while excluding negative regulators of Pol II (e.g. NELFE)[10]. This functional specificity of MED1[IDR] condensates required the sequence patterning of charged amino acids on disordered regions of partitioned partners into high local densities of either positive or negative charge, referred to as charge blocks. For example, alternating positive and negative charge blocks were found on the disordered region of SPT6 (SPT6[IDR]) but were absent from the disordered region of NELFE (NELFE[IDR])[10]. Experimental manipulation of SPT6[IDR] to remove charge blocks disrupted partitioning and manipulation of NELFE[IDR] to add charge blocks promoted partitioning with functional consequences for RNA Pol II transcription. While these experiments provided evidence for the importance of sequence charge patterning in selective partitioning, it remained unresolved whether the observed specificity required an unknown ordered-structure-mediated interaction or were largely accounted for by dynamic multivalent contacts among the IDRs. Given the connection between this experimental work and theoretical treatments of sequence-dependent co-mixing and de-mixing of condensed polyampholytes[13,15,16], here we test the extent to which a polymer theory-based statistical model for disordered chain molecules can explain our experimental data and make new testable predictions of MED1[IDR] condensate specificity.

In the context of experimental observations that charge-patterning is required for specificity of MED1[IDR] condensates, several theoretical approaches have been developed in recent years to account for various effects of sequence charge pattern on IDR behaviors. These efforts include using simulation and theory to investigate effects of sequence charge pattern on the conformational dimensions and phase separation propensities of polyampholytes (chain sequences with approximately equal numbers of positively and negatively charged units), leading to the recognition that sequence with more blocky charge patterns tend to adopt more compact conformations as an isolated chain[18,19], and they also have a higher propensity to phase separate. The two effects appear to be significantly correlated though this correlation is not without limitations[20]. Sequence charge pattern also affects chain dynamics within condensates in molecular dynamics simulation[21]. Of particular relevance to the present investigation of IDR partitioning is that under the general field-theoretic framework, both approximate analytical theory[22] and field-theoretic simulations[23-25] have been applied to study sequence-specific effects in the mixing of condensed IDR species[15,16]. On the basis of these advances, we now introduce a novel, computationally efficient, and high throughput method derived from analytical polymer theory for predicting the partitioning of charged IDR sequences into a biomolecular condensate scaffolded by another charged IDR.

As detailed below, this high throughput method can be used to scan libraries of protein sequences to make predictions about the extent to which a protein sequence will (or will not) be partitioned into MED1[IDR] condensates. After demonstrating that our method is capable of validating previously published experimental data, we make predictions for sequences for which there is no experimental data. Predictions of partitioning are then tested in cell-based assays where we can assess the degree of partitioning of any

arbitrary sequence into a condensate of MED1^{IDR} tethered to a specific genomic locus.
The cell-based assays show striking qualitative agreement with the model predictions and
for the cases where the model disagrees with experimental results, provide quantitative
insights into non-electrostatic contributions to selective partitioning in the MED1^{IDR}
system, which may be incorporated into a more refined treatment in future studies. Taken
together, these finding demonstrate, in the context of the experiments performed here,
that selective partitioning can largely be explained by dynamic multivalent contacts
among disordered regions without needing to invoke ordered-structure-mediated
interactions.

101
102

Results

Rationale
Partitioning of IDRs into MED1^{IDR} condensates or any other IDR condensate is governed
by multiple-chain interactions wherein the IDR and MED1^{IDR} chain conformations are
expected to be disordered. Accordingly, pertinent biophysical properties that are
thermodynamic in nature are determined by appropriate statistical averages over possible
configurations entailing IDR-IDR, MED1^{IDR}-MED1^{IDR}, and IDR-MED1^{IDR} interactions.
Based on the presumed predominance of these interactions in MED1^{IDR}-scaffolded
condensates, for simplicity our model does not account for possible interactions involving
other molecular species that are present inside the cell. Here, as a first step, we are
interested primarily in the effect of IDR sequence charge pattern on the electrostatics of
partitioning, in which case the IDR-MED1^{IDR} interaction energy U_{ij} (where the label i
stands for the IDR in question and the label j stands for MED1^{IDR} as in Eqs.1 and 2 below)
is the usual screened-Coulomb potential given in Materials and Methods [Eq.3]. In view
of the previously observed prominence of sequence charge pattern effects on IDR
partitioning into MED1^{IDR} condensates[10], it is conceptually worthwhile to first narrow our
focus on probing how much electrostatics alone can physically rationalize such
partitioning. Accordingly, while contributions from non-electrostatic interactions will clearly
need to be considered in a more comprehensive approach, they will only be touched upon
briefly in the present analysis.

Because the calculation of thermodynamic averages in principle entails considering all
possible chain configurations, it is computationally challenging to study such a multiple-
chain system using molecular dynamics simulations, especially if the goal is to assess
the predicted IDR-sequence-dependent behaviors for many such systems for different
IDRs. In lieu of taking into account all possible chain configurations, we found an intuitive
way to envision how the charge sequences of an IDR and MED1^{IDR} may affect IDR
partitioning, namely by paying attention to a set of "sliding-sequence" configurations as
representative of the underlying, much more complex IDR-MED1^{IDR} interaction. The
scheme is depicted in **Fig.1A** using natural disordered regions from NELFE and SPT6 as
examples (sequences for all regions used in this study are provided in **Table S1**). These
disordered regions are chosen as examples because we have previously determined
experimentally that this disordered region of SPT6 has high partitioning while the
disordered region of NELFE has low partitioning[10].

In **Fig.1A**, the essential physics of charge-sequence-dependent IDR-MED1[IDR]
interactions is illustrated by the position-dependent screened Coulomb energy
$U_{ij}(x)/k_{\mathrm{B}}T$ between the disordered regions of MED1 and NELFE (orange lines), and
separately with the N-terminal disordered region of SPT6 (blue lines) as each of the IDRs
slides along MED1[IDR] (indicated by the arrows) at a perpendicular distance of 3 C_α-C_α
virtual bond lengths (≈ 11.4 Å), wherein a lower (more negative) $U_{ij}(x)/k_{\mathrm{B}}T$ value means
a more favorable IDR-MED1[IDR] interaction. Here x is the horizontal distance variable in
units of virtual bond length, k_{B} is Boltzmann's constant, T is absolute temperature.

Each of the horizontal dotted lines in **Fig.1A** (same color code for the IDRs) is the
arithmetic mean $\sum_x U_{ij}(x)/k_{\mathrm{B}}T \sum_x 1$ over the range of horizontal variable x considered;
the thicker dashed curves are the Boltzmann-weighted thermodynamic averages
$\langle U_{ij}\rangle/k_{\mathrm{B}}T \equiv \sum_x U_{ij}(x)\exp(-U_{ij}(x)/k_{\mathrm{B}}T)/k_{\mathrm{B}}T \sum_x \exp(-U_{ij}(x)/k_{\mathrm{B}}T)$. The latter average,
which exhibits a larger difference between the disordered regions of SPT6 and NELFE,
is expected to be representative of the physical situation because a basic tenet of
statistical mechanics is that configurations that are energetically more favorable are
populated more prominently in a constant-temperature ensemble. Accordingly, the
sliding-sequence analysis offers a physical rationalization for the experimental
observation that SPT6[IDR] partitions significantly more strongly into MED1[IDR] (i.e., has a
higher partition coefficient, PC) than NELFE[IDR] [10] because the results in **Fig.1A** shows
clearly that SPT6 interacts significantly more favorably with MED1 than NELFE when
averaged over the sliding-sequence configurations.

Although the sliding-sequence analysis offers semi-quantitative predictions for PC, the
procedure is time consuming because for a pair of sequences of length ~N, it requires
~N^3 energy calculations. Molecular dynamics simulation of the partitioning process is
even more computationally expensive. Another possible theoretical avenue is to apply
random phase approximation (RPA) formalism[24,26,27] to analytically model the binary
phase separation of an isolated system containing an IDR species and MED1[IDR] on equal
footing. An exploratory calculation for wildtype (WT) and two variants of SPT6[IDR], which
were experimentally determined to have reduced partitioning[10], produced a trend of
predicted PCs consistent with the experimentally observed trend, supporting RPA as an
effective means to capture the effect of sequence charge pattern on IDP interactions (SI
text and **Fig.S1**). However, as it stands, the application of this method to PC prediction is
limited because the model assumptions are too restrictive for quantitative match with
experiments, and it is numerically costly for any task that requires a high throughput.

The FH-RPA model
As described in Materials and Methods and detailed in SI, we found that experimental
PCs can be rationalized and predicted quantitatively to a substantial degree through
capturing the physics of IDR-MED1[IDR] interaction approximately by treating MED1[IDR] as
the scaffold of a preformed condensate and the IDR as a client species. In this approach,
the theory-predicted PC for IDR type i into MED1[IDR] (labeled as j) is governed by a Flory
interaction parameter[28] χ_{ij}:

184
$$PC_i^{(\text{theory})} = c_1 + c_2 \chi_{ij} N_i \,,$$
[1]

186 where c_1 and c_2 are empirical coefficients to be determined by fitting this formula to
187 experimentally measured PCs. When only electrostatic interactions are taken into
188 account, χ_{ij} can be approximated as a sum of a zeroth order term $\left(\chi_\text{e}^{(0)}\right)_{ij}$ that depends
189 only on the net charges of the IDR and MED1$^{\text{IDR}}$, and a first order random phase
190 approximation (RPA) term

191
$$\left(\chi_\text{e}^{(1)}\right)_{ij} = 2\pi l_\text{B}^2 \rho_0 \int_0^\infty dk \frac{k^2}{(k^2+\kappa^2)^2} g_i(k)g_j(k)$$
[2]

that takes into account the sequence charge patterns of the IDR and MED1$^{\text{IDR}}$. In Eq.2,
l_B is the Bjerrum length that determines the strength of electrostatic interactions in a
thermal environment, ρ_0 is a reference density for the solution system, and $g_i(k)$, a
function of reciprocal space (Fourier-transformed) coordinate k, is dependent on the
sequence charge pattern and involves a smearing length a for the IDR residues in the
field-theoretic description. Charge patterns involving residue positions far apart along the
chain sequence (i.e., nonlocal patterns encompassing charges separated by long contour
lengths) are better captured by $g_i(k)$ with small k values, whereas charge patterns
involving residue positions close (local) to one another along the chain sequence are
better captured by $g_i(k)$ with large k values (expressions for l_B and $g_i(k)$ are provided in
Materials and Methods). Because this formalism combines Flory-Huggins (FH) mean-
field theory (MFT)[28] with RPA treatment of sequence-pattern-dependent electrostatic
interactions of charged polymers[24,26,27], we refer to it as the FH-RPA method (Eqs.1, 2, 4-
6).

As an initial validation test and parameter optimization, we used the experimentally
measured PCs of 30 IDR sequences into MED1$^{\text{IDR}}$ condensates (**Table S2** and **Fig.S2**)
to maximize their Pearson correlation coefficient r with the term $\chi_{ij} N_i$ in the theoretical
Eq.1 by optimizing l_B, κ, and a as fitting parameters (SI text and **Fig.S3**). The numerical
values of c_1 and c_2 (> 0), which do not affect r, were subsequently determined by a linear
fit to Eq.1 (SI text). This set of 30 partitioning measurements are compiled from our
previous publication[10] and include natural IDR sequences, variants of natural IDR
sequences, and synthetic designed sequences. These partitioning measurements are all
derived from the same cell-based fluorescence microscopy method using a U2OS cell
line with large repeats of LacO DNA elements integrated into its genome[29]. Due to the
high affinity interactions of LacI protein to LacO DNA, expression of a CFP-tagged LacI
fused to MED1$^{\text{IDR}}$ (CFP-LacI-MED1$^{\text{IDR}}$) leads to a high local concentration of MED1$^{\text{IDR}}$ at
the LacO repeat locus creating a synthetic condensate which can be detected using
confocal fluorescence microscopy as a bright CFP focus. To measure partitioning, we
introduce a second protein sequence fused to mCherry and measure mCherry
fluorescence signal inside and outside the CFP-LacI-MED1$^{\text{IDR}}$ focus. This strategy has
been used in many publications to assess partitioning in cells[10,12,30-35].

This first exercise with the FH-RPA model indicates that sequence charge pattern plays
a prominent role in determining IDR partitioning into MED1$^{\text{IDR}}$ condensates. **Fig.S4** shows
that if only the net charges of the IDRs are considered (MFT, using only the zeroth order

term), the Pearson correlation coefficient is merely $r \approx 0.35$ between theory-predicted and experimental PCs, whereas the correlation improves to $r \approx 0.67$ when sequence charge patterns are taken into account (RPA, using both the zeroth and first order terms). The correlation between experimental and theory-predicted PCs for the subset of 19 natural (WT) IDRs is shown in **Fig.1B**, again exhibiting improved correlation when sequence charge pattern is taken into account. These results demonstrate the utility of the FH-RPA model in predicting PC values for a relatively large library of IDRs. The results also demonstrate how combining mean field theory together with random phase approximation yields predictions that have greater agreement with experimental data.

FH-RPA model rationalizes effects of charge scrambling on IDR PCs

To illustrate how FH-RPA captures sequence charge pattern-dependent IDR-MED1^{IDR} interactions, we display the quantity $A_i(k) \equiv k g_i(k)/(k^2 + \kappa^2)$ in **Fig.2A** for MED1^{IDR} and for the IDRs of NELFE and SPT6. These IDRs are chosen as examples of well partitioned (SPT6) and poorly partitioned (NELFE) sequences. Since $\left(\chi_e^{(1)}\right)_{ij}$ is the main factor in the

FH-RPA-predicted PCs (Materials and Methods) and $\left(\chi_e^{(1)}\right)_{ij}$ is proportional to

$\int_0^\infty dk \, A_i(k) A_j(k)$ in accordance with the above Eq.2, the larger theory-predicted PC for SPT6 is seen here as a result of the larger $A_i(k)$ value of SPT6^{IDR} over that of NELFE^{IDR} for the entire range of k plotted in **Fig.2A**. How this difference arises from the two IDRs' sequence charge patterns is further illustrated by the heat maps in **Fig.2B** for the quantity $A_{\alpha\beta}(k) = [\widehat{\Gamma}(k)^2 k/(k^2 + \kappa^2)]\sigma_\alpha^{(i)}\sigma_\beta^{(i)}\exp(-|\alpha - \beta|k^2 b^2/6)$ at two k values. $A_{\alpha\beta}(k)$s are individual terms contributing to $A_i(k)$, viz., $A_i(k) = \sum_{\alpha=1}^{N_i} \sum_{\beta=1}^{N_i} A_{\alpha\beta}(k)/N_i$. The larger $A_i(k)$ values of SPT6^{IDR} than those of NELFE^{IDR} arise from more positive orange-colored areas in the heat maps for SPT6^{IDR} than those for NELFE^{IDR}.

As an illustration of how FH-RPA can discriminate between sequence patterning in qualitative agreement with previously published experimental data, we focused on sequence rearrangement experiments of SPT6^{IDR} or NELFE^{IDR}. In these experiments, overall sequence composition, and therefore average charge properties, are maintained but the position of residues is manipulated in either sequence to remove, maintain, or introduce charge patterning associated with partitioning into MED1^{IDR} condensates. For SPT6^{IDR} we made three different types of scrambled sequences (**Fig.2C**). Two of these remove blocky charge patterning by either scrambling all amino acids (full scramble) or scrambling only charged amino acids leaving non-charged amino acids unchanged (charge scramble). In the third type of sequence rearrangement, we scrambled the sequence of non-charged amino acids leaving charge amino acids unchanged thereby maintaining the wildtype charge patterning. This "non-charge scramble" was designed in three random iterations. Previously published[10] results compiled here are in qualitative agreement with changes to PC derived from FH-RPA, both showing that removing blocky charge pattern disfavors partitioning and manipulations that maintain blocky charge pattern do not affect partitioning (**Fig.2D**). For NELFE^{IDR}, we rearranged charged amino acids to introduce charge blocks (**Fig.2E**) leading to an increase in partitioning measured experimentally and to an increase in PC values derived from FH-RPA (**Fig.2F**). Taken

together, the FH-RPA model is able to explain changes in partitioning due to manipulations of sequence patterning even when the sequences have the same average charge properties. Notably, the MFT approach will fail in this regard because the MFT-predicted PC depends only on the amino acid composition of the sequence and therefore MFT always predicts the same PC for all scrambled versions of a given sequence.

FH-RPA model successfully identifies high-PC regions in proteins
In view of the significant dependence of predicted PCs with IDR chain length (SI text) and the possibility of IDR chain-length-dependent reduction in conformational entropy upon partitioning into a condensed polymeric environment[36], we next conducted a focused test of the FH-RPA model by considering IDR sequence segments sharing the same length. We do so by computing theory-predicted PCs for sliding sequence windows of 200 residues. The protocol, as described in **Fig.3A**, entails sliding a 200-residue window (grey bar, horizontal arrows) along a full-length sequence (human CTR9 in this example) and compute a theory-predicted PC at every step. This procedure produces a PC profile for the full-length sequence (black curve). To identify a high PC region within a full-length sequence, we first locate the position of the window with the maximum predicted PC (indicated by top horizontal red dashed line) and then take the two windows with 90% of the maximum predicted PC as the boundary of the high-PC region (horizontal orange bar in **Fig.3A**).

We used this method to create PC profiles of CTR9 and SPT6, two proteins partitioned into MED1[IDR] condensates and for which we have previously identified regions necessary and sufficient for partitioning[10]. Strikingly, the PC profile method identified regions within the N-terminal disordered region of SPT6 and the C-terminal disordered region of CTR9 (**Fig.3B and 3C**) which had been experimentally defined as necessary and sufficient for partitioning into CFP-LacI-MED1[IDR] foci[10]. By overlaying disorder predictions from Metapredict[37,38] (**Fig. 3B and 3C**, green line), plotted using the same average over sliding windows of 200 residues used for plotting the PC profile (orange line), we find that there is significant overlap between high PC and IDRs with high charge patterning (overlayed NCPR plot at the bottom). Interestingly, not all predicted disordered regions have high predicted PC (e.g. the C-terminal end of SPT6 in **Fig.3B**), as not all predicted disordered regions have the same charge or charge pattern characteristics. These results demonstrate the utility of the FH-RPA model in scanning full length proteins to identify regions with predicted high partitioning because those regions match experimentally validated regions responsible for partitioning of the protein.

PC profiles are conserved despite a reduction in positional sequence conservation
To investigate whether PC profiles are an evolutionary conserved feature of proteins,
we performed the PC profile analysis on SPT6 (**Fig.3D**) and CTR9 (**Fig.3F**) protein sequences from several animal species (human, mouse, chicken, frog and fish) (**Table S1**). The PC profiles of CTR9 or SPT6 from these different species were strikingly similar and all identified a high PC region in the same region of the protein (orange-shaded areas in **Fig.3D and 3F**). Sequence alignments of the high PC regions showed that % identity as defined by sequence alignment (Clustal omega) became lower relative to the human

sequence with evolutionary distance for both SPT6 (**Fig.3E,** left) and CTR9 (**Fig.3G,** left). The poor conservation of positional sequence is expected for a region predicted to be disordered[39-42]. Nonetheless, when PC profiles of the high PC regions of SPT6 (**Fig. 3E**, right) or CTR9 (**Fig. 3G**, right) from different species were quantitatively compared to one another (Materials and Methods, Eq.7), they were nearly identical—as is also quite clear from inspection of **Fig.3D** and **3F**, suggesting that the high PC value is evolutionary conserved even though there is a reduction in positional sequence conservation. To test this prediction, we expressed the high PC regions of SPT6 (**Fig.3H-3J**) or CTR9 (**Fig.3K-3M**) from the indicated species and measured their partitioning into CFP-LacI-MED1^{IDR} foci in Lac array cells. While there was some targeting of these mCherry tagged proteins to other regions of the nucleus (**Fig. 3I and 3L**), high PC regions of SPT6 from different species (**Fig.3I**) and CTR9 from different species (**Fig.3L**) all partitioned into the CFP-LacI-MED1^{IDR} foci. Quantification focused on mCherry partitioning into CFP-LacI-MED1^{IDR} foci relative to an mCherry alone control showed that all predicted high PC regions from SPT6 (**Fig.3J**) and CTR9 (**Fig.3M**) have significant enrichment. These results show that the FH-RPA PC profile method can identify regions of full-length proteins that partition into CFP-LacI-MED1^{IDR}, with PC profiles for SPT6 and CTR9 well conserved across species though the conservation of the amino acid residues along the sequence decreases significantly with evolutionary distance.

FH-RPA model successfully predicts low-PC and high-PC IDR windows

Given the utility of FH-RPA to scan through arbitrary protein sequences and the striking agreement of predictions with experimental data, we next sought to scan through the proteome to identify novel sequences with high or low predicted PC values and test these predictions in the Lac array cell system. We decided to perform the tiling analysis with 200 amino acid windows and focused this analysis on long ($\geq$ 200 amino acids) terminal disordered regions of nuclear proteins involved in transcription (**Fig.4A**). Starting with the UniProt reviewed human proteome, we first identified predicted disordered regions using Metapredict[37,38], filtered for regions $\geq$ 200 amino acids, filtered for regions found within 10% of the N or C terminus, and finally filtered for disordered regions from proteins annotated with keyword "transcription" by UniProt. This yielded 865 IDRs with lengths ranging from 200 to 2869 amino acids. We then calculated FH-RPA derived PC values for 200 amino acid tiles scanning all sequences. This represented calculation of PC for 165,621 regions.

Given the contribution of electrostatics in the FH-RPA model calculation, we sought to compare 200 amino acid regions with comparable average charge but with different PC scores. We reasoned that comparing regions predicted to have different PC values but which differed dramatically in charge properties would not be a good test of the prediction. We therefore only focused on regions with relatively high acidic (25%<acidic<33%) and basic (10%<basic<33%) amino acid fraction and roughly similar isoelectric points (4.0 - 4.65). This approach yielded a set of 4 regions predicted to have high PC and 7 predicted to have low PC. We included SPT6^{IDR} as our positive control and a 200 amino acid window of NELFE^{IDR} with the highest fraction of acidic amino acids as a negative control. Key parameters for SPT6^{IDR} and predicted high PC regions can be found in **Fig.4B** and those for NELFE and predicted low PC regions can be found in **Fig.4C**. All sequences

can be found in **Table S1**. We expressed these sequences as mCherry fusions together with CFP-LacI-MED1^{IDR} in Lac array cells and measured their partitioning into CFP-LacI-MED1^{IDR} foci in cells. In this way, we were testing sequences of the same length, comparable average charge properties, but different predicted PCs.

NCPR profile analysis of the 4 new high PC regions showed that these regions had high local density of charge (**Fig.4D**). In Lac array cells, all four of these high PC regions partitioned into CFP-LacI-MED1^{IDR} foci (**Fig.4E**). While there was some variability of where these regions were targeted elsewhere in the nucleus (**Fig.4E**), quantification focused on mCherry enrichment at CFP-LacI-MED1^{IDR} foci shows significant enrichment for the four high PC regions relative to mCherry alone (**Fig.4F**). These results demonstrate that the FH-RPA can successfully identify well partitioned regions from an arbitrary list of sequences.

While four out of four (100%) of the high PC predictions were validated experimentally, only four out of seven (57%) of the low PC regions matched expectations and were not partitioned into CFP-LacI-MED1^{IDR} foci (**Fig.4G-4I**). Surprisingly, three out of seven were partitioned into CFP-LacI-MED1^{IDR} foci (**Fig.4J-4L**), suggesting that the FH-RPA model was missing some feature of these protein regions that were contributing to partitioning. There was nothing obvious about the difference in NCPR profiles (**Fig. 4G and 4J**) which prompted us to examine potential non-electrostatic interactions for these 3 outliers.

FH-RPA model can be improved by including non-electrostatic effects
We examined possible reasons for why our currently implemented FH-RPA model predicted low PCs for regions of CDCA7L, ADNP, and ST18 that were shown experimentally to partition significantly into CFP-LacI-MED1^{IDR} foci (**Fig.4J-4L**). One limitation of our current FH-RPA model is that it does not account for non-electrostatic IDR-MED1^{IDR} interactions. We found one prominent difference in amino acid composition between the IDRs that are correctly predicted to have low PC (ATF7IP, BDP1, ZNF638, LRRHIP1) and the aforementioned three IDRs that are incorrectly predicted to have low PC: number of aromatic residues (F, Y, W). While the IDRs that are correctly predicted to have low PCs have at most 4 aromatic residues (ATF7IP: 4, BDP1: 1, ZNF638: 4, LRRHIP1: 2), the IDRs that are predicted to have low PCs but in fact have high or moderate experimental PCs have significantly more (≥10) aromatic residues (CDCA7L: 13, ADNP: 10, ST18: 10, NELFE: 10). Since MED1^{IDR} has a large number of cationic residues (75 K, 18 R), multivalent cation-π MED1^{IDR}-IDR interactions could be playing a role[43]. It follows that a likely explanation for the inability of the current FH-RPA model to correctly predict the moderate to high experimental PCs of CDCA7L, ADNP, ST18 (and to a much lesser extent NELFE) is the model's lack of consideration of non-electrostatic, and especially π-related interactions.

Aiming to explore whether incorporating π-related interactions can improve accuracy of PC prediction in general, we apply the sliding-sequence protocol in **Fig.1A** but now we compare the Boltzmann-averaged energies computed with only electrostatic (screened Coulomb) interactions against those computed with both electrostatic and non-

electrostatic interactions. Several pairwise amino acid interaction schemes were
developed recently to model IDR conformations and phase separations[44-47]. Here we
adopt, as an example, the Mpipi potential[46] because it has proven useful in accounting
for π-related IDR interactions[24,46]. The resulting changes in Boltzmann-averaged energies
for the 16 IDR sequences studied in **Fig.4** are shown in **Fig.4M**. Assuringly, the IDRs
predicted by FH-RPA to have high PCs (pink-shaded) have lower (more favorable)
electrostatics-only Boltzmann-averaged energies (left vertical scale), as a group, than
IDRs predicted by FH-RPA to have low PCs (blue-shaded), indicating that FH-RPA and
the sliding-sequence protocol consistently capture the electrostatic effects on IDR
partitioning into MED1[IDR]. When non-electrostatic effects are included, the Boltzmann-
averaged energies are shifted (right vertical scale). The IDRs with high predicted PCs
remain favorable (black and dark-gray bars in pink background), underscoring the
robustness of the FH-RPA predictions for this group of sequences. The IDRs with correctly
predicted low PCs remain similarly unfavorable or become even less favorable (black
bars in blue background shifting to higher or only very slightly lower Boltzmann-averaged
energies represented by the dark-gray bars), indicating that the FH-RPA predictions for
this group of sequences are also robust. Interestingly, the group of sequences incorrectly
predicted by FH-RPA to have low PCs (but in fact have moderate to high experimental
PCs) all become more favorable (red and green bars shifting to lower Boltzmann-
averaged energies), though not sufficiently favorable to match the favorability of those
sequences correctly predicted by FH-RPA to have high PCs. Nonetheless, this exercise
demonstrates that incorporation of non-electrostatic effects can improve accuracy of PC
prediction. A possible first step in the future development of such an improved prediction
scheme is to employ a mean-field approximation of non-electrostatic interactions[24].
Although such an approach does not account for the sequence pattern of uncharged
residues, it may still offer advantages because of its account of hydrophobic and aromatic
amino acid compositions (see SI for additional information).
437
To test this prediction from the FH-RPA + Mpipi correction, we mutated the aromatic
residues in the experimental high-PC CDCA7L, ADNP, and ST18 IDR sequences to
alanine. In **Fig.4M**, these aromatic to alanine mutant sequences are marked by an
asterisk and their Boltzmann-averaged energies are represented by magenta bars, all of
which remain high or become even slightly higher when non-electrostatic effects are
included, indicating that partitioning is disfavored. Consistently, in Lac array cells, all three
of the anomalous regions with aromatic to alanine substitutions had reduced partitioning
into CFP-LacI-MED1[IDR] foci compared to the wildtype sequence (**Fig.4N and 4O**).
Quantification focused on mCherry enrichment at CFP-LacI-MED1[IDR] foci showed that
while the WT regions had significant enrichment relative to the mCherry-alone control,
the aromatic to alanine substitutions reduced partitioning to levels comparable to or
indistinguishable from the mCherry-alone control.
450
Taken together, these results demonstrate that multivalent contacts among disordered
regions are sufficient to explain the selective partitioning observed for condensates
composed of the MED1[IDR]. The FH+RPA model allowed for calculating predicted PC
values for large libraries of sequences. The predictions had striking qualitative agreement

with previously published experimental results and predictions of the model were
experimentally validated.
457
458 **Discussion**
459
Disordered regions of proteins can promote the formation of biomolecular condensates
that partition specific sets of proteins[10,12-14,30,31,48-51], yet the mechanism underlying this
specificity has been challenging to investigate. We recently demonstrated that the pattern
of charged amino acids on IDRs was responsible for selective partitioning into
condensates composed of MED1[IDR], but we were unable to identify the mechanism of
this specificity. Here we demonstrate that the selective partitioning of intrinsically
disordered regions (IDRs) into MED1[IDR] condensates can be largely explained by
multivalent electrostatic interactions. Using a polymer physics-based approach, we
developed the FH-RPA model, which integrates Flory-Huggins mean-field theory with a
random phase approximation term to predict IDR partitioning based on sequence charge
patterning. Our model successfully rationalized and predicted partitioning behaviors
observed in previous experimental data and made predictions which were subsequently
validated by experiments. By systematically scanning protein sequences, we identified
novel high-partitioning IDRs and experimentally confirmed their behavior, highlighting the
utility of this model in identifying regions responsible for partitioning. While our approach
accurately predicted partitioning for most sequences, discrepancies for certain IDRs with
significant aromatic content revealed the need to incorporate non-electrostatic
interactions, such as cation-π interactions, to fully account for partitioning behavior. These
results suggest that biomolecular condensate composition can, at least in part, be
governed by statistical properties of disordered sequences.
480
Here we present theoretical effort focused largely on electrostatics, which plays important
roles in the assembly[22,52] and material properties[53] of biomolecular condensates, and
manifested also in salt-dependent phase separation of oligomeric peptides[54] and
biological IDRs[55-57] as well as the significant impact of sequence charge pattern on the
condensation of biological IDRs[52,58] and the complex coacervation of multiple species of
synthetic charged polymers[59,60]. Another exciting aspect of electrostatic contributions to
condensate composition is the potential for regulation by post-translational modifications
(PTMs). For example, multi-site phosphorylation has the potential to dramatically change
the charge properties of disordered proteins. Indeed, the hyper-phosphorylation of the
disordered C-terminal domain of RNA Polymerase II has been shown to change its
partitioning behavior[48,49]. Future efforts to investigate whether PTMs regulate the
partitioning of IDRs more generally will be an exciting future direction.
493
Taking a broader view, it should be noted that in the present implementation of our
methodology, the empirical parameters of the model are determined by fitting only
experimental data on (client) IDRs partitioning into MED1[IDR] condensates. As such, the
present implementation is not expected to address IDR partitioning into condensates
scaffolded by other disordered proteins. Intuitively, how is the partitioning of a given client
IDR into a MED1[IDR] condensate related to the IDR's partitioning into a condensate
scaffolded by another disordered protein will depend on the nature of the other scaffold.

For example, one may expect the degrees of client IDR partitioning will be similar if the
sequence of the other scaffolding protein entails similar electrostatic interactions as those
of MED1^{IDR}, but the degrees of partitioning can be quite different if the interactions
engaged by the other scaffolding protein are dominated by hydrophobic effects. The
manner in which the present PC prediction method can be generalized to apply across
condensates with different scaffolds remains to be investigated.
507
These limitations aside, our studies already highlight how the predicted PC values for
MED1^{IDR}-scaffolded condensates are conserved across different species even in the
absence of positional sequence conservation. These observations add to the growing
recognition of the evolutionary conservation of overall IDR properties such as charge,
aromatic and hydrophobic contents, sequence motifs and repeats, and other molecular
features that are sometimes not readily discernible by inspecting the primary amino acid
sequences alone because sequence alignments often reveal little similarity; but these
molecular features can be predictive of biological function[39,40,42]. For instance,
conservation of function of sequentially highly diverged IDRs across species has been
observed to be underpinned by conservation of simple electrostatic properties in the
adaptor protein Ste50[61] or binding affinities with a particular target in the case of
p53TAD/MDM2[62] without conservation of the primary amino acid sequence. The
experimental observations in **Fig.3H-M** verify that the predicted high-PC windows for
CTR9 and SPT6 indeed partition well into MED1 for IDRs from all five animal species we
tested. Further efforts to investigate the co-evolution of IDR pairs may reveal additional
features responsible for specificity.
524
The results described here demonstrate that multivalent contacts among disordered
regions are sufficient to explain the selective partitioning observed for condensates
composed of the MED1^{IDR}. Complementary to recent advance in applying forcefield
parameters to predict IDR interaction properties[63,64] our high throughput analytical
FH+RPA model allowed for efficient calculation of predicted PC values for large libraries
of sequences. The predictions had striking qualitative agreement with previously
published experimental results and predictions of the model were experimentally
validated. In conclusion, our findings demonstrate, in the context of the experiments
performed here, that selective partitioning can largely be explained by dynamic
multivalent contacts among disordered regions without considering possible ordered-
structure-mediated interactions.
536
537
Acknowledgements

This work was supported by Canadian Institutes of Health Research (CIHR) grant NJT-
155930, Natural Sciences and Engineering Research Council of Canada (NSERC) grants
RGPIN-2018-04351 and RGPIN-2024-04167, and the computational resources provided
by the Digital Research Alliance of Canada to H.S.C. and National Institutes of Health
(NIH) grant GM147583 to BRS.

544
545
Author contributions

Conceptualization, JW, HSC, and BRS, data curation, JW, ND, and HL, formal analysis,
JW and ND, funding acquisition, HSC and BRS, investigation, JW, ND, and HL,
methodology, JW, ND, HL, and HSC, software, JW and ND, supervision, HSC and BRS,
visualization, JW, ND, HSC, and BRS, writing – original draft, JW, HSC, and BRS, writing
– review & editing, JW, ND, HL, HSC, and BRS
552
553
554 **MATERIALS AND METHODS**
555
556 **A general screened electrostatic potential and the FH-RPA formulation**
557
558 In general, the IDR-MED1 interaction energy U_{ij} (where the label *i* stands for the IDR in
559 question and the label *j* stands for MED1) takes the usual screened-Coulomb form,

560
$$U_{ij}\left(\{\mathbf{R}^{(i)}\}, \{\mathbf{R}^{(j)}\}\right) = \frac{1}{4\pi\epsilon_0\epsilon_r} \sum_{\mu=1}^{n_p^{(i)}} \sum_{\nu=1}^{n_p^{(j)}} \sum_{\alpha=1}^{N_i} \sum_{\beta=1}^{N_j} \frac{\sigma_\alpha^{(i)}\sigma_\beta^{(j)}}{\left|\mathbf{R}_{\mu,\alpha}^{(i)}-\mathbf{R}_{\nu,\beta}^{(j)}\right|} \exp\left(-\kappa\left|\mathbf{R}_{\mu,\alpha}^{(i)} - \mathbf{R}_{\nu,\beta}^{(j)}\right|\right), \quad [3]$$

where ϵ_0 and ϵ_r are, respectively, vacuum and relative permittivity, $n_p^{(i)}$ and $n_p^{(j)}$ are,
respectively, the numbers of chains of type *i* and type *j* in the system (chains labeled by
μ, ν), N_i and N_j are the numbers of residues in sequence of types *i* and *j* (residue
positions labeled by α, β), $\sigma_\alpha^{(i)}$, $\sigma_\beta^{(j)}$ are, respectively, the charges at sequence positions
α, β for sequence types *i*, *j*, $\mathbf{R}_{\mu,\alpha}^{(i)}$ and $\mathbf{R}_{\nu,\beta}^{(j)}$ are the vectors for the spatial positions of (*i*, μ,
α)- and (*j*, ν, β)-labeled residues, and κ is the inverse Debye screening length.
567
Calculation of thermodynamic averages requires theoretically considering all possible
chain configurations or conducting sufficient sampling thereof computationally. The
intuitive "sliding-sequence" analysis (**Fig.1A**) surmises that the limited set of
configurations considered by sliding the IDR sequence over the MED1 sequence is
representative of the full IDR-MED1 interaction such that it can provide semi-quantitative
physical insights despite restricting to a particular $\{\mathbf{R}^{(i)}\}, \{\mathbf{R}^{(j)}\}$ instead of covering all
configurational possibilities rigorously. The results in **Fig.1A** were obtained using Bjerrum
length $l_B = e^2/4\pi\epsilon_0\epsilon_r k_B T = 7$ Å, where *e* is elementary (protonic) charge, $\epsilon_r \approx 80$ for bulk
water ($T \approx 300$ K), and $\kappa^1 = 10$ Å corresponding to [NaCl] $\approx$ 100mM.
577
578 In the present FH-RPA formulation, we consider only electrostatic contributions to χ_{ij} (as
579 indicated by the "e" subscript), viz.,

580
$$\chi_{ij} \to (\chi_e)_{ij} \approx \left(\chi_e^{(0)}\right)_{ij} + \left(\chi_e^{(1)}\right)_{ij} \quad\quad [4]$$

581 is composed of a zeroth order term

582
$$\left(\chi_e^{(0)}\right)_{ij} = -\frac{2\pi l_B \rho_0}{\kappa^2}\sigma^{(i)}\sigma^{(j)} \quad\quad [5]$$

that depends only on the IDR and MED1 net charges $\sigma^{(i)}$ and $\sigma^{(j)}$, where $\sigma^{(i)} = \sum_{\alpha=1}^{N_i}\sigma_\alpha^{(i)}$
is the net charge of sequence type *i* (e.g., the net charges for MED1, NELFE, and SPT6

are +43, −2, and −62, and their chain lengths N_i or N_j are 626, 242, and 201,
respectively). The first order term $\left(\chi_e^{(1)}\right)_{ij}$ is given by Eq.2 in terms of

$$g_i(k) = \frac{\hat{\Gamma}(k)^2}{N_i}\sum_{\alpha=1}^{N_i}\sum_{\beta=1}^{N_i}\sigma_\alpha^{(i)}\sigma_\beta^{(i)}\exp(-|\alpha-\beta|k^2 b^2/6) \qquad [6]$$

where k is reciprocal space (Fourier-transformed) coordinate variable, with small and
large k associating, respectively, with large and small (real) spatial distances, and
$\hat{\Gamma}(k) = \exp(-a^2 k^2/2)$ is a Gaussian smearing function (see SI for details).
591
For the NELFE and SPT6 examples in **Fig.2A** and **Fig.2B**, the theory-predicted PCs are
overwhelmingly contributed by the first-order sequence charge pattern term, with
$\left(\chi_e^{(0)}\right)_{ij}/\left(\chi_e^{(1)}\right)_{ij} \approx 10^{-5}$ for NELFE and $\approx 10^{-4}$ for SPT6. An illustration of how the RPA
term $\left(\chi_e^{(1)}\right)_{ij}$ as defined above is affected by the IDP's sequence charge pattern through
the quantity $g_i(k)$ in our formulation is provided in **Fig.2A,B** for the NELFE and SPT6
examples.
598
Calculating PC profiles
PC profiles are computed using 200-residue windows in the present study. For a given
sequence, we determine $\mathrm{PC}_i^{(\text{theory})}$ (Eq.1) for successive sequence windows comprising
of 200 residues. A PC profile is a collection of such $\mathrm{PC}_i^{(\text{theory})}$ values, wherein the
$\mathrm{PC}_i^{(\text{theory})}$ value of each 200-residue sequence window is plotted at the position of the
101th residue of the sequence windows (**Fig.3A**).
605
606   **Identifying high-PC regions**
607   In the present study, we identify a high-PC region within a sequence by first locating the
608   position of a 200-residue window with the maximum $\mathrm{PC}_i^{(\text{theory})}$ for the entire sequence. A
609   high-PC region is a collection of consecutive 200-residue windows with $\mathrm{PC}_i^{(\text{theory})}$ values
610   of at least 90% of the maximum $\mathrm{PC}_i^{(\text{theory})}$(**Fig.3A**).
611
PC profile similarity score
Let $\mathrm{PC}_{i,s}^{(\text{theory})}$ denote the theory-predicted partition coefficient (PC) of window s (Eq.1) of
sequence i (consisting of segment of residues $s, s+1, \ldots, s+M-1$ where M is the
sequence window size). We then define $x_s^{(i)} = \mathrm{PC}_{i,s}^{(\text{theory})} - 1$ and $x_s^{(i')} = \mathrm{PC}_{i',s}^{(\text{theory})} - 1$
for two sequences of equal lengths $N_i = N_{i'} \equiv N$. A similarity score between their PC profiles
may be constructed as
618

619 $$S_{i,i'} = 1 - \frac{\sum_{s=1}^{N-M+1}\left|x_s^{(i)}-x_s^{(i')}\right|}{\sum_{s=1}^{N-M+1}\left|x_s^{(i)}\right|+\left|x_s^{(i')}\right|} \qquad [7]$$

620
621   This similarity score is constructed such that $S_{i,i'} = 1$ for two identical PC profiles whereas any
622   sequence i has similarity $S_{i,i'} = 0$ against the trivial profile $\mathrm{PC}_{i',s}^{(\text{theory})} = 1$ for all s. For

623   sequences of different lengths, $N_i > N_{i'}$, we define the similarity score $S_{i,i'}$ as the maximum

624     $S_{i,i'}$ between i' and all sub-sequences of i with length $N_{i'}$, and vice versa if $N_i < N_{i'}$. %PC

625   similarity scores plotted in **Fig.3E** and **Fig.3G** is equal to 100 times $S_{i,i'}$.

Cell Culture

628   All cells were grown, as previously described[12], in full DMEM (Fisher Scientific 11995073)
629   supplemented with 10% FBS, penicillin-streptomycin (Thermo Fisher 15140122) and
630   GlutaMAX (Fisher Scientific 35050061) at 37 °C with 5% CO_2 in a humidified sterile
631   incubator.

Processing coverslips of co-transfected cells

As previously described[12], cells were seeded on coverslips (VWR 48366-067) in a 6-well
plate and let adhere overnight. Transfections using Lipofectamine™ 3000 Transfection
Reagent with designated plasmids were incubated overnight. After cells were transfected
and allowed to recover, cells were fixed in 4% paraformaldehyde (VWR BT140770) in
PBS for 10 min at room temperature (RT) in the dark (plate wrapped with foil). After three
washes in PBS for 10 min, samples were permeabilized with 0.5% triton X100 (Sigma
T9284) in PBS for 10 min at RT. Then cells were incubated with 1:5000 Hoechst 33342
(Thermo Fisher, 62249) in Milli-Q water at RT in the dark (5 min). After washing once
more with water, coverslips were mounted on slides (VWR 10144-820) with Vectashield
(VWR 101098-042). Coverslips were sealed with nail polish (VWR 100491-940) and
stored at 4°C.

Microscopy

647   Images were acquired, as previously described[12], using a Hamamatsu ORCA-Fusion
648   C14440 digital camera and a CSU-W1 Yokogawa Spinning Disk Field Scanning Confocal
649   System, and a 60x Plan Apo Lambda Oil Immersion objective (NA, 1.40). Thickness of Z-
650   slices were 0.2 μm. Exposure time and laser intensity were the same for samples imaged
651   in parallel. Laser lines used: 405nm, 488 nm, 561 nm, and 640 nm.

Microscopy analysis

mCherry partitioning was calculated, as previously described[12], by measuring the
integrated fluorescence at the Lac array locus core as defined by fluorophore fused to
LacI (highest intensity range on the Z-slice) and at a proximal spot in the nucleus
(background) using FIJI 2D measurement tools and ROI tools. The same ROI area was
used to measure the fluorescence at both background and focus. The quotient of these
values (arbitrary units) was calculated as raw intensity at locus divided by raw intensity in
the surrounding background.

NCPR profile plots

Representation of the position of positive and negative charge along a protein sequence
overlaid with net charge per residue (NCPR) profile provide a quick summary of the
charge patterning found in a protein region. The position of acidic or basic amino acids
are indicated as vertical red or blue lines, respectively. The horizontal black curve
represents the NCPR profile using a 10 amino acid sliding window, wherein the net charge
of successive 10-residue windows is plotted (10 times the net charge per residue).

Consequently, each NCPR profile has a possible max of +10 and a possible min of -10
centered at 0 charge. X-axis is the length of the protein and y-axis is charge index.
671
672

Statistics
GraphPad Software was used for statistical analysis. GraphPad notation for p values
was used in figure panels: (ns $p > 0.05$, $*p \leq 0.05$, $**p \leq 0.01$, $***p \leq 0.001$, $****p \leq$
0.0001). For figure 2D and 2F, data were compiled from various experiments previously
published[10] to compare qualitative trends to the model's predictions. For Fig.2D,
ordinary one-way ANOVA with Dunnet's multiple comparisons test to a "no IDR"
mCherry control (N, 11) was performed for WT SPT6^{IDR} (N, 10; p, <0.0001), full
scramble (N, 10; p, 0.6742), charge scramble (N, 10; p, >0.9999), and three
independent non-charge scrambles (each with N, 10; p, <0.0001). For Fig.2F, ordinary
one-way ANOVA with Dunnet's multiple comparisons test to a "no IDR" mCherry control
(N, 11) was performed for WT NELFE^{IDR} (N,10; p, 0.8805) and blocky (N, 10; p,
<0.0001). For Fig.3J, two-tailed unpaired t-tests relative to an mCherry control (N, 13)
were performed for high PC regions of SPT6 from human (N, 12; p, <0.0001), mouse
(N, 6; p, <0.0001), chicken (N, 8; p, <0.0001), frog (N, 8; p, 0.0003), or fish (N, 10; p,
<0.0001). For Fig.3M, two-tailed unpaired t-tests relative to an mCherry control (N, 13)
were performed for high PC regions of CTR9 from human (N, 9; p, <0.0001), mouse (N,
8; p, <0.0001), chicken (N, 11; p, <0.0001), frog (N, 8; p, <0.0001), or fish (N, 8; p,
<0.0001). For Fig.4F, two-tailed unpaired t-tests relative to an mCherry control (N, 9)
were performed for 200 amino acid windows predicted to have high PC from SPT6 (N,
21; p, <0.0001), KAT6B (N, 6; p, <0.0001), ATAD2 (N, 13; p, 0.0181), MYT1 (N, 9; p,
0.0032), or RSF1 (N, 8; p, 0.0001). For Fig.4I and 4L, two-tailed unpaired t-tests relative
to an mCherry control (N, 9) were performed for 200 amino acid windows predicted to
have low PC from ATF7IP (N, 11; p, 0.0083), BDP1 (N, 7; p, 0.0022), ZNF638 (N, 7; p,
0.0094), LRRFIP1 (N, 6; p, 0.2403), NEFLE (N, 8; p, 0.9475), CDCA7L (N, 5; p,
0.0003), ADNP (N, 13; p, 0.0345), or ST18 (N, 5; p, 0.0026). For Fig.4O, ordinary one-
way ANOVA with Dunnett's multiple comparisons test to a "no IDR" mCherry control (N,
9) was performed for WT and aromatic to alanine substitution mutants (aro-to-A) for
CDCA7L (WT: N, 5; p, <0.0001 and aro-to-A: N, 7; p, 0.1633), ADNP (WT: N, 13, p,
0.0499 and aro-to-A: N, 8; p, 0.9989), and ST18 (WT: N, 5; p, 0.0018 and aro-to-A: N,
16; p, 0.7019).
703
704

705 **Data Availability**
706 The data that support the findings of this study are available from the corresponding
707 authors upon reasonable request.

708
709
710

Figure 1

711
712

Fig. 1 | The FH-RPA model agrees with previously published experimental results
Fig.1A: IDPs with a blockier charge pattern interact more favorably with the IDR of
MED1. The vertical variable is $U_{ij}/k_\mathrm{B}T$ (potential energy given by Eq.[3] in units of
$k_\mathrm{B}T$). The horizontal variable is the distance between the centers of the MED1 and the
NELFE or SPT6 IDR sequence. Positively and negatively charged residues are
indicated by blue and red lines in the depiction of the MED1 and IDR sequences
(bottom, with the N and C termini marked). The NELFE and SPT6 sequences are
shown in their respective positions at which the interaction with MED1 is most favorable
(most negative energies marked by circles along the orange and blue curves). Each of
the dotted lines (same color code for the IDRs) is the arithmetic mean of the screened
Coulomb energy (vertical variable) over the horizontal range; the thicker dashed curves
are the Boltzmann-weighted average energies $\langle U_{ij}\rangle/k_\mathrm{B}T$.
Fig.1B: Scatter plots of published experimental versus theory-predicted PCs for the
natural sequences used in our parameter optimization. Correlation is significantly
enhanced in RPA (right) relative to MFT (left); *r* is Pearson correlation coefficient.
Sequence chain lengths are color coded. Experimental values are presented as the
average of 10 measurements.

730
731
732

Figure 2

733
734
735
Fig. 2 | FH-RPA model rationalizes effects of charge scrambling on IDR PCs
Fig.2A: IDR-MED1 interactions in RPA are accounted for by $A_i(k)$.
Fig.2B: The $A_{\alpha\beta}(k), \alpha \neq \beta$ heat maps for the smaller and larger *k* values [dashed lines
in (A)] are provided, respectively, by the lower and upper triangles. White areas
represent $A_{\alpha\beta}(k) = 0$ or $A_{\alpha\beta}(k) \approx 0$ (right scales). Sequence charge patterns of NELFE
and SPT6 [as in (A)] are shown along the heat map axes.
Fig.2C: Representation of the charge patterning and net charge per residue (NCPR)
profile of wildtype (WT) SPT6[IDR] and the indicated sequence variants. The position of
acidic or basic amino acids are indicated as vertical red or blue lines, respectively. The
horizontal black curve represents the NCPR profile using a 10 amino acid sliding
window, wherein the net charge of successive 10-residue windows is plotted (10 times
the net charge per residue). Consequently, each NCPR profile has a possible max of
+10 and a possible min of -10 centered at 0 charge.
Fig.2D: Left: compilation of previously published experimental data of the partitioning of
fluorescence signal from mCherry fused to the indicated control or SPT6[IDR] variant
inside and outside of CFP-LacI-MED1[IDR] foci in Lac array cells. Data are presented as a
boxplot (min-max). p-value, one-way ANOVA with multiple comparison test vs. the no
IDR control. These data were originally reported by[10]. Dashed horizontal line indicates
median of the "no IDR" control. Right: FH-RPA model predictions of partition coefficient
values (Eq.1) for indicated SPT6[IDR] variant into MED1[IDR] condensates.

Fig.2E: Similar to 2C, but for NELFE or a charge rearranged variant designed to mimic
the blocky charge patterning found in SPT6^{IDR}
Fig.2F: Similar to 2D, but for NELFE WT and charge rearranged variant (blocky).
759
760

Figure 3

Fig. 3 | FH-RPA model successfully identifies high-PC sequence windows in CTR9 and SPT6.

Fig.3A: Sliding-window analysis. Using human CTR9 as an example, PCs are computed for a 200-residue window (grey horizontal bar) sliding along (horizontal grey arrows) the full-length sequence and plotted at the midpoint of the sequence window (black curve in the main plot). The inset depicts the sequence charge pattern of the entire CTR9 as in Fig.1A, now with the black curve inside the inset showing sliding NCPR over consecutive 10-residue windows (inset vertical scale for charge ranges from −10 to +10 as in Fig.2C). The predicted high-PC region (orange bar) is constructed as

described in the text; the red bars are the first and last 200-residue windows in the high-
PC region.
Fig.3B: Sliding-window analysis for human SPT6. PC profile is plotted as in Fig.3A, now
in orange (left vertical scale), together with the Metapredict disorder profile computed for
the same set of 200-residue windows (green curve, right vertical scale). The disorder
regions, defined by disorder measure > 0.5, are shaded in green and indicated by the
horizontal green bar. The high-PC regions are indicated by the horizontal orange bar.
Fig.3C: Corresponding sliding-window analysis for human CTR9 (orange curve, same
data as in Fig.2A), now showing also the sequence disorder profile (green curve).
Fig.3D: PC profiles of the SPT6 IDR for different animal species are color-coded as
indicated, wherein the PC profile for the human sequence essentially overlaps with
those for the other sequences. The high-PC and high-disorder regions for the human
IDR sequence are marked as in Fig.3B using orange and green bars and shaded areas;
these colors do not share the meaning in the color code for the different sequences.
Fig.3E: Left: bar chart of % identity from Clustal omega of SPT6 high-PC regions
identified in different species relative to the human sequence. Right: % PC profile
similarities (as defined in Materials and Methods) between the same high-PC regions in
different species relative to the human sequence.
Fig.3F: Same as Fig.3D but for CTR9. The high-PC and high-disorder regions for the
human IDR sequence are marked as in Fig.3C.
Fig.3G: Same as Fig. 3E but for CTR9.
Fig.3H: Representation of the charge patterning and NCPR profile of high PC region
from SPT6 from the indicated species.
Fig.3I: Representative images of mCherry-tagged high PC regions of SPT6 from
indicated species co-transfected with CFP-LacI-MED1^{IDR} (human) in the Lac array cell
line. Top row shows mCherry fluorescence signal and bottom row shows CFP
fluorescence signal. White box designates the Lac array focus using the CFP channel.
Inset in top right corner of panels represents zoom in at CFP-LacI-MED1^{IDR} focus.
Dashed white line designates the cell nucleus. Scale, 5μm.
Fig.3J: Boxplot (min-max) of the partitioning as defined by the ratio of mCherry
fluorescence signal inside/outside of CFP-LacI-MED1^{IDR} foci for SPT6 high PC regions
from indicated species and an mCherry only control. Individual data points are plotted
on top of the box plot. Dashed horizontal line indicates median of the control. *p*-values
represent t-test for each high PC region relative to the control.
Fig.3K: Same as 3H but for CTR9 high PC regions from indicated species.
Fig.3L: Same as 3I but for CTR9 high PC regions from indicated species.
Fig.3M: Same as 3J but for CTR9 high PC regions from indicated species.

Figure 4

818 **Fig. 4 | FH-RPA model successfully predicts low-PC and high-PC IDR windows from**
819 **a large library of sequences.**

Fig.4A: Workflow to generate candidate set of 200 amino acid windows of IDRs from proteins involved in the same biological process (transcription) with comparable charge but either high or low predicted PC values.

Fig.4B: Key parameters of the candidate set of 200 amino acid windows with predicted high PC values selected for subsequent experiments. Protein name and UniProt IDs and the amino acid boundaries (window) that define the region are provided. For a given region, % acidic and basic amino acids and isoelectric point (pI) as well as FH-RPA derived PC values (PC) are provided.

Fig.4C: Same as Fig.4B but for the predicted low PC values

Fig.4D: Representation (as in Fig.2C) of the charge patterning and NCPR profile of 200 amino acid windows with predicted high PC.

Fig.4E: Representative images of mCherry-tagged predicted high PC windows of indicated protein co-transfected with CFP-LacI-MED1^{IDR} in the Lac array cell line. Top row shows mCherry fluorescence signal and bottom row shows CFP fluorescence signal. White box designates the Lac array focus using the CFP channel. Inset in top right corner of panels represents zoom in at CFP-LacI-MED1^{IDR} focus. Dashed white line designates the cell nucleus. Scale, 5μm.

Fig.4F: Boxplot (min-max) of the partitioning as defined by the ratio of mCherry fluorescence signal inside/outside of CFP-LacI-MED1^{IDR} foci for indicated 200 amino acid window and an mCherry only control. Individual data points are plotted on top of the box plot. Dashed horizontal line indicates median of the control. *p*-values represent t-test relative to the control.

Fig.4G: Representation of the charge patterning and NCPR profile of 200 amino acid windows with predicted low PC.

Fig.4H: Same as Fig.4E but for predicted low PC windows of indicated proteins.

Fig.4I: Same as Fig.4F but for predicted low PC windows of indicated proteins.

Fig.4J: Representation of the charge patterning and NCPR profile of 200 amino acid windows with predicted low PC but which show anomalous partitioning into CFP-LacI-MED1^{IDR} foci.

Fig.4K: Same as Fig.4E but for predicted low PC windows which show anomalous partitioning

Fig.4L: Same as Fig.4F but for predicted low PC windows which show anomalous partitioning

Fig.4M: Incorporation of non-electrostatic effects improve PC prediction. Sliding-sequence analysis is applied to IDRs with various RPA-predicted PCs (color coded). An intuitively chosen horizontal dashed line separates high (pink) and low (blue) predicted PCs. Boltzmann-averaged energies computed using only screened Coulomb potentials as in Fig.1A (solid bars, left vertical scale) are compared to those computed using the Mpipi potentials[46] (shaded bars, right vertical scale). Arrows indicate changes from screened Coulomb to Mpipi energies. Energies of IDRs with correctly predicted PC trends are shown in black, the 3 IDRs (in curly brackets) that were incorrectly predicted are in red, the one marginal case is in green, and the 3 aromatic to alanine mutants in Fig.4N,O are in magenta.

Fig.4N: Representative images of mCherry-tagged predicted low PC windows with aromatic residues substituted for alanine (aro to A mutants) co-transfected with CFP-LacI-MED1^{IDR} in the Lac array cell line. Compare results to the data for the wildtype

sequence presented in **Fig.4K**. Top row shows mCherry fluorescence signal and bottom
row shows CFP fluorescence signal. White box designates the Lac array focus using
the CFP channel. Inset in top right corner of panels represents zoom in at CFP-LacI-
MED1^{IDR} focus. Dashed white line designates the cell nucleus. Scale, 5μm.
Fig.4O: Boxplot (min-max) of the partitioning as defined by the ratio of mCherry
fluorescence signal inside/outside of CFP-LacI-MED1^{IDR} foci for indicated 200 amino
acid windows, the aromatic to alanine mutant (aro to A) and an mCherry only control.
Individual data points are plotted on top of the box plot. Dashed horizontal line indicates
median of the control. *p*-values represent one-way ANOVA with multiple comparisons
tests vs. control.

REFERENCES

Borg, M. *et al*. Polyelectrostatic interactions of disordered ligands suggest a physical basis for ultrasensitivity. *Proc National Acad Sci* **104**, 9650-9655 (2007). https://doi.org:10.1073/pnas.0702580104

Danielsson, J. *et al*. The Intrinsically Disordered RNR Inhibitor Sml1 Is a Dynamic Dimer. *Biochemistry* **47**, 13428-13437 (2008). https://doi.org:10.1021/bi801040b

Tompa, P. & Fuxreiter, M. Fuzzy complexes: polymorphism and structural disorder in protein–protein interactions. *Trends in Biochemical Sciences* **33**, 2-8 (2008). https://doi.org:10.1016/j.tibs.2007.10.003 PMID - 18054235

Mittag, T. *et al*. Structure/Function Implications in a Dynamic Complex of the Intrinsically Disordered Sic1 with the Cdc4 Subunit of an SCF Ubiquitin Ligase. *Structure* **18**, 494-506 (2010). https://doi.org:10.1016/j.str.2010.01.020

Mittag, T., Kay, L. E. & Forman-Kay, J. D. Protein dynamics and conformational disorder in molecular recognition. *Journal of Molecular Recognition* **23**, 105-116 (2010). https://doi.org:10.1002/jmr.961

Chen, T., Song, J. & Chan, H. S. Theoretical perspectives on nonnative interactions and intrinsic disorder in protein folding and binding. *Current Opinion in Structural Biology* **30**, 32-42 (2015). https://doi.org:10.1016/j.sbi.2014.12.002

Wu, S. *et al*. The Dynamic Multisite Interactions between Two Intrinsically Disordered Proteins. *Angewandte Chemie International Edition* **56**, 7515-7519 (2017). https://doi.org:10.1002/anie.201701883

Borgia, A. *et al*. Extreme disorder in an ultrahigh-affinity protein complex. *Nature* **19**, 31 (2018). https://doi.org:10.1038/nature25762 PMID - 29466338

Schuler, B. *et al*. Binding without folding – the biomolecular function of disordered polyelectrolyte complexes. *Current Opinion in Structural Biology* **60**, 66 - 76 (2020). https://doi.org:10.1016/j.sbi.2019.12.006

Lyons, H. *et al*. Functional partitioning of transcriptional regulators by patterned charge blocks. *Cell* **186**, 327-345 e328 (2023). https://doi.org:10.1016/j.cell.2022.12.013

Farag, M., Borcherds, W. M., Bremer, A., Mittag, T. & Pappu, R. V. Phase separation of protein mixtures is driven by the interplay of homotypic and heterotypic interactions. *Nat Commun* **14**, 5527 (2023). https://doi.org:10.1038/s41467-023-41274-x

De La Cruz, N. *et al*. Disorder-mediated interactions target proteins to specific condensates. *Mol Cell* **84**, 3497-3512 e3499 (2024). https://doi.org:10.1016/j.molcel.2024.08.017

Rana, U. *et al*. Asymmetric oligomerization state and sequence patterning can tune multiphase condensate miscibility. *Nature Chemistry*, 1-10 (2024). https://doi.org:10.1038/s41557-024-01456-6

Welles, R. M. *et al*. Determinants that enable disordered protein assembly into discrete condensed phases. *Nature Chemistry*, 1-11 (2024). https://doi.org:10.1038/s41557-023-01423-7

Pal, T., Wessén, J., Das, S. & Chan, H. S. Subcompartmentalization of polyampholyte species in organelle-like condensates is promoted by charge-pattern mismatch and

strong excluded-volume interaction. *Physical Review E* **103**, 042406 (2021). https://doi.org:10.1103/physreve.103.042406

Lin, Y.-H., Brady, J. P., Forman-Kay, J. D. & Chan, H. S. Charge pattern matching as a 'fuzzy' mode of molecular recognition for the functional phase separations of intrinsically disordered proteins. *New Journal of Physics* **19**, 115003 (2017). https://doi.org:10.1088/1367-2630/aa9369

Chew, P. Y., Joseph, J. A., Collepardo-Guevara, R. & Reinhardt, A. Thermodynamic origins of two-component multiphase condensates of proteins. *Chem Sci* **14**, 1820-1836 (2023). https://doi.org:10.1039/d2sc05873a

Das, R. K. & Pappu, R. V. Conformations of intrinsically disordered proteins are influenced by linear sequence distributions of oppositely charged residues. *Proc National Acad Sci* **110**, 13392 - 13397 (2013). https://doi.org:10.1073/pnas.1304749110 PMID - 23901099

Sawle, L. & Ghosh, K. A theoretical method to compute sequence dependent configurational properties in charged polymers and proteins. *The Journal of Chemical Physics* **143**, 085101 (2015). https://doi.org:10.1063/1.4929391 PMID - 26328871

Pal, T., Wessén, J., Das, S. & Chan, H. S. Differential Effects of Sequence-Local versus Nonlocal Charge Patterns on Phase Separation and Conformational Dimensions of Polyampholytes as Model Intrinsically Disordered Proteins. *The Journal of Physical Chemistry Letters* **15**, 8248-8256 (2024). https://doi.org:10.1021/acs.jpclett.4c01973

Hazra, M. K. & Levy, Y. Charge pattern affects the structure and dynamics of polyampholyte condensates. *Physical Chemistry Chemical Physics* **22**, 19368-19375 (2020). https://doi.org:10.1039/d0cp02764b PMID - 32822449

Lin, Y.-H., Forman-Kay, J. D. & Chan, H. S. Sequence-Specific Polyampholyte Phase Separation in Membraneless Organelles. *Phys Rev Lett* **117**, 178101 (2016). https://doi.org:10.1103/physrevlett.117.178101 PMID - 27824447

McCarty, J., Delaney, K. T., Danielsen, S. P. O., Fredrickson, G. H. & Shea, J.-E. Complete Phase Diagram for Liquid–Liquid Phase Separation of Intrinsically Disordered Proteins. *The Journal of Physical Chemistry Letters* **10**, 1644-1652 (2019). https://doi.org:10.1021/acs.jpclett.9b00099

Wessén, J., Das, S., Pal, T. & Chan, H. S. Analytical Formulation and Field-Theoretic Simulation of Sequence-Specific Phase Separation of Protein-Like Heteropolymers with Short- and Long-Spatial-Range Interactions. *The Journal of Physical Chemistry B* **126**, 9222-9245 (2022). https://doi.org:10.1021/acs.jpcb.2c06181

Lin, Y. H., Wessen, J., Pal, T., Das, S. & Chan, H. S. Numerical Techniques for Applications of Analytical Theories to Sequence-Dependent Phase Separations of Intrinsically Disordered Proteins. *Methods Mol Biol* **2563**, 51-94 (2023). https://doi.org:10.1007/978-1-0716-2663-4_3

Ermoshkin, A. V. & Olvera de la Cruz, M. Polyelectrolytes in the presence of multivalent ions: gelation versus segregation. *Phys Rev Lett* **90**, 125504 (2003). https://doi.org:10.1103/PhysRevLett.90.125504

Lin, Y.-H., Song, J., Forman-Kay, J. D. & Chan, H. S. Random-phase-approximation theory for sequence-dependent, biologically functional liquid-liquid phase

separation of intrinsically disordered proteins. *Journal of Molecular Liquids* **228**, 176-
193 (2017). https://doi.org:10.1016/j.molliq.2016.09.090

28   Flory, P. J. *Principles of Polymer Chemistry*. (Cornell University Press, 1953).

29   Janicki, S. M. *et al.* From Silencing to Gene Expression: Real-Time Analysis in Single
Cells. *Cell* **116**, 683 - 698 (2004). https://doi.org:10.1016/s0092-8674(04)00171-0

30   Chong, S. *et al.* Imaging dynamic and selective low-complexity domain interactions
that control gene transcription. *Science* **361** (2018).
https://doi.org:10.1126/science.aar2555 PMID - 29930090

31   Gan, P. *et al.* Coactivator condensation drives cardiovascular cell lineage
specification. *Science Advances* **10**, eadk7160 (2024).
https://doi.org:10.1126/sciadv.adk7160

32   Basu, S. *et al.* Unblending of Transcriptional Condensates in Human Repeat
Expansion Disease. *Cell* **181**, 1062 - 1079.e1030 (2020).
https://doi.org:10.1016/j.cell.2020.04.018 PMID - 32386547

33   Boija, A. *et al.* Transcription Factors Activate Genes through the Phase-Separation
Capacity of Their Activation Domains. *Cell* **175**, 1842 - 1855.e1816 (2018).
https://doi.org:10.1016/j.cell.2018.10.042 PMID - 30449618

34   Zamudio, A. V. *et al.* Mediator Condensates Localize Signaling Factors to Key Cell
Identity Genes. *Mol Cell* **76**, 753 - 766.e756 (2019).
https://doi.org:10.1016/j.molcel.2019.08.016 PMID - 31563432

35   Frazer, C. *et al.* Epigenetic cell fate in Candida albicans is controlled by transcription
factor condensates acting at super-enhancer-like elements. *Nature Microbiology* **5**, 1
- 16 (2020). https://doi.org:10.1038/s41564-020-0760-7

36   Chan, H. S. & Dill, K. A. Solvation: Effects of molecular size and shape. *The Journal of
Chemical Physics* **101**, 7007-7026 (1994). https://doi.org:10.1063/1.468327

37   Emenecker, R. J., Griffith, D. & Holehouse, A. S. Metapredict: a fast, accurate, and
easy-to-use predictor of consensus disorder and structure. *Biophysical Journal* **120**,
4312-4319 (2021). https://doi.org:10.1016/j.bpj.2021.08.039

38   Emenecker, R. J., Griffith, D. & Holehouse, A. S. Metapredict V2: An update to
metapredict, a fast, accurate, and easy-to-use predictor of consensus disorder and
structure. *Biorxiv*, 2022.2006.2006.494887 (2022).
https://doi.org:10.1101/2022.06.06.494887

39   Zarin, T. *et al.* Proteome-wide signatures of function in highly diverged intrinsically
disordered regions. *Elife* **8**, 1727 (2019). https://doi.org:10.7554/elife.46883

40   Zarin, T. *et al.* Identifying molecular features that are associated with biological
function of intrinsically disordered protein regions. *Elife* **10**, e60220 (2021).
https://doi.org:10.7554/elife.60220 PMID - 33616531

41   Pritišanac, I. *et al.* A Functional Map of the Human Intrinsically Disordered Proteome.
Biorxiv, 2024.2003.2015.585291 (2024). https://doi.org:10.1101/2024.03.15.585291

42   Lu, A. X. *et al.* Discovering molecular features of intrinsically disordered regions by
using evolution for contrastive learning. *Plos Comput Biol* **18**, e1010238 (2022).
https://doi.org:10.1371/journal.pcbi.1010238

43   Song, J., Ng, S. C., Tompa, P., Lee, K. A. W. & Chan, H. S. Polycation-π Interactions Are
a Driving Force for Molecular Recognition by an Intrinsically Disordered Oncoprotein

Family. *Plos Comput Biol* **9**, e1003239 (2013). https://doi.org:10.1371/journal.pcbi.1003239

Dignon, G. L., Zheng, W., Kim, Y. C., Best, R. B. & Mittal, J. Sequence determinants of protein phase behavior from a coarse-grained model. *Plos Comput Biol* **14**, e1005941 (2018). https://doi.org:10.1371/journal.pcbi.1005941

Das, S., Lin, Y.-H., Vernon, R. M., Forman-Kay, J. D. & Chan, H. S. Comparative roles of charge, π, and hydrophobic interactions in sequence-dependent phase separation of intrinsically disordered proteins. *Proc National Acad Sci* **117**, 28795-28805 (2020). https://doi.org:10.1073/pnas.2008122117 PMID - 33139563

Joseph, J. A. *et al*. Physics-driven coarse-grained model for biomolecular phase separation with near-quantitative accuracy. *Nature Computational Science* **1**, 732-743 (2021). https://doi.org:10.1038/s43588-021-00155-3

Tesei, G., Schulze, T. K., Crehuet, R. & Lindorff-Larsen, K. Accurate model of liquid–liquid phase behavior of intrinsically disordered proteins from optimization of single-chain properties. *Proc National Acad Sci* **118**, e2111696118 (2021). https://doi.org:10.1073/pnas.2111696118

Kwon, I. *et al*. Phosphorylation-Regulated Binding of RNA Polymerase II to Fibrous Polymers of Low-Complexity Domains. *Cell* **155**, 1049 - 1060 (2013). https://doi.org:10.1016/j.cell.2013.10.033

Guo, Y. E. *et al*. Pol II phosphorylation regulates a switch between transcriptional and splicing condensates. *Nature* **572**, 543 - 548 (2019). https://doi.org:10.1038/s41586-019-1464-0 PMID - 31391587

Lyons, H. *et al*. RNA polymerase II partitioning is a shared feature of diverse oncofusion condensates. *Cell* (2025). https://doi.org:10.1016/j.cell.2025.04.002

Ahn, J. H. *et al*. The phenylalanine-and-glycine repeats of NUP98 oncofusions form condensates that selectively partition transcriptional coactivators. *Mol Cell* **85**, 708-725.e709 (2025). https://doi.org:10.1016/j.molcel.2024.12.026

Nott, T. J. *et al*. Phase transition of a disordered nuage protein generates environmentally responsive membraneless organelles. *Mol Cell* **57**, 936 - 947 (2015). https://doi.org:10.1016/j.molcel.2015.01.013 PMID - 25747659

Welsh, T. J. *et al*. Surface Electrostatics Govern the Emulsion Stability of Biomolecular Condensates. *Nano Letters* **22**, 612-621 (2022). https://doi.org:10.1021/acs.nanolett.1c03138

Wang, Y., Lomakin, A., Kanai, S., Alex, R. & Benedek, G. B. Liquid–Liquid Phase Separation in Oligomeric Peptide Solutions. *Langmuir* **33**, 7715-7721 (2017). https://doi.org:10.1021/acs.langmuir.7b01693

Krainer, G. *et al*. Reentrant liquid condensate phase of proteins is stabilized by hydrophobic and non-ionic interactions. *Nat Commun* **12**, 1085 (2021). https://doi.org:10.1038/s41467-021-21181-9 PMID - 33597515

Lin, Y.-H. *et al*. Electrostatics of Salt-Dependent Reentrant Phase Behaviors Highlights Diverse Roles of ATP in Biomolecular Condensates. (2024). https://doi.org:10.7554/elife.100284.2

MacAinsh, M., Dey, S. & Zhou, H.-X. Direct and indirect salt effects on homotypic phase separation. *Elife* **13**, RP100282 (2024). https://doi.org:10.7554/elife.100282

58 Szabó, A. L., Sánta, A., Pancsa, R. & Gáspári, Z. Charged sequence motifs increase
the propensity towards liquid–liquid phase separation. *FEBS Letters* **596**, 1013-1028
(2022). https://doi.org:10.1002/1873-3468.14294
59 Chang, L.-W. *et al.* Sequence and entropy-based control of complex coacervates. *Nat*
Commun **8**, 1273 (2017). https://doi.org:10.1038/s41467-017-01249-1 PMID -
29097695
60 Sing, C. E. & Perry, S. L. Recent progress in the science of complex coacervation. *Soft*
Matter **16**, 2885-2914 (2020). https://doi.org:10.1039/d0sm00001a
61 Zarin, T., Tsai, C. N., Ba, A. N. N. & Moses, A. M. Selection maintains signaling function
of a highly diverged intrinsically disordered region. *Proc National Acad Sci* **114**,
E1450-E1459 (2017). https://doi.org:10.1073/pnas.1614787114
62 Mihalič, F. *et al.* Conservation of Affinity Rather Than Sequence Underlies a Dynamic
Evolution of the Motif-Mediated p53/MDM2 Interaction in Ray-Finned Fishes. *Mol Biol*
Evol **41**, msae018 (2024). https://doi.org:10.1093/molbev/msae018
63 Adachi, K. & Kawaguchi, K. Predicting Heteropolymer Interactions: Demixing and
Hypermixing of Disordered Protein Sequences. *Physical Review X* **14**, 031011 (2024).
https://doi.org:10.1103/physrevx.14.031011
64 Ginell, G. M. *et al.* Sequence-based prediction of intermolecular interactions driven
by disordered regions. *Science* **388**, eadq8381 (2025).
https://doi.org:10.1126/science.adq8381
1076