1

Peptides at vesicle and mineral prebiotic interfaces

Ivan Cherepashuk[1], Mikhail Makarov[1], Radko Soucek[2], Vaclav Verner[2], Robin Krystufek[2], Romana Hadravova[2], Valerio Guido Giacobelli[1,2], Kosuke Fujishima[3], Sean F. Jordan[5*], Liam M. Longo[3,6*], Klara Hlouchova[1,2*]

[1] *Department of Cell Biology, Faculty of Science, Charles University, Prague 12800, Czech Republic*

[2] *Institute of Organic Chemistry and Biochemistry, Czech Academy of Sciences, Prague 16610, Czech Republic*

[3] *Earth-Life Science Institute, Institute of Science Tokyo, Tokyo 1528550, Japan*

[4] *Graduate School of Media and Governance, Keio University, Fujisawa 2520882, Japan*

[5] *Life Sciences Institute, School of Chemical Sciences, Dublin City University, Glasnevin, Dublin, 9, Ireland*

[6] *Blue Marble Space Institute of Science, Seattle, Washington 98104, USA*

*Correspondence to: sean.jordan@dcu.ie, llongo@elsi.jp, havova1@natur.cuni.cz

Keywords: origins of life, peptides, minerals, vesicles, prebiotic compartments, amino acid alphabet evolution

Abstract

The origin of life likely involved a complex interplay between organic molecules and mineral surfaces, yet the molecular details of these interactions remain poorly understood. Over recent decades, considerable research has focused on the individual roles of key biomolecules - such as RNA, lipids, and proteins - in early abiogenesis. However, this reductionist view offers only a partial picture because the emergence of life likely involved networks of molecular interactions that collectively shaped early functional assemblies. In this study, we examine the ability of peptides - arguably one of the most abundant early polymers - to interact with mineral surfaces and lipid vesicles, prebiotic interfaces and compartments. Using peptide libraries constructed from either prebiotically plausible or contemporary amino acids, *we demonstrate that while acidic residues drive peptide binding to mineral surfaces (such as fluorapatite, studied here), the inclusion of arginine - a basic residue that may have been accessible in specific prebiotic environments – synergistically enhances the mobilization of bioavailable phosphate from geological reservoirs*. Furthermore, we observe a functional divergence in vesicle interactions: while prebiotic alphabets

promote dynamic membrane behaviours such as budding, libraries with 'late' canonical amino
acids can help preserve vesicle integrity against salt-induced collapse. Our finding supports the view
that interactions with peptides can elicit changes in both prebiotic minerals and vesicles,
underscoring the importance of studying these systems collectively.

Introduction

Mineral pores, coacervates, and vesicles (*enclosures bound by a lipid bilayer*) are believed to have
been among the earliest compartmentalization structures on Earth. These structures are thought to
have supported the emergence of life by influencing molecular assembly and stability, and serving
as catalysts (Aquino et al., 2004; Biondi et al., 2007; Deamer et al., 2002; Lambert, 2008; Morigaki
and Walde, 2007; Oparin, 1924; Rimola et al., 2013; West et al., 2017; Yu et al., 2013). While
extensive research has explored the physicochemical properties of putative early life
compartments, *including their formation, stability, and growth requirements* (Chen and Szostak,
2004; Cornell et al., 2019; Hentrich and Szostak, 2014), significantly less is known about their
interactions with peptides. Yet, binding of peptides to mineral surfaces and vesicles may have a
profound impact on their stability and catalytic properties. This knowledge gap reflects a long-
standing divide in origin-of-life research, where compartmentalization, peptide chemistry, and
mineral catalysis have often been investigated separately rather than as interconnected processes
(Preiner et al., 2020).

Peptides were likely the most abundant and functionally versatile polymers in prebiotic
environments (Fried et al., 2022). *Multiple studies have demonstrated the prebiotic accessibility of*
simple peptides, and ~15-residue oligopeptides have been produced by a range of prebiotically
relevant processes (Foden et al., 2020; Frenkel-Pinter et al., 2020; Kwiatkowski et al., 2021; Mayer et
al., 2018). Various lines of evidence demonstrate that even short peptides can play functionally
significant roles: For example, di- and tri-peptides have been shown to catalyse the hydrolysis of
proteins, nucleic acids, and esters (Cohen et al., 2024; Cornell et al., 2019; Gorlero et al., 2009).
Amino acids and short peptides are also able to bind to simple fatty acid vesicles, resulting in an
increase in vesicle stability to metal ions and high concentrations of salt (Cornell et al., 2019; Dalai
and Sahai, 2019) and greater lamellarity (Cornell et al., 2019). Likewise, the presence of acidic
residues can promote the binding of peptides onto mineral surfaces (Hoff et al., 2022). Taken
together, these studies suggest that peptides were active participants in the origin of life, even before
the emergence of templated protein synthesis.

Only about half of the canonical amino acids were likely to have been abundant on the early Earth or
producible by a primitive metabolism. These "early canonical" amino acids, inferred from
biosynthetic complexity, atmospheric discharge experiments, meteorite/asteroid analyses, and
hydrothermal vent studies (Cleaves, 2010; Copley et al., 2005; Higgs and Pudritz, 2009; Trifonov,
2000), are listed in **Figure 1** as the 10E set. In addition to the 10E set, a plethora of non-canonical
amino acids have been detected in prebiotic chemistry studies, and it is likely that both canonical
and non-canonical amino acids were incorporated into primitive peptides (Zaia et al., 2008). Recent
work has strongly favoured selection for protein foldability as the driving force for amino acid
incorporation into the genetic code (Giacobelli et al., 2022; Hlouchová, 2024; Longo et al., 2020;
Makarov et al., 2023; Solis, 2019; Tretyachenko et al., 2022). Here, we explore how primitive amino
acid alphabets may have structured (or been structured by) early peptide-mineral and peptide-

vesicle interactions. As the first peptides likely predated templated synthesis and had highly
statistical sequences, we have chosen to employ random-sequence peptide libraries (**Figure 1**),
which can better capture global differences between alphabets.

The presence of phosphate minerals and clays at the origin of life is virtually uncontested. Decades
of research into prebiotic membrane structures suggest that vesicles were also present on the early
Earth (Deamer and Pashley, 1989; Gebicki and Hicks, 1976; Hanczyc et al., 2003; Hargreaves and
Deamer, 1978; Jordan et al., 2019; Monnard and Deamer, 2011). *Membrane-forming molecules,*
such as single-chain amphiphiles, can be readily produced by Fischer-Tropsch chemistry (Cohen et
al., 2023; McCollom et al., 1999), particularly in hydrothermal systems, and these molecules can be
concentrated by association with mineral surfaces (Sahai et al., 2017). Moreover, the minimal fatty
acid concentration required for vesicle formation can be as low as the micromolar range (Jordan et
al., 2019). The observation that organic solvent-solubilized fractions of the Murchison meteorite self-
assemble into vesicles in water (Deamer and Pashley, 1989) provides additional support for the view
that different early Earth conditions could promote vesicle formation.

Here, we synthesize six random octapeptide libraries to explore the potential role of peptide-mineral
and peptide-vesicle interactions early in life's history. We consider the interactions of peptides built
from six different amino acid alphabets – five composed of prebiotically plausible amino acids and
one reflecting a contemporary protein composition – with the ubiquitous phosphate mineral
fluorapatite and simple decanoic acid:decanol (DA:DOH) vesicles. We find that primitive peptide
libraries preferentially interact with the mineral surface and promote release of orthophosphate.
These effects are likely due to an enrichment in acidic amino acids in the primitive amino acid
alphabet. With respect to vesicles, we find that later amino acid alphabets exert slightly stronger
vesicle-stabilizing effects in the presence of mono- and di-cations. We conclude that the bulk
properties of primitive peptides were likely well-suited for mineral interaction.

Materials and Methods

Chemicals. For the solid-phase synthesis of combinatorial peptide libraries, Fmoc-Rink Amide AM resin (100-200 mesh, 0.64 mmol/g loading) and Fmoc-protected amino acids were purchased from Iris Biotech (Germany). N,N'-dimethylformamide (DMF) was purchased from Carl Roth (Germany), dichloromethane and diethyl ether were purchased from PENTA (Czech Republic). Ceramic hydroxyapatite and fluorapatite (type II, 40 µm particle size) were purchased from Bio-Rad (USA). All other chemicals were purchased from Sigma Aldrich (USA). For experiments related to vesicle interactions, Rhodamine 6G, DA were purchased from ThermoFisher Scientific (USA), and DOH was purchased from Sigma Aldrich.

Chemical synthesis of combinatorial peptide libraries. 8-mer combinatorial peptide libraries were prepared manually using Fmoc-based solid-phase synthesis. Synthesis of the peptide mixtures was accomplished by coupling Fmoc-protected amino acids individually to the equal aliquots of the solid support that were merged after coupling (split and mix approach). Prior to synthesis, Fmoc-amino acid/Oxyma Pure stock solutions were prepared by simultaneously dissolving Fmoc-amino acids and Oxyma Pure to the final concentrations of 300 mM and 375 mM in DMF, respectively. 500 mg of Rink Amide AM resin (0.32 mmol) was swelled in 20 ml of DMF for 1 h and subsequently deprotected by incubation with 15 ml of 20 % (v/v) piperidine in DMF for 1 h at room temperature. The resin was washed four times with 10 ml of DMF, thoroughly resuspended in 10 ml of DMF and split into equal aliquots, depending on the number of amino acids (10 for "10E", 11 for "10U", "11R", "11D", "11K", and 19 for "19F" peptide libraries). For the synthesis of "10E" peptide library, 1.1 ml of Fmoc-amino acid/Oxyma Pure solutions and 64 µl of N,N'-diisopropylcarbodiimide were added to the resin aliquots, and the reaction mixture was incubated overnight at room temperature. The amounts of Fmoc-amino acid/Oxyma Pure solution and N,N'-diisopropylcarbodiimide used for coupling were scaled down for a higher number of amino acids (resin aliquots). Following the coupling, the resin aliquots were merged, washed five times with 10 ml of DMF and deprotected by incubation with 15 ml of 20% (v/v) piperidine in DMF for 1 hour at room temperature. The amino acid coupling and deprotection steps were performed eight times. All separation and washing steps were performed using 15 ml polypropylene syringes fitted with polypropylene frits. After the final coupling/deprotection step, the resin was washed five times with 10 ml of dichloromethane and dried *in vacuo*. The peptide mixture was cleaved off the dried resin by incubation with 10 ml of trifluoroacetic acid-triisopropylsilane-water (95:2.5:2.5, v/v) mixture for 2 hours at room temperature. The resin was filtered off, and the peptide mixture was precipitated with 150 ml of diethyl ether cooled down to 4 ℃ and submerged in liquid nitrogen for 10 min to maximize precipitation. The peptide mixture was subsequently centrifuged at 5000× g for 5 min at 4 ℃, and the peptide pellet was collected and lyophilized three times with 1 mM HCl overnight.

Additionally, 19F, 10E, and 11R libraries were synthesized with the inclusion of the TAMRA fluorescent dye for peptide co-localisation assays. Labelling was performed by overnight treatment with 3 eq. of 5(6)-carboxytetramethylrhodamine, 3.75 eq. Oxyma Pure and 3.75 eq. N,N'-diisopropylcarbodiimide in DMF after the final coupling/deprotection step.

Quality control of combinatorial peptide libraries. The molecular weight distributions of 8-mer combinatorial peptide libraries were confirmed by mass spectrometry using UltrafleXtreme MALDI-TOF/TOF mass spectrometer (Bruker Daltonics, Germany) according to the standard procedure. Prior to the MALDI-TOF analysis, peptide libraries were dissolved in acetonitrile:water (50:50, v/v) mixture and mixed with 2,5-dihydroxybenzoic acid (DHB) that was used as a matrix.

The amino acid composition of 8-mer combinatorial peptide libraries was estimated by HPLC amino acid analysis performed on Biochrom 30 system (Biochrom, United Kingdom) using automated ninhydrin derivatization. Prior to the HPLC amino acid analysis, the library samples were hydrolyzed in 6 M HCl at 110 °C for 20 h with the hydrolysate evaporated and reconstituted with 0.1 M HCl.

HPLC amino acid and mass spectrometry analyses of mineral binders from 8-mer combinatorial peptide libraries. 10 mg of fluorapatite was gently rotated with 0.5 ml of 1 mg/ml peptide library solution in HMA buffer (20 mM HEPES, 20 mM MES, 20 mM sodium acetate, pH 5.5) for 21 hours at room temperature. The mixture was centrifuged at 15,000× g for 3 min at room temperature to separate apatite from the peptide library solution, and the supernatant was subsequently discarded. The apatite was resuspended in 0.5 ml of HMA buffer (pH 5.5), incubated for 3 min at room temperature and centrifuged at 15000× g for 3 min. The washing step was repeated four times. The bound peptides were subsequently released from apatite by overnight incubation with 220 μl of 50 mM HCl at room temperature. Following the incubation, the mixture was centrifuged at 15000× g for 3 min, and 150 μl of supernatant was submitted to MALDI-TOF or HPLC amino acid analyses. Mass spectrometry was performed once, an HPLC analysis was performed six times.

Phosphate release by 8-mer combinatorial peptide libraries. Solubilization of fluorapatite by combinatorial peptide libraries at the pH of 5.5 and temperature of 55 ℃ conditions was estimated by molybdenum blue spectrophotometric assay (Murphy and Riley, 1962). Prior to analysis, molybdenum blue reagent was freshly prepared by mixing 2.5 M sulphuric acid, ammonium molybdate (40 g/l), 0.1 M ascorbic acid, and potassium antimonyl tartrate (1 mg Sb/ml) in 10:3:6:1 (v/v) ratio. 10 mg of apatite was gently rotated with 0.5 ml of 1 mg/ml peptide library solution in HMA buffer for 21 hours. The mixture was centrifuged at maximum speed (21.300× g) for 20 min to separate apatite from the peptide library solution, and 300 μl of supernatant was transferred to a new tube. The supernatant was centrifuged again at maximum speed (21.300× g) for 20 min to remove any residual apatite, and 250 μl of supernatant was transferred to a new tube. 5 μl of supernatant was mixed with MilliQ water to adjust the volume to 150 μl in a 96-well plate, and 30 μl of molybdenum blue reagent was added to each well. The reaction mixture was incubated for 20 min at room temperature, and the absorbance at 882 nm was measured using Infinite M Plex plate reader (TECAN, Switzerland). The amount of phosphate ions in the supernatant was quantified from the standard curve established using 2-fold serial dilutions of KH_2PO_4 solution in MilliQ water. The measurement was performed in technical triplicates. The analysis was performed three times.

Preparation of stock decanoic acid:decanol (DA:DOH) vesicle solution. Decanoic acid (DA) and
decanol (DOH) offer relative stability for vesicle experimentation and were observed in meteorite
materials [6]. All solutions were pre-warmed to 50 ℃ using a water bath. DA was dissolved in 340
volumes of acetate-phosphate-borate buffer (ABP, pH 7.4) or 5 mg/mL peptide library solution in ABP
(in experiments where peptide libraries were present in the lipid solution prior vesicle formation).
The mixture was mixed vigorously for 10 seconds, followed by the addition of 180 mM NaOH to
achieve a DA concentration of 53 mM. DOH was then added in an equimolar concentration to DA,
and the solution was mixed vigorously for 60 seconds to allow vesicle formation. *The pH was*
adjusted to 7.4 using HCl, resulting in a final concentration of 37.8 mM of DA and 38.4 mM of DOH.
The stock solution was then used in experiments downstream.

Preparation of peptide library solution for experimentation on DA:DOH vesicles. 5 mg of peptide
library was dissolved in ABP to a final concentration of 5 mg/ml. This solution, freshly upon mixing by
pipetting, was either added to DA (in experiments where peptide libraries were present in the lipid
solution prior vesicle formation) or diluted with ABP to a concentration of 2 mg/ml. The stock
solutions were used in consequent experiments shortly upon mixing by pipetting to ensure uniform
peptide concentration.

Amino acid analysis of peptide libraries upon incubation with DA:DOH vesicles. 150 µl of the
DA:DOH vesicle stock solution was mixed with 150 µl of 2 mg/ml solution of peptide libraries in ABP
("vesicle incubation" condition) or ABP alone (negative control) and incubated at 25 ℃ with shaking
at 1000 RPM, 2 mm orbit for 1 hour. Additionally, peptide libraries were incubated in the absence
of vesicle stock solution (replaced with ABP) in parallel ("vesicle control" condition in order to
elucidate peptide library binding to the Amicon centrifugal filter itself). 10 µl of the resulting solution
was dried out and submitted to amino acid analysis ("unfiltered" condition), and another 250 µl of
the were loaded into Amicon Ultra-4 3K centrifuge filters and centrifuged at 3000× g for 10
minutes. 10 µl of the flow-through from the Amicon centrifuge filters were also dried out and
submitted to amino acid analysis ("filtered" condition). Amino acid analysis was performed on a
Biochrom 30 amino acid analyzer. Results from the negative control were subtracted from the
results, and both relative concentrations of individual amino acids and whole peptide library
concentration were compared relatively within each condition. All experiments were conducted in 6
replicates.

Zeta-potential measurements. Measurements of zeta-potential of vesicle populations were
conducted using Zetasizer Nano ZS90 (Malvern Panalytical) with DTS1070 cuvettes at standard
manufacturer settings at 25 ℃. 350 µl of vesicle stock were mixed with 350 µl of either ABP or 2 mg/ml
peptide library solution in ABP, and 70 µl of 4M NaCl or 100 mM $MgCl_2$ solutions were also added,
depending on the experiment, to a final concentration of 363 and 8.9 mM, respectively. Experiments
were conducted in technical triplicates.

Dynamic light scattering. Dynamic light scattering measurements were conducted using Zetasizer
Nano ZS90 (Malvern Panalytical) with ZEN2112 cuvettes at 25 ℃. For this, 50 μl of vesicle-peptide
solution was mixed with 50 μl of ABP, and 10 μl of 4M NaCl or 100 mM $MgCl_2$ solutions were also
added, depending on the experiment, to a final concentration of 363 and 8.9 mM, respectively. Side
scattering measurements were then analysed to determine the size distribution of the samples. The
experiment was conducted in technical triplicates.

Spinning-disc confocal microscopy. 50 μl of stock vesicle solutions were mixed in a 1:1 ratio with 2
mg/mL solutions of peptide libraries in ABP buffer on Glass Bottom 18-Well μ-Slides (Ibidi,
Germany). Rhodamine 6G (excitation peak at 561 nm) was added to a final concentration of 1 μM.
Depending on the experiment, either 10 μl of 4M NaCl solution was added to a final concentration of
363 mM or 10 μl of 100 mM $MgCl_2$ solution to a final concentration of 9.1 mM. In alternative setups,
2 mg/ml of peptide library solution in ABP conjugated with TAMRA fluorescent dye (excitation at 561
nm) were mixed with their non-fluorescent counterparts (2 mg/ml in ABP) in a 1:50 ratio. In such
instances, DiD fluorescent dye (excitation at 638 nm) was used for membrane imaging instead of
Rhodamine 6G, which has an overlapping excitation range with TAMRA. Imaging was performed
using a Nikon CSU-W1 spinning-disc confocal microscope with a 100× oil-immersion objective and
a PRIME BSI camera (Teledyne Photometrics) at room temperature. Experiments were conducted
three times.

Transmission electron microscopy. 100 μl of DA:DOH vesicle solution was diluted in 900 μl PBS
buffer to a final concentration of 7.7 mM. Vesicle sample was then analysed in PBS buffer by
negative staining. Parlodion-carbon-coated grids were floated on the top of 10 μl drop of the sample
for 5 min. Then the grids were stained of 2% uranyl acetate, for 2 × 30 s and dried. The grids were
examined with a JEOL JEM-2100 Plus electron microscope operated at 200 kV (JEOL Ltd., Japan).

Turbidity measurement. Turbidity measurements were conducted in 96-well plates on a microplate
reader (Infinite M Plex, Tecan, Switzerland). Vesicle solutions were prepared in presence of peptide
libraries with free fatty acids as outlined above. 10 μl of 4 M NaCl or 10 μl of 100 mM $MgCl_2$ solutions
were added to final concentrations of 363 mM or 9.1 mM, respectively. OD_{490} measurement was then
conducted in triplicates, and absorbance was compared against average vesicle sample
absorbance at the absence of peptide libraries, which was put at 100% for every salt addition
condition screened (control (no salts added), 363 mM NaCl or 9.1 mM $MgCl_2$).

Critical vesicle concentration measurement. Critical vesicle concentration (CVC) was established
by preparing serial dilution of vesicle stock solution with or without peptide libraries (from 0.35 to
11.33 mM, 100 μl final sample volume in a 96-well plate), adding 2 μL of pyrene dye (pure, 1 mM stock
solution in methanol) to final concentration of 5 μM and measuring fluorescence emission at 384
and 373 nm using Infinite M Plex microplate reader (Tecan, Switzerland) at 25 ℃. The 384/373 ratio

curve was then plotted against vesicle concentration and the approximation of CVC was evaluated
at the 1.15 384/373 ratio cutoff point. The experiment was conducted in triplicates.

Statistical analysis. All data were analyzed using Python (version 3.11) with numpy, pandas, seaborn,
scipy, and matplotlib libraries. Normality of distributions was assessed using the Shapiro–Wilk test.
Pairwise comparisons between the Control group and peptide libraries were performed using the
non-parametric Mann–Whitney U test. To account for multiple testing, Holm-Bonferroni correction
was applied to adjust the p-values. Differences were considered statistically significant at $p < 0.05$
after correction. For CVC analysis, *group differences were assessed using a Kruskal–Wallis test*
followed by Holm-corrected Mann–Whitney post hoc comparisons. No statistically significant
differences were detected (p = 0.10).

Results and Discussion

Figure 1. Amino acid composition of the random octapeptide libraries. Canonical amino acids are indicated using single letter codes. Blue hexagons denote prebiotically available amino acids (Higgs and Pudritz, 2009); orange hexagons denote "late" amino acids that require more elaborate biosynthesis; green hexagons denote prebiotically available amino acids that were not incorporated into the genetic code (non-canonical amino acids). DAB, Nva, Nle, ABA stand for 2,4-diaminobutyric acid, norvaline, norleucine, and α-aminobutyric acid, respectively.

Peptide libraries: properties and motivations

Six different octapeptide libraries with fully random sequences were synthesized (**Figure 1**). Amino acids were selected based on their prebiotic plausibility (Higgs and Pudritz, 2009) and biophysical properties (Makarov et al., 2023): The 10E library consists of the ten early canonical amino acids (Longo et al., 2013). The 10U library replaces the three canonical branched amino acids (Val, Lue and Ile) with three prebiotically available, unbranched amino acids (norvaline, norleucine, and alpha-aminobutyric acid). Thus, by comparing the 10E and 10U libraries, the effect of amino acid branching, and the reduction in side-chain entropy it entails, can be assessed.

Both the 10E and 10U libraries are devoid of positively charged amino acids, making the N-terminus the only positively charged group on these peptides (McDonald and Storrie-Lombardi, 2010). *To study the role of basic amino acids, three alphabets based on the early canonicals (10E), with the addition of a single basic amino acid type, were constructed: 11D (addition of 2,4-diaminobutyric acid or DAB), 11R (addition of Arg), and 11K (additional of Lys).* DAB was likely present in prebiotic environments (Zaia et al., 2008) and is hypothesized to have been the first coded basic amino acid, present during an intermediate stage of the genetic code (Copley et al., 2005). Arg, although not typically detected in meteorites or atmospheric discharge experiments, may have been accessible via a cyanosulfidic protometabolism (Patel et al., 2015). In addition, conversion of prebiotically available ornithine into Arg by guanidination has been demonstrated, both for the free amino acid

and for ornithine-containing prebiotic peptides (Ariely et al., 1966; Longo et al., 2020). Lys is
hypothesized to be the most recent basic amino acid addition to the genetic code (Trifonov, 2000).

Finally, as a point of comparison to the primarily prebiotic alphabets above, the 19F library – which
includes all canonical amino acids except Cys – was constructed. The exclusion of Cys was due to
limitations of Fmoc-based peptide synthesis. Unlike the other libraries, 19F includes aromatic
amino acids (Phe, Tyr and Trp) and amide-containing side chains (Asn, Gln). Note also the presence
of His, which is often involved in metal binding (Li et al., 2024).

In each case, the amino acid types are approximately uniformly distributed within a peptide library
(**Figure S1**) and not biased by either prebiotic availability or frequency in proteins. Given the large
gaps in our understanding of primitive peptide synthesis, this decision is meant to retain as much
neutrality as possible. Assuming a perfectly uniform distribution of amino acids, the properties of
the bulk library can be estimated (Makarov et al., 2023): *the order of hydrophobicity, from highest to*
lowest, is 19F > 11K > 11R > 11D > 10U > 10E. The fraction of charged residues, from highest to lowest,
is 11R (0.27) = 11K (0.27) = 11D (0.27) > 19F (0.26) > 10E (0.2) = 10U (0.2). Finally, the estimated
median pI of each library is 10E (5.46) < 10U (5.49) < 11K (5.84) < 11D (5.85) < 11R (5.94) < 19F (6.13).

Peptide-vesicle and peptide-mineral interactions are influenced by library composition

First, we measured the extent to which each of the octapeptide libraries interacts with mineral and
vesicle surfaces. The mineral fluorapatite was selected due to its abundance and prebiotic
availability, as well as its potential role as a prebiotic source of phosphate (Paytan and McLaughlin,
2007). Fluorapatite is more stable than hydroxyapatite and found on both Earth and Mars (Hausrath
and Tschauner, 2013; Martínez et al., 2023). The n-decanoic acid and n-decanol (DA:DOH) vesicle
system was selected due its putative prebiotic availability (Cohen et al., 2024; Cornell et al., 2019).
*Peptide binding (**Figure 2A**) was measured after co-incubation with mineral powder or vesicles.*
Filtration (vesicles) and centrifugation (mineral grains) were used to separate the prebiotic substrate-
bound peptides from the bulk solution. Finally, peptides samples were subjected to amino acid
*analysis (see **Methodology**).*

Every peptide library screened exhibited some fluorapatite binding, with 5-20% of peptide molecules
retained on the surface, as measured by an experiment in which bound peptides were desorbed from
*the surface of the mineral (**Figure 2A** and **2B**, left). We find that the peptide libraries based on 10E*
*(10E, 10U, 11D, 11K, 11R) bind more strongly to fluorapatite than the 19F library (**Figure 2B**, left). In*
addition, amino acid analysis reveals that Asp and Glu residues were enriched in fluorapatite-bound
*peptides relative to the full library in each case (**Figure S2**).* These results are consistent with the
known affinity of acidic residues for apatite, as acidic side chains can coordinate calcium ions (Hoff
et al., 2022; Tavafoghi and Cerruti, 2016).

We note that several factors may influence the binding affinity of the peptide libraries to fluorapatite.
Foremost is pH: Our experiments were performed at pH 5.5, where the surface of fluorapatite is

expected to be positively charged based on a point of zero charge (PZC) of 6.8 (Bell et al., 1973). In
practice, however, the fluorapatite interfacial charge tends to be more negative than that predicted
by PZC (Chaïrat et al., 2007). By changing the pH, it may be possible to tune binding; however, as
long as Asp and Glu can coordinate calcium, we expect acidic amino acids to be the primary
mediators of binding. Second*, we note that higher sintering temperatures yield apatite samples with*
weaker peptide binding propensity (Jeyapalina et al., 2022). The fluorapatite used in our experiments
was sintered at a temperature of 700 °C (Trzaskowska et al., 2023).

*Next, we evaluated the binding of the peptide libraries to DA:DOH vesicles (**Figure 2B**, right). We first*
confirmed the presence of vesicles using transmission electron microscopy and fluorescent imaging
*(**Figure S3**). After incubation, Amicon® Ultra-4 3K centrifugal filters, which use a regenerated*
*cellulose membrane, were used to separate the vesicle-bound and unbound peptides (**Figure 2A**),*
as has been done previously (Black et al., 2013; Cornell et al., 2019). Although we observed
considerable non-specific peptide binding to the filter membrane itself, comparisons with vesicle-
*free controls demonstrated that this effect was uniform across all libraries (**Figure S4**).* Because this
background loss represents a systematic bias rather than a variable one, we report our findings as
relative comparisons of vesicle affinity.

We find that the 10E and 10U libraries have an 80% higher peptide concentration in the unbound
phase relative to 19F than libraries with positively charged amino acids (11D, 11R, and 11K). This
result suggests that positively charged amino acids play a role in peptide-vesicle association, in
agreement with previous reports on basic amino acid behaviour (Cohen et al., 2024; Kamat et al.,
2015; Kwiatkowski et al., 2021; Mitchell et al., 2000). Interestingly, the 11D library (containing DAB)
displayed a similar degree of vesicle binding to the 11R (containing Arg) and 11K (containing Lys)
libraries. This result suggests that fatty acid vesicle binding may not have exerted a strong selective
force on the amino acid alphabet, consistent with our earlier interpretation that foldability was the
primary property under selection. Moreover, we do not find substantial differences in the amino acid
composition of peptides filtered from the vesicle solution relative to the total library (**Figure S4**).

Figure 2. Peptide library association with prebiotic surfaces. A. Outline of the experimental pipeline. Since it is not feasible to centrifuge fatty acids down and alcohols interfere with absorbance at 220 nm, HPLC was used to establish the reduction in peptide concentration upon binding for both setups. B. Library binding to fluorapatite (left) or DA:DOH vesicles (right) calculated from amino acid analysis. Briefly, the amount of peptide bound to fluorapatite was estimated to be the sum of individual amino acid molar amounts in the sample. The same approach was taken in the vesicle-peptide assay but, due to limitations of the filtration-based assay, only relative concentrations are reported. Whiskers indicate the minimum and maximum values, boxes span the second to the third quartile, and the median value is indicated with a line. Statistical significance calculated using the Holm-Bonferroni-corrected U-test is indicated by asterisks: $p < 0.05$ (*), $p < 0.01$ (**), and $p < 0.001$ (***).

Peptide libraries promote phosphate release from fluorapatite

As phosphate is essential for energy conservation, information storage, and encapsulation, we determined whether the peptide libraries could promote the release of orthophosphate from fluorapatite. To detect released orthophosphate, we employed the molybdenum blue assay (**Figure 3B**; see **Figure S5** for a representative standard curve). We find that all screened libraries induced orthophosphate release from fluorapatite, increasing the concentration of soluble orthophosphate by up to a factor of two relative to the mineral alone (**Figure 3C**). *Notably, this effect appears to be composition-dependent. While all libraries promoted dissolution, those containing Arg (the **11R** and **19F** libraries) consistently elicited the highest levels of phosphate release. We propose a cooperative mechanism driving this efficiency. All libraries contain Asp and Glu, which likely coordinate calcium*

ions, destabilizing the surface lattice via calcium leaching, thereby shifting the equilibrium toward
dissolution (Matreux et al., 2025). However, the inclusion of Arg appears to synergize with this effect.
The guanidinium group of Arg is unique in its ability to form stable, bidentate hydrogen bonds with
phosphate and phosphonate groups (Woods and Ferré, 2005). This highlights a potential advantage
of the 'modern' basic amino acid Arg over simpler prebiotic basic amino acids in accessing
geochemical phosphate reservoirs.

Figure 3. A. Chemical structure of fluorapatite $Ca_{10}(PO_4)_6F_2$ (Hoff et al., 2022; Posner, 1985). B. Experimental workflow for
detecting orthophosphate release from fluorapatite. C. Orthophosphate release from fluorapatite. The "Control" sample
*lacks any peptides libraries, and incubation was performed in buffer (see **Methodology**). All peptide libraries except for 11K*
promoted significant orthophosphate release relative to the buffer control (p < 0.05 for all 5 comparisons to the control, as
*assessed by the Holm-Bonferroni-corrected U-test). Box and whisker plot as in **Figure 2**.*

Peptide libraries affect vesicle stability in high-salt conditions

We next determined whether there is a composition-dependent impact of peptide libraries on
vesicle structure, stability, or aggregation. We address these questions using optical density at 490
nm (OD_{490}; a measure of aggregation and lamellarity of fatty acid vesicles (Black et al., 2013; Cornell
et al., 2019; Wang et al., 2019)), spinning-disc fluorescent microscopy, and dynamic light scattering
(DLS) of vesicles created in the presence of peptide library solution. We find that vesicle morphology,
OD_{490}, and average particle size are largely unaffected by any of the six peptide libraries under no-
salt conditions (**Table S1**). However, upon addition of 363 mM NaCl or 8.9 mM $MgCl_2$, significant
differences were observed.

We find a large increase in OD_{490} upon addition of $MgCl_2$, if peptide libraries were present in the
sample (**Figure 4A**). *This effect can be partially explained by the formation of punctate structures*
*(Cornell et al., 2019) and/or multilamellar vesicles, depending on the peptide library used (**Figure 4B**,*
images O-U). We note, however, that the formation of $Mg(OH)_2$ can also result in an increase in
optical density. The destabilization of vesicles in the presence of both peptide libraries and divalent
cations may originate from increased osmolarity or disruption of hydrogen bonding networks. These

effects seem to be negated in part by the late canonical amino acids present in the 19F peptide
library.

Upon addition of NaCl to peptide-vesicle samples, we observe formation of filamentous structures
that resemble growing and budding fatty acid membranes (Hentrich and Szostak, 2014) (**Figure S6**)
in all but the 19F library. These filamentous structures are especially pronounced in the presence of
the 11R, 11D, and 10U libraries (**Figure 4B**, images J-K). Curiously, these libraries also show the
greatest disruption in vesicle formation upon addition of MgCl$_2$ (**Figure 4B**, images P, Q, and S). The
19F library, despite not showing definitive protective effects against flocculation, seems to preserve
vesicle integrity in the presence of MgCl$_2$, as indicated by microscopically observable membranes
*(**Figure 4B**, images O and U) and an increase in OD$_{490}$ compared to control conditions (which may*
*report on the degree of lamellarity or the formation of Mg(OH)$_2$; **Figure 4A**).*

As surface charge influences vesicle-peptide interactions (Chen and Szostak, 2004), we calculated
the ζ-potential of DA:OH vesicles in the presence and absence of salt. We find that 38.5 mM DA:DOH
vesicles have a ζ-potential of -61.5±4.2 mV, which drops to -17.6±9.2 mV and -31±0.4 mV in the
presence of 363 mM NaCl or 8.9 mM MgCl$_2$, respectively (**Table S1**). This reduction in ζ-potential
likely contributes to vesicle flocculation because vesicle surface charge is screened by the presence
of ions, decreasing vesicle-vesicle repulsion. Addition of peptide libraries only slightly reduces the
ζ-potential, with the greatest reduction observed for the 10U library, yielding a ζ-potential of -
49.9±0.9 mV. Thus, a change in ζ-potential upon peptide binding does not seem to mediate the effect
of the peptide libraries on vesicle structure and stability.

To test whether the peptide libraries alter membrane formation propensity, we measured the critical
vesicle concentration (CVC) using pyrene fluorescence. Peptide libraries caused only a minor, non-
significant (p = 0.10) increase in CVC from ~1 mM to ~1.2-1.4 mM of DA:DOH (calculated as a sum of
*molar concentrations of DA and DOH) or had no measurable impact on CVC (**Table S3**). Thus,*
changes in vesicle stability cannot be explained by an increase in CVC.

Taken together, our results suggest that the 19F library is the most beneficial for vesicle stability in
the presence of salts. This library is the most hydrophobic and least soluble (Makarov et al., 2023) of
the tested libraries, while also having the most basic isoelectric point. The extent to which the
benefits of 19F are due to general properties (hydrophobicity, charge) or specific interaction modes
with one or several of the late amino acids is unclear. If vesicle stabilization is specific to the
properties of the late amino acids, however, it may suggest a potential benefit to expanding the
genetic code, though whether this would be to stabilize membranes or augment the interaction of
peptides with membranes is similarly unclear. In addition, we find consistent differences between
the 10E and 10U libraries, with 10U having indications of stronger vesicle binding. This difference
may be due to the longer length of the unbranched sidechains of norvaline and norleucine relative to
their branched counterparts, facilitating fatty acid packing and/or membrane penetration. *We*
hypothesize that peptides with positively charged and unbranched amino acids may have promoted
lipid packing or cross-linking of lipid surfaces within primitive mixtures.

Figure 4. A. Optical density at 490 nm for vesicles-peptide library samples ("10E", "10U", "11D", "11R", "11K" and "19F").
Note that peptide libraries were added to free fatty acids before vesicle formation. Three conditions were tested: no salts
added, NaCl (363 mM final) and MgCl₂ (8.9 mM final). Box and whisker plot as in Figure 2. B. Micrographs of DA:DOH (1:1)
vesicles under different conditions. Vesicular flocculation can be observed upon addition of NaCl. A reduction in the
number of visible particles occurs upon addition of MgCl₂. Staining with Rhodamine 6G. Scale bar: 5 µm; Red arrow -
unilamellar vesicle; yellow arrow - possible multilamellar vesicle; blue arrow - punctate structure; green arrow –
flocculation, pink arrow - growing membranes.

Addition of MgCl₂ or NaCl enhances peptide library co-localisation with DA:DOH vesicles

Surface charge is of importance to the vesicle-peptide interactions and can mediate peptide binding
to membrane surfaces (Chen and Szostak, 2004). We used ζ-potential measurements to assess the
correlation between the vesicle behaviour and their relative charge. We find that 38.5 mM DA:DOH
vesicles have a ζ-potential of -61.5±4.2 mV. In the presence of 363 mM NaCl or 8.9 mM MgCl₂, the ζ
-potential drops to -17.6±9.2 mV and -31±0.4 mV, respectively (**Table S1**). This reduction in ζ -
potential likely contributes to vesicle flocculation because vesicle surface charge is negated by the
presence of ions, decreasing vesicle-vesicle repulsion. Addition of peptide libraries only slightly
reduces the ζ -potential,with the greatest reduction observed for the 10U library, yeilding a value of

-49.9±0.9 mV. We argue that the ζ-potential might not be a decisive factor mediating the effect of
the peptide libraries on vesicles.

To visualize peptide-vesicle co-localisation, we labelled peptide libraries with only canonical amino
acids (19F, 11R, and 10E) with *5-carboxytetramethylrhodamine* (TAMRA) and imaged them using
spinning-disc microscopy alongside *1,1'-dioctadecyl-3,3,3',3'-tetramethylindodicarbocyanine*
(DiD)-stained vesicles. We observed that even peptide libraries with weaker vesicle association in
the filtration assays exhibited marked co-localization with vesicle membranes upon the addition of
NaCl and Mg^{2+} (**Figure 5**), possibly due to the reduction in the ζ-potential of the vesicles. We propose
that the neutralisation of repelling forces on both membranes and peptides by NaCl and Mg^{2+}
increases co-localisation of peptides and fatty acids, and that Mg^{2+} can also act as a "bridging" agent
between these molecules. We argue that this co-localisation of peptide libraries indicates potential
association, binding, or even membrane integration, as recently reported by molecular dynamics
simulation (Cohen et al., 2024).

Notably, we find visual co-localisation of peptide libraries with DA:DOH membranes (but not within
the vesicle lumen) only if peptide libraries were added to free fatty acids prior vesicle formation.
When added to pre-formed vesicles, no observable co-localisation occurred between DA:DOH
vesicles and 10E and 10R libraries. We suggest that peptide libraries integrated into the forming
membranes and potentially modulate membrane formation, as indicated by the relative changes in
OD_{490} in earlier experiments. In contrast, peptides added to vesicle post-assembly only weakly
associated with the vesicle surface.

DA:DOH vesicles were reported to allow the passage of large organic molecules such as ribose and
5'-imidazole-activated dAMP through the membrane, measureable *on the timescale of hours* (Budin
and Szostak, 2011). Despite this result, we did not find substantial accumulation of fluorescently
labelled peptides in the vesicle lumen if peptides were added to pre-formed vesicles (**Figure 5**). We
hypothesise that, due to inclusion of basic amino acids, 19F and 11R libraries could potentially
permeate through the fatty acid membranes, but, due to the short time period in which the
measurements were conducted upon the formation of vesicles (~5-10 minutes), the membrane
permeability should not have been substantial enough to observe.

Figure 5. Fluorescence microscopy images of vesicles with peptide libraries either after (columns A-C, G-I, M-O) or before (columns D-F, J-L, P-R) vesicle formationunder three conditions: Control (top row), 363 mM NaCl (middle row), and 9.1 mM MgCl$_2$ (bottom row). Peptide libraries (19F, 11R, and 10E) are stained with TAMRA (red channel), while lipids are stained with DiD (blue channel). Addition of NaCl (middle row) induces vesicle aggregation and peptide co-localisation with the membranes (columns G-I), with similar, though less pronounced effects upon addition of MgCl$_2$ (columns M-O). Peptide libraries present in the solution before vesicle formation (columns D-F, J-L, P-R) consistently exhibit more stable co-localisation with vesicular membranes across conditions. Scale bars represent 5 µm.

Limitations of the study

Our study is subject to several limitations. First, the libraries used here are a uniform mix of random peptides. As such, individual sequences with exceptional properties are not accessible to the experiments conducted in this work. This feature explains why the effects of the studied libraries are often consistent with the properties of individual constituent amino acids. Moreover, because our libraries contain a mixture of peptides, positive synergistic effects (that enhance the properties of the bulk library) as well as quenching effects (that reduce the properties of the bulk library) are possible. In the latter case, for example, interaction between positively and negatively charged peptides may compete with interaction of positive charged peptides with vesicles.

Both mineral surfaces and fatty acid vesicles are complex systems, and caution must be taken when interpreting experimental results. Fluorapatite binding and orthophosphate relase, for example, are likely influenced by the surface heterogeneity of the sample and the amount of impurity within the mineral. Geologically relevant sample flow, as achieved in flow chambers or similar setups, likely also impacts the properties of mineral-peptide systems (Matreux et al., 2025). Vesicles, on the other hand, are extremely sensitive to sample handling and environmental conditions. We find that minor changes in the pH of the final solution or the duration of sample shaking can lead to observable changes in vesicle stability (**Figure S7**). As such, quantitative descriptions of vesicle stability and morphology should be approached with caution. To mitigate these effects, the same vesicle stock solution was used for every condition screened here.

523 Lastly, our study exclusively explored only a single variant of prebiotically plausible vesicles and
524 minerals as models for potential prebiotic interfaces. A great variety of possibilities exist for prebiotic
525 peptide-interface interactions depending on the model mineral or membrane forming molecule(s)
526 used.

Conclusions

As plausible predecessors of proteins, peptides likely bridged the gap between prebiotic
geochemistry and the emergence of biochemistry. In this study, we systematically explored how
peptide composition influences interactions with two critical prebiotic interfaces: mineral surfaces
and lipid vesicles.

We find that the 'early', prebiotically available amino acid repertoire was sufficient to mediate
interactions with *fluorapatite* and promote phosphate release. While early peptide compositions
promoted membrane dynamics (such as budding) in NaCl, they rendered vesicles susceptible to
salt-induced instability. Libraries containing later amino acids, however, seemed to promote vesicle
stability in the presence of salts.

Acknowledgements

This work was supported by a Research Grant from the Human Frontiers Science Program (HFSP) to
KH and KF (https://doi.org/10.52044/HFSP.RGEC272023.pc.gr.168579). MM and IC acknowledge
support from Grant Schemes at Charles University (CZ.02.2.69/0.0/0.0/19_073/0016935, project no.
START/SCI/148). KH, MM, and IC acknowledge the Biophysical techniques Core Facility of the Czech
Infrastructure for Integrative Structural Biology Instruct-CZ Centre, supported by Ministry of
Education, Youth and Sports of Czech Republic (MEYS CR) Infrastructure project LM2018127 and
European Regional Development Fund-Project, "UP CIISB" (No. CZ.02.1.01/0.0/0.0/18_046/
0015974). LML acknowledges support from HFSP (RGEC29/2025) and the NASA (80NSSC25K7873).
SFJ acknowledges support from the European Research Council (ERC) under the European Union's
Horizon Europe research and innovation programme (grant agreement No 1101114969) and the
Science Foundation Ireland (now Research Ireland) (SFI Pathway award 22/PATH-S/10692).

Authorship confirmation/contribution statement

Ivan Cherepashuk: investigation, validation, formal analysis, writing - original draft, review & editing,
visualisation (lead); conceptualisation, methodology, funding acquisition (equal).

Mikhail Makarov: conceptualisation, investigation, methodology, funding acquisition (equal).

Radko Soucek: investigation, resources, methodology (equal); writing - original draft, review &
editing (equal).

Václav Verner: investigation, resources, methodology (supporting); writing - original draft, review &
editing (supporting).

Robin Krystufek: resources, investigation (supporting).

Romana Hadravova: resources, investigation (supporting).

Valerio Guido Giacobelli: supervision, conceptualisation, project administration, writing - original
draft, review & editing, visualisation (supporting).

Kosuke Fujishima: supervision, conceptualisation, project administration, funding acquisition,
writing - review & editing (equal).

Sean F. Jordan: supervision, conceptualisation, methodology, writing - review & editing (equal).

Liam M. Longo: supervision, conceptualisation, methodology, writing - review & editing (lead).

Klara Hlouchova: supervision, conceptualisation, project administration, funding acquisition,
writing - review & editing (lead).

Authors' disclosure statement

The authors declare that there are no conflicts of interest related to this work.

References

Aquino AJA, Tunega D, Gerzabek MH, et al. Modeling Catalytic Effects of Clay Mineral Surfaces on Peptide Bond Formation. Journal of Physical Chemistry B 2004;108(28):10120–10130; doi: 10.1021/jp049149d.

Ariely S, Wilchek M and Patchornik A. Synthesis of poly-L-arginine. Biopolymers 1966;4(1):91–96; doi: 10.1002/bip.1966.360040110.

Bell LC, Posner AM and Quirk JP. The Point of Zero Charge of Hydroxyapatite and Fluorapatite in Aqueous Solutions. Journal of Colloid and Interface Science 1973;42(2):250–261; doi: 10.1016/0021-9797(73)90288-9.

Biondi E, Branciamore S, Maurel MC, et al. Montmorillonite Protection of an UV-Irradiated Hairpin Ribozyme: Evolution of the RNA World in a Mineral Environment. BMC Evolutionary Biology 2007;7(SUPPL. 2):1–7; doi: 10.1186/1471-2148-7-S2-S2.

Black RA, Blosser MC, Stottrup BL, et al. Nucleobases Bind to and Stabilize Aggregates of a Prebiotic Amphiphile, Providing a Viable Mechanism for the Emergence of Protocells. Proceedings of the National Academy of Sciences of the United States of America 2013;110(33):13272–13276; doi: 10.1073/pnas.1300963110.

Budin I and Szostak JW. Physical Effects Underlying the Transition from Primitive to Modern Cell Membranes. Proceedings of the National Academy of Sciences of the United States of America 2011;108(13):5249–5254; doi: 10.1073/pnas.1100498108.

Chaïrat C, Oelkers EH, Schott J, et al. Fluorapatite Surface Composition in Aqueous Solution Deduced from Potentiometric, Electrokinetic, and Solubility Measurements, and Spectroscopic Observations. Geochimica et Cosmochimica Acta 2007;71(24):5888–5900; doi: 10.1016/j.gca.2007.09.026.

Chen IA and Szostak JW. A Kinetic Study of the Growth of Fatty Acid Vesicles. Biophysical Journal 2004;87(2):988–998; doi: 10.1529/biophysj.104.039875.

Cleaves HJ. The Origin of the Biologically Coded Amino Acids. Journal of Theoretical Biology 2010;263(4):490–498; doi: 10.1016/j.jtbi.2009.12.014.

Cohen ZR, Todd ZR, Maibaum L, et al. Stabilization of Prebiotic Vesicles by Peptides Depends on Sequence and Chirality: A Mechanism for Selection of Protocell-Associated Peptides. Langmuir 2024;40(17):8971–8980; doi: 10.1021/acs.langmuir.4c00150.

Cohen ZR, Todd ZR, Wogan N, et al. Plausible Sources of Membrane-Forming Fatty Acids on the Early Earth: A Review of the Literature and an Estimation of Amounts. ACS Earth Space Chem 2023;7(1):11–27; doi: 10.1021/acsearthspacechem.2c00168.

Copley SD, Smith E and Morowitz HJ. A Mechanism for the Association of Amino Acids with Their Codons and the Origin of the Genetic Code. Proc Natl Acad Sci USA 2005;102(12):4442–4447; doi: 10.1073/pnas.0501049102.

Cornell CE, Black RA, Xue M, et al. Prebiotic Amino Acids Bind to and Stabilize Prebiotic Fatty Acid
Membranes. Proceedings of the National Academy of Sciences of the United States of America
2019;116(35):17239–17244; doi: 10.1073/pnas.1900275116.

Dalai P and Sahai N. Mineral–Lipid Interactions in the Origins of Life. Trends in Biochemical
Sciences 2019;44(4):331–341; doi: 10.1016/j.tibs.2018.11.009.

Deamer D, Dworkin JP, Sandford SA, et al. The First Cell Membranes. Astrobiology 2002;2(4):371–
381; doi: 10.1089/153110702762470482.

Deamer DW and Pashley RM. Amphiphilic Components of the Murchison Carbonaceous
Chondrite: Surface Properties and Membrane Formation. Origins of Life and Evolution of the
Biosphere 1989;19(1):21–38; doi: 10.1007/BF01808285.

Foden CS, Islam S, Fernández-García C, et al. Prebiotic Synthesis of Cysteine Peptides That
Catalyze Peptide Ligation in Neutral Water. Science 2020;370(6518):865–869; doi:
10.1126/science.abd5680.

Frenkel-Pinter M, Frenkel-Pinter M, Samanta M, et al. Prebiotic Peptides: Molecular Hubs in the
Origin of Life. Chemical Reviews 2020;120(11):4707–4765; doi: 10.1021/acs.chemrev.9b00664.

Fried SD, Fujishima K, Makarov M, et al. Peptides before and during the Nucleotide World: An
Origins Story Emphasizing Cooperation between Proteins and Nucleic Acids. Journal of the Royal
Society Interface 2022;19(187); doi: 10.1098/rsif.2021.0641.

Gebicki JM and Hicks M. Preparation and Properties of Vesicles Enclosed by Fatty Acid
Membranes. Chemistry and Physics of Lipids 1976;16(2):142–160; doi: 10.1016/0009-
3084(76)90006-2.

Giacobelli VG, Fujishima K, Lepšík M, et al. In Vitro Evolution Reveals Noncationic Protein–RNA
Interaction Mediated by Metal Ions. Echave J. ed. Molecular Biology and Evolution
2022;39(3):msac032; doi: 10.1093/molbev/msac032.

Gorlero M, Wieczorek R, Adamala K, et al. Ser-His Catalyses the Formation of Peptides and PNAs.
FEBS Letters 2009;583(1):153–156; doi: 10.1016/j.febslet.2008.11.052.

Hanczyc MM, Fujikawa SM and Szostak JW. Experimental Models of Primitive Cellular
Compartments: Encapsulation, Growth, and Division. Science 2003;302(5645):618–622; doi:
10.1126/science.1089904.

Hargreaves WR and Deamer DW. Liposomes from Ionic, Single-Chain Amphiphiles. ACS 1978;17.

Hausrath EM and Tschauner O. Natural Fumarolic Alteration of Fluorapatite, Olivine, and Basaltic
Glass, and Implications for Habitable Environments on Mars. Astrobiology 2013;13(11):1049–1064;
doi: 10.1089/ast.2013.0985.

Hentrich C and Szostak JW. Controlled Growth of Filamentous Fatty Acid Vesicles under Flow.
Langmuir 2014;30(49):14916–14925; doi: 10.1021/la503933x.

Higgs PG and Pudritz RE. A Thermodynamic Basis for Prebiotic Amino Acid Synthesis and the
Nature of the First Genetic Code. Astrobiology 2009;9(5):483–490; doi: 10.1089/ast.2008.0280.

Hlouchová K. Peptides En Route from Prebiotic to Biotic Catalysis. Accounts of Chemical Research
2024;57(15):2027–2037; doi: 10.1021/acs.accounts.4c00137.

Hoff SE, Liu J and Heinz H. Binding Mechanism and Binding Free Energy of Amino Acids and Citrate
to Hydroxyapatite Surfaces as a Function of Crystallographic Facet, pH, and Electrolytes. Journal of
Colloid and Interface Science 2022;605:685–700; doi: 10.1016/j.jcis.2021.07.109.

Jeyapalina S, Hillas E, Beck JP, et al. Fluorapatite and Fluorohydroxyapatite Apatite Surfaces Drive
Adipose-Derived Stem Cells to an Osteogenic Lineage. Journal of the Mechanical Behavior of
Biomedical Materials 2022;125:104950; doi: 10.1016/j.jmbbm.2021.104950.

Jordan SF, Rammu H, Zheludev IN, et al. Promotion of Protocell Self-Assembly from Mixed
Amphiphiles at the Origin of Life. Nat Ecol Evol 2019;3(12):1705–1714; doi: 10.1038/s41559-019-
1015-y.

Kamat NP, Tobé S, Hill IT, et al. Electrostatic Localization of RNA to Protocell Membranes by
Cationic Hydrophobic Peptides. Angewandte Chemie - International Edition 2015;54(40):11735–
11739; doi: 10.1002/anie.201505742.

Kwiatkowski W, Bomba R, Afanasyev P, et al. Prebiotic Peptide Synthesis and Spontaneous
Amyloid Formation Inside a Proto-Cellular Compartment. Angewandte Chemie - International
Edition 2021;60(10):5561–5568; doi: 10.1002/anie.202015352.

Lambert JF. Adsorption and Polymerization of Amino Acids on Mineral Surfaces: A Review. Origins
of Life and Evolution of Biospheres 2008;38(3):211–242; doi: 10.1007/s11084-008-9128-3.

Li Z, Bhowmik S, Sagresti L, et al. Simulating Metal-Imidazole Complexes. J Chem Theory Comput
2024;20(15):6706–6716; doi: 10.1021/acs.jctc.4c00581.

Longo LM, Despotović D, Weil-Ktorza O, et al. Primordial Emergence of a Nucleic Acid-Binding
Protein via Phase Separation and Statistical Ornithine-to-Arginine Conversion. Proceedings of the
National Academy of Sciences of the United States of America 2020;117(27):15731–15739; doi:
10.1073/pnas.2001989117.

Longo LM, Lee J and Blaber M. Simplified Protein Design Biased for Prebiotic Amino Acids Yields a
Foldable, Halophilic Protein. Proceedings of the National Academy of Sciences of the United
States of America 2013;110(6):2135–2139; doi: 10.1073/pnas.1219530110.

Makarov M, Rocha ACS, Krystufek R, et al. Early Selection of the Amino Acid Alphabet Was
Adaptively Shaped by Biophysical Constraints of Foldability. Journal of the American Chemical
Society 2023;145:5320–5329; doi: 10.1021/jacs.2c12987.

Martínez M, Shearer CK and Brearley AJ. Epitaxial Fluorapatite Vein in Northwest Africa 998 Host
Apatite: Clues on the Geochemistry of Late Hydrothermal Fluids on Mars. Meteorit & Planetary
Scien 2023;58(9):1229–1245; doi: 10.1111/maps.14042.

Matreux T, Schmid A, Rappold M, et al. Heat Flows Solubilize Apatite to Boost Phosphate
Availability for Prebiotic Chemistry. Nature Communications 2025;16(1); doi: 10.1038/s41467-025-
57110-3.

Mayer C, Schreiber U, Dávila MJ, et al. Molecular Evolution in a Peptide-Vesicle System. Life
2018;8(2); doi: 10.3390/life8020016.

McCollom TM, Ritter G and Simoneit BRT. Lipid Synthesis Under Hydrothermal Conditions by
Fischer- Tropsch-Type Reactions. Orig Life Evol Biosph 1999;29(2):153–166; doi:
10.1023/A:1006592502746.

McDonald GD and Storrie-Lombardi MC. Biochemical Constraints in a Protobiotic Earth Devoid of
Basic Amino Acids: The "BAA(-) World." Astrobiology 2010;10(10):989–1000; doi:
10.1089/ast.2010.0484.

Mitchell DJ, Steinman L, Kim DT, et al. Polyarginine Enters Cells More Efficiently than Other
Polycationic Homopolymers. Journal of Peptide Research 2000;56(5):318–325; doi:
10.1034/j.1399-3011.2000.00723.x.

Monnard P-A and Deamer DW. Membrane Self-Assembly Processes: Steps Toward the First
Cellular Life. In: The Minimal Cell. (Luisi PL and Stano P. eds) Springer Netherlands: Dordrecht;
2011; pp. 123–151; doi: 10.1007/978-90-481-9944-0_8.

Morigaki K and Walde P. Fatty Acid Vesicles. Current Opinion in Colloid and Interface Science
2007;12(2):75–80; doi: 10.1016/j.cocis.2007.05.005.

Murphy J and Riley JP. A Modified Single Solution Method for the Determination of Phosphate in
Natural Waters. Analytica Chimica Acta 1962;27(C):31–36; doi: 10.1016/S0003-2670(00)88444-5.

Oparin A. Proischogdenie Zhizni. Moscovsky Robotchii 1924.

Patel BH, Percivalle C, Ritson DJ, et al. Common Origins of RNA, Protein and Lipid Precursors in a
Cyanosulfidic Protometabolism. Nature Chemistry 2015;7(4):301–307; doi: 10.1038/nchem.2202.

Paytan A and McLaughlin K. The Oceanic Phosphorus Cycle. Chemical Reviews 2007;107(2):563–
576; doi: 10.1021/cr0503613.

Posner A. The Mineral of Bone. Clinical Orthopaedics and Related Research 1985;(200):87–99.

Preiner M, Asche S, Becker S, et al. The Future of Origin of Life Research: Bridging Decades-Old
Divisions. Life 2020;10(3); doi: 10.3390/life10030020.

Rimola A, Costa D, Sodupe M, et al. Silica Surface Features and Their Role in the Adsorption of
Biomolecules: Computational Modeling and Experiments. Chemical Reviews 2013;113(6):4216–
4313; doi: 10.1021/cr3003054.

Sahai N, Kaddour H, Dalai P, et al. Mineral Surface Chemistry and Nanoparticle-Aggregation
Control Membrane Self-Assembly. Sci Rep 2017;7(1):43418; doi: 10.1038/srep43418.

Solis AD. Reduced Alphabet of Prebiotic Amino Acids Optimally Encodes the Conformational
Space of Diverse Extant Protein Folds. BMC Evol Biol 2019;19(1):158; doi: 10.1186/s12862-019-
1464-6.

Tavafoghi M and Cerruti M. The Role of Amino Acids in Hydroxyapatite Mineralization. J R Soc
Interface 2016;13(123):20160462; doi: 10.1098/rsif.2016.0462.

Tretyachenko V, Vymětal J, Neuwirthová T, et al. Modern and Prebiotic Amino Acids Support
Distinct Structural Profiles in Proteins. Open Biology 2022;12(220040); doi:
https://doi.org/10.1098/rsob.220040.

Trifonov EN. Consensus Temporal Order of Amino Acids and Evolution of the Triplet Code. Gene
2000;261(1):139–151; doi: 10.1016/S0378-1119(00)00476-5.

Trzaskowska M, Vivcharenko V and Przekora A. The Impact of Hydroxyapatite Sintering
Temperature on Its Microstructural, Mechanical, and Biological Properties. IJMS 2023;24(6):5083;
doi: 10.3390/ijms24065083.

Wang A, Chan Miller C and Szostak JW. Core-Shell Modeling of Light Scattering by Vesicles: Effect
of Size, Contents, and Lamellarity. Biophysical Journal 2019;116(4):659–669; doi:
10.1016/j.bpj.2019.01.006.

West T, Sojo V, Pomiankowski A, et al. The Origin of Heredity in Protocells. Phil Trans R Soc B
2017;372(1735):20160419; doi: 10.1098/rstb.2016.0419.

Woods AS and Ferré S. Amazing Stability of the Arginine–Phosphate Electrostatic Interaction. J
Proteome Res 2005;4(4):1397–1402; doi: 10.1021/pr050077s.

Yu WH, Li N, Tong DS, et al. Adsorption of Proteins and Nucleic Acids on Clay Minerals and Their
Interactions: A Review. Applied Clay Science 2013;80–81:443–452; doi:
10.1016/j.clay.2013.06.003.

Zaia DAM, Zaia CTBV and De Santana H. Which Amino Acids Should Be Used in Prebiotic Chemistry
Studies? Orig Life Evol Biosph 2008;38(6):469–488; doi: 10.1007/s11084-008-9150-5.

739