Desert Hedgehog mediates stem Leydig cell differentiation through Ptch2/Gli1/Sf1 signaling axis

Changle Zhao, Yongxun Chen, Lei Liu, Xiang Liu, Hesheng Xiao, Feilong Wang, Qin Huang, Xiangyan Dai, Wenjing Tao, Deshou Wang[*], Jing Wei[*]

Author affiliations: Integrative Science Center of Germplasm Creation in Western China (CHONGQING) Science City, Key Laboratory of Freshwater Fish Reproduction and Development (Ministry of Education), Laboratory of Aquatic Science of Chongqing, School of Life Sciences, Southwest University, Chongqing 400715, China

Prof. Deshou Wang, E-mail address: wdeshou@swu.edu.cn. Tel.: (86) 023 68252702; Fax: (86) 023 68253005.

Prof. Jing Wei, E-mail address: lalsos@swu.edu.cn. Tel.: (86) 023 68252365; Fax: (86) 023 68253005.

Abstract

Desert Hedgehog (Dhh) mutations cause Leydig cell dysfunction, yet the mechanisms governing Leydig lineage commitment through Dhh-mediated receptor selectivity, transcriptional effector specificity, and steroidogenic coupling remain elusive. In this study, using CRISPR/Cas9-mediated gene knockout and stem Leydig cells (SLCs) transplantation, we identified a critical Dhh/Patched 2 (Ptch2)/Glioma-associated oncogene homolog 1 (Gli1)/steroidogenic factor 1 (Sf1) signaling axis essential for SLC differentiation in Nile tilapia *(Oreochromis niloticus)*. Dhh deficiency resulted in defective adult Leydig cells and androgen insufficiency. Rescue experiments involving 11-ketotestosterone administration and a Dhh agonist treatment, combined with SLCs transplantation, demonstrated that Dhh regulates SLC differentiation, not survival. *In vitro* knockout of *ptch1* and *ptch2* in SLCs revealed that Ptch2 likely acts as the functional receptor for Dhh. This was further supported by *in vivo* genetic rescue experiments, where *ptch2* mutation did not impair testicular development, yet completely rescued the testicular defects in *dhh* mutants—consistent with Ptch2 acting as an inhibitory receptor whose loss alleviates Dhh pathway suppression. Luciferase assays in Gli-knockout SLCs demonstrated that Gli1 acts as the primary transcriptional effector, and transactivates *sf1* expression. Additionally, functional transplantation assays confirmed that Sf1 is indispensable for SLCs differentiation, as Sf1-overexpressing SLCs rescued differentiation, whereas *sf1*-mutant SLCs failed. Overall, our work delineates the Dhh-Ptch2-Gli1-Sf1 axis and provides fundamental insights into the endocrine regulation of Leydig cell lineage development.

Keywords: Desert Hedgehog (Dhh), stem Leydig cell, Patched 2 (Ptch2), Glioma-associated oncogene homolog 1 (Gli1), Steroidogenic factor 1 (Sf1)

Highlights

- Dhh regulates the differentiation of SLCs rather than their survival
- Ptch2 acts as the functional receptor for Dhh signaling in SLCs
- Gli1 is the principal transcriptional activator for Dhh signaling in SLCs
- Sf1 is the critical downstream effector of Gli1 in SLCs differentiation

INTRODUCTION

Leydig cells are the primary source of androgens, producing approximately 95% of circulating testosterone to sustain spermatogenesis and male fertility (Inoue, Baba, and Morohashi, 2018; Jiang et al., 2014). Leydig cells originate from stem Leydig cells (SLCs), which differentiate into steroidogenic adult Leydig cells (ALCs) under paracrine regulation (P. Chen, Zirkin, and Chen, 2020; S. Yao et al., 2022). Steroidogenic factor 1 (Sf1), an orphan nuclear receptor transcription factor, serves as a master regulator of this process by activating steroidogenic enzymes (e.g., cytochrome P450, family 11, subfamily C, polypeptide 1 [Cyp11c1], 3β-hydroxysteroid dehydrogenase [3β-HSD]) to drive ALC maturation (Parker et al., 2002). In mice, mutations of *Sf1* exhibit arrested testicular development and abolished steroidogenesis (Houzelstein et al., 2024; Park, Tong, and Jameson, 2007; Smith et al., 2023), while human *SF1* mutations are associated with 46,XY disorders of sexual development (DSD), such as Swyer syndrome (Yu, Lee, Huerta-Saenz, and Allen, 2023). Similarly, in Nile tilapia (*Oreochromis niloticus*), our previous work reveals that *sf1* mutation disrupts testicular morphogenesis and downregulates the expression of key steroidogenic enzymes such as *cyp11c1* (Xie et al., 2016). Despite its established central role, the upstream signaling pathways that regulate Sf1 to commit the Leydig lineage remain poorly defined.

Hedgehog (Hh) signaling pathway is an evolutionarily conserved developmental regulator (Y. Zhang and Beachy, 2023). Canonical Hh signaling involves the binding of ligands to Patched (Ptch) receptors, which relieves the suppression on Smoothened (Smo) and ultimately activates Glioma-associated oncogene homolog (Gli) transcription factors to regulate target genes (Briscoe and Thérond, 2013; Finco, LaPensee, Krill, and Hammer, 2015; Ingham, Nakano, and Seger, 2011). In vertebrates, there are three Hh ligands (i.e. Sonic Hh, Indian Hh and Desert Hh [Dhh]), two Ptch receptors (Ptch1 and Ptch2) and three Gli homologs (Gli1-3) (Belgacem and Borodinsky, 2015; Carpenter et al., 1998; Cervantes, Lau, Cano, Borromeo-Austin, and Hebrok, 2010; Kothandapani et al., 2020). A large number of studies have well demonstrated that Dhh signaling pathway is critical in gonadal development (Finco et

al., 2015; Umehara et al., 2000). Dhh, secreted by Sertoli cells, is critical for Leydig cell commitment in mammals (Mehta et al., 2021; Pachernegg, Georges, and Ayers, 2022). In mice, mutations of *Dhh* lead to Leydig cell dysfunction, testosterone insufficiency, and infertility (Pierucci-Alves, Clark, and Russell, 2001), while human *DHH* mutations are associated with 46,XY DSD (Wei et al., 2022). Strikingly, these phenotypes mirror those caused by *Sf1* deficiencies, suggesting a potential regulatory nexus between Dhh and Sf1 signaling (Park et al., 2007). However, the molecular circuitry connecting Dhh signaling to Sf1 activation remains elusive.

Studies demonstrate functional divergence between Ptch1 and Ptch2. For instance, *Ptch1* knockout causes embryonic lethality due to constitutive Hh activation, whereas *Ptch2* deficiency yields no overt defects (Cong, Zhu, Chen, Chen, and Chong, 2025; Nieuwenhuis et al., 2006). During testicular development, Ptch1 displays high expression at early stage critical for gonadal primordium formation, while Ptch2 shows testis-enriched dynamic regulation (Kim, Lee, Seppala, Cobourne, and Kim, 2020; H. H. Yao, Whoriskey, and Capel, 2002). However, the distinct roles of Ptch1 and Ptch2 in Leydig lineage commitment remain unknown to date. Among Gli transcription factors, Gli1 acts as a primary activator, while Gli2 and Gli3 exhibit dual regulatory functions (Hui and Angers, 2011; Ruiz i Altaba, Sánchez, and Dahmane, 2002). In murine testes, co-expression of Gli1 and Gli2 in Leydig cells illustrates functional redundancy—single knockouts show normal development, but combined pharmacological inhibition (GANT61) abolishes Leydig cells and spermatogenesis (I. Barsoum and Yao, 2011). These findings underscore the complexity of Gli-mediated transcription in the testis (Kothandapani et al., 2020) and highlight the need to delineate the specific roles of individual Gli in SLC differentiation.

Nile tilapia (*Oreochromis niloticus*) is not only an important farm fish in global aquaculture but also a distinguished animal model for investigating gene function and endocrine regulation. This model organism possesses multiple advantageous attributes, including an XX/XY sex-determination system, early sexual maturity (approximately 3 months after hatching for males and 6 months after hatching for females), a short spawning cycle of approximately 14 days, year-round reproductive capacity under

laboratory conditions, well-characterized sex-linked genetic markers, and a body size amenable to physiological sampling (Li, Sun, Zhou, and Wang, 2024; Sun et al., 2014). Furthermore, a stem Leydig cell line (TSL) has been established from the testis of a 3-month-old Nile tilapia (Huang et al., 2020). TSL expresses platelet-derived growth factor receptor α (*pdgfrα*), *nestin*, and chicken ovalbumin upstream promoter transcription factor II (*coup-flIa*), which are usually considered as SLC-related markers in several other species (H. Chen, Wang, Ge, and Zirkin, 2017; Ge et al., 2006; Jiang et al., 2014). Notably, this cell line exhibits the capacity to differentiate into 11-ketotestosterone (11-KT)-producing Leydig cells both *in vitro* and *in vivo* (Huang et al., 2020). When cultured in a defined induction medium, TSL cells differentiate into a steroidogenic phenotype, expressing key steroidogenic genes including *star1*, *star2*, and *cyp11c1*, and producing 11-KT; upon transplantation into recipient testes, TSL cells successfully colonize the interstitial compartment, activate the expression of steroidogenic genes, and restore 11-KT production. In this study, by combining CRISPR/Cas9-mediated gene knockout with TSL transplantation technology, we unveil a Dhh-Ptch2-Gli1-Sf1 axis that is essential for Leydig cell lineage differentiation. Our findings provide fundamental insights into the endocrine regulation of Leydig cell development.

RESULTS

Mutation of *dhh* disrupts testicular organization and androgen synthesis

The Nile tilapia Dhh open reading frame (GenBank ID: 100707022, 1377 bp) was cloned and validated. The encoded 458-amino acid protein shares over 56% identity with vertebrate homologs (Fig. S1A) and clusters phylogenetically with mammalian DHH orthologs (Fig. S1B). Transcriptomic analyses conducted on various tissues and testes across different developmental stages revealed that *dhh* expression peaked in the testes, particularly at 90 days after hatching (dah) (Fig. S1C, D).

CRISPR/Cas9 targeting of *dhh* exon 1 from Nile tilapia generated homozygous

mutants (*dhh*$^{-/-}$) harboring a 13-bp (CAGGGATGCGGAC) frameshift deletion (Fig.

S2A–D), resulting in premature termination and loss of the conserved Hedge domain

(Fig. S2E). *dhh* mRNA levels in *dhh*$^{-/-}$ fish were significantly reduced compared to

WT (Fig. S2F). At 90 dah, *dhh*$^{-/-}$ testes exhibited marked atrophy (Fig. 1A, B) and

disorganized histology, with sparse spermatogonia, rare spermatids, and collapsed

interstitial architecture (Fig. 1C–D', K). Immunofluorescence confirmed significantly

reduced germ cell (Vasa$^+$), meiotic cell (Sycp3$^+$), and Leydig cell (Cyp11c1$^+$)

populations (Fig. 1E-J', L-N). Serum 11-KT levels in *dhh*$^{-/-}$ fish were significantly

lower than WT (Fig. 1O), indicating impaired androgen synthesis.

The differentiation of SLCs cannot be rescued by 11-KT, but by SAG

Given the established role of 11-KT in teleost testicular development and

spermatogenesis (Zheng et al., 2020), we first asked whether exogenous 11-KT could

ameliorate the testicular defects in *dhh*$^{-/-}$ mutants. Compared to untreated mutants,

dhh$^{-/-}$ individuals exposed to 11-KT exhibited a more organized testicular architecture,

with significantly increased populations of Vasa□ and Sycp3□, as evidenced by both

H&E staining and immunofluorescence (Fig. 2A–C', E–G', I–K', R, S). This

improvement was further supported by a recovery in GSI (Fig. 2Q). However, the

population of Cyp11c1-positive Leydig cells remained profoundly depleted in *dhh*$^{-/-}$

+11-KT testes (Fig. 2M–O', T). Taken together, these data indicate that 11-KT

treatment partially restored the development of germ cells but not Leydig cell lineage,

implying that the Leydig cell defect is androgen-independent.

We next investigated whether direct activation of the Dhh pathway could rescue

the *dhh*$^{-/-}$ phenotype. Treatment of *dhh*$^{-/-}$ mutants with SAG, a Hedgehog agonist,

significantly rescued testicular development. This was marked by a significant

recovery of germ cells (Vasa□), spermatocytes (Sycp3□), and critically, Leydig cells

(Cyp11c1□), alongside a restoration of testicular histology (Fig. 2D–D', H–H', L–L',

P–P', Q–T). These results demonstrate that *dhh* mutation does not disrupt survival of

endogenous Leydig cell lineage, but directly impairs their differentiation, leading to

secondary impairment in androgen production and germ cell development.

Mutation of *dhh* blocks SLCs differentiation *in vivo*

To dissect Leydig cell lineage impairment in $dhh^{-/-}$ testes, we transplanted the

TSL labeled with PKH26 (a fluorescent red hydrophobic membrane dye that enables

tracking of transplanted cells) into WT and $dhh^{-/-}$ testes (Fig. 3A). In WT recipients,

TSL cells colonized the interstitium and differentiated into Cyp11c1

positive steroidogenic cells (Fig. 3B1–B3). In $dhh^{-/-}$ testes, TSL engrafted equivalently

but failed to express Cyp11c1 (Fig. 3C1–C3), indicating a differentiation-specific

block. Rescue experiments showed that the TSL overexpressing tilapia Dhh (TSL-

OnDhh) or treated with SAG (TSL+SAG) restored Cyp11c1 expression in $dhh^{-/-}$ testes

(Fig. 3D1–E3, F) and a significant increase in endogenous 11-KT levels (Fig. 3G).

These results suggest that SLC differentiation is inhibited, whereas the survival and

engraftment of PKH26-labeled TSL cells were not affected in $dhh^{-/-}$ XY tilapia testes.

Collectively, these results demonstrate that Dhh signaling is specifically required for

the differentiation of SLCs into steroidogenic cells, but is dispensable for their

survival or initial recruitment into the testicular niche.

Ptch2 is the functional receptor for Dhh signaling in SLCs

RT-PCR results showed that both *ptch1* and *ptch2* were expressed in TSL (Fig.

S3). Fluorescence *in situ* hybridization (FISH) and IF revealed co-expression

of *ptch1* and *ptch2* in Cyp11c1 positive Leydig cells (Fig. 4A). To identify the Dhh-

driving receptor selectivity in SLCs, we established $ptch1^{-/-}$ and $ptch2^{-/-}$ TSL cell lines

(Figs. S4A, B) and evaluated pathway activity using an 8×GLI luciferase reporter

system (Fig. 4B). In WT TSL cells, the 8×GLI reporter demonstrated substantial

luciferase activity that was markedly enhanced by OnDhh overexpression (Fig. 4C),

confirming Dhh responsiveness in this system. Intriguingly, $ptch1^{-/-}$ cells maintained

Dhh-induced luciferase activation comparable to WT controls, whereas $ptch2^{-/-}$ cells

completely lost this responsiveness (Fig. 4D). This differential receptor requirement

implies that Ptch2 likely acts as the functional receptor for transducing Dhh signals in

TSL cells.

To investigate the function of Ptch2 *in vivo*, we generated $ptch2^{-/-}$ and $dhh^{-/-}$

;$ptch2^{-/-}$ mutant Nile tilapias (Figs. S5A-E). Compared to WT fish at 90 dah, $ptch2^{-/-}$

testes exhibited no significant differences in testicular histology (Fig. 4E–F'), GSI

(Fig. 4Q), germ cell (Vasa$^+$, Sycp3$^+$) and Leydig cell (Cyp11c1$^+$) populations (Fig.

4H-I', K-L', N-O', R-T), and serum 11-KT levels (Fig. 4U). Strikingly, the loss

of *ptch2* in the *dhh*$^{-/-}$ background (*dhh*$^{-/-}$;*ptch2*$^{-/-}$) fully rescued the testicular defects,

resulting in normal histology (Fig. 4G–G'), Leydig cell differentiation (Fig. 4P–P'),

and serum 11-KT levels (Fig. 4U). This genetic rescue is consistent with Ptch2

functioning as the inhibitory receptor for Dhh; its removal alleviates the suppression

on the pathway, thereby bypassing the requirement for the Dhh ligand.

Gli1 is the principal transcriptional effector of Dhh signaling in SLCs

In the Hh pathway, Gli transcription factors (Gli1/2/3) mediate diverse

developmental processes (26). RT-PCR results showed that *gli1*, *gli2*, and *gli3* were

all expressed in TSL (Fig. S3). FISH and IF revealed co-expression of *gli1*, *gli2*,

and *gli3* in Cyp11c1 positive Leydig cells (Fig. 5A). To identify Dhh-specific

effectors, we generated *gli1*$^{-/-}$, *gli2*$^{-/-}$, and *gli3*$^{-/-}$ TSL lines (Fig. S4C–E) and assessed

pathway activity using the 8×GLI luciferase reporter. While TSL-*gli2*$^{-/-}$ and TSL-*gli3*$^{-/-}$

cell lines maintained Dhh-induced luciferase activation comparable to WT controls,

the TSL-*gli1*$^{-/-}$ cell line completely lost this responsiveness (Fig. 5B), implying that

Gli1 is the principal transcriptional effector for Dhh signaling transduction in SLCs.

Sf1 is the critical downstream effector of Gli1 in SLCs differentiation

To identify key targets of the Dhh-Gli1 axis, we performed transcriptomic

profiling of TSL cells under conditions of pathway activation: Dhh overexpression

(TSL-*On*Dhh), Gli1 overexpression (TSL-OnGli1), and SAG treatment (TSL+SAG).

Comparative RNA-seq analysis identified a core set of 33 genes consistently

upregulated across all three conditions (Fig. 5C, S6A). Among these, *sf1* emerged as a

prominently upregulated candidate. Functional annotation of its promoter region

identified two conserved Gli1-binding motifs, B1 (AACCACCCA) and B2

(GAGCCACCCA) (Fig. 5D). A dual-luciferase reporter assay confirmed that Gli1

potently transactivates the *sf1* promoter (Fig. 5E). The specificity of this interaction

was further demonstrated by a cold probe competition assay: a wild-type

oligonucleotide containing the Gli-binding motif (GACCACCCA) competitively

inhibited Gli1-driven *sf1* transactivation in a dose-dependent manner, whereas a

mutated version (TTAATTAAA) had no effect (Fig. 5E).

The functional necessity of Sf1 was confirmed by transplantation assays using

sf1^{-/-} and Sf1-overexpressing (TSL-*On*Sf1) TSL cells. While TSL-*sf1^{-/-}* cells failed to

differentiate into Cyp11c1-positive cells even in WT testes, TSL-*On*Sf1 cells

successfully differentiated and expressed Cyp11c1 even in the *dhh^{-/-}* testicular

environment (Fig. 5F–H). These results definitively position Sf1 as the critical

downstream effector executing the Dhh-Ptch2-Gli1 signaling cascade during SLCs

differentiation.

DISCUSSION

This study delineates a critical Dhh-Ptch2-Gli1-Sf1 signaling axis essential for

SLCs differentiation in Nile tilapia. By combining *dhh^{-/-}* XY models with SLCs

transplantation, we definitively demonstrate that Dhh signaling is dispensable for

Leydig cell lineage survival but indispensable for its differentiation. Furthermore, our

integrated evidence from targeted gene knockout and functional assays suggests that

Ptch2 acts as the primary receptor for Dhh signaling in SLCs, and points to Gli1 as

the key transcriptional effector mediating the transactivation of *sf1*. Collectively, these

findings implicate Sf1 as the critical downstream target bridging Dhh signaling to

steroidogenic maturation, thereby resolving key ambiguities in Leydig cell lineage

development and providing evolutionary insights into vertebrate reproductive biology.

Consistent with its established role in mammals, our data in Nile tilapia confirm

that Dhh function in male testis development is evolutionarily conserved across

vertebrates. While mutations in *Dhh* lead to testosterone insufficiency and infertility

in mice (Pierucci-Alves et al., 2001) and are associated with 46,XY DSD in humans

(Wei et al., 2022), the precise cellular mechanism remained unresolved. Our study

demonstrates that *dhh^{-/-}* tilapia phenocopy these defects, exhibiting testicular

dysgenesis, a profound deficiency of Leydig cells, and drastically reduced 11-KT

levels (Fig. 1). To define the onset of Leydig cell differentiation, we performed a

developmental time-course analysis. This revealed that Cyp11c1-positive

steroidogenic cells first appear in wild-type testes at 30 dah, while being

conspicuously absent in *dhh^{-/-}* mutants at this same stage (Fig. S7). This clear

temporal pattern establishes ~30 dah as the developmental window when SLCs

initiate their differentiation program in the Nile tilapia. The functional indispensability

of Dhh at this specific stage was further confirmed by rescue experiments, in which

both SAG treatment and transplantation of Dhh-activated TSL cells restored

Cyp11c1-positive cell populations in $dhh^{-/-}$ testes (Fig. 3). Collectively, our findings

position Dhh as a crucial niche signal that acts at a defined developmental checkpoint

to drive the differentiation of SLCs into steroidogenic Leydig cells.

The functional divergence of Ptch1 and Ptch2 in Hh signaling represents an

intriguing aspect of pathway regulation. Structurally, both receptors share conserved

12-transmembrane domains vital for cholesterol transport but diverge in their

cytoplasmic C-terminal regions, which are implicated in differential Smoothened

inhibition efficiency (Fleet and Hamel, 2018; Qi, Schmiege, Coutavas, and Li, 2018).

In mammals, Ptch1 serves as the primary, broad-spectrum regulator of Hh signaling

during embryogenesis, whereas Ptch2 often assumes context-dependent roles, such as

in primordial germ cell migration via its co-receptor Gas1 (Kim et al., 2020).

However, their potential redundancy or specificity within the Leydig cell lineage has

remained an open question. Our study provides significant insight into this issue. We

confirmed that both *ptch1* and *ptch2* are expressed in Leydig cells (Fig. 4A).

Functional interrogation using TSL cell lines revealed a marked difference: genetic

ablation of *ptch2* profoundly attenuated Dhh-induced signaling, as measured by a Gli-

responsive luciferase reporter, whereas loss of *ptch1* had no detectable effect under

the same conditions (Fig. 4B-D). This *in vitro* data, pointing to a primary role for

Ptch2, was further supported by *in vivo* genetic interaction studies. The observation

that the $dhh^{-/-}$ phenotype was fully rescued in $dhh^{-/-};ptch2^{-/-}$ double mutants (Fig. 4E-U)

is consistent with Ptch2 acting as the inhibitory receptor whose loss alleviates

pathway suppression. The specificity for Ptch2 in this context might stem from unique

co-receptor interactions or expression patterns within the testicular niche. To

preliminarily assess potential compensatory regulation, we examined *ptch1*

expression in XY testes from WT, $ptch2^{-/-}$ and $dhh^{-/-};ptch2^{-/-}$ fish at 90 dah. No

significant differences in *ptch1* mRNA levels were detected among these genotypes

(Fig. S8), suggesting that loss of *ptch2* does not trigger compensatory upregulation of

ptch1 at the transcriptional level under the conditions examined. Nonetheless, global

ptch2 mutation affects multiple tissues, whereas our mechanistic focus is on SLC

differentiation within the testicular niche. Moreover, the early embryonic lethality of

global *ptch1* mutation in tilapia (Liu et al., 2024) precludes direct assessment of its

role in postnatal testis development. Therefore, although our findings strongly support

a predominant role for Ptch2 in mediating Dhh signaling in SLCs, definitive

resolution of receptor specificity will require future Leydig cell-specific conditional

knockout models.

Gli is a nuclear transcription factor, characterized by the presence of a zinc finger

domain located in the middle of the protein (Kinzler, Ruppert, Bigner, and Vogelstein,

1988). Downstream of the zinc finger domain is a phosphorylation cluster containing

phosphorylation sites for protein kinase A and glycogen synthase kinase 3, which

negatively regulate the transcriptional activity of Gli through phosphorylation

(Niewiadomski et al., 2014). In *Drosophila*, Ci is the sole Gli homolog, possessing

both transcriptional activation and repression functions (Huangfu and Anderson,

2006). In vertebrates, there are three Gli homologs: Gli1, Gli2, and Gli3. Gli1

functions solely as a transcriptional activator because it lacks proteolytic processing

domains. Gli2 has both activation and repression functions, while Gli3 primarily acts

as a transcriptional repressor (Niewiadomski et al., 2019; Pan and Wang, 2007).

However, there has been no research on the regulatory mechanism of Gli1/2/3 in

SLCs differentiation. In this study, we found that although *gli1*, *gli2*, and *gli3* are all

expressed in Leydig cell lineage (Figure 5A-C), the results from TSL cell lines with

mutations in *gli1*, *gli2*, and *gli3* indicated that TSL-*gli1$^{-/-}$*, rather than TSL-*gli2$^{-/-}$* or

TSL-*gli3$^{-/-}$*, failed to respond to Dhh signaling (Figure 5D). This suggests that Gli1 is

the key transcription factor downstream of the Dhh signaling cascade in SLCs.

Sf1 and Dhh play important roles in Leydig cell commitment and testicular

development (Parker et al., 2002; Y. Zhang and Beachy, 2023). Studies suggest a

potential interaction between these two factors. For example, in *Dhh$^{-/-}$* XY mice,

testicular development is impaired, with a significant reduction in the number of

Leydig cells and a marked downregulation of *Sf1* expression (H. H. Yao et al., 2002).

In XX mouse ovaries where Smo is constitutively activated, a large number of ectopic

Sf1-positive Leydig cells are generated (I. B. Barsoum, Bingham, Parker, Jorgensen,

and Yao, 2009). Similarly, in *sf1$^{-/-}$* XY tilapia, testicular development is impaired, with

a significant reduction in Leydig cell numbers and down regulation of *dhh* expression

(Xie et al., 2016). These findings imply that Dhh and Sf1 may interact through

feedback loops or co-regulatory mechanisms. However, direct evidence for the

regulation of Sf1 by the Hh signaling pathway remains lacking. This study elucidates

the molecular mechanism of Sf1 as a critical downstream effect of the Dhh-Ptch2-

Gli1 axis. Transcriptomic profiling revealed an increased expression of *sf1* by Dhh regulation signaling in TSL cells (Fig. S6), identifying it as a key downstream target of Dhh signaling. Promoter analysis identified two conserved Gli-binding motifs (Fig. 5F) within the *sf1* promoter, and luciferase assays confirmed that Gli1 transactivates *sf1* transcription (Fig. 5G). Functional validation demonstrated that TSL-*sf1*$^{-/-}$ cells failed to differentiate even in WT testes, whereas TSL-OnSf1 cells restored Leydig cell differentiation and steroidogenesis in *dhh*$^{-/-}$ testes (Fig. 5H-J). These results establish Sf1 as both necessary and sufficient for Leydig cell lineage differentiation.

In conclusion, our study systematically deciphers the signaling cascade governing SLCs differentiation, from the niche signal to the core steroidogenic regulator. We definitively show that the Dhh signal from the niche is indispensable for SLCs differentiation, but not their survival. Within this pathway, Ptch2 serves as the critical receptor, relaying the signal through the transcription factor Gli1. The pivotal outcome of this cascade is the activation of *sf1* by Gli1. Thus, we establish a complete Dhh-Ptch2-Gli1-Sf1 axis, resolving the long-standing question of how a key morphogen controls Leydig cell lineage development in vertebrates.

MATERIALS AND METHODS

Animals

Nile tilapias (*Oreochromis niloticus*) were kept in recirculating freshwater systems at 26□ under a 12-h light/dark photoperiod. All animal experiments were conducted in accordance with the regulations of the Guide for Care and Use of Laboratory Animals prescribed by the Committee of Laboratory Animal Experimentation at Southwest University, China (IACUC-20181015-12).

Generation of mutant lines by CRISPR/Cas9 in Nile tilapia

The sequences of tilapia *dhh* (GenBank ID: 100707022) and *ptch2* (GenBank ID: 100692939) were retrieved from the NCBI. CRISPR/Cas9-mediated mutagenesis was performed as described previously (Li et al., 2014). Guide RNAs (gRNAs) targeting the first exon of *dhh* (GCGGGCCCGGTCCGCATCCC) and *ptch2* (GTCCCAGGGGCCGGCGTATT) were designed using ZiFit

(http://zifit.partners.org/ZiFiT/). gRNA and Cas9 mRNA were synthesized following establish protocols (Li et al., 2014). The primers listed in Table S1. A mixture of synthetic gRNA (500 ng/μl) and Cas9 mRNA (1000 ng/μl) was microinjected into one-cell-stage fertilized embryos. Mutants were validated by polyacrylamide gel electrophoresis (PAGE) and Sanger sequencing. F0 chimeric XY males were crossed with WT XX females to generate heterozygous F1 offspring. F1 siblings carrying a 13-bp deletion in *dhh* or a 25-bp deletion in *ptch2* were intercrossed to generate homozygous F2 mutants. For double mutants, *dhh*[+/-] XY males were mated with *ptch2*[+/-] XX females to obtain *dhh*[+/-];*ptch2*[+/-] offspring. These double heterozygotes were then intercrossed to generate *dhh*[-/-];*ptch2*[-/-] mutants.

Generation of mutant lines by pCas9-NtU6sgRNA system in TSL Cells

The tilapia stem Leydig cell line TSL was maintained in ESM4 medium as described (Huang et al., 2020). Gene sequences for *ptch1* (GenBank ID: 100700194), *ptch2* (GenBank ID: 100692939), *gli1* (GenBank ID: 100710687), *gli2* (GenBank ID: 100704788), *gli3* (GenBank ID: 100707331), and *sf1* (GenBank ID: 100534561) were retrieved from NCBI. Knockout plasmids targeting these genes were designed following established protocols (Z. Zhang et al., 2023). gRNAs were generated using ZiFit, with the following exon-specific targets: *ptch1* (GGAGGCGCTCCTGCAGCACC, exon 3), *ptch2* (GTCCCAGGGGCCGGCGTATT, exon 1), *gli1* (TGACTCATGGATCAGGACCA, exon 2), *gli2* (GATTCGCCTGTCACCCCACG, exon 3), *gli3* (TGAGGAGCCCTCTACGTCTA, exon 2), and *sf1* (GGCTGTGTACTGGTACTGGG, exon 4; designed as previously described) (Z. Zhang et al., 2023). Primers are listed in Table S1. Plasmids were transfected into TSL cells using the MDMP E-1 microchannel transfection instrument (MENDGENE) with transwell cell compartments (BIOFIT), per manufacturer guidelines. Post-transfection (48 hours), cells were selected in ESM4 medium containing 500 μg/mL G418 for 7 days. Single-cell clones were isolated via dilution and serial passage, followed by genomic DNA (gDNA) extraction and Sanger sequencing to confirm

381 mutations.

Histological analyses of testes from WT, $dhh^{-/-}$, $ptch2^{-/-}$ and $dhh^{-/-};ptch2^{-/-}$ XY fish

GSI was calculated for WT, $dhh^{-/-}$, $ptch2^{-/-}$ and $dhh^{-/-}; ptch2^{-/-}$ XY fish at 90 dah

as: (gonad weight / body weight) $\times$ 100%. Testes were fixed in Bouin's solution for 24

hours at room temperature with agitation, dehydrated through an ethanol series, and

embedded in paraffin. Sections (5 μm thickness) were stained with hematoxylin and

eosin (H&E) and imaged using an Olympus BX53 microscope. Germ cell

classification (spermatogonia, spermatocytes) followed established criteria (Zhao et

al., 2023).

Immunofluorescence

Testes from WT, $dhh^{-/-}$, WT+11-KT, $dhh^{-/-}$+11-KT, $ptch2^{-/-}$ and $dhh^{-/-}; ptch2^{-/-}$ XY

fish at 90 dah were analyzed for germ cell marker Vasa, Leydig cell marker Cyp11c1,

and meiosis marker Sycp3. Tissue fixation and embedding. Sections (5 μm) were

deparaffinized, rehydrated, and subjected to antigen retrieval. After blocking with 5%

bovine serum albumin (BSA; 30 minutes, room temperature), sections were incubated

overnight at 4°C with rabbit polyclonal primary antibodies (Vasa, Cyp11c1, Sycp3;

1:1,000 dilution), validated previously (Dai et al., 2021). Alexa Fluor 488-conjugated

goat anti-rabbit secondary antibody (1:1,000; Invitrogen) was applied for 1 hour at

37°C, followed by DAPI nuclear counterstaining (1:1,000; Sigma-Aldrich).

Fluorescence signals were imaged using an Olympus Fv3000 confocal microscope.

For each biological replicate (n=5-6 fish per genotype), three non-serial, non-adjacent

testis sections were analyzed. From each section, three representative fields of view

were captured to ensure non-overlapping sampling. All positive cells number of Vasa,

Sycp3 and Cyp11c1 was quantified by Image J Pro 1.51 software using default

parameters.

Fluorescence *in situ* hybridization (FISH) and immunofluorescence

Testes from WT XY tilapia at 90 dah were fixed in 4% paraformaldehyde (Sigma)

overnight at 4°C with agitation, paraffin-embedded, and sectioned (5 μm). FISH was

performed as described (Zhao et al., 2023). Partial cDNA fragments

of *ptch1*, *ptch2*, *gli1*, *gli2*, and *gli3* were cloned into the pGEM-T Easy vector. Sense

(control) and antisense RNA probes were synthesized using a DIG RNA Labeling Kit

(Roche) and T7 mMESSAGE mMACHINE Kit (Ambion). Signals were amplified

with the TSA^{TM} Plus TMR system (PerkinElmer) per manufacturer protocols (primers

in Table S1). Concurrently, Leydig cell marker Cyp11c1 expression was assessed by

immunofluorescence.

11-KT and SAG rescue experiments

To rescue *dhh*$^{-/-}$ XY fish, offspring from *dhh*$^{+/-}$ XY male × *dhh*$^{+/-}$ XX female

crosses were divided into two groups. The experimental group received 11-KT via

immersion from 30 to 90 dah, with treatments refreshed every 2 days. 11-KT

concentrations increased incrementally from 57 ng/L at 30 dah to 1254 ng/L by 90

dah (39.9 ng/L increase per treatment cycle; see Fig. 2, 5, S7). For SAG rescue, fish

received intraperitoneal injections of 10 mg/kg SAG from 30 to 90 dah, with booster

doses administered weekly. Testes were collected at 90 dah for morphological and

histological evaluation.

TSL cell labeling, transplantation and IF

TSL, previously isolated from 3-month-old XY tilapia testes (Huang et al., 2020),

was used to assess stem Leydig cell differentiation in WT and *dhh*$^{-/-}$ XY fish. TSL

cells were treated as follows: (1) untreated TSL, (2) TSL + 0.5 μM SAG (Hh pathway

agonist), (3) TSL-*On*Dhh (overexpressing tilapia Dhh), (4) TSL-*On*Sf1

(overexpressing tilapia Sf1), and (5) TSL-*sf1*$^{-/-}$ (*sf1* mutation). For the SAG treatment

experiment, TSL cells were incubated with 0.5 μM SAG for 48 hours before

transplantation. Cells were labeled with PKH26 fluorescent dye (Sigma) and

transplanted (5,000 cells per testis) into WT or *dhh*$^{-/-}$ XY tilapia via the urogenital

tract (Huang et al., 2020). For TSL-*On*Dhh, the Dhh ORF (1377 bp) was amplified

from gDNA, cloned into pcDNA3.1 (Invitrogen), and transfected into TSL cells

(primers in Table S1). At 10 days post-transplantation, testes were analyzed by IF to

quantify Cyp11c1 positive cells among PKH26 positive populations.

GLI report assay

A firefly luciferase reporter plasmid (8×GLI) was generated by cloning eight

tandem copies of the GLI-binding motif (GACCACCCA) into the pGL4.23 vector

(minimal promoter). To validate 8×GLI functionality, TSL cells were transiently

transfected with: (1) pGL4.23 (negative control), (2) 8×GLI, or (3) 8×GLI plus Dhh-

overexpression plasmid (*On*Dhh), for 48 hours. The 8×GLI construct was further

transfected into TSL (WT), TSL-*ptch1$^{-/-}$*, TSL-*ptch2$^{-/-}$*, TSL-*gli1$^{-/-}$*, TSL-*gli2$^{-/-}$* or TSL-

gli3$^{-/-}$, with or without *On*Dhh. Transfections included pRL-TK as an internal control.

Luciferase activity was measured 48 hours post-transfection using the Dual Luciferase

Assay System (Promega) and a Luminoskan™ Ascent luminometer (Thermo Fisher

Scientific). Relative activity was calculated as firefly/Renilla luciferase ratios.

Western blot

The ORFs of Dhh (1377 bp) and Gli1 (4404 bp) were amplified from gDNA by

PCR. Two tandem FLAG tag sequences (GATTATAAAGATGATGATAAA) were

inserted flanking each ORF (after the start codon and before the stop codon) and

cloned into the pcDNA3.1 vector (Invitrogen), generating pcDNA3.1-*On*Dhh and

pcDNA3.1-*On*Gli1 (primers in Table S1). TSL cells were transiently transfected with

pcDNA3.1-*On*Dhh or pcDNA3.1-*On*Gli1 for 48 hours. Protein expression was

analyzed by Western blot following established methods (Dai et al., 2021). Briefly,

total protein lysates were separated by 15% SDS-PAGE, transferred to nitrocellulose

membranes, and blocked with 5% BSA in TBST (10 mM Tris pH 7.9, 150 mM NaCl,

0.1% Tween-20) for 1 hour at 37°C. Membranes were incubated overnight at 4°C

with rabbit anti-FLAG antibody (1:1000; Cell Signaling Technology), followed by

HRP-conjugated goat anti-rabbit secondary antibody (1:1000; Invitrogen) for 1 hour

at 37°C. Signals were detected using the BeyoECL Plus Kit (Beyotime) and imaged

on a Fusion FX7 system (Vilber Lourmat).

Measurement of 11-KT levels by ELISA

Blood was collected from the caudal vein of 90 dah XY fish (WT, *dhh$^{-/-}$*, *ptch2$^{-/-}$*,

dhh;ptch2$^{-/-}$, WT+TSL, *dhh*$^{-/-}$+TSL, *dhh*$^{-/-}$+TSL+SAG, *dhh*$^{-/-}$+TSL-*On*Dhh and *dhh*$^{-/-}$

+TSL-*sf1*$^{-/-}$), incubated at 4°C overnight, and centrifuged to isolate serum. Tissue fluid

samples from WT and *dhh*$^{-/-}$ XY fish at 5, 10, 20, and 30 dah were similarly prepared.

All samples were stored at −80°C until analysis. Serum and tissue fluid 11-KT levels

were quantified using a 11-keto Testosterone ELISA Kit (Cayman) validated for

teleost samples, per manufacturer instructions. Individual samples (serum or tissue

fluid; $n = 6$ per genotype) were analyzed in triplicate on a single 96-well plate, with

concentrations calculated from standard curves.

Transcriptome analyses

Tissue- and stage-specific expression of *dhh* was assessed using transcriptomic

data from tilapia tissues (brain, heart, head kidney, kidney, liver, muscle, ovary, testis)

and gonads at developmental stages spanning 5 to 360 dah (Tao et al., 2018; Tao et al.,

2013). For global gene expression profiling, RNA was extracted from TSL-WT, TSL-

*On*Dhh, TSL-*On*Gli1, and TSL + 0.5 µM SAG cells using RNAiso Plus (Takara). For

the SAG treatment experiment, TSL cells were incubated with 0.5 µM SAG for 48

hours before collection. For each genotype, cells from three independent culture wells

were pooled. Libraries were sequenced on an Illumina platform (GENEBOOK

Biotechnology), and clean reads were aligned to the reference genome with HISAT2.

Gene expression levels (FPKM: fragments per kilobase per million mapped reads)

were quantified using featureCounts. Differentially expressed genes (DEGs) were

identified for each condition (TSL-*On*Dhh, TSL-*On*Gli1, TSL+SAG) compared to

TSL-WT controls using edgeR (threshold: FDR < 0.05, |log2(foldchange)| ≥ 1.5).

Real-time PCR

Testes from WT and *dhh*$^{-/-}$ XY fish at 90 dah ($n = 3$ per genotype), along with

TSL-WT, TSL-*On*Dhh, TSL-*On*Gli1, and TSL + 0.5 µM SAG cells, were analyzed

for gene expression. Total RNA was extracted using RNAiso Plus (Takara), reverse-

transcribed into cDNA (PrimeScript II 1st Strand cDNA Synthesis Kit; Takara), and

amplified by real-time PCR on an ABI-7500 system (Applied Biosystems). Relative

mRNA levels were normalized to *β-actin* and calculated via the formula $R = 2^{-\Delta\Delta Ct}$

(primers in Table S1).

Dual luciferase report assay

The 5015-bp promoter region upstream of the *sf1* start codon was amplified and cloned into the pGL3-basic vector (Promega), generating pGL3-*sf1* (primers in Table S1). To assess Gli1 binding to the *sf1* promoter, HEK293 cells were co-transfected with pGL3-*sf1*, pcDNA3.1- *On*Gli1, cold probe (GACCACCCA), and mutant cold probe (TTAATTAAA) using the TransIT-X2 transfection reagent (Mirus Bio). Control transfections included empty pcDNA3.1 and pGL3 vectors, with pRL-TK for normalization. Luciferase activity was measured 48 hours post-transfection.

Statistical analyses

Data are presented as mean $\pm$ SD from at least three independent replicates. A two-tailed independent Student's t-test was used to determine the differences between the two groups. One-way ANOVA, followed by Tukey multiple comparison, was used to determine the significance of differences in more than two groups. $P < 0.05$ was used as a threshold for statistically significant differences.

Data availability

All data generated or analyzed during this study are included in the manuscript and its supporting files. Materials generated in this study are available from the corresponding author upon reasonable request.

Acknowledgements

This research was funded by the National Key Research and Development Program of China (Grant No. 2022YFD1201600), the National Natural Science Foundation of China (Grant No. 32172969, 31972776, 32102780, 32473159).

Author contributions

Changle Zhao: Writing – original draft, Software, Resources, Methodology, Investigation, Formal analysis, Data curation. **Yongxun Chen**: Writing – original draft, Methodology, Data curation. **Lei Liu**: Data curation, Conceptualization. **Xiang Liu**: Software, Investigation. **Hesheng Xiao**: Software, Data curation. **Feilong Wang**: Investigation, Formal analysis. **Qin Huang**: Resources, Software. **Xiangyan Dai**: Resources, Investigation. **Wenjing Tao**: Resources, Funding acquisition. **Deshou**

Wang: Supervision, Investigation, Funding acquisition. **Jing Wei**: Writing – review & editing, Validation, Supervision, Project administration, Funding acquisition, Data curation, Conceptualization.

Conflict of interest

The authors declare no competing interest.

References

Barsoum, I. B., Bingham, N. C., Parker, K. L., Jorgensen, J. S., & Yao, H. H. 2009. Activation of the Hedgehog pathway in the mouse fetal ovary leads to ectopic appearance of fetal Leydig cells and female pseudohermaphroditism. *Dev Biol, 329*(1), 96-103. doi:10.1016/j.ydbio.2009.02.025

Barsoum, I., & Yao, H. H. (2011). Redundant and differential roles of transcription factors Gli1 and Gli2 in the development of mouse fetal Leydig cells. *Biol Reprod, 84*(5), 894-899. doi:10.1095/biolreprod.110.088997

Belgacem, Y. H., & Borodinsky, L. N. (2015). Inversion of Sonic hedgehog action on its canonical pathway by electrical activity. *Proc Natl Acad Sci U S A, 112*(13), 4140-4145. doi:10.1073/pnas.1419690112

Briscoe, J., & Thérond, P. P. (2013). The mechanisms of Hedgehog signalling and its roles in development and disease. *Nat Rev Mol Cell Biol, 14*(7), 416-429. doi:10.1038/nrm3598

Carpenter, D., Stone, D. M., Brush, J., Ryan, A., Armanini, M., Frantz, G., . . . de Sauvage, F. J. (1998). Characterization of two patched receptors for the vertebrate hedgehog protein family. *Proc Natl Acad Sci U S A, 95*(23), 13630-13634. doi:10.1073/pnas.95.23.13630

Cervantes, S., Lau, J., Cano, D. A., Borromeo-Austin, C., & Hebrok, M. (2010). Primary cilia regulate Gli/Hedgehog activation in pancreas. *Proc Natl Acad Sci U S A, 107*(22), 10109-10114. doi:10.1073/pnas.0909900107

Chen, H., Wang, Y., Ge, R., & Zirkin, B. R. (2017). Leydig cell stem cells: Identification, proliferation and differentiation. *Mol Cell Endocrinol, 445*, 65-73. doi:10.1016/j.mce.2016.10.010

Chen, P., Zirkin, B. R., & Chen, H. (2020). Stem Leydig Cells in the Adult Testis: Characterization, Regulation and Potential Applications. *Endocr Rev, 41*(1), 22-32. doi:10.1210/endrev/bnz013

Cong, G., Zhu, X., Chen, X. R., Chen, H., & Chong, W. (2025). Mechanisms and therapeutic potential of the hedgehog signaling pathway in cancer. *Cell Death Discov, 11*(1), 40. doi:10.1038/s41420-025-02327-w

Dai, S., Qi, S., Wei, X., Liu, X., Li, Y., Zhou, X., . . . Li, M. (2021). Germline sexual fate is determined by the antagonistic action of dmrt1 and foxl3/foxl2 in tilapia. *Development, 148*(8). doi:10.1242/dev.199380

Finco, I., LaPensee, C. R., Krill, K. T., & Hammer, G. D. (2015). Hedgehog signaling and steroidogenesis. *Annu Rev Physiol, 77*, 105-129. doi:10.1146/annurev-physiol-061214-111754

Fleet, A. J., & Hamel, P. A. (2018). The protein-specific activities of the transmembrane modules of Ptch1 and Ptch2 are determined by their adjacent protein domains. *J Biol Chem, 293*(43), 16583-16595. doi:10.1074/jbc.RA118.004478

Ge, R. S., Dong, Q., Sottas, C. M., Papadopoulos, V., Zirkin, B. R., & Hardy, M. P. (2006). In search of rat
stem Leydig cells: identification, isolation, and lineage-specific development. *Proc Natl Acad*
Sci U S A, 103(8), 2719-2724. doi:10.1073/pnas.0507692103

Houzelstein, D., Eozenou, C., Lagos, C. F., Elzaiat, M., Bignon-Topalovic, J., Gonzalez, I., . . . McElreavey,
K. (2024). A conserved NR5A1-responsive enhancer regulates SRY in testis-determination.
Nat Commun, 15(1), 2796. doi:10.1038/s41467-024-47162-2

Huang, Q., Yang, Z., Wang, J., Luo, Y., Zhao, C., Li, M., . . . Wei, J. (2020). Establishment of a stem Leydig
cell line capable of 11-ketotestosterone production. *Reprod Fertil Dev, 32*(16), 1271-1281.
doi:10.1071/rd20171

Huangfu, D., & Anderson, K. V. (2006). Signaling from Smo to Ci/Gli: conservation and divergence of
Hedgehog pathways from Drosophila to vertebrates. *Development, 133*(1), 3-14.
doi:10.1242/dev.02169

Hui, C. C., & Angers, S. (2011). Gli proteins in development and disease. *Annu Rev Cell Dev Biol, 27,*
513-537. doi:10.1146/annurev-cellbio-092910-154048

Ingham, P. W., Nakano, Y., & Seger, C. (2011). Mechanisms and functions of Hedgehog signalling across
the metazoa. *Nat Rev Genet, 12*(6), 393-406. doi:10.1038/nrg2984

Inoue, M., Baba, T., & Morohashi, K. I. (2018). Recent progress in understanding the mechanisms of
Leydig cell differentiation. *Mol Cell Endocrinol, 468,* 39-46. doi:10.1016/j.mce.2017.12.013

Jiang, M. H., Cai, B., Tuo, Y., Wang, J., Zang, Z. J., Tu, X., . . . Xiang, A. P. (2014). Characterization of
Nestin-positive stem Leydig cells as a potential source for the treatment of testicular Leydig
cell dysfunction. *Cell Res, 24*(12), 1466-1485. doi:10.1038/cr.2014.149

Kim, Y., Lee, J., Seppala, M., Cobourne, M. T., & Kim, S. H. (2020). Ptch2/Gas1 and Ptch1/Boc
differentially regulate Hedgehog signalling in murine primordial germ cell migration. *Nat*
Commun, 11(1), 1994. doi:10.1038/s41467-020-15897-3

Kinzler, K. W., Ruppert, J. M., Bigner, S. H., & Vogelstein, B. (1988). The GLI gene is a member of the
Kruppel family of zinc finger proteins. *Nature, 332*(6162), 371-374. doi:10.1038/332371a0

Kothandapani, A., Lewis, S. R., Noel, J. L., Zacharski, A., Krellwitz, K., Baines, A., . . . Jorgensen, J. S.
(2020). GLI3 resides at the intersection of hedgehog and androgen action to promote male
sex differentiation. *PLoS Genet, 16*(6), e1008810. doi:10.1371/journal.pgen.1008810

Li, M., Sun, L., Zhou, L., & Wang, D. (2024). Tilapia, a good model for studying reproductive
endocrinology. *Gen Comp Endocrinol, 345,* 114395. doi:10.1016/j.ygcen.2023.114395

Li, M., Yang, H., Zhao, J., Fang, L., Shi, H., Li, M., . . . Wang, D. (2014). Efficient and heritable gene
targeting in tilapia by CRISPR/Cas9. *Genetics, 197*(2), 591-599.
doi:10.1534/genetics.114.163667

Liu, X., Zhao, C., Liu, L., Peng, X., Li, J., Tao, W., . . . Wei, J. (2024). Nile Tilapia (Oreochromis niloticus)
Patched1 Mutations Disrupt Cardiovascular Development and Vascular Integrity through
Smoothened Signaling. *Int J Mol Sci, 25*(6). doi:10.3390/ijms25063321

Mehta, P., Singh, P., Gupta, N. J., Sankhwar, S. N., Chakravarty, B., Thangaraj, K., & Rajender, S. (2021).
Mutations in the desert hedgehog (DHH) gene in the disorders of sexual differentiation and
male infertility. *J Assist Reprod Genet, 38*(7), 1871-1878. doi:10.1007/s10815-021-02140-1

Nieuwenhuis, E., Motoyama, J., Barnfield, P. C., Yoshikawa, Y., Zhang, X., Mo, R., . . . Hui, C. C. (2006).
Mice with a targeted mutation of patched2 are viable but develop alopecia and epidermal
hyperplasia. *Mol Cell Biol, 26*(17), 6609-6622. doi:10.1128/mcb.00295-06

Niewiadomski, P., Kong, J. H., Ahrends, R., Ma, Y., Humke, E. W., Khan, S., . . . Rohatgi, R. (2014). Gli

protein activity is controlled by multisite phosphorylation in vertebrate Hedgehog signaling.

Cell Rep, 6(1), 168-181. doi:10.1016/j.celrep.2013.12.003

Niewiadomski, P., Niedziółka, S. M., Markiewicz, Ł, Uśpieński, T., Baran, B., & Chojnowska, K. (2019).

Gli Proteins: Regulation in Development and Cancer. *Cells, 8*(2). doi:10.3390/cells8020147

Pachernegg, S., Georges, E., & Ayers, K. (2022). The Desert Hedgehog Signalling Pathway in Human

Gonadal Development and Differences of Sex Development. *Sex Dev, 16*(2-3), 98-111.

doi:10.1159/000518308

Pan, Y., & Wang, B. (2007). A novel protein-processing domain in Gli2 and Gli3 differentially blocks

complete protein degradation by the proteasome. *J Biol Chem, 282*(15), 10846-10852.

doi:10.1074/jbc.M608599200

Park, S. Y., Tong, M., & Jameson, J. L. (2007). Distinct roles for steroidogenic factor 1 and desert

hedgehog pathways in fetal and adult Leydig cell development. *Endocrinology, 148*(8), 3704-

3710. doi:10.1210/en.2006-1731

Parker, K. L., Rice, D. A., Lala, D. S., Ikeda, Y., Luo, X., Wong, M., . . . Schimmer, B. P. (2002).

Steroidogenic factor 1: an essential mediator of endocrine development. *Recent Prog Horm*

Res, 57, 19-36. doi:10.1210/rp.57.1.19

Pierucci-Alves, F., Clark, A. M., & Russell, L. D. (2001). A developmental study of the Desert hedgehog-

null mouse testis. *Biol Reprod, 65*(5), 1392-1402. doi:10.1095/biolreprod65.5.1392

Qi, X., Schmiege, P., Coutavas, E., & Li, X. (2018). Two Patched molecules engage distinct sites on

Hedgehog yielding a signaling-competent complex. *Science, 362*(6410).

doi:10.1126/science.aas8843

Ruiz i Altaba, A., Sánchez, P., & Dahmane, N. (2002). Gli and hedgehog in cancer: tumours, embryos

and stem cells. *Nat Rev Cancer, 2*(5), 361-372. doi:10.1038/nrc796

Smith, O. E., Morin, F., Roussel, V., Bertucci, M. C., Boyer, A., & Murphy, B. D. (2023). The role of

steroidogenic factor 1 (SF-1) in steroidogenic cell function of the testes and ovaries of mature

mice. *Reproduction, 165*(1), 1-17. doi:10.1530/rep-22-0049

Sun, Yun-Lv, Jiang, Dong-Neng, Zeng, Sheng, Hu, Chong-Jiang, Ye, Kai, Yang, Chao, . . . Wang, De-Shou.

(2014). Screening and characterization of sex-linked DNA markers and marker-assisted

selection in the Nile tilapia (*Oreochromis niloticus*). *Aquaculture, 433*, 19-27.

doi:10.1016/j.aquaculture.2014.05.035

Tao, W., Chen, J., Tan, D., Yang, J., Sun, L., Wei, J., . . . Wang, D. (2018). Transcriptome display during

tilapia sex determination and differentiation as revealed by RNA-Seq analysis. *BMC Genomics,*

19(1), 363. doi:10.1186/s12864-018-4756-0

Tao, W., Yuan, J., Zhou, L., Sun, L., Sun, Y., Yang, S., . . . Wang, D. (2013). Characterization of gonadal

transcriptomes from Nile tilapia (*Oreochromis niloticus*) reveals differentially expressed genes.

PLoS One, 8(5), e63604. doi:10.1371/journal.pone.0063604

Umehara, F., Tate, G., Itoh, K., Yamaguchi, N., Douchi, T., Mitsuya, T., & Osame, M. (2000). A novel

mutation of desert hedgehog in a patient with 46,XY partial gonadal dysgenesis accompanied

by minifascicular neuropathy. *Am J Hum Genet, 67*(5), 1302-1305. doi:10.1016/s0002-

9297(07)62958-9

Wei, J., Wu, J., Ru, W., Chen, G., Gao, L., & Tang, D. (2022). Novel compound heterozygous mutations

in the desert hedgehog (DHH) gene in cases of siblings with 46,XY disorders of sexual

development. *BMC Med Genomics, 15*(1), 178. doi:10.1186/s12920-022-01334-5

Xie, Q. P., He, X., Sui, Y. N., Chen, L. L., Sun, L. N., & Wang, D. S. (2016). Haploinsufficiency of SF-1

653   Causes Female to Male Sex Reversal in Nile Tilapia, Oreochromis niloticus. *Endocrinology,*

654   *157*(6), 2500-2514. doi:10.1210/en.2015-2049

655   Yao, H. H., Whoriskey, W., & Capel, B. (2002). Desert Hedgehog/Patched 1 signaling specifies fetal

656   Leydig cell fate in testis organogenesis. *Genes Dev, 16*(11), 1433-1440.

657   doi:10.1101/gad.981202

658   Yao, S., Wei, X., Deng, W., Wang, B., Cai, J., Huang, Y., . . . Wang, J. (2022). Nestin-dependent

659   mitochondria-ER contacts define stem Leydig cell differentiation to attenuate male

660   reproductive ageing. *Nat Commun, 13*(1), 4020. doi:10.1038/s41467-022-31755-w

661   Yu, Y., Lee, P. A., Huerta-Saenz, L., & Allen, N. G. (2023). A Novel Variant in NR5A1 Presenting as 46,XY

662   Difference of Sex Development. *JCEM Case Rep, 1*(5), luad103. doi:10.1210/jcemcr/luad103

663   Zhang, Y., & Beachy, P. A. (2023). Cellular and molecular mechanisms of Hedgehog signalling. *Nat Rev*

664   *Mol Cell Biol, 24*(9), 668-687. doi:10.1038/s41580-023-00591-1

665   Zhang, Z., Wang, J., Li, J., Liu, X., Liu, L., Zhao, C., . . . Wei, J. (2023). Establishment of an Integrated

666   CRISPR/Cas9 Plasmid System for Simple and Efficient Genome Editing in Medaka In Vitro and

667   In Vivo. *Biology (Basel), 12*(2). doi:10.3390/biology12020336

668   Zhao, C., Liu, X., Liu, L., Li, J., Liu, X., Tao, W., . . . Wei, J. (2023). Smoothened mediates medaka

669   spermatogonia proliferation via Gli1-Rgcc-Cdk1 axis†. *Biol Reprod, 109*(5), 772-784.

670   doi:10.1093/biolre/ioad090

671   Zheng, Q., Xiao, H., Shi, H., Wang, T., Sun, L., Tao, W., . . . Wang, D. (2020). Loss of Cyp11c1 causes

672   delayed spermatogenesis due to the absence of 11-ketotestosterone. *J Endocrinol, 244*(3),

673   487-499. doi:10.1530/joe-19-0438

Figure legends

Fig. 1. Mutation of *dhh* resulted in testicular developmental disorders.

(A-B) Morphological analysis of the testes from WT (*dhh*$^{+/+}$) and *dhh*$^{-/-}$ XY fish at 90

dah. Arrows indicate the location of the testes. (C-D') Histological analysis of

testicular sections from WT and *dhh*$^{-/-}$ XY fish at 90 dah. Sg, spermatogonia; Sc,

spermatocyte. (E-J') Representative immunofluorescence images showing the

expression of germ cell marker Vasa (E-F'), meiosis cell marker Sycp3 (G-H') and

Leydig cell marker Cyp11c1 (I-J'). (K) Gonadosomatic index (GSI) of the testes from

WT and *dhh*$^{-/-}$ XY fish at 90 dah (*n*=5 fish/genotype). (L-N) Quantification of the

percentage of cells positive for Vasa (E-F'), Sycp3 (G-H') and Cyp11c1 (I-J') among

all DAPI-positive cells in the WT and *dhh*$^{-/-}$ testes (*n*=5 fish/genotype). (O) Serum 11-

KT level of WT and *dhh*$^{-/-}$ XY fish at 90 dah (*n*=6 fish/genotype). Values were

presented as mean ± SD. Differences were determined by two-tailed independent

Student's t-test. **, $P < 0.01$. Scale bars, (A-B), 1 cm; (C-J), 50 μm.

Fig. 2. Rescue of testicular development in *dhh*$^{-/-}$ XY Nile tilapia by 11-KT and

SAG.

dhh^{-/-} XY Nile tilapias at 30 dah were treated either with 11-KT via water immersion

(with water changed every two days) or with 10 mg/kg SAG via intraperitoneal

injection (with supplemental injection every seven days). WT controls received 11-KT,

and *dhh*^{-/-} controls received an equivalent volume of the DMSO. Then, at 90 dah,

morphological and histological experiments were conducted. (A-D') Histological

analysis of testicular sections from 90 dah XY fish subjected to different treatments as

indicated. Sg, spermatogonia; Sc, spermatocyte. Scale bars, 50 μm. (E-P')

Representative immunofluorescence images showing the expression of Vasa (E-H'),

Sycp3 (I-L') and Cyp11c1 (M-P'). Scale bars, 50 μm. (Q) GSI of testes from the

different treatment groups at 90 dah (n=5 fish per group). (R-T) Quantification of the

percentage of cells positive for Vasa (E-H'), Sycp3 (I-L') and Cyp11c1 (M-P') among

all DAPI-positive cells in the testes (n=3 ∼ 5 fish per group). Values were presented as

mean ± SD. Different letters above the error bar indicate statistical differences at $P <$

0.05 as determined by one-way ANOVA followed by Tukey test.

Fig. 3. Dhh is required for SLCs differentiation *in vivo*.

TSL, Dhh-overexpressing TSL (TSL-*On*Dhh), or 0.5 μM SAG-treated TSL

(TSL+SAG) cells were labeled with PKH26 and transplanted into the testes of 90

dah WT or *dhh*^{-/-} recipient fish. Analyses were performed 10 days post-transplantation.

(A) Schematic diagram of the experimental design for SLCs transplantation and

analysis. (B1-E3) Representative immunofluorescence images of testicular sections

from recipient fish, showing the localization of transplanted PKH26-labeled SLCs

(red) and the expression of Cyp11c1 (green). Nuclei are stained with DAPI (blue).

Scale bars, 4 μm. (F) Quantification of the percentage of Cyp11c1-positive cells

among the transplanted PKH26-positive SLCs for each treatment group (n=5 fish per

group). (G) Serum 11-KT level in recipient fish following SLCs transplantation (n=5

fish per group). Values were presented as mean ± SD. Different letters above the error

bar indicate statistical differences at $P < 0.05$ as determined by one-way ANOVA

followed by Tukey test.

Fig. 4. Ptch2 mediates Dhh signaling in SLCs.

(A) Co-localization of Cyp11c1 (green, by immunofluorescence)

and *ptch1* or *ptch2* mRNA (red, by RNA-FISH) in adult (90 dah) testis sections.

Dashed lines outline representative Leydig cells. Scale bars, 4 μm. (B) Schematic

illustration of the experimental setup for the luciferase reporter assays shown in

panels C and D. (C) Luciferase activity in TSL cells co-transfected with a Gli-

responsive reporter (8xGLI) and overexpression of tilapia Dhh (*On*Dhh). A plasmid

lacking Gli-binding sites (pGL4.23) served as a negative control (*n*=3). (D) Luciferase

activity in TSL-WT, TSL-*ptch1*$^{-/-}$ and TSL-*ptch2*$^{-/-}$ cells transfected with the 8xGLI

reporter with or without *On*Dhh (*n*=4). (E-G') Histological analysis of testis sections

from the indicated genotypes at 90 dah. Sg, spermatogonia; Sc, spermatocyte. Scale

bars, 50 μm. (H-P') Immunofluorescence analysis of Vasa (H-J'), Sycp3 (K-M') and

Cyp11c1 (N-P') in testis sections from the indicated genotypes. Scale bars, 50 μm.

(Q) GSI of the testes from the indicated genotypes at 90 dah (*n*=4 fish per group).

(R-T) Quantification of the percentage of Vasa (H-J'), Sycp3 (K-M') and Cyp11c1

(N-P') positive cells among DAPI-positive cells (*n*=6 fish per group). (U) Serum 11-

KT levels in fish of the indicated genotypes at 90 dah (*n*=6 fish per group). Statistical

significance was determined by one-way ANOVA followed by Tukey's test (C, Q-U,

different letters above the error bar indicate statistical differences at $P < 0.05$) or

Student's *t*-test (D) (*, $P < 0.05$; **, $P < 0.01$; NS, no significant difference).

Fig. 5. Gli1 transactivates *sf1* to drive SLC differentiation.

(A) Co-localization of Cyp11c1 (green) and *gli1*, *gli2*, or *gli3* mRNA (red) in adult

(90 dah) testis sections by immunofluorescence and RNA-FISH. Dashed lines outline

representative Leydig cells. Scale bars, 4 μm. (B) Luciferase activity in TSL-WT,

TSL-*gli1*$^{-/-}$, TSL-*gli2*$^{-/-}$ and TSL-*gli3*$^{-/-}$ cells transfected with the 8xGLI reporter with

or without *On*Dhh (*n*=4). (C) Volcano plots of transcriptomic changes in TSL-*On*Dhh,

TSL-*On*Gli1 and TSL+ SAG, compared to TSL-WT. Red and blue dots represent

significantly up- and down-regulated genes, respectively. (D) Schematic of the

tilapia *sf1* gene promoter, indicating the two predicted Gli1 binding sites (B1 and B2).

(E) Transcriptional activation of the *sf1* promoter by Gli1. HEK293 cells were co-

transfected with pRL-TK, pGL3, pcDNA3.1, pGL3-*sf1*, pcDNA3.1-*On*Gli1, and the

indicated cold probe constructs, and luciferase activity was measured 48 hours post-

transfection. Competition assays were performed using unlabeled cold probe (Cold

probe, GACCACCCA, 10/100/250 ng/mL) or mutant unlabeled cold probe (mutant

Cold probe, TTAATTAAA, 10/100/250 ng/mL) (*n*=4). "+" indicates the addition of

the corresponding substance, while "-" indicates no addition, and the number

represents the amount added (ng/ml). (F-G) Representative immunofluorescence

images of testicular sections from recipient fish transplanted with *sf1*-deficient (TSL-*sf1*[-/-], F) or Sf1-overexpressing (TSL-*On*Sf1, G) SLCs, stained for Cyp11c1 (green) and PKH26 (red). Nuclei are stained with DAPI (blue). Scale bars, 4 μm. (H) Quantification of the percentage of Cyp11c1-positive cells among the transplanted PKH26-positive SLCs in the two transplantation groups (*n*=5 fish per group). Statistical significance was determined by one-way ANOVA followed by Tukey's test (E, different letters above the error bar indicate statistical differences at $P < 0.05$) or Student's *t*-test (B, H) (*, $P < 0.05$; **, $P < 0.01$; NS, no significant difference).

Fig. S1. Sequence analysis and mRNA expression profile of Nile tilapia *dhh*.

(A) Dhh amino acid sequence alignment. Amino acids are numbered in the right margin. Deletions are indicated by dashes, shaded areas indicate shared sequences. The amino-terminal hedge domain is underlined by dotted line, with the signal peptide region in the front and the autocatalytic carboxy-terminal domain (hog) in the back. The box indicates the autocatalytic site of an absolutely conserved Gly-Cys-Phe tripeptide. At the end of the alignment are percentage identity values of the full-length and hedge domain of tilapia Dhh to the orthologue from other species. (B) Phylogenetic analysis of Dhh. The phylogenetic tree was constructed using the neighbor-joining method within the MEGA7.0 program. Node values represent percent bootstrap confidence derived from 1000 replicates. (C-D) The RPKM (reads per kb per million reads) values of tilapia *dhh* in various adult tissues (C) and XY and XX gonads at 5, 7, 20, 30, 90, 180, 300 dah (days after hatching) (D) in transcriptome sequencing data which were sequenced using Illumina 2000 HiSeq technology in our previous study (Tao et al., 2013).

Fig. S2. Establishment of the Nile tilapia *dhh* mutant line by CRISPR/Cas9.

(A) Schematic representation of gRNA targeting the Nile tilapia *dhh* locus. The gRNA was designed to target exon 1. The translation start codon ATG and stop codon TAA were indicated by arrows. The PAM (protospacer adjacent motif) site was marked by green box. DNA sequence alignment of the three mutant lines with WT. The added sequences were marked by red letter, and the deleted sequences were indicated in red dotted lines. One had a 4-bp addition, one had an 8-bp deletion, and the other had a 13-bp deletion. (B) Schematic diagram showing the breeding plans of *dhh* F0 to F2 fish. (C) Homozygous mutants of F2 fish were screened by PAGE. The first lane is DNA marker, the second lane is *dhh*[+/+] fish, the third lane is *dhh*[+/-] fish, and the fourth

lane is *dhh*$^{-/-}$ fish. (D) Sanger sequencing results of *dhh* genes from WT and the 13-bp

deletion homozygous mutant fishes. (E) The *dhh* ORF sequence in WT was indicated

in black and the frameshift-altered sequence for the 13-bp deletion *dhh* ORF was

indicated in red. (F) The relative mRNA expression of *dhh* in WT and the 13-bp

deletion homozygous mutant fishes by RT-qPCR. Values were presented as mean ±

SD (*n*=3). Differences were determined by two-tailed independent Student's t-test. **,

P < 0.01.

Fig. S3. RT-PCR analyses of Hh pathway genes in TSL cells. *β-actin* served as a

loading control, and RNAs from TSL cells were used as templates for negative

controls. The numbers in parentheses indicate the PCR cycles.

Fig. S4. Establishment of the TSL *ptch1*, *ptch2*, *gli1*, *gli2*, *gli3* and *sf1* mutant

lines.

(A-F) Schematic representation of gRNAs targeting the *ptch1*, *ptch2*, *gli1*, *gli2*, *gli3*

and *sf1* locus. The gRNAs were designed to target ORF (open reading frame). The

PAM (protospacer adjacent motif) sites were marked by red letter. Sanger sequencing

of *ptch1*, *ptch2*, *gli1*, *gli2*, *gli3* and *sf1* single mutant alleles in single cell clones. PCR

amplicons from DNA templates of the six cell clones were directly used for

sequencing.

Fig. S5. Establishment of the Nile tilapia *ptch2* mutant line by CRISPR/Cas9

system.

(A) Schematic representation of gRNA targeting the Nile tilapia *ptch2* locus. The

gRNA was designed to target exon 1. The translation start codon ATG and stop codon

TGA were indicated by arrows. The PAM site was marked by green box. DNA

sequence alignment of the five mutant lines with WT. The added sequences were

marked by red letter, and the deleted sequences were indicated in red dotted lines. (B)

Schematic diagram showing the breeding plans of *ptch2* F0 to F2 fish. (C) DNA

sequencing showed the 25-bp deletion within the *ptch2* ORF in the homozygous

mutant compared to the WT. (D) Schematic of the prediction of the intact Ptch2

protein in the WT and the truncated Ptch2 protein in the homozygous mutants. (E)

Homozygous mutants of F2 fish were screened by PAGE. The first lane is DNA

marker, the second lane is *ptch2*$^{+/+}$ fish, the third lane is *ptch2*$^{+/-}$ fish, and the fourth

lane is *ptch2*$^{-/-}$ fish. "*" indicates heteroduplex, and arrows indicate homoduplex.

Fig. S6. Transcriptome data analyses and verification by RT-qPCR.

(A) Hierarchical clustering analysis of global gene expression patterns in TSL-WT,

TSL-*On*Dhh, TSL-*On*Gli1 and TSL+ 0.5 μM SAG. Blue indicates decreased

expression, and red indicates increased expression. (B) Heatmap showing the

expression patterns of upregulated differentially expressed genes (DEGs) identified in

Fig. 5C. The FPKM value for each gene in each sample is indicated within the

squares. The color gradient from blue to red reflects low to high expression levels per

row (gene). (C) qRT-PCR validation. Values were presented as mean ± SD (n = 3).

Fig. S7. The expression levels of Cyp11c1 in the testis and the levels of 11-KT in

tissue fluid from WT and *dhh*$^{-/-}$ XY fish at 5, 10, 20, and 30 dah.

(A-E) Representative immunofluorescence images showing Cyp11c1 expression

(green) in testis sections from WT fish at 5, 10, 20, 30 dah and *dhh*$^{-/-}$ mutants at 30

dah. Scale bars, 4 μm. (F) Tissue fluid 11-KT levels in WT and *dhh*$^{-/-}$ XY fish at

indicated time points (*n*=6 fish per group). Values were presented as mean ± SD.

Differences were determined by two-tailed independent Student's t-test. NS, no

significant difference.

Fig. S8. Expression analysis of *ptch1* in WT and mutant XY testes.

Relative mRNA expression levels of *ptch1* in XY testes from WT, *ptch2*$^{-/-}$ and *dhh*$^{-/-}$

;*ptch2*$^{-/-}$ fish at 90 dah by RT-qPCR. Values were presented as mean ± SD (*n*=3 fish

per group). Statistical significance was determined by one-way ANOVA followed by

Tukey's test. NS, no significant difference.

Fig. S9. Validation of key gene expression changes in mutant and overexpression

TSL cell lines. (only for reviewer).

(A) Relative mRNA expression levels of *sf1* in TSL-WT, TSL-*gli1*$^{-/-}$, TSL-*gli2*$^{-/-}$ and

TSL-*gli3*$^{-/-}$ cells by RT-qPCR. Values were presented as mean ± SD (*n*=3). Different

letters above the error bar indicate statistical differences at *P* < 0.05 as determined by

one-way ANOVA followed by Tukey test. (B) Relative mRNA expression of *dhh* in

TSL-WT and TSL-*On*Dhh cells by RT-qPCR. Values were presented as mean ± SD

(*n*=3). Differences were determined by two-tailed independent Student's t-test. **, *P*

< 0.01. (C) Relative mRNA expression of *sf1* in TSL-WT and TSL-*On*Sf1 cells by

RT-qPCR. Values were presented as mean ± SD (*n*=3). Differences were determined

by two-tailed independent Student's t-test. **, *P* < 0.01.

Fig. S10. Negative control experiments for RNA fluorescence in situ

hybridization (FISH). (only for reviewer)

Co-staining of Cyp11c1 (by immunofluorescence) and sense RNA probes

for *ptch1*, *ptch2*, *gli1*, *gli2*, or *gli3* (by FISH) on testis sections from 90 dah fish. The

sense probes serve as negative controls to confirm the specificity of the antisense

probe hybridization signals shown in the main figures. Scale bars, 4 μm.

| | WT | *dhh⁻/⁻* | | |
| | 11-KT | DMSO | 11-KT | SAG |

H.E.

A A' Sg Sc   B B' Sg   C C' Sg Sc   D D' Sg Sc

Vasa/DAPI

E E'   F F'   G G'   H H'

Sycp3/DAPI

I I'   J J'   K K'   L L'

Cyp11c1/DAPI

M M'   N N'   O O'   P P'

Q
GSI (%)
0.20
0.15 c
0.10   b   b
0.05   a
0.00
WT +11-KT   *dhh⁻/⁻* +DMSO   *dhh⁻/⁻* +11-KT   *dhh⁻/⁻* +SAG

R
Vasa⁺ / DAPI⁺ (%)
100
80
60   c
40   b   b
20   a
0
WT +11-KT   *dhh⁻/⁻* +DMSO   *dhh⁻/⁻* +11-KT   *dhh⁻/⁻* +SAG

S
Sycp3⁺ / DAPI⁺ (%)
30
c
20   b   b
10
a
0
WT +11-KT   *dhh⁻/⁻* +DMSO   *dhh⁻/⁻* +11-KT   *dhh⁻/⁻* +SAG

T
Cyp11c1⁺ / DAPI⁺ (%)
20
15 c
10   b
5   a   a
0
WT +11-KT   *dhh⁻/⁻* +DMSO   *dhh⁻/⁻* +11-KT   *dhh⁻/⁻* +SAG