Desert Hedgehog mediates stem Leydig cell differentiation through Ptch2/Gli1/Sf1 signaling axis

Changle Zhao, Yongxun Chen, Lei Liu, Xiang Liu, Hesheng Xiao, Feilong Wang, Qin Huang, Xiangyan Dai, Wenjing Tao, Deshou Wang[*], Jing Wei[*]

Author affiliations: Integrative Science Center of Germplasm Creation in Western China (CHONGQING) Science City, Key Laboratory of Freshwater Fish Reproduction and Development (Ministry of Education), Laboratory of Aquatic Science of Chongqing, School of Life Sciences, Southwest University, Chongqing 400715, China

Prof. Deshou Wang, E-mail address: wdeshou@swu.edu.cn. Tel.: (86) 023 68252702; Fax: (86) 023 68253005.

Prof. Jing Wei, E-mail address: lalsos@swu.edu.cn. Tel.: (86) 023 68252365; Fax: (86) 023 68253005.

Abstract

Desert Hedgehog (Dhh) mutations cause Leydig cell dysfunction, yet the mechanisms governing Leydig lineage commitment through Dhh-mediated receptor selectivity, transcriptional effector specificity, and steroidogenic coupling remain elusive. In this study, using CRISPR/Cas9-mediated gene knockout and stem Leydig cells (SLCs) transplantation, we identified a critical Dhh/Patched 2 (Ptch2)/Glioma-associated oncogene homolog 1 (Gli1)/steroidogenic factor 1 (Sf1) signaling axis essential for SLC differentiation in Nile tilapia *(Oreochromis niloticus)*. Dhh deficiency resulted in defective adult Leydig cells and androgen insufficiency. Rescue experiments involving 11-ketotestosterone administration and a Dhh agonist treatment, combined with SLCs transplantation, demonstrated that Dhh regulates SLC differentiation, not survival. *In vitro* knockout of *ptch1* and *ptch2* in SLCs revealed that Ptch2 likely acts as the functional receptor for Dhh. This was further supported by *in vivo* genetic rescue experiments, where *ptch2* mutation did not impair testicular development, yet completely rescued the testicular defects in *dhh* mutants—consistent with Ptch2 acting as an inhibitory receptor whose loss alleviates Dhh pathway suppression. Luciferase assays in Gli-knockout SLCs demonstrated that Gli1 acts as the primary transcriptional effector, and transactivates *sf1* expression. Additionally, functional transplantation assays confirmed that Sf1 is indispensable for SLCs differentiation, as Sf1-overexpressing SLCs rescued differentiation, whereas *sf1*-mutant SLCs failed. Overall, our work delineates the Dhh-Ptch2-Gli1-Sf1 axis and provides fundamental insights into the endocrine regulation of Leydig cell lineage development.

Keywords: Desert Hedgehog (Dhh), stem Leydig cell, Patched 2 (Ptch2), Glioma-associated oncogene homolog 1 (Gli1), Steroidogenic factor 1 (Sf1)

Highlights

- Dhh regulates the differentiation of SLCs rather than their survival
- Ptch2 acts as the functional receptor for Dhh signaling in SLCs
- Gli1 is the principal transcriptional activator for Dhh signaling in SLCs
- Sf1 is the critical downstream effector of Gli1 in SLCs differentiation

INTRODUCTION

Leydig cells are the primary source of androgens, producing approximately 95% of circulating testosterone to sustain spermatogenesis and male fertility [1, 2]. Leydig cells originate from stem Leydig cells (SLCs), which differentiate into steroidogenic adult Leydig cells (ALCs) under paracrine regulation [3, 4]. Steroidogenic factor 1 (Sf1), an orphan nuclear receptor transcription factor, serves as a master regulator of this process by activating steroidogenic enzymes (e.g., cytochrome P450, family 11, subfamily C, polypeptide 1 [Cyp11c1], 3β-hydroxysteroid dehydrogenase [3β-HSD]) to drive ALC maturation [5]. In mice, mutations of *Sf1* exhibit arrested testicular development and abolished steroidogenesis [6-8], while human *SF1* mutations are associated with 46,XY disorders of sexual development (DSD), such as Swyer syndrome [9]. Similarly, in Nile tilapia (*Oreochromis niloticus*), our previous work reveals that *sf1* mutation disrupts testicular morphogenesis and downregulates the expression of key steroidogenic enzymes such as *cyp11c1* [10]. Despite its established central role, the upstream signaling pathways that regulate Sf1 to commit the Leydig lineage remain poorly defined.

Hedgehog (Hh) signaling pathway is an evolutionarily conserved developmental regulator [11]. Canonical Hh signaling involves the binding of ligands to Patched (Ptch) receptors, which relieves the suppression on Smoothened (Smo) and ultimately activates Glioma-associated oncogene homolog (Gli) transcription factors to regulate target genes [12-14]. In vertebrates, there are three Hh ligands (i.e. Sonic Hh, Indian Hh and Desert Hh [Dhh]), two Ptch receptors (Ptch1 and Ptch2) and three Gli homologs (Gli1-3) [15-18]. A large number of studies have well demonstrated that Dhh signaling pathway is critical in gonadal development [14, 19]. Dhh, secreted by Sertoli cells, is critical for Leydig cell commitment in mammals [20, 21]. In mice, mutations of *Dhh* lead to Leydig cell dysfunction, testosterone insufficiency, and infertility [22], while human *DHH* mutations are associated with 46,XY DSD [23]. Strikingly, these phenotypes mirror those caused by *Sf1* deficiencies, suggesting a potential regulatory nexus between Dhh and Sf1 signaling. However, the molecular circuitry connecting Dhh signaling to Sf1 activation remains elusive.

Studies demonstrate functional divergence between Ptch1 and Ptch2. For instance, *Ptch1* knockout causes embryonic lethality due to constitutive Hh activation, whereas *Ptch2* deficiency yields no overt defects [24, 25]. During testicular development, Ptch1 displays high expression at early stage critical for gonadal primordium formation, while Ptch2 shows testis-enriched dynamic regulation [26, 27]. However, the distinct roles of Ptch1 and Ptch2 in Leydig lineage commitment remain unknown to date. Among Gli transcription factors, Gli1 acts as a primary activator, while Gli2 and Gli3 exhibit dual regulatory functions [28, 29]. In murine testes, co-expression of Gli1 and Gli2 in Leydig cells illustrates functional redundancy—single knockouts show normal development, but combined pharmacological inhibition (GANT61) abolishes Leydig cells and spermatogenesis [30]. These findings underscore the complexity of Gli-mediated transcription in the testis and highlight the need to delineate the specific roles of individual Gli in SLC differentiation.

Nile tilapia (*Oreochromis niloticus*) is not only an important farm fish in global aquaculture but also a distinguished animal model for investigating gene function and endocrine regulation. This model organism possesses multiple advantageous attributes, including an XX/XY sex-determination system, early sexual maturity (approximately 3 months after hatching for males and 6 months after hatching for females), a short spawning cycle of approximately 14 days, year-round reproductive capacity under laboratory conditions, well-characterized sex-linked genetic markers, and a body size amenable to physiological sampling [31, 32]. Furthermore, a stem Leydig cell line (TSL) has been established from the testis of a 3-month-old Nile tilapia [33]. TSL expresses platelet-derived growth factor receptor α (*pdgfrα*), *nestin*, and chicken ovalbumin upstream promoter transcription factor II (*coup-flla*), which are usually considered as SLC-related markers in several other species [2, 34, 35]. Notably, this cell line exhibits the capacity to differentiate into 11-ketotestosterone (11-KT)-producing Leydig cells both *in vitro* and *in vivo* [33]. When cultured in a defined induction medium, TSL cells differentiate into a steroidogenic phenotype, expressing key steroidogenic genes including *star1*, *star2*, and *cyp11c1*, and producing 11-KT; upon transplantation into recipient testes, TSL cells successfully colonize the

105   interstitial compartment, activate the expression of steroidogenic genes, and restore

106   11-KT production. In this study, by combining CRISPR/Cas9-mediated gene

107   knockout with TSL transplantation technology, we unveil a Dhh-Ptch2-Gli1-Sf1 axis

108   that is essential for Leydig cell lineage differentiation. Our findings provide

109   fundamental insights into the endocrine regulation of Leydig cell development.

RESULTS

Mutation of *dhh* disrupts testicular organization and androgen synthesis

The Nile tilapia Dhh open reading frame (GenBank ID: 100707022, 1377 bp)

was cloned and validated. The encoded 458-amino acid protein shares over 56%

identity with vertebrate homologs (Fig. S1A) and clusters phylogenetically with

mammalian DHH orthologs (Fig. S1B). Transcriptomic analyses conducted on

various tissues and testes across different developmental stages revealed that *dhh*

expression peaked in the testes, particularly at 90 days after hatching (dah) (Fig. S1C,

D).

CRISPR/Cas9 targeting of *dhh* exon 1 from Nile tilapia generated homozygous

mutants ($dhh^{-/-}$) harboring a 13-bp frameshift deletion (Fig. S2A–D), resulting in

premature termination and loss of the conserved Hedge domain (Fig.

S2E). *dhh* mRNA levels in $dhh^{-/-}$ fish were significantly reduced compared to WT (Fig.

S2F). At 90 dah, $dhh^{-/-}$ testes exhibited marked atrophy (Fig. 1A, B) and disorganized

histology, with sparse spermatogonia, rare spermatids, and collapsed interstitial

architecture (Fig. 1C–D', K). Immunofluorescence confirmed significantly reduced

germ cell ($Vasa^{+}$), meiotic cell ($Sycp3^{+}$), and Leydig cell ($Cyp11c1^{+}$) populations (Fig.

1E-J', L-N). Serum 11-KT levels in $dhh^{-/-}$ fish were significantly lower than WT (Fig.

1O), indicating impaired androgen synthesis.

The differentiation of SLCs cannot be rescued by 11-KT, but by SAG

131   Given the established role of 11-KT in teleost testicular development and

132   spermatogenesis [36], we first asked whether exogenous 11-KT could ameliorate the

133   testicular defects in $dhh^{-/-}$ mutants. Compared to untreated mutants, $dhh^{-/-}$ individuals

exposed to 11-KT exhibited a more organized testicular architecture, with significantly increased populations of Vasa☐ and Sycp3☐, as evidenced by both H&E staining and immunofluorescence (Fig. 2A–C', E–G', I–K', R, S). This improvement was further supported by a recovery in GSI (Fig. 2Q). However, the population of Cyp11c1-positive Leydig cells remained profoundly depleted in *dhh*[-/-] +11-KT testes (Fig. 2M–O', T). Taken together, these data indicate that 11-KT treatment partially restored the development of germ cells but not Leydig cell lineage, implying that the Leydig cell defect is androgen-independent.

We next investigated whether direct activation of the Dhh pathway could rescue the *dhh*[-/-] phenotype. Treatment of *dhh*[-/-] mutants with SAG, a Hedgehog agonist, significantly rescued testicular development. This was marked by a significant recovery of germ cells (Vasa☐), spermatocytes (Sycp3☐), and critically, Leydig cells (Cyp11c1☐), alongside a restoration of testicular histology (Fig. 2D–D', H–H', L–L', P–P', Q–T). These results demonstrate that *dhh* mutation does not disrupt survival of endogenous Leydig cell lineage, but directly impairs their differentiation, leading to secondary impairment in androgen production and germ cell development.

Mutation of *dhh* blocks SLCs differentiation *in vivo*

To dissect Leydig cell lineage impairment in *dhh*[-/-] testes, we transplanted the TSL labeled with PKH26 (a fluorescent red hydrophobic membrane dye that enables tracking of transplanted cells) into WT and *dhh*[-/-] testes (Fig. 3A). In WT recipients, TSL cells colonized the interstitium and differentiated into Cyp11c1 positive steroidogenic cells (Fig. 3B1–B3). In *dhh*[-/-] testes, TSL engrafted equivalently but failed to express Cyp11c1 (Fig. 3C1–C3), indicating a differentiation-specific block. Rescue experiments showed that the TSL overexpressing tilapia Dhh (TSL-*On*Dhh) or treated with SAG (TSL+SAG) restored Cyp11c1 expression in *dhh*[-/-] testes (Fig. 3D1–E3, F) and a significant increase in endogenous 11-KT levels (Fig. 3G). These results suggest that SLC differentiation is inhibited in *dhh*[-/-] XY tilapia testes. In addition, the survival and engraftment of PKH26-labeled TSL cells were unaffected in the *dhh*[-/-] environment (Fig. 3C3). Collectively, these results demonstrate that Dhh signaling is specifically required for the differentiation of SLCs into steroidogenic

164 cells, but is dispensable for their survival or initial recruitment into the testicular niche.

Ptch2 is the functional receptor for Dhh signaling in SLCs

RT-PCR results showed that both *ptch1* and *ptch2* were expressed in TSL (Fig.

S3). Fluorescence *in situ* hybridization (FISH) and IF revealed co-expression

of *ptch1* and *ptch2* in Cyp11c1 positive Leydig cells (Fig. 4A). To identify the Dhh-

driving receptor selectivity in SLCs, we established *ptch1*$^{-/-}$ and *ptch2*$^{-/-}$ TSL cell lines

(Figs. S4A, B) and evaluated pathway activity using an 8×GLI luciferase reporter

system (Fig. 4B). In WT TSL cells, the 8×GLI reporter demonstrated substantial

luciferase activity that was markedly enhanced by *On*Dhh overexpression (Fig. 4C),

confirming Dhh responsiveness in this system. Intriguingly, *ptch1*$^{-/-}$ cells maintained

Dhh-induced luciferase activation comparable to WT controls, whereas *ptch2*$^{-/-}$ cells

completely lost this responsiveness (Fig. 4D). This differential receptor requirement

implies that Ptch2 likely acts as the functional receptor for transducing Dhh signals in

TSL cells.

To investigate the function of Ptch2 *in vivo*, we generated *ptch2*$^{-/-}$ and *dhh*$^{-/-}$

;*ptch2*$^{-/-}$ mutant Nile tilapias (Figs. S5A-E). Compared to WT fish at 90 dah, *ptch2*$^{-/-}$

testes exhibited no significant differences in testicular histology (Fig. 4E–F'), GSI

(Fig. 4Q), germ cell (Vasa^{+}, Sycp3^{+}) and Leydig cell (Cyp11c1^{+}) populations (Fig.

4H-I', K-L', N-O', R-T), and serum 11-KT levels (Fig. 4U). Strikingly, the loss

of *ptch2* in the *dhh*$^{-/-}$ background (*dhh*$^{-/-}$;*ptch2*$^{-/-}$) fully rescued the testicular defects,

resulting in normal histology (Fig. 4G–G'), Leydig cell differentiation (Fig. 4P–P'),

and serum 11-KT levels (Fig. 4U). This genetic rescue is consistent with Ptch2

functioning as the inhibitory receptor for Dhh; its removal alleviates the suppression

on the pathway, thereby bypassing the requirement for the Dhh ligand.

Gli1 is the principal transcriptional effector of Dhh signaling in SLCs

In the Hh pathway, Gli transcription factors (Gli1/2/3) mediate diverse

developmental processes (26). RT-PCR results showed that *gli1*, *gli2*, and *gli3* were

all expressed in TSL (Fig. S3). FISH and IF revealed co-expression of *gli1*, *gli2*,

and *gli3* in Cyp11c1 positive Leydig cells (Fig. 5A). To identify Dhh-specific

effectors, we generated *gli1*$^{-/-}$, *gli2*$^{-/-}$, and *gli3*$^{-/-}$ TSL lines (Fig. S4C–E) and assessed

pathway activity using the 8×GLI luciferase reporter. While TSL-$gli2^{-/-}$ and TSL-$gli3^{-/-}$

cell lines maintained Dhh-induced luciferase activation comparable to WT controls,

the TSL-$gli1^{-/-}$ cell line completely lost this responsiveness (Fig. 5B), implying that

Gli1 is the principal transcriptional effector for Dhh signaling transduction in SLCs.

Sf1 is the critical downstream effector of Gli1 in SLCs differentiation

To identify key targets of the Dhh-Gli1 axis, we performed transcriptomic

profiling of TSL cells under conditions of pathway activation: Dhh overexpression

(TSL-*On*Dhh), Gli1 overexpression (TSL-OnGli1), and SAG treatment (TSL+SAG).

Comparative RNA-seq analysis identified a core set of 33 genes consistently

upregulated across all three conditions (Fig. 5C, S6A). Among these, *sf1* emerged as a

prominently upregulated candidate. Functional annotation of its promoter region

identified two conserved Gli1-binding motifs (Fig. 5D). A dual-luciferase reporter

assay confirmed that Gli1 potently transactivates the *sf1* promoter (Fig. 5E). The

specificity of this interaction was further demonstrated by a cold probe competition

assay: a wild-type oligonucleotide containing the Gli-binding motif (GACCACCCA)

competitively inhibited Gli1-driven *sf1* transactivation in a dose-dependent manner,

whereas a mutated version (TTAATTAAA) had no effect (Fig. 5E).

The functional necessity of Sf1 was confirmed by transplantation assays using

sf1$^{-/-}$ and Sf1-overexpressing (TSL-*On*Sf1) TSL cells. While TSL-*sf1*$^{-/-}$ cells failed to

differentiate into Cyp11c1-positive cells even in WT testes, TSL-*On*Sf1 cells

successfully differentiated and expressed Cyp11c1 even in the *dhh*$^{-/-}$ testicular

environment (Fig. 5F–H). These results definitively position Sf1 as the critical

downstream effector executing the Dhh-Ptch2-Gli1 signaling cascade during SLCs

differentiation.

DISCUSSION

219 This study delineates a critical Dhh-Ptch2-Gli1-Sf1 signaling axis essential for

220 SLCs differentiation in Nile tilapia. By combining *dhh*$^{-/-}$ XY models with SLCs

221 transplantation, we definitively demonstrate that Dhh signaling is dispensable for

222 Leydig cell lineage survival but indispensable for its differentiation. Furthermore, our

integrated evidence from targeted gene knockout and functional assays suggests that Ptch2 acts as the primary receptor for Dhh signaling in SLCs, and points to Gli1 as the key transcriptional effector mediating the transactivation of *sf1*. Collectively, these findings implicate Sf1 as the critical downstream target bridging Dhh signaling to steroidogenic maturation, thereby resolving key ambiguities in Leydig cell lineage development and providing evolutionary insights into vertebrate reproductive biology.

Consistent with its established role in mammals, our data in Nile tilapia confirm that Dhh function in male testis development is evolutionarily conserved across vertebrates. While mutations in *Dhh* lead to testosterone insufficiency and infertility in mice [22] and are associated with 46,XY DSD in humans [23], the precise cellular mechanism remained unresolved. Our study demonstrates that $dhh^{-/-}$ tilapia phenocopy these defects, exhibiting testicular dysgenesis, a profound deficiency of Leydig cells, and drastically reduced 11-KT levels (Fig. 1). To define the onset of Leydig cell differentiation, we performed a developmental time-course analysis. This revealed that Cyp11c1-positive steroidogenic cells first appear in wild-type testes at 30 dah, while being conspicuously absent in $dhh^{-/-}$ mutants at this same stage (Fig. S7). This clear temporal pattern establishes ~30 dah as the developmental window when SLCs initiate their differentiation program in the Nile tilapia. The functional indispensability of Dhh at this specific stage was further confirmed by rescue experiments, in which both SAG treatment and transplantation of Dhh-activated TSL cells restored Cyp11c1-positive cell populations in $dhh^{-/-}$ testes (Fig. 3). Collectively, our findings position Dhh as a crucial niche signal that acts at a defined developmental checkpoint to drive the differentiation of SLCs into steroidogenic Leydig cells.

The functional divergence of Ptch1 and Ptch2 in Hh signaling represents an intriguing aspect of pathway regulation. Structurally, both receptors share conserved 12-transmembrane domains vital for cholesterol transport but diverge in their cytoplasmic C-terminal regions, which are implicated in differential Smoothened inhibition efficiency [37, 38]. In mammals, Ptch1 serves as the primary, broad-spectrum regulator of Hh signaling during embryogenesis, whereas Ptch2 often assumes context-dependent roles, such as in primordial germ cell migration via its co-receptor Gas1 [27]. However, their potential redundancy or specificity within the Leydig cell lineage has remained an open question. Our study provides significant insight into this issue. We confirmed that both *ptch1* and *ptch2* are expressed in

Leydig cells (Fig. 4A). Functional interrogation using TSL cell lines revealed a marked difference: genetic ablation of *ptch2* profoundly attenuated Dhh-induced signaling, as measured by a Gli-responsive luciferase reporter, whereas loss of *ptch1* had no detectable effect under the same conditions (Fig. 4B-D). This *in vitro* data, pointing to a primary role for Ptch2, was further supported by *in vivo* genetic interaction studies. The observation that the $dhh^{-/-}$ phenotype was fully rescued in $dhh^{-/-};ptch2^{-/-}$ double mutants (Fig. 4E-U) is consistent with Ptch2 acting as the inhibitory receptor whose loss alleviates pathway suppression. The apparent specificity for Ptch2 in this context might stem from unique co-receptor interactions or expression patterns within the testicular niche. Nonetheless, the early embryonic lethality of global *ptch1* mutation in tilapia [39] precluded a direct assessment of its potential role in postnatal testis development. Therefore, while our findings strongly nominate Ptch2 as the principal receptor for Dhh in SLCs, a definitive exclusion of a role for Ptch1 will require future studies employing Leydig cell-specific conditional knockout models.

Gli is a nuclear transcription factor, characterized by the presence of a zinc finger domain located in the middle of the protein [40]. Downstream of the zinc finger domain is a phosphorylation cluster containing phosphorylation sites for protein kinase A and glycogen synthase kinase 3, which negatively regulate the transcriptional activity of Gli through phosphorylation [41]. In *Drosophila*, Ci is the sole Gli homolog, possessing both transcriptional activation and repression functions [42]. In vertebrates, there are three Gli homologs: Gli1, Gli2, and Gli3. Gli1 functions solely as a transcriptional activator because it lacks proteolytic processing domains. Gli2 has both activation and repression functions, while Gli3 primarily acts as a transcriptional repressor [43, 44]. However, there has been no research on the regulatory mechanism of Gli1/2/3 in SLCs differentiation. In this study, we found that although *gli1*, *gli2*, and *gli3* are all expressed in Leydig cell lineage (Figure 5A-C), the results from TSL cell lines with mutations in *gli1*, *gli2*, and *gli3* indicated that TSL-$gli1^{-/-}$, rather than TSL-$gli2^{-/-}$ or TSL-$gli3^{-/-}$, failed to respond to Dhh signaling (Figure 5D). This suggests that Gli1 is the key transcription factor downstream of the Dhh signaling cascade in SLCs.

Sf1 and Dhh play important roles in Leydig cell commitment and testicular development [5, 11]. Studies suggest a potential interaction between these two factors. For example, in $Dhh^{-/-}$ XY mice, testicular development is impaired, with a significant

reduction in the number of Leydig cells and a marked downregulation of *Sf1* expression [26]. In XX mouse ovaries where Smo is constitutively activated, a large number of ectopic Sf1-positive Leydig cells are generated [45]. Similarly, in *sf1*$^{-/-}$ XY tilapia, testicular development is impaired, with a significant reduction in Leydig cell numbers and down regulation of *dhh* expression [10]. These findings imply that Dhh and Sf1 may interact through feedback loops or co-regulatory mechanisms. However, direct evidence for the regulation of Sf1 by the Hh signaling pathway remains lacking. This study elucidates the molecular mechanism of Sf1 as a critical downstream effect of the Dhh-Ptch2-Gli1 axis. Transcriptomic profiling revealed an increased expression of *sf1* by Dhh regulation signaling in TSL cells (Fig. S6), identifying it as a key downstream target of Dhh signaling. Promoter analysis identified two conserved Gli-binding motifs (Fig. 5F) within the *sf1* promoter, and luciferase assays confirmed that Gli1 transactivates *sf1* transcription (Fig. 5G). Functional validation demonstrated that TSL-*sf1*$^{-/-}$ cells failed to differentiate even in WT testes, whereas TSL-OnSf1 cells restored Leydig cell differentiation and steroidogenesis in *dhh*$^{-/-}$ testes (Fig. 5H-J). These results establish Sf1 as both necessary and sufficient for Leydig cell lineage differentiation.

In conclusion, our study systematically deciphers the signaling cascade governing SLCs differentiation, from the niche signal to the core steroidogenic regulator. We definitively show that the Dhh signal from the niche is indispensable for SLCs differentiation, but not their survival. Within this pathway, Ptch2 serves as the critical receptor, relaying the signal through the transcription factor Gli1. The pivotal outcome of this cascade is the activation of *sf1* by Gli1. Thus, we establish a complete Dhh-Ptch2-Gli1-Sf1 axis, resolving the long-standing question of how a key morphogen controls Leydig cell lineage development in vertebrates.

MATERIALS AND METHODS

Animals

Nile tilapias (*Oreochromis niloticus*) were kept in recirculating freshwater systems at 26□ under a 12-h light/dark photoperiod. All animal experiments were conducted in accordance with the regulations of the Guide for Care and Use of Laboratory Animals prescribed by the Committee of Laboratory Animal

Experimentation at Southwest University, China (IACUC-20181015-12).

Generation of mutant lines by CRISPR/Cas9 in Nile tilapia

The sequences of tilapia *dhh* (GenBank ID: 100707022) and *ptch2* (GenBank ID: 100692939) were retrieved from the NCBI. CRISPR/Cas9-mediated mutagenesis was performed as described previously [46]. Guide RNAs (gRNAs) targeting the first exon of *dhh* (GCGGGCCCGGTCCGCATCCC) and *ptch2* (GTCCCAGGGGCCGGCGTATT) were designed using ZiFit (http://zifit.partners.org/ZiFiT/). gRNA and Cas9 mRNA were synthesized following established protocols [46]. The primers listed in Table S1. A mixture of synthetic gRNA (500 ng/μl) and Cas9 mRNA (1000 ng/μl) was microinjected into one-cell-stage fertilized embryos. Mutants were validated by polyacrylamide gel electrophoresis (PAGE) and Sanger sequencing. F0 chimeric XY males were crossed with WT XX females to generate heterozygous F1 offspring. F1 siblings carrying a 13-bp deletion in *dhh* or a 25-bp deletion in *ptch2* were intercrossed to generate homozygous F2 mutants. For double mutants, $dhh^{+/-}$ XY males were mated with $ptch2^{+/-}$ XX females to obtain $dhh^{+/-};ptch2^{+/-}$ offspring. These double heterozygotes were then intercrossed to generate $dhh^{-/-};ptch2^{-/-}$ mutants.

Generation of mutant lines by pCas9-NtU6sgRNA system in TSL Cells

The tilapia stem Leydig cell line TSL was maintained in ESM4 medium as described [33]. Gene sequences for *ptch1* (GenBank ID: 100700194), *ptch2* (GenBank ID: 100692939), *gli1* (GenBank ID: 100710687), *gli2* (GenBank ID: 100704788), *gli3* (GenBank ID: 100707331), and *sf1* (GenBank ID: 100534561) were retrieved from NCBI. Knockout plasmids targeting these genes were designed following established protocols [47]. gRNAs were generated using ZiFit, with the following exon-specific targets: *ptch1* (GGAGGCGCTCCTGCAGCACC, exon 3), *ptch2* (GTCCCAGGGGCCGGCGTATT, exon 1), *gli1* (TGACTCATGGATCAGGACCA, exon 2), *gli2* (GATTCGCCTGTCACCCCACG, exon

3), *gli3* (TGAGGAGCCCTCTACGTCTA, exon 2),

and *sf1* (GGCTGTGTACTGGTACTGGG, exon 4; designed as previously described)

[47]. Primers are listed in Table S1. Plasmids were transfected into TSL cells using

the MDMP E-1 microchannel transfection instrument (MENDGENE) with transwell

cell compartments (BIOFIT), per manufacturer guidelines. Post-transfection (48

hours), cells were selected in ESM4 medium containing 500 μg/mL G418 for 7 days.

Single-cell clones were isolated via dilution and serial passage, followed by genomic

DNA (gDNA) extraction and Sanger sequencing to confirm mutations.

Histological analyses of testes from WT, *dhh*$^{-/-}$, *ptch2*$^{-/-}$ and *dhh*$^{-/-}$;*ptch2*$^{-/-}$ XY fish

GSI was calculated for WT, *dhh*$^{-/-}$, *ptch2*$^{-/-}$ and *dhh*$^{-/-}$; *ptch2*$^{-/-}$ XY fish at 90 dah

as: (gonad weight / body weight) $\times$ 100%. Testes were fixed in Bouin's solution for 24

hours at room temperature with agitation, dehydrated through an ethanol series, and

embedded in paraffin. Sections (5 μm thickness) were stained with hematoxylin and

eosin (H&E) and imaged using an Olympus BX53 microscope. Germ cell

classification (spermatogonia, spermatocytes) followed established criteria [48].

Immunofluorescence

Testes from WT, *dhh*$^{-/-}$, WT+11-KT, *dhh*$^{-/-}$+11-KT, *ptch2*$^{-/-}$ and *dhh*$^{-/-}$; *ptch2*$^{-/-}$ XY

fish at 90 dah were analyzed for germ cell marker Vasa, Leydig cell marker Cyp11c1,

and meiosis marker Sycp3. Tissue fixation and embedding. Sections (5 μm) were

deparaffinized, rehydrated, and subjected to antigen retrieval. After blocking with 5%

bovine serum albumin (BSA; 30 minutes, room temperature), sections were incubated

overnight at 4°C with rabbit polyclonal primary antibodies (Vasa, Cyp11c1, Sycp3;

1:1,000 dilution), validated previously [49]. Alexa Fluor 488-conjugated goat anti-

rabbit secondary antibody (1:1,000; Invitrogen) was applied for 1 hour at 37°C,

followed by DAPI nuclear counterstaining (1:1,000; Sigma-Aldrich). Fluorescence

signals were imaged using an Olympus Fv3000 confocal microscope. For each

biological replicate (*n*=5-6 fish per genotype), three non-serial, non-adjacent testis

sections were analyzed. From each section, three representative fields of view were

captured to ensure non-overlapping sampling. All positive cells number of Vasa, Sycp3 and Cyp11c1 was quantified by Image J Pro 1.51 software using default parameters.

Fluorescence *in situ* hybridization (FISH) and immunofluorescence

Testes from WT XY tilapia at 90 dah were fixed in 4% paraformaldehyde (Sigma) overnight at 4°C with agitation, paraffin-embedded, and sectioned (5 μm). FISH was performed as described [48]. Partial cDNA fragments of *ptch1*, *ptch2*, *gli1*, *gli2*, and *gli3* were cloned into the pGEM-T Easy vector. Sense (control) and antisense RNA probes were synthesized using a DIG RNA Labeling Kit (Roche) and T7 mMESSAGE mMACHINE Kit (Ambion). Signals were amplified with the TSA^{TM} Plus TMR system (PerkinElmer) per manufacturer protocols (primers in Table S1). Concurrently, Leydig cell marker Cyp11c1 expression was assessed by immunofluorescence.

11-KT and SAG rescue experiments

To rescue *dhh$^{-/-}$* XY fish, offspring from *dhh$^{+/-}$* XY male × *dhh$^{+/-}$* XX female crosses were divided into two groups. The experimental group received 11-KT via immersion from 30 to 90 dah, with treatments refreshed every 2 days. 11-KT concentrations increased incrementally from 57 ng/L at 30 dah to 1254 ng/L by 90 dah (39.9 ng/L increase per treatment cycle; see Fig. 2, 5, S7). For SAG rescue, fish received intraperitoneal injections of 10 mg/kg SAG from 30 to 90 dah, with booster doses administered weekly. Testes were collected at 90 dah for morphological and histological evaluation.

TSL cell labeling, transplantation and IF

TSL, previously isolated from 3-month-old XY tilapia testes [33], was used to assess stem Leydig cell differentiation in WT and *dhh$^{-/-}$* XY fish. TSL cells were treated as follows: (1) untreated TSL, (2) TSL + 0.5 μM SAG (Hh pathway agonist), (3) TSL-*On*Dhh (overexpressing tilapia Dhh), (4) TSL-*On*Sf1 (overexpressing tilapia Sf1), and (5) TSL-*sf1$^{-/-}$* (*sf1* mutation). For the SAG treatment experiment, TSL

cells were incubated with 0.5 μM SAG for 48 hours before transplantation. Cells were

labeled with PKH26 fluorescent dye (Sigma) and transplanted (5,000 cells per testis)

into WT or $dhh^{-/-}$ XY tilapia via the urogenital tract [33]. For TSL-OnDhh,

the Dhh ORF (1377 bp) was amplified from gDNA, cloned into pcDNA3.1

(Invitrogen), and transfected into TSL cells (primers in Table S1). At 10 days post-

transplantation, testes were analyzed by IF to quantify Cyp11c1 positive cells among

PKH26 positive populations.

GLI report assay

A firefly luciferase reporter plasmid (8×GLI) was generated by cloning eight

tandem copies of the GLI-binding motif (GACCACCCA) into the pGL4.23 vector

(minimal promoter). To validate 8×GLI functionality, TSL cells were transiently

transfected with: (1) pGL4.23 (negative control), (2) 8×GLI, or (3) 8×GLI plus Dhh-

overexpression plasmid (OnDhh), for 48 hours. The 8×GLI construct was further

transfected into TSL (WT), TSL-$ptch1^{-/-}$, TSL-$ptch2^{-/-}$, TSL-$gli1^{-/-}$, TSL-$gli2^{-/-}$ or TSL-

$gli3^{-/-}$, with or without OnDhh. Transfections included pRL-TK as an internal control.

Luciferase activity was measured 48 hours post-transfection using the Dual Luciferase

Assay System (Promega) and a Luminoskan™ Ascent luminometer (Thermo Fisher

Scientific). Relative activity was calculated as firefly/Renilla luciferase ratios.

Western blot

The ORFs of Dhh (1377 bp) and Gli1 (4404 bp) were amplified from gDNA by

PCR. Two tandem FLAG tag sequences (GATTATAAAGATGATGATAAA) were

inserted flanking each ORF (after the start codon and before the stop codon) and

cloned into the pcDNA3.1 vector (Invitrogen), generating pcDNA3.1-OnDhh and

pcDNA3.1-OnGli1 (primers in Table S1). TSL cells were transiently transfected with

pcDNA3.1-OnDhh or pcDNA3.1-OnGli1 for 48 hours. Protein expression was

analyzed by Western blot following established methods [49]. Briefly, total protein

lysates were separated by 15% SDS-PAGE, transferred to nitrocellulose membranes,

and blocked with 5% BSA in TBST (10 mM Tris pH 7.9, 150 mM NaCl, 0.1%

Tween-20) for 1 hour at 37°C. Membranes were incubated overnight at 4°C with

rabbit anti-FLAG antibody (1:1000; Cell Signaling Technology), followed by HRP-conjugated goat anti-rabbit secondary antibody (1:1000; Invitrogen) for 1 hour at 37°C. Signals were detected using the BeyoECL Plus Kit (Beyotime) and imaged on a Fusion FX7 system (Vilber Lourmat).

Measurement of 11-KT levels by ELISA

Blood was collected from the caudal vein of 90 dah XY fish (WT, $dhh^{-/-}$, $ptch2^{-/-}$, $dhh;ptch2^{-/-}$, WT+TSL, $dhh^{-/-}$+TSL, $dhh^{-/-}$+TSL+SAG, $dhh^{-/-}$+TSL-OnDhh and $dhh^{-/-}$+TSL-$sf1^{-/-}$), incubated at 4°C overnight, and centrifuged to isolate serum. Tissue fluid samples from WT and $dhh^{-/-}$ XY fish at 5, 10, 20, and 30 dah were similarly prepared. All samples were stored at −80°C until analysis. Serum and tissue fluid 11-KT levels were quantified using a 11-keto Testosterone ELISA Kit (Cayman) validated for teleost samples, per manufacturer instructions. Individual samples (serum or tissue fluid; $n = 6$ per genotype) were analyzed in triplicate on a single 96-well plate, with concentrations calculated from standard curves.

Transcriptome analyses

Tissue- and stage-specific expression of dhh was assessed using transcriptomic data from tilapia tissues (brain, heart, head kidney, kidney, liver, muscle, ovary, testis) and gonads at developmental stages spanning 5 to 360 dah [50, 51]. For global gene expression profiling, RNA was extracted from TSL-WT, TSL-OnDhh, TSL-OnGli1, and TSL + 0.5 μM SAG cells using RNAiso Plus (Takara). For the SAG treatment experiment, TSL cells were incubated with 0.5 μM SAG for 48 hours before collection. For each genotype, cells from three independent culture wells were pooled. Libraries were sequenced on an Illumina platform (GENEBOOK Biotechnology), and clean reads were aligned to the reference genome with HISAT2. Gene expression levels (FPKM: fragments per kilobase per million mapped reads) were quantified using featureCounts. Differentially expressed genes (DEGs) were identified for each condition (TSL-OnDhh, TSL-OnGli1, TSL+SAG) compared to TSL-WT controls using edgeR (threshold: FDR < 0.05, |log2(foldchange)| ≥ 1.5).

Real-time PCR

Testes from WT and $dhh^{-/-}$ XY fish at 90 dah ($n = 3$ per genotype), along with

TSL-WT, TSL-*On*Dhh, TSL-*On*Gli1, and TSL + 0.5 μM SAG cells, were analyzed

for gene expression. Total RNA was extracted using RNAiso Plus (Takara), reverse-

transcribed into cDNA (PrimeScript II 1st Strand cDNA Synthesis Kit; Takara), and

amplified by real-time PCR on an ABI-7500 system (Applied Biosystems). Relative

mRNA levels were normalized to *β-actin* and calculated via the formula $R = 2^{-\Delta\Delta Ct}$

(primers in Table S1).

Dual luciferase report assay

The 5015-bp promoter region upstream of the *sf1* start codon was amplified and

cloned into the pGL3-basic vector (Promega), generating pGL3-*sf1* (primers in Table

S1). To assess Gli1 binding to the *sf1* promoter, HEK293 cells were co-transfected

with pGL3-*sf1*, pcDNA3.1- *On*Gli1, cold probe (GACCACCCA), and mutant cold

probe (TTAATTAAA) using the TransIT-X2 transfection reagent (Mirus Bio). Control

transfections included empty pcDNA3.1 and pGL3 vectors, with pRL-TK for

normalization. Luciferase activity was measured 48 hours post-transfection.

Statistical analyses

480 Data are presented as mean ± SD from at least three independent replicates. A

481 two-tailed independent Student's t-test was used to determine the differences between

482 the two groups. One-way ANOVA, followed by Tukey multiple comparison, was used

483 to determine the significance of differences in more than two groups. $P < 0.05$ was

484 used as a threshold for statistically significant differences.

Data availability statement

486 The data that support the findings of this study are available from the

487 corresponding author upon reasonable request.

Acknowledgements

489 This research was funded by the National Key Research and Development

490 Program of China (Grant No. 2022YFD1201600), the National Natural Science

491 Foundation of China (Grant No. 32172969, 31972776, 32102780, 32473159).

Author contributions

493 **Changle Zhao**: Writing – original draft, Software, Resources, Methodology,

Investigation, Formal analysis, Data curation. **Yongxun Chen**: Writing – original draft, Methodology, Data curation. **Lei Liu**: Data curation, Conceptualization. **Xiang Liu**: Software, Investigation. **Hesheng Xiao**: Software, Data curation. **Feilong Wang**: Investigation, Formal analysis. **Qin Huang**: Resources, Software. **Xiangyan Dai**: Resources, Investigation. **Wenjing Tao**: Resources, Funding acquisition. **Deshou Wang**: Supervision, Investigation, Funding acquisition. **Jing Wei**: Writing – review & editing, Validation, Supervision, Project administration, Funding acquisition, Data curation, Conceptualization.

Conflict of interest

The authors declare no competing interest.

References

1. Inoue, M., Baba, T., and Morohashi, K.I. (2018). Recent progress in understanding the mechanisms of Leydig cell differentiation. Mol. Cell. Endocrinol. *468*, 39-46.

2. Jiang, M.H., Cai, B., Tuo, Y., Wang, J., Zang, Z.J., Tu, X., Gao, Y., Su, Z., Li, W., Li, G., et al. (2014). Characterization of Nestin-positive stem Leydig cells as a potential source for the treatment of testicular Leydig cell dysfunction. Cell. Res. *24*, 1466-1485.

3. Yao, S., Wei, X., Deng, W., Wang, B., Cai, J., Huang, Y., Lai, X., Qiu, Y., Wang, Y., Guan, Y., et al. (2022). Nestin-dependent mitochondria-ER contacts define stem Leydig cell differentiation to attenuate male reproductive ageing. Nat. Commun. *13*, 4020.

4. Chen, P., Zirkin, B.R., and Chen, H. (2020). Stem Leydig Cells in the Adult Testis: Characterization, Regulation and Potential Applications. Endocr. Rev. *41*, 22-32.

5. Parker, K.L., Rice, D.A., Lala, D.S., Ikeda, Y., Luo, X., Wong, M., Bakke, M., Zhao, L., Frigeri, C., Hanley, N.A., et al. (2002). Steroidogenic factor 1: an essential mediator of endocrine development. Recent. Prog. Horm. Res. *57*, 19-36.

6. Houzelstein, D., Eozenou, C., Lagos, C.F., Elzaiat, M., Bignon-Topalovic, J., Gonzalez, I., Laville, V., Schlick, L., Wankanit, S., Madon, P., et al. (2024). A conserved NR5A1-responsive enhancer regulates SRY in testis-determination. Nat. Commun. *15*, 2796.

7. Smith, O.E., Morin, F., Roussel, V., Bertucci, M.C., Boyer, A., and Murphy, B.D. (2023). The role of steroidogenic factor 1 (SF-1) in steroidogenic cell function of the testes and ovaries of mature mice. Reproduction (Cambridge, England). *165*, 1-17.

8. Park, S.Y., Tong, M., and Jameson, J.L. (2007). Distinct roles for steroidogenic factor 1 and desert hedgehog pathways in fetal and adult Leydig cell development. Endocrinology. *148*, 3704-3710.

9. Yu, Y., Lee, P.A., Huerta-Saenz, L., and Allen, N.G. (2023). A Novel Variant in NR5A1 Presenting as 46,XY Difference of Sex Development. JCEM. Case. Rep. *1*, luad103.

10. Xie, Q.P., He, X., Sui, Y.N., Chen, L.L., Sun, L.N., and Wang, D.S. (2016). Haploinsufficiency of SF-1 Causes Female to Male Sex Reversal in Nile Tilapia, Oreochromis niloticus. Endocrinology.

157, 2500-2514.

11.   Zhang, Y., and Beachy, P.A. (2023). Cellular and molecular mechanisms of Hedgehog signalling.
Nat. Rev. Mol. Cell. Biol. *24*, 668-687.

12.   Briscoe, J., and Thérond, P.P. (2013). The mechanisms of Hedgehog signalling and its roles in
development and disease. Nat. Rev. Mol. Cell. Biol. *14*, 416-429.

13.   Ingham, P.W., Nakano, Y., and Seger, C. (2011). Mechanisms and functions of Hedgehog
signalling across the metazoa. Nat. Rev. Genet. *12*, 393-406.

14.   Finco, I., LaPensee, C.R., Krill, K.T., and Hammer, G.D. (2015). Hedgehog signaling and
steroidogenesis. Annu. Rev. Physiol. *77*, 105-129.

15.   Carpenter, D., Stone, D.M., Brush, J., Ryan, A., Armanini, M., Frantz, G., Rosenthal, A., and de
Sauvage, F.J. (1998). Characterization of two patched receptors for the vertebrate hedgehog
protein family. Proc. Natl. Acad. Sci. USA. *95*, 13630-13634.

16.   Belgacem, Y.H., and Borodinsky, L.N. (2015). Inversion of Sonic hedgehog action on its
canonical pathway by electrical activity. Proc. Natl. Acad. Sci. USA. *112*, 4140-4145.

17.   Cervantes, S., Lau, J., Cano, D.A., Borromeo-Austin, C., and Hebrok, M. (2010). Primary cilia
regulate Gli/Hedgehog activation in pancreas. Proc. Natl. Acad. Sci. USA. *107*, 10109-10114.

18.   Kothandapani, A., Lewis, S.R., Noel, J.L., Zacharski, A., Krellwitz, K., Baines, A., Winske, S.,
Vezina, C.M., Kaftanovskaya, E.M., Agoulnik, A.I., et al. (2020). GLI3 resides at the intersection
of hedgehog and androgen action to promote male sex differentiation. PLoS. Genet. *16*,
e1008810.

19.   Umehara, F., Tate, G., Itoh, K., Yamaguchi, N., Douchi, T., Mitsuya, T., and Osame, M. (2000). A
novel mutation of desert hedgehog in a patient with 46,XY partial gonadal dysgenesis
accompanied by minifascicular neuropathy. Am. J. Hum. Genet. *67*, 1302-1305.

20.   Pachernegg, S., Georges, E., and Ayers, K. (2022). The Desert Hedgehog Signalling Pathway in
Human Gonadal Development and Differences of Sex Development. Sex. Dev. *16*, 98-111.

21.   Mehta, P., Singh, P., Gupta, N.J., Sankhwar, S.N., Chakravarty, B., Thangaraj, K., and Rajender, S.
(2021). Mutations in the desert hedgehog (DHH) gene in the disorders of sexual
differentiation and male infertility. J. Assist. Reprod. Genet. *38*, 1871-1878.

22.   Pierucci-Alves, F., Clark, A.M., and Russell, L.D. (2001). A developmental study of the Desert
hedgehog-null mouse testis. Biol. Reprod. *65*, 1392-1402.

23.   Wei, J., Wu, J., Ru, W., Chen, G., Gao, L., and Tang, D. (2022). Novel compound heterozygous
mutations in the desert hedgehog (DHH) gene in cases of siblings with 46,XY disorders of
sexual development. BMC. Med. Genomics. *15*, 178.

24.   Cong, G., Zhu, X., Chen, X.R., Chen, H., and Chong, W. (2025). Mechanisms and therapeutic
potential of the hedgehog signaling pathway in cancer. Cell. Death. Discov. *11*, 40.

25.   Nieuwenhuis, E., Motoyama, J., Barnfield, P.C., Yoshikawa, Y., Zhang, X., Mo, R., Crackower,
M.A., and Hui, C.C. (2006). Mice with a targeted mutation of patched2 are viable but develop
alopecia and epidermal hyperplasia. Mol. Cell. Biol. *26*, 6609-6622.

26.   Yao, H.H., Whoriskey, W., and Capel, B. (2002). Desert Hedgehog/Patched 1 signaling specifies
fetal Leydig cell fate in testis organogenesis. Genes. Dev. *16*, 1433-1440.

27.   Kim, Y., Lee, J., Seppala, M., Cobourne, M.T., and Kim, S.H. (2020). Ptch2/Gas1 and Ptch1/Boc
differentially regulate Hedgehog signalling in murine primordial germ cell migration. Nature.
Commun. *11*, 1994.

28.   Ruiz i Altaba, A., Sánchez, P., and Dahmane, N. (2002). Gli and hedgehog in cancer: tumours,

576   embryos and stem cells. Nat. Rev. Cancer. *2*, 361-372.

577   29.   Hui, C.C., and Angers, S. (2011). Gli proteins in development and disease. Annu. Rev. Cell. Dev.

578   Biol. *27*, 513-537.

579   30.   Barsoum, I., and Yao, H.H. (2011). Redundant and differential roles of transcription factors

580   Gli1 and Gli2 in the development of mouse fetal Leydig cells. Biol. Reprod. *84*, 894-899.

581   31.   Li, M., Sun, L., Zhou, L., and Wang, D. (2024). Tilapia, a good model for studying reproductive

582   endocrinology. Gen. Comp. Endocrinol. *345*, 114395.

583   32.   Sun, Y.-L., Jiang, D.-N., Zeng, S., Hu, C.-J., Ye, K., Yang, C., Yang, S.-J., Li, M.-H., and Wang, D.-S.

584   (2014). Screening and characterization of sex-linked DNA markers and marker-assisted

585   selection in the Nile tilapia (*Oreochromis niloticus*). Aquaculture. *433*, 19-27.

586   33.   Huang, Q., Yang, Z., Wang, J., Luo, Y., Zhao, C., Li, M., Xiao, H., Tao, W., Wang, D., and Wei, J.

587   (2020). Establishment of a stem Leydig cell line capable of 11-ketotestosterone production.

588   Reprod. Fertil. Dev. *32*, 1271-1281.

589   34.   Ge, R.S., Dong, Q., Sottas, C.M., Papadopoulos, V., Zirkin, B.R., and Hardy, M.P. (2006). In

590   search of rat stem Leydig cells: identification, isolation, and lineage-specific development.

591   Proc. Natl. Acad. Sci. USA. *103*, 2719-2724.

592   35.   Chen, H., Wang, Y., Ge, R., and Zirkin, B.R. (2017). Leydig cell stem cells: Identification,

593   proliferation and differentiation. Mol. Cell. Endocrinol. *445*, 65-73.

594   36.   Zheng, Q., Xiao, H., Shi, H., Wang, T., Sun, L., Tao, W., Kocher, T.D., Li, M., and Wang, D. (2020).

595   Loss of Cyp11c1 causes delayed spermatogenesis due to the absence of 11-ketotestosterone.

596   J. Endocrinol. *244*, 487-499.

597   37.   Qi, X., Schmiege, P., Coutavas, E., and Li, X. (2018). Two Patched molecules engage distinct

598   sites on Hedgehog yielding a signaling-competent complex. Science (New York, N.Y.). *362*.

599   38.   Fleet, A.J., and Hamel, P.A. (2018). The protein-specific activities of the transmembrane

600   modules of Ptch1 and Ptch2 are determined by their adjacent protein domains. J. Biol. Chem.

601   *293*, 16583-16595.

602   39.   Liu, X., Zhao, C., Liu, L., Peng, X., Li, J., Tao, W., Wang, D., and Wei, J. (2024). Nile Tilapia

603   (*Oreochromis niloticus*) Patched1 Mutations Disrupt Cardiovascular Development and

604   Vascular Integrity through Smoothened Signaling. Int. J. Mol. Sci. *25*.

605   40.   Kinzler, K.W., Ruppert, J.M., Bigner, S.H., and Vogelstein, B. (1988). The GLI gene is a member

606   of the Kruppel family of zinc finger proteins. Nature. *332*, 371-374.

607   41.   Niewiadomski, P., Kong, J.H., Ahrends, R., Ma, Y., Humke, E.W., Khan, S., Teruel, M.N., Novitch,

608   B.G., and Rohatgi, R. (2014). Gli protein activity is controlled by multisite phosphorylation in

609   vertebrate Hedgehog signaling. Cell. Rep. *6*, 168-181.

610   42.   Huangfu, D., and Anderson, K.V. (2006). Signaling from Smo to Ci/Gli: conservation and

611   divergence of Hedgehog pathways from Drosophila to vertebrates. Development (Cambridge,

612   England). *133*, 3-14.

613   43.   Niewiadomski, P., Niedziółka, S.M., Markiewicz, Ł., Uśpieński, T., Baran, B., and Chojnowska, K.

614   (2019). Gli Proteins: Regulation in Development and Cancer. Cells. *8*.

615   44.   Pan, Y., and Wang, B. (2007). A novel protein-processing domain in Gli2 and Gli3 differentially

616   blocks complete protein degradation by the proteasome. J. Biol. Chem. *282*, 10846-10852.

617   45.   Barsoum, I.B., Bingham, N.C., Parker, K.L., Jorgensen, J.S., and Yao, H.H. (2009). Activation of

618   the Hedgehog pathway in the mouse fetal ovary leads to ectopic appearance of fetal Leydig

619   cells and female pseudohermaphroditism. Dev. Biol. *329*, 96-103.

46.   Li, M., Yang, H., Zhao, J., Fang, L., Shi, H., Li, M., Sun, Y., Zhang, X., Jiang, D., Zhou, L., et al.
(2014). Efficient and heritable gene targeting in tilapia by CRISPR/Cas9. Genetics. *197*, 591-
599.
47.   Zhang, Z., Wang, J., Li, J., Liu, X., Liu, L., Zhao, C., Tao, W., Wang, D., and Wei, J. (2023).
Establishment of an Integrated CRISPR/Cas9 Plasmid System for Simple and Efficient Genome
Editing in Medaka In Vitro and In Vivo. Biology. *12*.
48.   Zhao, C., Liu, X., Liu, L., Li, J., Liu, X., Tao, W., Wang, D., and Wei, J. (2023). Smoothened
mediates medaka spermatogonia proliferation via Gli1-Rgcc-Cdk1 axis†. Biol. Reprod. *109*,
772-784.
49.   Dai, S., Qi, S., Wei, X., Liu, X., Li, Y., Zhou, X., Xiao, H., Lu, B., Wang, D., and Li, M. (2021).
Germline sexual fate is determined by the antagonistic action of dmrt1 and foxl3/foxl2 in
tilapia. Development (Cambridge, England). *148*.
50.   Tao, W., Yuan, J., Zhou, L., Sun, L., Sun, Y., Yang, S., Li, M., Zeng, S., Huang, B., and Wang, D.
(2013). Characterization of gonadal transcriptomes from Nile tilapia (*Oreochromis niloticus*)
reveals differentially expressed genes. PloS. One. *8*, e63604.
51.   Tao, W., Chen, J., Tan, D., Yang, J., Sun, L., Wei, J., Conte, M.A., Kocher, T.D., and Wang, D.
(2018). Transcriptome display during tilapia sex determination and differentiation as revealed
by RNA-Seq analysis. BMC. Genomics. *19*, 363.

Figure legends

Fig. 1. Mutation of *dhh* resulted in testicular developmental disorders.

(A-B) Morphological analysis of the testes from WT (*dhh*$^{+/+}$) and *dhh*$^{-/-}$ XY fish at 90

dah. Arrows indicate the location of the testes.

(C-D') Histological analysis of testicular sections from WT and *dhh*$^{-/-}$ XY fish at 90

dah. Sg, spermatogonia; Sc, spermatocyte.

(E-J') Representative immunofluorescence images showing the expression of germ

cell marker Vasa (E-F'), meiosis cell marker Sycp3 (G-H') and Leydig cell marker

Cyp11c1 (I-J').

(K) Gonadosomatic index (GSI) of the testes from WT and *dhh*$^{-/-}$ XY fish at 90 dah

(*n*=5 fish/genotype).

(L-N) Quantification of the percentage of cells positive for Vasa (E-F'), Sycp3 (G-H')

and Cyp11c1 (I-J') among all DAPI-positive cells in the WT and *dhh*$^{-/-}$ testes (*n*=5

fish/genotype).

(O) Serum 11-KT level of WT and *dhh*$^{-/-}$ XY fish at 90 dah (*n*=6 fish/genotype).

Values were presented as mean ± SD. Differences were determined by two-tailed

independent Student's t-test. **, *P* < 0.01. Scale bars, (A-B), 1 cm; (C-J), 50 μm.

Fig. 2. Rescue of testicular development in *dhh*$^{-/-}$ XY Nile tilapia by 11-KT and

SAG.

dhh^{-/-} XY Nile tilapias at 30 dah were treated either with 11-KT via water immersion (with water changed every two days) or with 10 mg/kg SAG via intraperitoneal injection (with supplemental injection every seven days). WT controls received 11-KT, and *dhh*^{-/-} controls received an equivalent volume of the DMSO. Then, at 90 dah, morphological and histological experiments were conducted.

(A-D') Histological analysis of testicular sections from 90 dah XY fish subjected to different treatments as indicated. Sg, spermatogonia; Sc, spermatocyte. Scale bars, 50 μm.

(E-P') Representative immunofluorescence images showing the expression of Vasa (E-H'), Sycp3 (I-L') and Cyp11c1 (M-P'). Scale bars, 50 μm.

(Q) GSI of testes from the different treatment groups at 90 dah ($n=5$ fish per group).

(R-T) Quantification of the percentage of cells positive for Vasa (E-H'), Sycp3 (I-L') and Cyp11c1 (M-P') among all DAPI-positive cells in the testes ($n=3 \sim 5$ fish per group).

Values were presented as mean ± SD. Different letters above the error bar indicate statistical differences at $P < 0.05$ as determined by one-way ANOVA followed by Tukey test.

Fig. 3. Dhh is required for SLCs differentiation *in vivo*.

TSL, Dhh-overexpressing TSL (TSL-*On*Dhh), or 0.5 μM SAG-treated TSL (TSL+SAG) cells were labeled with PKH26 and transplanted into the testes of 90 dah WT or *dhh*^{-/-} recipient fish. Analyses were performed 10 days post-transplantation.

(A) Schematic diagram of the experimental design for SLCs transplantation and analysis.

(B1-E3) Representative immunofluorescence images of testicular sections from recipient fish, showing the localization of transplanted PKH26-labeled SLCs (red) and the expression of Cyp11c1 (green). Nuclei are stained with DAPI (blue). Scale bars, 4 μm.

(F) Quantification of the percentage of Cyp11c1-positive cells among the transplanted PKH26-positive SLCs for each treatment group ($n=5$ fish per group).

(G) Serum 11-KT level in recipient fish following SLCs transplantation ($n=5$ fish per group).

Values were presented as mean ± SD. Different letters above the error bar indicate

statistical differences at $P < 0.05$ as determined by one-way ANOVA followed by

Tukey test.

Fig. 4. Ptch2 mediates Dhh signaling in SLCs.

(A) Co-localization of Cyp11c1 (green, by immunofluorescence)

and *ptch1* or *ptch2* mRNA (red, by RNA-FISH) in adult (90 dah) testis sections.

Dashed lines outline representative Leydig cells. Scale bars, 4 μm.

(B) Schematic illustration of the experimental setup for the luciferase reporter assays

shown in panels C and D

(C) Luciferase activity in TSL cells co-transfected with a Gli-responsive reporter

(8xGLI) and overexpression of tilapia Dhh (*On*Dhh). A plasmid lacking Gli-binding

sites (pGL4.23) served as a negative control (*n*=3).

(D) Luciferase activity in TSL-WT, TSL-*ptch1*$^{-/-}$ and TSL-*ptch2*$^{-/-}$ cells transfected

with the 8xGLI reporter with or without *On*Dhh (*n*=4).

(E-G') Histological analysis of testis sections from the indicated genotypes at 90 dah.

Sg, spermatogonia; Sc, spermatocyte. Scale bars, 50 μm.

(H-P') Immunofluorescence analysis of Vasa (H-J'), Sycp3 (K-M') and Cyp11c1 (N-

P') in testis sections from the indicated genotypes. Scale bars, 50 μm.

(Q) GSI of the testes from the indicated genotypes at 90 dah (*n*=4 fish per group).

(R-T) Quantification of the percentage of Vasa (H-J'), Sycp3 (K-M') and Cyp11c1

(N-P') positive cells among DAPI-positive cells (*n*=6 fish per group).

(U) Serum 11-KT levels in fish of the indicated genotypes at 90 dah (*n*=6 fish per

group).

Statistical significance was determined by one-way ANOVA followed by Tukey's test

(C, Q-U, different letters above the error bar indicate statistical differences at $P < 0.05$)

or Student's *t*-test (D) (*, $P < 0.05$; **, $P < 0.01$; NS, no significant difference).

Fig. 5. Gli1 transactivates *sf1* to drive SLC differentiation

(A) Co-localization of Cyp11c1 (green) and *gli1*, *gli2*, or *gli3* mRNA (red) in adult

(90 dah) testis sections by immunofluorescence and RNA-FISH. Dashed lines outline

representative Leydig cells. Scale bars, 4 μm.

(B) Luciferase activity in TSL-WT, TSL-*gli1*$^{-/-}$, TSL-*gli2*$^{-/-}$ and TSL-*gli3*$^{-/-}$ cells

transfected with the 8xGLI reporter with or without *On*Dhh (*n*=4).

(C) Volcano plots of transcriptomic changes in TSL-*On*Dhh, TSL-*On*Gli1 and TSL+

SAG, compared to TSL-WT. Red and blue dots represent significantly up- and down-

regulated genes, respectively.

(D) Schematic of the tilapia *sf1* gene promoter, indicating the two predicted Gli1

binding sites (B1 and B2).

(E) Transcriptional activation of the *sf1* promoter by Gli1. Competition assays were

performed using unlabeled cold probe (Cold probe, GACCACCCA, 10/100/250

ng/mL) or mutant unlabeled cold probe (mutant Cold probe, TTAATTAAA,

10/100/250 ng/mL) (*n*=4). "+" indicates the addition of the corresponding substance,

while "-" indicates no addition, and the number represents the amount added (ng/ml).

(F-G) Representative immunofluorescence images of testicular sections from recipient

fish transplanted with *sf1*-deficient (TSL-*sf1*$^{-/-}$, F) or Sf1-overexpressing (TSL-*On*Sf1,

G) SLCs, stained for Cyp11c1 (green) and PKH26 (red). Nuclei are stained with

DAPI (blue). Scale bars, 4 μm.

(H) Quantification of the percentage of Cyp11c1-positive cells among the

transplanted PKH26-positive SLCs in the two transplantation groups (*n*=5 fish per

group).

Statistical significance was determined by one-way ANOVA followed by Tukey's test

(E, different letters above the error bar indicate statistical differences at $P < 0.05$) or

Student's *t*-test (B, H) (*, $P < 0.05$; **, $P < 0.01$; NS, no significant difference).

Fig. S1. Sequence analysis and mRNA expression profile of Nile tilapia *dhh*.

(A) Dhh amino acid sequence alignment. Amino acids are numbered in the right

margin. Deletions are indicated by dashes, shaded areas indicate shared sequences.

The amino-terminal hedge domain is underlined by dotted line, with the signal

peptide region in the front and the autocatalytic carboxy-terminal domain (hog) in the

back. The box indicates the autocatalytic site of an absolutely conserved Gly-Cys-Phe

tripeptide. At the end of the alignment are percentage identity values of the full-length

and hedge domain of tilapia Dhh to the orthologue from other species.

(B) Phylogenetic analysis of Dhh. The phylogenetic tree was constructed using the

neighbor-joining method within the MEGA7.0 program. Node values represent

percent bootstrap confidence derived from 1000 replicates.

(C-D) The RPKM (reads per kb per million reads) values of tilapia *dhh* in various

adult tissues (C) and XY and XX gonads at 5, 7, 20, 30, 90, 180, 300 dah (days after

hatching) (D) in transcriptome sequencing data which were sequenced using Illumina

2000 HiSeq technology in our previous study (Tao et al., 2013).

Fig. S2. Establishment of the Nile tilapia *dhh* mutant line by CRISPR/Cas9.

(A) Schematic representation of gRNA targeting the Nile tilapia *dhh* locus. The gRNA

was designed to target exon 1. The translation start codon ATG and stop codon TAA

were indicated by arrows. The PAM (protospacer adjacent motif) site was marked by

green box. DNA sequence alignment of the three mutant lines with WT. The added

sequences were marked by red letter, and the deleted sequences were indicated in red

dotted lines. One had a 4-bp addition, one had an 8-bp deletion, and the other had a

13-bp deletion.

(B) Schematic diagram showing the breeding plans of *dhh* F0 to F2 fish.

(C) Homozygous mutants of F2 fish were screened by PAGE. The first lane is DNA

marker, the second lane is $dhh^{+/+}$ fish, the third lane is $dhh^{+/-}$ fish, and the fourth lane

is $dhh^{-/-}$ fish.

(D) Sanger sequencing results of *dhh* genes from WT and the 13-bp deletion

homozygous mutant fishes.

(E) The *dhh* ORF sequence in WT was indicated in black and the frameshift-altered

sequence for the 13-bp deletion *dhh* ORF was indicated in red.

(F) The relative mRNA expression of *dhh* in WT and the 13-bp deletion homozygous

mutant fishes by RT-qPCR. Values were presented as mean ± SD (*n*=3). Differences

were determined by two-tailed independent Student's t-test. **, $P < 0.01$.

Fig. S3. RT-PCR analyses of Hh pathway genes in TSL cells. *β-actin* served as a

loading control, and RNAs from TSL cells were used as templates for negative

controls. The numbers in parentheses indicate the PCR cycles.

Fig. S4. Establishment of the TSL *ptch1*, *ptch2*, *gli1*, *gli2*, *gli3* and *sf1* mutant

lines.

(A-F) Schematic representation of gRNAs targeting the *ptch1*, *ptch2*, *gli1*, *gli2*, *gli3*

and *sf1* locus. The gRNAs were designed to target ORF (open reading frame). The

PAM (protospacer adjacent motif) sites were marked by red letter. Sanger sequencing

of *ptch1*, *ptch2*, *gli1*, *gli2*, *gli3* and *sf1* single mutant alleles in single cell clones. PCR

amplicons from DNA templates of the six cell clones were directly used for

sequencing.

Fig. S5. Establishment of the Nile tilapia *ptch2* mutant line by CRISPR/Cas9

system.

(A) Schematic representation of gRNA targeting the Nile tilapia *ptch2* locus. The

gRNA was designed to target exon 1. The translation start codon ATG and stop codon

TGA were indicated by arrows. The PAM site was marked by green box. DNA

sequence alignment of the five mutant lines with WT. The added sequences were

marked by red letter, and the deleted sequences were indicated in red dotted lines.

(B) Schematic diagram showing the breeding plans of *ptch2* F0 to F2 fish.

(C) DNA sequencing showed the 25-bp deletion within the *ptch2* ORF in the

homozygous mutant compared to the WT.

(D) Schematic of the prediction of the intact Ptch2 protein in the WT and the

truncated Ptch2 protein in the homozygous mutants.

(E) Homozygous mutants of F2 fish were screened by PAGE. The first lane is DNA

marker, the second lane is $ptch2^{+/+}$ fish, the third lane is $ptch2^{+/-}$ fish, and the fourth

lane is $ptch2^{-/-}$ fish. "*" indicates heteroduplex, and arrows indicate homoduplex.

Fig. S6. Transcriptome data analyses and verification by RT-qPCR.

(A) Hierarchical clustering analysis of global gene expression patterns in TSL-WT,

TSL-*On*Dhh, TSL-*On*Gli1 and TSL+ 0.5 μM SAG. Blue indicates decreased

expression, and red indicates increased expression.

(B) Heatmap showing the expression patterns of upregulated differentially expressed

genes (DEGs) identified in Fig. 5C. The FPKM value for each gene in each sample is

indicated within the squares. The color gradient from blue to red reflects low to high

expression levels per row (gene).

(C) qRT-PCR validation. Values were presented as mean ± SD (n = 3).

Fig. S7. The expression levels of Cyp11c1 in the testis and the levels of 11-KT in

tissue fluid from WT and $dhh^{-/-}$ XY fish at 5, 10, 20, and 30 dah.

(A-E) Representative immunofluorescence images showing Cyp11c1 expression

(green) in testis sections from WT fish at 5, 10, 20, 30 dah and $dhh^{-/-}$ mutants at 30

dah. Scale bars, 4 μm.

(F) Tissue fluid 11-KT levels in WT and $dhh^{-/-}$ XY fish at indicated time points (*n*=6

fish per group). Values were presented as mean ± SD. Differences were determined by

two-tailed independent Student's t-test. NS, no significant difference.

Fig. S8. Validation of key gene expression changes in mutant and overexpression

TSL cell lines. (only for reviewer)

(A) Relative mRNA expression of *sf1* in TSL-WT, TSL-*gli1*$^{-/-}$, TSL-*gli2*$^{-/-}$ and TSL-

gli3^{-/-} cells by RT-qPCR. Values were presented as mean ± SD (*n*=3). Different letters

above the error bar indicate statistical differences at *P* < 0.05 as determined by one-

way ANOVA followed by Tukey test.

(B) Relative mRNA expression of *dhh* in TSL-WT and TSL-*On*Dhh cells by RT-

qPCR. Values were presented as mean ± SD (*n*=3). Differences were determined by

two-tailed independent Student's t-test. **, *P* < 0.01.

(C) Relative mRNA expression of *sf1* in TSL-WT and TSL-*On*Sf1 cells by RT-qPCR.

Values were presented as mean ± SD (*n*=3). Differences were determined by two-

tailed independent Student's t-test. **, *P* < 0.01.

Fig. S9. Negative control experiments for RNA fluorescence in situ hybridization

(FISH). (only for reviewer)

Co-staining of Cyp11c1 (by immunofluorescence) and sense RNA probes

for *ptch1*, *ptch2*, *gli1*, *gli2*, or *gli3* (by FISH) on testis sections from 90 dah fish. The

sense probes serve as negative controls to confirm the specificity of the antisense

probe hybridization signals shown in the main figures. Scale bars, 4 μm.

| | WT | | dhh⁻/⁻ | | |