# Assessing the Suitability of Deubiquitylases As Substrates For Targeted Protein Degradation

Joel Tong[1,3], J. Monty Watkins[2,3], James M. Burke[2,3], Thomas Kodadek[1,3*]

[1]Department of Chemistry, The Herbert Wertheim UF Scripps Institute for Biomedical Innovation & Technology, Jupiter, FL, USA
[2]Department of Molecular Medicine, The Herbert Wertheim UF Scripps Institute for Biomedical Innovation & Technology, Jupiter, FL, USA
[3]Skaggs Graduate School of Chemical and Biological Sciences, Jupiter, FL, USA
*Correspondence: kodadek@ufl.edu

## Significance

The development of selective inhibitors of Deubiquitylase enzymes (DUBs) is difficult due to a high level of homology in the active sites of the ≈ 100 such enzymes in the human proteome. A potentially attractive strategy to achieve this goal would be to develop PROTACs or molecular glues that engage the target DUB in a less conserved region outside of the catalytic domain. However, this raises the concern that auto-deubiquitylation would make DUBs poor substrates for this modality. Here we describe a chemical genetics system to evaluate this issue. We find that some Dubs are readily degradable via the Ubiquitin-proteasome pathway and some are not. Of the latter category, some resist turnover through auto-deubiquitylation and some are simply poor proteasome substrates.

## Highlights

- Some active DUBs, including USP11 are vulnerable to targeted protein degradation by a PROTAC.
- Other DUBs, including USP4 and USP15, resist PROTAC-mediated degradation via auto-deubiquitylation.
- Some DUBs are poor substrates for targeted degradation because they are processed inefficiently by the proteasome.

## Summary

Deubiquitylases (DUBs) are a family of specialized proteases that hydrolyze the isopeptide bond between a lysine and the C-terminal carboxylate of Ubiquitin. DUBs are involved in a myriad of cellular processes and many are attractive drug targets. However, it has proven extremely difficult to develop selective inhibitors due to the high degree of homology between DUB active sites. Targeted protein degradation using a proteolysis-targeting chimera (PROTAC) that recognizes the DUB in a less conserved region outside of the catalytic domain constitutes an attractive alternative strategy for selectively inhibiting a given DUB. Such ligands are unlikely to block the catalytic activity of the enzyme, raising the concern that auto-deubiqtuiylation will make DUBs inherently poor substrates for PROTACs of this type. Since drug-like ligands that engage DUBs outside of the active site are extremely rare, this issue is difficult to address in a straightforward fashion. In this study we establish a generally applicable chemical genetics workflow to evaluate the degradability of DUBs by a PROTAC. The data indicate that some DUBs are readily degradable and some are not. In particular, USP11, an attractive drug target in various cancers and Alzheimer's disease, is shown to be rapidly degradable, while its paralogs, USP4 and USP15 resist degradation through auto-deubiquitylation.

**Keywords**

PROTAC • Deubiquitylase • targeted protein degradation • selective inhibitor • chemical genetics

**Introduction**

Ubiquitylation is a key post-translational modification that has diverse functional outcomes, the most well-known being K48-linked ubiquitylation that targets a protein for proteasomal degradation[1]. This is a reversible post-translational modification (PTM). Ubiquitylation is mediated by E1, E2 and E3 ubiquitin ligases, which create an isopeptide bond between a lysine side chain and the C-terminal carboxylate of Ubiquitin, while deubiquitylases (DUBs) are proteases that cleave these iso-peptide bonds[2]. There are almost 100 DUBs encoded in the human proteome, and many of these are critical players in disease progression. For example, USP7 reduces the stability of the tumor suppressor p53 by deubiquitylating its E3 ligase MDM2[3,4]. USP11 has been shown to promote tumorigenesis by deubiquitylating and hence stabilizing oncogene products such as c-Myc and PGAM5[5,6]. Beyond oncology, USP30 has been linked to Parkinson's disease due to its role in counteracting Parkin-mediated mitophagy[7], while inhibition of USP14 was demonstrated to enhance turnover of the aggregation-prone Tau

protein that is relevant in Alzheimer's disease[8]. Hence, many DUBs represent attractive drug targets.

As most DUBs are cysteine proteases, it is relatively straightforward to develop a small molecule inhibitor targeting the active site. However, achieving selectivity, particularly across the large USP family, is a major challenge as these DUBs share a high degree of homology in their catalytic domain. This obstacle has been overcome in a few cases, for example the development of highly selective, active site-targeted inhibitors of USP7, but these were the result of major medicinal chemistry efforts[9-11]. Recently, selective inhibitors for USP9X[12] and USP30[13] have been reported, but these were also highly labor- and time-intensive endeavors. There is therefore a pressing need for more facile development of DUB therapeutics.

Buhrlarge and co-workers reported an interesting chemoproteomics platform to address this selectivity challenge[14]. Compounds in their library had an electrophilic warhead to target the catalytic cysteine and various heterocyclic building blocks to target the DUB at a less conserved surface some distance away from the active site. Different linkers were used to connect the two. Through this work, they identified a selective inhibitor of VCPIP1 with an $IC_{50}$ of 70 nM. However, it is notable that most of the compounds screened had poor selectivity, especially across the USP family.

An alternative approach that would avoid this daunting selectivity problem would be to employ a targeted protein degradation strategy using a proteolysis-targeting chimera (PROTAC) that engages the DUB of interest in a less conserved domain away from the catalytic site. For example, USP11 has a DUSP domain that is present in only seven out of the approximately 100 DUBs. However, it is reasonable to anticipate ligands that engage a DUB away from the active site will not inhibit its catalytic activity. This, in turn, raises the obvious concern that DUBs will be poor substrates for this type of PROTAC due to auto-deubiquitylation, which is a known phenomenon for some DUBs.[15-19] Unfortunately, a straightforward assessment of this issue is complicated by the fact that there are almost no DUB-binding small molecules that engage the protein outside of the active site. Therefore, we sought to develop a chemical genetics-based workflow to address this issue and "de-risk" PROTAC development campaigns against a DUB of interest. In this study, we use the dTAG system[20] to investigate the degradability of catalytically active DUBs and specifically assess the degree to which auto-deubiquitylation acts to shield these enzymes from targeted protein degradation. We show that a FKBP12$^{F36V}$-USP11 fusion

protein is highly degradable by the PROTAC dTAG-13, while FKBP12[F36V] fusions of USP4, USP15 and UCHL1 are only modestly degraded. For USP4 and USP15, that this resistance to degradation is largely due to auto-deubiquitylation, whereas UCHL1 appears to simply be a poor substrate for proteasome-mediated proteolysis due to its compact structure. Based on these results, we propose three main categories for DUBs with respect to their suitability as PROTAC substrates and discuss the possibility of exploiting auto-deubiquitylation to selectively degrade USP11 over its paralogs USP4 and USP15.

Results

**FKBP*-USP11 is readily degradable by the PROTAC dTAG-13**

To interrogate the degradable DUB-ome, one approach would be to utilize a pan-DUB PROTAC that recruit DUBs to an E3 ubiquitin ligase and assess degradation via a chemoproteomic workflow. This strategy has previously been adopted to probe the degradability of various kinases[21-23]. However, all currently available pan-DUB ligands are active site-targeted inhibitors[24,25], which would preclude the assessment of DUB degradability when the DUB is not inhibited. In addition, as mentioned above, there are almost no selective, non-inhibiting DUB ligands that would allow us to test each DUB individually.

Due to the dearth of suitable tool compounds, we decided to adopt the dTAG technology for our investigation. Briefly, a mutant FKBP tag (FKBP12[F36V]), henceforth referred to as FKBP*, is fused to a protein of interest (POI). The chemical dimerizer dTAG-13 binds to the FKBP* and the E3 ligase Cereblon, inducing proximity between the two. This leads to the poly-ubiquitylation and subsequent proteasomal degradation of the POI[20]. In this way, the POI is targeted for degradation without disrupting its catalytic activity. FKBP* was fused to the DUBs USP11, USP4, USP15 and UCHL1. An FKBP fusion with HDAC1 was also constructed. All of the constructs also included an N-terminal FLAG tag. The DUBs USP11, USP4, USP15 and UCHL1 were chosen because they are known to play important roles in cancer.[5,6,26-31] HDAC1 was chosen as a control because it is an amide hydrolase, but not a DUB, and is readily degradable using the dTAG system.[20] These FKBP- and FLAG-tagged fusion proteins were stably expressed in Flp-In 293 cells. Upon treatment of these cells with dTAG-13, degradation of the FLAG-FKBP*-DUB constructs was evaluated by Western blotting using anti-FLAG antibody.

As expected, there was significant degradation of the control construct FKBP*-HDAC1 at an optimal concentrations of dTAG-13 (167 nM; Figure 1A). At higher dTAG-13 concentrations,

degradation was much less efficient due to the hook effect, where the PROTAC primarily forms unproductive binary complexes instead of a ternary complex.[32] FKBP*-HDAC1 degradation by dTAG-13 was also fast, with depletion to 25 % of DMSO levels observed within two hours (Figures 1B).

The FKBP*-USP11 fusion protein was also degraded efficiently. Across all the concentrations of dTAG-13 tested, protein levels were less than 25 % of that in vehicle-treated cells after four hours, reaching as low as 2.5 % with 55.6 nM dTAG-13 (Figure 1C). This degradation was rapid. DUB levels were about 17 % relative to those seen in vehicle-treated cells after just two hours of dTAG-13 treatment (Figure 1D). Co-addition of the E1 inhibitor TAK-243 (Figure 1E) or the proteasome inhibitor Bortezomib (Figure 1F) with dTAG-13 abolished FKBP*-USP11 turnover. These data indicate that USP11, at least in the context of this system, is readily degradable by a PROTAC via the Ubiquitin-proteasome pathway.

**FKBP*-USP11 is active**

The fact that FKBP*-USP11 was so readily degradable is significant as it is a high priority cancer target. However, a concern is that the fusion protein has reduced catalytic activity, allowing it to be degraded more readily than would be the case for native USP11. The deubiquitylase activity of USP11 plays an important role in DNA double-stranded break (DSB) repair.[33,34] Cortez and co-workers have shown that knocking down USP11 alone, even in the absence of ionizing radiation, was sufficient to cause a three- to four-fold increase in $\gamma$-H2AX foci[33], which is a marker of unrepaired DSBs[35]. We thus assessed if FKBP*-USP11 complements the loss of USP11 in DSB repair after knockdown of the endogenous protein.

Endogenous USP11 was knocked down using siRNA targeting the untranslated region of USP11 (UTR1 siRNA). In one set of cells, FKBP*-USP11 was also depleted by treatment with dTAG-13, while in a parallel set of cells FKBP*-USP11 levels were left unperturbed. If FKBP*-USP11 is able to function in DSB repair, we would expect to see a higher level of $\gamma$-H2AX foci in the cells where both proteins were knocked down than in cells where only endogenous USP11 was depleted. We confirmed by Western Blot that endogenous USP11 species was depleted in each sample (Figures 2A and B). We found that there was a marked increase in $\gamma$-H2AX foci when both endogenous USP11 and FKBP*-USP11 levels were depleted, as expected. In addition, the more compact nuclei in these cells could be attributed to chromatin condensation,

which is an integral component of the DNA damage response[36]. However, in cells where only endogenous USP11, but not FKBP*-USP11, was depleted, this effect was attenuated (Figures 2C and D, Figures S1A and B). These data show that FKBP*-USP11 can buffer the deleterious loss of endogenous USP11, suggesting that it is a functional protein with respect to DSB repair.

Additionally, the interactome of FKBP*-USP11 was probed to assess if some of the known interaction partners of endogenous USP11 also associate with FKBP*-USP11. To do this, FLAG-tagged FKBP*-USP11 was immunoprecipitated from a cell lysate using anti-FLAG antibody. The precipitated proteins were eluted from the matrix and, after trypsinization, the resultant peptide mixture was analyzed by mass spectrometry. An equivalent experiment was done with FKBP*-UCHL1 as a control. Across two biological replicates, USP7, which is a known interactor of USP11[37], co-precipitated with FKBP*-USP11 but not FKBP*-UCHL1. PRKDC was also pulled down, which is of interest because like USP11, it is also involved in DNA double-strand break (DSB) repair, being part of the kinase complex responsible for the phosphorylation of histone H2AX[38] (Data S1).

Taken together, these data argue that FKBP*-USP11 retains some of the known functions of USP11, suggesting that it is a reasonable surrogate for the endogenous protein.

**USP4, USP15 and UCHL1 fusion proteins are degraded by dTAG-13 less efficiently**
Next, the degradability of the FKBP*-USP4, -USP15 and -UCHL1 fusions was evaluated. In contrast to USP11, there was only moderate degradation of these constructs after four hours across different concentrations of dTAG-13 (Figures 3A, 3C and 3E). In cells treated with 500 nM dTAG-13, FKBP*-USP4 levels were about 63 % that of vehicle-treated cells (Figure 3A). For FKBP*-USP15 and FKBP*-UCHL1, protein levels of dTAG-13-treated cells were about 65 % and 47 % that of vehicle-treated cells respectively (Figures 3C and 3E). A time course experiment revealed slow degradation of all three constructs (Figures 3B, 3D and 3F). Both FKBP*-USP4 and FKBP*-USP15 levels were greater than 80 % that of vehicle-treated cells after two hours, in stark contrast to the rapid degradation of FKBP*-USP11. Only minimal degradation of FKBP*-UCHL1 was observed two hours after dTAG-13 treatment. More significant degradation was observed at longer times. After 24 hours, FKBP*-USP4, FKBP*-USP15 and FKBP*-UCHL1 levels had dropped to around 9 %, 23 % and 28 %, respectively, of that observed in vehicle-treated cells (Figures 3G – L). Overall, these data indicate that the

FKBP* fusions to USP4, USP15 and UCHL1 can be degraded by PROTACs but less rapidly and efficiently than FKBP*-USP11.

**FKBP*-USP4 and -USP15 actively resist degradation through auto-deubiquitylation**

To investigate whether the less robust degradation of USP4, USP15 and UCHL1 was due to auto-deubiquitylation, we made mutant constructs in which the catalytic cysteine was replaced with a serine or alanine residue (FKBP*-USP4(C311S), FKBP*-USP15(C298A) and FKBP*-UCHL1(C90A)). These mutant DUBs were then also stably expressed in Flp-In 293 cells.

Treatment of both FKBP*-USP4(C311S) and FKBP*-UCHL1(C90A)-expressing cells with dTAG-13 resulted in greater depletion of the mutant construct compared to wild type DUB across all the concentrations tested (Figure 4 A and 4C). At 500 nM dTAG-13, mutant FKBP*-USP4(C311S) levels were about 8 % of that in vehicle-treated cells. In contrast the level of the wild-type fusion protein was reduced to 63% of that seen in vehicle-treated cells under the same conditions. Similarly, mutant FKBP*-USP15(C298A) levels were about 19% of that in vehicle-treated cells, compared to 65 % for the wild-type construct. We also observed a more rapid rate of degradation of both mutant fusions (Figures 4B and 4D). After two hours, FKBP*-USP4(C311S) levels were approximately 39 % of that observed in vehicle-treated cells, whereas levels of wild-type FKBP*-USP4 were about 86 % of the vehicle control. For FKBP*-USP15(C298A), protein levels were about 27 % of that in vehicle-treated cells compared to about 85 % for wild-type FKBP*-USP15.  These data strongly support the idea that catalytically active FKBP*-USP4 and FLBP*-USP15 resist PROTAC-mediated degradation through auto-deubiquitylation.

**UCHL1 is degraded slowly because it is a poor proteasome substrate**

When FKBP*-UCHL1(C90A)-expressing cells were treated with dTAG-13, we observed that after four hours the levels of the wild-type and mutant fusion proteins were quite similar (Figure 5A). The time course also revealed a comparable rate of degradation between the wild type and catalytically inactive DUB (Figure 5B). Hence, unlike USP4 and USP15, auto-deubiquitylation does not seem to be the main factor contributing to poor degradation of UCHL1.

An alternative reason for the poor degradability of FKBP*-UCHL1 could be the lack of an unstructured tail on this protein, a feature important for rapid turnover of proteins by the proteasome.[39,40] An unstructured region allows facile engagement of the substrate by the proteasomal ATPases that mediate protein unwinding and feed the resultant unstructured chain into the catalytic cavity of the complex. UCHL1 is a structurally characterized, tightly folded protein lacking such an unstructured region[41].

To test this hypothesis, a 35 amino acid disordered region derived from *S. cerevisiae* Cytochrome b$_2$ was added to the C-terminus of FKBP*-UCHL1. This tail has been shown previously to significantly improve the degradability of hard-to-unwind substrates[42]. Indeed, this FKBP*-UCHL1-35 construct was degraded more efficiently than the parent construct. For example, at 55.6 nM dTAG-13, the levels of FKBP*-UCHL1 and FKBP*-UCHL1-35, were 47% and 21%, respectively, of that observed in vehicle-treated cells (Figure 5C). The time course also revealed that the FKBP*-UCHL1-35 construct was degraded more quickly. After just two hours, the level of FKBP*-UCHL1-35 was about 50 % that of vehicle-treated cells, compared to 95 % for unmodified FKBP*-UCHL1 (Figure 5D). These data strongly suggest that the lack of a disordered tail is a significant contributor to sluggish FKBP*-UCHL1 degradation.

To further probe this model, we examined the effect of a p97 inhibitor on the turnover of FKBP*-UCHL1. For protein substrates that lack a disordered tail, the p97 unfoldase complex is responsible for generating the required unstructured region for proteasomal engagement.[43,44] We therefore reasoned that the slower, but measurable, degradation of FKBP*-UCHL1 might require p97 activity, whereas the more rapid degradation of FKBP*-UCHL1-35 may not. Thus, we co-treated FKBP*-UCHL1- and FKBP*-UCHL1-35-expressing cells with dTAG-13 and the p97 inhibitor CB-5083 and assessed the levels of FKBP* fusion protein. As expected, FKBP*-UCHL1 degradation was abolished by the addition of CB-5083, indicating a strong dependence on p97 for degradation (Figure 6). Unexpectedly however, the degradation of FKBP*-UCHL1-35 was also abolished when CB-5083 was added, suggesting that the degradation of FKBP*-UCHL1-35 is also dependent on p97 (Figure 6). While tangential to the main point of this study, this finding challenges the current paradigm that the main role of p97 in degrading tightly folded proteins and that it is not essential for turnover of substrates with an intrinsically disordered tail.[43-45] This is considered further in the discussion section.

**Discussion**

In this work, we employed dTAG technology[20] to evaluate the degradability of the DUBs USP11, USP4, UCHL1 and USP15 when engaged by a PROTAC that does not block the active site. Based on our results, we propose three main categories for DUBs with respect to their suitability as targets of PROTAC-style degraders. The first are DUBs that, like USP11, are highly degradable. We found that FKBP*-USP11 levels are reduced rapidly to <10% that observed in vehicle-treated cells (Figure 1). The second comprises DUBs, such as USP4 and USP15, that actively resist degradation through auto-deubiquitylation. These fusion proteins were degraded more slowly and less efficiently than USP11. However, inactivation of their protease activity through mutation of the active site cysteine rendered them more susceptible to degradation (Figures 3 and 4). The third category includes UCHL1, which is degraded inefficiently (Fig. 3) but not because of its catalytic activity. In this case, the protein is simply a poor proteasome substrate due to the absence of an unstructured tail that can be easily engaged by the AAA class ATPases in the proteasome that are responsible for substrate unfolding.  This view is supported by the observation that addition of a 35 amino acid unstructured tail to FKBP*-UCHL1 strongly stimulated its degradation upon exposure of cells to dTAG-13 (Fig. 5).

A side note that is peripheral to this investigation, but of interest with respect to the fundamental mechanism of protein turnover in cells, is our observation that even FKBP*-UCHL1-35 is highly dependent on p97 for its degradation (Figure 6). As mentioned above, this runs counter to current dogma in the field that suggests p97 is dispensable for the turnover of proteins with an unstructured tail. This could be explained by UCHL1's rather unusual structure. Its backbone forms a 5-2 knot[46], such that pulling on either terminus will still lead to a knotted structure. Thus, even if the AAA-ATPases of the proteasome engage FKBP*-UCHL1-35, the unwinding process is likely to stall and require p97 to unknot the protein.

Returning to the main theme of this study, we believe that this workflow is useful to de-risk a PROTAC development against a particular DUB of interest since the development of high quality ligands to these targets is a significant undertaking. In this vein, USP11-class DUBS, which are readily degraded, are especially interesting as these are presumably the preferred targets for non-inhibitory PROTACs. However, as is the case in any chemical genetics experiment, care needs to be taken in applying conclusions reached through analysis of a fusion construct to the native protein. In this case, a particular concern is that the fusion of the

FKBP mutant to USP11 compromises the catalytic activity of the protein and that its sensitivity to PROTAC-mediated degradation is an artifact.

This seems unlikely for several reasons. First, we demonstrated that FKBP*-USP11 is able to substitute for the endogenous protein in the repair of DNA double-stranded breaks, as monitored by an accumulation of γ-H2AX foci that are a marker of unrepaired breaks (Fig. 2). The DUB activity of USP11 is essential for this process, so this observation argues that FKBP*-USP11 is an active DUB. Second, we showed that FKBP*-USP11 associates with USP7, a known interaction partner of USP11, as well as PRKDC, a component of the DSB repair complex. We had considered also evaluating immunoprecipitated FKBP*-USP11 for DUB activity using a fluorogenic assay but this is complicated by the presence of other DUBs, including USP7 in the immunoprecipitate. Finally, USP11 is a paralog of USP4 and USP15, sharing 45 % and 46 % identity with these proteins, respectively. We found that both USP4 and USP15 resist turnover through auto-deubiquitylation, so it seems unlikely that the identical fusion to USP11 would compromise its catalytic activity. Taken together, these data strongly suggest that FKBP*-USP11 is an active DUB. However, we acknowledge that it is impossible to say unequivocally that the sensitivity of FKBP*-USP11 to PROTAC-mediated turnover will indeed reflect that of native USP11 once appropriate ligands for the construction of non-inhibitory PROTACs become available. Our sole contention is that the findings shown here de-risk this activity considerably.

Again, with the caveat that we are looking at fusion proteins, the significant difference observed in the degradability of the FKBP*-USP11, -USP4 and -USP15 is interesting from the point of view of developing selective degraders of USP11. These proteins are very closely related paralogs and it may be difficult to identify compounds that distinguish cleanly between them even using ligands that engage the protein outside the active site. However, the data reported here suggest the possibility that even a PROTAC that engages all three paralogs might display significant selectivity for USP11 degradation if it is the least efficient of three in deubiquitylating itself. There are several examples of large divergences between proteins that are engaged by a PROTAC and those that are degraded[21,22]. But these are attributed to differences in the efficiency of poly-Ubiquitylation of the substrate by the E3 ligase. This issue of more or less efficient auto-deubiquitylation would represent a new mechanism for achieving selectivity.

**Resource Availability**

Lead contact

Requests for further information should be directed to the lead contact, Thomas Kodadek (kodadek@ufl.edu)

Materials availability

All plasmids and stable cell lines generated in this study are available from the lead contact upon request.

Data and code availability

All data are available in the main text or supporting information. Any additional information is available from the lead contact upon request. This paper does not report any original code.

**Acknowledgements**

This work was supported by a grant from the U.S. National Cancer Institute (RO1 CA290247). We would like to thank the Mass Spectrometry and Proteomics Core Facility at the Herbert Wertheim UF Scripps Institute for Biomedical Innovation & Technology for assistance with the FKBP*-USP11 interactome experiment.

**Author contributions**

Both T.K. and J.T. conceived this study. J.T. performed most of the experiments, with help from J.M.W. for the imaging experiment and the Proteomics Core for the FKBP*-USP11 interactome experiment. Both T.K. and J.T. wrote and edited the manuscript. The entire study was supervised by T.K. and J.M.B.

**Declaration of interests**

T.K. is a significant shareholder in Triana Biomedicines.

**References**

1.    Rape, M. (2018). Ubiquitylation at the crossroads of development and disease. Nat Rev Mol Cell Biol *19*, 59-70. 10.1038/nrm.2017.83.
2.    Lange, S.M., Armstrong, L.A., and Kulathu, Y. (2022). Deubiquitinases: From mechanisms to their inhibition by small molecules. Mol Cell *82*, 15-29. 10.1016/j.molcel.2021.10.027.

3.      Li, M., Brooks, C.L., Kon, N., and Gu, W. (2004). A Dynamic Role of HAUSP in the p53-Mdm2 Pathway. Molecular Cell *13*, 879–886. https://doi.org/10.1016/S1097-2765(04)00157-1.

4.      Cummins, J.M., Rago, C., Kohli, M., Kinzler, K.W., Lengauer, C., and Vogelstein, B. (2004). Tumour suppression: disruption of HAUSP gene stabilizes p53. Nature *428*, 1 p following 486. 10.1038/nature02501.

5.      Pornour, M., Jeon, H.Y., Ryu, H., Khadka, S., Xu, R., Chen, H., Hussain, A., Lam, H.M., Zhuang, Z., Oo, H.Z., et al. (2024). USP11 promotes prostate cancer progression by up-regulating AR and c-Myc activity. Proc Natl Acad Sci U S A *121*, e2403331121. 10.1073/pnas.2403331121.

6.      Zhang, N., Wang, Q., Lu, Y., Wang, F., and He, Z. (2024). The deubiquitinating enzyme USP11 regulates breast cancer progression by stabilizing PGAM5. Breast Cancer Res *26*, 135. 10.1186/s13058-024-01892-9.

7.      Bingol, B., Tea, J.S., Phu, L., Reichelt, M., Bakalarski, C.E., Song, Q., Foreman, O., Kirkpatrick, D.S., and Sheng, M. (2014). The mitochondrial deubiquitinase USP30 opposes parkin-mediated mitophagy. Nature *510*, 370-375. 10.1038/nature13418.

8.      Lee, B.H., Lee, M.J., Park, S., Oh, D.C., Elsasser, S., Chen, P.C., Gartner, C., Dimova, N., Hanna, J., Gygi, S.P., et al. (2010). Enhancement of proteasome activity by a small-molecule inhibitor of USP14. Nature *467*, 179-184. 10.1038/nature09299.

9.      Turnbull, A.P., Ioannidis, S., Krajewski, W.W., Pinto-Fernandez, A., Heride, C., Martin, A.C.L., Tonkin, L.M., Townsend, E.C., Buker, S.M., Lancia, D.R., et al. (2017). Molecular basis of USP7 inhibition by selective small-molecule inhibitors. Nature *550*, 481-486. 10.1038/nature24451.

10.     Kategaya, L., Di Lello, P., Rouge, L., Pastor, R., Clark, K.R., Drummond, J., Kleinheinz, T., Lin, E., Upton, J.P., Prakash, S., et al. (2017). USP7 small-molecule inhibitors interfere with ubiquitin binding. Nature *550*, 534-538. 10.1038/nature24006.

11.     Schauer, N.J., Liu, X., Magin, R.S., Doherty, L.M., Chan, W.C., Ficarro, S.B., Hu, W., Roberts, R.M., Iacob, R.E., Stolte, B., et al. (2020). Selective USP7 inhibition elicits cancer cell killing through a p53-dependent mechanism. Sci Rep *10*, 5324. 10.1038/s41598-020-62076-x.

12.     Clancy, A., Heride, C., Pinto-Fernandez, A., Elcocks, H., Kallinos, A., Kayser-Bricker, K.J., Wang, W., Smith, V., Davis, S., Fessler, S., et al. (2021). The deubiquitylase USP9X controls ribosomal stalling. J Cell Biol *220*. 10.1083/jcb.202004211.

13.     Fang, T.Z., Sun, Y., Pearce, A.C., Eleuteri, S., Kemp, M., Luckhurst, C.A., Williams, R., Mills, R., Almond, S., Burzynski, L., et al. (2023). Knockout or inhibition of USP30 protects dopaminergic neurons in a Parkinson's disease mouse model. Nat Commun *14*, 7295. 10.1038/s41467-023-42876-1.

14.     Chan, W.C., Liu, X., Magin, R.S., Girardi, N.M., Ficarro, S.B., Hu, W., Tarazona Guzman, M.I., Starnbach, C.A., Felix, A., Adelmant, G., et al. (2023). Accelerating inhibitor discovery for deubiquitinating enzymes. Nat Commun *14*, 686. 10.1038/s41467-023-36246-0.

15.     Meray, R.K., and Lansbury, P.T., Jr. (2007). Reversible monoubiquitination regulates the Parkinson disease-associated ubiquitin hydrolase UCH-L1. J Biol Chem *282*, 10567-10575. 10.1074/jbc.M611153200.

16. Denuc, A., Bosch-Comas, A., Gonzalez-Duarte, R., and Marfany, G. (2009). The UBA-UIM domains of the USP25 regulate the enzyme ubiquitination state and modulate substrate recognition. PLoS One *4*, e5571. 10.1371/journal.pone.0005571.

17. Wijnhoven, P., Konietzny, R., Blackford, A.N., Travers, J., Kessler, B.M., Nishi, R., and Jackson, S.P. (2015). USP4 Auto-Deubiquitylation Promotes Homologous Recombination. Mol Cell *60*, 362-373. 10.1016/j.molcel.2015.09.019.

18. Shen, C., Ye, Y., Robertson, S.E., Lau, A.W., Mak, D.O., and Chou, M.M. (2005). Calcium/calmodulin regulates ubiquitination of the ubiquitin-specific protease TRE17/USP6. J Biol Chem *280*, 35967-35973. 10.1074/jbc.M505220200.

19. Peng, Y., Liao, Q., Tan, W., Peng, C., Hu, Z., Chen, Y., Li, Z., Li, J., Zhen, B., Zhu, W., et al. (2019). The deubiquitylating enzyme USP15 regulates homologous recombination repair and cancer cell response to PARP inhibitors. Nat Commun *10*, 1224. 10.1038/s41467-019-09232-8.

20. Nabet, B., Roberts, J.M., Buckley, D.L., Paulk, J., Dastjerdi, S., Yang, A., Leggett, A.L., Erb, M.A., Lawlor, M.A., Souza, A., et al. (2018). The dTAG system for immediate and target-specific protein degradation. Nat Chem Biol *14*, 431-441. 10.1038/s41589-018-0021-8.

21. Huang, H.T., Dobrovolsky, D., Paulk, J., Yang, G., Weisberg, E.L., Doctor, Z.M., Buckley, D.L., Cho, J.H., Ko, E., Jang, J., et al. (2018). A Chemoproteomic Approach to Query the Degradable Kinome Using a Multi-kinase Degrader. Cell Chem Biol *25*, 88-99 e86. 10.1016/j.chembiol.2017.10.005.

22. Bondeson, D.P., Smith, B.E., Burslem, G.M., Buhimschi, A.D., Hines, J., Jaime-Figueroa, S., Wang, J., Hamman, B.D., Ishchenko, A., and Crews, C.M. (2018). Lessons in PROTAC Design from Selective Degradation with a Promiscuous Warhead. Cell Chem Biol *25*, 78-87 e75. 10.1016/j.chembiol.2017.09.010.

23. Donovan, K.A., Ferguson, F.M., Bushman, J.W., Eleuteri, N.A., Bhunia, D., Ryu, S., Tan, L., Shi, K., Yue, H., Liu, X., et al. (2020). Mapping the Degradable Kinome Provides a Resource for Expedited Degrader Development. Cell *183*, 1714-1731 e1710. 10.1016/j.cell.2020.10.038.

24. Tian, X., Isamiddinova, N.S., Peroutka, R.J., Goldenberg, S.J., Mattern, M.R., Nicholson, B., and Leach, C. (2011). Characterization of selective ubiquitin and ubiquitin-like protease inhibitors using a fluorescence-based multiplex assay format. Assay Drug Dev Technol *9*, 165-173. 10.1089/adt.2010.0317.

25. Conole, D., Cao, F., Am Ende, C.W., Xue, L., Kantesaria, S., Kang, D., Jin, J., Owen, D., Lohr, L., Schenone, M., et al. (2023). Discovery of a Potent Deubiquitinase (DUB) Small-Molecule Activity-Based Probe Enables Broad Spectrum DUB Activity Profiling in Living Cells. Angew Chem Int Ed Engl *62*, e202311190. 10.1002/anie.202311190.

26. Zhang, H., Han, Y., Xiao, W., Gao, Y., Sui, Z., Ren, P., Meng, F., Tang, P., and Yu, Z. (2023). USP4 promotes the proliferation, migration, and invasion of esophageal squamous cell carcinoma by targeting TAK1. Cell Death Dis *14*, 730. 10.1038/s41419-023-06259-0.

27.     Zhang, X., Berger, F.G., Yang, J., and Lu, X. (2011). USP4 inhibits p53 through deubiquitinating and stabilizing ARF-BP1. EMBO J *30*, 2177-2189. 10.1038/emboj.2011.125.

28.     Chen, X.S., Wang, K.S., Guo, W., Li, L.Y., Yu, P., Sun, X.Y., Wang, H.Y., Guan, Y.D., Tao, Y.G., Ding, B.N., et al. (2020). UCH-L1-mediated Down-regulation of Estrogen Receptor alpha Contributes to Insensitivity to Endocrine Therapy for Breast Cancer. Theranostics *10*, 1833-1848. 10.7150/thno.39814.

29.     Yao, J., Reyimu, A., Sun, A., Duoji, Z., Zhou, W., Liang, S., Hu, S., Wang, X., Dai, J., and Xu, X. (2022). UCHL1 acts as a potential oncogene and affects sensitivity of common anti-tumor drugs in lung adenocarcinoma. World J Surg Oncol *20*, 153. 10.1186/s12957-022-02620-3.

30.     Chen, L.L., Smith, M.D., Lv, L., Nakagawa, T., Li, Z., Sun, S.C., Brown, N.G., Xiong, Y., and Xu, Y.P. (2020). USP15 suppresses tumor immunity via deubiquitylation and inactivation of TET2. Science Advances *6*. 10.1126/sciadv.abc9730.

31.     Huangfu, L., Zhu, H., Wang, G., Chen, J., Wang, Y., Fan, B., Wang, X., Yao, Q., Guo, T., Han, J., et al. (2024). The deubiquitinase USP15 drives malignant progression of gastric cancer through glucose metabolism remodeling. J Exp Clin Cancer Res *43*, 235. 10.1186/s13046-024-03152-2.

32.     Cecchini, C., Pannilunghi, S., Tardy, S., and Scapozza, L. (2021). From Conception to Development: Investigating PROTACs Features for Improved Cell Permeability and Successful Protein Degradation. Front Chem *9*, 672267. 10.3389/fchem.2021.672267.

33.     Wiltshire, T.D., Lovejoy, C.A., Wang, T., Xia, F., O'Connor, M.J., and Cortez, D. (2010). Sensitivity to poly(ADP-ribose) polymerase (PARP) inhibition identifies ubiquitin-specific peptidase 11 (USP11) as a regulator of DNA double-strand break repair. J Biol Chem *285*, 14565-14571. 10.1074/jbc.M110.104745.

34.     Ting, X., Xia, L., Yang, J., He, L., Si, W., Shang, Y., and Sun, L. (2019). USP11 acts as a histone deubiquitinase functioning in chromatin reorganization during DNA repair. Nucleic Acids Res *47*, 9721-9740. 10.1093/nar/gkz726.

35.     Mah, L.J., El-Osta, A., and Karagiannis, T.C. (2010). gammaH2AX: a sensitive molecular marker of DNA damage and repair. Leukemia *24*, 679-686. 10.1038/leu.2010.6.

36.     Burgess, R.C., Burman, B., Kruhlak, M.J., and Misteli, T. (2014). Activation of DNA damage response signaling by condensed chromatin. Cell Rep *9*, 1703-1717. 10.1016/j.celrep.2014.10.060.

37.     Maertens, G.N., El Messaoudi-Aubert, S., Elderkin, S., Hiom, K., and Peters, G. (2010). Ubiquitin-specific proteases 7 and 11 modulate Polycomb regulation of the INK4a tumour suppressor. EMBO J *29*, 2553-2565. 10.1038/emboj.2010.129.

38.     Park, E.J., Chan, D.W., Park, J.H., Oettinger, M.A., and Kwon, J. (2003). DNA-PK is activated by nucleosomes and phosphorylates H2AX within the nucleosomes in an acetylation-dependent manner. Nucleic Acids Res *31*, 6819-6827. 10.1093/nar/gkg921.

39. Inobe, T., Fishbain, S., Prakash, S., and Matouschek, A. (2011). Defining the geometry of the two-component proteasome degron. Nat Chem Biol *7*, 161-167. 10.1038/nchembio.521.

40. Prakash, S., Tian, L., Ratliff, K.S., Lehotzky, R.E., and Matouschek, A. (2004). An unstructured initiation site is required for efficient proteasome-mediated degradation. Nat Struct Mol Biol *11*, 830-837. 10.1038/nsmb814.

41. Das, C., Hoang, Q.Q., Kreinbring, C.A., Luchansky, S.J., Meray, R.K., Ray, S.S., Lansbury, P.T., Ringe, D., and Petsko, G.A. (2006). Structural basis for conformational plasticityof the Parkinson's disease-associatedubiquitin hydrolase UCH-L1. Proceedings of the National Academy of Sciences of the United States of America *103*, 4675– 4680. https://doi.org/10.1073/pnas.0510403103.

42. Martinez-Fonts, K., Davis, C., Tomita, T., Elsasser, S., Nager, A.R., Shi, Y., Finley, D., and Matouschek, A. (2020). The proteasome 19S cap and its ubiquitin receptors provide a versatile recognition platform for substrates. Nat Commun *11*, 477. 10.1038/s41467-019-13906-8.

43. Beskow, A., Grimberg, K.B., Bott, L.C., Salomons, F.A., Dantuma, N.P., and Young, P. (2009). A conserved unfoldase activity for the p97 AAA-ATPase in proteasomal degradation. J Mol Biol *394*, 732-746. 10.1016/j.jmb.2009.09.050.

44. Olszewski, M.M., Williams, C., Dong, K.C., and Martin, A. (2019). The Cdc48 unfoldase prepares well-folded protein substrates for degradation by the 26S proteasome. Commun Biol *2*, 29. 10.1038/s42003-019-0283-z.

45. Li, H., Ji, Z., Paulo, J.A., Gygi, S.P., and Rapoport, T.A. (2024). Bidirectional substrate shuttling between the 26S proteasome and the Cdc48 ATPase promotes protein degradation. Mol Cell *84*, 1290-1303 e1297. 10.1016/j.molcel.2024.01.029.

46. Virnau, P., Mirny, L.A., and Kardar, M. (2006). Intricate knots in proteins: Function and evolution. PLoS Comput Biol *2*, e122. 10.1371/journal.pcbi.0020122.

**Figure 1. USP11 is readily degradable by the PROTAC dTAG-13.** (A and C) Representative western blot and quantification of FKBP*-HDAC1 (A) or FKBP*-USP11 (C) levels after HEK293 cells stably expressing the indicated FKBP* construct were treated for 4 h with a range of dTAG-13 concentrations (55.6 nM – 4.5 $\mu$M). Data are presented as mean ± SD (n = 3 biological replicates). (B and D) Representative western blot and quantification of FKBP*-HDAC1 (B) or FKBP*-USP11 (D) levels after HEK293 cells stably expressing the indicated FKBP* construct were treated with 500 nM dTAG-13 for 2 h, 4 h, 8 h or 24 h. Data are presented as mean ± SD (n = 3 biological replicates). (E and F) Representative western blot and quantification of FKBP*-USP11 levels after HEK293 cells stably expressing the indicated FKBP* construct were treated with 500 nM dTAG-13 ± 10 $\mu$M TAK-243 (E) or 10 $\mu$M Bortezomib (F) for 4 h. Data are presented as mean ± SD (n = 2 biological replicates).

**Figure 2. FKBP\*-USP11 is active.** (A) Western blot of FKBP\*-USP11 and endogenous USP11 levels in HEK293 cells stably expressing FKBP\*-USP11 after the indicated treatment. Cells were transfected with siRNA targeting the untranslated region of USP11 (UTR1 siRNA) or non-targeting siRNA (NT siRNA) (72 h incubation post-transfection) and treated with 500 nM dTAG-13 or vehicle control (24 h treatment). Blot is representative of two biological replicates. (B) Quantification of (A). Data are presented as mean ± SD (n = 2 biological replicates). (C and D) Representative images (C) and violin plot (D) of $\gamma$-H2AX levels in HEK293 cells stably expressing FKBP\*-USP11 treated as described in (A). Data are representative of two biological replicates.

**Figure 3. USP4, USP15 and UCHL1 are only modestly degraded by dTAG-13**
(A, C and E) Representative western blot and quantification of FKBP*-USP4 (A), FKBP*-USP15 (C) or FKBP*-UCHL1 (E) levels after HEK293 cells stably expressing the indicated FKBP* construct were treated for 4 h with a range of dTAG-13 concentrations (55.6 nM – 4.5 µM). Data are presented as mean ± SD (n = 3 biological replicates). (B, D and F) Representative western blot and quantification of FKBP*-USP4 (B), FKBP*-USP15 (D) or FKBP*-UCHL1 (F) levels after HEK293 cells stably expressing the indicated FKBP* construct were treated with 500 nM dTAG-13 for 2 h, 4 h, 8 h or 24 h. Data are presented as mean ± SD (n = 3 biological replicates).

**Figure 4. USP4 and USP15 actively resist degradation through auto-deubiquitylation**
(A and C) Representative western blot of FKBP*-USP4(C311S) (A) or FKBP*-USP15(C298A) (C) levels after HEK293 cells stably expressing the indicated FKBP* construct were treated for 4 h with a range of dTAG-13 concentrations (55.6 nM – 4.5 µM). Mutant DUB fusion levels are compared to the wild-type FKBP*-DUB construct. Data are presented as mean ± SD (n = 3 biological replicates), unpaired t test with Welch correction (*q < 0.05). (B and D) Representative western blot of FKBP*-USP4(C311S) (B) or FKBP*-USP15(C298A) (D) levels after HEK293 cells stably expressing the indicated FKBP* construct were treated with 500 nM dTAG-13 for 2 h, 4 h, 8 h or 24 h. Mutant DUB fusion levels are compared to the wild-type FKBP*-DUB construct. Data are presented as mean ± SD (n = 3 biological replicates), unpaired t test with Welch correction (*q < 0.05, nd: q ≥ 0.05).

**Figure 5. UCHL1 is degraded slowly because it is a poor proteasome substrate**

(A and C) Representative western blot of FKBP*-UCHL1(C90A) (A) or FKBP*-UCHL1-35 (C) levels after HEK293 cells stably expressing the indicated FKBP* construct were treated for 4 h with a range of dTAG-13 concentrations (55.6 nM – 4.5 μM). Protein levels are compared to FKBP*-UCHL1. Data are presented as mean ± SD (n = 3 biological replicates), unpaired t test with Welch correction (*q < 0.05, nd: q ≥ 0.05). (B and D) Representative western blot of FKBP*-UCHL1(C90A) (B) or FKBP*-UCHL1-35 (D) levels after HEK293 cells stably expressing the indicated FKBP* construct were treated with 500 nM dTAG-13 for 2 h, 4 h, 8 h or 24 h. Protein levels are compared to FKBP*-UCHL1. Data are presented as mean ± SD (n = 3 biological replicates), unpaired t test with Welch correction (*q < 0.05, nd: q ≥ 0.05).

**Figure 6. The degradation of both FKBP*-UCHL1 and FKBP*-UCHL1-35 is dependent on p97.**
(A and B) Western blot of FKBP*-UCHL1 (A) or FKBP*-UCHL1-35 (B) levels after HEK293 cells stably expressing the indicated FKBP* construct were treated with 500 nM dTAG-13 ± 10 µM CB-5083 for 8 h. Blots are representative of three biological replicates. (C) Quantification of (A) and (B). Data are presented as mean ± SD (n = 3 biological replicates), unpaired t test with Welch correction (*q < 0.05).

**'Assessing the Suitability of Deubiquitylases As Substrates For Targeted Protein Degradation'**
**STAR Methods**

## EXPERIMENTAL MODEL AND STUDY PARTICIPANT DETAILS

### Cell lines
Flp-In 293 cells were purchased from ThermoFisher. Flp-In 293 cells and stable Flp-In 293 cell lines expressing FKBP* constructs were grown in in DMEM (4.5 g/L D-Glucose) supplemented with 10 % FBS, 2 mM GlutaMAX and 100 $\mu$g/mL Zeocin (Flp-In 293) or 100 $\mu$g/mL Hygromycin B (stable Flp-In 293 cell lines). Cells were maintained at 37 $^{\circ}$C with 5 % $CO_2$.

## METHOD DETAILS

### Plasmid construction
The pcDNA5-FRT plasmid backbone was PCR amplified with indicated primers (IDT, see table S1) from the pcDNA5-FRT-HaloTag7 plasmid[1]. gBlocks Gene Fragments encoding FKBP* constructs (including overhangs) were purchased from IDT and inserted into the pcDNA5-FRT plasmid backbone by In-Fusion cloning (Takara). For site-directed mutagenesis, the template plasmid was PCR amplified with the indicated primers (IDT, see table S1) and re-circularized by In-Fusion cloning (Takara).

### Generation of Flp-In 293 cells stably expressing FKBP* constructs
Flp-In 293 cells in a 10 cm dish were co-transfected with 21.6 $\mu$g of pOG44 plasmid (ThermoFisher) and 2.4 $\mu$g of FKBP* construct plasmid using Lipofectamine 2000 (ThermoFisher). Media was refreshed after 24 hours. After an additional 24 hours, cells were split into four 10 cm dishes in media supplemented with 100 $\mu$g/mL Hygromycin B (ThermoFisher). After seven days, Hygromycin B-resistant cells were pooled and expanded. The stable cell lines were validated by Western Blotting (anti-FLAG).

### Degradation assay
Stable Flp-In 293 cells were seeded in 6-well plates with complete media lacking antibiotics and incubated at 37 $^{\circ}$C overnight. The next day, cells were treated with varying concentrations of dTAG-13 (4.5 $\mu$M – 55.6 nM) and incubated at 37 $^{\circ}$C for four hours before collection. For the time course, cells were treated with 500 nM dTAG-13 and incubated at 37 $^{\circ}$C for two, four, eight or 24 hours before collection.

### Evaluating dependence on the proteasome, ubiquitin or p97 for degradation
Stable Flp-In 293 cells were seeded in 6-well plates with complete media lacking antibiotics and incubated at 37 $^{\circ}$C overnight. The next day, cells were co-treated with 500 nM dTAG-13 and 10 $\mu$M Bortezomib (proteasome inhibitor), TAK-243 (ubiquitin activating enzyme UBA1 inhibitor) or CB-5083 (p97 inhibitor) and incubated at 37 $^{\circ}$C for four (Bortezomib and TAK-243) or eight hours (CB-5083) before collection.

**Western blotting**
Cells were lysed in M-PER Mammalian Protein Extraction Reagent + 1X Halt Protease
Inhibitor Cocktail (ThermoFisher) for 12 minutes and clarified by centrifugation at 14,000
xg for 15 minutes. Protein concentration was determined by Bradford assay and
normalized gel samples were prepared with 4X Laemmli sample buffer (Bio-Rad)
supplemented with 2-mercaptoethanol, DPBS and cell lysate. Gel samples were loaded
on 4 – 20 % TGX gels (Bio-Rad) and run at 180 V for 35 – 40 minutes. Gels were
transferred to a nitrocellulose membrane (Bio-Rad) using the TransBlot Turbo semi-dry
transfer system (mixed MW setting). Membranes were blocked with 5 % Blot-
QuickBlocker (G-Biosciences) in PBST for one hour and then incubated with primary
antibodies (see key resources table) in the same blocking buffer at 4 °C overnight. The
membranes were then washed with PBST (3 X 15 min) and incubated with secondary
antibodies (see key resources table) in Intercept Blocking Buffer (Li-Cor) at 25 °C for
one hour. Membranes were washed again with PBST (3 X 15 min) and imaged on a Li-
Cor Odyssey M scanner.

**USP11 depletion for γ-H2AX assay**
Flp-In 293 cells stably expressing FKBP*-USP11 were seeded in 6-well plates with
complete media lacking antibiotics and incubated at 37 °C overnight. The next day, the
cells were transfected with a final concentration of 50 nM custom USP11 UTR1 siRNA,
or siGENOME Non-Targeting siRNA (Horizon Discovery, see table S1) using
DharmaFECT 1 transfection reagent (Horizon Discovery). Media was refreshed after 24
hour incubation at 37 °C. After an additional 24 hours of incubation at 37 °C, 15 mm
round coverslips were placed into a 12-well plate and pre-treated with 70 % ethanol,
followed by Poly-L-Lysine solution (Sigma-Aldrich). The cells were then split into the 12-
well plate, with two wells for each experimental condition (one for imaging, one for
Western Blot analysis), and treated with 500 nM dTAG-13 or vehicle control. After 24
hours of incubation at 37 °C, the cells were either collected for western blot analysis or
used for immunofluorescent staining.

**Immunofluorescence imaging**
Cells in a 12-well plate were washed with PBS (X1) and fixed with 4 % formaldehyde in
PBS for 10 minutes at 25 °C. The cells were then washed with PBS (X1) and
permeabilized with cold 70 % ethanol for two hours at 4 °C. Cells were washed with
PBS (X3) and then incubated with γ-H2AX antibody (Cell Signaling Technology) in 1 %
BSA (PBS) at 4 °C overnight. The cells were then washed with PBS (3 X 15 min) and
incubated with α-rabbit secondary antibody (Abcam) in in 1 % BSA (PBS) at 4 °C for
two hours. Cells were then washed again with PBS (3 X 15 min). Cover slips were
mounted onto glass slides with VECTASHIELD PLUS mounting medium with DAPI
(Vector Laboratories) and dried. Cells were imaged on a Nikon Eclipse Ti2 with a CFI60
Plan Apochromat Lambda D 100x Oil Immersion Objective Lens, N.A. 1.45, W.D.
0.13mm, F.O.V. 25mm, DIC, Spring Loaded.

**FKBP*-USP11 / FKBP*-UCHL1 pulldown**
Flp-In 293 cells stably expressing either FKBP*-USP11 or FKBP*-UCHL1 were lysed
and clarified (see method details for western blotting). Protein concentration was

determined by Bradford assay and normalized samples were prepared with TBS. 60 ul of anti-FLAG M2 magnetic beads (Millipore) was equilibrated into TBS. 600 ug of protein/lysate was added to the beads and tumbled at 4 °C for two hours. The beads were washed with TBS (3 X 400 ul) and proteins were eluted with 2X Laemmli sample buffer (Bio-Rad). Eluted proteins were loaded on 4 – 20 % TGX gels (Bio-Rad) and run at 180 V for about five minutes. The gel was cut out and used for LC-MS/MS analysis.

**LC-MS/MS**
Gel fractionated proteins were in-gel digested with trypsin (Pierce Biotechnology) for 1 hour at 50 °C using ProteaseMax™ Surfactant trypsin enhancer following reduction and alkylation with dithiothreitol and iodoacetamide, respectively, according to the manufacturer's instructions (Promega). LC-MS/MS analysis of extracted peptides was subsequently carried out using an Orbitrap Fusion Tribrid mass spectrometer, following 2 mg capacity ZipTip (Millipore) C18 sample clean-up according to the manufacturer's instructions. Peptides were eluted from an EASY PepMapTM RSLC C18 column (Thermo) into the mass spectrometer using a gradient of 5 – 25 % solvent B (80/20 acetonitrile/water, 0.1% formic acid) in 90 minutes, followed by 25 – 44 % solvent B in 30 minutes, 44 – 80 % solvent B in 0.10 minute, a 10 minute-hold of 80 % solvent B, a return to 5 % solvent B in 3 minutes, and finally with another 3-minute hold of 5 % solvent B. The gradient was then extended for the purpose of cleaning the column by increasing solvent B to 98 % in 3 minutes, a 98 % solvent B hold for 10 minutes, a return to 5 % solvent B in 3 minutes, a 5 % solvent B fold for 3 minutes, an increase of solvent B to 98 % in 3 minutes, a 98 % solvent B hold for 10 minutes, a return to 5 % solvent B in 3 minutes and a 5 % solvent B hold for 3 minutes and finally, another increase to 98 % solvent B in 3 minutes and a hold of 98 % solvent B for 10 minutes. All flow rates were 250 nL/min delivered using an nEasy-LC1000 nano liquid chromatography system (Thermo). Solvent A consisted of 0.1% formic acid. Ions were created with an EASY Spray source (Thermo) held at 45 °C using a voltage of 2.5kV. Data dependent scanning was performed by the Xcalibur v 4.7.69.37 software using a survey scan at 120, 000 resolution in the Orbitrap analyzer scanning mass/charge (m/z) 350-2000 followed by higher-energy collisional dissociation (HCD) tandem mass spectrometry (MS/MS) at a normalized collision energy of 30% of the most intense ions at maximum speed, at an automatic gain control of 1.0E4. Precursor ions were selected by the monoisotopic precursor selection (MIPS) setting to peptide and MS/MS was performed on charged species of 2-8 charges at a resolution of 30,000. Dynamic exclusion was set to exclude ions once within a 25 second window. All scan events occurred within a 2-second specified cycle time.

**Protein/peptide identification**
Tandem mass spectra were searched against the human proteome protein sequences from Uniprot (UP000005640) downloaded on July 09, 2023, and common contaminant proteins available with Proteome Discoverer v 2.5.0.400. At the time of the search the UP0000005640 contained 20523 sequences, and the contaminant protein database contained an additional 298 sequences. All MS/MS spectra were searched using Thermo Proteome Discoverer 2.5.0.400 (Thermo) considering fully tryptic peptides with up to 2 missed cleavage sites. Variable modifications considered during

the search included methionine oxidation (15.995 Da), and asparagine and qlutamine deamidation (0.984 Da). Cysteine carbamidomethylation (57.021 Da) was considered as a static modification. Proteins were identified at 99% confidence with XCorr score cut-offs[2] as determined by a reversed database search. The Minora Feature Detector node which detects chromatographic peaks and features as also used in Proteome Discoverer at default settings. The protein and peptide identification results were also visualized with Scaffold v 5.0.0 (Proteome Software Inc), a program that relies on various search engine results (i.e.: Sequest, X!Tandem, MASCOT) and which uses Bayesian statistics to reliably identify more spectra[3]. Proteins were accepted that passed a minimum of two peptides identified at 1% peptide and protein FDR, within Scaffold. The mass spectrometry analysis was performed at The Herbert Wertheim UF Scripps Institute for Biomedical Innovation & Technology, Mass Spectrometry and Proteomics Core Facility (RRID:SCR_023576).

## QUANTIFICATION AND STATISTICAL ANALYSIS

Western blots were quantified using the Emperia Studio software (Li-Cor). Statical analysis was performed with Graphpad Prism 10; details can be found in the figure legends. For the microscopy experiment, two fields of view were used for quantification. $\gamma$-H2AX signal for each cell was quantified using Fiji software. For analysis of tandem mass spec data, details can be found in the method details (under protein/peptide identification).

### References

1.     Balzarini, M., Tong, J., Gui, W., Jayalath, I.M., Schell, B.B., and Kodadek, T. (2024). Recruitment to the Proteasome Is Necessary but Not Sufficient for Chemically Induced, Ubiquitin-Independent Degradation of Native Proteins. ACS Chem Biol *19*, 2323-2335. 10.1021/acschembio.4c00422.
2.     Qian, W.J., Liu, T., Monroe, M.M., Strittmatter, E.F., Jacobs, J.M., Kangas, L.J., Petritis, K., Camp, D.G., and Smith, R.D. (2005). Probability-Based Evaluation of Peptide and Protein Identifications from Tandem Mass Spectrometry and SEQUEST Analysis: The Human Proteome. Journal of Proteome Research *4*, 53-62. 10.1021/pr0498638.
3.     Keller, A., Nesvizhskii, A.I., Kolker, E., and Aebersold, R. (2002). Empirical Statistical Model To Estimate the Accuracy of Peptide Identifications Made byMS/MS and Database Search. Analytical Chemistry *74*. https://doi.org/10.1021/ac025747h.

**Table S1: List of oligonucleotides used in this study,**

**Primers (IDT)**

**Vector Amplification**
pcDNA5-FRT Vec Fwd
pcDNA5-FRT Vec Rev
pcDNA5-FRT-FKBP*-XX-FLAG Vec Fwd
pcDNA5-FRT-FKBP*-XX-FLAG Vec Rev

**Mutagenesis**
USP4(C311S) Fwd
USP4(C311S) Rev
UCHL1(C90A) Fwd
UCHL1(C90A) Rev
USP15(C298A) Fwd
USP15(C298A) Rev

**siRNAs (Horizon Discovery)**
Custom USP11 Duplex: Custom siRNA, Standard
0.025 µmol ON-TARGET (UTR1 siRNA)
siGENOME Non-Targeting siRNA
#1, 5 nmol (NT siRNA)

**related to STAR Methods**

**Sequence**

GGATCCGACTACAAGGACGATGA
GGTACCAAGCTTAAGTTTA
GATTATAAAGACGATGACGACAA
GGATCCGGACCCTGA

TAACACAAGCTTTATGAATAGTGCGCTCCAGTG
ATAAAGCTTGTGTTACCGAGGTTGCCC
GAATAGCGCCGGCACCATTGGGCTTATCC
GTGCCGGCGCTATTCCCAATCGTCTGC
GAATACAGCATTTATGAACAGTGCAATCCAATGC
ATAAATGCTGTATTCCCAAGGTTGCTCAAAC

| Sequence | Cat# |
|---|---|
| Sense: 5' G.U.G.U.A.U.A.U.A.U.A.A.A.G.C.A.C.C.A.U.U 3'<br>Antisense: 5' U.G.G.U.G.C.U.U.U.A.U.A.U.A.U.A.C.A.C.U.U 3' | CTM-1075346<br><br>D-001210-01-05 |

**Data S1: Proteomics data from pulldown experiment**

**Proteins that co-precipitated with FKBP\*-USP11 but not FKBP\*-UCHL1**
**(USP7 and PRKDC are highlighted)**

**Identified Protein**
Large ribosomal subunit protein uL6 OS=Homo sapiens OX=9606 GN=RPL9P9 PE=1 SV=1
26S proteasome non-ATPase regulatory subunit 2 OS=Homo sapiens OX=9606 GN=PSMD2 PE=1 SV=3
CAD protein OS=Homo sapiens OX=9606 GN=CAD PE=1 SV=3
DNA-dependent protein kinase catalytic subunit OS=Homo sapiens OX=9606 GN=PRKDC PE=1 SV=3
Small ribosomal subunit protein eS8 OS=Homo sapiens OX=9606 GN=RPS8 PE=1 SV=2
Ornithine aminotransferase, mitochondrial OS=Homo sapiens OX=9606 GN=OAT PE=1 SV=1
26S proteasome non-ATPase regulatory subunit 3 OS=Homo sapiens OX=9606 GN=PSMD3 PE=1 SV=2
Large ribosomal subunit protein eL14 OS=Homo sapiens OX=9606 GN=RPL14 PE=1 SV=4
Ribonucleoside-diphosphate reductase subunit M2 OS=Homo sapiens OX=9606 GN=RRM2 PE=1 SV=1
Cell division cycle and apoptosis regulator protein 1 OS=Homo sapiens OX=9606 GN=CCAR1 PE=1 SV=2
rRNA 2'-O-methyltransferase fibrillarin OS=Homo sapiens OX=9606 GN=FBL PE=1 SV=2
Microtubule-associated protein 1B OS=Homo sapiens OX=9606 GN=MAP1B PE=1 SV=2
RNA-binding protein 5 OS=Homo sapiens OX=9606 GN=RBM5 PE=1 SV=2
Large ribosomal subunit protein uL4 OS=Homo sapiens OX=9606 GN=RPL4 PE=1 SV=5
OTU domain-containing protein 4 OS=Homo sapiens OX=9606 GN=OTUD4 PE=1 SV=4
Ubiquitin carboxyl-terminal hydrolase 7 OS=Homo sapiens OX=9606 GN=USP7 PE=1 SV=2
Splicing factor 3A subunit 3 OS=Homo sapiens OX=9606 GN=SF3A3 PE=1 SV=1
Gem-associated protein 2 OS=Homo sapiens OX=9606 GN=GEMIN2 PE=1 SV=1
Tripeptidyl-peptidase 2 OS=Homo sapiens OX=9606 GN=TPP2 PE=1 SV=4
Endoplasmin OS=Homo sapiens OX=9606 GN=HSP90B1 PE=1 SV=1
Large ribosomal subunit protein uL11 OS=Homo sapiens OX=9606 GN=RPL12 PE=1 SV=1
Large ribosomal subunit protein uL13 OS=Homo sapiens OX=9606 GN=RPL13A PE=1 SV=2

| Accession Number | Alternate ID |
|---|---|
| P32969 | RPL9P9 |
| Q13200 | PSMD2 |
| P27708 | CAD |
| P78527 | PRKDC |
| P62241 | RPS8 |
| P04181 | OAT |
| O43242 | PSMD3 |
| P50914 | RPL14 |
| P31350 | RRM2 |
| Q8IX12 | CCAR1 |
| P22087 | FBL |
| P46821 | MAP1B |
| P52756 | RBM5 |
| P36578 | RPL4 |
| Q01804 | OTUD4 |
| Q93009 | USP7 |
| Q12874 | SF3A3 |
| O14893 | GEMIN2 |
| P29144 | TPP2 |
| P14625 | HSP90B1 |
| P30050 | RPL12 |
| P40429 | RPL13A |