Targeted Inhibition of Translation Initiation via 2'O-Methylation of Start Codons

Emmely A. Patrasso[1,6], Hannah Serio[1,6], Adam Suh[1], Stephanie Mou[2], Kevin Vasquez[1,3], Siqi Wu[4],

Yang Yi[4,5], Daniel Arango[1,5,*]

[1]Department of Pharmacology. Feinberg School of Medicine. Northwestern University. Chicago,

IL. USA

[2]Driskill Graduate Program in Life Sciences. Feinberg School of Medicine. Northwestern

University. Chicago, IL. USA.

[3]Department of Chemical & Biological Engineering. McCormick School of Engineering.

Northwestern University. Evanston, IL. USA.

[4]Department of Urology. Feinberg School of Medicine. Northwestern University. Chicago, IL. USA.

[5]Robert H. Lurie Comprehensive Cancer Center. Feinberg School of Medicine. Northwestern

University. Chicago, IL. USA.

[6] These authors contributed equally to this work

[*] To whom correspondence may be addressed: daniel.arangotamayo@northwestern.edu

Abstract

While protein synthesis typically initiates at an optimal AUG start codon, the 5' untranslated region

(5'UTR) of mRNAs harbors non-canonical start codons that result in the translation of upstream

Open Reading Frames (uORFs). However, the mechanisms underlying the selection of non-

canonical start codons remain poorly understood. Structural analysis of translation pre-initiation

complexes showed that the 2'OH group of the first nucleotide within start codons is monitored by

18S rRNA, allowing optimal translation initiation. We identified a chemical modification of mRNA,

2'O-methylation (N_m), that represses translation initiation by interfering with the interaction of

mRNA and 18S rRNA, thereby preventing the pre-initiation complex from recognizing start

codons. Significantly, we uncovered the presence of 2'O-methylation in the first nucleotide of non-

canonical initiation sites within 5'UTRs, where it repressed upstream translation initiation. These

findings revealed a potent regulatory function of 2'O-methylation in translation initiation and uORF

expression, providing new insights into the mechanisms of non-canonical start codon selection in

human cells.

Keywords

2'O-methylation, N_m, non-AUG translation, uORF, epitranscriptome

Main

During translation initiation, a 43S ribonucleoprotein particle containing the small ribosome subunit (40S), the initiator tRNA (tRNA$_i^{met}$), and translation initiation factors, binds the 5' end of mRNAs to form a 48S pre-initiation complex.[1] This complex scans through the 5'UTR until it encounters an optimal start codon.[1] The optimality of a start codon is typically determined by the Kozak sequence (RccAUGG, where R is either adenosine or guanosine).[2] Once the start codon is recognized, GTP hydrolysis and rearrangement of the 48S complex enable the large 60S subunit to bind and form the 80S initiator ribosome.[1] Subsequently, the 80S ribosome moves into elongation and continues decoding through an Open Reading Frame (ORF) until a stop codon is reached.[1]

While the Kozak sequence defines the optimal translation initiation site, translation can also initiate at suboptimal or cryptic start sites characterized by the presence of AUGs in non-Kozak sequences or near-cognate start codons, which differ from AUG by one nucleotide (CUG, GUG, UUG, AAG, ACG, AGG, AUA, AUC, and AUU).[3] While these cryptic start codons are usually bypassed by the 48S pre-initiation complex through a process known as leaky scanning,[4] translation initiation at upstream sites results in the expression of upstream Open Reading Frames (uORFs).[3] Initially thought to be mere cis-regulatory sequences that inhibit the expression of downstream canonical ORFs, recent studies demonstrated that uORFs also yield stable micropeptides with diverse biological functions, many of which are involved in neurological or metabolic diseases and cancers.[5-7] However, the mechanisms underlying the selection of non-canonical start codons and uORF expression under normal homeostasis and in disease remain unclear.

One poorly characterized layer of translation regulation is the epitranscriptome, defined as the set of more than 150 chemical modifications of ribonucleotides that alter RNA function. Several RNA modifications have been reported to regulate translation initiation. For instance, the hypermodified nucleotide 6-threonyl-carbamoyl-adenosine (t^{6}A) in position 37 of tRNA$_i^{met}$ directly interacts with Kozak sequences, where it stabilizes the translation pre-initiation complex.[8,9] Moreover, the mRNA cap is characterized by a non-templated 7-methylguanosine (m^{7}G), followed by one or two templated nucleosides with methylation in the 2'OH of the ribose, a modification called 2'O-methylation (also referred to as A$_m$, C$_m$, G$_m$, U$_m$, or simply N$_m$).[10] When the first nucleotide is an adenosine, it is usually further modified into N^6,2'O-dimethyladenosine (m^{6}A$_m$).[11] The m^{7}G cap is essential for translation, whereas the subsequent N$_m$ modifications in the first or second

nucleotide regulate mRNA stability.[10-12] Beyond the cap, several mRNA modifications are found

internally in the 5'UTRs, including m^6A,[13-15] N4-acetylcytidine (ac^4C),[8,16] and 2′O-methylation.[17-20]

While m^6A regulates cap-independent translation by recruiting 'reader' proteins,[10-12] and ac^4C

weakens the interaction of $tRNA_i^{met}$ with the Kozak sequence,[8] the function of 2′O-methylation

within 5'UTRs is unknown.

Using synthetic mRNAs with 2′O-methylation in every position of the Kozak sequence and re-

analysis of cryo-EM structures of translation pre-initiation complexes, we found that 2'O-

methylation in the first nucleotide of the start codon completely abolishes protein synthesis.

Mechanistically, 2'O-methylation interferes with the interactions between mRNA and 18S rRNA,

thereby preventing the pre-initiation complex from recognizing start codons. Notably, we

uncovered the presence of 2′O-methylation sites in upstream non-canonical start codons within

human 5'UTRs, where N_m inhibits upstream translation initiation in a position-dependent manner.

Collectively, this study revealed a potent regulatory role of 2'O-methylation in translation initiation,

providing novel insights into the mechanisms of non-canonical start codon selection in human

transcriptomes.

Results

2'OH hydrogen bonds within translation initiation complexes

To understand what defines an optimal start codon, several groups solved cryo-EM structures of

mammalian pre-initiation and initiation complexes.[8,9,21,22] These efforts provided a snapshot of the

crucial intermolecular interactions surrounding Kozak sequences (Fig. 1a). Besides the

recognition of start codons by the anticodon loop in $tRNA_i^{met}$, the interactions of positions A/G[-3]

and G[+4] with translation initiation factors are required for translation initiation (Fig. 1a). In

revisiting several mammalian translation initiation structures[8,9,21], we observed that 2'OH groups

of mRNA nucleotides within the Kozak sequence appeared to be monitored by other components

of the pre-initiation complex. For instance, t^6A in position 37 of $tRNA_i^{met}$ forms a hydrogen (H)-

bond with the 2'OH group of the nucleotide preceding the start codon, C[-1] (Fig. 1b and Extended

Data Fig. 1a). The 2'OH group of the first nucleotide, A[+1], in the start codon interacts with U1830

of 18S rRNA, either with the phosphate (PO_4^{3-}) in the backbone (Fig. 1c and Extended Data Figs.

1b-f), or the keto (C=O) group in position C2, especially in the late stages of the pre-initiation

complex (Fig. 1d and Extended Data Figs. 1e-f). Two additional H-bonds were observed between

the 2'OH group of G[+3] with lysine seven (K7) of the translation initiation factor 1A (eIF1A, Fig.

1e) and between the 2'OH group of nucleotide A[-3] in the mRNA with G961 of 18S rRNA (Fig.

1f). Together, these observations suggest that 2'OH groups in mRNA are monitored by 18S rRNA,

tRNA$_i^{met}$, and eIF1A, potentially playing a crucial role in start codon recognition.

2'O-methylation of start codons inhibits translation

To probe the significance of the H-bonds between 2'OH groups in mRNA with tRNA$_i^{met}$, 18S rRNA,

and eIF1A, we implemented a three-way splint ligation approach (Fig. 2a).[23] Briefly, a 5'UTR

lacking a start codon was transcribed and capped *in vitro* (Extended Data Figs. 2a,b). In parallel,

the ORF of NanoLuciferase (NanoLuc)—also lacking a start codon—was transcribed and

polyadenylated *in vitro* (Extended Data Fig. 2c). These pieces were ligated to a chemically

synthesized RNA fragment containing the Kozak sequence using a DNA splint that hybridizes to

the three independent RNA pieces (Fig. 2a and Extended Data Fig. 2d). The ligated NanoLuc

mRNAs with a Kozak sequence, the ligated mRNAs that lack a start codon, and the unligated

RNA pieces were transfected into HEK293T cells. Crucially, the ligated mRNAs containing the

Kozak sequence exhibited an ~100-fold increase in NanoLuc activity compared to the unligated

RNA pieces or the ligated mRNAs lacking a start codon (Fig. 2b).

To control for possible confounding artifacts, wherein a high proportion of unligated products

affect translation, we transfected increasing concentrations of full-length NanoLuc mRNA with

constant amounts of unligated RNA (Fig. 2c). Reciprocally, we transfected constant amounts of

full-length NanoLuc mRNA with increasing concentrations of unligated RNAs (Extended Data Fig.

2e). In either case, NanoLuc activity was unaffected by the unligated RNAs, exhibiting linear

behavior across a broad range of mRNA concentrations.

Leveraging the ability of 2'O-methylation to disrupt the donor capacity of 2'OH groups, we

systematically methylated the 2'OH groups of every nucleotide within the Kozak sequence using

the three-way splint ligation approach (Fig. 1d and Extended Data Fig. 2f). The ligation efficiency

was verified through gel electrophoresis and RT-qPCR, obtaining similar ligation efficiencies

across multiple replicates (Extended Data Fig. 2g). Equivalent amounts of ligated mRNAs were

transfected into HEK293T cells, and translation efficiency was measured six hours after

transfection through a luminescence assay. Remarkably, methylation of the 2'OH group in the

first position of the AUG codon completely abolished NanoLuc activity (Fig. 2e). Significant

luminescence reduction was also observed when methylation occurred on positions G[+3] and

G[+4] (Fig. 2e). In contrast, no significant effects were observed in positions A[-3], C[-2], C[-1],

and U[+2] (Fig. 2e). The complete inhibition of NanoLuc activity by A_m[+1] remained after
normalization of NanoLuc activity by transfected mRNA levels (Extended Data Fig. 2h).

To guarantee that the inhibitory effect of A_m[+1] on NanoLuc activity is transcript-specific and not
a global perturbation of protein synthesis, we co-transfected unmodified or A_m[+1]-modified
NanoLuc mRNAs with unmodified Firefly luciferase mRNAs. We observed that A_m[+1] abolished
NanoLuc activity without significantly altering Firefly luciferase activity (Fig. 2f). In addition,
evaluation of mRNA levels at different time intervals post-transfection showed no significant
differences in mRNA stability between unmodified and 2'O-methylated transcripts (Fig. 2g),
indicating that the effect of 2'O-methylation on protein output occurs via the inhibition of
translation.

Next, we evaluated the reproducibility of these results across different mRNAs and cell types.
First, we transfected the ligated NanoLuc mRNAs into HeLa cells. Like HEK293T cells, A_m[+1]
completely abolished NanoLuc activity in HeLa cells (Extended Data Fig. 2i). G_m[+3] and G_m[+4]
significantly reduced NanoLuc activity, albeit to a lesser extent than A_m[+1] (Extended Data Fig.
2i). Second, we generated green fluorescent protein (GFP) mRNA reporters, each one containing
2'O-methylation at a different position in the Kozak sequence. Ligation efficiency was generally
higher than 50% for GFP constructs (Extended Data Fig. 2j), which include a much larger 5'UTR
than the NanoLuc mRNA reporters (Extended Data Fig. 2k and Supplementary Table 1).
Following transfection of the GFP mRNA reporters into HeLa cells, protein expression was
determined by flow cytometry. Consistently, the GFP fluorescence results mirror those of
NanoLuc activity: A_m[+1] completely abolished GFP expression; G_m[+3] and G_m[+4] reduced
protein expression, whereas A_m[-3], C_m [-2], C_m [-1], and U_m [+2] had no effect on GFP expression
(Figs. 2h,i).

Collectively, these findings demonstrate that 2'O-methylation of Kozak sequences inhibits
translation in a position-specific manner. 2'O-methylation of position A[+1] has the most potent
inhibitory effect, suggesting that H-bonds between the 2'OH group of nucleotide A[+1] and U1830
in 18S rRNA are required for translation initiation.

2'O-methylation inhibits translation from non-AUG start codons
Over 99% of human mRNAs contain alternative AUGs or near-cognate codons in their 5'UTRs
(Fig. 3a and Extended Data Fig. 3a-c). We previously performed Ribo-seq in the presence of

harringtonine.[8] This translation inhibitor allows the mapping of translation initiation sites

transcriptome-wide (Extended Data Fig. 3d).[8] These experiments identified alternative translation

initiation sites in ~70% of cellular mRNAs (Extended Data Fig. 3e). Distribution of upstream start

codons shows a higher prevalence of CUG, AUG, and GUG, with evidence for upstream initiation

in all near-cognate start codons (Fig. 3b). Building on these observations, we sought to determine

whether 2'O-methylation affects translation initiation in the most prevalent near-cognate start

codons: CUG and GUG (Fig. 3c).

Like canonical AUG codons, CUG, GUG, and their methylated versions (C_mUG and G_mUG)

exhibited comparable ligation efficiencies (Extended Data Fig. 3f). Notably, the unmodified CUG

and GUG near-cognate codons induced a ~50-fold reduction in NanoLuc activity compared to the

unmodified AUG codon (Fig. 3d). Yet, 2'O-methylation at position N[+1] reduced the remaining

activity (Fig. 3d). To better assess the effect of 2'O-methylation in near-cognate initiation, we

increased the time of luminescence acquisition to enhance the signal-to-noise ratio. As expected,

2'O-methylation significantly reduced translation from near-cognate start codons (Fig. 3e and

Extended Data Fig. 3g). These findings confirm that AUG is the optimal start codon while near-

cognate codons such as CUG and GUG can also start translation, albeit inefficiently (Fig. 3f).

Significantly, 2'O-methylation at position N[+1] abolishes translation independently on whether

the start codon is a canonical AUG or a near-cognate start codon (Fig. 3f).

2'O-methylation is present in near-cognate codons within 5'UTRs

The finding that N_m[+1] in cognate and near-cognate start codons represses protein synthesis

raised the possibility that 2'O-methylation may be a regulatory mechanism of translation initiation

in cellular mRNAs. Using chemical biology methods and Nanopore direct RNA sequencing (DRS)

approaches, previous studies identified N_m as a prevalent mRNA modification.[17-20] While the

highest proportion of internal N_m sites was found in coding sequences (CDS) and 3'UTRs, 2'O-

methylation sites were also identified in 5'UTRs (Extended Data Fig. 4a). Notably, the Nanopore-

DRS approach yielded the highest number of N_m sites, including sites mapped to 5'UTRs

(Extended data Fig. 4a).[17]

To determine whether N_m is present in canonical start codons, we extracted the N_m genomic

coordinates from four different studies[17-20] and intersected them with the genomic coordinates of

canonical start codons. To increase confidence in the candidate N_m sites, we analyzed sites with

an estimated modification ratio of more than 10% per position. We found a near complete absence

of N_m in A[+1] (Fig. 4a). For instance, in HeLa cells, a cell line used in all N_m mapping studies,

only one site in the A_m[+1] position of canonical start codons was obtained (Extended Data Fig.

4b), despite thousands of N_m sites reported in coding sequences (Extended Data Fig. 4a). A total

of six A_m[+1] sites in canonical start codons were obtained in all studies (Fig. 4a), with two sites

being reproduced across different cell lines (Extended Data Table 1). These findings suggest a

negative selection constraint for 2'O-methylation in canonical start codons.

To determine whether N_m is present in upstream AUGs or near-cognate codons, we extracted the

N_m sites that mapped to 5'UTRs of any annotated Ensembl transcript and examined whether

these sites were located in position N[+1] of an upstream AUG or an upstream near-cognate

codon (Extended Data Table 2). This analysis identified >600 N_m[+1] sites across all cell lines

(Fig. 4b and Extended Data Fig. 4c). While CUG had the highest prevalence of 2'O-methylation

in position N[+1] among all near-cognates, upstream AUGs are the least prevalent codons with

only nine A_m[+1] sites identified (Fig. 4b). Notably, the number of C_mUG, C_mUG, A_mUC, and A_mCG

sites were significantly enriched compared to the distribution of these codons in the 5'UTRs of

protein-coding genes (Fig. 4c), whereas AUGs were significantly de-enriched (Fig. 4c), reinforcing

the notion that A_mUG is under a strong negative selection constraint.

Next, we biochemically validated the presence of N_m[+1] in endogenous 5'UTRs. We employed

Reverse Transcription at Low dNTPs (RTL-P),[24] a sensitive method that induces reverse

transcriptase stops (RT-STOP) at low concentrations of dNTPs (Fig. 4d), resulting in a reduction

in amplicon density after PCR amplification of the RT product. While there are more than eleven

human 2'O-methyltransferase enzymes (Extended Data Fig. 5a),[25] Fibrillarin (FBL) is consistently

identified as the predominant enzyme responsible for introducing N_m at internal sites in mRNA.[17-

20,26] Thus, we performed RTL-P in total RNA isolated from wildtype HeLa cells transfected with

two different siRNAs against Fibrillarin (FBL) or a scramble siRNA control (Fig. 4e). Candidate

sites were selected based on the highest predicted modification ratio from the Nanopore-DRS

approach in HeLa cells (Extended Data Table 3). Ten of the twelve candidate sites tested exhibited

a substantial loss in amplicon density under limited dNTP conditions (Figs. 4f-h and Extended

Data Figs. 5b-h). Importantly, the RTL-P signature was lost upon knockdown of FBL (Figs. 4f-h

and Extended Data Figs. 5b-h), demonstrating the presence of FBL-dependent 2'O-methylation

sites in human 5'UTRs. Notably, these findings suggest that 2'O-methylation is preferentially

located in upstream near-cognate codons with significant enrichment and higher prevalence at

CUG and GUG codons.

2'O-methylation inhibits upstream translation initiation

The presence of N_m in upstream AUGs and near-cognate codons raises the possibility that human

cells use 2'O-methylation to inhibit upstream translation initiation. To evaluate this possibility, we

re-analyzed a previously published HR-Ribo-seq[8] in HeLa cells to compare the density of initiator

ribosomes over start codons. As expected, ribosome density around canonical AUGs is higher

than upstream AUGs or near cognate codons (Fig. 5a). To determine the effect of N_m in upstream

translation initiation, we computed the ribosome density at upstream AUGs or near-cognate

codons relative to the ribosome density at the canonical AUG within the same transcripts. Sites

with an RPKM value ≥ 1 were binned based on the methylation status (unmethylated versus

N_m[+1]) and their codon identity. Remarkably, A_mUG codons exhibited a significant reduction in

ribosome density compared to non-methylated AUGs (Fig. 5b and Extended Data Table 4) despite

the low number of A_mUG sites. We next focused on the near-cognate codons CUG, GUG, and

AUC, as those codons showed a significant N_m enrichment in human 5'UTRs (Fig. 4c). The

relative ribosome density at methylated C_mUG and G_mUG was significantly decreased compared

to the unmethylated codons, reaching similar levels found in methylated A_mUG (Fig. 5b and

Extended Data Table 4). Although methylated AUC showed a similar tendency, the differences

between A_mUC and the unmethylated codons were not significant (Fig. 5b).

Next, we compared the ribosome density over a window of 29 nucleotides, the ribosome-

protected mRNA length, on 2'O-methylated codons at position N_m[+1] compared to 2'O-

methylation in positions N_m[-3], N_m[-2], N_m[-1], and N_m[+2]. We reasoned that this comparison

would recapitulate the mRNA reporter studies, which established that A_m[+1] completely abolished

translation, whereas positions N_m[-3], N_m[-2], N_m[-1], and N_m[+2] had no effect in translation

initiation (Figs. 2e,i). Consistent with the mRNA reporters, A_mUG, C_mUG, and G_mUG exhibited

reduced ribosome density compared to methylated codons in positions N_m[-3], N_m[-2], N_m[-1], and

N_m[+2] (Figs. 5c-e and Extended Data Figs. 6a-c).

Though we observed only a handful of 2'O-methylation sites in canonical AUG start codons, we

evaluated whether these methylation events inhibited the translation of the canonical ORF. For

this purpose, we assessed differences in translation efficiency (TE) obtained from traditional Ribo-

seq in HeLa cells.[27] While TE is decreased in A_mUG compared to unmethylated canonical AUG

codons, the differences were not statistically significant, probably due to a lack of statistical power

in the A_mUG group (n=5, Extended Data Fig. 6d).

Overall, these findings demonstrate that 2'O-methylation in the upstream start codons AUG, CUG,

and GUG inhibits upstream translation initiation in cellular mRNAs. These upstream codons with

$N_m[+1]$ are located in genes predominantly involved in cellular components, biological processes,

and molecular functions related to ribosome biogenesis and mRNA translation (Extended Data

Fig. 7).

2'O-methylation inhibits start codon recognition

Once the start codon is recognized, a 48S pre-initiation complex hydrolyzes GTP, resulting in the

binding of the large 60S subunit to form the 80S initiator ribosome (Fig. 6a). The inhibitory effect

of 2'O-methylation in position $A_m[+1]$ suggests that the H-bond between the 2'OH group of

nucleotide A[+1] and U1830 in 18S rRNA is required for translation initiation. There are several

possible explanations for this phenomenon. First, 2'O-methylation in position A[+1] stalls the 48S

pre-initiation before or after recognizing the start codon. Second, $A_m[+1]$ precludes the 80S

initiation complex from proceeding into elongation. Third, 2'O-methylation in position A[+1] makes

the start codon non-optimal or unrecognizable by the pre-initiation complex, thus inducing leaky

scanning. To test these possibilities, we performed *in vitro* translation with reticulocyte lysates

using a Cy5-labeled mRNA reporter in the presence of non-hydrolyzable GTP (GMPPNP) or

harringtonine (HR). Pre-treatment with GMPPNP stalls the 48S pre-initiation complex after codon

recognition (Fig. 6b). Relatedly, HR stalls the 80S initiation complex before proceeding into

elongation (Fig. 6c).

After *in vitro* translation, reactions were loaded on a 5-30% sucrose density gradient followed by

ultracentrifugation, fractionation, and fluorescence (Cy5) recording. As expected, treatment with

GMPPNP increases the abundance of the 48S peak compared to vehicle control (Figs. 6d,e, blue

lines), whereas harringtonine treatment increases the 80S peak in the Cy5-labeled unmodified

mRNA reporter (Fig. 6d,f, blue line). In contrast, $A_m[+1]$ decreases the formation of both the 48S

peak and 80S peaks in the presence of GMPPNP or HR, respectively, as well as in the untreated

sample (Figs. 6d-f, orange arrows).

Collectively, these findings indicate that codon recognition by the 48S pre-initiation complex and

subsequent formation of an 80S initiator ribosome did not occur when the start codon was 2'O-

methylated in position N[+1]. Thus, $N_m[+1]$ leads to leaky scanning, and the 48S subunit can

continue scanning until it encounters another downstream start codon or, alternatively,

dissociates from the mRNA (Fig. 6g).

Discussion

We uncovered a potent regulatory function of 2'O-methylation in translation initiation and uORF

expression, providing new insights into the mechanisms of start codon selection in human cells.

Three significant findings support this mechanism. First, H-bonds between the 2'OH group of

nucleotide A[+1] in mRNA with U1830 in 18S rRNA are required for translation initiation. Second,

A_mUG, C_mUG, and G_mUG inhibit translation initiation at canonical and upstream start codons.

Third, 2'O-methylation of position N[+1] of the start codon is preferentially deposited at upstream

near-cognate codons (C_mUG and G_mUG), exhibiting negative selection for canonical and

upstream AUGs in human transcriptomes.

The movement of the 48S subunit along the 5′ UTR is characterized by a conformational

intermediate between the closed state, in which the tRNA$_i^{met}$ is fully inserted into the P site trying

to make an interaction with the mRNA codon, and the open state, in which the anticodon in

tRNA$_i^{met}$ is distant from the mRNA.[22,28] This conformational intermediate enables the initiator

tRNA$_i^{met}$ to translocate along the mRNA while transiently sampling codons for base-pair

complementarity. The purpose of this transient sampling is to recognize an optimal start codon

while rejecting non-cognate codons and, in most cases, near-cognate codons or AUGs under

poor sequence contexts.[29-32] Once the start codon is entirely recognized, a fully closed and

arrested 48S subunit signals a conformational change that allows the 60S subunit to join.[9,21,28,32,33]

Since GMPPNP arrests the 48S after the start codon is recognized but before the 60S subunit

joins, the lack of GMPPNP-induced 48S arrest in mRNAs carrying A_mUG suggests that 2'O-

methylation inhibits start codon recognition (Fig. 6f), potentially preventing tRNA$_i^{met}$ from fully

inserting into the P site and thereby precluding the 48S subunits from forming a fully closed

conformation. Consequently, 2'O-methylation makes an otherwise optimal translation initiation

site look like a non-cognate codon, and the 48S subunit continues scanning until it encounters a

downstream start codon or, alternatively, dissociates from the mRNA. In either case, the $A_m[+1]$

modification does not impair mRNA stability or globally affect protein synthesis (Fig. 2f,g).

The intermolecular interactions of mRNA with tRNA$_i^{met}$, 18S rRNA, and translation initiation

factors, as well as the orientation and pairing of the codon:anticodon, are essential for start codon

selection (Fig. 1a).[9,22,28,32] Chemically, the 2'-OH group in RNA can act as both a donor and

acceptor in H-bonds. Methylation replaces the hydrogen atom of the 2'-OH with a methyl group (-

CH3), effectively eliminating the donor capability of the 2'-OH group. Since the 2'OH group of the

A[+1] position acts as a donor in the H-bond with U1830, either with the phosphate (PO_4^{3-})

backbone or the keto (C=O) group in position C2, our findings suggest that this interaction is

required for proper start codon recognition. Notably, while the 2'OH of position A[+1] is near PO_4^{3-}

in U1830 along all steps of codon recognition, rearrangement, and 60S joining (Figs. 1c,d and

Extended Data Figs. 1b-f), the 2'OH of position A[+1] is only near the keto (C=O) group in the late

stages of the pre-initiation complex (Extended Data Figs. 1e-f). We speculate that 18S rRNA

monitors for proper pairing and orientation of the codon:anticodon interaction. Thus, blocking the

A[+1]^{mRNA}:U1830^{18S} interaction precludes the ribosome from properly proofreading start codon

recognition. Alternatively, since 2'O-methylation is known for altering RNA structure,[34,35] a

structural shift induced by 2'O-methylation may alter the orientation of A[+1] in the mRNA, thereby

preventing it from properly pairing with the anticodon base. Future structural analyses would be

necessary to determine the exact mechanism by which A_m[+1] inhibits start codon recognition.

Notably, the effect of 2'O-methylation in translation initiation is site-specific. 2'O-methylation of

position N[-3], N[-2], N[-1], and N[+2] had no effect in protein output (Figs. 2e,i). An unexpected

finding was the lack of effect in position N[-1], as previous studies reported a significant role of

this C[-1]^{mRNA}:t^{6}A^{tRNA} interaction in stabilizing the initiation complex.[8,9] For instance, we previously

reported that replacing cytosine with ac^{4}C at position N[-1] weakens the C[-1]^{mRNA}:t^{6}A^{tRNA}

interaction, resulting in reduced translation initiation.[8] Additionally, studies in yeast reported that

inhibition of t^{6}A biosynthesis increases upstream non-AUG translation.[36] While our findings using

2'O-methylation could suggest that the C[-1]^{mRNA}:t^{6}A^{tRNA} interaction may not be required for

translation initiation, it is important to note that 2'O-methylation does not alter the acceptor

capacity of the C-O-CH3 group. Thus, the C[-1]^{mRNA}:t^{6}A^{tRNA} hydrogen bonds may be the result of

the carboxyl group of t^{6}A acting as a donor and the C-O-H group of C[-1] acting as an acceptor.

Under these circumstances, the incorporation of a methyl group at position 2'OH would not disrupt

the C[-1]^{mRNA}:t^{6}A^{tRNA} hydrogen bond.

2'O-methylation in positions N[+3] and N[+4] significantly reduced protein synthesis, albeit to a

lesser extent than N[+1] (Figs. 2e,i). While positions N[+3] and N[+4] are also involved in crucial

intermolecular interactions within the translation initiation complex (Figs. 1a,e,f), the effect in

protein synthesis may be mediated through impairing elongation rather than translation initiation.

For instance, 2'O-methylation is known to be a potent repressor of translation elongation by

precisely inhibiting the interaction of mRNA with the monitoring bases of rRNA during the

proofreading steps of translation elongation.[26,37]

It is well established that 2′-O-methylation is highly abundant in noncoding RNAs, including rRNA

and tRNA, and occurs in the 5′ cap of all mRNAs in human transcriptomes.[25] The presence of N_m

at internal mRNA sites is more controversial,[38] with some studies suggesting that N_m does not

exist internally in mRNA, while others report thousands of N_m sites in mRNA.[17-20,39,40] 2′O-

methylation confers several chemical properties that can be exploited for the site-specific mapping

of N_m, including resistance to alkaline hydrolysis (RiboMeth-seq),[40] periodate oxidation (N_m-seq),[20]

enzymatic cleavage (NJU-seq),[19] specific sensitivity to engineered reverse transcriptase (N_m-mut-

seq),[18] steric hindrance during reverse transcription (2OMe-seq),[39] and unique electrical

signatures in Nanopore-DRS.[17] Despite the variety of approaches that can detect N_m sites in

mRNA, there is little to no reproducibility between the studies.[38] The large inconsistencies across

multiple studies could be due to technical factors, as some deep sequencing technologies are

more sensitive than others but are also prone to artifacts. For example, in the initially published

N_m-seq method,[20] a mispriming artifact led to many false positives—although this artifact was later

corrected in a corrigendum. Alternatively, inconsistencies across studies may have a biological

basis if N_m deposition in mRNA is context-dependent. Nevertheless, low throughput validations of

N_m sites,[26] including ours (Figs. 4d-h and Extended Data Figs. b-h), have undoubtedly identified

the presence of N_m at internal mRNA sites.

Our findings that 2'O-methylation is significantly enriched at near-cognate codons, especially

CUG, suggest that 2'O-methylation is leveraged by cells to control non-AUG translation. Such a

mechanism could be beneficial under stress conditions when canonical cap-dependent

translation is repressed, allowing non-AUG translation to serve as a stress response mechanism.

In this manner, cells could block unwanted non-canonical initiation sites through 2'O-methylation

while leaving desired CUG codons unmethylated to facilitate their initiation. If so, deposition of

2'O-methylation in near-cognate codons might be induced under stress conditions. Future

research outside the scope of this study is necessary to test these hypotheses.

Our findings also have a significant impact on RNA therapeutics. Aberrant translation initiation

has been reported in numerous diseases, with hundreds of pathogenic genetic variants in 5'UTRs

linked to the introduction or destruction of uORFs.[5,41-44] These pathological conditions would

benefit from strategies that inhibit the aberrant initiation site while enhancing the expression of

the canonical ORF. Since 2'O-methylation effect on translation initiation is transcript and position-

specific, our findings open the door to investigate approaches capable of guiding 2'O-methylation

to specific positions within aberrant mRNAs,[45-47] for the targeted inhibition of pathogenic

translation initiation in human disease.

In sum, we identified a chemical modification of mRNA, 2'O-methylation, that unambiguously

inhibits start codon recognition in a position-specific manner. Remarkably, we found that human

cells use 2'O-methylation to inhibit non-canonical translation. These findings uncovered a potent

regulatory mechanism of translation initiation with potential applications in RNA therapeutics

aimed at targeting aberrant translation initiation in human diseases.

Author contributions

Conceptualization, D.A; Methodology: D.A.; Investigation, E.A.P., S.M., A.S., K.V, S.W. and Y.Y;

Formal Analysis, D.A., H.S., and Y.Y.; Data Curation, D.A. and H.S.; Writing – Original Draft, D.A.;

Writing – Review & Editing, D.A. and H.S.; Visualization, D.A. and H.S.; Supervision, D.A., Project

Administration, D.A.; Funding Acquisition, D.A.

Acknowledgments

This study was supported with funds from the U.S. Department of Defense grant HT9425-24-1-

0352 and the Kinship Foundation (SSP-2023-102) to D.A. Y.Y. was supported by the U.S.

Department of Defense grant HT9425-23-1-0661 and NIH P50CA180995 SPORE in Prostate

Cancer Career Enhancement Award. K.V. was supported by the National Institutes of Health

Training Grant (T32GM008449) through Northwestern University's Biotechnology Training

Program.

Competing Interest Statement

Nothing to disclose

Materials and Methods

Cell culture

HEK293T cells were cultured in Dulbeccos' Modified Eagle Medium (DMEM, Gibco, Cat. #:

11965092) supplemented with 10% Fetal bovine serum (Corning, Cat. #: 30-015-CV) and 1%

Penicillin/Streptomycin (Dot scientific, Cat. #: DSP93560-50). HeLa cells were cultured in DMEM

(Gibco, Cat. #: 11965092) supplemented with 10% bovine calf serum (Cytiva, Cat. #:

SH30073.04) and 1% Penicillin/Streptomycin.

Synthesis and capping of 5' RNA fragments

To produce the NanoLuc 5'UTR, an antisense DNA template containing a T7 promoter sequence

was purchased from Integrated DNA Technologies (IDT) and annealed to a sense T7 primer

(Supplementary Table 1). To produce the GFP 5'UTR, we generated the DNA template by PCR

using primers that annealed to a plasmid containing the coding sequence of micro red

fluorescence protein, making sure to exclude the canonical start codon from the amplicon product.

The forward primer contained the T7 sequence as an overhang. PCR was performed using the

Q5 Hot Start High-Fidelity 2X Master Mix (New England Biolabs, Cat. #: M0494). Primers and

resulting DNA sequences are described in Supplementary Table 1.

DNA duplexes were *in vitro* transcribed using the HiScribe® T7 High Yield RNA Synthesis Kit

(New England Biolabs, Cat. #: E2040), following the manufacturer's suggestions. RNA was

purified using 1.8X SPRI beads (CleanNGS Cat. #: CNGS-0001). Purified RNA was capped *in*

vitro using the Vaccinia Capping System (New England Biolabs, Cat. #: M2080) and re-purified

using 1.8X SPRI beads. The integrity of RNA fragments was verified using polyacrylamide gels

prepared in a tris-borate-urea buffer (2 mL of 40% PAA [19:1], 5 g of Urea, 1mL of 10X TBE,

complete to 10 mL ddH2O). Electrophoresis was run in a 0.5X TBE running buffer and stained

with a 1X SYBR gold solution (ThermoFisher, Cat. #: S11494). Gels were visualized in a Sapphire

Biomolecular Imager (Azure Biosystems).

Dot blots

Dot blots were performed to verify the capping of *in vitro* transcribed RNAs. Briefly, 500 ng RNA

was denatured at 65°C for 5 min and loaded onto Hybond-N+ membranes. Membranes were

crosslinked with 120 mJ^2/cm^2 in the UV Crosslinker. Membranes were first stained in a 0.2%

methylene blue solution (0.2% methylene blue in 0.4M sodium acetate and 0.4M acetic acid),

then washed three times in DI water, and imaged in a Sapphire Biomolecular Imager.

Membranes were then blocked with 1% bovine serum albumin (BSA) in PBS containing 0.05%

Tween-20 (PBST) for 15 min at room temperature and probed overnight with mouse anti-m^{7}G

antibodies (1:1000, MBL International Corporation, Cat. #: RN016M) in 1% BSA at 4°C.

Membranes were next washed three times with 0.1% PBST, incubated with HRP-conjugated

secondary anti-mouse IgG (1:10000 dilution, Cell Signaling Technologies, Cat. # 7076S) in 1%

BSA at room temperature for 1 hr, washed three times with 0.1% PBST and developed with

enhanced chemiluminescent (ECL) western blotting substrate (Promega, Cat #: W1001) on a

Sapphire Biomolecular Imager.

Synthesis and polyadenylation of 3' RNA fragments

To produce the ORF and 3'UTR of NanoLuc, we generated the DNA template by PCR using

primers that annealed to a plasmid containing the coding sequence of NanoLuc, making sure to

exclude the canonical start codon from the amplicon product. The forward primer contained the

T7 sequence as an overhang. To produce the ORF and 3'UTR of GFP, we generated the DNA

template by PCR using primers that annealed to a plasmid containing the coding sequence of

GFP, making sure to exclude the canonical start codon from the amplicon product. The forward

primer contained the T7 sequence as an overhang. PCR was performed using the Q5 Hot Start

High-Fidelity 2X Master Mix (New England Biolabs, Cat. #: M0494). Primers and resulting DNA

sequences are described in Supplementary Table 1.

DNA duplexes were *in vitro* transcribed using the HiScribe® T7 High Yield RNA Synthesis Kit

(New England Biolabs, Cat. #: E2040), following the manufacturer's suggestions. RNA was

purified using 1X SPRI beads (CleanNGS Cat. #: CNGS-0001). RNA fragments (80 μg) were

dephosphorylated with 100 U of calf intestinal alkaline phosphatase (CIAP, Promega, Cat. #:

M2825), followed by mono-phosphorylation of the 5' end using 15 U of T4 Polynucleotide Kinase

(PNK, New England Biolabs, Cat. #: M0201L) in the presence of ATP. Phosphorylated RNA

products were purified using 1X SPRI beads. RNA products (10 μg) were polyadenylated using 5

U of *E. coli* Poly(A) Polymerase (New England Biolabs, Cat. #: M0276). The integrity of

polyadenylated RNA fragments was verified using TBU-PAGE. Electrophoresis was run in a 0.5X

TBE running buffer and stained with a 1X SYBR gold solution.

Three-way splint ligations

507
Phosphorylated RNA oligonucleotides (12 nt) containing the Kozak sequence with an AUG, CUG,
or GUG start codons were purchased from IDT. Likewise, the 2'O-methylated versions were
purchased from IDT (Supplementary Table 1). These RNA oligos were combined with 5' and 3' *in*
vitro transcribed RNA fragments at equimolar concentrations (15 pmol). An antisense DNA
oligonucleotide (15 pmol) was added to splint the different RNA pieces. Reactions were denatured
at 65 °C for 2 minutes and ramped down to room temperature for annealing.

Ligations were performed using 10 U of the T4 RNA ligase 2 (dsRNA ligase, New England
Biolabs, Cat. #: M0239) for 4 hr at 37 °C according to the manufacturer's suggestions. Reactions
were stopped with 2 U of Turbo DNase I (ThermoFisher Scientific, Cat. #: AM2238) for 30 min at
37°C and purified using the RNA Clean & Concentrator-5 kit (Zymo Research, Cat. #: R1014). To
verify the integrity of splint ligated NanoLuc RNA, 500 ng of ligated RNAs were separated by 4%
TBU-PAGE and stained with 1X SYBR gold (Invitrogen, Cat. #: S11494). To verify the integrity of
splint ligated GFP RNA, 500 ng of ligated RNAs were separated by 1% denaturing agarose gels
and stained with 1X SYBR gold (Invitrogen, Cat. #: S11494). Gels were visualized in a Sapphire
Biomolecular Imager (Azure Biosystems).

NanoLuc and Firefly luciferase assays

525
HEK293T or HeLa cells were seeded at a density of 30,000 cells in 100 μL complete DMEM
medium in 96-well plates. Cells were cultured for 24 hr before transfecting with 300 ng of capped
and polyadenylated splint ligated RNA or unligated RNA control. Transfection reactions were
prepared using the TransIT-mRNA Transfection Kit (Mirus, Cat. #: MIR2225) following the
manufacturer's instructions. Following the addition of the transfection mixtures, cells were
incubated for 6 hrs at 37°C. For luminescence detection, cells were assayed with Nano-Glo®
Luciferase Assay System (Promega, Cat. #: N1110) according to the manufacturer's instructions
for 96-well plates. For the dual transfection experiments, cells were assayed using the Nano-Glo
Dual-Luciferase Reporter Assay System (Promega, Cat. #: N1610) according to the
manufacturer's instructions for 96-well plates. Luminescence was read in a BioTek Synergy LX
Multimode Reader (Agilent Technologies).

Flow cytometry

HEK293T or HeLa cells were seeded at a density of 150,000 cells in 1 mL medium in 12-well

plates. Cells were transfected with 600 ng of capped and polyadenylated splint ligated mRNA or

unligated RNA control. Transfection reactions were prepared using the TransIT-mRNA

Transfection Kit (Mirus) following the manufacturer's instructions for 12-well plates. Following the

addition of the transfection mixtures, cells were incubated for 18 hrs at 37°C. Cells were

trypsinized and washed with PBS three times before they were resuspended in 500 μL PBS and

immediately analyzed via Flow cytometry in an LSR Fortessa 1 Analyzer or a BD FACSymphony

A5-Laser Analyzer. The percentage of GFP-positive cells was obtained using FlowJo/v.10.10.0.

RNA isolation and RT-qPCR

Total RNA was isolated from the transfection reaction at time 0 or transfected cells at different

time points post-transfection using the Trizol method. RT-qPCR was performed to evaluate the

efficiency of ligation before transfection or the stability of ligated mRNA at different time points

post-transfection. Briefly, the extracted RNA was reverse transcribed using SuperScript™ IV

Reverse Transcriptase (ThermoFisher Scientific, Cat. #: 18090010) using gene-specific RT

primers for NanoLuc, GFP, and 18S rRNA (Supplementary Table 1). qPCR was performed using

the 2X PowerUp SYBR Master Mix (ThermoFisher Scientific, Cat. #: A25742) and primers that

amplify the ligated mRNAs, total mRNAs (ligated plus unligated), and 18S rRNA (Supplementary

Table 1). To obtain the efficiency of ligation, we applied the formula $100 * 2^{\wedge}[-(Ct^{ligated}/Ct^{total.mRNA})}$.

To measure mRNA stability, we applied the formula $final.time(2^{\wedge}[-(Ct^{ligated}/Ct^{18S.rRNA})}/$

$initial.time(2^{\wedge}[-(Ct^{ligated}/Ct^{18S.rRNA})}$.

In vitro **translation and sucrose density gradients**

Cy5-labeled synthetic mRNAs were purchased from Genescript (Supplementary Table 1).

Reticulocyte lysates (37.5 μL) were pre-incubated with vehicle DMSO, 1 mM GMPPNP

(Guanosine 5′-[β,γ-imido]triphosphate trisodium salt hydrate, Millipore Sigma, Cat. #: G0635), or

100 μg/ml Harringtonine (Abcam, Cat. #: ab141941) prior to adding 150 nM of Cy5-labeled

synthetic mRNAs in a total volume of 50 μL. Reactions were incubated for 10 min at 30 °C and

immediately stopped on ice. Reactions were loaded on 5-30% sucrose density gradient (50 mM

Tris-acetate [pH 7.0], 50 mM NH_4Cl, 12 mM $MgCl_2$, 1 mM DTT) poured using a BioComp Gradient

Master. Gradients were spun at 42,000 rpm in a Beckmann Coulter SW41 Ti rotor for 2:26 hr at

4°C. Fractionation and recording of Cy5 fluorescence were performed using a BioComp Density

Gradient Fractionation system.

siRNA transfection

The FBL siRNAs used in this study were purchased from Thermo Fisher (siFBL-1: s4820, siFBL-

2: s4821). siRNAs were transfected into HeLa cells using the Lipofectamine RNAiMAX

Transfection Reagent (ThermoFisher Scientific, Cat. #: 13778150) by following the manufacturer's

manual. Total RNA and cell lysates were collected three days post-transfection. The efficiency of

FBL knockdown was verified via Western blots using specific anti-FBL antibodies (Abcam, Cat. #:

ab5821) and anti-β-actin antibodies (Cell Signaling Technologies, Cat. #: 3700) as a loading

control.

RTL-P assay

To confirm the N_m occurrence along with its alteration upon FBL depletion, the RTL-P assay was

conducted as previously described.[17] First, total RNAs from control and FBL-knockdown HeLa

cells were isolated using RNeasy Plus Mini Kit (Qiagen). Reverse transcription (RT) was then

performed using specific reverse primers targeting mRNA sequences downstream of the 2'O-

methylation site in the presence of either a low (1 µM) or a high (1 mM) concentration of dNTPs.

The obtained cDNA was subjected to two separate amplification reactions using one pair of

primers targeting upstream and downstream of the methylation site and another pair of primers

that amplified a downstream region that would not be interrupted by the 2'O-methylation site

(Supplementary Table 1). The PCR mixture consisted of 10 µL AmfiSure PCR master mix

(GenDEPOT), 2 µL cDNA, and 1 mM PCR primers. The PCR products were then separated using

1% agarose gels and photographed by a Bio-Rad imaging system. DNA band signal intensities

were calculated using the ImageJ software. The methylation ratio of each group was determined

from the densities of the PCR bands obtained using high and low dNTP concentrations. The RT

and PCR primers designed for the RTL-P assay are listed in Supplementary Table 1.

Identification of 2'OH hydrogen bonds in translation pre-initiation complexes

The following structures were downloaded from the Protein Data Bank: 7UCJ (rabbit reticulocyte),

6YAL (rabbit reticulocyte), 8PJ2, 8PJ3, 8PJ4, 8PJ5, and 8PJ6 (reconstituted human 48S).

Structural visualization was performed in Pymol. Briefly, the background was set to white, and all

elements were initially hidden. Specific residues of the mRNA, centered around nucleotides N[-

3], N[-2], N[-1], N[+1], N[+2], N[+3], and N[+3] were selected and visualized as both ribbons (with

increased width) and sticks. Specific residues within tRNA$_i^{met}$, 18S rRNA, eIF1A, and eIF2α that

interact with mRNA were also selected. Ribbon elements were colored blue-white for mRNA,

green for tRNA, yellow-orange for 18S rRNA, and black for eIF1A or eIF2α. Atomic elements were

colored red (oxygen), blue (nitrogen), and orange (phosphorus). Lighting parameters were
adjusted to enhance visualization, and hydrogen bond calculations between mRNA, tRNA$_i^{met}$, 18S
rRNA, eIF1A, and eIF2α were included.
610

Identification of N$_m$ sites in AUG and near-cognate codons

Maps of 2'O-methylation were extracted from four publications using either biochemical or
Nanopore DRS coupled with machine learning approaches. These methods are named N$_m$-seq,[20]
N$_m$-mut-seq,[18] NJU-seq,[19] and Nanopore-DRS[17]. Publicly available BED files from these studies
were retrieved from the GEO and supplementary tables (GSE90164, GSE174518, Supplemental
Tables 4 and 5 from Li et al. 2024,[17] and Supplemental Table 2 from Tang et al. 2024).[19] Notably,
for the N$_m$-seq study,[20] we downloaded the N$_m$ sites reported in the corrigendum (GSE90164),
which eliminated the mispriming artifacts of the original publication. Three datasets (NJU-seq,
Nanopore-DRS, and N$_m$-mut-seq) reported the stoichiometry of N$_m$ sites; therefore, we chose to
analyze N$_m$ sites with a 2'O-methylation ratio above ten percent.
621

622 To identify N$_m$ sites present in canonical start codons, we extracted the genomic information for
623 Ensembl canonical start codons from a human genome annotation file (GRCh38). Then,
624 bedtools/2.29.2 was used to identify N$_m$ sites around canonical start codons. N$_m$ sites were then
625 binned by their location relative to the canonical AUG; N$_m$[-3], N$_m$[-2], N$_m$[-1], N$_m$[+1], N$_m$[+2],
626 N$_m$[+3], and N$_m$[+4]. The number of sites was plotted based on the position, cell line, and study.
627

To identify N$_m$ sites in upstream AUG and near-cognate codons, we selected N$_m$ sites within
protein-coding genes that aligned to 5'UTR in any possible transcript. Then, we extracted the
genomic sequence flanking every N$_m$ site. We filtered for sequences containing AUG and near-
cognate codons. Each N$_m$ site location was derived relative to the AUG or near-cognate codons
(N$_m$[-3], N$_m$[-2], N$_m$[-1], N$_m$[+1], N$_m$[+2], N$_m$[+3], and N$_m$[+4]). Note that N$_m$ sites can be binned into
multiple positions when located near more than one AUG or near-cognate codon. This analysis
was performed with custom python/3.9.16 scripts using biopython/1.85 scripts.
635

Analysis of HR-Ribo-seq

Raw sequencing files from previously published HR-Ribo-seq in HeLa cells were downloaded
from GSE162043.[8] Adapter sequences were removed from sequence files using cutadapt/4.2.
Reads mapping to tRNA and rRNA sequences were removed with bowtie2/2.4.5. The remaining

reads were mapped to the human genome (GRCh38) using hisat2/2.1.0. Alignments were
normalized by reads per kilobase per million mapped reads (RPKM) using deeptools/3.5.1.

To compare ribosome density between the canonical AUG, upstream AUGs, and upstream near-
cognate codons, we obtained a BED file containing the genomic coordinates of canonical start
codons from Ensembl/113 using biomart. A custom python script extracted RPKMs in a 30-
nucleotide window, 13 nucleotides downstream and 16 nucleotides upstream of the +1 position
in the canonical start codon. Similarly, we used a custom python script to generate a BED file
containing the genomic coordinates of AUGs and near-cognate codons within 5'UTRs. RPKMs
were extracted in the 30-nucleotide window surrounding upstream AUGs and near-cognate
codons. RPKMs of two replicates were averaged together. To remove sites containing extremely
low RPKM values, sites with at least one position where the RPKM is less than 1 within the 30-
nucleotide range were removed from the analysis. An R script was used to visualize the
differences in average RPKMs as a line plot over the range of 30 nucleotides around start codons
or methylated positions.

To analyze the relative ribosome density at upstream codons compared to the canonical start
codon in the presence and absence of N_m[+1], we extracted the RPKM at position +1 for upstream
near-cognate codons and canonical AUGs. RPKMs were averaged for two replicates, and an R
script was used to remove N_m sites with an RPKM less than 1. The ratio of the upstream RPKM
to the canonical RPKM in the presence and absence of upstream N_m[+1] was visualized as box
plots using the ggpubr and ggplot2 packages in R. A Wilcoxon test was performed for each
upstream N_m[+1] codon and the corresponding non-methylated codon. For genes containing
multiple upstream near-cognate codons, each ratio was plotted independently.

Codon bias analysis
Using a list of genes that were expressed in HeLa HR-Ribo-seq, 5'UTR sequences of the longest
transcript per gene were extracted from Ensembl/113 with biomart. This was also performed for
genes containing N_m[+1]. The frequencies of AUG and near-cognate codons were calculated for
Modified codons in the 5'UTRs and for the presence of these codons in the 5'UTRs of expressed
protein-coding genes. Codon frequencies were analyzed by performing a chi-square test of
proportion with a Benjamini-Hochberg correction.

Gene ontology analysis

Genes containing A_mUG, C_mUG, and G_mUG in the 5'UTR were used for gene ontology (GO)

analysis for three ontologies: biological process, cellular component, and molecular function. An

R script was used to perform and visualize this analysis using the clusterProfiler/4.14.6 and

org.Hs.eg.db/3.20.0 packages. The top 5 categories for each ontology were displayed by

descending *p-value*.

Analysis of translation efficiency

To evaluate the impact of 2'O-methylation of cognate AUGs on the translation of the canonical

ORF, we downloaded the translation efficiency (TE) values of HeLa mRNAs reported in Arango

et al., 2018 (GSE102113).[27] mRNAs were binned based on the methylation status as AUG or

A_mUG and visualized as box plots in Rstudio/v4.4.0 using packages ggpubr and ggplot2.

Statistical significance between groups was assessed using a Wilcoxon rank-sum test.

Other statistical analysis

All other statistical tests and plotting were performed in Rstudio/v4.4.0. The specific statistical

tests, graph type, variables, number of observations per variable, and *p-values* are indicated in

the figures or their legends.

Data availability

This study did not generate original sequencing or structural data.

References

1. Brito Querido, J., Diaz-Lopez, I. & Ramakrishnan, V. The molecular basis of translation

initiation and its regulation in eukaryotes. *Nat Rev Mol Cell Biol* **25**, 168-186 (2024).

2. Kozak, M. Point mutations define a sequence flanking the AUG initiator codon that

modulates translation by eukaryotic ribosomes. *Cell* **44**, 283-92 (1986).

3. Kearse, M.G. & Wilusz, J.E. Non-AUG translation: a new start for protein synthesis in

eukaryotes. *Genes Dev* **31**, 1717-1731 (2017).

4. Kozak, M. Pushing the limits of the scanning mechanism for initiation of translation.

Gene **299**, 1-34 (2002).

5. Lee, D.S.M. et al. Disrupting upstream translation in mRNAs is associated with human

disease. *Nat Commun* **12**, 1515 (2021).

6.   Calvo, S.E., Pagliarini, D.J. & Mootha, V.K. Upstream open reading frames cause
widespread reduction of protein expression and are polymorphic among humans. *Proc*
Natl Acad Sci U S A **106**, 7507-12 (2009).

7.   Dasgupta, A. & Prensner, J.R. Upstream open reading frames: new players in the
landscape of cancer gene regulation. *NAR Cancer* **6**, zcae023 (2024).

8.   Arango, D. et al. Direct epitranscriptomic regulation of mammalian translation initiation
through N4-acetylcytidine. *Mol Cell* **82**, 2797-2814 e11 (2022).

9.   Simonetti, A., Guca, E., Bochler, A., Kuhn, L. & Hashem, Y. Structural Insights into the
Mammalian Late-Stage Initiation Complexes. *Cell Rep* **31**, 107497 (2020).

10. Galloway, A. & Cowling, V.H. mRNA cap regulation in mammalian cell function and fate.
Biochim Biophys Acta Gene Regul Mech **1862**, 270-279 (2019).

11. Mauer, J. et al. Reversible methylation of m(6)A(m) in the 5' cap controls mRNA stability.
Nature **541**, 371-375 (2017).

12. Akichika, S. et al. Cap-specific terminal N (6)-methylation of RNA by an RNA polymerase
II-associated methyltransferase. *Science* **363**(2019).

13. Guca, E. et al. N(6)-methyladenosine in 5' UTR does not promote translation initiation.
Mol Cell **84**, 584-595 e6 (2024).

14. Zhou, J. et al. Dynamic m(6)A mRNA methylation directs translational control of heat
shock response. *Nature* **526**, 591-4 (2015).

15. Meyer, K.D. et al. 5' UTR m(6)A Promotes Cap-Independent Translation. *Cell* **163**, 999-
1010 (2015).

16. Beiki, H. et al. Detection of ac4C in human mRNA is preserved upon data reassessment.
Mol Cell **84**, 1611-1625 e3 (2024).

17. Li, Y. et al. 2'-O-methylation at internal sites on mRNA promotes mRNA stability. *Mol Cell*
84, 2320-2336 e6 (2024).

18. Chen, L. et al. Nm-Mut-seq: a base-resolution quantitative method for mapping
transcriptome-wide 2'-O-methylation. *Cell Res* **33**, 727-730 (2023).

19. Tang, Y. et al. An integrative platform for detection of RNA 2'-O-methylation reveals its
broad distribution on mRNA. *Cell Rep Methods* **4**, 100721 (2024).

20. Dai, Q. et al. Nm-seq maps 2'-O-methylation sites in human mRNA with base precision.
Nat Methods **14**, 695-698 (2017).

21. Petrychenko, V. et al. Structural basis for translational control by the human 48S
initiation complex. *Nat Struct Mol Biol* **32**, 62-72 (2025).

22. Brito Querido, J. et al. Structure of a human 48S translational initiation complex. *Science*
369, 1220-1227 (2020).

23. Hertler, J. et al. Synthesis of point-modified mRNA. *Nucleic Acids Res* **50**, e115 (2022).

24. Dong, Z.W. et al. RTL-P: a sensitive approach for detecting sites of 2'-O-methylation in
RNA molecules. *Nucleic Acids Res* **40**, e157 (2012).

25. Ayadi, L., Galvanin, A., Pichot, F., Marchand, V. & Motorin, Y. RNA ribose methylation (2'-
O-methylation): Occurrence, biosynthesis and biological functions. *Biochim Biophys Acta*
Gene Regul Mech **1862**, 253-269 (2019).

26. Elliott, B.A. et al. Modification of messenger RNA by 2'-O-methylation regulates gene
expression in vivo. *Nat Commun* **10**, 3401 (2019).

27. Arango, D. et al. Acetylation of Cytidine in mRNA Promotes Translation Efficiency. *Cell*
175, 1872-1886 e24 (2018).

28. Llacer, J.L. et al. Conformational Differences between Open and Closed States of the
Eukaryotic Translation Initiation Complex. *Mol Cell* **59**, 399-412 (2015).

29. Zhou, F., Zhang, H., Kulkarni, S.D., Lorsch, J.R. & Hinnebusch, A.G. eIF1 discriminates
against suboptimal initiation sites to prevent excessive uORF translation genome-wide.
RNA **26**, 419-438 (2020).

30. Thakur, A. & Hinnebusch, A.G. eIF1 Loop 2 interactions with Met-tRNA(i) control the
accuracy of start codon selection by the scanning preinitiation complex. *Proc Natl Acad*
Sci U S A **115**, E4159-E4168 (2018).

31. Thakur, A., Marler, L. & Hinnebusch, A.G. A network of eIF2beta interactions with eIF1
and Met-tRNAi promotes accurate start codon selection by the translation preinitiation
complex. *Nucleic Acids Res* **47**, 2574-2593 (2019).

32. Hussain, T. et al. Structural changes enable start codon recognition by the eukaryotic
translation initiation complex. *Cell* **159**, 597-607 (2014).

33. Yi, S.H. et al. Conformational rearrangements upon start codon recognition in human
48S translation initiation complex. *Nucleic Acids Res* **50**, 5282-5298 (2022).

34. Abou Assi, H. et al. 2'-O-Methylation can increase the abundance and lifetime of
alternative RNA conformational states. *Nucleic Acids Res* **48**, 12365-12379 (2020).

35. Kawai, G. et al. Conformational rigidity of specific pyrimidine residues in tRNA arises
from posttranscriptional modifications that enhance steric interaction between the base
and the 2'-hydroxyl group. *Biochemistry* **31**, 1040-6 (1992).

36. Thiaville, P.C. et al. Global translational impacts of the loss of the tRNA modification
t(6)A in yeast. *Microb Cell* **3**, 29-45 (2016).

37.   Choi, J. et al. 2'-O-methylation in mRNA disrupts tRNA decoding during translation
elongation. *Nat Struct Mol Biol* **25**, 208-216 (2018).
38.   Zhou, K.I., Pecot, C.V. & Holley, C.L. 2'-O-methylation (Nm) in RNA: progress,
challenges, and future directions. *RNA* **30**, 570-582 (2024).
39.   Incarnato, D. et al. High-throughput single-base resolution mapping of RNA 2´-O-
methylated residues. *Nucleic Acids Res* **45**, 1433-1441 (2017).
40.   Marchand, V., Blanloeil-Oillo, F., Helm, M. & Motorin, Y. Illumina-based RiboMethSeq
approach for mapping of 2'-O-Me residues in RNA. *Nucleic Acids Res* **44**, e135 (2016).
41.   Filatova, A. et al. Annotation of uORFs in the OMIM genes allows to reveal pathogenic
variants in 5'UTRs. *Nucleic Acids Res* **51**, 1229-1244 (2023).
42.   Soukarieh, O. et al. Common and Rare 5'UTR Variants Altering Upstream Open Reading
Frames in Cardiovascular Genomics. *Front Cardiovasc Med* **9**, 841032 (2022).
43.   Whiffin, N. et al. Characterising the loss-of-function impact of 5' untranslated region
variants in 15,708 individuals. *Nat Commun* **11**, 2523 (2020).
44.   Martin-Geary, A.C. et al. Systematic identification of disease-causing promoter and
untranslated region variants in 8040 undiagnosed individuals with rare disease. *Genome*
Med **17**, 40 (2025).
45.   Zhao, J. et al. Generating snoRNA-guided Programmable 2'- O -methylation. *bioRxiv*
(2024).
46.   Huang, C. & Yu, Y.T. Targeted 2'-O methylation at a nucleotide within the pseudoknot of
telomerase RNA reduces telomerase activity in vivo. *Mol Cell Biol* **30**, 4368-78 (2010).
47.   Zhao, X. & Yu, Y.T. Targeted pre-mRNA modification for gene silencing and regulation.
Nat Methods **5**, 95-100 (2008).
796
797

Figure 1. 2'OH hydrogen bonds within translation initiation complexes.

a, schematic representation of the intermolecular interactions within translation pre-initiation
complexes. **b-d,** Cryo-EM visualization of rabbit 80S initiator ribosomes captured in complexed
with harringtonine (PDB:7UCJ, Arango et al., 2022)[8] and centered around the interaction of the
2'OH group in nucleotide C[-1]^{mRNA} with t^{6}A in tRNA$_i^{met}$ (b); centered around the interaction of the
2'OH group in nucleotide A[+1]^{mRNA} with the PO_4^{3-} group of U1830 in 18S rRNA (c); or centered
around the interaction of the 2'OH group in nucleotide A[+1]^{mRNA} with the C[2]=O group of U1830
in 18S rRNA. **e-f,** cryo-EM visualization of human 48S pre-initiation complexes (PDB:8PJ3
Petrychenko et al., 2025)[21] centered around the interaction of the 2'OH group in nucleotide
G[+3]^{mRNA} with lysin 7 in eIF1A; or centered around the interaction of the 2'OH group in nucleotide
A[-3]^{mRNA} with G961 in 18S rRNA. All structures were downloaded from PDB and visualized using
Pymol.

Figure 2. 2'O-methylation of start codons inhibits protein synthesis.

a, experimental design of a three-way splint ligation approach to insert 2'O-methylation in single
positions within mRNA start codons. **b,** relative light units (RLU) were measured in cell lysates
from HEK293T cells transfected with ligated NanoLuc mRNAs, with or without a start codon, or
transfected with unligated mRNAs as control. Mean ± SEM, n = 3. p = One-way ANOVA with
Tukey's post hoc analysis. **c,** relative light units were measured in lysates from HEK293T cells
co-transfected with increasing amounts of full-length NanoLuc mRNA reporters and constant
amounts (50 ng) of unligated mRNA. Mean ± SEM, n = 3. A linear regression demonstrates the
linearity of the assay. **d,** chemical structures of unmethylated and 2'O-methylated nucleosides. **e,**
relative light units were measured in lysates from HEK293T cells transfected with ligated NanoLuc

mRNAs carrying 2'O-methylation in different positions around the start codon. Unmethylated
(blue) and unligated mRNAs were used as controls. Mean ± SEM, n = 4. p = One-way ANOVA
with Tukey's post hoc analysis. **f,** relative light units were measured in lysates from HEK293T
cells co-transfected with unmodified mRNAs encoding Firefly luciferase and unmodified or A_m[+1]-
modified mRNAs encoding NanoLuc. Mean ± SEM, n = 4. **g,** NanoLuc mRNA levels were
measured using RT-qPCR at different time intervals post-transfection. Primers were designed to
ensure that only ligated mRNA is detected. 18S rRNA was used as a normalizing control. Mean
± SEM, n = 3. **h,** representative histograms of fluorescence intensity measured by flow cytometry
in HeLa cells transfected with ligated GFP mRNA reporters carrying 2'O-methylation in different
positions around the start codon. Unmethylated (blue) and unligated mRNAs were used as
controls. **i,** quantification of data from (h). Mean ± SEM, n = 3. p = One-way ANOVA with Tukey's
post hoc analysis.

Figure 3. Inhibition of non-AUG translation by 2'O-methylation of near-cognate start codons.

a, schematic representation of canonical and upstream translation initiation. **b,** distribution of
upstream initiation codons obtained from HR-Ribo-seq (Arango et al., 2022). 8 c, three-way splint
ligation approach to insert 2'O-methylation a position N[+1] of CUG and GUG start codons. **d,**
relative light units were measured in lysates from HeLa cells transfected with unmodified, C_mUG,
or G_mUG-modified NanoLuc mRNA reporters. Mean ± SEM, n = 2. **e,** normalized RLU relative to
unmodified mRNA. Mean ± SEM, n = 4. p = two-tailed *Student's t-test*. **f,** schematic representation
of translation initiation efficiency by AUG, CUG, GUG, and their respective 2'O-methylated
versions.

843
844
845

 Figure 4. Identification of 2'O-methylation in AUG and near-cognate codons.

a, 2'O-methylation maps from four different studies, N_m-seq,[20] N_m-mut-seq,[18] NJU-seq,[19] and
Nanopore-DRS[17] were compiled and intersected with the coordinates for canonical AUG start
codons in human transcriptomes. Data from four different cell lines is shown in the dot plots. **b,**
sites with 2'O-methylation in position N[+1] of upstream AUG, or any near-cognate codon in
5'UTRs. Data from all different studies and cell lines described in (a) are included in the bar plots.
c, the fraction of each N_m[+1]-methylated codon was compared to the distribution of AUG and all
near cognate codons in the 5'UTR of human mRNAs. p = chi-squared test for proportions. * p <
0.01, ** p < 0.001, *** p < 0.0001. **d,** schematic of RTL-P. Reverse transcription was performed
in the presence of an optimal dNTP concentration (1 mM) or suboptimal dNTP concentration
(1 μM) that induces a premature RT-STOP. A pair of primers surrounding the 2'O-methylation is
used during PCR amplification. In this manner, the amplicon signal is an indicator of RT-STOP
efficiency and N_m stoichiometry. Another pair of primers, located downstream of the 2'O-
methylation site, serves as an internal control for mRNA levels. **e,** Western blot validation of FBL
knockdown using two different siRNAs in HeLa cells. **e-h,** *left:* representative agarose gels of
amplicon density for the 2'O-methylated site (top) and the loading control (bottom). *Right:*
quantification of the relative N_m ratio dependent on FBL. Mean ± SEM, n = 3. p = One-way ANOVA
with Tukey's post hoc analysis.

Figure 5. 2'O-methylation inhibits upstream translation initiation.

a, averaged HR-Ribo-seq density (RPKMs) over a window of -13 to +16 nucleotides around
position N[+1] of canonical AUG or upstream AUG, CUG, GUG, and AUC. **b,** Relative HR-Ribo-
seq density was obtained through the $\log_2$ of the fold change between the RPKMs at the
unmodified (blue) or N_m[+1]-modified (red) upstream codon relative to the RPKMs at the
downstream canonical AUG within the same transcript. Only sites with RPKM values ≥1 were
included. Box plots summarize the median across all sites. p = two-sided Wilcoxon rank sum test.
c-d, averaged HR-Ribo-seq density (RPKMs) over a window of -13 to +16 nucleotides around
2'O-methylated codons at position N_m[+1] (red) or 2'O-methylated codons at positions N_m[-3, -2,
-1, and +2] (blue). Only sites with RPKM values ≥1 across all positions were included.

Figure 6. 2'O-methylation inhibits start codon recognition.

a-c, schematic representation of translation inhibition (a), mechanisms of 48S inhibition by
GMPPNP (b), and mechanisms of initiator 80S inhibition by HR (c). **d-f,** Cy5-labeled synthetic
mRNAs containing unmodified or A_mUG-modified start codons were incubated with untreated
reticulocyte lysates (c), treated with GMPPNP (d), or treated with HR (e). Reactions were
fractionated through a sucrose density gradient, and fluorescence (Cy5) intensity was recorded
concomitantly with fractionation. **g,** model: 2'O-methylation inhibits start codon recognition by
making an otherwise optimal translation initiation site look like a non-cognate codon.
Consequently, the 48S subunit continues scanning until encountering a downstream start codon
or, alternatively, dissociates from the mRNA.

886
887
888

889 **Extended Data Figure 1.** H-bonds between the 2'OH groups of mRNA nucleotides with
890 tRNA$_i^{met}$ and 18S rRNA. Related to Fig. 1.

891
a, Cryo-EM visualization of late-stage rabbit 48S pre-initiation complexes (PDB:6YAL, Simonetti
et al., 2022)[9] centered around the interaction of the 2'OH group in nucleotide C[-1]^{mRNA} with t^{6}A in
tRNA$_i^{met}$. This interaction was reproduced in rabbit 80S initiator ribosomes in complex with
Harringtonine (Fig, 1B). **b-f,** Cryo-EM visualization of 48S pre-initiation complexes (Petrychenko
et al., 2025)[21] centered around the interaction of the 2'OH group in nucleotide A[+1]^{mRNA} with
U1830 in 18S rRNA across different stages of 48S pre-initiation complexes, including the stage
of codon recognition and GTP hydrolysis (b, PDB: 8PJ2), binding of eIF5B (c, PDB: 8PJ3),
followed by sequential release of eIF5B (d, PDB: 8PJ4) and eIF2 (e, PDB: 8PJ5), and the late-
stage readiness for 60S subunit joining (f, PDB: 8PJ6). All structures were downloaded from PDB
and visualized using Pymol.

Extended Data Figure 2. Quality controls for the three-way splint ligation and translation
assays. Related to Fig. 2.

904
a, the 5' end of the mRNA reporters, which lacked a start codon, was transcribed and capped *in*
vitro. The integrity of the RNA was verified by polyacrylamide gel electrophoresis (PAGE) and
SYBR gold staining. **b,** successful capping of the *in vitro* transcribed 5' ends was verified by dot
blot using anti-m^{7}G antibodies. Methylene blue staining was used as a loading control. **c,** the 3'

end of the mRNA reporters, containing the ORF and the 3'UTR but lacking a start codon, was
transcribed and polyadenylated *in vitro*. The integrity of the RNA and the efficiency of
polyadenylation were verified using PAGE and SYBR gold staining. **d,** the ligation efficiency of
NanoLuc mRNA reporters was verified using PAGE and SYBR gold staining. Ligated mRNA
shows a shift in molecular weight compared to unligated mRNA. **e,** relative light units were
measured in lysates from HEK293T cells co-transfected with increasing amounts of unligated
RNA pieces and constant amounts (20 ng) of full-length NanoLuc mRNA. Mean ± SEM, n = 3.
The line plot shows stable NanoLuc activity across different amounts of unligated RNA. **f,** Integrity
of the synthetic 5' phosphorylated RNA oligos containing unmodified or 2'O-methylation in
different positions around the start codon. **g,** the efficiency of ligation was verified by RT-qPCR.
Two sets of primers were used (Supplementary Table 1). The first set specifically recognizes the
ligated mRNA, while the second set binds NanoLuc ORF to obtain total levels of mRNA. The
efficiency of ligation was calculated as: (ligated mRNA)/(total RNA)*100. **h,** NanoLuc activity
normalized by transfected mRNAs, which was obtained using RT-qPCR from the RNA transfected
into HEK293T cells. Mean ± SEM, n = 3. **i,** relative light units were measured in lysates from HeLa
cells transfected with ligated NanoLuc mRNAs carrying 2'O-methylation in different positions
around the start codon. Unmethylated (blue) and unligated mRNAs were used as controls. Mean
± SEM, n = 4. p = One-way ANOVA with Tukey's post hoc analysis. **j,** denaturing agarose gel
electrophoresis indicating the ligation efficiencies for the GFP mRNA reporters. These RNA
pieces are larger than the ones used for NanoLuc reporters and do not resolve well in PAGE gels.
Thus, agarose gels are better suited for their separation. **k,** schematic indicating the relative
lengths of 5'UTRs in the NanoLuc and GFP mRNA reporters.
931

932 **Extended Data Figure 3.** Upstream AUGs and near-cognate codons are prevalent in human
933 transcriptomes. Related to Fig. 3.

934
a-b, the 5'UTR sequence of all canonical protein-coding transcripts was downloaded from Biomart
and analyzed to estimate the percentage of transcripts with one or more upstream AUGs (a) and
the percentage of transcripts with one or more upstream near-cognate codon (b). This analysis
indicated that 99% of human transcripts contain at least one near-cognate codon in their 5'UTRs,
whereas 50% of human transcripts contain at least one AUG in their 5'UTR sequences. **c,**
distribution of upstream near-cognate or AUG codons per transcript. CUG is the most prevalent
near-cognate codon in human transcripts. **d,** while virtually all human transcripts contain cryptic
initiation sites in their 5'UTRs, not all are used at a given moment or physiological condition. To
determine the upstream near-cognate or AUG codons used for initiation, we previously performed
HR-Ribo-seq in HeLa cells.[8] In this method, the inhibitor Harringtonine (HR) stalls the initiator
ribosome while letting the elongator ribosomes slide off of mRNAs. Subsequent RNAse
footprinting and isolation of ribosome-protected fragments (RPFs) allow the identification of
initiation sites transcriptome-wide. **e,** HR-Ribo-seq indicates that ~70% of transcripts initiate
translation from at least one upstream site in wildtype HeLa cells. **f,** the integrity of ligated
NanoLuc mRNAs carrying near-cognate codons was verified using PAGE and SYBR gold
staining. **g,** relative light units were measured in lysates from HeLa cells transfected with
unmodified, C_mUG, or G_mUG-modified NanoLuc mRNAs. Mean ± SEM, n = 4. p = two-tailed
Student's t-test.
953

Extended Data Figure 4. Distribution of 2'O-methylation sites in human mRNAs. Related to Fig. 4.

a

Distribution of N_m sites per study per cell line

Nanopore-DRS
Li et. al. 2024

NJU-seq
Tang et. al. 2024

N_m-mut-seq
Chen et al. 2023

N_m-seq
Dai et. al. 2018

Cell Line
● HeLa
● HEK293T
● HepG2
● A549

b

N_m sites around canonical start codons

Nanopore-DRS
Li et. al. 2024

NJU-seq
Tang et. al. 2024

N_m-mut-seq
Chen et al. 2023

N_m-seq
Dai et. al. 2018

$A_m[+1]$

Position relative to canonical AUG

Cell Line
● HeLa
● HEK293T
● HepG2
● A549

c

Prevalence of $N_m[+1]$ per codon

HEK293

HeLa

HepG2

A549

a, 2'O-methylation maps from four different studies, N_m-seq, N_m-mut-seq, NJU-seq, and Nanopore-DRS were compiled and binned based on the transcript location (5'UTR, CDS, or 3'UTR) within mRNA transcripts. Data from four different cell lines is shown in the dot plots. **b,** the genomic coordinate of 2'O-methylation sites were intersected with the coordinates for canonical AUG start codon in human transcriptomes. Sites within a window of -3 to +4 nucleotides are reported in four different cell lines. **c,** the prevalence of 2'O-methylation sites in upstream AUGs or near-cognate codons per cell line. Data from all different studies and cell lines described in (a) are included in the bar plots.

Extended Data Figure 5. Validation of 2'O-methylation sites in near-cognate codons. Related to Fig. 4.

a, list of human 2'O-methyltransferase enzymes. **b,** list of validated sites. All tested sites were upstream AUGs or near-cognate codons with 2'O-methylation in the first nucleotide of the codon. Sites were selected based on the estimated modification ratio (the top sites) from the Nanopore-DRS method[17]. **c-h,** *left:* representative agarose gels of amplicon density for the 2'O-methylated site (top) and the loading control (bottom). *Right:* quantification of the relative N_m ratio dependent on FBL. Mean ± SEM, n = 3. One-way ANOVA with Tukey's post hoc analysis.

975 **Extended Data Figure 6.** Ribosome density in upstream 2'O-methylated AUG, CUG, and GUG
976 codons. Related to Fig. 5.

977
978 **a-c,** averaged HR-Ribo-seq density (RPKMs) over a window of -13 to +16 nucleotides around
979 2'O-methylated codons at position $N_m[+1]$ (red) or 2'O-methylated codons at positions $N_m[-3]$, $N_m[-2]$, $N_m[-1]$, and $N_m[+2]$. **d,** summarized translation efficiency (TE) in canonical A_mUG codons (n=5)
981 and canonical unmethylated AUG (n=7,091) start sites in HeLa cells. Translation efficiencies were
982 obtained from Arango et al., 2018.[27]
983

984   **Extended Data Figure 7.** Gene ontology analysis. Related to Fig. 5.
985

a

b

c

986   **a-c,** gene ontology analysis of all genes containing A_mUG, C_mUG, and G_mUG (a), containing only
987   C_mUG (b), or containing only G_mUG (c) codons in their 5'UTRs.