Title
Embracing the power of genomics to inform evolutionary significant units

Authors
Jayna C. Bergman[1*], Rebecca S. Taylor[1], Micheline Manseau[1]
[1]*Landscape Science and Technology, Environment and Climate Change Canada, Colonel By*
Drive, Ottawa, ON K1S 5B6, Canada
8
9 [*]Corresponding author. Email: jaynabergman@gmail.com
10

11 **Keywords**
12 conservation units, conservation genomics, genetics, discreteness, adaptive variation,
13 phylogeographic history
14

Acknowledgements

This research was supported by Environment and Climate Change Canada (ECCC).

Conflicts of Interest

The authors declare no conflicts of interest.

Data Availability Statement

The authors have nothing to report.

Abstract

Appropriate identification of evolutionary significant units (ESUs) is essential for effective conservation planning. Genomic data has emerged as a key tool to inform ESU decisions due to the increased resolution of information, yet little is known about how genomic data is being used in practice to inform these decisions. To address this, we conducted a systematic literature review and found that genomic data is increasingly being used to inform ESUs globally and across plant and animal taxa. We identified a lack of consistency in the methods used to analyze the genomic data, although with reduced representation sequencing and whole genome sequencing improving the ability to infer evolutionary significance. Adaptive loci, a key advantage of genomic data, need to be interpreted with caution and simply identifying these loci may lead to the inflation of ESUs. Our review further revealed inconsistencies in how ESUs are defined and that conclusions do not always meet the criteria outlined in the definition provided. For instance, the amount of gene flow determined rarely influenced the number of ESUs recommended. Overall, 67% of the studies included in this review suggested an increase in the number of ESUs regardless of vast differences across studies. A unifying definition of what constitutes an ESU and a detailed workflow that can provide reliable support to assess the evolutionary significance of these units is still needed. The unparalleled depth of information obtained from genomic data, and the application of a proven workflow has the potential to improve the identification of ESUs globally.

Introduction

Identifying intraspecific units for conservation is a critical strategy for protecting diversity

within species (Funk et al., 2012; Coates et al., 2018; Hohenlohe et al., 2020; Allendorf et al.,

2022; Forester et al., 2022; Turbek et al., 2023; Schmidt et al., 2024). One widely used concept is

Evolutionary Significant Units (ESU), which are biologically informed and often legally recognized

groups of individuals below the species level. ESUs are broadly recognized as being (1)

geographically or genetically discrete and (2) representative of an important component of a

species evolutionary history (see reviews by Crandall et al., 2000; Fraser and Bernatchez, 2001;

Funk et al., 2012; Turbek et al., 2023). The identification of ESUs can have direct implications for

the legal status and management of populations (Coates et al., 2018), and inappropriate

identification can waste limited conservation resources (Forester et al., 2022). Therefore,

carefully considering how ESUs are described is essential for effective conservation planning.

However, no single definition of ESUs is universally used and the general concept has

evolved over time due to ongoing debate about which criteria should be emphasized (e.g.

Ryder, 1986; Waples, 1991; Moritz, 1994; Fraser and Bernatchez, 2001; Funk et al., 2012). Ryder

(1986) first introduced the term ESU for conservation purposes and described them as

populations that "possess genetic attributes significant for the present and future generations

of the species in question". Other proposed definitions have highlighted different criteria. For

example, Waples (1991) emphasized the importance of reproductive isolation, while Fraser and

Bernatchez (2001) focused on requiring highly restrictive gene flow. Moritz (1994) argued the

use of specific genetic markers and determined that ESUs should be "reciprocally monophyletic

for mitochondrial DNA alleles and show significant divergence of allele frequencies at nuclear

loci". More recently, Funk et al., (2012) proposed that ESUs should be "historically isolated from

each other and likely have important adaptive differences among them", integrating both

neutral and adaptive variation for this process. In contrast, Crandall et al., (2000) questioned the

utility of the ESU concept altogether, and suggested "that the terminology [ESU] is abandoned

and replaced with a more holistic concept of species". In addition to these biological definitions,

there are also legally defined units designated by governing bodies to implement policy for legal

protection and conservation status that overlaps conceptually with ESUs (Coates et al., 2018;

Lehnert et al., 2023). For example, Designatable Units (DUs) are delineated by the Committee

on the Status of Endangered Wildlife in Canada (COSEWIC, 2023), and Distinct Population

Segments (DPS) are delineated by the U.S. Endangered Species Act (US Fish and Wildlife

Services, 1996). These legal units tend to use separate definitions with different criteria to

inform the designations. Despite these varying perspectives, ESUs are widely used in

conservation, underscoring the need for greater clarity and standardization in how they are

identified.

Genomic data are emerging as a key tool in the complex decision-making process of

identifying ESUs (Funk et al., 2012; Barbosa et al., 2018; Waples and Lindsey, 2018; Funk et al.,

2019; Turbek et al., 2023). Traditional genetic markers, such as mitochondrial DNA (mtDNA) and

microsatellites, have been commonly used to identify ESUs (e.g. Moritz, 1994; Kanthaswamy et

al., 2006; Jensen et al., 2014). However, these markers typically target only a small number of

loci (e.g. <25), providing limited genetic information that largely reflects neutral evolutionary

processes (Funk et al., 2019; Barbosa et al., 2020; Ghildiyal et al., 2023; Chinna et al., 2025). In

contrast, genomic data generated through high throughput sequencing can sample thousands

to millions of loci across the entire genome, offering much higher resolution on both neutral

and adaptive variation (Coates et al., 2018; Ghildiyal et al., 2023; see Box 1 for an overview of

genomic sequencing approaches). This increased analytical power facilitates a wide range of

downstream analyses (Schmidt et al., 2024) that can be useful for identifying ESUs.

The unparalleled volume of information provided by genomic data has significantly

advanced the ability to accurately identify ESUs (Funk et al., 2012; Coates et al., 2018). Genome-

wide patterns of variation can reveal phylogenetic relationships with high precision (Barbosa et

al., 2020), demographic models can infer past evolutionary processes (Schmidt et al., 2024), and

putatively adaptive loci showing pronounced signatures of selection can be identified (Funk et

al., 2012; Barbosa et al., 2018). However, some types of genomic data may miss highly localized

signatures of adaptation (e.g. RADseq; Lou et al., 2021), that could lead to overlooking genetic

variation essential to a population's persistence. A more comprehensive approach is whole-

genome sequencing (WGS), which provides deep insights into a species' evolutionary history

(e.g. Ellegren, 2014; Jensen et al., 2021; Taylor et al., 2024b). When high coverage data are

available, WGS enables accurate genotype calling and the detection of regions under selection

(Fuentes-Pardo and Ruzzante, 2017; Lou et al., 2021). Despite the clear advantages of genomic

data in identifying ESUs (Funk et al., 2012; Ghildiyal et al., 2023; Schmidt et al., 2024), and the

availability of several published workflows to help guide analyses (e.g. Funk et al., 2012; Turbek

et al., 2023; Lehnert et al., 2023), it remains unclear how genomic data are being used in

practice to support these decisions.

Aims of the study

The aim of this study was to determine how genomic data is being used when directly

applied to identify ESUs across taxa. To achieve this, we reviewed published literature that uses

genomic data to inform ESU identification. Specifically, we aimed to identify the variation in data

reporting among conservation genomic studies, the methods used to analyze the genomic data,

and the definitions and criteria used to define what an ESU is. We further aimed to uncover how

these decisions directly impacted the conclusions drawn on the number of ESUs suggested.

Subsequently, we examined how WGS are used in ESU identification to understand if the

increased amount of information available with WGS, when compared to other genomics

methods, is being utilized to uncover additional variation. Finally, we offer recommendations for

considerations to make when using genomic information to identify ESUs.

Literature Review

We conducted a literature search in Web of Science for studies that use genomic data to

help inform the identification of ESUs. We included studies that were published until January

30, 2024, by searching the title, abstract, or keywords with the search string (genom* OR (WGS

OR "whole genome" OR *seq OR GBS OR "genotyping-by-sequencing" OR SNP* OR "single

nucleotide polymorphism*" OR "target capture" OR "target DNA" OR ddRAD OR "reduced

representation sequencing" OR "next-generation sequencing" OR NGS OR transcript* OR *omics

OR "adaptive loci" OR "adaptive locus") AND ("conservation unit*" OR "evolutionary significant

unit*" OR "evolutionarily significant unit*" OR "adaptive unit*" OR "distinct population

segment*" OR "designatable unit*"). We supplemented our search with articles from Scopus

using the same set of search terms. Resulting studies from both databases were combined and

duplicate studies were removed. The titles and abstracts of the retrieved articles were first

manually screened with the assistance of ASReview (ASReview LAB developers, 2023), followed

by full-text review for those that met the initial criteria. We excluded studies that did not use

genomic data in their analyses (e.g., irrelevant methods such as using only mtDNA,

microsatellites, or cpDNA) and those that did not make direct conclusions about the number of

ESUs (e.g., irrelevant topics such as generating a reference genome, assigning individuals to

existing ESUs, or species delineation without a direct link to ESUs). For synthesis papers that did

not include analysis of original data, we reviewed their cited references to identify additional

relevant studies. We included studies on all vertebrate, invertebrate, and plant species and did

not apply any geographic restrictions. In total, our search terms identified 415 articles of

potential relevance (Table S1). Of these, 154 met our inclusion criteria and were evaluated in

the literature review (Table S2).

144   For each study we recorded whether a written definition of what is considered an ESU

145   was provided, and which reference was used to support that definition. We then extracted

146   information on sampling details and the genomic datasets used from both the main text and

147   supplementary materials as needed. If a study included separate datasets for more than one

148   species, each species was treated independently and assigned a separate row in the data table.

149   This resulted in 161 data entries in the final dataset used (Table S2).

150   The methods used to analyze the genomic data were recorded as either done or not

done, with additional columns noting the specific programs used. All analytical methods were

manually classified into two broad groups: methods assessing "discreteness" and those

assessing "evolutionary significance". Methods classified under "discreteness" included

clustering and ancestry-based approaches, F_{st}/AMOVA, assessments of gene flow, tests for

isolation-by-distance, and network analyses. Methods under "evolutionary significance"

included phylogenetic analyses to uncover deeper evolutionary relationships, assessments of

demographic history, tests for separate glacial refugia, and estimating divergence times.

Because we were particularly interested in identifying trends related to adaptive loci, we

separately categorized studies that identified them. For these studies, we noted whether

adaptive loci were detected using F_{ST} outlier analyses or genotype-environment association

(GEA) approaches. We also recorded whether these loci were used as a dataset to explore

population structure or were simply identified as being present. In addition, for studies that

assessed gene flow, we determined if gene flow was directly calculated (e.g. migration rates, D-

statistic, f-statistics) or inferred (e.g. Admixture plots, presence of hybrids, amount of genetic

differentiation). Based on the study's conclusions, we categorized the amount of gene flow

determined into three groups: lots of gene flow, limited gene flow, and no gene flow.

Overall, we aimed to understand the impact of genomic data on the suggested number

of ESUs. To do this, we extracted each study's final conclusions regarding the number of ESUs

proposed and whether this represented an increase or decrease compared to the previously

identified number of ESUs. Additional details are provided in the supplementary material on

how data for each column was extracted from the studies (Table S3). All plots were generated in

R (R Core Team, 2024) using *ggplot2* (Wickham, 2016) and *patchwork* (Pedersen, 2024).

Results and Discussion

Our review showed that genomic data is used to inform the identification of ESUs across

taxonomic groups around the globe and this use is steadily increasing. All classes of animals

were represented in the collected studies (Fig. 1A), with fish having the most representation

(n=42) and amphibians having the least (n=9). We also found that all continents, excluding

Antarctica, were present in the dataset (Fig. 1B). However, there was a geographical bias in the

studies with 42% done on species in North America, and of those studies 61% focused on

populations within the United States (Fig. S1). Comparatively, some continents were severely

underrepresented. For example, only 2% of the studies were done on species in Africa and 5%

on species in South America (Fig. 1B). Plants were well represented in the collected studies

(n=28), but a geographic bias was identified with 64% of these studies done on species in Asia

(Table S2). Additionally, we identified a rapid increase in the use of genomic data to identify

ESUs starting in 2020 (Fig. 1C). This increased use of genomic data across the world and across

taxa highlights the importance of understanding how this data is produced, analyzed and

interpreted to inform ESUs globally.

Figure 1. (A) The taxonomic groups represented by the 161 data entries included in the review, (B) the continents that the included studies had collected data from, and (C) the year that the study was published in. Our literature search ended in January 2024, and six studies are included in our literature review from that year. These studies are used in all other analyses and conclusions, but the year 2024 is excluded from panel C because it is not representative of the number of studies published in the entire year.

Data quality and Sequencing methods

We found a lack of consistent reporting on key details that are needed to assess the

quality of genomic data used for identifying ESU. In particular, information on filtering decisions

and their effects on the dataset is critical for evaluating the reliability of data (O'Leary et al.,

2018). For example, read depth (also referred to as depth of coverage) is influenced by filtering

decisions and can affect the accuracy of genotype calls, which in turn impacts downstream

analyses (Hemstrom et al., 2024). Strikingly, only 40% of the studies in this review reported

either the average depth of coverage or the range of coverage values across samples (Table S2).

Of the remaining studies, 32% reported the minimum coverage filter that was used to call

genotypes, while 28% did not report any information to indicate coverage (Table S2). We

strongly advocate for all studies to report depth of coverage, in addition to the settings used to

filter genomic datasets.

We also observed inconsistencies in how geographic sampling information was reported.

Some studies did not clearly indicate the number of geographic sampling locations, or the

number of individuals sampled per location. These details are important, as the number of

geographic sampling locations is strongly associated with the ability to detect patterns of

isolation by distance (Hillis and Chambers, 2020), calculate reliable summary statics (e.g. F_{st}

values; Holsinger and Weir, 2009), and identify candidate loci for adaptation (Aguirre-Liguori et

al., 2020), all of which can be used to inform ESU decisions. Sampling information was not

reported in nine of the studies included in our review (Table S2). While full sampling tables were

often provided in the supplementary material, several studies did not summarize this

information in the main text, reducing transparency and making it harder to assess potential

biases or sources of error in the dataset.

We grouped the sequencing methods used in the studies included in our review into six

broad categories (see Box 1; Fig. S2). Reduced representation sequencing was the most

commonly used approach with 103 studies (RRS; inclusive of restriction-site associated

sequencing: RAD-seq, double digest restriction-site associated sequencing: ddRAD-seq,

genotype by sequencing: GBS, and multiplexed inter-simple sequence repeats: MIG-seq; Fig.

S2). The next most common methods were single nucleotide polymorphism (SNP)-based

genotyping (e.g. SNP arrays, SNP assays, SNP chips, and SNP panels) and whole genome

sequencing (WGS), both used in 22 studies (Fig. S2). The remaining sequencing approaches

included targeted gene sequencing (e.g. major histocompatibility complex [MHC] genes and

exon sequencing; n=10), transcriptomics (n=3), and pooled sequencing (pool-seq; n=1). Notably,

most of these less commonly used methods focus on functional genomic regions and are

therefore less suited for analyses to identify population structure or evolutionary history.

Furthermore, pool-seq is often confounded with sequencing errors and individual genotypes are

lost during the pooling process (Fuentes-Pardo and Ruzzante, 2017). Overall, RRS was used

nearly five times more often than any other sequencing approach, but SNP-based genotyping

and WGS were also common methods for generating genomic data to inform ESU identification.

Methods used to analyze genomic data

Among the various analyses available for genomic data, certain methods were more

common across studies (Fig. 2A). For example, 128 studies used a clustering or ordination

approach (e.g. PCA, DAPC) and 133 studies used an ancestry-based approach (e.g. Admixture,

Structure; Fig. 2A). Both these types of analyses are useful to explore population structure and

are recommended as starting points by recent workflows to determine ESUs (see Lehnert et al.,

2023; Turbek et al., 2023). However, not as many studies assessed the demographic history

(n=61), estimated divergence times (n=37), or identified if groups were in separate glacial

refugia (n=9; Fig. 2A). This is surprising, as assessments of demographic history, particularly

those using estimates of effective population size (Ne), have recently been highlighted as

important metrics for tracking genetic diversity (e.g., Hoban et al., 2022; Waples, 2024; Waples,

2025). Although tracking genetic diversity through Ne is not directly linked to the identification

of ESUs, and we do not recommend using Ne alone to define ESUs (Taylor et al., 2024a), it can

serve as an informative metric to help understand the evolutionary history of populations

(Chhina et al., 2025).

When we categorized methods into assessing for either "discreteness" or "evolutionary

significance" (see methods) to inform ESU identification, we found the majority of studies used

methods that could assess both (n=108; Fig. 2B). In contrast, 29 studies assessed only

discreteness and of these studies, six relied solely on clustering analyses and/or ancestry-based

methods (Fig. 2B). Interestingly, 9% of the studies recommended increasing the number of ESUs

even though they assessed only discreteness (Fig. 2B). This raises concerns, as determining

population structure using PCA and admixture-based analyses alone may produce unreliable

results and inaccurately reflect a population's evolutionary history (Lawson et al., 2018; Elhaik,

2022), leading to the misidentification of ESUs. For example, inbred individuals can appear as

distinct clusters on a PCA, which may be misinterpreted as evidence of discrete genetic groups

(Taylor et al., 2024b). We recommend using multiple analyses to thoroughly explore population

structure in ways that assess both discreteness and evolutionary significance when identifying

ESUs. Clear and detailed examples of how to use several methods to provide strong support for

ESU identification, including analyses addressing both discreteness and evolutionary

significance, are provided in Box 2.

Figure 2. Out of the 161 data entries included in our literature review: (A) The number of

studies using different genomic analysis methods to determine how many ESUs should

be suggested. (B) The number of studies that applied methods testing for genetic

discreteness and/or evolutionary significance to inform ESU identification. Methods

classified under "discreteness" included clustering and ancestry-based approaches,

F_{st}/AMOVA, assessments of gene flow, tests for isolation-by-distance, and network

analyses. Methods under "evolutionary significance" included phylogenetic analyses to

uncover deeper evolutionary relationships, assessments of demographic history, tests

for separate glacial refugia, and divergence time estimation. Bar colors indicate the

proportion of studies in each category that recommended an increase, decrease, or no

change in the number of ESUs.

To understand whether the sequencing type influenced the choice of analyses, we

compared the methods used to analyze the genomic data across studies employing SNP-based

genotyping, RRS, and WGS. SNP-based studies accounted for all the studies that solely used

clustering and/or ancestry-based approaches, and they had the lowest proportion of studies to

assess both discreteness and evolutionary significance compared to RRS and WGS (Fig. 3). There

was a large range in the number of SNPs included in these datasets (25 to 95,043 SNPs; Table

S2), but 55% of these studies had fewer than 400 SNPs possibly resulting in a lack of resolution

required to carry out analyses needed to uncover deeper divergence across the genome. SNP-

based genotyping often leave large portions of the genome unsampled (Dokan et al., 2021).

They rely on targeted approaches that require prior knowledge of where variation is occurring

(Chhina et al., 2025), limiting the ability to detect cryptic population structure or assess

evolutionary significance, both of which are critical for informing ESU delineation. Despite these

limitations, the largest proportion of studies using SNP-based genotyping still recommended

increasing the number of ESUs.

In contrast, studies using RRS and WGS tended to use similar and more comprehensive

analyses to identify ESUs (Fig. 3; Fig. S3). In both cases, most studies assessed both discreteness

and evolutionary significance and likewise the majority suggested increasing the number of

ESUs (Fig. 3). However, WGS studies consistently included assessments of evolutionary

significance or identified adaptive loci, whereas a subset of RRS studies assessed only

discreteness (Fig. 3). This may be due to an advantage of WGS, which provides a more

comprehensive view of a species' evolutionary history by sequencing the entire genome rather

than sampling portions of it (Fuentes-Pardo and Ruzzante, 2017). This advantage may be further

highlighted, as a higher proportion of WGS studies estimated demographic histories and

divergence times compared to those using RRS or SNP-based genotyping (Fig. S3). We observed

a rising trend in the use of WGS, which is expected to continue in the coming years (Fig. S4). Our

review included studies published up to January 2024, and of the six studies from one month in

2024, half used WGS (Fig. S4). Altogether this suggests that WGS is becoming increasingly

common, and that the greater genomic resolution may facilitate the use of analyses that

provide stronger support for ESU identification in future studies.

A)

SNP-based genotyping

B)

Reduced representation sequencing

C)

Whole genome sequencing

Figure 3. The proportion of studies that analyzed the genomic data with methods that

tested for genetic discreteness and/or evolutionary significance, separated into different

sequencing methods of: (A) SNP-based genotyping (inclusive of SNP arrays, SNP assays,

and SNP chips; n=22) (B) Reduced representation sequencing (inclusive of restriction-site

associated sequencing: RAD-seq, double digest restriction-site associated sequencing:

ddRAD-seq, Genotype by sequencing: GBS, and Multiplexed Inter-simple sequence

repeat genotype by sequencing: MIG-seq; n = 103), and (C) Whole genome sequencing

(n=22). Methods classified under "discreteness" included clustering and ancestry-based

approaches, F_{st}/AMOVA, assessments of gene flow, tests for isolation-by-distance, and

network analyses. Methods under "evolutionary significance" included phylogenetic

analyses to uncover deeper evolutionary relationships, assessments of demographic

history, tests for separate glacial refugia, and estimated divergence times. The filled color

of the bars show the proportion of studies in each of these categories that suggested

either increasing, decreasing, or no change to occur to the number of ESUs.

Gene flow

Gene flow is widely recognized as a critical evolutionary force that shapes genetic

structure within and among species (Tigano and Friesen, 2016; Aitken et al., 2024; Chhina et al.,

2025). It is typically defined as the movement of genetic information from one genetic group to

another that often follows the migration of individuals but is not limited to just migration events

(Funk et al., 2012; Tigano and Friesen, 2016). Several definitions of ESUs suggest that there

should be a lack of gene flow or reproductive isolation between groups (i.e. no gene flow) as

criteria for defining distinct evolutionary units (e.g. Waples, 1991; Fraser and Bernatchez, 2001;

COSEWIC, 2023). These perspectives may be based on the concern that gene flow could erode

genetic differences, potentially resulting in the loss of local adaptation (Lenormand, 2002). This,

in turn, could reduce the number of ESUs by diminishing the unique genetic diversity that

defines them. Alternatively, it is increasingly recognized that gene flow can occur between

populations adapted to different local conditions, and that local adaptation can be maintained

despite high levels of gene flow (Tigano and Friesen, 2016; Chhina et al., 2025). However,

whether gene flow increases or decreases local adaptation is largely population dependent

(Aitken et al., 2024). Other perspectives on ESUs have pointed out that it may be important to

balance the maintenance of gene flow to preserve adaptive differences (Crandall et al., 2000).

Thus, these different views offer no unified approach and leave substantial room for subjective

interpretation on how to consider gene flow when identifying ESUs.

We assessed the uncertainty surrounding gene flow by analyzing the methods that were

used to determine gene flow by the studies included in our review. We categorized assessments

into directly calculating gene flow (e.g. migration rates, D-statistics, f-statistics) and inferring

gene flow (e.g. admixture plots, amount of genetic differentiation, presence of hybrids). Results

showed that 15% of studies directly calculated gene flow, while 47% inferred it, and 35% did not

assess gene flow (Fig. 4A). Notably, the majority of studies recommended increasing the

number of ESUs, regardless of how gene flow was determined or if it was assessed at all (Fig.

4A). Studies that calculated gene flow never recommended decreasing the number of ESUs,

whereas those inferring gene flow or not assessing it had a portion of studies suggesting a

decrease (14% and 11%, respectively; Fig. 4A).

We then examined whether the amount of gene flow found influenced the suggested

number of ESUs. Based on the language used in the conclusions of the reviewed studies, we

categorized gene flow into three levels: lots of gene flow, limited gene flow, and no gene flow.

These categories were somewhat arbitrary, as there was no consistent threshold that could be

applied across the various methods that were used to assess gene flow. Surprisingly, the

amount of gene flow reported did not appear to influence whether a study suggested increasing

or decreasing the number of ESUs (Fig. 4B). Indeed, even among studies that described "lots of

gene flow," the largest proportion still recommended increasing the number of ESUs (Fig. 4B).

We also found that whether gene flow was calculated or inferred did not affect the level of gene

flow reported (Fig. S5). Overall, these results were unexpected, due to the several definitions of

ESUs that emphasize the importance of limited gene flow or reproductive isolation between

groups (e.g. Waples, 1991; Fraser and Bernatchez, 2001; COSEWIC, 2023). However, we suggest

that some level of gene flow should be allowed between two ESUs, especially if this is the

natural state of the populations. Gene flow among populations can maintain and even increase

genetic diversity (Aitken and Whitlock, 2013; Aitken et al., 2024; Chhina et al., 2025). In such

cases, conservation management should aim to maintain this gene flow, even if populations are

considered as separate ESUs (Chhina et al., 2025). Our results suggest that this perspective may

be shared by several other individuals.

Figure 4. (A) The number of studies that assessed gene flow when determining how

many ESUs to recognize, either by directly calculating gene flow (e.g. migration rates, D-

statistics, f-statistics) or inferring gene flow (e.g. admixture plots, amount of genetic

differentiation, presence of hybrids), as well as the number of studies that did not assess

gene flow. (B) Among studies that assessed gene flow, the author's conclusion regarding

the level of gene flow observed between the proposed ESUs. In both panels, bar colors

indicate the proportion of studies in each category that suggested increasing,

decreasing, or no change in the number of ESUs.

Adaptive loci

A significant advantage of using genomic data over mtDNA or microsatellites to identify

ESUs is the ability to determine adaptive regions of the genome through the detection of

adaptive (or putatively adaptive) loci (Funk et al., 2012; Barbosa et al., 2018; Waples and

Lindsey, 2018). Two common approaches that are often used to identify adaptive loci are F_{st}

outlier analyses, and genotype-environment association (GEA) approaches. F_{st} outlier analyses

(e.g. BayScan, Lositan) detect loci that fall outside the range of the expected distribution of F_{st}

values (Funk et al., 2012). These loci are often assumed to be under divergent selection and

used as proxies for local adaptation (Waples and Lindley, 2018) and therefore thought to be

representative of evolutionary significance. Alternatively, GEA approaches (e.g. RDA, LFMM)

identify associations between allele frequencies and environmental variables, to reveal loci that

are associated with specific environmental conditions (Hoban et al., 2016; Forester et al., 2018).

Workflows to help guide ESU designation often incorporate the identification of adaptive loci

(e.g. Funk et al., 2012; Lehnert et al., 2023; Turbek et al., 2023). However, interpreting adaptive

loci needs to be handled cautiously (Hoban et al., 2016; Allendorf, 2017). F_{st} outliers can result

from processes other than local adaptation (Bierne et al., 2013), such as a species demographic

history (Whitlock and Lotterhos, 2015) or genetic drift (Lotterhos and Whitlock, 2014). Similarly,

loci identified through GEA methods often include false positives and may not reliably indicate

local adaptation (Nadeau et al., 2016). In some cases, GEA analyses can be paired with common

garden experiments or genome-wide association studies (GWAS) to help verify which loci are

associated with climate adaptation (e.g. Faske et al., 2021; Aitken et al., 2024).

We determined the number of studies in our review that identified adaptive loci and

examined whether they used F_{st} outlier analyses or GEA analyses, as these were the two

primary approaches used. In total, 59 studies identified adaptive loci, and of those studies, 22

used GEA approaches and 37 used F_{st} outlier analyses (Fig. 5A). Across both methods, a greater

proportion of studies recommended increasing the number of ESUs (GEA: 77%, F_{st} outlier: 65%;

Fig. 5A). However, even among studies that did not identify adaptive loci, the largest proportion

still suggested an increase in ESUs (66%; Fig. 5A). Interestingly, studies that used GEA

approaches only recommended either increasing the number of ESUs or maintaining the

current number, none suggested a decrease. These results may reflect greater confidence in the

adaptive potential of loci identified through GEA methods, as associations with environmental

conditions can strengthen the inference that these loci are truly adaptive (Hoban et al., 2016).

We also examined whether the way in which adaptive loci were used in a study influenced

conclusions about the number of ESUs suggested. We categorized studies with adaptive loci as

those that simply identified the presence of adaptive loci, and those that incorporated these

loci into downstream analyses that assessed population structure based on adaptive variation.

Among studies using GEA approaches, there was roughly an even split between those that only

identified loci and those that used them as a dataset (n=10 and n=12, respectively; Fig. 5B).

However, studies that additionally used adaptive loci as a dataset were more likely to suggest no

change in the number of ESUs (Fig. 5B). In contrast, for F_{st} outliers there were more studies that

only identified adaptive loci (n=25) than used them in downstream analyses (n=10) and these

studies were more likely to suggest an increase in the number of ESUs (Fig. 5C). This pattern

may suggest that identifying adaptive loci alone, without additionally examining their

population structure, can inflate the number of ESUs identified, possibly due to the high rate of

false positives associated with these methods (Bierne et al., 2013; Nadeau et al., 2016).

However, further analyzing adaptive loci for population structure may mitigate this effect.

Ultimately, while loci identified through F_{st} outliers and GEA approaches are a useful proxy for

local adaptation, their interpretation should be approached with caution. Additional analyses

may be useful to confirm a group's evolutionary significance beyond identifying adaptive loci, as

these loci do not always provide reliable evidence for adaptive potential (Bierne et al., 2013;

Chhina et al., 2024) and thus for identifying ESUs.

Figure 5. (A) The number of studies that determined adaptive loci using genotype-

environment association (GEA; n=22) or F_{ST} outlier analyses (n=37), as well as those that did

not identify adaptive loci (n=102). Bar colors indicate the number of studies in each category

that recommended an increase, decrease, or no change in the number of ESUs. Among

studies that identified adaptive loci, they were further categorized as "dataset" if the loci

were used in downstream analyses, or as "identified" if the loci were only reported but not

used in further analyses. (B) Results for studies using GEA approaches; (C) Results for studies

using F_{ST} outlier analyses.

Definitions and criteria used to define ESUs

One significant challenge we found in our review was the inconsistency in how ESUs are

defined. A few definitions were more commonly cited in the empirical studies we reviewed, and

we highlight these definitions in Table 1. Although ESUs are broadly recognized as being both

genetically discrete and evolutionary significant, multiple definitions are available, each offering

a different perspective on how to identify them (Table 1; also see reviews by Coates et al., 2018;

Turbek et al., 2023). However, studies often reference multiple definitions simultaneously,

resulting in merging the separate criteria of these definitions. In addition, ESU definitions often

have subjective language, and this has been suggested to limit their practical application to real

world situations (Fraser and Bernatchez, 2001). This ambiguity appears to be a persistent issue

in the current studies we reviewed.

Table 1. Definitions of Evolutionary Significant Units (ESUs) from previous publications. This is not an exhaustive list but rather shows the reference and the corresponding definitions that were cited most frequently by the 154 publications included in this review.

Reference	Definition	Number of times cited*
Ryder, 1986	"a subset of the more inclusive entity species, which possess genetic attributes significant for the present and future generations of the species in question"	16
Waples, 1991	"A population must satisfy two criteria to be considered an ESU: (1) it must be substantially reproductively isolated from other conspecific	8

	population units, and (2) it must represent an important component in the evolutionary legacy of the species."	
Moritz, 1994	"ESUs should be (1) reciprocally monophyletic for mtDNA alleles and (2) show significant divergence of allele frequencies at nuclear loci."	29
Crandall et al., 2000	"...suggest that the terminology [ESU] is abandoned and replaced with a more holistic concept of species, consisting of populations with varying levels of gene flow evolving through drift and natural selection"	8
Fraser and Bernatchez, 2001	"A lineage demonstrating highly restricted gene flow from other such lineages within the higher organizational level (lineage) of the species"	12
Funk et al., 2012	"Major, intraspecific units that (1) have been historically isolated from each other and (2) that likely have important adaptive differences among them"	11
Committee on the Status of Endangered Wildlife in Canada (Designatable Units: COSEWIC, 2023)	"A unit below the level of a taxonomic species if it has attributes that make it both (1) discrete, and (2) evolutionary significant, where discrete means that there is currently very little transmission of heritable information from other such units and evolutionarily significant means that the unit harbors heritable adaptive traits or an evolutionary history not found elsewhere in Canada."	7
Other citations	NA	20
Not reported	NA	103

*This column sums to more than the total number of studies due to several studies citing multiple references

We found that only 36% of studies explicitly stated how they defined an ESU, including

providing at least one reference for the given definition (Fig. 6A; Table 1). This raises concerns

because of the multiple, and sometimes conflicting definitions of what constitutes an ESU.

Interestingly, we found that studies that provided a definition were just as likely as those that

did not to suggest an increase in the number of recognized ESUs (69% and 66%, respectively),

and this outcome was more common than suggesting a decrease in the number of ESUs or no

change (Fig. 6A). Additionally, in some cases, the definition of ESU cited appeared to influence

whether a study recommended increasing or decreasing the number of ESUs (Fig. 6B). For

example, studies citing Waples (1991), more often suggested no change in the number of ESUs,

a pattern not observed with any other reference (Fig. 6B). In contrast, studies citing Moritz

(1994) and Funk et al. (2012) were proportionally more likely than those citing any other

definition to recommend an increase in the number of ESUs, rather than a decrease or no

change. Providing a clear definition for how an ESU is being considered is essential not only for

transparency, but also for having consistent and meaningful comparisons across studies.

Without providing a definition, it is unclear what evidence is being used to separate populations

into distinct ESUs, or alternatively, justify grouping them together.

Among the studies that provided a definition, the few key sources that were commonly

referenced (Table 1; Fig. 6B) differed in the main criteria required to determine separate ESUs.

We found that conclusions about increasing or decreasing the number of ESUs were often made

independently of the criteria outlined in the cited definition. For instance, 38% of studies citing

Waples (1991) and 33% citing Fraser and Bernatchez (2001) did not assess gene flow (Fig. S6),

despite the role of gene flow being specifically stated in those definitions. Similarly, 66% of

studies referencing Moritz (1994) did not include analyses with mtDNA (Table S2), even though

the definition clearly states the requirement of mtDNA to show reciprocal monophyletic groups.

This suggests that conclusions are often made independently of the criteria applied in an ESU

definition, raising concerns about the consistency and comparability of ESU identification across

studies, taxa, and regions.

Figure 6. (A) The number of studies that provided a written definition of how they

recognized an evolutionary significant unit, accompanied by a cited reference for that

definition (n=58), and the number of studies that did not provide a definition (n=103). Bar

colors indicate the study conclusion on whether they recommended an increase, decrease,

or no change in the number of evolutionary significant units. (B) The most frequently cited

references to define ESUs among the 161 data entries in our review, and the number of

studies that suggested an increase, decrease, or no change in the number of ESUs.

Changes in the number and delineation of ESUs

The studies in our review more frequently suggested increasing the number of ESUs

than maintaining the current number or decreasing them (Fig. 7). Specifically, 67% of the

studies suggested an increase in ESUs, 22% suggested no change, and only 11% proposed a

decrease (Fig. 7). We found this trend to be consistent regardless of sequencing type, the

downstream methods used to analyze the genomic data, or the way ESU were defined. Thus,

these findings indicate that genomic data is generally leading to the identification of more ESUs

within species, despite the many differences across studies.

Figure 7. The overall conclusions on the suggested number of evolutionary significant

units (increase, decrease, or no change the number of conservation units) across the 161

data entries included in this literature review.

An important finding is the inconsistency in how studies determined whether to increase

the number of ESUs. One major source of this inconsistency may be due to the lack of a

common threshold to determine if groups are different enough to be considered as separate

ESUs. For example, F_{ST} values are one of the most widely used statistics in population genetics,

but this statistic may give highly different estimates of genetic differences based on the number

of individuals analyzed, the number of geographic sampling locations, and the type of genetic

marker assessed (Holsinger and Weir, 2009). Implementing one common threshold may be

extremely difficult, and a discussion on how to do this is long overdue (e.g. also recommended

by Funk et al., 2012). Second, there is no unifying workflow available to help make these

decisions in a reliable and consistent way. Notably, there are a couple of recent workflows that

have been published to help guide the analyses for identifying separate conservation units. For

example, Turbek et al., (2023) has developed a decision tree for species exhibiting patterns of

Isolation-by-distance, and Lehnert et al., (2023) aimed to delineate Designatable Units in

Canada (legal units often considered to have similar concepts to ESUs). Most likely due to their

recent publication dates at the time of this literature review, these decision trees have yet to be

widely used (but see Miller et al., 2024 which follows the approach of Lehnert et al., 2023).

However, these workflows do not fully address the broader concept of evolutionary significance

past identifying adaptive loci. Thus, a more detailed workflow to help guide studies to assess

genetic discreteness and fully evaluate evolutionary significance is still needed to provide much-

needed consistency for identifying ESUs.

Conclusion and Recommendations

The vast amount of information that genomic data provides greatly improves our ability

to inform the identification of ESUs (Funk et al., 2012; Coates et al., 2018; Funk et al., 2019;

Turbek et al., 2023). Genomic data facilitates a deeper understanding of the genetic differences

within a species and can reliably uncover their evolutionary history, both of which are essential

to accurately identify intraspecific ESUs. In this study, we conducted a comprehensive literature

review to explore how genomic data has been used to inform ESUs globally. We found that key

details about the genomic data being used, such as sequencing depth, are frequently not

reported. Additionally, we found that there is a lack of consistency in the methods used to

analyze genomic data or how ESUs are defined. Despite these inconsistencies, the majority of

studies recommended increasing the number of ESUs. Here we identify some gaps in how

genomic data is currently being used to identify ESUs and highlight some recommendations that

could be used to improve future studies.

We strongly recommend clear and consistent reporting of genomic data, study details, and

how ESUs are being defined. We found that an overarching problem when using genomic data

to inform ESUs is the lack of a unifying definition and an accompanying unifying framework, and

that future work is needed to agree upon a single definition of what constitutes an ESU which

can be used across all taxa. This lack of unification is causing inconsistencies in the analyses

being used, making it challenging to have comparable conclusions across different studies.

While some studies relied solely on measures of genetic discreteness (e.g. PCA), we recommend

that additional lines of evidence, such as demographic reconstructions, be incorporated to

assess evolutionary significance (see Box 2 for examples). Genetic structure identified through

analyses based on clustering and ancestry-based frequencies can be highly influenced by recent

isolation due to habitat fragmentation, genetic drift, or recent inbreeding, all of which can be

due to anthropogenic causes (Schlaepfer et al., 2018), and not a result of significant differences

evolved over time. Similarly, analyses used to detect adaptive loci, such as F_{ST} outliers, must be

interpreted cautiously, as they can be influenced by patterns other than adaptation (Bierne et

al., 2013; Lotterhos and Whitlock, 2014; Nadeau et al., 2016). Importantly, we suggest that

identifying adaptive loci is not always necessary if divergent evolutionary histories can be

demonstrated through other analyses.

We advocate for identifying ESUs in a manner which preserves the evolutionary legacy of a

species. We therefore recommend the use of phylogenomic and demographic history

reconstruction as important analyses to use when identifying ESUs. Ideally, phylogeny

reconstruction using genome-wide data should be used to provide a powerful and reliable

inference of evolutionary history. With the increase of accessibility to WGS (Fuentes-Pardo and

Ruzzante, 2017; Lou et al., 2021), this is a practical solution and could provide the much-needed

consistency for identifying ESUs across different taxa. Beyond improving phylogenetic

resolution, by using whole genomes we can improve the ability to test for local adaptation

beyond F_{st} outliers and GEA approaches. More sophisticated analyses for this need to be

developed and incorporated into workflows for ESUs enabling studies to fully utilize the large

amount of information that can be gained when using whole genomes. However, we recognize

that WGS is still not feasible for all systems, particularly those with large or complex genomes.

In such cases, RRS has been shown to provide reliable results to infer the demographic history

of a species (Elleouet and Aitken, 2018) and could be used as an alternative.

Despite these suggestions for improvement, significant progress has been made over the

past decade to better utilize the power of genomics to inform the identification of ESUs

(Flanagan, et al., 2018; Funk et al., 2019; Allendorf et al., 2022). Genomic data, including WGS,

is more accessible than ever before (Lou et al., 2021; Schmidt et al., 2024) and can provide

valuable insights into genome-wide variation inclusive of neutral, adaptive, and deleterious

variation that may be critical for current and future conservation management (Flanagan et al.,

2018; Chhina et al., 2024; Schmidt et al., 2024). Moving forward, it is essential that both

researchers and practitioners not only understand how to assess genomic data but also how to

apply it effectively for identifying ESUs. By fully embracing the power of genomics with more

standardization and thoughtful interpretation, we can improve species conservation and ensure

that decisions regarding the identification of ESUs are made accurately and informatively for

evolutionary relevance.

Literature cited

Aguirre-Liguori JA, Luna-Sánchez JA, Gasca-Pineda J, Eguiarte LE (2020). Evaluation of the Minimum Sampling Design for Population Genomic and Microsatellite Studies: An Analysis Based on Wild Maize. *Front Genet* **11**: 870.

Aitken SN, Jordan R, Tumas HR (2024). Conserving Evolutionary Potential: Combining Landscape Genomics with Established Methods to Inform Plant Conservation. *Annual Review of Plant Biology* **75**: 707–736.

Aitken SN, Whitlock MC (2013). Assisted Gene Flow to Facilitate Local Adaptation to Climate Change. *Annu Rev Ecol Evol Syst* **44**: 367–388.

Allendorf FW (2017). Genetics and the conservation of natural populations: allozymes to genomes. *Mol Ecol* **26**: 420–430.

Allendorf FW, Funk C, Aitken SN, Byrne M, Luikart G (2022). Conservation and the genomics of populations. 3rd ed. Oxford (UK): Oxford University Press.

Barbosa S, Hendricks SA, Funk WC, Rajora OP, Hohenlohe PA (2020). Wildlife Population Genomics: Applications and Approaches. In: Hohenlohe PA, Rajora OP (eds) *Population Genomics: Wildlife*, Population Genomics. Springer International Publishing: Cham, pp 3–59.

Barbosa S, Mestre F, White TA, Paupério J, Alves PC, Searle JB (2018). Integrative approaches to guide conservation decisions: Using genomics to define conservation units and functional corridors. *Mol Ecol* **27**: 3452–3465.

Bierne N, Roze D, Welch JJ (2013). Pervasive selection or is it…? why are F_{ST} outliers sometimes so frequent? *Molecular Ecology* **22**: 2061–2064.

Chambers EA, Hillis DM (2020). The Multispecies Coalescent Over-Splits Species in the Case of Geographically Widespread Taxa (R Glor, Ed.). *Systematic Biology* **69**: 184–193.

Chhina AK, Abhari N, Mooers A, Lewthwaite JMM (2024). Linking the spatial and genomic structure of adaptive potential for conservation management: a review. *Genome* **67**: 403–423.

Coates DJ, Byrne M, Moritz C (2018). Genetic Diversity and Conservation Units: Dealing With the Species-Population Continuum in the Age of Genomics. *Front Ecol Evol* **6**: 165.

Committee on the Status of Endangered Wildlife in Canada, November (2023). COSEWIC guidelines for recognizing designatable units. COSEWIC. https://cosewic.ca/index.php/en/reports/preparing-status-reports/guidelines-recognizing-designatable-units.html

Crandall KA, Bininda-Emonds ORP, Mace GM, Wayne RK (2000). Considering evolutionary processes in conservation biology. *Trends in Ecology & Evolution* **15**: 290–295.

Dokan K, Kawamura S, Teshima KM (2021). Effects of single nucleotide polymorphism ascertainment on population structure inferences (Y Kim, Ed.). *G3 Genes|Genomes|Genetics* **11**: jkab128.

Elhaik E (2022). Principal Component Analyses (PCA)-based findings in population genetic studies are highly biased and must be reevaluated. *Sci Rep* **12**: 14683.

Ellegren H (2014). Genome sequencing and population genomics in non-model organisms. *Trends in Ecology & Evolution* **29**: 51–63.

Elleouet JS, Aitken SN (2018). Exploring Approximate Bayesian Computation for inferring recent demographic history with genomic markers in nonmodel species. *Molecular Ecology Resources* **18**: 525–540.

Faske TM, Agneray AC, Jahner JP, Sheta LM, Leger EA, Parchman TL (2021). Genomic and common garden approaches yield complementary results for quantifying environmental drivers of local adaptation in rubber rabbitbrush, a foundational Great Basin shrub. *Evolutionary Applications* **14**: 2881–2900.

Flanagan SP, Forester BR, Latch EK, Aitken SN, Hoban S (2018). Guidelines for planning genomic assessment and monitoring of locally adaptive variation to inform species conservation. *Evol Appl* **11**: 1035–1052.

Forester BR, Lasky JR, Wagner HH, Urban DL (2018). Comparing methods for detecting multilocus adaptation with multivariate genotype–environment associations. *Molecular Ecology* **27**: 2215–2233.

Forester BR, Murphy MA, Mellison C, Petersen J, Pilliod DS, Van Horne R, *et al.* (2022). Genomics-informed delineation of conservation units in a desert amphibian. *Molecular Ecology*: 1–21.

Fraser DJ, Bernatchez L (2001). Adaptive evolutionary conservation: towards a unified concept for defining conservation units. *Mol Ecol* **10**: 2741–2752.

Fuentes-Pardo A, Ruzzante D (2017). Whole-genome sequencing approaches for conservation biology: Advantages, limitations and practical recommendations. *Molecular Ecology* **26**: 5369–5406.

Funk WC, Forester BR, Converse SJ, Darst C, Morey S (2019). Improving conservation policy with genomics: a guide to integrating adaptive potential into U.S. Endangered Species Act decisions for conservation practitioners and geneticists. *Conserv Genet* **20**: 115–134.

Funk WC, McKay JK, Hohenlohe PA, Allendorf FW (2012). Harnessing genomics for delineating conservation units. *Trends in Ecology & Evolution* **27**: 489–496.

Gallego-García N, Vargas-Ramírez M, Shaffer HB (2023). The importance of cryptic diversity in the conservation of wide-ranging species: The red-footed tortoise Chelonoidis carbonarius in Colombia. *Molecular Ecology* **32**: 4531–4545.

Garrigan D, Hedrick PW (2003). Perspective: Detecting adaptive molecular polymorphism:
Lessions from the MHC. *Evolution* **57**: 1707–1722.

Ghildiyal K, Nayak SS, Rajawat D, Sharma A, Chhotaray S, Bhushan B, *et al.* (2023). Genomic
insights into the conservation of wild and domestic animal diversity: A review. *Gene* **886**:
147719.

Hemstrom W, Grummer JA, Luikart G, Christie MR (2024). Next-generation data filtering in the
genomics era. *Nat Rev Genet*.

Hoban S, Archer FI, Bertola LD, Bragg JG, Breed MF, Bruford MW, *et al.* (2022). Global genetic
diversity status and trends: towards a suite of Essential Biodiversity Variables (EBVs) for
genetic composition. *Biological Reviews* **97**: 1511–1538.

Hoban S, Kelley JL, Lotterhos KE, Antolin MF, Bradburd G, Lowry DB, *et al.* (2016). Finding the
Genomic Basis of Local Adaptation: Pitfalls, Practical Solutions, and Future Directions. *The*
American Naturalist **188**: 379–397.

Hohenlohe PA, Funk WC, Rajora OP (2021). Population genomics for wildlife conservation and
management. *Mol Ecol* **30**: 62–82.

Holsinger KE, Weir BS (2009). Genetics in geographically structured populations: defining,
estimating and interpreting FST. *Nat Rev Genet* **10**: 639–650.

Jensen EL, Gaughran SJ, Garrick RC, Russello MA, Caccone A (2021). Demographic history and
patterns of molecular evolution from whole genome sequencing in the radiation of
Galapagos giant tortoises. *Molecular Ecology* **30**: 6325–6339.

Jensen EL, Govindarajulu P, Russello MA (2014). When the shoe doesn't fit: applying
conservation unit concepts to western painted turtles at their northern periphery. *Conserv*
Genet **15**: 261–274.

Jorge RF, Lima AP, Stow AJ (2022). Selection and localised genetic structure in the threatened
Manauense Harlequin Frog (Bufonidae: Atelopus manauensis). *Conservation Genetics* **23**:
559–574.

Kanthaswamy S, Kurushima JD, Smith DG (2006). Inferring Pongo conservation units: a
perspective based on microsatellite and mitochondrial DNA analyses. *Primates* **47**: 310–
321.

Lawson DJ, Van Dorp L, Falush D (2018). A tutorial on how not to over-interpret STRUCTURE
and ADMIXTURE bar plots. *Nat Commun* **9**: 3258.

Lehnert SJ, Bradbury IR, Wringe BF, Van Wyngaarden M, Bentzen P (2023). Multifaceted
framework for defining conservation units: An example from Atlantic salmon (*Salmo salar*)
in Canada. *Evolutionary Applications* **16**: 1568–1585.

Lenormand T (2002). Gene flow and the limits to natural selection. *Trends in Ecology &*
Evolution **17**: 183–189.

Lotterhos KE, Whitlock MC (2014). Evaluation of demographic history and neutral
parameterization on the performance of F_{ST} outlier tests. *Molecular Ecology* **23**: 2178–
2192.

Lou RN, Jacobs A, Wilder AP, Therkildsen NO (2021). A beginners guide to low-coverage whole
genome sequencing for population genomics. *Molecular Ecology* **30**: 5966–5993.

Miller CV, Bossu CM, Sarraco JF, Toews DPL, Rushing CS, Roberto-Charron A, *et al.* (2024).
Genomics-informed conservation units reveal spatial variation in climate vulnerability in a
migratory bird. *Molecular Ecology* **33**.

Moritz C (1994). Defining 'Evolutionarily Significant Units' for conservation. *Trends in Ecology &*
Evolution **9**: 373–375.

Nadeau S, Godbout J, Lamothe M, Gros-Louis M-C, Isabel N, Ritland K (2015). Contrasting
patterns of genetic diversity across the ranges of *Pinus monticola* and *P. strobus* : A
comparison between eastern and western North American postglacial colonization
histories. *American Journal of Botany* **102**: 1342–1355.

O'Leary SJ, Puritz JB, Willis SC, Hollenbeck CM, Portnoy DS (2018). These aren't the loci you'e
looking for: Principles of effective SNP filtering for molecular ecologists. *Molecular Ecology*
27: 3193–3206.

Pederson T (2024). _patchwork: The Composer of Plots_. R package version 1.3.0,
<https://CRAN.R-project.org/package=patchwork>.

Peterson BK, Weber JN, Kay EH, Fisher HS, Hoekstra HE (2012). Double Digest RADseq: An
Inexpensive Method for De Novo SNP Discovery and Genotyping in Model and Non-Model
Species (L Orlando, Ed.). *PLoS ONE* **7**: e37135.

R Core Team (2024). *R: A language and environment for statistical computing*. R Foundation for
Statistical Computing, Vienna, Austria.

Ryder OA (1986). Species conservation and systematics: the dilemma of subspecies. Trends in
Ecology and Evolution **1**: 9-10.

Schlaepfer DR, Braschler B, Rusterholz H, Baur B (2018). Genetic effects of anthropogenic
habitat fragmentation on remnant animal and plant populations: a meta-analysis.
Ecosphere **9**: e02488.

Schmidt TL, Thia JA, Hoffmann AA (2024). How Can Genomics Help or Hinder Wildlife
Conservation? *Annu Rev Anim Biosci* **12**: annurev-animal-021022-051810.

Suyama Y, Matsuki Y (2015). MIG-seq: an effective PCR-based method for genome-wide single-
nucleotide polymorphism genotyping using the next-generation sequencing platform. *Sci*
Rep **5**: 16963.

Taylor RS, Manseau M, Keobouasone S, Liu P, Mastromonaco G, Solmundson K, *et al.* (2024b). High genetic load without purging in caribou, a diverse species at risk. *Current Biology* **34**: 1234-1246.e7.

Taylor RS, Manseau M, Wilson PJ (2024a). Delineating conservation units should be independent of effective population size. *Trends in Ecology & Evolution*: S0169534723003270.

Tigano A, Friesen VL (2016). Genomics of local adaptation with gene flow. *Molecular Ecology* **25**: 2144–2164.

Turbek SP, Funk WC, Ruegg KC (2023). Where to draw the line? Expanding the delineation of conservation units to highly mobile taxa (W Murphy, Ed.). *Journal of Heredity* **114**: 300–311.

U.S. Fish and Wildlife Service & National Marine Fisheries Service (1996). Policy regarding the recognition of distinct vertebrate population segments under the Endangered Species Act. *Federal Register*, **61**: 4722–4725. https://www.federalregister.gov/documents/1996/02/07/96-2639/policy-regarding-the-recognition-of-distinct-vertebrate-population-segments-under-the-endangered

Van de Schoot R, de Bruin J, Schram R, Zahedi P, de Boer J, Weijdema F et al. (2021). ASReview: open-source machine learning software for systematic reviews. *Nat Mach Intell* 3: 125–126.

Waples RS (1991). Pacific Salmon, Oncorhynchus spp., and the Definition of 'Species' Under the Endangered Species Act. *Marine Fisheries Review*.

Waples RS (2024). The N_e / N ratio in applied conservation. *Evolutionary Applications* **17**: e13695.

Waples RS (2025). The Idiot's Guide to Effective Population Size. *Molecular Ecology*: e17670.

Waples RS, Lindley ST (2018). Genomics and conservation units: The genetic basis of adult migration timing in Pacific salmonids. *Evolutionary Applications* **11**: 1518–1526.

Wickham (2016). ggplot2: Elegant Graphics for Data Analysis. Springer-Verlag, New York.

Whitlock MC, Lotterhos KE (2015). Reliable Detection of Loci Responsible for Local Adaptation: Inference of a Null Model through Trimming the Distribution of F_{ST}. *The American Naturalist* **186**: S24–S36.

Conflicts of Interest

The authors declare no conflicts of interest.

Data Availability Statement

The authors have nothing to report.

Box 1

Genomic data are DNA sequences or molecular markers that encompass a large and

representative set of regions across the genome to provide high resolution information

(Schmidt et al., 2024). Genomic data can allow for the identification of the structure and/or

function of genetic variation across the genome.

Genomic data can be generated by the following sequencing methods which are grouped into

several categories:

1. **SNP-based genotyping** – Refers to techniques that have a set of predefined and
targeted probes with specific position of single nucleotide polymorphisms (SNPs). These
SNPs target variants that are based on relevance to specific individuals/populations,
adaptive loci, or particular traits. We have grouped SNP chips, SNP array, SNP assays and
SNP panels together because these are all types of targeted SNP-based genotyping.

2. **Reduced representation sequencing (RRS)** – Is an approach to generate genome-wide
sequencing data by sequencing only digested fragments of the genome. Fragments are
from across the entire genome, but only a fraction of the genome is sequenced. The
following sequencing types are RRS methods:

   - ***Double digest restriction-site associated DNA sequencing (ddRAD)*** – Uses two
   restriction enzymes to ensure more uniform fragments of the whole genome by
   having two cut sites (Peterson et al., 2012). We have considered **DArTseq** methods as
   ddRAD because they both deploy double enzyme digestion across the genome.
   DArTseq uses a different system for commercial services.

- **_Genotype-by-sequencing (GBS)_** – Single enzyme digestion of the genome that uses
PCR for fragment size selection.

- **_Restriction-site associated DNA sequencing (RADseq)_** – uses one enzyme to perform
a single digestion of the whole genome and then randomly interrupts the digested
fragments.

- **_Multiplexed Inter-simple sequence repeat genotype by sequencing (MIG-seq_**) – This
method does not use restriction-enzyme digestion but rather PCR-based markers
that amplify a large number of genome-wide regions without prior genetic
information (Suyama and Matsuki, 2015).

3. **Whole genome sequencing** – sequencing the entire genome of an organism to get a
comprehensive view of the genome. There are two types of whole genome sequencing
that differ by the depth of the sequencing:

- **_High coverage whole genome sequencing_** – Whole genome sequencing that
typically has coverage high enough to allow genotypes to be accurately called.

- **_Low coverage whole genome sequencing_** – Whole genome sequencing resulting in a
small number of reads covering the genomic loci (e.g. <10X coverage). Genotype
likelihood-based methods are used with low coverage data.

4. **Targeted gene sequencing** – This sequencing approach focuses on specific genes or
genomic regions of interest, rather than sequencing the entire genome.

- **_Major histocompatibility complex genes (MHC)_** – These functional genes play a vital
role in an organism's immune system and are highly polymorphic. MHC diversity can

801   be used as an indicator of a population's adaptive ability (Garrigan and Hedrick,

802   2003).

803   • *Exome sequencing* – Exons are protein-coding regions of the genome. Exome

804   sequencing is a targeted sequencing approach to capture variation within these

805   protein-coding regions.

806   5. **Pooled sequencing (pool-seq)** – Multiple individual DNA samples are pooled together

807   and then sequenced in a single run. Pooled sequencing typically has a high depth of

808   coverage (e.g. >50X) and can be cost effective (Fuentes-Pardo and Ruzzante, 2021).

809   However individual genotypes are undistinguishable due to the process of pooling the

810   DNA (Fuentes-Pardo and Ruzzante, 2021).

811   6. **Transcriptomics** – Is the characterization of RNA transcripts through RNA sequencing

812   that is typically used to understand gene expression and regulation.

Box 2

Case study 1: The importance of cryptic diversity in the conservation of wide-ranging species:

The red-footed tortoise (*Chelonoidis carbonarius*) in Columbia

Gallego-Garcia et al., 2023

The red-footed tortoise (*Chelonoidis carbonarius*) is a widespread, vulnerable species found in

Central America, South America, and several Caribbean islands. Gallego-Garcia et al. (2023)

sequenced 233 individuals from 25 sites across the species' range in Colombia using triple

enzyme restriction-site associated DNA sequencing (3RAD), generating a dataset of 30,327

SNPs. This summary focuses on the analyses conducted across all individuals to identify

Evolutionarily Significant Units (ESUs), although additional analyses were performed within each

identified ESU to determine additional smaller management units. To assess genetic

discreteness, the authors used Principal Component Analysis (PCA), STRUCTURE, and F_{ST} values.

All three methods supported the presence of distinct eastern and western lineages (i.e. two

ESUs). The lack of admixture in the STRUCTURE plot and high F_{ST} values (~0.42) were used to

infer no gene flow between the two groups. To assess evolutionary significance, a phylogenetic

tree was generated and revealed a basal split between the lineages, supporting deep

divergence and a lack of ancestral gene flow. Additionally, effective population sizes and

divergence times were estimated for both lineages to further support their evolutionary

significance. Finally, putatively adaptive loci were identified through genotype-environment

associations (GEA) analysis using a redundancy analysis and an F_{ST} outlier analysis using

PCAdapt. Overall, Gallego-Garcia et al., (2023) suggest that the eastern and western lineages

should be recognized as two separate ESUs based on analyses that support both genetic

discreteness and evolutionary significance, thereby increasing the number of ESUs for the

species.

Case study 2: Selection and localised genetic structure in the threatened Manauense Harlequin

frog (Bufonidae: *Atelopus manauensis*)

Jorge et al., 2022

The Manauense Harlequin frog (*Atelopus manauensis*) is a threaten, stream-breeding

amphibian with a narrow distribution in the Amazon. Jorge et al., (2022) sequenced 94

individual males from 21 streams using a modified protocol of double digest restriction-site

associated DNA sequencing (ddRAD), generating a dataset of 3,859 SNPs. To identify putatively

adaptive loci, the authors used F_{ST} outlier analyses (PCAdapt and ARLEQUIN) and a genotype-

environment association (GEA) approach using Latent Factor Mixed Models (LFMM). The GEA

analysis detected significant associations between genetic variation and environmental

variables, suggesting local adaptation. A neutral dataset of 3,121 SNPs was then created by

removing the adaptive loci identified through the three analyses, as well as loci not in Hardy-

Weinberg equilibrium. Genetic discreteness was assessed using the neutral dataset through

Discriminant Analysis of Principal Components (DAPC), sparse non-negative matrix factorization

(SNMf), STRUCTURE, and F_{ST} values. All four analyses identified six discrete genetic clusters, with

high F_{ST} values (0.3-0.5) between them. To assess evolutionary significance, in addition to

identifying the adaptive loci, the authors constructed a phylogenetic tree, and estimated

divergence times, which ranged from approximately 40,000 to 2,000 years ago depending on

the cluster. Finally, a stairway plot indicated population declines around 100,000 years ago, and

current estimated effective population sizes per group ranged from 20 to 100 individuals.

Overall, Jorge et al., (2022) suggest an increase in the number of ESUs, recognizing a total of six

separate ESUs that show both genetic discreteness and evolutionary significance.