Dynamic hotspots in the Uba7 UFD direct UbcH8 recognition

Çağdaş Dağ[1,2,3,4,*], Mahil Lambert[1,4], Alp E. Kazar[1], Kerem Kahraman[1], Oktay Göcenler[1],

Woonghee Lee[5], Cansu D. Tozkoparan Ceylan[1], Frank Löhr[4], Jin Gon Shim[6], Arthur L. Haas[7],

Volker Dötsch[5], Joshua J. Ziarek[6], Emine Sonay Elgin[3,*]

[1]Nanofabrication and Nanocharacterization Center for Scientific and Technological Advanced Research (n[2]STAR), Koç University, İstanbul, Turkiye

[2]Koç University Isbank Center for Infectious Diseases (KUISCID), Koç University, Istanbul, Turkiye

[3] Department of Chemistry, College of Sciences, Muğla Sıtkı Koçman University, Muğla, 48000, Turkiye

[4]Centre for Biomolecular Magnetic Resonance, Institute for Biophysical Chemistry, Goethe-University of Frankfurt/Main, Germany

[5]Department of Chemistry, University of Colorado Denver, Denver, CO 80204, USA

[6]Department of Pharmacology, Feinberg School of Medicine, Northwestern University, 320 East Superior Avenue, Chicago, IL, 460611, USA

[7]Department of Biochemistry and Molecular Biology, LSUHSC-School of Medicine, 1901 Perdido Street, New Orleans, LA, 70112, USA

*Corresponding author: cdag@ku.edu.tr selgin@mu.edu.tr

Abstract

The ISGylation pathway, a key post-translational modification, plays a pivotal role in the innate

immune response by covalently attaching ISG15 to target proteins. This cascade involves a series

of enzymatic steps, including activation by the E1 enzyme Uba7, conjugation by the E2 enzyme

UbcH8, and ligation by an E3 ligase. Central to this process is the ubiquitin-fold domain (UFD)

of Uba7, which facilitates the transfer of ISG15 to UbcH8. However, the structural and mechanistic

details of this interaction remain poorly understood. In this work, we present the solution NMR

structure and functional analysis of the ubiquitin-fold domain of human Uba7, the E1 enzyme in

the ISGylation cascade. Through detailed NMR titration experiments and mutational studies, we

mapped the interaction surface of Uba7-UFD with UbcH8, identifying key residues and their

contributions to the E1-E2 interaction. We found that Uba7-UFD is flexible in its free form; this

flexibility is conserved across ubiquitin-like systems but shows a regional shift that may play an

important role in correct E2 and Ubl selection. Chemical shift perturbation and mutational analysis

further demonstrate the importance of specific residues, particularly Cys996, in maintaining UFD's

structural integrity and binding capacity. Additionally, mutations designed to alter the flexibility

and length of the loop region between UFD and UbcH8 show significant effects on binding,

indicating that these regions are crucial for efficient E2 recruitment. These findings provide new

insights into the mechanistic basis of E2 enzyme selection in ISGylation and underscore the

functional relevance of dynamic structural transitions in E1-E2 complex formation.

Statement of Significance

ISGylation is a key ubiquitin-like modification pathway essential for antiviral defense, immune

regulation, and protein quality control. However, the molecular principles that govern

communication between the E1 and E2 enzymes in this pathway remain poorly defined. Here, we

present the first NMR structure of the human Uba7 ubiquitin-fold domain (UFD) and its interaction

with UbcH8. Our findings reveal how structural flexibility within the UFD is crucial for specific

and efficient E2 recognition, providing fundamental insight into the dynamic mechanism of

ISGylation and advancing our understanding of ubiquitin-like enzyme cascades relevant to cellular

regulation and disease.

Introduction

ISGylation is a post translational modification (PTM) that results in the binding of interferon-stimulated gene 15 (ISG15) to the target protein. ISG15 is a 15 kDa ubiquitin-like protein (Ubl) containing two ubiquitin-like domains connected by a short linker peptide (Haas et al., 1987). Expression of ISG15 is primarily induced by type I interferon (IFN α/β) signaling (Farrell et al., 1979), through interactions between IFN regulatory factors (IRFs) and interferon-stimulated response element (ISRE)-containing promoters (Sadler and Williams, 2018). Innate immune response on a molecular level is generated via host pattern-recognition receptors and expression of IFN-I through downstream signaling. In this regard, ISG15 expression coincides with initiation of the innate immune response which serves as the first line of defense against pathogens. After expression, ISG15 takes part in innate immune response either as an intracellular regulatory protein or through ISGylation of primarily pathogen proteins (Zhang et al., 2021). In the context of the innate immune response, effectiveness of ISGylation against pathogens varies between species. For instance, ISGylation in mice is shown to be a highly effective antiviral mechanism for inhibiting viral assembly slowing down pathogenesis (Durfee et al., 2010). ISGylation of viral proteins also interferes with localization, host-viral protein interactions, and protease activity thereby strengthening the antiviral response (Rahnefeld et al., 2014). Whereas in humans, inherited ISG15 deficiency only results in susceptibility to mycobacterial, not viral diseases indicating differing effectiveness against pathogens and possible differences in ISGylation pathway between species (Bogunovic et al., 2012).

Similar to ubiquitin, ISG15 is also covalently attached via its C-terminal LRLRGG motif to a lysine residue on the target protein through an analogous E1-E2-E3 enzymatic cascade (Narasimhan, Potter and Haas, 1996). ISGylation enzymatic cascade includes an activating E1

enzyme (Uba7/Ube1L), a conjugating E2 enzyme (UbcH8), and an E3 ligase (HERC5) which their

expression is simultaneously induced with ISG15 through ISRE containing elements. ISGylation

begins with an ATP dependent reaction of sequential adenylation at the C-terminal motif

"LRLRGG" generating an AMP-ISG15 intermediate which forms a thioester bond with the

catalytic cysteine of the E1 activation enzyme, Uba7. In this step, active site cysteine half domains,

the first catalytic cysteine half-domain (FCCH) and the second catalytic cysteine half-domain

(SSCH) of the E1 form a thioester bond with the adenylated C-terminal glycine of ISG15. The

ubiquitin folding domain (UFD) of Uba7 functions as the recognizing agent of the E2 enzyme and

ISG15 where ISG15 is transferred to the catalytic cysteine of the E2 conjugating enzyme, UbcH8,

through a transthioesterification reaction. E2 functions as a bridge between the Uba7 (E1), and

HERC5 (E3), enzymes containing catalytic domain interaction sites of both E1 and E3. In the

subsequent reaction, ISG15 is transferred to the catalytic cysteine of E3 through another

transthioesterification reaction then ligated to a lysine residue of the target protein in a nonspecific

manner (Haas, 2007). Despite its similarities, ISGylation differs considerably from ubiquitylation

in terms of variability and prevalence in the cell. Unlike its counterpart, ISGylation cascade

contains only one set of E1-E2 enzymes and therefore regulation of ISGylation is further

influenced through intrinsic interactions between enzymes of the cascade in addition to external

pathways regulating the ISGylation.

As the cornerstone of Ub/Ubl conjugation pathways, E1 enzymes are responsible for initiating the

activation of the corresponding Ub/Ubl along with its transfer to the relevant E2 enzymes. This

crucial function helps to ensure the accuracy and reliability of Ub/Ubl-mediated conjugation

pathways (Capili and Lima 2007, Streich and Haas, 2010) highlighting the crucial role that E1

enzymes play in maintaining the integrity of ubiquitylation pathways. Previous studies have

managed to identify crystal structures of Ub E1 along with E1s for Ub-like modifiers (Ubls) such

as SUMO and Nedd8 (Figure 1), both singularly and in complex states with their corresponding

Ubls (Olsen and Lima, 2013; Huang et al., 2005, Lois and Lima, 2005). Structures of the SUMO

E1 provide mechanistic insights into SUMO activation and E2 recruitment to E1. A discovery

from these studies was the revealing of the multidomain structure inherent to canonical E1s. The

domains include a pseudodimeric adenylation domain which plays a crucial role in Ubl activation,

a Cys domain which contains the catalytic cysteine residue and a ubiquitin-fold domain (UFD)

involved in the recruitment of E2 (Figure 1). To better contextualize the structural findings of the

present study, we first conducted a comparative analysis of E1 enzyme ubiquitin-fold domains

(UFDs) from several ubiquitin and ubiquitin-like (Ubl) modification systems (Figure 1). The aim

of this figure is to illustrate that in most available crystal structures of free (apo) E1 enzymes, the

E2-binding regions of the UFD domains lack interpretable electron density, indicating substantial

conformational flexibility in these segments. This absence of density across multiple E1–UFD

structures—including Uba1, Uba2, Uba3, and Uba6—suggests that these regions are inherently

disordered or mobile in solution until they engage their cognate E2 enzymes. Such disorder-to-

order transitions have been previously proposed to facilitate accurate E2 selection and dynamic

repositioning of the catalytic domains during transthiolation. Our NMR-derived structure of the

human Uba7-UFD (Figure 1K) provides complementary insight into this phenomenon by

capturing the domain in its solution state, where flexibility is preserved rather than averaged out

as in crystal lattices. In constructing Figure 1, we intentionally combined representative E1–UFD

structures, both apo and complexed with E2 or Ubl partners, to highlight how the flexible β-grasp

and loop regions adopt defined conformations upon complex formation.

Furthermore, these prior structural studies have unveiled the considerable conformational shifts

E1s undergo to execute their roles effectively. In light of conformational changes revealing the

UFD, a hidden E2 binding area on the E1 is exposed, assisting in facilitating contact between the

Ubl thioester and E2 (Olsen and Lima, 2013). Upon the Ubl thioester's transfer from E1 to E2, it

is proposed that the UFD reverts to its originally locked conformation. This switch aids in the

release of the E2-Ub thioester product through a spatial mechanism (Olsen and Lima, 2013).

Similarly, E1-E2 interaction during ISGylation marks a crucial regulatory step of the ISGylation

pathway however while considerable insights have been attained regarding the structural insights

into E1 activity, the specific process at the atomic level has yet to be determined of ISGylation

pathway. The importance of flexible regions and flexibility in protein-protein interactions has

become increasingly evident with the advancement of structural methods. In this context,

"disorder-to-order transition" in protein binding regions is being extensively studied using NMR

spectroscopy and plays a crucial role in elucidating protein-protein interactions (Jubb, Blundell

and Ascher, 2015). Specific mechanistic and interactive characteristics of ISGylation are still

poorly   understood.

In this study, we explore the NMR structure of Uba7-UFD, interaction dynamics and the critical

role of the disorder-to-order transition within the E1-E2 complex of the ISGylation cascade at the

residual level using NMR spectroscopy. The structural flexibility observed for Uba7-UFD in the

unbound state is consistent with a general feature of E1 enzymes, in which localized dynamic

elements are maintained within an otherwise well-folded UFD scaffold to support productive E2

engagement. Upon interaction with UbcH8, the relaxation profiles indicate a redistribution of

backbone dynamics within the UFD—most prominently around the highlighted β2/β4 surface—

suggesting that E2 binding modulates pre-existing motional properties rather than invoking a

global conformational switch. Notably, while flexibility at the UFD–E2 interface appears

conserved across ubiquitin-like activation systems, the specific segments exhibiting the strongest

dynamic signatures show a measurable regional shift between different E1–E2 pairs, providing a

plausible mechanism for fine-tuning E2 specificity and ensuring efficient ISG15 transfer. Through

detailed NMR titration experiments, we mapped the extensive interaction surfaces between Uba7-

UFD and UbcH8, identifying over 30 residues involved in this dynamic binding process. The study

also highlighted the significance of electrostatic complementarities between the interacting regions

of Uba7-UFD and UbcH8, which contribute to their stable complex formation. Furthermore,

mutational analysis underscored the importance of key residues, such as cysteine 996, in

maintaining the structural integrity and binding capacity of Uba7-UFD. These findings provide

novel insights into the conformational plasticity of E1 enzymes and their regulatory role in

ISGylation, paving the way for a deeper understanding of this post-translational modification

process at the molecular level.

Materials and Methods

Construction of Uba7-UFD expression plasmid

The DNA sequence coding for Uba7-UFD (amino acid residues 903-1011) was amplified from

pGEX-full length HsUba7 by polymerase chain reaction (PCR). Taq polymerase (Thermo) and

primers 5' Uba7-UFD-BglII (5'-GCGCAGATCTCAGACGTTCCATCACCTG-3') and 3' Uba7-

UFD-HindIII 5'-GCGCAAGCTTACAGCTCATAGTGCAGAGG-3' were used in the reaction.

Amplified PCR product was cleaned up using Omega PCR clean-up kit and digested with BglII

and HindIII while the pQE30-GBI expression vector was digested with BamHI and HindIII. The

digested PCR product and plasmid ligated with T4 DNA ligase. Accuracy of the cloned gene was

confirmed by DNA sequencing (Macrogen, Korea) using reverse and forward pQE30 primers.

Recombinant pQE-30-GBI-Uba7-UFD plasmid transformed into the chemically competent

SG13009 expression strain of *E. coli* using heat shock.

Site-Directed mutagenesis to Uba7-UFD

For site-directed mutagenesis PCR reaction, Pfu polymerase was mixed with template pQE-GBI-

Ube1-UFD plasmid and forward and reverse primers containing the desired mutation. Primer

names and sequences used for mutations are given below.

5' C996A 5' GTGCTAGAGCTGAGCGCTGAGGGTGACGACG 3'

3' C996A 5' CGTCGTCACCCTCAGCGCTCAGCTCTAGCAC 3'

5' C996S 5' GTGCTAGAGCTGAGCAGCGAGGGTGACGACG 3'

3' C996S 5' CGTCGTCACCCTCTCGGCTCAGCTCTAGCAC 3'

5' A1004C 5' GGGTGACGACGAGGACACTTGCTTCCCACCTCTGCACTATG 3'

3' A1004C 5' CATAGTGCAGAGGTGGGAAGCAAGTGTCCTCGTCGTCACCC 3'

Once completed, PCR reaction mixture was incubated with 10 U DpnI restriction enzyme for 2

hours at 37 °C to digest the template plasmid. Resulting reaction mixture containing amplified

mutant plasmid was transformed into Novablue competent *E. coli* cells. The existence of the

desired mutation was confirmed by DNA sequencing. Plasmids containing target mutation were

selected and transformed into competent SG13009 (pREP4) *E. coli* cells. Plasmids encoding

mutant R944L and 998GGG proteins were purchased from GENSCRIPT.

Expression of recombinant GBI-Uba7-UFD fusion protein in *E. Coli*

The freshly-streaked single colony of SG13009 strain of *E. coli* containing PQE30-GBI-Uba7-

UFD plasmid was inoculated in 10 mL Luria Broth (LB) media containing 150 μg/mL ampicillin

and 50 μg/mL kanamycin. Inoculated culture was grown overnight with shaking at 250 rpm at 37

°C. Next morning, 10 mL overnight culture was added to 1 L fresh LB media containing the same

concentrations of antibiotics and grown at 37 °C and 250 rpm until OD_{600} reached 0.8. The protein

expression was induced with 1 mM IPTG. After induction, cells were grown for 5 more hours at

37 °C. Cells were then centrifuged at 4500 x g for 25 minutes at 4 °C and stored at – 80 °C until

protein purification.

Refolding and purification of fusion protein with affinity chromatography

For purification of GBI-Uba7-UFD fusion protein, cell paste was resuspended in 30 mL Lysis

buffer containing 50 mM Tris (pH 7.5), 150 mM NaCl, 5 mM DTT and sonicated at 4 °C, then

centrifuged at 15000 x g for 60 minutes at 4 °C. Proteins in pellet and supernatant were checked

by SDS-PAGE and supernatant were discarded. Inclusion bodies in pellet were washed two times

with 50 mL 0.5% Triton X-100 in 50 mM Tris (pH 7.5), 150 mM NaCl to solubilize membranes

and membrane proteins and then centrifuged at 15000 x g for 60 minutes at 4 °C. Cleaned inclusion

bodies were solubilized in 100 mL denaturation buffer containing 7 M Guanidium-HCl, 50 mM

Tris, 150 mM NaCl, 10 mM DTT overnight with stirring at 20 °C. The next day, the mixture was

centrifuged at 15000 x g for 60 minutes at 15 °C to separate insoluble components from the

solution. Refolding of fusion protein was achieved by dialysis. For this, supernatant was

transferred into a dialysis membrane (MWCO value 3500) and dialyzed twice against 4 liters of

50 mM Tris, 150 mM NaCl, 10 mM DTT pH 7.5 buffer for 8 hours at 15 °C. After second dialysis,

precipitated components were removed by centrifugation at 15000 x g for 60 minutes at 15 °C.

Refolded fusion protein was purified by immobilized metal ion affinity (IMAC) chromatography

using a 5 mL HisTrap HP column (GE HealthSciences) precharged with nickel and attached on an

Akta Prime Plus protein purification system. The dialyzed sample was bound to a HisTrap HP

column at a flow rate of 3 mL/min. Unbound proteins were washed away with a buffer containing

50 mM Tris-HCl, 150 mM NaCl, 5 mM DTT, 20 mM imidazole pH 7.5 until the absorbance at

280 nm was below 10 mAbs. Elution of HisTagged fusion protein was achieved by linearly

increasing the amount of elution buffer (50 mM Tris, 150 mM NaCl, 5 mM DTT, 500 mM

Imidazole pH 7.5) from 0 to 100 % in 30 mL volume. All fractions were analyzed by SDS-PAGE

and those containing the purified GBI-Uba7-UFD fusion protein were pooled, concentrated and

desalted with 20 mM phosphate Buffer pH: 7.4 to 200 μM volume in a pressure concentrator

(Stirred cell, Amicon) using a membrane with 3500 Da MWCO.

Isotopically labeled protein expression

For 1 L of M9 minimal media 200 mL sterile 5x M9 salts lacking ^{15}N-NH$_4$Cl (64 g Na$_2$HPO$_4$

7H$_2$O, 15 g KH$_2$PO$_4$, 2.5 g NaCl), were diluted to 950 mL with dd H$_2$O. 1 g ^{15}N-NH$_4$Cl (dissolved

in 10 mL ddH$_2$O and filter sterilized), 2 mL sterile 1 M MgSO$_4$, 20 mL 20% filter sterilized glucose

(for double labeled protein samples 2 g/L ^{13}C glucose), and 0.1 mL sterile 1M CaCl$_2$ was added to

M9 salts and the volume was completed to 1 L with sterile ddH$_2$O. Media was then supplemented

with trace metals and vitamins.

Freshly transformed cells were used to inoculate 20 mL of LB media containing ampicillin and

kanamycin. Cultures were grown overnight at 37 °C, 250 rpm. Cells were pelleted the next day by

centrifugation at 1000 x g for 10 min at 4 °C, resuspended in 10 mL of M9 minimal media and

added to the 1L of either ^{15}N-labeled M9 culture for ^{15}N-labeled protein production or ^{15}N-^{13}C

labeled M9 culture for double-labeled protein production. Protein expression was carried out as

described above with the exception that after induction with 0.5 mM IPTG, cells were incubated

at 25 °C for 15 hours to allow for expression of target proteins. Cells were then harvested by

centrifugation at 4000 x g for 30 minutes. Cell paste was kept at -20 °C until further processing.

NMR data collection

2D ^{1}H-^{15}N HSQC, 3D HNCA, 3D SE HNCO, 3D SE HNCACB, 3D SE HN(CO)CA, 3D SE

CCONH, 3D SE HCCONH, 3D SE HBHACONH, 3D HNCACO, 3D SE HNCOCACB, 3D

HCCH-TOCSY 3D SE ^{15}N-NOESY-HSQC, 3D SE ^{13}C-NOESY-HSQC-aliphatics and 3D ^{13}C-

NOESY-HSQC-aromatics were performed at 20 ^{0}C on Bruker Avance II 600 MHz spectrometer

with a triple resonance z-axis gradient CryoProbe. All NMR samples were prepared in 20 mM

sodium phosphate, pH 7.4 containing 5-10% D_2O and 0.05% sodium azide.

NMR data processing, analysis and structure calculation

All multidimensional NMR datasets were processed using NMRPipe (Delaglio et al., 1995). Raw

free-induction decay (FID) data were subjected to zero-filling to at least twice the acquired

complex points, apodized with a shifted sine-bell window, and Fourier-transformed in both

dimensions. Phase and baseline corrections were applied manually. Processed spectra were

visualized and analyzed using XEASY and the POKY suite (Lee et al., 2021). Backbone and side-

chain resonance assignments were obtained from standard 3D triple-resonance experiments

(HNCA, HNCO, HNCACB, HN(CO)CA, CCONH, HCCONH, HBHACONH, HCCH-TOCSY,

15N-NOESY-HSQC, and 13C-NOESY-HSQC). Secondary-structure elements were identified

from chemical-shift indices using CSI and torsion-angle restraints derived with TALOS+ (Shen et

al., 2009). Inter-residue NOE distance restraints were extracted from NOESY spectra and

combined with dihedral-angle restraints for structure calculations in POKY/CNS. The final

ensemble of 20 lowest-energy conformers was refined against all experimental restraints and

validated using standard NMR structure-quality metrics (RMSD, Ramachandran, and NOE

statistics).

All NMR titration experiments and HSQC spectra were acquired using isotopically labeled protein

samples at a concentration of 200 μM. HSQC cross-peak movements were quantified as a

combined ^1H-^{15}N chemical shift perturbation calculated from $\Delta\delta AV = [(\Delta\delta\ ^1H*5)^2 + (\Delta\delta\ ^{15}N)^2]^{1/2}$

equation. The structure calculations were performed using POKY (Lee et al., 2021) programs on

a computer running the MacOS operating system.

Docking Analysis

The docking process in HADDOCK2.4 docking protocol (de Vries et al., 2010) is divided into

three stages. In the first stage, the proteins are treated as rigid bodies, and their interaction energy

is calculated for different orientations. The orientations with the lowest energy are then selected

for further refinement. In the second stage, simulated annealing is used to refine the interface

between the two proteins. Simulated annealing is a Monte Carlo method that allows the system to

escape from local energy minima. In the context of protein-protein docking, simulated annealing

allows the proteins to explore different conformations at the interface and identify the

conformation with the most favorable energy. In the final stage, the complex is immersed in

solvent to improve the energetics. The solvent molecules are explicitly included in the simulation,

and their interactions with the proteins are taken into account. This stage helps to refine the

structure of the complex and ensure that it is stable in a solvated environment. By combining these

three stages, HADDOCK2.4 outputs a prediction of the structure of protein-protein complex.

NMR relaxation measurements

Backbone dynamics of Uba7-UFD and UbcH8 were characterized using ^{15}N relaxation

experiments (T_1, T_2, and {^{1}H}-^{15}N heteronuclear NOE) to assess conformational flexibility and

dynamic changes upon complex formation. Relaxation measurements were performed for the free

^{15}N-labeled Uba7-UFD and free ^{15}N-labeled UbcH8 under identical buffer conditions (20 mM

phosphate buffer, pH 7.4, 150 mM NaCl, 5 mM DTT, 5–10% D$_2$O).

Experiments for Uba7-UFD were carried out at 20 °C on a Bruker Avance III 900 MHz

spectrometer equipped with a triple-resonance cryogenic probe, while UbcH8 relaxation data were

acquired under identical conditions on a Bruker Avance II 600 MHz spectrometer.

Standard Bruker pulse sequences were used for all relaxation measurements. T_1 relaxation times

were obtained using eight relaxation delays (10, 50, 100, 200, 400, 600, 800, and 1000 ms), and

T_2 relaxation times were recorded using eight delays (10, 30, 50, 70, 90, 110, 130, and 150 ms).

{^{1}H}-^{15}N heteronuclear NOE spectra were collected with and without proton presaturation (3 s)

using identical acquisition parameters. All spectra were processed with NMRPipe and analyzed

using Bruker Dynamics Center software.

Results

Backbone and side-chain resonance assignments of native human Uba7-UFD

Uba7-UFD residues 903-1011 were recombinantly produced with an N-terminal GBI fusion in *E.*

coli cells and purified using Ni^{2+} affinity chromatography. Despite efforts to produce a soluble

construct, the GBI-Uba7-UFD expressed into inclusion bodies similar to our previous study with

Uba3-UFD (Elgin et al., 2012). The most effective approach to refold GBI-Uba7-UFD was

denaturation with 7 M guanidinium chloride and 5 mM DTT followed by dialysis into a solution

containing 50 mM Tris, 150 mM NaCl, 5 mM DTT, pH 7.5. Unfortunately, Uba7-UFD possessed

very low stability/solubility once the GBI solubility tag was removed resulting in precipitation

within ~4 h (data not shown). We next evaluated the possibility of using GBI-Uba7-UFD fusion

for NMR structure and binding studies. Comparison of 2D ^{1}H-^{15}N HSQC spectra for both the free

domain and the GBI fusion showed no detectable difference in the amide resonances

demonstrating the fusion does not affect the Uba7-UFD tertiary structure (Supp Fig. 1).

Comparison of ^{1}H-^{15}N HSQC spectra collected before (day 1) and after (day 9) a suite of 3D

assignment experiments reveals no significant time-dependent changes (Supp Fig. 2). Therefore,

GBI-Uba7-UFD was found to be suitable for further structure data collection and titration

experiments.

The chemical shift dispersion of [U-^{15}N]-GBI-Uba7-UFD amide resonances are consistent with a

well-folded protein (Fig. 2A); however, only 71 of the 100 expected backbone N-H resonances

were observed (109 total residues including eight prolines). These were assigned to residues 919-

1011 using a suite of conventional 3D triple-resonance spectra from which secondary structure

was predicted using TALOS+ (Shen et al., 2009) (Fig. 2A). A stretch of seven residues following

the third beta-strand were unassignable. The homologous region of the two E1 enzyme UFDs,

Uba3 and Uba2 (Wang et al., 2009; Elgin et al., 2012), were similarly unassigned due to low signal

intensity in 3D experiments which suggests these missing peaks result from inherent plasticity of

these regions (Fig. 2B).

NMR Structure determination of Human Uba7-UFD

The three-dimensional structure of the human Uba7-UFD was successfully determined using

multi-dimensional NMR spectroscopy (Fig. 2C). A total of 1616 unique NOE distance and 153

(phi/psi) dihedral angle constraints are used in the final structure calculations; refinement statistics

for the ensemble of 20 lowest energy conformers are shown in Table 1. The final ensemble

exhibited good convergence with an average pairwise root mean square deviation (RMSD) of 0.9

Å for backbone atoms and 1.4 Å for all heavy atoms. The secondary structure consists of three α-

helices and four β-strands arranged in a ubiquitin fold domain (Fig. 2B, 2C; Supp Fig. 4) consistent

with previously reported ubiquitin-like domains (Wang et al., 2010; Lv et at., 2018). After the

second β-strand, a partially folded intermediate third β-strand is present, with distinct loops located

before and after it. These loops and β-strand may play a role in the interaction with the E2 enzyme

in the ISGylation cascade, similar to the mechanism observed in ubiquitination (Lv et al., 2018).

The crystal structures of apo E1 enzymes (PDB:7ZH9, 6CWZ, 7PVB) lack electron density in

certain portions of the UFDs (Walden, Podgorski, and Schulman, 2003) (Figure 1). This is

consistent with NMR analyses that the UFDs contain intrinsically disordered regions in their free

form (Wang et al., 2009; Elgin et al., 2012) that adopt stable secondary structures upon E1-E2

complexation (Huang et al., 2005; Wang et al., 2010) (Figure 1). Comparison of NMR structures

of UFD's from Uba7 and Uba3 shows that Uba7-UFD NMR structure shows a well-folded second

alpha helix (Figure 2B-Uba7_960-968), which is largely unfolded in Uba3-UFD (Figure 2B-

Uba3_397-403) (PDB 2LQ7; Fig. 1J-K,2B). In contrast, the third beta strand of Uba7-UFD (Figure

2B-Uba7_950-960), which is well defined in Uba3-UFD (Figure 2B-Uba3_388-392), is relatively

flexible (Fig. 1H-K). In addition, the NMR-derived structure showed that the acidic loop region

located between the fourth and fifth beta strands also exhibits much greater flexibility compared

to the corresponding loop region in Uba3-UFD (Fig. 1H-K). Although the overall ubiquitin fold

structure is conserved in both Uba7- and Uba3-UFDs, NMR ensembles clearly demonstrate that

the location of conformational flexibility is quite unique to each domain.

Chemical shift mapping reveals the Uba7-UbcH8 interface

To test the hypothesis that E1-UFD conformational disorder plays a role in E2

binding/selection/specificity, we performed an NMR titration to map UbcH8 onto the Uba7-UFD

surface. ^{1}H-^{15}N HSQC spectra were collected on [U-^{15}N]-Uba7-UFD mixed with increasing

concentrations of unlabeled-UbcH8 in molar ratios of 1:0.25, 1:0.5, 1:1 and 1:1.75 (Supp Fig. 5).

Inspection of these spectra revealed a subset of peaks that were dose-dependently perturbed upon

addition of UbcH8 (Figure 3A); combined ^{1}H/^{15}N chemical shift changes were calculated and the

most perturbed residues were highlighted onto the Uba7-UFD structure (Figure 3B, 3C). UbcH8

binding surface of Uba7-UFD appears to be quite large, involving about 18 residues. Interestingly,

some of the peaks displaying the largest chemical shift perturbations upon addition of UbcH8 were

weaker in intensity compared to the average peak intensities in the spectrum of the free Uba7-

UFD. In fact, four peaks with pronounced chemical shift perturbations corresponded to those

residues in the unassigned fragments and were absent in 3D spectra. One of the simplest

explanations for the low intensity and absence in 3D spectra is that these residues may undergo

conformational exchange on the microsecond-millisecond timescale (Supp Fig. 6).

We next mapped the binding interface onto the UbcH8 surface by titrating increasing

concentrations of Uba7-UFD into [U-^{15}N]-UbcH8. Substantial chemical shift perturbations

clustered to the UbcH8 N-terminal region, helix 1, strands 1-3, and helix 3 (Figure 3D-F). Several

peaks were also noted to disappear in the N-terminal region, helix 1, strands 1-3, and helix 3 upon

addition of unlabeled Uba7-UFD (Supp Fig. 7). This result is consistent with the mutational study

showing that the N-terminal region of UbcH8 is important in Uba7 interaction (Huibregtse et al.

2008).

NMR data driven docking of UbcH8 and Uba7-UFD complex

Next, we used the most substantial CSPs from our NMR titrations on the isolated domain

interactions to restrain a model of the full-length Uba7-HFB/UbcH8 complex using the

HADDOCK 2.4 NMR-based docking protocol (Vries et al., 2010). The AlphaFold Uba7 model

(APSD: AF-P41226-F1-v4) and the UbcH8 crystal structure (PDB: 1WZV) were used as inputs

and the resulting cluster with the lowest HADDOCK score was analyzed (Figure 4). The

HADDOCK model illustrates how complexation is facilitated by two sets of charge-charge

interactions; specifically, Uba7-UFD ^{999}DDEDTA^{1003} and ^{950}SALLYAAGWSPEK^{962}

complement UbcH8 ^{5}MRVVKE^{10} and ^{25}LSSDD^{29}, respectively. Furthermore, a series of

hydrophobic and charged residues surrounding the active site cysteine undergoes conformational

changes during the transthiolation stage (Figure 4). The spatial arrangement of these regions

precisely orients UbcH8 to ensure efficient ISG transfer between catalytic cysteine residues. We

also collected relaxation datasets for the Uba7-UFD:UbcH8 complex at a 1:2 molar ratio; however,

several resonances disappeared upon complex formation, preventing reliable fitting of relaxation

curves and thus precluding publication-quality analysis.

Dynamics of Uba7 UFD

To characterize the internal motions and structural rigidity of the Uba7 UFD in solution, backbone

^{15}N relaxation measurements were obtained, including longitudinal relaxation times (T_1),

transverse relaxation times (T_2), T_1/T_2 ratios, and {^1H}-^{15}N heteronuclear NOE values (Figure 5).

The overall relaxation profiles demonstrate that Uba7-UFD adopts a well-folded and dynamically

stable β-grasp architecture, with relaxation parameters typical for a compact monomeric protein

of this size (~11 kDa). Regions corresponding to the β2 (residues 944-951) and β4 (residues 984-

989) strands, which form the E2-binding surface are highlighted in orange. These segments show

slightly lower {^1H}-^{15}N NOE values and β2 segment show slightly higher T_2 relaxation times,

indicating enhanced local flexibility relative to the β-sheet core. Such mobility is often observed

in loop regions or solvent-exposed β-strand edges that mediate transient interactions. The loop

region spanning residues 996-1008 exhibits low {^1H}-^{15}N NOE values and prolonged T_2 relaxation

times, indicating enhanced backbone flexibility compared with the rest of the Uba7-UFD domain.

This flexible segment forms part of the surface that interacts with ISG15 and likely contributes to

its recognition. The observed mobility suggests that this loop can undergo transient conformational

adjustments, allowing optimal positioning of ISG15 during E1-E2-substrate transfer. Such

dynamic behavior may therefore play a crucial role in facilitating ISG15 conjugation and ensuring

efficient catalysis within the ISGylation pathway.

To assess how E2 recruitment reshapes UFD dynamics, equivalent relaxation measurements were

collected for ^{15}N-labeled Uba7-UFD in the presence of unlabeled UbcH8 (Figure 5E–H).

Consistent with complex formation, the relaxation behavior shifts toward values expected for a

higher effective molecular size, including increased T_1 and elevated T_1/T_2 ratios across the

observable residues (Figure 5E,G). At the same time, residue-specific heterogeneity becomes more

pronounced within the highlighted β2/β4 surface and adjacent segments, consistent with binding-

coupled changes in local mobility and/or conformational exchange at the interface (Figure 5F,H).

We note that complex formation can lead to the disappearance of a subset of resonances, which

limits quantitative analysis for all residues and is consistent with intermediate-timescale exchange

contributions in portions of the binding surface. The elevated mobility observed in β2/β4 and the

996–1008 loop provides a plausible physical basis for the disorder-to-order or adaptive-fit

behavior inferred from binding and structural analyses, helping to reconcile how Uba7-UFD can

remain stably folded while retaining the conformational plasticity required for efficient E2

recruitment and productive transthiolation during ISGylation.

Together, these data reveal that Uba7-UFD possesses a hierarchical dynamic profile, where

structural rigidity within the β-sheet framework coexists with localized flexibility at the potential

binding surface features likely critical for efficient recognition and alignment with its cognate E2

enzyme during ISGylation.

Dynamics of E1 UFD binding surface on UbcH8

The dynamical behavior of the UbcH8 was also probed using ^{15}N T1, ^{15}N T2, and {^{1}H}-^{15}N

heteronuclear NOE relaxation experiments (Lakomek, Ying and Bax, 2013). Figures 6A and 6B

show the T1 and T2 relaxation rates plotted against the residue number and color mapped on the

crystal structure of UbcH8 (PDB:1WZW), respectively. Residues with high T1 and low T2

relaxation rates corresponding to relatively rigid regions are colored in green whereas those

displaying lower T1 and higher T2 relaxation rate hence higher flexibility are shown in yellow,

orange, and red in the order of increasing flexibility. Gray color corresponds to the unassigned

region. Figure 6C shows the heteronuclear ^{15}N-^{1}H NOE values plotted as a function of UbcH8

amino acid sequence. Fitting of data was done with the help of Bruker dynamics center and which

showed more than 95% confidence level. All together, T1 and T2 relaxation rates and

heteronuclear NOE values measured for UbcH8 consistently show that while the core of the

protein is rigid, the outward-facing residues on the conserved N-terminal α-helix show enhanced

flexibility, consistent with their proposed role in the UbcH8-binding interface (Figure 6D). Our

docking studies show that this region of UbcH8 is directly involved in the interaction with Uba7-

UFD. This relatively higher mobility may contribute to the E2's interaction and function by

facilitating conformational changes necessary for E1–E2 transthiolation.

Conformational Flexibility of Uba7-UFD and Its Role in E2 Binding: Insights from Mutational

Analysis

NMR solution structure of free Uba7-UFD indicated that the putative E2 binding region might be

flexible and experiencing conformational plasticity, transitioning among multiple conformational

states. Comparably, conformational flexibility/heterogeneity has been observed in E2 binding

surfaces of SUMO (Wang et al., 2009) and Nedd8 E1 UFD (Elgin et al., 2012). Notably this

conformational flexibility has been shown to be functionally important in SUMO conjugation

(Wang et al., 2009).

To further explore this phenomenon, we designed and analyzed a series of mutants to determine

how these alterations would affect E2 binding to Uba7-UFD. Specifically, we aimed to identify

the role of individual amino acids in maintaining the structural integrity and interaction capabilities

of the UFD domain. Given the high degree of plasticity observed in the putative E2 binding region,

understanding how mutations influence this flexibility is crucial. This investigation not only

provides insight into the specific amino acid interactions that are critical for binding but also sheds

light on the overall mechanism by which Uba7-UFD achieves its conformational adaptability. By

mapping these interactions and comparing the behavior of wild-type and mutant proteins, we

sought to delineate the structural elements that enable effective binding and subsequent functional

outcomes.

Role C996 in Uba7-UFD conformational integrity and UbcH8 binding

In order to determine the effect of mutations on E2 binding to Uba7-UFD, seven mutants were

created and five of them also titrated with UbcH8. Full length Uba7 protein contains a total of 17

cysteine residues, only one of which is in the UFD. Since this domain does not contain any other

cysteine residue to form disulfide bridge with C996, this single cysteine probably remains reduced.

C996 is located in the end of the second beta sheet, and while the subsequent amino acids show a

very flexible behavior in the structure, cysteine and previous amino acids have formed a stable

beta sheet secondary structure form.

We first mutated the free cysteine to alanine, which is the most preferred amino acid in mutation

scans. Overlay of ^{1}H-^{15}N HSQC spectra of wild-type and C996A variant shows that the overall

conformations of the mutant and wild-type proteins remain largely the same (Supp Figure 9). In

contrast, changes in chemical shift values were observed around the mutated region, indicating a

systematic reorganization in the preceding and subsequent beta sheets (Figure 7A). Figure 7B

shows the residues affected by C996A mutation mapped onto the Uba7-UFD structure. Residues

with chemical shift perturbations larger than 0.4 and 0.2 ppm colored in orange and merigold,

respectively, are mostly clustered in neighboring B1, and B4 strands and alpha 1 helix. In order to

determine the effect of mutation on E2 binding to Uba7-UFD, mutant protein was titrated with

UbcH8. Supp Figure 10 shows the overlay of ^{1}H-^{15}N HSQC spectra of Uba7-UFD-C996A before

and after addition of various amounts of UbcH8. No significant ^{1}H-^{15}N chemical shift

perturbations in C996A mutant upon addition of UbcH8 was observed and addition of more and

more of UbcH8 resulted in disappearance of some of the peaks involved in binding. These titration

studies clearly indicate that C996 is important for Uba7-UFD-UbcH8 interaction (Supp Figure 10).

Mutation of C996 to alanine appeared to have effect on UbcH8 binding. Since mutation of cysteine

to alanine results in truncation of the side chain at the C-beta, the goal of C996S mutation was to

replace –SH group with –OH group to see if this would restore the UbcH8 binding. The

comparison of ^{1}H-^{15}N HSQC spectrum of C996S with that of wild-type shows that the overall

conformation of C996S mutant is very similar to that of wild-type (Supp Figure 11) and as in

C996A mutant, significant chemical shift perturbations are located around the mutation site

(Figure 7C-7D). NMR binding studies carried out using ^{15}N labeled Ubc7-UFD C996S further

confirmed the role of C996 in UbcH8 binding since there were no significant chemical shift

perturbations in the ^{1}H-^{15}N HSQC spectra of C996S mutant upon addition of UbcH8(Supp Figure

12). The failure of the -OH group in restoring the UbcH8 binding suggests that free -SH might not

merely be involved in hydrogen bonding but might be important in providing necessary flexibility

for UbcH8 interaction (Bhopatkar et al., 2020).

After determining that free cysteine was necessary for UbcH8 binding/for keeping Uba7-UFD in

a suitable conformation for the UbcH8 binding we searched for amino acids in close proximity to

C996 within Uba7-UFD to test the effect of breaking C996 intermolecular interactions on the

conformation of the domain as well as on UbcH8 binding. Amino acids directly interacting with

C996 were identified through NOE analysis. NOE peaks observed between C996 and R944

indicated that there is an interaction between these two amino acids. Therefore, an R944L mutant

protein was generated, produced, and purified after refolding. Gel filtration chromatography of

purified ^{15}N-GBI-Uba7-UFD resulted in two separate peaks with the same molecular weight

confirmed by SDS-PAGE analysis. Peaks were collected in separate fractions and analyzed by

NMR. ^{1}H-^{15}N HSQC spectra showed that the peak eluted first from the gel filtration

column was completely misfolded/aggregated (Supp Figure 13A) while the fraction

corresponding to the second peak consisted of a mixture of both folded and misfolded

fusion protein (Supp Figure 13B). Gel filtration chromatography and NMR results

clearly show that R944 and its interaction with C996 are important for conformational

stability of Uba7–UFD and mutation of R944 to alanine makes domain prominent to

aggregation. The role of free cysteine amino acid in protein structure and internal

dynamics is often overlooked. However, a study conducted by Mazmanian et al. in 2016

examined the participation and significance of free cysteine amino acids in hydrogen

bonding networks in detail (Mazmanian et al., 2016). In the case of the Uba7–UFD

complex, when a cysteine atom forms a hydrogen bond with arginine (R944), it exhibits

distinct interaction patterns depending on the charge of the hydrogen bond donor.

Specifically, cysteine forms shorter and more linear hydrogen bonds with positively

charged residues such as histidine (His$^+$), lysine (Lys$^+$), and arginine (Arg$^+$), where the

hydrogen–sulfur distance (H$\cdots$S) is less than 2.5 Å, and the bond angle (N–H$\cdots$S) exceeds

150° (Mazmanian et al., 2016). In contrast, hydrogen bonds with neutral donors are

longer (H$\cdots$S > 2.5 Å) and less linear (N–H$\cdots$S < 150°). Notably, Arg$^+$ donates two

hydrogen bonds to the cysteine thiol (CysSH), a characteristic not observed in other

residues (Mazmanian et al., 2016). These findings align with previous studies

demonstrating that positively charged hydrogen bond donors preferentially form

stronger and more directional interactions with thiol groups, contributing to protein

stability and conformational flexibility. Thus, the Uba7–UFD complex exemplifies the

critical role of free cysteines in fine–tuning the stability and plasticity of protein

structures.

Effect of acidic loop flexibility and length on Uba7-UFD interaction with UbcH8

As mentioned above, our NMR solution structure of Uba7-UFD shows that the loop region that

comes right after the C996 is relatively flexible. Since this loop is involved in interaction with

UbcH8, one question is whether restricting this flexibility would prevent binding. To test this,

another mutant was designed aiming to form a disulfide bond between the single cysteine 996 and

alanine 1004. In this mutant, A1004 was individually replaced by cysteine. This variant was also

expressed as inclusion bodies, solubilized with 7M GdnHCl and refolded by dialysis as described

previously. Supp Figure 14 shows an overlay of ^{1}H-^{15}N HSQC spectra of A1004C mutant in the

absence and presence of various amounts of UbcH8. Figure 7E shows the chemical shift

perturbation analysis from the titration experiment. Unlike the previous mutations, this mutant

variant still showed intrinsically interacted with UbcH8. As seen in Figure 7F, the interaction

surface is also partially compatible with the interaction surface of the WT protein. Opposite to

WT, A1004C mutant's 1st beta sheet region did not show any interaction with UbcH8.

The acidic loop interacting with UbcH8 has been extended with a glycine amino acid as a final

mutation approach (998GGG). The 998GGG mutant protein has been produced, purified, and

subjected to ^{1}H-^{15}N HSQC spectrum collection. Although there were regional differences in the

obtained spectrum, no significant changes were observed in the overall distribution (Supp Figure

15A). Unlabeled UbcH8 has been added to the obtained protein in a 1:1 ratio. Upon examination

of the spectra obtained after mixing the apo form of 998GGG with UbcH8, no changes were

observed in the chemical shift values indicating that integrity of the loop was necessary for UbcH8

binding. (Supp Figure 15B).

Discussion

Ubiquitin and Ubiquitin-like post-translational modification systems are of great importance to

organisms due to their extensive regulatory roles within cells. Elucidating the mechanisms at the

atomic level within the enzymatic cascades that form the basis of Ub and Ubl post-translational

modification systems has been a significant topic in biochemistry and structural biology for

decades. Although the atomic-level details of the mechanisms of systems such as ubiquitination,

neddylation, and sumoylation have been examined, structural studies on the ISGylation system

were not available until recent years. Specifically, due to the high plasticity of the ubiquitin fold

domain of the E1 enzyme, determining its structure has taken a considerable amount of time.

During the preparation of our manuscript, two additional papers aimed at elucidating the details of

ISGylation have been published (Wallace et al., 2023; Afsar et al., 2023). Although Cryo-EM and

crystal structures are excellent tools for visualizing the overall mechanism of systems, they are

still static and frozen structures. This makes them less effective for examining weak interaction

surfaces and low-affinity complex structures. At this point, NMR spectroscopy continues to be an

important structural biology method, providing atomic-level information to complement the

remarkable findings from other methods and to elucidate the side effects caused by freezing. In

the first part of our study, we determined the NMR structure of the ISGylation E1 enzyme and

observed that the plasticity and unstable beta-sheet surfaces exhibited in solution are believed to

play a significant role in the system's operation and selectivity.

Contemporary Cryo-EM structures, especially those examining the E1-E2 interaction, present

differing results regarding the amino acids and interaction angles involved in the interaction

between the first alpha helix of the E2 enzyme and the high plasticity regions of the E1 UF domain

(Wallace et al., 2023; Afsar et al., 2023). Despite the high overall resolutions of these structures,

it is notable that the resolution in these domains sometimes drops regionally to 4-5 Å. The direct

interactions obtained from our NMR data-driven model show greater alignment with the findings

of Wallace et al. 2023.

Findings from T1 and T2 relaxation experiments, coupled with titration and docking data, provide

a coherent picture suggesting that the N terminus region of Ubch8 is more flexible and that this

flexibility is relevant to the interaction with the Uba7-UFD. NMR characterization studies of

Uba7-UFD variants showed that both wild type and mutant domains have a high tendency to

aggregate. None of the designed mutant Uba7-UFDs was more stable than the wild type Uba7-

UFD. Our NMR mapping study showed that UbcH8 binding surface of Uba7-UFD is substantial

and there are over 30 amino acids directly involved in binding. This study also showed that a single

reduced cysteine residue at position 996 may play a major role in domain stability and fine tuning

of plasticity.

Conclusion

Our study has reiterated the importance of examining protein structures and dynamics in solution.

It has been demonstrated that the UFD domain of the Uba7 protein undergoes conformational

changes upon binding to UbcH8, consistent with the induced fit model. Structural analysis of the

Uba7 UFD and complex structures obtained from other systems have shown that a large surface

area of the UFD interacts with E2, highlighting the significance of this interaction in E1-E2 binding

and proper E2 selection. Additionally, structural investigations have revealed that a region on the

E1 UFD interacts with Ubiquitin/Ubiquitin-like proteins. It is important to conduct new studies to

demonstrate the importance of this interaction in structural and functional regulation.

Author Contributions

Çağdaş Dağ: Conceptualization, Methodology, Supervision Funding acquisition, Investigation, Formal analysis, Writing- Reviewing and Editing, Writing-Original draft preparation **Alp Eren Kazar:**Investigation, Formal analysis, Visualization, Writing- Reviewing and Editing, Writing- Original draft preparation **Emine Sonay Elgin:** Conceptualization, Methodology, Supervision Funding acquisition, Investigation, Formal analysis, Writing- Reviewing and Editing, Writing- Original draft preparation **Arthur L. Haas:** Conceptualization, Supervision **Kerem Kahraman**: Investigation, Formal analysis, Visualization. **Oktay Göcenler:** Formal analysis, Investigation Writing - Original Draft, Visualization **Mahil Lambert:** Formal analysis, Investigation, Writing - Original Draft, Visualization **Volker Dötsch**: Conceptualization, Supervision, Writing- Reviewing and Editing **Cansu D. Tozkoparan:**Visualization, Formal analysis **Joshua J. Ziarek:** Conceptualization, Writing- Reviewing and Editing

ACKNOWLEDGMENT

We gratefully acknowledge Prof. Brian F. Volkman (Medical College of Wisconsin) for generously opening his laboratory to us and providing access to critical instrumentation and consumables (GM64598-03). His support and willingness to share resources were invaluable for the successful execution of this work. We sincerely thank him for his scientific generosity and hospitality. ESE acknowledges support from TÜBİTAK (Project No: 104T193). CD acknowledges support from TÜBİTAK (Project No: 120Z594, 122Z747). JJZ acknowledges support from National Institutes of Health grant R35GM143054. The authors acknowledge the use of the services and facilities of n^2STAR-Koç University Nanofabrication and Nanocharacterization Center for Scientific and Technological Advanced Research. This study made use of the National Magnetic Resonance Facility at Madison, which is supported by NIH grant R24GM141526 and

P41GM103399. This work benefited from access to [CERM and BMRZ] and has been supported

by iNEXT-Discovery, project number 871037, funded by the Horizon 2020 program of the

European Commission. Financial support by the Access to Research Infrastructures activity in the

7th Framework Programme of the EC (Project number: 261863, Bio-NMR) for conducting the

research is gratefully acknowledged. This study made use of NMRbox: National Center for

Biomolecular NMR Data Processing and Analysis, a Biomedical Technology Research Resource

(BTRR), which is supported by NIH grant P41GM111135 (NIGMS).

REFERENCES

Afsar, M., Liu, G., Jia, L., Ruben, E. A., Nayak, D., Sayyad, Z., Bury, P. D. S., Cano, K. E., Nayak, A., Zhao, X. R.,

Shukla, A., Sung, P., Wasmuth, E. V., Gack, M. U., & Olsen, S. K. (2023). Cryo-EM structures of Uba7 reveal the

molecular basis for ISG15 activation and E1-E2 thioester transfer. *Nature communications*, *14*(1), 4786.

https://doi.org/10.1038/s41467-023-39780-z

Bogunovic, D., Byun, M., Durfee, L. A., Abhyankar, A., Sanal, O., Mansouri, D., Salem, S., Radovanovic, I., Grant,

A. V., Adimi, P., Mansouri, N., Okada, S., Bryant, V. L., Kong, X. F., Kreins, A., Velez, M. M., Boisson, B.,

Khalilzadeh, S., Ozcelik, U., Darazam, I. A., … Casanova, J. L. (2012). Mycobacterial disease and impaired IFN-γ

immunity in humans with inherited ISG15 deficiency. *Science (New York, N.Y.)*, *337*(6102), 1684–1688.

https://doi.org/10.1126/science.1224026

Bhopatkar AA, Uversky VN, Rangachari V. Disorder and cysteines in proteins: A design for orchestration of

conformational see-saw and modulatory functions. Prog Mol Biol Transl Sci. https://10.1016/bs.pmbts.2020.06.001.

Capili, A. D., Lima, C. D. (2007). Structure and analysis of a complex between SUMO and Ubc9 illustrates features

of a conserved E2-Ubl interaction. *Journal of molecular biology*, *369*(3), 608–618.

https://doi.org/10.1016/j.jmb.2007.04.006

Durfee, L. A., Lyon, N., Seo, K., Huibregtse, J. M. (2010). The ISG15 conjugation system broadly targets newly

synthesized proteins: implications for the antiviral function of ISG15. *Molecular cell*, *38*(5), 722–732.

https://doi.org/10.1016/j.molcel.2010.05.002

de Vries, S. J., van Dijk, M., Bonvin, A. M. (2010). The HADDOCK web server for data-driven biomolecular docking.

Nature protocols, *5*(5), 883–897. https://doi.org/10.1038/nprot.2010.32

Elgin, E. S., Sökmen, N., Peterson, F. C., Volkman, B. F., Dağ, C., Haas, A. L. (2012). E2-binding surface on Uba3

β-grasp   domain   undergoes   a   conformational   transition.   *Proteins*,   *80*(10),   2482–2487.

https://doi.org/10.1002/prot.24148

Farrell, P. J., Broeze, R. J., Lengyel, P. (1979). Accumulation of an mRNA and protein in interferon-treated Ehrlich

ascites tumour cells. *Nature*, *279*(5713), 523–525. https://doi.org/10.1038/279523a0.

Haas, A.L. (2007). ISG15-dependent Regulation. In Protein Degradation Series (eds R.J. Mayer, A.J. Ciechanover

and M. Rechsteiner). https://doi.org/10.1002/9783527619320.ch5b

Huang, D. T., Paydar, A., Zhuang, M., Waddell, M. B., Holton, J. M., & Schulman, B. A. (2005). Structural basis for

recruitment of Ubc12 by an E2 binding domain in NEDD8's E1. Molecular cell, 17(3), 341-350

Jubb, H., Blundell, T. L., Ascher, D. B. (2015). Flexibility and small pockets at protein-protein interfaces: New

insights   into   druggability.   *Progress   in   biophysics   and   molecular   biology*,   *119*(1),   2–9.

https://doi.org/10.1016/j.pbiomolbio.2015.01.009

Lakomek, N. A., Ying, J., Bax, A. (2012). Measurement of ^{15}N relaxation rates in perdeuterated proteins by TROSY-

based methods. *Journal of biomolecular NMR*, *53*(3), 209–221. https://doi.org/10.1007/s10858-012-9626-5

Lee, W., Rahimi, M., Lee, Y., Chiu, A. (2021). POKY: a software suite for multidimensional NMR and 3D structure

calculation   of   biomolecules,   *Bioinformatics*,   Volume   37,   Issue   18,   Pages   3041–3042,

https://doi.org/10.1093/bioinformatics/btab180

Loeb, K. R., Haas, A. L. (1992). The interferon-inducible 15-kDa ubiquitin homolog conjugates to intracellular

proteins. *The Journal of biological chemistry*, *267*(11), 7806–7813.

Lois, L. M., & Lima, C. D. (2005). Structures of the SUMO E1 provide mechanistic insights into SUMO activation

and E2 recruitment to E1. *The EMBO journal*, *24*(3), 439-451.

Lv, Z., Williams, K.M., Yuan, L., Atkison, JH, Olsen SK. Crystal structure of a human ubiquitin E1-ubiquitin complex

reveals conserved functional elements essential for activity. J Biol Chem. 2018 Nov 23;293(47):18337-18352. doi:

10.1074/jbc.RA118.003975

Mazmanian, K., Sargsyan, K., Grauffel, C., Dudev, T., & Lim, C. (2016). Preferred Hydrogen-Bonding Partners of

Cysteine: Implications for Regulating Cys Functions. *The journal of physical chemistry. B*, *120*(39), 10288–10296.

https://doi.org/10.1021/acs.jpcb.6b08109

Narasimhan, J., Potter, J. L., Haas, A. L. (1996). *Conjugation of the 15-kDa Interferon-induced Ubiquitin Homolog*

Is Distinct from That of Ubiquitin. Journal of Biological Chemistry, 271(1), 324–330. doi:10.1074/jbc.271.1.324

Olsen, S. K., & Lima, C. D. (2013). Structure of a ubiquitin E1-E2 complex: insights to E1-E2 thioester transfer.

Molecular cell, 49(5), 884-896.

Poller, W., Knobeloch, K. P., Voigt, A. (2014). Ubiquitin-like protein ISG15 (interferon-stimulated gene of 15 kDa)

in host defense against heart failure in a mouse model of virus-induced cardiomyopathy. *Circulation*, *130*(18), 1589–

1600. https://doi.org/10.1161/CIRCULATIONAHA.114.009847

Rahnefeld, A., Klingel, K., Schuermann, A., Diny, N. L., Althof, N., Lindner, A., … Voigt, A. (2014). *Ubiquitin-Like*

Protein ISG15 (Interferon-Stimulated Gene of 15 kDa) in Host Defense Against Heart Failure in a Mouse Model of

Virus-Induced   CardiomyopathyCLINICAL   PERSPECTIVE.   Circulation,   130(18),   1589–1600.

doi:10.1161/circulationaha.114

Sadler, A. J., Williams, B. R. (2008). Interferon-inducible antiviral effectors. *Nature reviews. Immunology*, *8*(7), 559–

568. https://doi.org/10.1038/nri2314

Shen, Y., Delaglio, F., Cornilescu, G. et al. TALOS+: a hybrid method for predicting protein backbone torsion angles

from NMR chemical shifts. J Biomol NMR 44, 213–223 (2009).

Streich, F.C., Haas, A.L. (2010). Activation of Ubiquitin and Ubiquitin-Like Proteins. In: Groettrup, M. (eds)

Conjugation and Deconjugation of Ubiquitin Family Modifiers. Subcellular Biochemistry, vol 54. Springer, New

York, NY. https://doi.org/10.1007/978-1-4419-6676-6_1

Wang, J., Taherbhoy, A. M., Hunt, H. W., Seyedin, S. N., Miller, D. W., Miller, D. J., Huang, D. T., & Schulman, B.

A. (2010). Crystal structure of UBA2(ufd)-Ubc9: insights into E1-E2 interactions in Sumo pathways. *PloS one*, *5*(12),

e15805. https://doi.org/10.1371/journal.pone.0015805

Zhang, M., Li, J., Yan, H., Huang, J., Wang, F., Liu, T., Zeng, L., Zhou, F. (2021). ISGylation in Innate Antiviral

Immunity and Pathogen Defense Responses: A Review. *Frontiers in cell and developmental biology*, *9*, 788410.

https://doi.org/10.3389/fcell.2021.788410

Figure 1. Ubiquitin and Ubiquitin-like post-translational modification systems E1 enzymes' ubiquitin fold

domain structures. Regions on the structures that have high plasticity and directly interact with E2 are

marked in red, and regions thought to interact with Ubl are marked in orange. Electron density maps are

also shown over the colored regions. A. Saccharomyces cerevisiae Uba1 Apo form (PDB: 7ZH9) B. Homo

sapiens Uba1 complex with ubiquitin (PDB: 6DC6) C. Saccharomyces cerevisiae complex with Ubc13

(E2) (PDB: 6ZHS) D. Schizosaccharomyces pombe complex with Ubc15 (E2) and ubiquitin (PDB: 5KNL)

E. Homo sapiens Uba2 apo form (6CWZ) F. Homo sapiens Uba2 complex with SUMO (3KYC) G.

Saccharomyces cerevisiae Uba2-UFD complex with Ubc9 (3ONG) H. Homo sapiens Uba3 apo form

(1YOV) I. Homo sapiens Uba3-UFD complex with Ubc12 (1Y8X) J. Homo sapiens Uba3 UFD apo form

(2LQ7) K. Homo sapiens Uba7-UFD apo form (8WWX) L. Homo sapiens apo Uba6 (7PVN) M. Homo

sapiens Uba6 complex with FAT10 (7PYV) N. Homo sapiens Uba6 complex with FAT10 (7SOL)

Figure 2. NMR structure of Human Uba7 ubiquitin fold domain A. Human Uba7 enzyme ubiquitin fold

domain ^{1}H-^{15}N HSQC spectrum and N-H assignments. B. Human Ubiquitin-like systems activating

enzymes ubiquitin fold domain sequential and secondary structure alignment C. Superposition of the 20

lowest-energy NMR structures of the Uba7 ubiquitin fold domain

Figure 3. Mapping of interactions between UbcH8 and Uba7-UFD at the amino acid level and NMR data

driven modeling of the Uba7-UbcH8 complex structure A. ¹⁵N-Uba7-UFD vs UbcH8 titration Chemical

Shift Perturbation graph ¹H-¹⁵N chemical shift perturbations for Uba7-UFD were calculated for each

residue. The residues colored orange possessed chemical shift perturbations larger than 0.4 ppm threshold

(red line). The residue colored merigold possessed chemical shift perturbations larger than the 0.3 ppm

threshold (merigold line). B. ¹⁵N-Uba7-UFD vs UbcH8 titration Chemical Shift Perturbation map Residues

with chemical shift perturbations larger than the 0.4 ppm threshold (red), residues with chemical shift

perturbations larger than 0.3 ppm threshold (merigold) and peak intensity changes that were larger than the

threshold (blue) were mapped onto the Uba7-UFD structure C. Surface electrostatic potential of Uba7-UFD

Red and blue coloured regions denote negative and positive charges, respectively. D. ¹⁵N-UbcH8 vs Uba7-

UFD Titration Chemical Shift Perturbation graph ¹H-¹⁵N chemical shift perturbations for UbcH8 were

calculated for each residue. The residues colored orange possessed chemical shift perturbations larger than

0.4 ppm threshold (red line). The residues colored merigold possessed chemical shift perturbations larger

than the 0.2 ppm threshold (merigold line). E. ^{15}N-UbcH8 vs Uba7-UFD Titration Chemical Shift

Perturbation map Residues with chemical shift perturbations larger than the 0.4 ppm threshold (orange),

residues with chemical shift perturbations larger than 0.2 ppm threshold (merigold) and peak intensity

changes that were larger than the threshold (blue) were mapped onto the UbcH8 crystal structure (PDB

1WZV), red residue represents the active-site cysteine (Cys85) of UbcH8 involved in transthiolation. F.

Surface electrostatic potential of UbcH8 Red and blue coloured regions denote negative and positive

charges, respectively.

Figure 4. NMR data driven model of the Uba7-UbcH8 complex structure

Figure 5. [15]N Relaxation analysis for Uba7 UFD and Uba7 UFD-UbcH8 complex A. [15]N T1 relaxation rate

as function of Uba7 UFD sequence B. [15]N T2 relaxation rate as function of Uba7 UFD sequence C. T1/T2

values of Uba7 UFD protein as function of Uba7 UFD sequence D. [1]H-[15]N NOE relaxation analysis for

Uba7 UFD E. [15]N T1 relaxation rates of [15]N-labeled Uba7 in complex with unlabeled UbcH8 F. [15]N T2

relaxation rates of [15]N-labeled Uba7 in complex with unlabeled UbcH8 G. T1/T2 relaxation values of [15]N-

labeled Uba7 UFD in complex with unlabeled UbcH8, plotted as a function of the Uba7 UFD sequence H.

[1]H-[15]N NOE relaxation analysis for Uba7 UFD in complex with unlabeled UbcH8

Figure 6. ^{15}N Relaxation analysis for UbcH8 A. ^{15}N T1 relaxation rate as function of protein sequence and

mapped on crystal structure (PDB:1WZW) B. ^{15}N T2 relaxation rate as function of protein sequence and

mapped on crystal structure (PDB:1WZW) C. ^{1}H-^{15}N NOE relaxation analysis for UbcH8 and mapped on

crystal structure (PDB:1WZW) D. T1/T2 values of UbcH8 protein as function of protein sequence

Figure 7. Investigation of the Effects of Mutations on ^{1}H-^{15}N Chemical Shift Values of Uba7-UFD A. ^{15}N-
Uba7-UFD WT vs ^{15}N-Uba7-UFD C996A Delta Chemical Shift Map The residues colored orange
possessed delta chemical shift larger than 0.4 ppm threshold (red line). The residues colored merigold
possessed delta chemical shift larger than the 0.2 ppm threshold (merigold line). B.^{15}N-Uba7-UFD WT vs
^{15}N-Uba7-UFD C996A significant Delta Chemical Shifts Mapped on protein structure Residues with delta
chemical shift larger than 0.4 ppm threshold (orange), residues with chemical shift perturbations larger than
0.2 ppm threshold (merigold) were mapped onto the Uba7-UFD structure C. ^{15}N-Uba7-UFD WT vs ^{15}N-
Uba7-UFD C996S Delta Chemical Shift Map The residues colored orange possessed delta chemical shift

larger than 1 ppm threshold (red line). The residues colored merigold possessed delta chemical shift larger

than the 0.5 ppm threshold (merigold line). D. ^{15}N-Uba7-UFD WT vs ^{15}N-Uba7-UFD C996S significant

Delta Chemical Shifts Mapped on protein structure Residues with delta chemical shift larger than 1 ppm

threshold (orange), residues with chemical shift perturbations larger than 0.5 ppm threshold (merigold)

were mapped onto the Uba7-UFD structure E. ^{15}N-Uba7-UFD A1004C vs UbcH8 titration Chemical Shift

Perturbation Map ^{1}H-^{15}N chemical shift perturbations for Uba7-UFD were calculated for each residue. The

residues colored orange possessed chemical shift perturbations larger than 0.1 ppm threshold (red line). The

residues colored merigold possessed chemical shift perturbations larger than the 0.06 ppm threshold

(merigold line). F. ^{15}N-Uba7-UFD A1004C significant Chemical Shift Perturbations Mapped on protein

structure Residues with chemical shift perturbations larger than the 0.1 ppm threshold (orange), residues

with chemical shift perturbations larger than 0.06 ppm threshold (merigold) and peak intensity changes that

were larger than the threshold (blue) were mapped onto the Uba7-UFD structure

800   Supplementary Figure 1. Human Uba7 enzyme ubiquitin fold domain [1]H-[15]N HSQC spectrum A. [1]H-[15]N

801   HSQC spectrum of Uba7-UFD after GBI fusion protein cut. B. [1]H-[15]N HSQC spectrum of GBI-Uba7-UFD

802   C. [1]H-[15]N HSQC spectrum of GBI-Uba7-UFD overlaid with [1]H-[15]N HSQC spectrum of Uba7-UFD after

803   GBI fusion protein cut D. Chemical shift perturbations (CSPs) analysis of Uba7-UFD with and without the

804   GB1 fusion tag.

810   Supplementary Figure 2. ^{1}H-^{15}N HSQC spectrum of 0.4 mM ^{15}N-GBI-Uba7-UFD (wt) protein (yellow on

811   day 1, red on day 9)

Supplementary Figure 3. Overlay of ^{15}N-GBI-Uba7-UFD ^{1}H-^{15}N HSQC spectra at different temperatures.

(green 25°C, black 20°C, dark gray 15°C, light gray 10°C)

823   Supplementary Figure 4. NMR chemical-shift-based TALOS-N secondary-structure prediction of Uba7-

824   UFD

Supplementary Figure 5. Superposition of the ^{1}H-^{15}N HSQC spectra of free (black) and various amounts of
UbcH8 bound forms of wild type ^{15}N-GBI-Uba7-UFD. GBI-Uba7-UFD:UbcH8 molar ratios in the bound
forms are indicated in yellow for 1:0.25, red for 1:0.5, blue for 1:1, and green for 1:1.75. Peaks that
disappear upon addition of 1:0.25 molar ratio of UbcH8 are indicated with red circles. Residues that show
combined chemical shift perturbations of larger than 0.2 are indicated with an arrow and black one letter
code followed by the residue number.

840 Supplementary Figure 6. ^{15}N-Uba7-UFD vs UbcH8 titration Delta Chemical Shift Intensity Graph ^{1}H-^{15}N

841 HSQC peak intensity changes for Uba7-UFD were calculated for each residue. The residues colored blue

842 possessed peak intensity changes that were larger than the threshold (blue line)

854   Supplementary Figure 7. ^{15}N-UbcH8 vs Uba7-UFD titration Delta Chemical Shift Intensity Graph ^{1}H-^{15}N

855   HSQC peak intensity changes for UbcH8 were calculated for each residue. The residues colored blue

856   possessed peak intensity changes that were larger than the threshold (blue line)

863   Supplementary Figure 8. Sequence alignment of UbcH8 from different organisms.

868   Supplementary Figure 9. Superposition of 2D ^{1}H-^{15}N HSQCs of wt ^{15}N-GBI-Uba7-UFD (black) with,

869   C996A mutant (green).

876   Supplementary Figure 10. Superposition of the ^{1}H-^{15}N HSQC spectra of free (black) and various amounts

877   of UbcH8 bound forms of ^{15}N-GBI-Uba7-UFD, C996A mutant. GBI-Uba7-UFD:UbcH8 molar ratios in

878   the bound forms are indicated in yellow for 1:0.25, red for 1:0.5, blue for 1:1, and green for 1:1.5.

885 Supplementary Figure 11. Superposition of 2D ^{1}H-^{15}N HSQCs of wt ^{15}N-GBI-Uba7-UFD (black) with,

886 C996S mutant (green).

891   Supplementary Figure 12. Superposition of the ^{1}H-^{15}N HSQC spectra of free (black) and various amounts

892   of UbcH8 bound forms of ^{15}N-GBI-Uba7-UFD, C996S mutant. GBI-Uba7-UFD:UbcH8 molar ratios in the

893   bound forms are indicated in yellow for 1:0.25, red for 1:0.5, blue for 1:1, and green for 1:1.5.

C   **Peak 1**   **Peak 2**

Supplementary Figure 13. ^{1}H-^{15}N HSQC spectra of free form of R944L mutant ^{15}N-GBI-Uba7-UFD. A.

^{1}H-^{15}N HSQC spectrum from first gel filtration peak. B. ^{1}H-^{15}N HSQC spectrum from second gel filtration

peak. C. S75 GF chromatogram

Supplementary Figure 14. Superposition of the ^{1}H-^{15}N HSQC spectra of free (black) and various amounts

of UbcH8 bound forms of ^{15}N-GBI-Uba7-UFD, A1004C mutant. GBI-Uba7-UFD:UbcH8 molar ratios in

the bound forms are indicated in yellow for 1:0.25, red for 1:0.5.

A

B

921 Supplementary Figure 15. 2D NMR analysis of Uba7-UFD 998GGG mutant A. ^{1}H-^{15}N HSQC spectra of

922 free form of 998GGG mutant ^{15}N-GBI-Uba7-UFD B. Superposition of the ^{1}H-^{15}N HSQC spectra of free

923 (red) and 1:1 ratio of UbcH8 bound form of ^{15}N-GBI-Uba7-UFD, 998GGG mutant (Green).

Distance and dihedral restraints	
Short range (|i-j| ≤ 1)	224
Medium range (|i-j| < 5)	319
Long range (|i-j| > 5)	358
Dihedral restraints (φ/ψ)	
Dihedral restrains	153
Total number of restricting constraints	1616
Average RMSD (Å) against the lowest energy model for ordered residues (PSVS)	
Backbone	0.9
All heavy atoms	1.4
RMSD from ideal geometry	
Bond lengths (Å)	0.015
Bond angles (°)	1.4
Consistent violations	
Distance constraints (>0.5 Å)	0
Dihedral angle constraints (>5°)	14.6
Van der Waals constraints (>0.2 Å)	1.8
Ramachandran analysis	
Favoured regions (%)	93.7
Allowed regions (%)	6.2
Generously allowed	0.1
Disallowed	0

925   Table 1. Restrains and statistics for 20 best solution NMR structures of Uba7-UFD