Role of Uba7-UFD in E1-E2 ISGylation: NMR-Based Insights into E2 Binding Dynamics

Çağdaş Dağ[1,2,3,], Alp E. Kazar[1], Kerem Kahraman[1], Oktay Göcenler[1], Woonghee Lee[4], Mahil Lambert[5], Cansu D. Tozkoparan Ceylan[1], Jin Gon Shim[6], Arthur L. Haas[7], Volker Dötsch[5], Joshua J. Ziarek[6], Emine Sonay Elgin[3,*]*

[1]Nanofabrication and Nanocharacterization Center for Scientific and Technological Advanced Research (n[2]STAR), Koç University, İstanbul, Turkiye

[2]Koç University Isbank Center for Infectious Diseases (KUISCID), Koç University, Istanbul, Turkiye

[3] Department of Chemistry, College of Sciences, Muğla Sıtkı Koçman University, Muğla, 48000, Turkiye

[4]Department of Chemistry, University of Colorado Denver, Denver, CO 80204, USA

[5]Centre for Biomolecular Magnetic Resonance, Institute for Biophysical Chemistry, Goethe-University of Frankfurt/Main, Germany

[6]Department of Pharmacology, Feinberg School of Medicine, Northwestern University, 320 East Superior Avenue, Chicago, IL, 460611, USA

[7]Department of Biochemistry and Molecular Biology, LSUHSC-School of Medicine, 1901 Perdido Street, New Orleans, LA, 70112, USA

*Corresponding author: cdag@ku.edu.tr selgin@mu.edu.tr

Abstract

The ISGylation pathway, a key post-translational modification, plays a pivotal role in the innate

immune response by covalently attaching ISG15 to target proteins. This cascade involves a series

of enzymatic steps, including activation by the E1 enzyme Uba7, conjugation by the E2 enzyme

UbcH8, and ligation by an E3 ligase. Central to this process is the ubiquitin-fold domain (UFD)

of Uba7, which facilitates the transfer of ISG15 to UbcH8. However, the structural and

mechanistic details of this interaction remain poorly understood. In this work, we present the

solution NMR structure and functional analysis of the ubiquitin-fold domain of human Uba7, the

E1 enzyme in the ISGylation cascade. Through detailed NMR titration experiments and

mutational studies, we mapped the interaction surface of Uba7-UFD with UbcH8, identifying

key residues and their contributions to the E1-E2 interaction. Our results reveal that Uba7-UFD

exhibits flexibility in its free form, adopting a more ordered conformation upon binding UbcH8.

Chemical shift perturbation and mutational analysis further demonstrate the importance of

specific residues, particularly Cys996, in maintaining UFD's structural integrity and binding

capacity. Additionally, mutations designed to alter the flexibility and length of the loop region

between UFD and UbcH8 show significant effects on binding, indicating that these regions are

crucial for efficient E2 recruitment. These findings provide new insights into the mechanistic

basis of E2 enzyme selection in ISGylation and underscore the functional relevance of dynamic

structural transitions in E1-E2 complex formation.

Introduction

ISGylation is a post translational modification (PTM) that results in the binding of interferon-stimulated gene 15 (ISG15) to the target protein. ISG15 is a 15 kDa ubiquitin-like protein (Ubl) containing two ubiquitin-like domains connected by a short linker peptide (Haas et al., 1987). Expression of ISG15 is primarily induced by type I interferon (IFN α/β) signaling (Farrell et al., 1979), through interactions between IFN regulatory factors (IRFs) and interferon-stimulated response element (ISRE)-containing promoters (Sadler and Williams, 2018). Innate immune response on a molecular level is generated via host pattern-recognition receptors and expression of IFN-I through downstream signaling. In this regard, ISG15 expression coincides with initiation of the innate immune response which serves as the first line of defense against pathogens. After expression, ISG15 takes part in innate immune response either as an intracellular regulatory protein or through ISGylation of primarily pathogen proteins (Zhang et al., 2021). In the context of the innate immune response, effectiveness of ISGylation against pathogens varies between species. For instance, ISGylation in mice is shown to be a highly effective antiviral mechanism for inhibiting viral assembly slowing down pathogenesis (Durfee et al., 2010). ISGylation of viral proteins also interferes with localization, host-viral protein interactions, and protease activity thereby strengthening the antiviral response (Rahnefeld et al., 2014). Whereas in humans, inherited ISG15 deficiency only results in susceptibility to mycobacterial, not viral diseases indicating differing effectiveness against pathogens and possible differences in ISGylation pathway between species (Bogunovic et al., 2012).

Similar to ubiquitin, ISG15 is also covalently attached via its C-terminal LRLRGG motif to a lysine residue on the target protein through an analogous E1-E2-E3 enzymatic cascade (Narasimhan, Potter and Haas, 1996). ISGylation enzymatic cascade includes an activating E1

enzyme (Uba7/Ube1L), a conjugating E2 enzyme (UbcH8), and an E3 ligase (HERC5) which

their expression is simultaneously induced with ISG15 through ISRE containing elements.

ISGylation begins with an ATP dependent reaction of sequential adenylation at the C-terminal

motif "LRLRGG" generating an AMP-ISG15 intermediate which forms a thioester bond with the

catalytic cysteine of the E1 activation enzyme, Uba7. In this step, active site cysteine half

domains, the first catalytic cysteine half-domain (FCCH) and the second catalytic cysteine half-

domain (SSCH) of the E1 form a thioester bond with the adenylated C-terminal glycine of

ISG15. The ubiquitin folding domain (UFD) of Uba7 functions as the recognizing agent of the

E2 enzyme and ISG15 where ISG15 is transferred to the catalytic cysteine of the E2 conjugating

enzyme, UbcH8, through a transthioesterification reaction. E2 functions as a bridge between the

Uba7 (E1), and HERC5 (E3), enzymes containing catalytic domain interaction sites of both E1

and E3. In the subsequent reaction, ISG15 is transferred to the catalytic cysteine of E3 through

another transthioesterification reaction then ligated to a lysine residue of the target protein in a

nonspecific manner (Haas, 2007). Despite its similarities, ISGylation differs considerably from

ubiquitylation in terms of variability and prevalence in the cell. Unlike its counterpart,

ISGylation cascade contains only one set of E1-E2 enzymes and therefore regulation of

ISGylation is further influenced through intrinsic interactions between enzymes of the cascade in

addition to external pathways regulating the ISGylation.

As the cornerstone of Ub/Ubl conjugation pathways, E1 enzymes are responsible for initiating

the activation of the corresponding Ub/Ubl along with its transfer to the relevant E2 enzymes.

This crucial function helps to ensure the accuracy and reliability of Ub/Ubl-mediated conjugation

pathways (Capili and Lima 2007, Streich and Haas, 2010) highlighting the crucial role that E1

enzymes play in maintaining the integrity of ubiquitylation pathways. Previous studies have

managed to identify crystal structures of Ub E1 along with E1s for Ub-like modifiers (Ubls) such

as SUMO and Nedd8 (Figure 1), both singularly and in complex states with their corresponding

Ubls (Olsen and Lima, 2013; Huang et al., 2005, Lois and Lima, 2005). Structures of the SUMO E1

provide mechanistic insights into SUMO activation and E2 recruitment to E1. *The EMBO journal*, *24*(3),

439-451.). A discovery from these studies was the revealing of the multidomain structure

inherent to canonical E1s. The domains include a pseudodimeric adenylation domain which

plays a crucial role in Ubl activation, a Cys domain which contains the catalytic cysteine residue

and a ubiquitin-fold domain (UFD) involved in the recruitment of E2 (Figure 1). Furthermore,

these prior structural studies have unveiled the considerable conformational shifts E1s undergo to

execute their roles effectively. In light of conformational changes revealing the UFD, a hidden

E2 binding area on the E1 is exposed, assisting in facilitating contact between the Ubl thioester

and E2 (Olsen and Lima, 2013). Upon the Ubl thioester's transfer from E1 to E2, it is proposed

that the UFD reverts to its originally locked conformation. This switch aids in the release of the

E2-Ub thioester product through a spatial mechanism (Olsen and Lima, 2013). Similarly, E1-E2

interaction during ISGylation marks a crucial regulatory step of the ISGylation pathway however

while considerable insights have been attained regarding the structural insights into E1 activity,

the specific process at the atomic level has yet to be determined of ISGylation pathway. The

importance of flexible regions and flexibility in protein-protein interactions has become

increasingly evident with the advancement of structural methods. In this context, "disorder-to-

order transition" in protein binding regions is being extensively studied using NMR spectroscopy

and plays a crucial role in elucidating protein-protein interactions (Jubb, Blundell and Ascher,

2015). Specific mechanistic and interactive characteristics of ISGylation are still poorly

understood.

In this study, we explore the NMR structure of Uba7-UFD, interaction dynamics and the critical

role of the disorder-to-order transition within the E1-E2 complex of the ISGylation cascade at the

residual level using NMR spectroscopy. The structural flexibility of Uba7-UFD in its unbound

state is essential for its function, transitioning into a more ordered conformation upon interaction

with UbcH8. This disorder-to-order transition is crucial for the specific recognition and selection

of the E2 enzyme, facilitating the efficient transfer of ISG15. Through detailed NMR titration

experiments, we mapped the extensive interaction surfaces between Uba7-UFD and UbcH8,

identifying over 30 residues involved in this dynamic binding process. The study also

highlighted the significance of electrostatic complementarities between the interacting regions of

Uba7-UFD and UbcH8, which contribute to their stable complex formation. Furthermore,

mutational analysis underscored the importance of key residues, such as cysteine 996, in

maintaining the structural integrity and binding capacity of Uba7-UFD. These findings provide

novel insights into the conformational plasticity of E1 enzymes and their regulatory role in

ISGylation, paving the way for a deeper understanding of this post-translational modification

process at the molecular level.

Materials and Methods

Construction of Uba7-UFD expression plasmid

The DNA sequence coding for Uba7-UFD (amino acid residues 903-1011) was amplified from

pGEX-full length HsUba7 by polymerase chain reaction (PCR). Taq polymerase (Thermo) and

primers 5' Uba7-UFD-BglII (5'-GCGCAGATCTCAGACGTTCCATCACCTG-3') and 3' Uba7-

UFD-HindIII 5'-GCGCAAGCTTACAGCTCATAGTGCAGAGG-3' were used in the reaction.

Amplified PCR product was cleaned up using Omega PCR clean-up kit and digested with BglII

and HindIII while the pQE30-GBI expression vector was digested with BamHI and HindIII. The

digested PCR product and plasmid ligated with T4 DNA ligase. Accuracy of the cloned gene was

confirmed by DNA sequencing (Macrogen, Korea) using reverse and forward pQE30 primers.

Recombinant pQE-30-GBI-Uba7-UFD plasmid transformed into the chemically competent

SG13009 expression strain of *E. coli* using heat shock.

Site-Directed mutagenesis to Uba7-UFD

For site-directed mutagenesis PCR reaction, Pfu polymerase was mixed with template pQE-GBI-

Ube1-UFD plasmid and forward and reverse primers containing the desired mutation. Primer

names and sequences used for mutations are given below.

5' C996A 5' GTGCTAGAGCTGAGCGCTGAGGGTGACGACG 3'

3' C996A 5' CGTCGTCACCCTCAGCGCTCAGCTCTAGCAC 3'

5' C996S 5' GTGCTAGAGCTGAGCAGCGAGGGTGACGACG 3'

3' C996S 5' CGTCGTCACCCTCTCGGCTCAGCTCTAGCAC 3'

5' A1004C 5' GGGTGACGACGAGGACACTTGCTTCCCACCTCTGCACTATG 3'

3' A1004C 5' CATAGTGCAGAGGTGGGAAGCAAGTGTCCTCGTCGTCACCC 3'

Once completed, PCR reaction mixture was incubated with 10 U DpnI restriction enzyme for 2

hours at 37 °C to digest the template plasmid. Resulting reaction mixture containing amplified

mutant plasmid was transformed into Novablue competent *E. coli* cells. The existence of the

desired mutation was confirmed by DNA sequencing. Plasmids containing target mutation were

selected and transformed into competent SG13009 (pREP4) *E. coli* cells. Plasmids encoding

mutant R944A and 998GGG proteins were purchased from GENSCRIPT.

Expression of recombinant GBI-Uba7-UFD fusion protein in *E. Coli*

The freshly-streaked single colony of SG13009 strain of *E. coli* containing PQE30-GBI-Uba7-

UFD plasmid was inoculated in 10 mL Luria Broth (LB) media containing 150 μg/mL ampicillin

and 50 μg/mL kanamycin. Inoculated culture was grown overnight with shaking at 250 rpm at 37

°C. Next morning, 10 mL overnight culture was added to 1 L fresh LB media containing the

same concentrations of antibiotics and grown at 37 °C and 250 rpm until OD_{600} reached 0.8. The

protein expression was induced with 1 mM IPTG. After induction, cells were grown for 5 more

hours at 37 °C. Cells were then centrifuged at 4500 x g for 25 minutes at 4 °C and stored at – 80

°C until protein purification.

Refolding and purification of fusion protein with affinity chromatography

For purification of GBI-Uba7-UFD fusion protein, cell paste was resuspended in 30 mL Lysis

buffer containing 50 mM Tris (pH 7.5), 150 mM NaCl, 5 mM DTT and sonicated at 4 °C, then

centrifuged at 15000 x g for 60 minutes at 4 °C. Proteins in pellet and supernatant were checked

by SDS-PAGE and supernatant were discarded. Inclusion bodies in pellet were washed two

times with 50 mL 0.5% Triton X-100 in 50 mM Tris (pH 7.5), 150 mM NaCl to solubilize

membranes and membrane proteins and then centrifuged at 15000 x g for 60 minutes at 4 °C.

Cleaned inclusion bodies were solubilized in 100 mL denaturation buffer containing 7 M

Guanidium-HCl, 50 mM Tris, 150 mM NaCl, 10 mM DTT overnight with stirring at 20 °C. The

next day, the mixture was centrifuged at 15000 x g for 60 minutes at 15 °C to separate insoluble

components from the solution. Refolding of fusion protein was achieved by dialysis. For this,

supernatant was transferred into a dialysis membrane (MWCO value 3500) and dialyzed twice

against 4 liters of 50 mM Tris, 150 mM NaCl, 10 mM DTT pH 7.5 buffer for 8 hours at 15 °C.

After second dialysis, precipitated components were removed by centrifugation at 15000 x g for

60 minutes at 15 °C.

Refolded fusion protein was purified by immobilized metal ion affinity (IMAC) chromatography

using a 5 mL HisTrap HP column (GE HealthSciences) precharged with nickel and attached on

an Akta Prime Plus protein purification system. The dialyzed sample was bound to a HisTrap HP

column at a flow rate of 3 mL/min. Unbound proteins were washed away with a buffer

containing 50 mM Tris-HCl, 150 mM NaCl, 5 mM DTT, 20 mM imidazole pH 7.5 until the

absorbance at 280 nm was below 10 mAbs. Elution of HisTagged fusion protein was achieved by

linearly increasing the amount of elution buffer (50 mM Tris, 150 mM NaCl, 5 mM DTT, 500

mM Imidazole pH 7.5) from 0 to 100 % in 30 mL volume. All fractions were analyzed by SDS-

PAGE and those containing the purified GBI-Uba7-UFD fusion protein were pooled,

concentrated and desalted with 20 mM phosphate Buffer pH: 7.4 to 200 μM volume in a pressure

concentrator (Stirred cell, Amicon) using a membrane with 3500 Da MWCO.

Isotopically labeled protein expression

For 1 L of M9 minimal media 200 mL sterile 5x M9 salts lacking ^{15}N-NH$_4$Cl (64 g Na$_2$HPO$_4$

7H$_2$O, 15 g KH$_2$PO$_4$, 2.5 g NaCl), were diluted to 950 mL with dd H$_2$O. 1 g ^{15}N-NH$_4$Cl

(dissolved in 10 mL ddH$_2$O and filter sterilized), 2 mL sterile 1 M MgSO$_4$, 20 mL 20% filter

sterilized glucose (for double labeled protein samples 2 g/L ^{13}C glucose), and 0.1 mL sterile 1M

CaCl$_2$ was added to M9 salts and the volume was completed to 1 L with sterile ddH$_2$O. Media

was then supplemented with trace metals and vitamins.

Freshly transformed cells were used to inoculate 20 mL of LB media containing ampicillin and

kanamycin. Cultures were grown overnight at 37 °C, 250 rpm. Cells were pelleted the next day

by centrifugation at 1000 x g for 10 min at 4 °C, resuspended in 10 mL of M9 minimal media

and added to the 1L of either ^{15}N-labeled M9 culture for ^{15}N-labeled protein production or ^{15}N-

^{13}C labeled M9 culture for double-labeled protein production. Protein expression was carried out

as described above with the exception that after induction with 0.5 mM IPTG, cells were

incubated at 25 °C for 15 hours to allow for expression of target proteins. Cells were then

harvested by centrifugation at 4000 x g for 30 minutes. Cell paste was kept at -20 °C until further

processing.

NMR data collection

2D ^{1}H-^{15}N HSQC, 3D HNCA, 3D SE HNCO, 3D SE HNCACB, 3D SE HN(CO)CA, 3D SE

CCONH, 3D SE HCCONH, 3D SE HBHACONH, 3D HNCACO, 3D SE HNCOCACB, 3D

HCCH-TOCSY 3D SE ^{15}N-NOESY-HSQC, 3D SE ^{13}C-NOESY-HSQC-aliphatics and 3D ^{13}C-

NOESY-HSQC-aromatics were performed at 20 ^{0}C on Bruker Avance II 600 MHz spectrometer

with a triple resonance z-axis gradient CryoProbe. All NMR samples were prepared in 20 mM

sodium phosphate, pH ?? containing 5-10% D_2O and 0.05% sodium azide.

NMR data processing, analysis and structure calculation

All NMR data were processed using the NMRPipe software package. The analysis of NMR data

was carried out with the XEASY software. The chemical shift index (CSI) program package was

used to identify secondary structure elements from chemical shift values. The ^{1}Hα, ^{13}Cα, ^{13}Cβ,

^{13}C', and ^{15}N chemical shift values were converted into angle restraints using the TALOS+

program. HSQC cross-peak movements were quantified as a combined ^{1}H-^{15}N chemical shift

perturbation calculated from $\Delta\delta AV = [(\Delta\delta\ ^1H\ *5)^2 + (\Delta\delta\ ^{15}N)^2]^{1/2}$ equation. The structure

calculations were performed using POKY (Lee et al., 2021) programs on a computer running the

MacOS operating system.

Docking Analysis

The docking process in HADDOCK2.4 docking protocol (de Vries et al., 2010) is divided into

three stages. In the first stage, the proteins are treated as rigid bodies, and their interaction energy

is calculated for different orientations. The orientations with the lowest energy are then selected

for further refinement. In the second stage, simulated annealing is used to refine the interface

between the two proteins. Simulated annealing is a Monte Carlo method that allows the system

to escape from local energy minima. In the context of protein-protein docking, simulated

annealing allows the proteins to explore different conformations at the interface and identify the

conformation with the most favorable energy. In the final stage, the complex is immersed in

solvent to improve the energetics. The solvent molecules are explicitly included in the

simulation, and their interactions with the proteins are taken into account. This stage helps to

refine the structure of the complex and ensure that it is stable in a solvated environment. By

combining these three stages, HADDOCK2.4 outputs a prediction of the structure of protein-

protein complex.

Results

Backbone and side-chain resonance assignments of native human Uba7-UFD

Uba7-UFD residues 903-1011 were recombinantly produced with an N-terminal GBI fusion in

E. coli cells and purified using Ni^{2+} affinity chromatography. Despite efforts to produce a soluble

construct, the GBI-Uba7-UFD expressed into inclusion bodies similar to our previous study with

Uba3-UFD (Elgin et al., 2012). The most effective approach to refold GBI-Uba7-UFD was

denaturation with 7 M guanidinium chloride and 5 mM DTT followed by dialysis into a solution

containing 50 mM Tris, 150 mM NaCl, 5 mM DTT, pH 7.5. Unfortunately, Uba7-UFD

possessed very low stability/solubility once the GBI solubility tag was removed resulting in

precipitation within ~4 h (data not shown). We next evaluated the possibility of using GBI-

Uba7-UFD fusion for NMR structure and binding studies. Comparison of 2D ^{1}H-^{15}N HSQC

spectra for both the free domain and the GBI fusion showed no detectable difference in the

amide resonances demonstrating the fusion does not affect the Uba7-UFD tertiary structure

(Supp Fig. 1). Comparison of ^{1}H-^{15}N HSQC spectra collected before (day 1) and after (day 9) a

suite of 3D assignment experiments reveals no significant time-dependent changes (Supp Fig. 2).

Therefore, GBI-Uba7-UFD was found to be suitable for further structure data collection and

titration experiments.

The chemical shift dispersion of [U-^{15}N]-GBI-Uba7-UFD amide resonances are consistent with a

well-folded protein (Fig. 2A); however, only 71 of the 100 expected backbone N-H resonances

were observed (109 total residues including eight prolines). These were assigned to residues 919-

1011 using a suite of conventional 3D triple-resonance spectra from which secondary structure

was predicted using TALOS+ (Shen et al., 2009) (Fig. 2A). A stretch of seven residues following

the third beta-strand were unassignable. The homologous region of the two E1 enzyme UFDs,

Uba3 and Uba2 (Wang et al., 2009; Elgin et al., 2012), were similarly unassigned due to low

signal intensity in 3D experiments which suggests these missing peaks result from inherent

plasticity of these regions (Fig. 2B).

NMR Structure determination of Human Uba7-UFD

The three-dimensional structure of the human Uba7-UFD was successfully determined using

multi-dimensional NMR spectroscopy (Fig. 2C). A total of 1616 unique NOE distance and 153

(phi/psi) dihedral angle constraints are used in the final structure calculations; refinement

statistics for the ensemble of 20 lowest energy conformers are shown in Table 1. The final

ensemble exhibited good convergence with an average pairwise root mean square deviation

(RMSD) of 0.9 Å for backbone atoms and 1.4 Å for all heavy atoms. The secondary structure

consists of three α-helices and four β-strands arranged in a ubiquitin fold domain (Fig. 2B,C;

Supp Fig. 4) consistent with previously reported ubiquitin-like domains (Wang et al., 2010; Lv et

at., 2018). After the second β-strand, a partially folded intermediate third β-strand is present,

with distinct loops located before and after it. These loops and β-strand may play a role in the

interaction with the E2 enzyme in the ISGylation cascade, similar to the mechanism observed in

ubiquitination (Lv et al., 2018).

The crystal structures of apo E1 enzymes (PDB:7ZH9, 6CWZ, 7PVB) lack electron density in

certain portions of the UFDs (Walden, Podgorski, and Schulman, 2003) (Figure 1). This is

consistent with NMR analyses that the UFDs contain intrinsically disordered regions in their free

form (Wang et al., 2009; Elgin et al., 2012) that adopt stable secondary structures upon E1-E2

complexation (Huang et al., 2005; Wang et al., 2010) (Figure 1). Comparison of NMR structures

of UFD's from Uba7 and Uba3 shows that Uba7-UFD NMR structure shows a well-folded

second alpha helix, which is largely unfolded at the C-terminal in Uba3-UFD (PDB 2LQ7; Fig.

1J-K,2B). In contrast, the third beta strand of Uba7-UFD, which is well defined in Uba3-UFD, is

relatively flexible (Fig. 1H-K). In addition, the NMR-derived structure showed that the acidic

loop region located between the fourth and fifth beta strands also exhibits much greater

flexibility compared to the corresponding loop region in Uba3-UFD (Fig. 1H-K). Although the

overall ubiquitin fold structure is conserved in both Uba7- and Uba3-UFDs, NMR ensembles

clearly demonstrate that the location of conformational flexibility is quite unique to each domain.

Chemical shift mapping reveals the Uba7-UbcH8 interface

To test the hypothesis that E1-UFD conformational disorder plays a role in E2

binding/selection/specificity, we performed an NMR titration to map UbcH8 onto the Uba7-UFD

surface. ^{1}H-^{15}N HSQC spectra were collected on [U-^{15}N]-Uba7-UFD mixed with increasing

concentrations of unlabeled-UbcH8 in molar ratios of 1:0.25, 1:0.5, 1:1 and 1:1.75 (Supp Fig. 5).

Inspection of these spectra revealed a subset of peaks that were dose-dependently perturbed upon

addition of UbcH8 (Figure 3A); combined ^{1}H/^{15}N chemical shift changes were calculated and the

most perturbed residues were highlighted onto the Uba7-UFD structure (Figure 3B,3C). UbcH8

binding surface of Uba7-UFD appears to be quite large, involving about 18 residues.

Interestingly, some of the peaks displaying the largest chemical shift perturbations upon addition

of UbcH8 were weaker in intensity compared to the average peak intensities in the spectrum of

the free Uba7-UFD. In fact, four peaks with pronounced chemical shift perturbations

corresponded to those residues in the unassigned fragments and were absent in 3D spectra. One

of the simplest explanations for the low intensity and absence in 3D spectra is that these residues

may undergo conformational exchange on the microsecond-millisecond timescale (Supp Fig. 6).

We next mapped the binding interface onto the UbcH8 surface by titrating increasing

concentrations of Uba7-UFD into [U-^{15}N]-UbcH8. Substantial chemical shift perturbations

clustered to the UbcH8 N-terminal region, helix 1, strands 1-3, and helix 3 (Figure 3D-F).

Several peaks were also noted to disappear in the N-terminal region, helix 1, strands 1-3, and

helix 3 upon addition of unlabeled Uba7-UFD (Supp Fig. 7). This result is consistent with the

mutational study showing that the N-terminal region of UbcH8 is important in Uba7 interaction

(Huibregtse et al. 2008).

NMR data driven docking of UbcH8 and Uba7-UFD complex

Next, we used the most substantial CSPs from our NMR titrations on the isolated domain

interactions to restrain a model of the full-length Uba7-HFB/UbcH8 complex using the

HADDOCK 2.4 NMR-based docking protocol (Vries et al., 2010). The AlphaFold Uba7 model

(APSD: AF-P41226-F1-v4) and the UbcH8 crystal structure (PDB: 1WZV) were used as inputs

and the resulting cluster with the lowest HADDOCK score was analyzed (Figure 4). The

HADDOCK model illustrates how complexation is facilitated by two sets of charge-charge

interactions; specifically, Uba7-UFD ^{999}DDEDTA^{1003} and ^{950}SALLYAAGWSPEK^{962}

complement UbcH8 ^{5}MRVVKE^{10} and ^{25}LSSDD^{29}, respectively. Furthermore, a series of

hydrophobic and charged residues surrounding the active site cysteine undergoes conformational

changes during the transthiolation stage (Figure 4). The spatial arrangement of these regions

precisely orients UbcH8 to ensure efficient ISG transfer between catalytic cysteine residues.

Dynamics of E1 UFD binding surface on UbcH8

The dynamical behavior of the UbcH8 was also probed using ^{15}N T1, ^{15}N T2, and {^{1}H}-^{15}N

heteronuclear NOE relaxation experiments (Lakomek, Ying and Bax, 2013). Figures 5A and 5B

show the T1 and T2 relaxation rates plotted against the residue number and color mapped on the

crystal structure of UbcH8 (PDB:1WZW), respectively. Residues with high T1 and low T2

relaxation rates corresponding to relatively rigid regions are colored in green whereas those

displaying lower T1 and higher T2 relaxation rate hence higher flexibility are shown in yellow,

orange, and red in the order of increasing flexibility. Gray color corresponds to the unassigned

region. Figure 5C shows the heteronuclear ^{15}N-^{1}H NOE values plotted as a function of UbcH8

amino acid sequence. Fitting of data was done with the help of Bruker dynamics center and

which showed more than 95% confidence level. All together, T1 and T2 relaxation rates and

heteronuclear NOE values measured for UbcH8 consistently show that while the core of the

protein is rigid, the N-terminal alpha-helix which is highly conserved among various species

(Supp Fig. 8) exhibits increased flexibility (Figure 5D). Interestingly, our docking studies show

that this region of UbcH8 is directly involved in the interaction with Uba7-UFD. This flexibility

could be crucial for the E2's function, potentially facilitating conformational changes required for

E1-E2 transthiolation.

Conformational Flexibility of Uba7-UFD and Its Role in E2 Binding: Insights from Mutational

Analysis

NMR solution structure of free Uba7-UFD indicated that the putative E2 binding region might be

flexible and experiencing conformational plasticity, transitioning among multiple conformational

states. Comparably, conformational flexibility/heterogeneity has been observed in E2 binding

surfaces of SUMO (Wang et al., 2009) and Nedd8 E1 UFD (Elgin et al., 2012). Notably this

conformational flexibility has been shown to be functionally important in SUMO conjugation

(Wang et al., 2009).

To further explore this phenomenon, we designed and analyzed a series of mutants to determine

how these alterations would affect E2 binding to Uba7-UFD. Specifically, we aimed to identify

the role of individual amino acids in maintaining the structural integrity and interaction

capabilities of the UFD domain. Given the high degree of plasticity observed in the putative E2

binding region, understanding how mutations influence this flexibility is crucial. This

investigation not only provides insight into the specific amino acid interactions that are critical

for binding but also sheds light on the overall mechanism by which Uba7-UFD achieves its

conformational adaptability. By mapping these interactions and comparing the behavior of wild-

type and mutant proteins, we sought to delineate the structural elements that enable effective

binding and subsequent functional outcomes.

Role C996 in Uba7-UFD conformational integrity and UbcH8 binding

In order to determine the effect of mutations on E2 binding to Uba7-UFD, seven mutants were

created and five of them also titrated with UbcH8. Full length Uba7 protein contains a total of 17

cysteine residues, only one of which is in the UFD. Since this domain does not contain any other

cysteine residue to form disulfide bridge with C996, this single cysteine probably remains

reduced. C996 is located in the end of the second beta sheet, and while the subsequent amino

acids show a very flexible behavior in the structure, cysteine and previous amino acids have

formed a stable beta sheet secondary structure form.

We first mutated the free cysteine to alanine, which is the most preferred amino acid in mutation

scans. Overlay of ^{1}H-^{15}N HSQC spectra of wild-type and C996A variant shows that the overall

conformations of the mutant and wild-type proteins remain largely the same (Supp Figure 9). In

contrast, changes in chemical shift values were observed around the mutated region, indicating a

systematic reorganization in the preceding and subsequent beta sheets (Figure 6A). Figure 5B

shows the residues affected by C996A mutation mapped onto the Uba7-UFD structure. Residues

with chemical shift perturbations larger than 0.4 and 0.2 ppm colored in orange and merigold,

respectively, are mostly clustered in neighboring B1, and B4 strands and alpha 1 helix. In order

to determine the effect of mutation on E2 binding to Uba7-UFD, mutant protein was titrated with

UbcH8. Supp Figure 10 shows the overlay of ^{1}H-^{15}N HSQC spectra of Uba7-UFD-C996A before

and after addition of various amounts of UbcH8. No significant ^{1}H-^{15}N chemical shift

perturbations in C996A mutant upon addition of UbcH8 was observed and addition of more and

more of UbcH8 resulted in disappearance of some of the peaks involved in binding. These

titration studies clearly indicate that C996 is important for Uba7-UFD-UbcH8 interaction (Supp

Figure 10).

Mutation of C996 to alanine appeared to have effect on UbcH8 binding. Since mutation of

cysteine to alanine results in truncation of the side chain at the C-beta, the goal of C996S

mutation was to replace –SH group with –OH group to see if this would restore the UbcH8

binding. The comparison of ^{1}H-^{15}N HSQC spectrum of C996S with that of wild-type shows that

the overall conformation of C996S mutant is very similar to that of wild-type (Supp Figure 11)

and as in C996A mutant, significant chemical shift perturbations are located around the mutation

site (Figure 6C-6D). NMR binding studies carried out using ^{15}N labeled Ubc7-UFD C996S

further confirmed the role of C996 in UbcH8 binding since there were no significant chemical

shift perturbations in the ^{1}H-^{15}N HSQC spectra of C996S mutant upon addition of UbcH8(Supp

Figure 12). The failure of the -OH group in restoring the UbcH8 binding suggests that free -SH

might not merely be involved in hydrogen bonding but might be important in providing

necessary flexibility for UbcH8 interaction (Bhopatkar et al., 2020).

After determining that free cysteine was necessary for UbcH8 binding/for keeping Uba7-UFD in

a suitable conformation for the UbcH8 binding we searched for amino acids in close proximity to

C996 within Uba7-UFD to test the effect of breaking C996 intermolecular interactions on the

conformation of the domain as well as on UbcH8 binding. Amino acids directly interacting with

C996 were identified through NOE analysis. NOE peaks observed between C996 and R944

indicated that there is an interaction between these two amino acids. Therefore, an R944A

mutant protein was generated, produced, and purified after refolding. Gel filtration

chromatography of purified ^{15}N-GBI-Uba7-UFD resulted in two separate peaks with the same

molecular weight confirmed by SDS-PAGE analysis. Peaks were collected in separate fractions

and analyzed by NMR. ^{1}H-^{15}N HSQC spectra showed that the peak eluted first from the gel

filtration column was completely misfolded/aggregated (Supp Figure 13A) while the fraction

corresponding to the second peak consisted of a mixture of both folded and misfolded fusion

protein (Supp Figure 13B). Gel filtration chromatography and NMR results clearly show that

R944 and its interaction with C996 are important for conformational stability of Uba7-UFD and

mutation of R944 to alanine makes domain prominent to aggregation. The role of free cysteine

amino acid in protein structure and internal dynamics is often overlooked. However, a study

conducted by Mazmanian et al. in 2016 examined the participation and significance of free

cysteine amino acids in hydrogen bonding networks in detail (Mazmanian et al., 2016). In the

case of the Uba7-UFD complex, when a cysteine atom forms a hydrogen bond with arginine

(R944), it exhibits distinct interaction patterns depending on the charge of the hydrogen bond

donor. Specifically, cysteine forms shorter and more linear hydrogen bonds with positively

charged residues such as histidine (His□), lysine (Lys□), and arginine (Arg□), where the

hydrogen–sulfur distance (H···S) is less than 2.5 Å, and the bond angle (N−H···S) exceeds 150°

(Mazmanian et al., 2016). In contrast, hydrogen bonds with neutral donors are longer (H···S > 2.5

Å) and less linear (N−H···S < 150°). Notably, Arg□ donates two hydrogen bonds to the cysteine

thiol (CysSH), a characteristic not observed in other residues (Mazmanian et al., 2016). These

findings align with previous studies demonstrating that positively charged hydrogen bond donors

preferentially form stronger and more directional interactions with thiol groups, contributing to

protein stability and conformational flexibility. Thus, the Uba7-UFD complex exemplifies the

critical role of free cysteines in fine-tuning the stability and plasticity of protein structures.

Effect of acidic loop flexibility and lenght on Uba7-UFD interaction with UbcH8

As mentioned above, our NMR solution structure of Uba7-UFD shows that the loop region that

comes right after the C996 is relatively flexible. Since this loop is involved in interaction with

UbcH8, one question is whether restricting this flexibility would prevent binding. To test this,

another mutant was designed aiming to form a disulfide bond between the single cysteine 996

and alanine 1004. In this mutant, A1004 was individually replaced by cysteine. This variant was

also expressed as inclusion bodies, solubilized with 7M GndHCl and refolded by dialysis as

described previously. Supp Figure 14 shows an overlay of ^{1}H-^{15}N HSQC spectra of A1004C

mutant in the absence and presence of various amounts of UbcH8. Figure 6E shows the chemical

shift perturbation analysis from the titration experiment. Unlike the previous mutations, this

mutant variant still showed intrinsically interacted with UbcH8. As seen in Figure 6F, the

interaction surface is also partially compatible with the interaction surface of the WT protein.

Opposite to WT, A1004C mutant's 1st beta sheet region did not show any interaction with

UbcH8.

The acidic loop interacting with UbcH8 has been extended with a glycine amino acid as a final

mutation approach (998GGG). The 998GGG mutant protein has been produced, purified, and

subjected to ^{1}H-^{15}N HSQC spectrum collection. Although there were regional differences in the

obtained spectrum, no significant changes were observed in the overall distribution (Supp Figure

15A). Unlabeled UbcH8 has been added to the obtained protein in a 1:1 ratio. Upon examination

of the spectra obtained after mixing the apo form of 998GGG with UbcH8, no changes were

observed in the chemical shift values indicating that integrity of the loop was necessary for UbcH8

binding. (Supp Figure 15B).

Discussion

Ubiquitin and Ubiquitin-like post-translational modification systems are of great importance to

organisms due to their extensive regulatory roles within cells. Elucidating the mechanisms at the

atomic level within the enzymatic cascades that form the basis of Ub and Ubl post-translational

modification systems has been a significant topic in biochemistry and structural biology for

decades. Although the atomic-level details of the mechanisms of systems such as ubiquitination,

neddylation, and sumoylation have been examined, structural studies on the ISGylation system

were not available until recent years. Specifically, due to the high plasticity of the ubiquitin fold

domain of the E1 enzyme, determining its structure has taken a considerable amount of time.

During the preparation of our manuscript, two additional papers aimed at elucidating the details

of ISGylation have been published (Wallace et al., 2023; Afsar et al., 2023). Although Cryo-EM

and crystal structures are excellent tools for visualizing the overall mechanism of systems, they

are still static and frozen structures. This makes them less effective for examining weak

interaction surfaces and low-affinity complex structures. At this point, NMR spectroscopy

continues to be an important structural biology method, providing atomic-level information to

complement the remarkable findings from other methods and to elucidate the side effects caused

by freezing. In the first part of our study, we determined the NMR structure of the ISGylation E1

enzyme and observed that the plasticity and unstable beta-sheet surfaces exhibited in solution are

believed to play a significant role in the system's operation and selectivity.

Contemporary Cryo-EM structures, especially those examining the E1-E2 interaction, present

differing results regarding the amino acids and interaction angles involved in the interaction

between the first alpha helix of the E2 enzyme and the high plasticity regions of the E1 UF

domain (Wallace et al., 2023; Afsar et al., 2023). Despite the high overall resolutions of these

structures, it is notable that the resolution in these domains sometimes drops regionally to 4-5 Å.

The direct interactions obtained from our NMR data-driven model show greater alignment with

the findings of Wallace et al. 2023.

Findings from T1 and T2 relaxation experiments, coupled with titration and docking data,

provide a coherent picture suggesting that the N terminus region of Ubch8 is more flexible and

that this flexibility is relevant to the interaction with the Uba7-UFD. NMR characterization

studies of Uba7-UFD variants showed that both wild type and mutant domains have a high

tendency to aggregate. None of the designed mutant Uba7-UFDs was more stable than the wild

type Uba7-UFD. Our NMR mapping study showed that UbcH8 binding surface of Uba7-UFD is

substantial and there are over 30 amino acids directly involved in binding. This study also

showed that a single reduced cysteine residue at position 996 may play a major role in domain

stability and fine tuning of plasticity.

Conclusion

Our study has reiterated the importance of examining protein structures and dynamics in

solution. It has been demonstrated that the UFD domain of the Uba7 protein undergoes

conformational changes upon binding to UbcH8, consistent with the induced fit model.

Structural analysis of the Uba7 UFD and complex structures obtained from other systems have

shown that a large surface area of the UFD interacts with E2, highlighting the significance of this

interaction in E1-E2 binding and proper E2 selection. Additionally, structural investigations have

revealed that a region on the E1 UFD interacts with Ubiquitin/Ubiquitin-like proteins. It is

important to conduct new studies to demonstrate the importance of this interaction in structural

and functional regulation.

Author Contributions

Çağdaş Dağ: Conceptualization, Methodology, Supervision Funding acquisition, Investigation,

Formal analysis, Writing- Reviewing and Editing, Writing-Original draft preparation **Alp Eren**

Kazar: Investigation, Formal analysis, Visualization, Writing- Reviewing and Editing, Writing-

Original draft preparation **Emine Sonay Elgin:** Conceptualization, Methodology, Supervision

Funding acquisition, Investigation, Formal analysis, Writing- Reviewing and Editing, Writing-

Original draft preparation **Arthur L. Haas:** Conceptualization, Supervision **Kerem Kahraman**:

Investigation, Formal analysis, Visualization. **Oktay Göcenler:** Formal analysis, Investigation

Writing - Original Draft, Visualization **Mahil Lambert:** Formal analysis, Investigation, Writing

- Original Draft, Visualization **Volker Dötsch**: Conceptualization, Supervision, Writing-

Reviewing and Editing **Cansu D. Tozkoparan:**Visualization, Formal analysis **Joshua J.**

Ziarek: Conceptualization, Writing- Reviewing and Editing

ACKNOWLEDGMENT

We gratefully acknowledge Prof. Brian F. Volkman (Medical College of Wisconsin) for

generously opening his laboratory to us and providing access to critical instrumentation and

consumables (GM64598-03). His support and willingness to share resources were invaluable for

the successful execution of this work. We sincerely thank him for his scientific generosity and

hospitality. ESE acknowledges support from TU☐BI☐TAK (Project No: 104T193). CD

acknowledges support from TU☐BI☐TAK (Project No: 120Z594, 122Z747). JJZ acknowledges

support from National Institutes of Health grant R35GM143054. The authors acknowledge the

use of the services and facilities of n^2STAR-Koç University Nanofabrication and

Nanocharacterization Center for Scientific and Technological Advanced Research. This study

made use of the National Magnetic Resonance Facility at Madison, which is supported by NIH

grant R24GM141526 and P41GM103399. This work benefited from access to [CERM and

BMRZ] and has been supported by iNEXT-Discovery, project number 871037, funded by the

Horizon 2020 program of the European Commission. Financial support by the Access to

Research Infrastructures activity in the 7th Framework Programme of the EC (Project number:

261863, Bio-NMR) for conducting the research is gratefully acknowledged. This study made use

of NMRbox: National Center for Biomolecular NMR Data Processing and Analysis, a

Biomedical Technology Research Resource (BTRR), which is supported by NIH grant

P41GM111135 (NIGMS).

REFERENCES

Afsar, M., Liu, G., Jia, L., Ruben, E. A., Nayak, D., Sayyad, Z., Bury, P. D. S., Cano, K. E., Nayak, A., Zhao, X. R.,

Shukla, A., Sung, P., Wasmuth, E. V., Gack, M. U., & Olsen, S. K. (2023). Cryo-EM structures of Uba7 reveal the

molecular basis for ISG15 activation and E1-E2 thioester transfer. *Nature communications*, *14*(1), 4786.

https://doi.org/10.1038/s41467-023-39780-z

Bogunovic, D., Byun, M., Durfee, L. A., Abhyankar, A., Sanal, O., Mansouri, D., Salem, S., Radovanovic, I, Grant,

A. V., Adimi, P., Mansouri, N., Okada, S., Bryant, V. L., Kong, X. F., Kreins, A., Velez, M. M., Boisson, B.,

Khalilzadeh, S., Ozcelik, U., Darazam, I. A., … Casanova, J. L. (2012). Mycobacterial disease and impaired IFN-γ

immunity in humans with inherited ISG15 deficiency. *Science (New York, N.Y.)*, *337*(6102), 1684–1688.

https://doi.org/10.1126/science.1224026

Bhopatkar AA, Uversky VN, Rangachari V. Disorder and cysteines in proteins: A design for orchestration of

conformational see-saw and modulatory functions. Prog Mol Biol Transl Sci. https://10.1016/bs.pmbts.2020.06.001.

Capili, A. D., Lima, C. D. (2007). Structure and analysis of a complex between SUMO and Ubc9 illustrates features

of a conserved E2-Ubl interaction. *Journal of molecular biology*, *369*(3), 608–618.

https://doi.org/10.1016/j.jmb.2007.04.006

Durfee, L. A., Lyon, N., Seo, K., Huibregtse, J. M. (2010). The ISG15 conjugation system broadly targets newly

synthesized proteins: implications for the antiviral function of ISG15. *Molecular cell*, *38*(5), 722–732.

https://doi.org/10.1016/j.molcel.2010.05.002

de Vries, S. J., van Dijk, M., Bonvin, A. M. (2010). The HADDOCK web server for data-driven biomolecular

docking. *Nature protocols*, *5*(5), 883–897. https://doi.org/10.1038/nprot.2010.32

Elgin, E. S., Sökmen, N., Peterson, F. C., Volkman, B. F., Dağ, C., Haas, A. L. (2012). E2-binding surface on Uba3

β-grasp domain undergoes a conformational transition. *Proteins*, *80*(10), 2482–2487.

https://doi.org/10.1002/prot.24148

Farrell, P. J., Broeze, R. J., Lengyel, P. (1979). Accumulation of an mRNA and protein in interferon-treated Ehrlich

ascites tumour cells. *Nature*, *279*(5713), 523–525. https://doi.org/10.1038/279523a0.

Haas, A.L. (2007). ISG15-dependent Regulation. In Protein Degradation Series (eds R.J. Mayer, A.J. Ciechanover

and M. Rechsteiner). https://doi.org/10.1002/9783527619320.ch5b

Huang, D. T., Paydar, A., Zhuang, M., Waddell, M. B., Holton, J. M., & Schulman, B. A. (2005). Structural basis

for recruitment of Ubc12 by an E2 binding domain in NEDD8's E1. Molecular cell, 17(3), 341-350

Jubb, H., Blundell, T. L., Ascher, D. B. (2015). Flexibility and small pockets at protein-protein interfaces: New

insights into druggability. *Progress in biophysics and molecular biology*, *119*(1), 2–9.

https://doi.org/10.1016/j.pbiomolbio.2015.01.009

Lakomek, N. A., Ying, J., Bax, A. (2012). Measurement of ^{15}N relaxation rates in perdeuterated proteins by

TROSY-based methods. *Journal of biomolecular NMR*, *53*(3), 209–221. https://doi.org/10.1007/s10858-012-9626-5

Lee, W., Rahimi, M., Lee, Y., Chiu, A. (2021). POKY: a software suite for multidimensional NMR and 3D

structure calculation of biomolecules, *Bioinformatics*, Volume 37, Issue 18, Pages 3041–3042,

https://doi.org/10.1093/bioinformatics/btab180

Loeb, K. R., Haas, A. L. (1992). The interferon-inducible 15-kDa ubiquitin homolog conjugates to intracellular

proteins. *The Journal of biological chemistry*, *267*(11), 7806–7813.

Lois, L. M., & Lima, C. D. (2005). Structures of the SUMO E1 provide mechanistic insights into SUMO

activation and E2 recruitment to E1. *The EMBO journal*, *24*(3), 439-451.

Lv, Z., Williams, K.M., Yuan, L., Atkison, JH, Olsen SK. Crystal structure of a human ubiquitin E1-ubiquitin

complex reveals conserved functional elements essential for activity. J Biol Chem. 2018 Nov 23;293(47):18337-

18352. doi: 10.1074/jbc.RA118.003975

Mazmanian, K., Sargsyan, K., Grauffel, C., Dudev, T., & Lim, C. (2016). Preferred Hydrogen-Bonding Partners of

Cysteine: Implications for Regulating Cys Functions. *The journal of physical chemistry. B*, *120*(39), 10288–10296.

https://doi.org/10.1021/acs.jpcb.6b08109

Narasimhan, J., Potter, J. L., Haas, A. L. (1996). *Conjugation of the 15-kDa Interferon-induced Ubiquitin Homolog*

Is Distinct from That of Ubiquitin. Journal of Biological Chemistry, 271(1), 324–330. doi:10.1074/jbc.271.1.324

Olsen, S. K., & Lima, C. D. (2013). Structure of a ubiquitin E1-E2 complex: insights to E1-E2 thioester transfer.

Molecular cell, 49(5), 884-896.

Poller, W., Knobeloch, K. P., Voigt, A. (2014). Ubiquitin-like protein ISG15 (interferon-stimulated gene of 15 kDa)

in host defense against heart failure in a mouse model of virus-induced cardiomyopathy. *Circulation*, *130*(18),

1589–1600. https://doi.org/10.1161/CIRCULATIONAHA.114.009847

Rahnefeld, A., Klingel, K., Schuermann, A., Diny, N. L., Althof, N., Lindner, A., … Voigt, A. (2014). *Ubiquitin-*

Like Protein ISG15 (Interferon-Stimulated Gene of 15 kDa) in Host Defense Against Heart Failure in a Mouse

Model of Virus-Induced CardiomyopathyCLINICAL PERSPECTIVE. Circulation, 130(18), 1589–1600.

doi:10.1161/circulationaha.114

Sadler, A. J., Williams, B. R. (2008). Interferon-inducible antiviral effectors. *Nature reviews. Immunology*, *8*(7),

559–568. https://doi.org/10.1038/nri2314

Shen, Y., Delaglio, F., Cornilescu, G. et al. TALOS+: a hybrid method for predicting protein backbone torsion

angles from NMR chemical shifts. J Biomol NMR 44, 213–223 (2009).

Streich, F.C., Haas, A.L. (2010). Activation of Ubiquitin and Ubiquitin-Like Proteins. In: Groettrup, M. (eds)

Conjugation and Deconjugation of Ubiquitin Family Modifiers. Subcellular Biochemistry, vol 54. Springer, New

York, NY. https://doi.org/10.1007/978-1-4419-6676-6_1

Wang, J., Taherbhoy, A. M., Hunt, H. W., Seyedin, S. N., Miller, D. W., Miller, D. J., Huang, D. T., & Schulman,

B. A. (2010). Crystal structure of UBA2(ufd)-Ubc9: insights into E1-E2 interactions in Sumo pathways. *PloS one*,

5(12), e15805. https://doi.org/10.1371/journal.pone.0015805

Zhang, M., Li, J., Yan, H., Huang, J., Wang, F., Liu, T., Zeng, L., Zhou, F. (2021). ISGylation in Innate Antiviral

Immunity and Pathogen Defense Responses: A Review. *Frontiers in cell and developmental biology*, *9*, 788410.

https://doi.org/10.3389/fcell.2021.788410

Figure 1. Ubiquitin and Ubiquitin-like post-translational modification systems E1 enzymes' ubiquitin fold

domain structures. Regions on the structures that have high plasticity and directly interact with E2 are

marked in red, and regions thought to interact with Ubl are marked in orange. Electron density maps are

also shown over the colored regions. A. Saccharomyces cerevisiae Uba1 Apo form (PDB: 7ZH9) B.

Homo sapiens Uba1 complex with ubiquitin (PDB: 6DC6) C. Saccharomyces cerevisiae complex with

Ubc13 (E2) (PDB: 6ZHS) D. Schizosaccharomyces pombe complex with Ubc15 (E2) and ubiquitin

(PDB: 5KNL) E. Homo sapiens Uba2 apo form (6CWZ) F. Homo sapiens Uba2 complex with SUMO

(3KYC) G. Saccharomyces cerevisiae Uba2-UFD complex with Ubc9 (3ONG) H. Homo sapiens Uba3

apo form (1YOV) I. Homo sapiens Uba3-UFD complex with Ubc12 (1Y8X) J. Homo sapiens Uba3 UFD

apo form (2LQ7) K. Homo sapiens Uba7-UFD apo form (8WWX) L. Homo sapiens apo Uba6 (7PVN)

M. Homo sapiens Uba6 complex with FAT10 (7PYV) N. Homo sapiens Uba6 complex with FAT10

(7SOL)

Figure 2. NMR structure of Human Uba7 ubiquitin fold domain A. Human Uba7 enzyme ubiquitin fold

domain ^{1}H-^{15}N HSQC spectrum and N-H assignments. B. Human Ubiquitin-like systems activating

enzymes ubiquitin fold domain sequential and secondary structure alignment C. Superposition of the 20

lowest-energy NMR structures of the Uba7 ubiquitin fold domain

Figure 3. Mapping of interactions between UbcH8 and Uba7-UFD at the amino acid level and NMR data

driven modeling of the Uba7-UbcH8 complex structure A. ^{15}N-Uba7-UFD vs UbcH8 titration Chemical

Shift Perturbation graph ^{1}H-^{15}N chemical shift perturbations for Uba7-UFD were calculated for each

residue. The residues colored orange possessed chemical shift perturbations larger than 0.4 ppm threshold

(orange line). The residues colored merigold possessed chemical shift perturbations larger than the 0.3

ppm threshold (merigold line). B. ^{15}N-Uba7-UFD vs UbcH8 titration Chemical Shift Perturbation map

Residues with chemical shift perturbations larger than the 0.4 ppm threshold (orange), residues with

chemical shift perturbations larger than 0.3 ppm threshold (merigold) and peak intensity changes that

were larger than the threshold (blue) were mapped onto the Uba7-UFD structure C. Surface electrostatic

potential of Uba7-UFD Red and blue coloured regions denote negative and positive charges,

respectively.D. ^{15}N-UbcH8 vs Uba7-UFD Titration Chemical Shift Perturbation graph ^{1}H-^{15}N chemical

shift perturbations for UbcH8 were calculated for each residue. The residues colored orange possessed

chemical shift perturbations larger than 0.4 ppm threshold (orange line). The residues colored merigold

possessed chemical shift perturbations larger than the 0.2 ppm threshold (merigold line). E. [15]N-UbcH8 vs

Uba7-UFD Titration Chemical Shift Perturbation map Residues with chemical shift perturbations larger

than the 0.4 ppm threshold (orange), residues with chemical shift perturbations larger than 0.2 ppm

threshold (merigold) and peak intensity changes that were larger than the threshold (blue) were mapped

onto the UbcH8 crystal structure (PDB 1WZV). F. Surface electrostatic potential of UbcH8 Red and blue

coloured regions denote negative and positive charges, respectively.

654   Figure 4. NMR data driven model of the Uba7-UbcH8 complex structure

Figure 5. [15]N Relaxation analysis for UbcH8 A. [15]N T1 relaxation rate as function of protein sequence and

mapped on crystal structure (PDB:1WZW) B. [15]N T2 relaxation rate as function of protein sequence and

mapped on crystal structure (PDB:1WZW) C. [1]H-[15]N NOE relaxation analysis for UbcH8 and mapped on

crystal structure (PDB:1WZW) D. T1/T2 values of UbcH8 protein as function of protein sequence

^{15}N-Uba7-UFD WT vs ^{15}N-Uba7-UFD C996A Delta Chemical Shift

^{15}N-Uba7-UFD WT vs ^{15}N-Uba7-UFD C996S Delta Chemical Shift

^{15}N-Uba7-UFD A1004C vs UbcH8 titration

Figure 6. Investigation of the Effects of Mutations on ^{1}H-^{15}N Chemical Shift Values of Uba7-UFD A.

^{15}N-Uba7-UFD WT vs ^{15}N-Uba7-UFD C996A Delta Chemical Shift Map The residues colored orange

possessed delta chemical shift larger than 0.4 ppm threshold (orange line). The residues colored merigold

possessed delta chemical shift larger than the 0.2 ppm threshold (merigold line). B. ^{15}N-Uba7-UFD WT vs

^{15}N-Uba7-UFD C996A significant Delta Chemical Shifts Mapped on protein structure Residues with

delta chemical shift larger than 0.4 ppm threshold (orange), residues with chemical shift perturbations

larger than 0.2 ppm threshold (merigold) were mapped onto the Uba7-UFD structure C. ^{15}N-Uba7-UFD

WT vs ^{15}N-Uba7-UFD C996S Delta Chemical Shift Map The residues colored orange possessed delta

chemical shift larger than 1 ppm threshold (orange line). The residues colored merigold possessed delta

chemical shift larger than the 0.5 ppm threshold (merigold line). D. ^{15}N-Uba7-UFD WT vs ^{15}N-Uba7-

UFD C996S significant Delta Chemical Shifts Mapped on protein structure Residues with delta chemical

shift larger than 1 ppm threshold (orange), residues with chemical shift perturbations larger than 0.5 ppm

threshold (merigold) were mapped onto the Uba7-UFD structure E. ^{15}N-Uba7-UFD A1004C vs UbcH8

titration Chemical Shift Perturbation Map ^{1}H-^{15}N chemical shift perturbations for Uba7-UFD were

calculated for each residue. The residues colored orange possessed chemical shift perturbations larger

than 0.1 ppm threshold (orange line). The residues colored merigold possessed chemical shift

perturbations larger than the 0.06 ppm threshold (merigold line). F. ^{15}N-Uba7-UFD A1004C significant

Chemical Shift Perturbations Mapped on protein structure Residues with chemical shift perturbations

larger than the 0.1 ppm threshold (orange), residues with chemical shift perturbations larger than 0.06

ppm threshold (merigold) and peak intensity changes that were larger than the threshold (blue) were

mapped onto the Uba7-UFD structure

695 Supplementary Figure 1. Human Uba7 enzyme ubiquitin fold domain ^{1}H-^{15}N HSQC spectrum A. ^{1}H-^{15}N

696 HSQC spectrum of Uba7-UFD after GBI fusion protein cut. B. ^{1}H-^{15}N HSQC spectrum of GBI-Uba7-

697 UFD C. ^{1}H-^{15}N HSQC spectrum of GBI-Uba7-UFD overlaid with ^{1}H-^{15}N HSQC spectrum of Uba7-UFD

698 after GBI fusion protein cut

710 Supplementary Figure 2. ^{1}H-^{15}N HSQC spectrum of 0.4 mM ^{15}N-GBI-Uba7-UFD (wt) protein (green on

711 day 1, red on day 9)

716   Supplementary Figure 3. Overlay of ^{15}N-GBI-Uba7-UFD ^{1}H-^{15}N HSQC spectraat different temperatures.

717   (green 25°C, black 20°C, dark gray 15°C, light gray 10°C)

723   Supplementary Figure 4. NMR chemical-shift-based TALOS-N secondary-structure prediction of Uba7-

724   UFD

Supplementary Figure 5. Superposition of the ^{1}H-^{15}N HSQC spectra of free (black) and various amounts

of UbcH8 bound forms of wild type ^{15}N-GBI-Uba7-UFD. GBI-Uba7-UFD:UbcH8 molar ratios in the

bound forms are indicated in yellow for 1:0.25, red for 1:0.5, blue for 1:1, and green for 1:1.75. Peaks that

disappear upon addition of 1:0.25 molar ratio of UbcH8 are indicated with red circles. Residues that show

combined chemical shift perturbations of larger than 0.2 are indicated with an arrow and black one letter

code followed by the residue number.

Supplementary Figure 6. ^{15}N-Uba7-UFD vs UbcH8 titration Delta Chemical Shift Intensity Graph ^{1}H-^{15}N

HSQC peak intensity changes for Uba7-UFD were calculated for each residue. The residues colored blue

possessed peak intensity changes that were larger than the threshold (blue line)

753   Supplementary Figure 7. ^{15}N-UbcH8 vs Uba7-UFD titration Delta Chemical Shift Intensity Graph ^{1}H-^{15}N

754   HSQC peak intensity changes for UbcH8 were calculated for each residue. The residues colored blue

755   possessed peak intensity changes that were larger than the threshold (blue line)

762   Supplementary Figure 8. Sequence alignment of UbcH8 from different organisms.

767 Supp Fig. 9. Superposition of 2D ^{1}H-^{15}N HSQCs of wt 15N-GBI-Uba7-UFD (black) with, C996A mutant

768 (green).

Supp Fig. 10. Superposition of the ^{1}H-^{15}N HSQC spectra of free (black) and various amounts of UbcH8

bound forms of ^{15}N-GBI-Uba7-UFD, C996A mutant. GBI-Uba7-UFD:UbcH8 molar ratios in the bound

forms are indicated in yellow for 1:0.25, red for 1:0.5, blue for 1:1, and green for 1:1.5.

784 Supp Fig. 11. Superposition of 2D ^{1}H-^{15}N HSQCs of wt 15N-GBI-Uba7-UFD (black) with, C996S

785 mutant (green).

790 Supp Fig. 12. Superposition of the ^{1}H-^{15}N HSQC spectra of free (black) and various amounts of UbcH8

791 bound forms of 15N-GBI-Uba7-UFD, C996S mutant. GBI-Uba7-UFD:UbcH8 molar ratios in the bound

792 forms are indicated in yellow for 1:0.25, red for 1:0.5, blue for 1:1, and green for 1:1.5.

809 Supp Fig. 13. ^{1}H-^{15}N HSQC spectra of free form of R944A mutant ^{15}N-GBI-Uba7-UFD. A. ^{1}H-^{15}N

810 HSQC spectrum from first gel filtration peak. B. ^{1}H-^{15}N HSQC spectrum from second gel filtration peak.

813   Supp Fig. 14. Superposition of the ^{1}H-^{15}N HSQC spectra of free (black) and various amounts of UbcH8

814   bound forms of ^{15}N-GBI-Uba7-UFD, A1004C mutant. GBI-Uba7-UFD:UbcH8 molar ratios in the bound

815   forms are indicated in yellow for 1:0.25, red for 1:0.5.

A

B

819   Supp Fig. 15. 2D NMR analysis of Uba7-UFD 998GGG mutant A. ^{1}H-^{15}N HSQC spectra of free form of

820   998GGG mutant ^{15}N-GBI-Uba7-UFD B. Superposition of the ^{1}H-^{15}N HSQC spectra of free (red) and 1:1

821   ratio of UbcH8 bound form of ^{15}N-GBI-Uba7-UFD, 998GGG mutant (Green).

Distance and dihedral restraints	
Short range (|i-j| ≤ 1)	224
Medium range (|i-j| < 5)	319
Long range (|i-j| > 5)	358
Dihedral restraints (φ/ψ)	
Dihedral restrains	153
Total number of restricting constraints	1616
Average RMSD (Å) against the lowest energy model for ordered residues (PSVS)	
Backbone	0.9
All heavy atoms	1.4
RMSD from ideal geometry	
Bond lengths (Å)	0.015
Bond angles (°)	1.4
Consistent violations	
Distance constraints (>0.5 Å)	0
Dihedral angle constraints (>5°)	14.6
Van der Waals constraints (>0.2 Å)	1.8
Ramachandran analysis	
Favoured regions (%)	93.7
Allowed regions (%)	6.2
Generously allowed	0.1
Disallowed	0

823   Table 1. Restrains and statistics for 20 best solution NMR structures of Uba7-UFD