Sequential Membrane Remodeling by Cholesterol Distinctly Modulates HCN Channels in Naïve and Neuropathic DRG Neurons

Lucas J. Handlin[a], Clémence Gieré[b], Nicolas L.A. Dumaire[b], Lyuba Salih[b], Aubin Moutal[b], and Gucan Dai[a,1]

Author Affiliations

[a]Edward A. Doisy Department of Biochemistry and Molecular Biology, Saint Louis University School of Medicine

[b]Department of Pharmacology and Physiology, Saint Louis University School of Medicine

[1]Correspondence: Gucan Dai (email: gucan.dai@health.slu.edu)

ABSTRACT

Cholesterol, abundantly present in distinct plasma membrane pools, is a critical modulator of ion channel function, including hyperpolarization-activated cyclic nucleotide-gated (HCN) channels that regulate the excitability of dorsal root ganglion (DRG) nociceptor neurons. However, the relative contributions of cholesterol that organizes ordered membrane domains (OMDs) versus freely accessible cholesterol pools to HCN channel modulation remain unknown. Using advanced FRET techniques, we examined how supplementing cholesterol alters nociceptor DRG neurons in the context of a rat model of nerve injury. We uncovered a process of sequential, stepwise membrane remodeling: an initial phase with OMD expansion and a rapid rise in free cholesterol, followed by continued accumulation of free cholesterol without further OMD expansion. In injured DRG neurons with low cholesterol and reduced OMD size, both mechanisms contribute, whereas in naïve DRG neurons—characterized by high cholesterol and large OMDs—modulation occurs mostly via increased free cholesterol. These findings deepen our understanding of cholesterol's role in modulating ion channels and contributing to neuropathic pain.

INTRODUCTION

Cholesterol is a fundamental component of the eukaryotic plasma membrane, accounting for about 30-50 mol% of total lipids in mammalian cells and exerting profound effects on membrane structure and function (*1-3*). Its rigid sterol ring structure embeds between the acyl chains of membrane lipids, reducing their motional freedom and promoting lipid packing, thereby increasing membrane order and thickness while decreasing permeability to small polar molecules (*4, 5*). Beyond these physical effects, cholesterol contributes to the lateral heterogeneity of the membrane by driving the formation of lipid-ordered membrane domains, which serve as organizing platforms for proteins and signaling assemblies (*2, 6*). The asymmetric distribution of cholesterol between the inner and outer leaflets, as well as its dynamic exchange between membrane pools, adds further complexity to its regulatory functions (*3, 7*). Given its central role in shaping membrane architecture and influencing protein–lipid interactions, cholesterol has emerged as a key regulator of cellular excitability and signal transduction (*8-11*).

Cholesterol-mediated modulation of ion channel function remains an area of intense investigation, with longstanding debates about whether its effects arise from direct molecular interactions or from alterations in membrane organization (*1, 12-14*). Emerging evidence suggests that cholesterol-rich, lipid-ordered membrane compartments—also known as ordered membrane domains (OMDs)—play a crucial role in organizing membrane proteins and signaling complexes, thereby influencing the functional properties of ion channels (*1, 6, 10, 15, 16*). Among these, OMD-mediated modulation of hyperpolarization-activated cyclic nucleotide-gated (HCN) channels has emerged as a key mechanism in the pathophysiology of neuropathic pain (*10, 14, 17-20*). HCN channels regulate neuronal excitability through generating "I_h current", which is mostly sodium inward current in physiological conditions that promotes membrane depolarization (*21-24*). For somatosensory neurons, small-diameter nociceptor dorsal root ganglion (DRG) neurons predominantly express the HCN2 isoform and exhibit relatively slow HCN currents that are highly sensitive to cAMP (*10, 17, 25, 26*). In contrast, larger DRG neurons display faster HCN currents and primarily express the HCN1 isoform, which shows limited sensitivity to cAMP (*17, 25*). Aberrant upregulation or altered gating of HCN channels in sensory neurons has been linked to increased excitability and spontaneous firing, hallmarks of chronic pain states (*17, 27, 28*).

The voltage-sensor activity and function of HCN channels are modulated by their localization within OMDs, suggesting a lipid-dependent mechanism of regulation (*10, 20*). We recently uncovered that disruption of OMDs dramatically alters native HCN currents, pointing to OMDs as key modulators of HCN channel activity in the context of neuropathic pain (*10, 29*). Cholesterol extraction and

supplementation modulate action potential firing in small nociceptor DRG neurons, in part by altering OMD size and consequently affecting HCN channel gating (*10*). Understanding how cholesterol and OMDs regulate HCN channels could reveal novel therapeutic strategies for targeting pathological hyperexcitability in neuropathic pain.

This study provides evidence that cholesterol dynamically regulates HCN channel gating through a dual mechanism: by modifying OMDs and by altering free, accessible cholesterol levels in the inner leaflet of the plasma membrane. By integrating fluorescence lifetime imaging (FLIM), Förster resonance energy transfer (FRET), genetically encoded cholesterol sensors, and homo-FRET-based anisotropy imaging, we reveal a sequential response to cholesterol enrichment at the plasma membrane: an initial OMD expansion, followed by a sustained increase in inner and outer leaflet cholesterol. This distinction allows us to temporally separate the impact of OMDs from that of cholesterol accumulation, addressing a persistent challenge in membrane biophysics. Our findings have direct implications for the pathophysiology of neuropathic pain. In a nerve injury model, we show that DRG neurons exhibit smaller OMDs and reduced inner-leaflet cholesterol. In vivo, cholesterol depletion induces mechanical hypersensitivity in naïve rats while cholesterol supplementation alleviates mechanical hypersensitivity in a model of chronic neuropathic pain. These changes correlate with altered HCN channel gating, providing a mechanistic link between lipid remodeling and hyperexcitability in chronic pain states. By elucidating how cholesterol dynamically regulates HCN channels, this study opens new avenues for investigating cholesterol-mediated modulation of ion channels in excitable cells.

RESULTS

Temporally resolved steps of membrane remodeling during cholesterol enrichment

To define how cholesterol modulates membrane functions, we first examined changes in plasma membrane properties following cholesterol enrichment. Because cholesterol is an inherent component of OMDs, distinguishing its specific effects from those of OMDs on membrane proteins remains challenging. Altering one without affecting the other was not feasible. To quantify the OMD size, we used FRET pairs based on cholera toxin subunit B (CTxB), specifically Alexa Fluor 488 (AF-488) CTxB conjugates as FRET donors and Alexa Fluor 647 (AF-647) CTxB conjugates as FRET acceptors. As previously demonstrated, CTxB binds to the GM1 lipids in OMDs with high affinity (*30, 31*), and the FRET efficiency between CTxB-AF-488/CTxB-AF-647 can report the size of OMDs in cultured cells (*10*). To understand the roles of cholesterol itself versus the OMDs, in combination with the OMD probes, we used a genetically encoded cholesterol sensor eGFP-GRAM-W that labels the inner leaflet accessible cholesterols (*32*). The GRAM-W sensor is derived from the GRAM (Glucosyltransferases, Rab-like GTPase Activators, and Myotubularins) cholesterol-binding domain of the cholesterol transport protein $GRAMD_{1b}$, with a glycine mutated to tryptophan (G187W) (*32-34*). This mutation enhances its localization to the plasma membrane. Compared to traditional cholesterol sensors, such as those derived from perfringolysin O (PFO-D4) or anthrolysin O (ALO-D4), which often bind to cytosolic cholesterol, GRAM-W offers improved plasma membrane targeting and specificity (*32*). This makes GRAM-W a more reliable tool for studying cholesterol dynamics at the plasma membrane (PM).

Live-cell FLIM-FRET imaging using CTxB-conjugated FRET pairs was performed after applying CTxB AF-488 (FRET donor) and AF-647 (FRET acceptor) to label plasma membrane OMDs in tsA-201 cells (Fig. 1a). Fluorescence lifetimes were analyzed using phasor plots (*10*), which map each pixel's lifetime and allow identification of FRET-induced shifts, reflecting an increase in the OMD size (Fig. 1 a and b). FLIM-FRET efficiency was calculated along a phasor plot-based FRET trajectory, with phasor points farther from the donor-alone position indicating higher FRET (Fig. 1a) (*10*). We found that the FLIM-FRET increased rapidly following the addition of water-soluble cholesterol (WSC, 0.5 mg/mL cholesterol pre-complexed with methyl-β-cyclodextrin) to the culture medium, reaching saturation within 2 minutes (Fig. 1 b and c). In parallel, membrane-localized fluorescence intensity of eGFP-GRAM-W also increased after WSC treatment, but with significantly slower and double-exponential kinetics (Fig. 1c and Supplementary Fig. 1). This difference suggests that OMD expansion occurs rapidly and reaches a plateau soon after cholesterol enrichment, whereas the accumulation of free cholesterol in the inner

leaflet proceeds more gradually, reflecting a two-phase effect of cholesterol enrichment on PM properties and distinguishing OMD changes from those of free cholesterol. This distinction is likely due to the limited availability of other lipid species, such as sphingolipids, for establishing OMDs, and enables us to use this approach to functionally isolate the contributions of OMDs and free cholesterol to the PM in cultured cells.

Fig. 1

Fig. 1. Modulation of plasma membrane (PM) properties by cholesterol enrichment.
(**a-b**) Representative phasor plots of membrane-localized fluorescence of tsA201 cells from donor alone (CTxB AF-488) or with CTxB AF-647 before (a) and after (b) the water-soluble cholesterol (WSC) treatment. The FRET efficiency was calculated using the FRET trajectory (in red). The insets show amplified views of the phasor plots. A confocal image of membrane localized fluorescence of a tsA cell labeled with CTxB is also shown. (**c**) Time course of the change in normalized fluorescence intensity at the PM and the CTxB-based FLIM-FRET efficiency post 0.5 mg/mL WSC application in tsA201 cells. Data shown are mean + s.e.m., WSC was added to the culture medium at the start of imaging, which was performed at intervals of ~27 s for FRET (n = 4) and ~20 s for GRAM-W (n = 3). Time zero refers to the state prior to WSC addition. The purple bar denotes the duration of WSC pipetting into the recording chamber. (**d**) Strategies implemented to report the PM properties after WSC-mediated cholesterol supplementation. A mCherry-labeled ostreolysin A (OlyA) was used to recognize outer leaflet cholesterol/SM complexes, whereas GRAM-W-eGFP was used to label free cholesterol levels in the inner leaflet of the PM. The OMD size was probed using FRET-based reporters, either CTxB-conjugated or peptide (L10)-based FRET pairs. (**e**) Time course of the change in normalized fluorescence intensity and the FLIM-FRET efficiency at the PM post 0.5 mg/mL WSC application in tsA201 cells, using four different probes as illustrated in d. Data were collected every 5 mins to minimize photobleaching, and shown as mean ± s.e.m., n = 4 - 10 cells. (**f-g**) Summary time course of the change in normalized GRAM-W fluorescence intensity at the PM and the FLIM efficiency of CTxB-AF488/AF647 FRET post 0.5 mg/mL WSC application for naïve DRG neurons (f) and for nociceptor DRG neurons of the SNI model animals (g). Data shown are mean ± s.e.m., n = 4 - 6 cells. The CTxB-based FRET level showed a significant increase after WSC for SNI DRG neurons (*p < 0.05 using paired t-test) but no change for naïve neurons (n.s.: no statistical significance).

To further demonstrate the two phases of PM modulation following cholesterol enrichment, a mCherry-tagged ostreolysin A (OlyA) was used to report levels of outer leaflet cholesterol/sphingomyelin (SM) complexes (*35, 36*) (Fig. 1d). Cholesterols sequestered by SM cannot be probed by conventional cholesterol sensors like PFO-D4 (*7*). In addition, we employed a previously established FRET-based OMD probe to report OMD size, using a peptide (Lck 10 or L10)-based FRET pair (*20, 31, 37, 38*), and the CTxB-conjugated probes as described above (*10*). Applying the all four types of probes individually and overlaying their time courses following the same WSC treatment in tsA-201 cells consolidated the two-step effect of cholesterol enrichment on plasma membrane properties (Fig. 1e). Imaging was performed at 5-minute intervals to focus on distinguishing changes in OMD-associated versus free cholesterol, rather than tracking the precise kinetics of the FRET change caused by *in vitro* membrane enrichment with high cholesterol concentrations. This approach particularly minimized photobleaching, which could otherwise obscure intensity-based fluorescence measurements. Consistently, we observed an initial increase in FRET between CTxB-AF-488/CTxB-AF-647 and a decrease in FRET between L10-CFP/L10-YFP. For the AF-488/AF-647 pair, the FRET efficiency increased from 9% to 11.8% (p = 0.02), whereas for the L10 FRET pair, the FRET efficiency decreased from 8.3% to 4.5% (p = 0.039) (Fig. 1e). As previously demonstrated, these opposing FRET changes both indicate OMD expansion (*10*) (Fig. 1e). The FRET changes plateaued within 5 minutes as we have shown using the higher frequency imaging in Fig. 1c. In contrast, GRAM-W and OlyA fluorescence at the plasma membrane continued to increase beyond the first 2–5 minutes, extending throughout the 15-20-minute timeframe (*32*). These results suggest that the initial phase of cholesterol enrichment involves OMD expansion along with increases in

inner leaflet cholesterol and outer leaflet cholesterol/SM complexes. The second phase primarily reflects a continued accumulation of cholesterol, either in free form or bound to SM, in tsA-201 cells. Since the second phase does not involve OMD expansion, any functional effects likely arise from the free cholesterols. The limiting factor for the OMD expansion could be the amount of SM available. Collectively, this methodology provides a potential strategy to temporally distinguish the effects of OMD expansion from cholesterol itself on ion channel gating in DRG neurons.

We observed distinct patterns in the effects of WSC supplementation on changes in OMD and free cholesterol levels when comparing small-diameter nociceptor DRG neurons from naïve rats and those subjected to the spared nerve injury (SNI) model of neuropathic pain (39). As we previously demonstrated, nociceptor DRG neurons in the SNI model exhibit significantly smaller OMDs compared to the large OMDs observed in naïve nociceptor DRG neurons (10). In both conditions, membrane-localized fluorescence intensity of the GRAM-W sensor showed a sustained increase over the 15-minute WSC treatment (Fig. 1 f and g). However, the CTxB-based FRET reporter of OMD size exhibited minimal changes in FLIM-FRET efficiency for naïve DRG neurons, whereas SNI DRG neurons showed a substantial increase within 5 minutes (Fig. 1 f and g, and Supplementary Fig. 2). The initial FLIM-FRET efficiency measured using CTxB-based probes was significantly lower in SNI DRG neurons (6.3%) than in naïve neurons (17.3%) and slightly lower than in tsA-201 cells (9%). Notably, following WSC supplementation, the OMD changes detected by CTxB-based FRET in SNI DRG neurons closely mirrored those observed in tsA-201 cells, reaching an increase of 12.7%.

Homo-FRET based fluorescence anisotropy measurement of membrane cholesterol levels.

We performed homo-FRET-based live-cell anisotropy for imaging membrane-localized GRAM-W fluorescence (Fig. 2) (40). Fluorescence anisotropy measures the polarization of emitted light by comparing perpendicular and parallel emission components, enabling the quantification of homo-FRET by detecting depolarization resulting from energy transfer between nearby fluorophores (Fig. 2 a, b). A decrease in anisotropy reflects an increase in homo-FRET efficiency, indicating closer proximity or clustering between the probes (40). Homo-FRET-based anisotropy is sensitive to the organization and density of fluorophores while being less influenced by factors such as photobleaching of fluorophores (41). This sensitivity enables a more accurate quantification of inner-leaflet cholesterol density, which would be challenging to resolve using intensity-based fluorescence alone. Following WSC supplementation, we observed a continued decrease in GRAM-W fluorescence anisotropy, accompanied by an increase in homo-FRET efficiency (Fig. 2 c-e). Notably, the initial homo-FRET efficiency was

highest in naïve DRG neurons but lower in tsA-201 cells and DRG neurons of the SNI model. These results suggest that free, accessible cholesterol in the inner leaflet is also reduced after nerve injury, adding to the previously reported decrease in OMD size observed in neuropathic pain (*10*).

Fig. 2. Fluorescence anisotropy reports homo-FRET between cholesterol sensors.
(**a**) Representative confocal images showing membrane-localized eGFP-GRAW-W fluorescence emission in a naïve DRG neuron, acquired in perpendicular and parallel directions. (**b**) Corresponding anisotropy-based heatmaps derived from the images in a, shown for both pre- and post-WSC conditions, illustrating a reduction in anisotropy after WSC treatment. (**c-e**) Summary time course of the changes in the GRAM-W fluorescence anisotropy and estimated homo-FRET efficiency for tsA cells (c, n = 10), naïve DRG neurons (d, n = 6), and DRG neurons of SNI model animals (e, n = 5). Since the homo-FRET data are derived from the measured anisotropy, the statistical significance is only shown for the anisotropy and remains the same for the homo-FRET. (**f-h**) Summary time course of changes in the OlyA fluorescence anisotropy and estimated homo-FRET efficiency for tsA

cells (f, n = 4), naïve DRG neurons (g, n = 6), and DRG neurons of SNI model animals (h, n = 4). Data shown are mean ± s.e.m., *p < 0.05, **p < 0.01 using paired t-test.

We also performed anisotropy experiments using OlyA-mCherry and observed distinct differences in homo-FRET responses to WSC treatment across tsA-201 cells, naïve DRG neurons, and SNI DRG neurons (Fig. 2 f-h). In tsA cells, anisotropy and homo-FRET revealed small but significant changes, aligning with the modest increase in outer leaflet cholesterol/SM complexes as concluded from the intensity-based measurements. In naïve DRG neurons, anisotropy and homo-FRET values remained unchanged after WSC treatment, in contrast to the noticeable decrease in anisotropy and corresponding increase in homo-FRET observed in SNI DRG neurons. Notably, baseline homo-FRET values were highest in naïve DRG neurons and reduced in SNI DRG neurons, consistent with a decrease in OMD size following nerve injury. Given that OlyA binds to cholesterol/SM complexes enriched in OMDs, the high baseline homo-FRET and lack of response to cholesterol supplementation in naïve DRG neurons suggest that these cells already contain high or saturated levels of cholesterol/SM complexes that cannot be further increased by adding free cholesterol. This again implies that the availability of SM, rather than cholesterol, may be the limiting factor for the OMD expansion.

Inhibitory effects of cholesterol on HCN channel gating in nociceptor DRG neurons

Previously, we modulated cholesterol levels in DRG neurons for electrophysiological measurements using methyl-β-cyclodextrin (mβCD) for extraction or water-soluble cholesterol (WSC) for supplementation, achieved by preloaded methyl-β-CD-cholesterol complexes (*10, 42, 43*). These treatments were brief (30 sec to 1 min) and applied prior to whole-cell patch-clamp recordings on separate control and treated cells. Here, we applied these reagents directly to the same patch during whole-cell recordings and compared hyperpolarization-activated HCN currents in nociceptor DRG neurons before and after a more prolonged treatment (Fig. 3). We found that 0.5 mg/mL WSC supplementation significantly inhibited HCN current, reducing the amplitude of currents elicited by a maximal hyperpolarizing voltage, suggesting a decrease in channel open probability (Fig. 3 a and b), and shifting the half-maximal activation voltage leftward for the conductance-voltage (G-V) relationship ($\Delta V_{1/2} = 8.4$ mV after 10 mins of WSC treatment) (Fig. 3 d and e). These effects on the G-V relationship and current amplitude persisted in the presence of additional cAMP at a supersaturating concentration of 0.5 mM in the pipette solution, with a $\Delta V_{1/2}$ of ~ 6 mV and a ~25% reduction in current amplitude following 10 minutes of WSC treatment. The use of a supersaturating 0.5 mM concentration of cAMP (well above the ~1 μM EC_{50} for HCN channels) ensures rapid diffusion into the cytosol and a saturating effect on HCN channel activity (*44*). In contrast, cholesterol extraction (5 mM mβCD) produced nearly opposite effects,

increasing current amplitude (Fig. 3c and Supplementary Fig. 2), though it had minimal impact on $V_{1/2}$, with or without the additional 0.5 mM cAMP (*10*). Instead, cholesterol extraction increased the slope factor (V_s) of the G-V relationship (*10*), whereas WSC treatment caused only a negligible change in V_s for naïve nociceptor neurons either with or without the additional 0.5 mM cAMP (Fig. 3f). Since the V_s reflects voltage sensor movement and is influenced by the OMD localization of HCN channels, it could serve as a reporter of OMD dimensions of DRG neurons (*10, 20*). We propose that these effects arise from the naturally high cholesterol content and large OMDs in naïve DRG neurons. In this context, further increases in cholesterol levels may primarily elevate free accessible cholesterol, leading to a shift in $V_{1/2}$ and a decrease in open probability without affecting the V_s. Our findings lead to a model in which cholesterol modulates HCN channels through a dual mechanism: by altering OMD size and potentially interacting directly with the channels at the inner leaflet of the membrane. Moreover, this substantial modulation of HCN current amplitude (and open probability) is physiologically significant and was validated through the same patch-clamp experiments. Compared to changes in the slope factor, alterations in current amplitude may have a greater impact on action potential firing and may contribute more directly to neuronal excitability.

Beyond the equilibrium parameters of HCN gating, we found that the activation kinetics of HCN channels in nociceptor DRG neurons were modulated by cholesterol in the same patch-clamp experiments. The most prominent effect was a significant acceleration of channel activation following treatment with 5 mM mβCD, which shortened the primary activation time constant (τ_1) from an average of 80 ± 4.3 ms to 52 ± 2.9 ms, at the saturating hyperpolarizing voltage of –140 mV (Fig. 3g). This effect was also present at other hyperpolarizing voltages (Fig. 3h). In contrast, cholesterol enrichment with 0.5 mg/mL WSC for 5 mins slowed channel activation, increasing τ_1 from an average of 72 ± 9.6 ms to 88 ± 8.7 ms at –140 mV, although this effect was less pronounced than that of mβCD (Fig. 3 i and j). The mβCD-induced acceleration of activation persisted in the presence of 0.5 mM cAMP in the pipette solution (Supplementary Fig. 3). However, under the same cAMP condition, the effect of WSC treatment on τ_1 was no longer significant (τ_1 from an average of 59 ± 4.5 ms to 65.3 ± 13.5 ms, n = 4), likely because the cAMP potentiation of channel activation is predominant, and there is likely a convergent effect between cAMP and cholesterol. Together, these findings suggest that HCN channel activation is highly sensitive to cholesterol levels. In addition, our recent findings on cholesterol-mediated modulation of current amplitude and activation kinetics (*10*) were further substantiated in this study, offering new insights into the mechanisms by which HCN channels influence neuronal excitability.

Fig. 3

Fig. 3. Effects of manipulating cholesterol levels on the equilibrium and kinetic properties of HCN channel gating in nociceptor DRG neurons.
(**a**) Application of WSC decreased the current amplitude of HCN currents in the same-patch experiment. Double-exponential fittings (in red) of the current traces at -140 mV are highlighted. (**b**) Amplitude of HCN currents elicited

by a saturating hyperpolarizing voltage was used for the graph comparing the before and after treatments. Data shown are mean + s.e.m., n = 10 patches, *p = 0.01, paired t-test. (**c**) Effects of the acute mβCD treatment on current amplitude of HCN currents, recorded from the same patch, showing opposite directional changes as in panel B. Data shown are mean ± s.e.m., n = 5 patches, **p = 0.005, paired t-test. (**d**) Representative G-V relationship of HCN channel activation recorded from same-patch experiments before and after 5 minutes of 0.5 mg/mL WSC application, in the presence or absence of 0.5 mM additional cAMP. (**e-f**) Summary of the parameters derived from Boltzmann fitting of the G-V relationship of HCN channel activation before and after the WSC treatment: $V_{1/2}$ in panel e and slope factors (V_s) in panel f, n.s.: no statistical significance. n = 7 cells without added cAMP and n = 5 with added 0.5 mM cAMP, *p < 0.05, **p < 0.01, paired t-test. (**g**, **i**) HCN channel currents elicited by different hyperpolarizing voltages were shown with double-exponential fits to highlight the acceleration of channel activation by mβCD (g) and the slowdown of channel activation by WSC (i). (**h**, **j**) Summary graph illustrating the effects of mβCD (h, n = 5) and WSC (j, n = 8) on the τ_1 of channel activation at different voltages. Data shown are mean ± s.e.m, *p < 0.05, **p < 0.01, paired t-test.

Sequential modulation of HCN channel gating by cholesterol enrichment in DRG neurons

Based on our FLIM-FRET and anisotropy experiments indicating that the effects of cholesterol enrichment can be temporally distinguished using the WSC (0.5 mg/mL) supplementation protocol, we applied this approach to examine whether cholesterol-mediated modulation of membrane excitability following HCN activation could exhibit a biphasic feature, using a protocol in which the HCN activation in small DRG neurons was assessed from the recovery in membrane potential after a hyperpolarizing current pulse. A –90 pA current injection induced pronounced membrane hyperpolarization, activating endogenous HCN channels, which then depolarized the membrane and restored the potential over a few hundred milliseconds to a steady state (Fig. 4a). The percentage of voltage recovery at steady state primarily reflects the extent of HCN channel opening. We found that the percentage of voltage recovery decreased after same-patch WSC treatment, reaching a maximal effect within ~5 minutes with an exponential time constant of 128 s (Fig. 4b). No decrease was observed in the control experiments without WSC treatment (Fig. 4b). Furthermore, similar results were observed in SNI nociceptor DRG neurons (Fig. 4c); however, compared to naïve DRG neurons, the WSC-induced reduction showed an initial delay with a slower time constant of 176 s, consistent with a sequential cholesterol-mediated modulation of HCN function. Presumably, in SNI neurons, voltage recovery after hyperpolarization was relatively resistant to WSC treatment during the first 200 s, likely because a substantial portion of cholesterol was initially allocated to restoring OMDs. This could result in a smaller apparent increase in the free cholesterol pool (Fig. 1 f and g) that plays a substantial role in maintaining HCN channel opening.

Fig. 4

Fig. 4. Effects of cholesterol enrichment on HCN channel function in nociceptor DRG neurons from naïve and SNI animals.
(**a**) Representative protocol and membrane potential recordings using hyperpolarizing current injection (−90 pA) to activate HCN channels, leading to membrane depolarization. The ratio of the recovered potential ($V_{rec.}$) relative to the total hyperpolarization (V_{peak}) reflects the percent of recovery in membrane potential mediated by HCN activation. (**b**) Time course of changes in the percent recovery ($V_{rec.} / V_{peak}$) after WSC treatment for naïve nociceptor DRG neurons (n = 4 patches), compared with untreated controls (n = 3 patches), mean ± s.e.m.; red trace shows a single-exponential fit. (**c**) Time course of the change in the percent recovery for SNI DRG neurons (n = 3 patches, mean ± s.e.m.) using the same protocol as in panel a and b. (**d-f**) Time course of the change in the $V_{1/2}$ and slope factor (V_s) derived from the G-V relationship (d), the current amplitude (e), and τ_1 of channel activation (f) after 0.5 mg/mL WSC application for naïve small DRG neurons, n.s.: no statistical significance for the slope factor over time after WSC treatments, n = 7 - 10 patches. (**g-i**) Summary time course of the change in the $V_{1/2}$ and slope factor (V_s)

(g), the current amplitude (h), and τ_1 of channel activation (i) after 0.5 mg/mL WSC application for SNI small DRG neurons. Compared to naïve neurons, the slope factor in panel h showed significant changes in SNI neurons following WSC treatment. Data shown are mean ± s.e.m., n = 4 - 7 patches, *p < 0.05, **p < 0.01, paired t-test.

Next, to examine in greater detail how cholesterol supplementation modulates HCN channel gating and to test whether this modulation occurs sequentially, we performed same-patch time-course experiments. We quantified the effects of WSC (0.5 mg/mL) supplementation on HCN channel gating parameters by recording native HCN currents and plotting the G-V relationships, from both naïve DRG neurons and DRG neurons of the SNI model (Fig. 4 d-i). These HCN function measurements were performed every 5 minutes post WSC treatment, as each voltage protocol series requires 2–3 minutes to complete. In naïve DRG neurons, the slope factor V_s remained unchanged (Fig. 4d). In contrast, the $V_{1/2}$ ($\Delta V_{1/2}$ = ~10 mV), current amplitude, and the primary time-constant τ_1 of activation exhibited significant alterations throughout the ~15 minutes of WSC treatment, with the most pronounced effects observed within the first 5 minutes (Fig. 4 d-i). Control experiments without WSC treatment showed only minor changes, likely due to gradual dilution of endogenous cytosolic cAMP ($\Delta V_{1/2}$ = ~3 mV and little to no change in slope factor, current amplitude, and τ_1 of activation, Supplementary Fig. 4). Moreover, control experiments with WSC treatment, in the presence of supersaturating 0.5 mM added cAMP, showed a $\Delta V_{1/2}$ = ~7 mV and no change in the slope factor, post WSC treatment. Results from these control experiments indicate that the observed effects were specifically mediated by cholesterol enrichment.

DRG neurons from the SNI model displayed changes in all gating parameters over the course of WSC exposure (Fig. 4 g-i). The key difference compared to naïve neurons was the significant change in the V_s, which occurred within the first 5 minutes (Fig. 4g). Additionally, the change in the $V_{1/2}$ for SNI neurons ($\Delta V_{1/2}$ = ~6 mV) was less pronounced compared to that in naïve neurons (Fig. 4 d and g, and Supplementary Fig. 4), which partially explains why the WSC-induced reduction in the recovery of membrane potential in Fig 4c was slower in SNI neurons. These findings suggest that the reduction in the slope factor primarily reflects OMD expansion, whereas the changes in $V_{1/2}$, current amplitude, and τ_1 for activation likely result from both OMD expansion and an increased accumulation of accessible cholesterols in the plasma membrane. Similarly, all gating parameters changed more markedly within the first 5 minutes than between 5 and 15 minutes (Fig. 4 d-i). Cholesterols could exert direct binding effects on HCN channels in DRG neurons, as suggested by a putative cholesterol binding site of HCN3 channels (*45*), which is conserved across all HCN isoforms.

We previously demonstrated that the voltage sensor S4 helix of the HCN channel is sensitive to changes in OMD size in DRG neurons (*10, 20*). Notably, the unusually long S4 segment in HCN channels (*20, 46, 47*) may underlie their unique susceptibility to dual modulation by cholesterol and OMDs, a feature not commonly observed in other voltage-gated ion channels. To assess whether this mechanism extends to other key ion channels involved in DRG neuron excitability, we recorded sodium currents from small nociceptive DRG neurons following cholesterol depletion (Supplementary Fig. 5). Sodium channels in DRG neurons play important roles in controlling action potential firing and pain sensation (*48, 49*). Similar to previous findings (*50, 51*), mβCD treatment caused a slight hyperpolarizing shift in the activation curve ($\Delta V_{1/2}$ = ~ 5 mV), without affecting the slope factor of activation or steady-state parameters of the fast inactivation. In contrast, as we emphasized earlier, cholesterol depletion increases the slope factor of the G-V relationship for HCN channels in nociceptor DRG neurons, suggesting a decrease in the effective gating charge movement associated with channel activation (*10*). Sodium channels may not preferentially localize to OMDs, or their shorter S4 segments may lack sensitivity to changes in membrane domain compartmentalization (*52*). These findings suggest that the dual modulatory effect of cholesterol on HCN channels may represent a channel-specific mechanism not shared by voltage-gated sodium channels.

Cholesterol levels modulate membrane excitability and mechanical pain sensitivity

Cholesterol enrichment reduced the hyperexcitability of SNI nociceptor DRG neurons *in vitro*, consistent with a suppressive effect on HCN channel activity in our experiments (*10*). We observed considerable effects in the same patch experiment, consistent with our previous findings using β-cyclodextrin (β-CD) or WSC pretreatments of DRG neurons (*10*). In non-spontaneously firing naïve DRG neurons, cholesterol extraction induced spontaneous firing (Fig. 5a), whereas in spontaneously firing neurons, it increased firing frequency (Fig. 5b). In the same-patch experiment, WSC treatment markedly suppressed action potential firing, as demonstrated by current-clamp recordings with incremental current injections (Fig. 5c and d). These same-patch treatment data highlight the critical role of cholesterol in regulating the membrane excitability of sensory neurons.

Fig. 5

Fig. 5. Cholesterol regulates membrane excitability and mechanical hypersensitivity in naïve and neuropathic pain rats.

(**a-b**) Representative action potential firing of non-spontaneously (a) and spontaneously (b) firing naïve DRG neurons before and after β-cyclodextrin (β-CD) treatment from the same-patch experiments (**c**) Representative action potential firing of a naïve DRG neuron before and after the WSC treatment from the same-patch experiment. (**d**) Summary graph showing the effect of WSC on the action potential firing in relation to the injected currents for naïve DRG neurons (n = 11, mean ± s.e.m.). (**e-f**) Naïve rats were injected intraplantarly with β-CD or its ineffective analog, α-cyclodextrin (α-CD) (20 mM each in 50 μL), and paw withdrawal threshold (PWT) was measured. In d, graph showing the hind-paw withdrawal threshold of naïve rats injected as indicated. The threshold for the injected paw versus the un-injected paw for each group is shown. β-CD, but not α-CD, reduced the threshold in the injected paw compared to the control paw and α-CD-injected groups. In e, bar graph with scatter plot showing the integrated area under the curves corresponding to the marked lines in panel a (from 0 to 48 h), demonstrating a significant reduction in the PWT in the β-CD-injected paw group (n = 6 rats per group). (**g-h**) Following the spared nerve injury (SNI) surgery, rats developed chronic neuropathic pain. Rats then received an intraplantar injection of WSC (50 μL). In f, graph showing the PWT of rats with SNI injected as indicated. For each rat, both the injured and non-injured paws were injected, and withdrawal thresholds were measured. WSC injection significantly increased the PWT and reversed mechanical hypersensitivity on the injured side of SNI rats while having no effect on the control non-injured side. In g, bar graph with scatter plot showing the integrated area of the data (lines with markers) in panel f (from 0 to 24 h), demonstrating a significant anti-hypersensitivity effect of cholesterol compared to saline-treated SNI rats (n = 10 - 11 rats per group). Data are presented as mean ± SEM. *p < 0.05 compared to the control.

To examine whether cholesterol manipulation influences pain behaviors *in vivo* and to assess its physiological impact, we tested if a local administration of cholesterol-depleting or cholesterol-supplementing agents in naïve or SNI neuropathic pain rats resulted in changes of mechanical hypersensitivity. Intraplantar injection of β-CD, but not its inactive analog α-cyclodextrin, induced significant mechanical hypersensitivity in naïve rats, as evidenced by a time-dependent reduction in paw withdrawal threshold on the injected paw compared to the contralateral control paw and α-cyclodextrin–injected animals (Fig. 5e). This effect persisted for 24 hours and was quantitatively confirmed (Fig. 5f), highlighting a clear pronociceptive effect of acute cholesterol chelation in naïve animals. We next examined whether cholesterol supplementation could alleviate established mechanical hypersensitivity in rats with spared nerve injury, a well-characterized model of neuropathic pain (*53, 54*). Intraplantar injection of water-soluble cholesterol partially reversed mechanical hypersensitivity in the injured paw, while having no effect on the non-injured contralateral paw (Fig. 5 g and h). The anti-hypersensitivity effect was sustained for 4 hours. Together, these findings demonstrate that cholesterol homeostasis plays a critical role in regulating nociceptive thresholds under both physiological and neuropathic pain conditions. Moreover, these findings provide *in vivo* evidence supporting the therapeutic potential of cholesterol enrichment and are consistent with our observations in cultured nociceptor DRG neurons, where WSC treatment reduces membrane hyperexcitability (*10*). Cholesterol depletion enhances pain sensitivity, whereas cholesterol supplementation reverses mechanical hypersensitivity, supporting a model in which membrane cholesterol acts as a potent modulator of pain signaling.

DISCUSSION

Our approach investigates the effects of cholesterol on ion channels and membrane properties. By combining FLIM-FRET-based imaging of OMDs, cholesterol sensors, and homo-FRET-based anisotropy imaging, we have developed a strategy to temporally distinguish the contributions of OMDs from those of free, accessible cholesterol in the plasma membrane. This approach can be broadly applied to study other membrane proteins regulated by lipid microenvironments, such as voltage-gated potassium channels, G-protein-coupled receptors, and mechanosensitive ion channels. Many of these proteins localize to or interact with OMDs, yet their regulation by cholesterol remains poorly understood. By dissecting dynamic changes in compartmentalized nanodomains and cholesterol distribution, our methodology enables future studies to explore how cholesterol modulates ion channels and impinges on neuronal excitability.

In neuropathic pain, upregulation of HCN channels and shifts in their voltage dependence contribute to spontaneous firing and heightened pain sensitivity (*17, 28, 55*). By identifying a dual mechanism by which cholesterol regulates HCN channels—via both OMD expansion and increased inner-leaflet cholesterol—we provide a mechanism for understanding how membrane lipid composition alters excitability in disease states. HCN channel hyperactivity is a well-established facilitator of neuronal firing, its inhibition—such as through cholesterol-mediated modulation—can effectively dampen the pathological hyperexcitability. However, the role of HCN is highly context-dependent: when activated near resting potential, opening HCN channels can often shunt excitability by increasing membrane conductance, effectively suppressing depolarizing currents (*28, 55-60*). This suggests that cholesterol may not uniformly mitigate excitability in all neurons; its effect likely depends on the specific cellular context, including the resting membrane potential, HCN isoform expression, and concurrent activation of other voltage-gated channels. This context dependence determines whether the dominant effect of HCN opening is excitatory or inhibitory for membrane excitability. Specifically, our results demonstrate that a decrease in free cholesterol levels at the membrane of nociceptor DRG neurons following nerve injury may be functionally linked to pain behaviors. We demonstrate that cholesterol depletion sensitizes naïve animals to mechanical stimuli, whereas cholesterol supplementation attenuates mechanical hypersensitivity in a neuropathic pain model, consistent with an inhibitory role of cholesterol. These pain-related phenotypes are accompanied by reduced homo-FRET between cholesterol sensors and distinct alterations in HCN channel gating, including shifts in voltage sensitivity and activation kinetics. The ability to temporally separate the effects of OMD expansion from those of free cholesterol accumulation using biophysical and imaging tools allowed us to link specific gating changes to different stages of membrane

remodeling. Notably, changes in slope factor (V_s) were more closely associated with OMD expansion, while shifts in half-maximal activation voltage ($V_{1/2}$) and time constants were more sensitive to increased cholesterol content in the inner leaflet.

This dual mechanism of HCN channel modulation likely arises from their distinctive structural features. The long S4 voltage-sensing helix in HCN channels, which spans a considerable portion of the membrane electric field, is particularly responsive to changes in membrane order and lipid packing (*20, 46, 47*). Furthermore, the presence of putative cholesterol-binding motifs in HCN channel subtypes raises the possibility of direct cholesterol-channel interactions that modulate gating. In addition, cholesterols could regulate other membrane lipids and indirectly modulate channel activities (*61-64*), which remains an area of future investigation. These features make HCN channels highly susceptible to lipid-mediated modulation, offering a mechanistic insight for their dynamic behavior in sensory neurons under inflammation or injury. From a translational perspective, these findings raise the possibility that manipulating membrane cholesterol—either pharmacologically or genetically—could serve as a strategy for tuning HCN channel function and alleviating chronic pain. While current therapies targeting HCN channels have largely focused on pharmacological pore-blockers, our work suggests that modulating the HCN voltage sensor/OMD interactions or the lipid-bilayer nano-environment could achieve similar outcomes.

In summary, this study provides insights into the biophysical mechanisms by which cholesterol and OMDs cooperatively regulate ion channel activity, with relevance for HCN channel function in nociceptors. By integrating fluorescence imaging, electrophysiology, and in vivo behavior, we present a view of how sequential cholesterol-driven membrane lipid remodeling contributes to neuronal hyperexcitability and pain. This integrative framework offers new insights for understanding and treating neuropathic pain by targeting the membrane environment.

MATERIALS AND METHODS

Cell Culture and Molecular Reagents

The tsA-201 cell line (catalog number: 96121229), a variant of human embryonic kidney (HEK293) cells, was sourced from Sigma-Aldrich (St. Louis, MO), and was authenticated and confirmed using STR profiling by Sigma. These cells were maintained in Dulbecco's modified Eagle's medium (DMEM; Gibco) supplemented with 10% fetal bovine serum (FBS; Gibco) and 1% penicillin/streptomycin (Gibco) under humidified conditions at 37°C with 5% CO_2 in tissue culture dishes (CellTreat, Pepperell, MA). Transfection of tsA-201 cells at 70%-90% confluency was performed using the Lipofectamine 3000 Kit (Invitrogen, Carlsbad, CA, #L30000080) following established protocols. MycoFluor™ Mycoplasma Detection Kit (Invitrogen) was used to test for contamination.

Cholera toxin subunit B (CTxB) conjugated to Alexa Fluor™ 488 and 647 was obtained from Thermo Fisher Scientific and applied to cultured cells at a concentration of 20 nM for approximately 5 - 10 minutes before imaging. CTxB binds to GM1-enriched OMDs with a picomolar affinity. β-cyclodextrin, α-cyclodextrin and water-soluble cholesterol (WSC) were purchased from Sigma-Aldrich (St. Louis, MO). For same-cell or same-patch cholesterol enrichment experiments, equal volume of 1 mg/mL WSC in external solution was pipetted into the recording chamber, resulting in a final cholesterol concentration of 0.5 mg/mL. The eGFP-GRAM-W plasmid was obtained from Addgene (#211701). The C-terminal mCherry tagged OlyA was synthesized and purified by GenScript (Piscataway, NJ) and was applied to cultured cells at a concentration of 1 uM for ~1 hour. Before imaging, OlyA-treated cells were incubated in regular DMEM medium for 30 mins to wash away unbound OlyA.

FLIM-FRET and Fluorescence Anisotropy

Fluorescence microscopy and FLIM experiments was performed like previous research (*10*). A laser scanning confocal system with a FastFLIM data acquisition module (ISS, Inc.) and two hybrid PMT detectors (Hamamatsu) was used for frequency-domain FLIM imaging. A supercontinuum laser (YSL Photonics) provided excitation at specific wavelengths for CFP, GFP, AF-488, and mCherry with consistent laser intensity across samples. Emission was detected using dichroic cubes and filters optimized for CFP/YFP and AF-488/AF-555 FRET pairs. Confocal images (256 × 256 pixels) were acquired using a 100 μm pinhole and pixel dwell times of 0.1 – 0.4 ms. Image acquisition and processing were performed with VistaVision software (ISS, Inc.), enabling phasor plotting and FLIM analysis (*10, 65-67*). A smoothing filter and intensity-threshold filter were applied to isolate membrane-localized lifetime

signals. The phasor FLIM approach distinguished membrane fluorescence from cytosolic and background signals, allowing separation of endocytosed and membrane-localized CTxB. Phase delays (φ) and modulation ratios (m) were determined using Fourier transforms, accounting for the instrument response function calibrated with fluorophores of known lifetimes (Atto 425, rhodamine 110, rhodamine B) (*66*). FLIM-FRET analysis was conducted with the VistaVision FRET trajectory function, optimizing background and donor contributions for accuracy, maintaining background levels below 5% to minimize interference.

Fluorescence anisotropy was used to assess the membrane localization and dynamics of eGFP-GRAM-W and mCherry-tagged OlyA proteins. Anisotropy measurements were performed using a polarization module integrated with a confocal microscope, enabling simultaneous acquisition. The system included a halfwave plate and a linear polarizer assembly at the laser entrance, both of which were rotatable with 1-degree precision over 360 degrees to define the excitation polarization state. Emission fluorescence was separated into parallel ($I_\parallel$) and perpendicular ($I_\perp$) components relative to the excitation polarization using a polarization beam splitter assembly, allowing for precise anisotropy measurements. Cells labeled with membrane-localized GFP- or mCherry-tagged constructs were cultured on glass-bottom petri-dishes and maintained in imaging buffer to preserve physiological conditions. Anisotropy was determined using a software package in VistaVision that acquired fluorescence intensity in the two detection channels, with intensity-based background subtraction applied to correct for autofluorescence and cytosolic fluorescence. The fluorescence anisotropy (r) at the plasma membrane was calculated using the equation: $r = (I_\parallel - I_\perp) / (I_\parallel + 2*I_\perp)$ (*41, 68*).

The homo-FRET efficiency of GRAM-W was estimated using fluorescence anisotropy measurements at the plasma membrane and in the cytoplasmic region following β-cyclodextrin treatment. After 5 minutes of cholesterol extraction with β-cyclodextrin, GRAM-W exhibited near-complete translocation to the cytosol, where the measured anisotropy of 0.375 was assumed to represent a state minimally affected by homo-FRET and was designated as r'. This assumption is based on the typical cytosolic concentration of constructs following overexpression, which ranges from 1-10 µM, corresponding to average intermolecular distances of approximately 118 nm (for 1 µM) to 55 nm (for 10 µM) in a 3-D space. Given that the Förster distance (R_0) for homo-FRET between GFP molecules is approximately 5 nm, these large intermolecular distances in the cytosol are expected to preclude significant homo-FRET interactions. The homo-FRET efficiency was then calculated using the equation $E_{homo\text{-}FRET} = 1 - r/r'$, where r is the measured anisotropy at the plasma membrane and r' is the cytosolic anisotropy (0.375), serving as the reference value in the absence of homo-FRET (*41, 69*). The homo-FRET efficiency of OlyA-mCherry was estimated using a similar approach. The anisotropy of OlyA-

mCherry in the absence of homo-FRET was measured to be 0.251 using purified OlyA-mCherry dissolved (5 µM) in dPBS.

Rat Studies and Primary Dorsal Root Ganglion (DRG) Neurons

Pathogen-free adult Sprague-Dawley rats (RRID: RGD_70508, females ~100 g weight, males ~250 g weight; Envigo, Placentia, CA) were maintained in a controlled environment with a 12-hour light/dark cycle (lights on at 07:00 h) and a regulated temperature of 23 ± 3°C. Food and water were provided ad libitum. All experimental procedures were conducted in accordance with National Institutes of Health guidelines and were approved by the Institutional Animal Care and Use Committee (IACUC) of Saint Louis University (protocol number: 3014). Behavioral testing was performed by researchers blinded to experimental conditions.

Dorsal root ganglia (DRG) from the ipsilateral side of nerve injuries were isolated from 100 g female Sprague-Dawley rats. The dissection involved removing dorsal skin and muscle layers, followed by precise cutting of the vertebral bone processes to expose the DRGs. The excised DRGs were trimmed at their roots and enzymatically digested in 3 mL sterile, bicarbonate-free, serum-free DMEM (Catalog# 11965, Thermo Fisher Scientific, Waltham, MA) supplemented with neutral protease (3.125 mg/mL, Catalog# LS02104, Worthington, Lakewood, NJ) and collagenase Type I (5 mg/mL, Catalog# LS004194, Worthington, Lakewood, NJ). The tissue was incubated for 45 minutes at 37°C with gentle agitation. After digestion, the dissociated DRG neurons (~1.5 × 10^6 cells) were collected by centrifugation, washed with DRG culture medium (DMEM supplemented with 1% penicillin/streptomycin sulfate from a 10,000 µg/mL stock, 30 ng/mL nerve growth factor, and 10% fetal bovine serum), and plated onto 12-mm coverslips coated with poly-D-lysine and laminin.

Spared nerve injury (SNI) Surgery and Von Frey test

Animals were anesthetized with isoflurane (5%) and placed on a heated blanket to preserve body temperature. Anesthesia was maintained (2.5%) throughout the surgery. Areflexia was confirmed before starting the surgery. Chlorhexidine was applied twice on the right leg, and a small incision (2-3 cm) was performed on the thigh. The biceps femoris muscle was bluntly dissected to expose the three terminal branches of the sciatic nerve. The common peroneal and tibial branches were tightly ligated with 4–0 silk and axotomized, leaving the sural branch intact. The muscle was sutured with 5–0 absorbable suture

and skin was auto clipped. Antibiotic ointment was applied on the skin. Animals were allowed to recover for 7 days before any testing.

Mechanical allodynia was measured by assessing the withdrawal threshold of the paw in response to probing with a series of calibrated fine (von Frey) filaments. Animals were placed in Plexiglas boxes on an elevated mesh screen and were given 15 minutes to acclimate. Filaments were applied on the plantar surface of both hind-paws in ascending forces until a withdrawal reflex or until the cut-off filament of 15g. The withdrawal threshold was determined by sequentially increasing and decreasing the stimulus strength (the "up and down" method), and data were analyzed with the nonparametric method of Dixon (*70*) and expressed as the mean withdrawal threshold.

Patch-Clamp Electrophysiology

Whole-cell patch-clamp recordings were conducted using an EPC10 amplifier in conjunction with PATCHMASTER (RRID:SCR_000034) software (HEKA), employing a 5 kHz sampling rate. Borosilicate glass pipettes were fabricated with a P1000 micropipette puller (Sutter Instrument, Novato, CA), yielding an initial resistance of 2.5–4 MΩ. A Sutter MP-225A motorized micromanipulator was utilized for precise electrode positioning. Experiments were carried out at room temperature.

To measure HCN currents in DRG neurons, the internal pipette solution was composed of (in mM): 10 NaCl, 137 KCl, 4 Mg-ATP, 10 HEPES, 1 EGTA, and adjusted to pH 7.3 with KOH. The 4 mM Mg-ATP is sufficient to prevent $PI(4,5)P_2$ hydrolysis-mediated current rundown during the whole-cell patch-clamp experiments (*71, 72*). The extracellular solution consisted of (in mM): 154 NaCl, 5.6 KCl, 1 MgCl2, 1 CaCl2, 8 HEPES, and 10 D-glucose, with pH set to 7.4 using NaOH. Series resistance was consistently maintained below 10 MΩ, with compensation adjusted between 40% and 70%. The conductance-voltage (G-V) relationship of endogenous HCN currents was derived from instantaneous tail currents measured at -60 mV following test pulses ranging from -40 mV to between 0 and -140 mV. Leak-subtracted tail currents at -60 mV were normalized to the maximum tail current amplitude to determine relative conductance (G/G_{max}). These values were plotted as a function of the test pulse voltage and fitted to the Boltzmann equation: $G/G_{max} = 1/(1 + exp[(V − V_{1/2})/V_S])$, where V represents membrane potential, $V_{1/2}$ denotes the voltage for half-maximal activation, and Vs is the slope factor. The slope factor was defined as Vs = RT/zeF, where R is the gas constant, T is the absolute temperature (297 K), F is the Faraday constant, *e* is the elementary charge, and z represents the estimated gating charge per channel required for activation. Action potentials were recorded in PATCHMASTER using current-clamp mode rather than voltage clamp (*10*). Resting membrane potential was set with zero current injection. Absolute current

injections were then applied to induce action potentials. DRG neurons showed current-dependent adaptation, with firing rates decreasing as injected current increased.

For whole-cell Na^+ channel current recordings, we used a reduced extracellular sodium concentration, as established in previous studies, to decrease the electrochemical driving force and enable more effective voltage clamping of fast Na^+ currents (73). The extracellular solution consisted of (in mM): 35 NaCl, 65 Choline-Cl, 30 TEA (tetraethylammonium)-Cl, 0.1 $CaCl_2$, 5 $MgCl_2$, 0.1 $CdCl_2$, 10 HEPES, and 10 glucose, pH was adjusted to 7.4. The internal pipette solution contained (in mM): 100 Cs-MeS (cesium methanesulfonate), 40 TEA-Cl, 5 NaCl, 1 $CaCl_2$, 11 EGTA, 10 HEPES, 2 Mg-ATP, and 1 GTP, pH was adjusted to 7.4. A p/−4 leak subtraction method was used to cancel passive capacitive transients. The peak current amplitude at each voltage step was used to construct the G-V relationship for channel activation, after correcting for the voltage-dependent driving force. To estimate the driving force at each membrane potential, we determined the reversal potential by linearly extrapolating peak current amplitudes recorded at depolarized voltages to the voltage at which the net current was zero. Steady-state inactivation was assessed using a voltage protocol consisting of pre-pulse steps from –70 mV to 0 mV, followed by a main test pulse at 0 mV. Peak current amplitudes during the main pulse were normalized to the maximal response and used to estimate post-inactivation conductance.

Statistics and Reproducibility

Data parameters are expressed as mean ± s.e.m. from independent cells or patches. All successful single-cell imaging and patch clamp recordings were included in the analysis; no successful datasets were excluded. Sample size was based on previously published results using patch-clamp recordings of cell lines and primary cells, rat behavior tests, or applying ensemble FRET that generated sufficient statistical power. All experiments were conducted randomly and confirmed to be reproducible. Key experiments were conducted with at least two independent cell transfections or surgical preparations from animals. The specific number of independent replicates is provided in the figure legends. For single-cell imaging or patch-clamp experiments, statistical significance was assessed using paired t-test for comparisons between before and after treatments. For behavior experiments, two-way ANOVA or Mann-Whitney tests were used. Statistical thresholds were set at $^*p < 0.05$ and $^{**}p < 0.01$, respectively.

ACKNOWLEDGEMENTS

This research is supported by National Institute of General Medical Sciences R35GM154778 grant (to G.D.), and by the Doisy Fund (to G.D. and L.J.H.) of the Edward A. Doisy Department of Biochemistry and Molecular Biology at Saint Louis University School of Medicine. The Moutal lab is supported by startup funds from the Department of Pharmacology and Physiology, Saint Louis University, SLU Institute for Drug & Biotherapeutic Innovation seed grant, National Institute of Neurological Disorders and Stroke R01NS119263 grant, and Department of Defense CPMRP #CP220080P1 grant. Cartoons in figures were created using BioRender.

DATA AVAILABILITY

Source data are provided as a Source Data file (also in figshare.com as DOI:10.6084/m9.figshare.29983183), and additional data are in the Supplementary Information.

AUTHOR CONTRIBUTIONS

G.D. conceptualized the research and designed experiments. L. J. H. performed electrophysiology experiments, G.D. performed fluorescence imaging. C.G., N.L.A.D., L.S., and A.M. performed and analyzed behavior experiments on rats and prepared neuronal samples. G.D. and A.M. wrote the paper.

COMPETING INTERESTS

The authors declare no competing interests.

REFERENCES

1. S. L. Veatch, S. L. Keller, Organization in lipid membranes containing cholesterol. *Phys Rev Lett* **89**, 268101 (2002).
2. K. Simons, R. Ehehalt, Cholesterol, lipid rafts, and disease. *J Clin Invest* **110**, 597-603 (2002).
3. S. L. Liu *et al.*, Orthogonal lipid sensors identify transbilayer asymmetry of plasma membrane cholesterol. *Nat Chem Biol* **13**, 268-274 (2017).
4. E. Corvera, O. G. Mouritsen, M. A. Singer, M. J. Zuckermann, The permeability and the effect of acyl-chain length for phospholipid bilayers containing cholesterol: theory and experiment. *Biochim Biophys Acta* **1107**, 261-270 (1992).
5. W. K. Subczynski, M. Pasenkiewicz-Gierula, J. Widomska, L. Mainali, M. Raguz, High Cholesterol/Low Cholesterol: Effects in Biological Membranes: A Review. *Cell Biochem Biophys* **75**, 369-385 (2017).
6. D. Lingwood, K. Simons, Lipid rafts as a membrane-organizing principle. *Science* **327**, 46-50 (2010).
7. A. Das, M. S. Brown, D. D. Anderson, J. L. Goldstein, A. Radhakrishnan, Three pools of plasma membrane cholesterol and their relation to cholesterol homeostasis. *Elife* **3**, (2014).
8. M. Kinnebrew *et al.*, Cholesterol accessibility at the ciliary membrane controls hedgehog signaling. *Elife* **8**, (2019).
9. H. Wang *et al.*, Regulation of β-amyloid production in neurons by astrocyte-derived cholesterol. *Proc Natl Acad Sci U S A* **118**, (2021).
10. L. J. Handlin *et al.*, Membrane lipid nanodomains modulate HCN pacemaker channels in nociceptor DRG neurons. *Nat Commun* **15**, 9898 (2024).
11. C. V. Robinson, T. Rohacs, S. B. Hansen, Tools for Understanding Nanoscale Lipid Regulation of Ion Channels. *Trends Biochem Sci* **44**, 795-806 (2019).
12. I. Levitan, D. K. Singh, A. Rosenhouse-Dantsker, Cholesterol binding to ion channels. *Front Physiol* **5**, 65 (2014).
13. K. M. Beverley, I. Levitan, Cholesterol regulation of mechanosensitive ion channels. *Front Cell Dev Biol* **12**, 1352259 (2024).
14. O. Furst, N. D'Avanzo, Isoform dependent regulation of human HCN channels by cholesterol. *Sci Rep* **5**, 14270 (2015).
15. I. Levental, E. Lyman, Regulation of membrane protein structure and function by their lipid nano-environment. *Nat Rev Mol Cell Biol* **24**, 107-122 (2023).
16. D. W. Hilgemann *et al.*, Lipid signaling to membrane proteins: From second messengers to membrane domains and adapter-free endocytosis. *J Gen Physiol* **150**, 211-224 (2018).
17. E. C. Emery, G. T. Young, E. M. Berrocoso, L. Chen, P. A. McNaughton, HCN2 ion channels play a central role in inflammatory and neuropathic pain. *Science* **333**, 1462-1466 (2011).
18. E. C. Emery, G. T. Young, P. A. McNaughton, HCN2 ion channels: an emerging role as the pacemakers of pain. *Trends Pharmacol Sci* **33**, 456-463 (2012).
19. S. Lainez, C. Tsantoulas, M. Biel, P. A. McNaughton, HCN3 ion channels: roles in sensory neuronal excitability and pain. *J Physiol* **597**, 4661-4675 (2019).
20. L. J. Handlin, G. Dai, Direct regulation of the voltage sensor of HCN channels by membrane lipid compartmentalization. *Nat Commun* **14**, 6595 (2023).
21. D. DiFrancesco, Characterization of single pacemaker channels in cardiac sino-atrial node cells. *Nature* **324**, 470-473 (1986).
22. A. Saponaro *et al.*, Gating movements and ion permeation in HCN4 pacemaker channels. *Mol Cell* **81**, 2929-2943 e2926 (2021).
23. W. N. Zagotta *et al.*, Structural basis for modulation and agonist specificity of HCN pacemaker channels. *Nature* **425**, 200-205 (2003).
24. K. B. Craven, W. N. Zagotta, CNG and HCN channels: two peas, one pod. *Annu Rev Physiol* **68**, 375-401 (2006).

25. A. Momin, H. Cadiou, A. Mason, P. A. McNaughton, Role of the hyperpolarization-activated current Ih in somatosensory neurons. *J Physiol* **586**, 5911-5929 (2008).
26. A. Saponaro, G. Thiel, A. Moroni, Structural and functional approaches to studying cAMP regulation of HCN channels. *Biochem Soc Trans* **49**, 2573-2579 (2021).
27. S. R. Chaplan *et al.*, Neuronal hyperpolarization-activated pacemaker channels drive neuropathic pain. *J Neurosci* **23**, 1169-1178 (2003).
28. G. T. Young, E. C. Emery, E. R. Mooney, C. Tsantoulas, P. A. McNaughton, Inflammatory and neuropathic pain are rapidly suppressed by peripheral block of hyperpolarisation-activated cyclic nucleotide-gated ion channels. *Pain* **155**, 1708-1719 (2014).
29. A. Barbuti *et al.*, Localization of pacemaker channels in lipid rafts regulates channel kinetics. *Circ Res* **94**, 1325-1331 (2004).
30. T. Matsubara *et al.*, Heterogeneous Ganglioside-Enriched Nanoclusters with Different Densities in Membrane Rafts Detected by a Peptidyl Molecular Probe. *Langmuir* **37**, 646-654 (2021).
31. R. Sachl *et al.*, On multivalent receptor activity of GM1 in cholesterol containing membranes. *Biochim Biophys Acta* **1853**, 850-857 (2015).
32. D. H. Z. Koh *et al.*, Visualization of accessible cholesterol using a GRAM domain-based biosensor. *Nat Commun* **14**, 6773 (2023).
33. B. Ercan, T. Naito, D. H. Z. Koh, D. Dharmawan, Y. Saheki, Molecular basis of accessible plasma membrane cholesterol recognition by the GRAM domain of GRAMD1b. *EMBO J* **40**, e106524 (2021).
34. T. Naito *et al.*, Movement of accessible plasma membrane cholesterol by the GRAMD1 lipid transfer protein complex. *Elife* **8**, (2019).
35. S. Endapally *et al.*, Molecular Discrimination between Two Conformations of Sphingomyelin in Plasma Membranes. *Cell* **176**, 1040-1053 e1017 (2019).
36. M. Skocaj *et al.*, Tracking cholesterol/sphingomyelin-rich membrane domains with the ostreolysin A-mCherry protein. *PLoS One* **9**, e92783 (2014).
37. J. Myeong, C. G. Park, B. C. Suh, B. Hille, Compartmentalization of phosphatidylinositol 4,5-bisphosphate metabolism into plasma membrane liquid-ordered/raft domains. *Proc Natl Acad Sci U S A* **118**, (2021).
38. G. Dai, Neuronal KCNQ2/3 channels are recruited to lipid raft microdomains by palmitoylation of BACE1. *J Gen Physiol* **154**, (2022).
39. I. Decosterd, C. J. Woolf, Spared nerve injury: an animal model of persistent peripheral neuropathic pain. *Pain* **87**, 149-158 (2000).
40. A. N. Bader *et al.*, Homo-FRET imaging as a tool to quantify protein and lipid clustering. *Chemphyschem* **12**, 475-483 (2011).
41. A. Joshi *et al.*, Intermolecular energy migration via homoFRET captures the modulation in the material property of phase-separated biomolecular condensates. *Nat Commun* **15**, 9215 (2024).
42. S. Mahammad, I. Parmryd, Cholesterol depletion using methyl-β-cyclodextrin. *Methods Mol Biol* **1232**, 91-102 (2015).
43. A. E. Christian, M. P. Haynes, M. C. Phillips, G. H. Rothblat, Use of cyclodextrins for manipulating cellular cholesterol content. *J Lipid Res* **38**, 2264-2272 (1997).
44. J. Kusch *et al.*, Interdependence of receptor activation and ligand binding in HCN2 pacemaker channels. *Neuron* **67**, 75-85 (2010).
45. B. Yu *et al.*, Cryo-EM structure of human HCN3 channel and its regulation by cAMP. *J Biol Chem* **300**, 107288 (2024).
46. C. H. Lee, R. MacKinnon, Structures of the Human HCN1 Hyperpolarization-Activated Channel. *Cell* **168**, 111-120 e111 (2017).
47. C. H. Lee, R. MacKinnon, Voltage Sensor Movements during Hyperpolarization in the HCN Channel. *Cell* **179**, 1582-1589 e1587 (2019).
48. M. Alsaloum *et al.*, Voltage-gated sodium channels in excitable cells as drug targets. *Nat Rev Drug Discov*, (2025).

49. W. A. Catterall, Structure and function of voltage-gated sodium channels at atomic resolution. *Exp Physiol* **99**, 35-51 (2014).
50. S. Albani *et al.*, Depletion of membrane cholesterol modifies structure, dynamic and activation of Na(v)1.7. *Int J Biol Macromol* **278**, 134219 (2024).
51. M. Amsalem, C. Poilbout, G. Ferracci, P. Delmas, F. Padilla, Membrane cholesterol depletion as a trigger of Nav1.9 channel-mediated inflammatory pain. *EMBO J* **37**, (2018).
52. G. Wisedchaisri *et al.*, Resting-State Structure and Gating Mechanism of a Voltage-Gated Sodium Channel. *Cell*, (2019).
53. I. Decosterd, C. J. Woolf, Spared nerve injury: an animal model of persistent peripheral neuropathic pain. *Pain* **87**, 149-158 (2000).
54. A. Moutal *et al.*, CRISPR/Cas9 editing of Nf1 gene identifies CRMP2 as a therapeutic target in neurofibromatosis type 1-related pain that is reversed by (S)-Lacosamide. *Pain* **158**, 2301-2319 (2017).
55. C. Tsantoulas *et al.*, Hyperpolarization-activated cyclic nucleotide-gated 2 (HCN2) ion channels drive pain in mouse models of diabetic neuropathy. *Sci Transl Med* **9**, eaam6072 (2017).
56. N. P. Poolos, M. Migliore, D. Johnston, Pharmacological upregulation of h-channels reduces the excitability of pyramidal neuron dendrites. *Nat Neurosci* **5**, 767-774 (2002).
57. W. Hu, B. P. Bean, Differential Control of Axonal and Somatic Resting Potential by Voltage-Dependent Conductances in Cortical Layer 5 Pyramidal Neurons. *Neuron* **97**, 1315-1326 e1313 (2018).
58. K. W. Ko, M. N. Rasband, V. Meseguer, R. H. Kramer, N. L. Golding, Serotonin modulates spike probability in the axon initial segment through HCN channels. *Nat Neurosci* **19**, 826-834 (2016).
59. D. V. Vasylyev, S. Liu, S. G. Waxman, I_h current stabilizes excitability in rodent DRG neurons and reverses hyperexcitability in a nociceptive neuron model of inherited neuropathic pain. *J Physiol* **601**, 5341-5366 (2023).
60. M. S. George, L. F. Abbott, S. A. Siegelbaum, HCN hyperpolarization-activated cation channels inhibit EPSPs by interactions with M-type K^+ channels. *Nat Neurosci* **12**, 577-584 (2009).
61. D. O. Kwarteng *et al.*, Cholesterol affects the binding of proteins to phosphatidic acid without influencing its ionization properties. *J Lipid Res* **66**, 100749 (2025).
62. M. Delgado-Ramirez, A. L. Lopez-Serrano, S. Sanchez-Armass, U. Meza, A. A. Rodriguez-Menchaca, Crosstalk between cholesterol and PIP_2 in the regulation of Kv7.2/Kv7.3 channels. *Biol Chem* **405**, 161-165 (2024).
63. D. M. Taglieri, D. A. Delfin, M. M. Monasky, Cholesterol regulation of PIP(2): why cell type is so important. *Front Physiol* **3**, 492 (2012).
64. L. J. Pike, J. M. Miller, Cholesterol depletion delocalizes phosphatidylinositol bisphosphate and inhibits hormone-stimulated phosphatidylinositol turnover. *J Biol Chem* **273**, 22298-22304 (1998).
65. M. A. Digman, V. R. Caiolfa, M. Zamai, E. Gratton, The phasor approach to fluorescence lifetime imaging analysis. *Biophys J* **94**, L14-16 (2008).
66. L. Malacrida, S. Ranjit, D. M. Jameson, E. Gratton, The Phasor Plot: A Universal Circle to Advance Fluorescence Lifetime Analysis and Interpretation. *Annu Rev Biophys* **50**, 575-593 (2021).
67. S. Ranjit, L. Malacrida, D. M. Jameson, E. Gratton, Fit-free analysis of fluorescence lifetime imaging data using the phasor approach. *Nat Protoc* **13**, 1979-2004 (2018).
68. G. Dai, Z. M. James, W. N. Zagotta, Dynamic rearrangement of the intrinsic ligand regulates KCNH potassium channels. *J Gen Physiol* **150**, 625-635 (2018).
69. T. S. van Zanten, G. P. S, S. Mayor, Quantitative fluorescence emission anisotropy microscopy for implementing homo-fluorescence resonance energy transfer measurements in living cells. *Mol Biol Cell* **34**, tp1 (2023).
70. S. R. Chaplan, F. W. Bach, J. W. Pogrel, J. M. Chung, T. L. Yaksh, Quantitative assessment of tactile allodynia in the rat paw. *J Neurosci Methods* **53**, 55-63 (1994).

71.   G. Dai, H. Yu, M. Kruse, A. Traynor-Kaplan, B. Hille, Osmoregulatory inositol transporter SMIT1 modulates electrical activity by adjusting PI(4,5)P$_2$ levels. *Proc Natl Acad Sci U S A* **113**, E3290-3299 (2016).
72.   P. Pian, A. Bucchi, A. Decostanzo, R. B. Robinson, S. A. Siegelbaum, Modulation of cyclic nucleotide-regulated HCN channels by PIP$_2$ and receptors coupled to phospholipase C. *Pflugers Arch* **455**, 125-145 (2007).
73.   X. Zhang, B. T. Priest, I. Belfer, M. S. Gold, Voltage-gated Na$^+$ currents in human dorsal root ganglion neurons. *Elife* **6**, (2017).