Enzymes define pathways and metabolic relationships

Jiada Zhan[1,2], Zachery R. Jarrell[2], Melissa L. Kemp[3], Jaclyn Weinberg[2], Young-Mi Go[1,2],
Greg S. Martin[2], Dean P. Jones[1,2*]
1: Nutrition and Health Sciences, Laney Graduate School, Emory University, Atlanta, Georgia, USA
2: Division of Pulmonary, Allergy, Critical Care and Sleep Medicine, Department of Medicine, Emory University,
Atlanta, Georgia, USA
3: The Wallace H. Coulter Department of Biomedical Engineering, Georgia Institute of Technology & Emory
University, Atlanta, Georgia, USA
*: corresponding author

Abstract

Gene-centric pathway mapping tools, widely used to interpret untargeted liquid chromatography

mass spectrometry (LC-MS) metabolomics data, may underperform because a single metabolite

can generate multiple spectral features, inflating false positive rates. Classic enzymology, which

established metabolite flow before gene sequencing, offers experimentally validated precursor-

product relationships that could overcome these ambiguities. We evaluated whether

enzymology-defined precursor-product correlations are consistently detectable in human plasma

LC-MS data. We detected amino acids, carnitine-related, TCA cycle, and pentose phosphate

pathway metabolites in one individual sampled eight times over five years and in 50 adults

sampled 6 to 8 times each. In the single participant repeated measures, strong positive

correlations were observed for most direct precursor-product pairs. The longitudinal and cross-

sectional analyses reproduced these patterns. Precursor-product proportionality, a fundamental

principle of enzymology, is detectable in LC-MS datasets and remains consistent across studies.

Applying these correlations to metabolomics workflows can improve pathway analysis, help

metabolite identification, and reveal how genetic variations, diets, therapeutic drugs, and

environmental exposures jointly impact metabolic pathways.

Introduction

Recently, five biochemical and analytical challenges have been identified that can cause gene-centric tools to underperform when applied to metabolomics data.[1] Untargeted liquid chromatography mass spectrometry (LC-MS) metabolomics-based pathway analysis often relies on pathway-mapping tools,[2,3] but ambiguities in mass spectral peak assignment – multiple peaks arising from the same metabolite – can generate high false positive rates.[4,5] Most metabolic pathways were established through enzymology before gene sequencing became available, and these studies offer direct, experimentally validated maps of metabolite flow. Thus, enzymology offers a simple but powerful way forward to enhance pathway analysis.

Enzymology established that metabolites flow from precursors to products and form dynamic linear, cyclical, and branched systems with regulation occurring at committed steps and branch points (**Fig. 1a**).[6] Here, we define a precursor-product relationship as a one-step biochemical conversion. Biochemical analyses showed that under steady-state conditions, the concentrations of metabolites often vary proportionately (see legend, **Fig. 1a**).[6,7] This principle was demonstrated in the pathway-level identification of xenobiotics and xenobiotic metabolites in blood[9] and endogenous metabolites in plant tissue.[7]

Results

Here, we show examples of precursor-product correlations observed from hydrophilic interaction liquid chromatography (HILIC) coupled to mass spectrometry with positive electrospray ionization (ESI) for amino acids (glutamine, glutamate, ornithine, citrulline, and arginine) and carnitine-related metabolites (butyrobetaine, carnitine, propionylcarnitine, O-acetylcarnitine,

palmitoylcarnitine) in one person's plasma sampled eight times across five years (**Fig. 1b**).[9]

These metabolites were confirmed by matching retention time (RT) and ion dissociation spectra

(MS/MS) to authentic standards. Among these examples, the lack of proportional change of

palmitoylcarnitine with carnitine illustrated that a positive correlation is not always present,

especially for metabolites at branch points and regulated sites.

To test whether positive precursor-product correlations exist at a population level, we analyzed

LC-MS plasma data from 50 individuals, each with six to eight visits.[9] Consistent with the data

found in the individual (**Fig. 1b**), amino acids and carnitine-related metabolites exhibit positive

correlations in longitudinal analyses across the 50 individuals (**Fig. 1c**), except for the lack of

proportional variation of palmitoylcarnitine with carnitine. Unlike these longitudinal

comparisons, however, most human metabolomics studies use a cross-sectional design with only

single LC-MS snapshots of individuals. To determine whether enzymology-defined precursor-

product correlations are present in a cross-sectional sampling, we restricted comparisons of the

50 individuals to include only baseline samples, and results showed the same relationships were

present (**Fig. 1d**).

To determine if these trends are upheld with reversed phase chromatography coupled to negative

ESI, we performed additional analyses for the TCA cycle (isocitric acid and alpha-ketoglutaric

acid) and pentose phosphate pathway metabolites (D-gluconate and 6-phosphogluconic acid)

with more subjects included. Isocitric acid and alpha-ketoglutaric acid showed significant

positive correlations in the longitudinal and cross-sectional analyses (**Fig. 2a**). A significant

correlation was observed between 6-phosphogluconic acid and D-gluconate (**Fig. 2b**), but not

between D-gluconate and D-glucono-1,5-lactone (**Fig. 2b**).

As an external validation, we analyzed published LC-MS human serum data from a study

consisting of 271 serum samples from 91 subjects with metastatic renal cell carcinoma treated

with nivolumab.[10] Nine RT and MS/MS confirmed precursor-product pairs were selected,

similar to the pairs in Figure 1. Consistent with Figure 1, many precursor-product pairs were

significantly correlated, while the pairs of decanoylcarnitine and carnitine and glutamate and

glutamine were not significantly correlated (**Fig. 2c**).

In pathway analysis with tools such as Mummichog[2], metabolite annotations have many-to-many

relationships that complicate their interpretation. To explore how precursor-product relationships

may enhance the pathway analysis, we performed untargeted pathway analysis with

Mummichog[2] in MetaboAnalyst[3] on participants with obesity in the CHDWB cohort (CHDWB

obesity).[11] Using one-way ANOVA analysis with the *limma* algorithm,[12] 1659 features (raw p-

value < 0.05) out of 13112 total features were used for pathway analysis (**Fig. 3a**). The

significantly altered valine, leucine, and isoleucine metabolism was selected for additional

analysis. A correlation heatmap of all features in the Mummichog algorithm revealed multiple

blocks of highly correlated features, with one block related to valine and leucine/isoleucine (**Fig.**

3c). The relevant network graph suggested three most relevant features associated with the

confirmed valine feature (m/z 118.0866, 65 s, [M+H]$^+$) (**Fig. 3c**). Among the two annotated

isoleucine features, one (m/z 132.1022, 51 s, [M+H]$^+$) showed a higher correlation (Spearman

correlation = 0.66, p < 0.0001) with the confirmed valine feature (m/z 118.0866, 65 s, [M+H]$^+$)

compared to the other correlations (**Fig. 3d**). Similarly, among the three annotated valine features,

the confirmed valine feature (m/z 118.0866, 65 s, [M+H]$^+$) showed a higher correlation with the

confirmed isoleucine feature (m/z 132.1022, 51 s, [M+H]$^+$) compared to the other correlations

(**Fig. 3d**). These metabolites do not have enzymatic precursor-product relationships but are

exchanged between cells and plasma by common transport systems. This correlation approach

can be used to supplement pathway enrichment-guided annotations for improved accuracy.

Discussion

As metabolomics moves beyond static pathway maps, enzymology-defined precursor-product

correlations provide a mechanistically grounded view of metabolic pathways. Even though most

metabolism occurs in organ systems other than blood,[1] the plasma data (**Fig. 1, 2, 3**) showed that

precursor-product relationships are preserved despite variable contributions of organ systems

(**Fig. 1e**). Consistent precursor-product relationships on two in-house LC-MS datasets and an

external metabolomics dataset show these correlations can be used across study designs, study

populations, and various LC-MS methods.

Batch correction is often used to decrease the impact of analytical variation in population studies.

We found that precursor-product correlations remained significant, although slightly attenuated,

by the addition of batch correction by the ComBat algorithm (xMSanalyzer[13]) for the

longitudinal and cross-sectional analyses of the 50 CHDWB subjects (**Supplementary Fig. 1**).

Given that the correlations in the study with 1 subject across 8 visits were mostly unchanged and

all were analyzed in the same batch, it appears that ComBat batch correction may unintentionally

disrupt biological covariations in the precursor-product relationships.

In comparison of glutamine and glutamate correlations among studies, we found that glutamine

and glutamate were positively correlated CHDWB healthy plasma study, but not in the cancer

serum study. This can occur because each of them is the precursor and product of many other

metabolites, or it could occur because disease can distort the precursor-product correlation. For

instance, cancer changes inter-organ glutamine trafficking by increasing the expression of

glutaminases,[14,15] which may explain the negative correlation result between glutamate and

glutamine in the cancer study. A negative correlation between glutamate and glutamine also

occurred in the CHDWB obesity study (data not shown), which may occur because glutamine is

low in the subjects with obesity[16,17]. These findings suggest that precursor-product relationships

in metabolomics analysis can have value for biological understanding even when precursors and

products are not positively correlated.

In pathway enrichment analyses, such as those provided by Mummichog, one feature can

represent many metabolites and complicate interpretation, even though the pathway-level results

are reliable. Inclusion of precursor-product relationships provides a useful approach to improve

reliability in the interpretation of the metabolite annotations. Importantly, the results with

leucine, isoleucine, valine pathway metabolites show that metabolites which share cell

transporter mechanisms can also display precursor-product like relationships even when they do

not have enzymatic precursor-product relationships. This demonstrates the potential to extend

the precursor-product relationship concept to metabolites sharing common transport systems,

which was also described in another study.[18]

One must be cautious about enzyme-based relationships;[6] some precursors and products do not

show proportional behavior, and some distantly related metabolites have strong correlations.[19]

Thus, it may be arbitrary to connect two features together and claim they form a precursor-

product relationship even if their chemical formula, chemical structures, and mass difference

seem to support the claim. This is especially true for computational metabolomics, where data-

driven metabolic network algorithms and in silico biochemical reactions were used for

metabolite annotation. As demonstrated by **Figure 3c**, false positives will be mixed with true

positives, and the number of annotation errors will increase via an iterative propagation process.

From our perspective, known one-step precursor-product and transporter relationships should be

prioritized compared to the data-driven metabolic network and in silico biochemical reaction

approaches.

Overall, incorporating the precursor-product relationships into metabolomics analysis can

improve pathway analysis and help metabolite identification. Extension to include metabolite

transport systems and development of tools to improve capabilities will further realize the

potential for comprehensive metabolomics to be a central hub for systems biology.

Metabolomics reflects all endogenous and xenobiotic alterations, genetic variations, enzyme

activities, post-transcriptional regulation and protein modifications, diets, therapeutic drugs,

diseases, and environmental exposures derived from biologic precursor-product relationships.[20–

22] The present results show that precursor-product relationships are highly preserved under

different conditions and provide a valuable resource to enhance LC-MS-based metabolomics

research.

Resource availability

Lead contact

Further information and requests for resources should be directed to the corresponding author,

Dean P. Jones (dpjones@emory.edu).

Data and code availability

The datasets illustrated in this study were included in Supplementary Data 1 for Figure 1,

Supplementary Data 2 for Figure 2, and Supplementary Data 3 for Figure 3. Raw data is

available at the Metabolomics Workbench[23]:

https://dev.metabolomicsworkbench.org:22222/data/DRCCMetadata.php?Mode=Study&StudyI

D=ST004055&Access=GecQ2393. The study is scheduled to be released on 2026-07-01 or

before publication.

Acknowledgments

This work was supported by the National Institute of Environmental Health Sciences (R01

ES031980 and P30 ES019776) and the National Institute on Aging (AG080247, AG085279, and

U01 AG088658).

Author contributions

G.S.M. established the study cohort. D.P.J. provided access to the metabolomics data. J.Z.

analyzed the data. J.Z. and D.P.J. wrote the manuscript. Z.R.J., M.L.K., J.W., Y.M.G., G.S.M.,

and D.P.J. revised the manuscript. D.P.J. supervised all aspects of the project.

Declaration of interests

The authors declare no competing interests.

Figure Legends

Figure 1. Precursor-product relationships in human plasma. (**a**). Metabolism occurs in linear (left,

top), cyclic (left, middle), and branched pathways (left, bottom). For sequential enzyme reactions in an

open system at steady state, rates of precursor (S) input, formation of intermediate products (A, B, C, D),

and formation and elimination of end product (P) are equivalent. With the same reaction rates for all

steps, steady-state concentrations of intermediates [A, B, C, D] are related by the respective pseudo-first

order rate constants (k_j): $k_1[S] = k_2[A] = k_3[B] = k_4[C] = k_5[D]$. Because regulation typically occurs only

at branch points or at sites committing a hydrocarbon to a specific fate, this means that proportional

relationships of precursors and one-step intermediate products are expected. Noted, when there is no

intermediate product between a precursor and end product, such as glutamine and glutamate, the notation

of intermediate products should be disregarded. (**b**) Proportional intensities for precursor and product

metabolites in different pathways are evident in eight repeated blood samplings from an individual

collected over five years (blue). Examples are provided for plasma amino acids and carnitine-related

metabolites, measured by hydrophilic interaction liquid chromatography (HILIC) and mass spectrometry

(LC-MS) with positive ESI (See Reference 10 for methodological details)[24]. All the metabolites on the x-

axis are the precursors, and all the metabolites on the y-axis are the products. Glutamine, glutamate,

butyrobetaine, and carnitine are considered part of a linear pathway, ornithine, citrulline, and arginine are

considered part of a cyclical pathway (urea cycle), and carnitine, propionylcarnitine, O-acetylcarnitine,

and palmitoylcarnitine are considered part of a branched pathway (carnitine shuttle). Sample collection

involved overnight fasting for the first blood draw, but not for the others. Note that an example of a

precursor-product pair without proportional concentrations, palmitoylcarnitine and carnitine (bottom row),

is provided. (**c**) Proportional intensities for the same precursor and product pairs as in Panel b for 50

individuals with multiple repeated blood samplings across five years (red). (**d**) Cross-sectional analysis of

baseline samples for the 50 individuals in Panel **c** (black) shows that precursor-product correlations are

preserved for the same precursor and product pairs. (**e**) In multicompartment models, most metabolism

occurs in organ compartments rather than the blood. Despite this, the data in Panels **b** to **d** show that

proportional relationships of precursors and products are observed in blood plasma samples.

Figure 2. Additional precursor-product correlation tests in human plasma and serum. (**a**) Precursor-

product correlation analyses in the TCA cycle pathway. Isocitric acid and alpha-ketoglutaric acid were

tested for correlation. Data included eight repeated blood samplings from an individual collected over five

years (blue, left), 220 individuals with multiple repeated blood samplings across five years (red, middle),

and cross-sectional analysis of baseline samples for the 213 individuals in Panel **c** (black, right). (**b**)

Precursor-product correlation analyses in the pentose phosphate pathway for D-gluconate and D-glucono-

1,5-lactone (upper panel) and for 6-phosphoglyconic acid and D-gluconate (lower panels). Color schemes

were the same as Figure 2a. (**c**) Precursor-product correlation analyses in multiple pathways (linear,

cyclical, and branched) in an external validation dataset. This dataset represents a longitudinal study (3

timepoints) of 271 serum samples from 91 subjects with metastatic renal cell carcinoma treated with

nivolumab.

Figure 3. Untargeted pathway analysis in the CHDWB obesity study. (a) One-way ANOVA analysis

with the *limma* algorithm was used to find significantly altered metabolomics features among subjects

with metabolically healthy non-obesity (MHN), metabolically healthy obesity (MHO), and metabolically

unhealthy obesity (MUO). Only the log2 fold changes between subjects with MHN and MUO were used

in the volcano plot. A raw p-value of 0.05 was used to select significantly altered features. **(b)** Untargeted

pathway analysis using the Mummichog algorithm. Untargeted pathway analysis was performed using the

Mummichog version 1 (functional analysis without using retention time)[2] in the MetaboAnalyst[3]. In the

analysis, 1659 significantly altered features were used. **(c)** Correlation heatmap and network analysis

among all valine, leucine, and isoleucine metabolism-related features in the Mummichog algorithm

outputs. Spearman correlation was used in the correlation heatmap using the heatmap.2 function in the

gplots R package (left). Partial Least Squares analysis in the mixOmics package was used between the

confirmed valine feature and other features, and then a relevance network graph was drawn with a cutoff

of 0.1 (right). **(d)** Precursor-product correlation analysis between annotated valine and annotated

isoleucine features. Valine (*m/z* 118.0866, 65s, $[M+H]^+$) and isoleucine (*m/z* 132.1022, 51 s $[M+H]^+$)

were confirmed by RT and MS/MS of the authentic standard.

Supplementary Figure 1. Precursor-product relationships in human plasma using ComBat batch

corrected metabolomics data. (**a**) Precursor-product correlation analyses among metabolites in multiple

pathways for 8 blood collections for 1 subject across give years (blue). (**b**) Precursor-product correlation

analyses among metabolites in multiple pathways among 50 subjects with repeated blood collections (6-

8). (**c**) Precursor-product correlation analyses among metabolites in multiple pathways among 50 subjects

with a baseline blood sampling. Data from 50 individuals with baseline blood collection (black) were

used.

STAR☐Methods

Study population

The metabolomics data are from the Center for Health Discovery and Wellbeing (CHDWB)

study, a longitudinal cohort.[9] CHDWB enrolled 668 participants who are essentially healthy

employees at Emory University, and we selected 50 subjects with at least 6 visits in the first set

of analyses (Fig. 1), and 213 subjects with only one visit for the second set of analyses (CHDWB

obesity). Individuals with poorly controlled chronic disease or acute illness were excluded from

the study enrollment. The first visit required overnight fasting for the blood draw, while the rest

of the visits did not have the fasting requirement. The CHDWB study was reviewed and

approved under Emory Investigational Review Board (IRB approval No. 00007243).

Sample Preparation

50 μL of plasma sample was combined with 100 μL acetonitrile containing 11 stable-isotope

internal standards added ([13C6]-D-glucose, [15N]-indole, [15N,13C5]-L-methionine, [2-15N]-

L-lysine dihydrochloride, [15N]-choline chloride, [13C5]-L-glutamic acid, [13C7]-benzoic acid,

[15N]-L-tyrosine, [15N2]-uracil, [3,3-13C2]-cystine, and [trimethyl-13C3]-caffeine). The

mixture was kept on ice for 30 minutes, then spun at $14,000 \times g$ for 10 minutes at 4 °C to

precipitate proteins. The clear supernatant was transferred into autosampler vials and

immediately loaded into a 4 °C autosampler for analysis.

Metabolomics Instrumental Analysis

Metabolite profiling was performed using a Thermo Scientific Vanquish Duo HPLC system

coupled to a Thermo Scientific ID-X high-resolution mass spectrometer operating in positive‐

mode electrospray ionization. Each sample was injected in triplicate, with pooled plasma

references analyzed at the beginning, middle, and end of each sample batch to monitor analytical

drift. Separation employed a Waters BEH Amide column (2.1 × 100 mm, 1.7 µm) with mobile

phase A (0.1% formic acid, 1 mM ammonium acetate in water) and mobile phase B (95%

acetonitrile, 0.1% formic acid, 1 mM ammonium acetate). The flow rate was 0.3 mL/min,

starting at 90% B, linearly decreasing to 20% B between 0.6 and 3.15 min, then held for an

additional 5 min. The mass spectrometer was set to 120k resolution, scanning from *m/z* 85–1275

on the Orbitrap detector. Sheath, auxiliary, and sweep gas flows were 50, 10, and 1 arbitrary

units, respectively, with spray voltages of 3.5 kV. MS/MS spectra were acquired in a separate

technical replicate using parallel reaction monitoring with targeted inclusion lists. Higher‑

energy collisional dissociation was performed at 35% normalized collision energy, collecting

MS[1] scans at 60k resolution and MS/MS scans at 30k resolution. For the CHDWB obesity study,

in brief, the samples were analyzed by the HILIC chromatography paired with a Q Exactive HF

Hybrid Quadrupole-Orbitrap mass spectrometer with positive ionization mode.[11] For the cancer

study used for external validation, in brief, the samples were analyzed with HILIC paired to a Q

Exactive Orbitrap with positive ionization mode. Details were described elsewhere.[10]

Metabolite Identification

Metabolites included in this study were confirmed by accurate mass MS[1] signal (± 5 ppm),

coelution with authentic standards (± 30 s), and ion dissociation spectra (MS/MS) matching

authentic standards.[24] These metabolites were the level 1 identifications according to the

Metabolomics Standards Initiative and Schymanski criteria.[25,26]

Statistical Analysis

Relative intensities of the metabolites were used for the correlation analyses. The intensities were

median-summarized across triplicates, and non-detects were imputed using the half minimum

method during this process. The intensities were log10 transformed and then used for the

Spearman correlations between precursors and products. All precursor-product correlation

analysis results were presented regardless of their correlation coefficients and p-values. A raw p-

value of 0.05 was used in the Mummichog pathway analysis without p-value adjustment. A

cutoff of 0.1 was used in the relevant network graph analyses.

References

1. Lee, K.S., Su, X., and Huan, T. (2025). Metabolites are not genes — avoiding the misuse of
pathway analysis in metabolomics. Nat. Metab., 1–4. https://doi.org/10.1038/s42255-025-
01283-0.

2. Li, S., Park, Y., Duraisingham, S., Strobel, F.H., Khan, N., Soltow, Q.A., Jones, D.P., and
Pulendran, B. (2013). Predicting Network Activity from High Throughput Metabolomics.
PLOS Comput. Biol. *9*, e1003123. https://doi.org/10.1371/journal.pcbi.1003123.

3. Pang, Z., Lu, Y., Zhou, G., Hui, F., Xu, L., Viau, C., Spigelman, A.F., MacDonald, P.E.,
Wishart, D.S., Li, S., et al. (2024). MetaboAnalyst 6.0: towards a unified platform for
metabolomics data processing, analysis and interpretation. Nucleic Acids Res., gkae253.
https://doi.org/10.1093/nar/gkae253.

4. Giera, M., Aisporna, A., Uritboonthai, W., and Siuzdak, G. (2024). The hidden impact of in-
source fragmentation in metabolic and chemical mass spectrometry data interpretation. Nat.
Metab. *6*, 1647–1648. https://doi.org/10.1038/s42255-024-01076-x.

5. Domingo-Almenara, X., Montenegro-Burke, J.R., Benton, H.P., and Siuzdak, G. (2018).
Annotation: A Computational Solution for Streamlining Metabolomics Analysis. Anal. Chem.
90, 480–489. https://doi.org/10.1021/acs.analchem.7b03929.

6. Steuer, R., Kurths, J., Fiehn, O., and Weckwerth, W. (2003). Observing and interpreting
correlations in metabolomic networks. Bioinforma. Oxf. Engl. *19*, 1019–1026.
https://doi.org/10.1093/bioinformatics/btg120.

7. Rose, I.A., and Warms, J.V. (1966). Control of glycolysis in the human red blood cell. J. Biol.
Chem. *241*, 4848–4854.

8. Liu, K.H., Lee, C.M., Singer, G., Bais, P., Castellanos, F., Woodworth, M.H., Ziegler, T.R.,
Kraft, C.S., Miller, G.W., Li, S., et al. (2021). Large scale enzyme based xenobiotic
identification for exposomics. Nat. Commun. *12*, 5418. https://doi.org/10.1038/s41467-021-
25698-x.

9. Tabassum, R., Cunningham, L., Stephens, E.H., Sturdivant, K., Martin, G.S., Brigham, K.L.,
and Gibson, G. (2014). A Longitudinal Study of Health Improvement in the Atlanta CHDWB
Wellness Cohort. J. Pers. Med. *4*, 489–507. https://doi.org/10.3390/jpm4040489.

10. Li, H., Bullock, K., Gurjao, C., Braun, D., Shukla, S.A., Bossé, D., Lalani, A.-K.A., Gopal,
S., Jin, C., Horak, C., et al. (2019). Metabolomic adaptations and correlates of survival to
immune checkpoint blockade. Nat. Commun. *10*, 4346. https://doi.org/10.1038/s41467-019-
12361-9.

11. Lim, G., Jarrell, Z.R., Go, Y.-M., and Jones, D.P. (2025). Amino Acid Associations in
Healthy and Unhealthy Obesity. J. Nutr. *155*, 2643–2652.
https://doi.org/10.1016/j.tjnut.2025.04.037.

12. limma powers differential expression analyses for RNA-sequencing and microarray studies |
Nucleic Acids Research | Oxford Academic
https://academic.oup.com/nar/article/43/7/e47/2414268.

13. xMSanalyzer: automated pipeline for improved feature detection and downstream analysis of
large-scale, non-targeted metabolomics data - PMC
https://pmc.ncbi.nlm.nih.gov/articles/PMC3562220/.

14. Hensley, C.T., Wasti, A.T., and DeBerardinis, R.J. (2013). Glutamine and cancer: cell biology,
physiology, and clinical opportunities. J. Clin. Invest. *123*, 3678–3684.
https://doi.org/10.1172/JCI69600.

15. Ling, H.H., Pan, Y.-P., Fan, C.-W., Tseng, W.-K., Huang, J.-S., Wu, T.-H., Chou, W.-C.,
Wang, C.-H., Yeh, K.-Y., and Chang, P.-H. (2019). Clinical Significance of Serum Glutamine
Level in Patients with Colorectal Cancer. Nutrients *11*, 898.
https://doi.org/10.3390/nu11040898.

16. Petrus, P., Lecoutre, S., Dollet, L., Wiel, C., Sulen, A., Gao, H., Tavira, B., Laurencikiene, J.,
Rooyackers, O., Checa, A., et al. (2020). Glutamine Links Obesity to Inflammation in
Human White Adipose Tissue. Cell Metab. *31*, 375-390.e11.
https://doi.org/10.1016/j.cmet.2019.11.019.

17. Lecoutre, S., Maqdasy, S., Rizo-Roca, D., Renzi, G., Vlassakev, I., Alaeddine, L.M., Higos,
R., Jalkanen, J., Zhong, J., Zareifi, D.S., et al. (2024). Reduced adipocyte glutaminase
activity promotes energy expenditure and metabolic health. Nat. Metab. *6*, 1329–1346.
https://doi.org/10.1038/s42255-024-01083-y.

18. Deo, R.C., Hunter, L., Lewis, G.D., Pare, G., Vasan, R.S., Chasman, D., Wang, T.J., Gerszten,
R.E., and Roth, F.P. (2010). Interpreting metabolomic profiles using unbiased pathway
models. PLoS Comput. Biol. *6*, e1000692. https://doi.org/10.1371/journal.pcbi.1000692.

19. Go, Y.-M., Walker, D.I., Soltow, Q.A., Uppal, K., Wachtman, L.M., Strobel, F.H., Pennell, K.,
Promislow, D.E.L., and Jones, D.P. (2015). Metabolome-wide association study of
phenylalanine in plasma of common marmosets. Amino Acids *47*, 589–601.
https://doi.org/10.1007/s00726-014-1893-x.

20. Baillie, T.A. (2008). Metabolism and Toxicity of Drugs. Two Decades of Progress in
Industrial Drug Metabolism. Chem. Res. Toxicol. *21*, 129–137.
https://doi.org/10.1021/tx7002273.

21. Playdon, M.C., Tinker, L.F., Prentice, R.L., Loftfield, E., Hayden, K.M., Van Horn, L.,
Sampson, J.N., Stolzenberg-Solomon, R., Lampe, J.W., Neuhouser, M.L., et al. (2024).
Measuring diet by metabolomics: a 14-d controlled feeding study of weighed food intake.
Am. J. Clin. Nutr. *119*, 511–526. https://doi.org/10.1016/j.ajcnut.2023.10.016.

22. Humphrey, S.J., James, D.E., and Mann, M. (2015). Protein Phosphorylation: A Major
Switch Mechanism for Metabolic Regulation. Trends Endocrinol. Metab. *26*, 676–687.
https://doi.org/10.1016/j.tem.2015.09.013.

23. Sud, M., Fahy, E., Cotter, D., Azam, K., Vadivelu, I., Burant, C., Edison, A., Fiehn, O.,
Higashi, R., Nair, K.S., et al. (2016). Metabolomics Workbench: An international repository
for metabolomics data and metadata, metabolite standards, protocols, tutorials and training,
and analysis tools. Nucleic Acids Res. *44*, D463–D470. https://doi.org/10.1093/nar/gkv1042.

24. Liu, K.H., Nellis, M., Uppal, K., Ma, C., Tran, V., Liang, Y., Walker, D.I., and Jones, D.P.
(2020). Reference Standardization for Quantification and Harmonization of Large-Scale
Metabolomics. Anal. Chem. *92*, 8836–8844. https://doi.org/10.1021/acs.analchem.0c00338.

25. Sumner, L.W., Amberg, A., Barrett, D., Beale, M.H., Beger, R., Daykin, C.A., Fan, T.W.-M.,
Fiehn, O., Goodacre, R., Griffin, J.L., et al. (2007). Proposed minimum reporting standards
for chemical analysis Chemical Analysis Working Group (CAWG) Metabolomics Standards
Initiative (MSI). Metabolomics Off. J. Metabolomic Soc. *3*, 211–221.
https://doi.org/10.1007/s11306-007-0082-2.

26. Schymanski, E.L., Jeon, J., Gulde, R., Fenner, K., Ruff, M., Singer, H.P., and Hollender, J.
(2014). Identifying Small Molecules via High Resolution Mass Spectrometry:
Communicating Confidence. Environ. Sci. Technol. *48*, 2097–2098.
https://doi.org/10.1021/es5002105.

a

1 Subject
8 visits

220 Subjects
6 to 8 visits

213 Subjects
1 visit

TCA Cycle

$r_s = 0.36$, p = 4.4e-01

$r_s = 0.52$, p < 1e-04

$r_s = 0.62$, p < 1e-04

Isocitric acid vs *Alpha-ketoglutaric acid*

b **Pentose Phosphate Pathway**

$r_s = -0.11$, p = 8.4e-01

$r_s = -0.14$, p = 1.3e-04

$r_s = -0.03$, p = 6.2e-01

D-gluconate vs *D-glucono-1,5-lactone*

$r_s = 0.47$, p = 2.8e-01

$r_s = 0.47$, p < 1e-04

$r_s = 0.41$, p < 1e-04

6-phosphogluconic acid vs *D-gluconate*

c **External Validation**

Linear Pathway

$r_s = -0.19$, p = 2e-03

$r_s = 0.32$, p < 1e-04

Glutamate vs *Glutamine*

Carnitine vs *Butyrobetaine*

Cyclical Pathway

$r_s = 0.5$, p < 1e-04

$r_s = 0.53$, p < 1e-04

Citrulline vs *Ornithine*

Ornithine vs *Arginine*

Branched Pathway

$r_s = 0.66$, p < 1e-04

$r_s = 0.43$, p < 1e-04

$r_s = 0.13$, p = 3.3e-02

Propionylcarnitine vs *Carnitine*

Butanoylcarnitine vs *Carnitine*

Octanoylcarnitine vs *Carnitine*

$r_s = 0.04$, p = 5.4e-01

$r_s = 0.13$, p = 2.8e-02

Decanoylcarnitine vs *Carnitine*

Palmitoylcarnitine vs *Carnitine*

a

MUO < MHN   MUO > MHN

b Pathway

Tyrosine
Valine, leucine and isoleucine
Tryptophan
Inositol phosphate
Caffeine
Lysine degradation
One carbon pool by folate
Phenylalanine
Ascorbate and aldarate

Pathway Overlap Size
8
20
25
41

Overlap Size Percentage
1.00
0.99
0.98
0.97
0.96

c Valine, leucine and isoleucine metabolism

Spearman correlation

Color key
-0.67   -0.1   0.67

Leucine/Isoleucine
Valine

Valine   Leucine/Isoleucine

d

$r_s = 0.66$, $p < 1e-04$

$r_s = 0.03$, $p = 6.5e-01$

$r_s = 0.21$, $p = 6.8e-04$

$r_s = 0.14$, $p = 2.8e-02$